

Holomorphic Explosions

Aslı Deniz¹ and Carsten Lunde Petersen²

¹American University of the Middle East, CBA, Department of
Mathematics and Statistics, Kuwait

²Roskilde University,
Department of Science and Environment, Denmark

Abstract

This paper concerns the problem of extending the parameter domain of holomorphic motions to include isolated boundary points, punctures of the domains. Supposing that the parameter domain has an isolated boundary point λ^* , we explore the extension properties of the holomorphic motion to λ^* .

1 Introduction

The notion of holomorphic motion was introduced by Mañé-Sad-Sullivan, in their work [M-S-S] on the dynamics of rational maps in order to construct conjugacies based on perturbations of the inclusion map of a subset of the sphere. Since then, this concept has been applied as a powerful tool in the field of holomorphic dynamics. A holomorphic motion in $\widehat{\mathbb{C}}$ can be described as a family of holomorphic functions with disjoint graphs over some domain $\Lambda \subset \widehat{\mathbb{C}}$ parametrized by the fiber over some point $\lambda_0 \in \Lambda$. More precisely, for a domain Λ in $\widehat{\mathbb{C}}$, a point $\lambda_0 \in \Lambda$, and a subset E of $\widehat{\mathbb{C}}$, the map $H : \Lambda \times E \rightarrow \widehat{\mathbb{C}}$ is called a *holomorphic motion of E* , parametrized by Λ , with base point λ_0 , if and only if the following conditions hold:

- i. $H(\lambda_0, z) = z$ for all $z \in E$,

- ii. H is vertically injective, i.e., for every fixed $\lambda \in \Lambda$, $H_\lambda : z \mapsto H(\lambda, z)$ is injective, and
- iii. H is horizontally holomorphic, i.e., for every fixed $z \in E$, $H^z : \lambda \mapsto H(\lambda, z)$ is holomorphic.

There are two famous extension results for holomorphic motions: *the λ -Lemma*, states that any given holomorphic motion $H : \Lambda \times E \rightarrow \widehat{\mathbb{C}}$ has a unique extension to a holomorphic motion of the closure \overline{E} , parametrized by Λ . The second well known result, called *the extended λ -Lemma* or *the Ślodkowski's Theorem* states that for simply connected $\Lambda \subset \widehat{\mathbb{C}}$, any holomorphic motion $H : \Lambda \times E \rightarrow \widehat{\mathbb{C}}$, extends to a holomorphic motion $H : \Lambda \times \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$. Moreover, for Λ a hyperbolic disk and for every parameter $\lambda \in \Lambda$, the extension H_λ is a quasiconformal homeomorphism, whose dilatation is bounded by $e^{d_\Lambda(\lambda_0, \lambda)}$, where $d_\Lambda(\cdot, \cdot)$ denotes the hyperbolic distance in Λ [Sł].

Let $\Lambda^* \subseteq \mathbb{C}$ be a domain with a puncture at λ^* , i.e., having an isolated boundary point $\lambda^* \in \mathbb{C}$. Given a holomorphic motion $H : \Lambda^* \times E \rightarrow \mathbb{C}$, when we consider the extension of H to λ^* , either holomorphicity of the functions H^z , or injectivity of H_{λ^*} may fail. Moreover, dynamical information may be lost in such an extension, if H is compatible with some dynamics. In this paper, we investigate the extension properties of H to $\Lambda := \Lambda^* \cup \{\lambda^*\}$, when E is connected. Moreover, we study the local representation $H^z(\lambda) = \sum_{n=0}^{\infty} a_n(z)(\lambda - \lambda^*)^n$ near λ^* , when the functions H^z extend holomorphically to λ^* and thus define an extension of H to Λ . We see that if $z \mapsto a_0(z)$ is injective, then the extension of H is a holomorphic motion parametrized by Λ . We show that if a_0 is not injective, then it is necessarily constant and hence the moving set E shrinks to a point at λ^* . We call the latter type of extension a *holomorphic explosion* and we study the properties of holomorphic explosions. We prove that for any holomorphic explosion there exists an underlying holomorphic motion $\widehat{H} : \Lambda \times E \rightarrow \mathbb{C}$, $n \geq 0$, and a polynomial P of degree $d \leq n - 1$ such that the local representation of a holomorphic explosion is $H_z(\lambda) = P(\lambda - \lambda^*) + (\lambda - \lambda^*)^n \widehat{H}(\lambda, z)$.

The organization of this article is as follows: In Section 2, we discuss extension properties of holomorphic motions to isolated boundary points of the parameter domain. This discussion leads to the introduction of the notion of holomorphic explosions. In the following two sections we give applications of this new concept to holomorphic dynamics.

This work was supported by Marie Curie RTN 035651-CODY and Roskilde University.

2 Main Extension Results of a Holomorphic Motion

Suppose a holomorphic motion H in \mathbb{C} is parametrized by a domain Λ^* , which has isolated boundary point λ^* such that $\Lambda = \Lambda^* \cup \{\lambda^*\} \subset \mathbb{C}$. We show that under mild conditions, the map H extends horizontally holomorphically to Λ . That is, each function H^z extends to a holomorphic function $H^z : \Lambda \rightarrow \mathbb{C}$ and so defines an extension of H to Λ . This is the subject of the following theorem.

Theorem 2.1. *Let $E \subset \mathbb{C}$ be connected, and Λ^* be a domain with an isolated boundary point λ^* such that $\Lambda = \Lambda^* \cup \{\lambda^*\} \subset \mathbb{C}$. Let $H : \Lambda^* \times E \rightarrow \mathbb{C}$ be a holomorphic motion over Λ^* with base point $\lambda_0 \in \Lambda^*$. Suppose there exist two distinct points $z_0, z_1 \in E$, such that H^{z_0} and H^{z_1} extend holomorphically to Λ . Then H extends horizontally holomorphically to Λ .*

Proof. Define

$$\tilde{H}(\lambda, z) := z_0 + (z_1 - z_0) \frac{H(\lambda, z) - H(\lambda, z_0)}{H(\lambda, z_1) - H(\lambda, z_0)}. \quad (1)$$

Then $\tilde{H}(\lambda, z_1) \neq \tilde{H}(\lambda, z_0)$ for all $\lambda \in \Lambda^*$, as $z_1 \neq z_0$. Thus for every $z \in E$, $\tilde{H}^z = \tilde{H}(\cdot, z)$ is holomorphic in Λ^* . Moreover, for every $\lambda \in \Lambda^*$, $\tilde{H}_\lambda = \tilde{H}(\lambda, \cdot)$ is injective, since \tilde{H}_λ is obtained from H_λ by post-composition with an injective affine map, which is the identity for $\lambda = \lambda_0$. So \tilde{H} is also a holomorphic motion in \mathbb{C} of E , parametrized by Λ^* and with base point λ_0 . Also $\tilde{H}(\lambda, z_i) \equiv z_i$ for $i = 0, 1$ by construction. Then \tilde{H} fixes the two distinct points z_0, z_1 in \mathbb{C} . Hence, for every $z \in E \setminus \{z_0, z_1\}$ the holomorphic function \tilde{H}^z takes its values in $\mathbb{C} \setminus \{z_0, z_1\}$. And so the point λ^* is a removable singularity for \tilde{H}^z by Picard's Theorem. That is, \tilde{H}^z extends meromorphically to Λ for every $z \in E$. This defines a horizontally meromorphic extension of \tilde{H} to $\Lambda \times E$.

We show that this extension of \tilde{H} is in fact horizontally holomorphic. That is, there is no point $z \in E$, such that $\tilde{H}^z(\lambda^*) = \infty$. Set $U := \{z \in E; \tilde{H}(\lambda^*, z) = \infty\}$. We prove that U is the empty set. To this end, we will show that U is both open and closed in the topology of E . Then as E is connected by hypothesis and $z_0, z_1 \in E \setminus U$ we conclude that $U = \emptyset$. For each j the subset $E_j := E \setminus \{z_j\}$ is a relatively open subset of E containing U . Let

γ denote a negatively oriented closed circular arc with center λ^* bounding a disk $\Gamma \subset \Lambda$. Define two functions $n_j : E_j \rightarrow \mathbb{Z}$ for $j = 0, 1$ by

$$n_j(z) := \frac{1}{2\pi i} \oint_{\gamma} \frac{\frac{\partial}{\partial \lambda} \tilde{H}(\lambda, z)}{\tilde{H}(\lambda, z) - z_j} d\lambda$$

Then each n_j is continuous, since $\gamma \subset \Lambda^*$, $z_j = \tilde{H}(\lambda, z_j)$ for all λ and \tilde{H}_λ is injective for all $\lambda \in \Lambda^*$. Hence each n_j is locally constant in E_j as it takes values in the discrete set \mathbb{Z} . Moreover, by the Argument Principle, the function n_j counts the number of poles minus the number of zeroes (with multiplicity) of $\lambda \mapsto \tilde{H}(\lambda, z) - z_j$ in Γ . And the only possible zero or pole is at λ^* . So that $n_j(z)$ is the order of λ^* as a pole of \tilde{H}^z if $n_j(z) > 0$, $-n_j(z)$ is the order of λ^* as a zero of \tilde{H}^z if $n_j(z) < 0$ and $n_j(z) = 0$ precisely if $z_j \neq \tilde{H}^z(\lambda^*) \in \mathbb{C}$.

To see that U is open, let $z \in U$ be arbitrary. Then $z \neq z_0$, $n_0(z) > 0$ and there exists a relatively open neighbourhood $\omega \subset E_0$ of z such that $n_0(w) = n_0(z) > 0$ for all $w \in \omega$, since n_0 is locally constant. As $z \in U$ was arbitrary, the set U is open in $E \setminus \{z_0\}$ and thus in E .

To see that U is closed, we prove that $E \setminus U$ is open in E . Since each E_j is open in E and $E = E_0 \cup E_1$ it suffices to prove that each subset $E_j \setminus U$ is open. Let $z \in E_j \setminus U$ for some j . Then $n_j(z) \leq 0$ and there exists a relatively open neighbourhood $\omega \subset E_0$ of z such that $n_0(w) = n_0(z) \leq 0$ for all $w \in \omega$, since n_j is locally constant. As $z \in E_j \setminus U$ was arbitrary the latter is open in E_j and thus in E . So $E \setminus U = (E_0 \setminus U) \cup (E_1 \setminus U)$ is also open.

To conclude, since \tilde{H} has a horizontally holomorphic extension to λ^* , so does H . \square

Complement 2.2. *With the hypotheses of Theorem 2.1 the map \tilde{H} defined by (1) extends to a holomorphic motion of E over Λ , that is \tilde{H}_{λ^*} is injective. Moreover there exist two holomorphic functions $f, g : \Lambda \rightarrow \mathbb{C}$ with $f(\lambda_0) = 0$, and $g(\lambda_0) = 1$, g never vanishing in Λ^* , such that*

$$H(\lambda, z) = f(\lambda) + g(\lambda) \cdot \tilde{H}(\lambda, z). \quad (2)$$

Proof. Let us first prove that the horizontally holomorphic extension of \tilde{H} to Λ , i.e., across λ^* is continuous on $\Lambda \times E$. Since it is an extension of a holomorphic motion over Λ^* of E we only need to prove continuity at the

points (λ^*, z) , where $z \in E$. However this follows from \tilde{H} being horizontally holomorphic. More precisely with Γ a round disk in Λ around λ^* with negatively oriented boundary γ as in the proof above we have

$$\tilde{H}(\lambda, z) = -\frac{1}{2\pi i} \oint_{\gamma} \frac{\tilde{H}(\lambda', z)}{\lambda' - \lambda} d\lambda'$$

Thus \tilde{H} is continuous on $\Gamma \times E$, since it is continuous on $\gamma \times E$.

Suppose towards a contradiction that there exist two distinct points $w_0, w_1 \in E$ such that $\tilde{H}_{\lambda^*}(w_0) = \tilde{H}_{\lambda^*}(w_1) = w$. Our aim is to show that $U_w := \{z \in E \mid \tilde{H}_{\lambda^*}(z) = w\}$ is the empty set. To do so, we shall show that U_w is both open and closed in the topology on E . Since \tilde{H}_{λ^*} is continuous it is closed. Thus we only need to prove it is also open.

Suppose U_w is not open. Then there exists a point $w_{\infty} \in U_w$ and a sequence $\{w_k\} \subset E \setminus U_w$ such that $w_k \rightarrow w_{\infty}$. Moreover either $w_{\infty} \neq w_0$ or $w_{\infty} \neq w_1$. We shall consider the first case, the second being similar. Define

$$B(\lambda, z) := \tilde{H}(\lambda, z) - \tilde{H}(\lambda, w_0).$$

on $\Lambda \times E$. Clearly, $\tilde{H}(\lambda, z)$ being a holomorphic motion over Λ^* , $B(\lambda, z)$ vanishes at λ^* only for $z \in U_w \setminus \{w_0\}$. Let

$$\epsilon = \inf_{\lambda \in \gamma} |B(\lambda, w_{\infty})|. \quad (3)$$

Since $B(\lambda, z)$ is continuous, and since $w_k \rightarrow w_{\infty}$, there exists $N \in \mathbb{N}$, such that for all $j \geq N$,

$$\forall \lambda \in \gamma : \quad |B(\lambda, w_{\infty}) - B(\lambda, w_j)| < \epsilon. \quad (4)$$

By assumption, $B(\lambda, w_j)$ has no zero at λ^* for all $j \geq N$. Considering (3) and (4), $B(\lambda, w_j)$ and $B(\lambda, w_{\infty})$ have the same number of zeroes in Γ , counted with multiplicity, by the Rouché's Theorem. This assures that, $B(\lambda, w_j)$ has at least one zero in Γ , which is not λ^* . This implies there exists $\lambda' \neq \lambda^*$ in Γ , such that $B(\lambda', w_j) = 0$, i.e., $\tilde{H}(\lambda', w_j) = \tilde{H}(\lambda', w_0)$. But this contradicts the fact that the map $\tilde{H}_{\lambda'}$ is injective for $\lambda' \neq \lambda^*$, being a holomorphic motion in $\Lambda^* \times E$. This proves that there is no sequence $\{w_k\} \subset E \setminus U_w$, with $w_k \rightarrow w_{\infty} \in U_w \setminus \{w_0\}$. Therefore, $U_w \setminus \{w_0\}$ is open. In case $w_{\infty} = w_0$, we apply the same procedure for $w_{\infty} \neq w_1$, and we similarly show that $U_w \setminus \{w_1\}$ is open. Therefore $U_w = (U_w \setminus \{w_0\}) \cup (U_w \setminus \{w_1\})$ is open.

Obviously, the set $E \setminus U_w$ is nonempty, as it contains either z_0 , z_1 , or both. Moreover it is open and closed by the above. And since E is connected, $U_w = \emptyset$.

Now defining

$$\begin{aligned} f(\lambda) &:= H(\lambda, z_0) - \frac{z_0}{z_1 - z_0} (H(\lambda, z_1) - H(\lambda, z_0)) \\ g(\lambda) &:= \frac{H(\lambda, z_1) - H(\lambda, z_0)}{z_1 - z_0} \end{aligned} \quad (5)$$

in (1), we obtain (2). □

Corollary 2.3. Given $z_0, z_1 \in E$:

- i. If $H(\lambda^*, z_0) \neq H(\lambda^*, z_1)$, then $g(\lambda^*) \neq 0$, and hence the extension of H to $\Lambda \times E$ is a holomorphic motion.
- ii. If $H(\lambda^*, z_0) = H(\lambda^*, z_1) = z^* \in \mathbb{C}$, then $g(\lambda^*) = 0$ and $f(\lambda^*) = z^*$, that is, for all $z \in E$, $H(\lambda^*, z) = z^*$.

Remark 2.4. In the form (2) neither of the functions f or g , nor the holomorphic motion \tilde{H} in the representation of H in (2) is unique. Also the map $\tilde{H}(\lambda^*, E)$ is only unique up to post-composition by a non-constant affine map. However let n denote the order of λ^* as a zero of g . Then n is unique and the Taylor polynomial of f around λ^* is unique. Indeed, g can be written as $g(\lambda) = (\lambda - \lambda^*)^n \cdot \hat{g}(\lambda)$, where \hat{g} is a nonvanishing holomorphic function on Λ . Let $P(\lambda - \lambda^*)$ denote the first n terms in the Taylor series of $f(\lambda)$ around λ^* . Then $f(\lambda) = P(\lambda - \lambda^*) + (\lambda - \lambda^*)^n \cdot \hat{f}(\lambda)$, where \hat{f} holomorphic in Λ . Then defining $\hat{H}(\lambda, z) := \hat{f}(\lambda) + \hat{g} \cdot \tilde{H}(\lambda, z)$

$$H(\lambda, z) = P(\lambda - \lambda^*) + (\lambda - \lambda^*)^n \cdot \hat{H}(\lambda, z) \quad (6)$$

Obviously \hat{H} can be reparametrized as a holomorphic motion over Λ with base point λ^* , of the set $E^* = \hat{H}(\lambda^*, E)$. Observe that the polynomial P of degree less than n and the holomorphic motion \hat{H} are unique.

Remark 2.5. The assumption that the set E is connected is essential for the conclusion of the Theorem 2.1 and its complement. In fact let $\Lambda = \mathbb{D}$, $\lambda^* = 0$,

so that $\Lambda^* = \mathbb{D}^*$. Let $\lambda_0 = 1/e$, set $E_H = \{0, 1/e, e\}$, $E_G = \{0, 1/e, 1/e^2\}$ and define two holomorphic motions over Λ^*

$$H(\lambda, z) = \begin{cases} 0, & z = 0, \\ \lambda, & z = 1/e, \\ 1/\lambda, & z = e. \end{cases} \quad \text{and} \quad G(\lambda, z) = \begin{cases} 0, & z = 0, \\ \lambda, & z = 1/e, \\ \lambda^2, & z = 1/e^2. \end{cases}$$

Then both H and G satisfies all the hypotheses in Theorem 2.1 except that the sets E_H and E_G are not connected. Moreover the function H^e has a pole at 0. Thus H extends horizontally meromorphically, but not horizontally holomorphically to \mathbb{D} . That is H does not satisfy the conclusion in Theorem 2.1. And G does satisfy the conclusion of Theorem 2.1, but does not satisfy the conclusion of Complement 2.2.

Remark 2.6. The assumption in Theorem 2.1 that for two distinct points $z_0, z_1 \in E$ the holomorphic functions H^{z_0} and H^{z_1} extend holomorphically to Λ is essential for the conclusion of the Theorem. Let $E = \mathbb{C}$, $\Lambda = \mathbb{D}$, $\lambda^* = 0$, so that $\Lambda^* = \mathbb{D}^*$ and define two holomorphic motions over Λ^*

$$H(\lambda, z) = (e^{1/\lambda} - e^{1/\lambda^*}) + z \quad \text{and} \quad G(\lambda, z) = e^{(1/\lambda - 1/\lambda^*)} \cdot z$$

Then neither H nor G satisfies the conclusion of Theorem 2.1.

Proposition 2.7. *Let $E \subset \mathbb{C}$ be a connected set containing at least two points, and let $H : \mathbb{C} \times E \rightarrow \mathbb{C}$ be a holomorphic motion of E over \mathbb{C} with base point λ_0 . Then there exist two holomorphic functions $f, g : \mathbb{C} \rightarrow \mathbb{C}$ with $f(\lambda_0) = 0$, g non-vanishing and $g(\lambda_0) = 1$, such that H has an expression*

$$H(\lambda, z) = f(\lambda) + g(\lambda) \cdot z \tag{7}$$

Proof. Let $z_0, z_1 \in E$ be distinct points. Consider the equation (2), where \tilde{H} is defined as in (1) from z_0, z_1 . Recall that \tilde{H} fixes both points z_0, z_1 . Hence for any $z \in E \setminus \{z_0, z_1\}$ the horizontal map $\tilde{H}^z : \mathbb{C} \rightarrow \mathbb{C}$ avoids the two points z_0, z_1 and thus is constant by the Picard's Little Theorem. Hence $\tilde{H}(\lambda, z) = z$ on $E \times \mathbb{C}$. Taking f and g as in (5) we obtain (7). \square

Motivated by Theorem 2.1 and its Complement 2.2 we define the notion of a *Holomorphic Explosion*, which is a relative of holomorphic motions.

Definition 2.8. (Holomorphic Explosion) Let $\Lambda \subsetneq \widehat{\mathbb{C}}$ be a domain and $E^* \subset \mathbb{C}$, let $(\lambda^*, z^*) \in \Lambda \times \mathbb{C}$ and let $n > 0$ be an integer. An order n holomorphic explosion of E^* in \mathbb{C} from (λ^*, z^*) is a horizontally holomorphic map $H : \Lambda \times E^* \rightarrow \mathbb{C}$ which can be written as

$$H(\lambda, z) = f(\lambda) + g(\lambda) \cdot \widehat{H}(\lambda, z), \quad (8)$$

where \widehat{H} is a holomorphic motion of E^* over Λ with base point λ^* , and $f, g : \Lambda \rightarrow \mathbb{C}$ are holomorphic functions such that $f(\lambda^*) = z^*$ and g admits λ^* as a zero of order n . We call λ^* , *the explosion parameter*.

Note that in the definition above

- The motion domain Λ can contain more than just one explosion parameter λ^* .
- The explosion parameter λ^* can be infinity. In this case the formula (8) can be replaced by (see also Remark 2.4)

$$H(\lambda, z) = P\left(\frac{1}{\lambda - \lambda'}\right) + \frac{1}{(\lambda - \lambda')^n} \cdot \widehat{H}(\lambda, z).$$

where P is a polynomial of degree at most $n - 1$ and $\lambda' \notin \Lambda$. In particular if $0 \notin \Lambda$ we can take $\lambda' = 0$ and have

$$H(\lambda, z) = P\left(\frac{1}{\lambda}\right) + \frac{1}{\lambda^n} \cdot \widehat{H}(\lambda, z). \quad (9)$$

- Another more complicated, but also more flexible formulation of an order n holomorphic explosion H of the set E^* is as follows. The map H is a horizontally analytic map $H : \Lambda \times E^* \rightarrow \mathbb{C}$, which can be written as

$$H(\lambda, z) = f(\lambda) + g(\lambda) \cdot \widehat{H}(\lambda, \psi(z)), \quad (10)$$

where \widehat{H} is a holomorphic motion of a set $E_0 \subset \mathbb{C}$ over Λ with base point $\lambda_0 \in \Lambda$, $\widehat{H}(\lambda^*, E_0) = E^*$, $\psi(z) = (\widehat{H}_{\lambda^*})^{-1}$ and $f, g : \Lambda \rightarrow \mathbb{C}$ are holomorphic functions such that $f(\lambda^*) = z^*$, g admits λ^* as a zero of order n and $g(\lambda_0) \neq 0$.

A simple application of the classical λ -Lemma shows that at similar conclusion holds for holomorphic explosions, that is the closure $\overline{E^*}$ of E^* in \mathbb{C} also exhibits a holomorphic explosion.

Lemma 2.9. (*λ -Lemma for holomorphic explosions*) *Let $E^* \subset \mathbb{C}$ and let $\Lambda \subsetneq \widehat{\mathbb{C}}$ be a domain. Suppose $H : \Lambda \times E^* \rightarrow \mathbb{C}$ is a holomorphic explosion from (λ^*, z^*) . Then H uniquely extends to a holomorphic explosion from (λ^*, z^*) of the closure $\overline{E^*}$ and parametrized by Λ .*

3 Application I

Consider the one parameter family of transcendental entire maps

$$f_a(z) = a(e^z(z-1) + 1), \quad a \in \mathbb{C}^*.$$

This family parametrizes the space of affine conjugacy classes of entire transcendental maps with two singular values, one of which coincides with a critical fixed point and the other being asymptotic.

We denote by A_a the basin of the superattracting fixed point 0. And we denote by A_a^0 the immediate basin, that is the connected component of A_a which contains 0. The asymptotic value of f_a is at $z = a$. The main hyperbolic component \mathcal{C}^0 in parameter space is the set of parameters for which the asymptotic value is contained in A_a^0 , that is,

$$\mathcal{C}^0 = \{a, a \in A_a^0\}.$$

It is proven in [De] that \mathcal{C}^0 is bounded, connected and $\mathcal{C}^0 \cup \{0\}$ is simply connected.

The Böttcher coordinate ϕ_a -associated to the fixed point $z = 0$ is the unique conformal conjugacy between f_a and $z \mapsto z^2$ in some neighborhood of 0. It has the form

$$\phi_a(z) = \frac{a}{2}z + O(z^2). \tag{11}$$

The conjugacy ϕ_a depends holomorphically on (a, z) and extends to the whole immediate basin A_a^0 if and only if $a \notin A_a^0$, or equivalently $a \notin \mathcal{C}^0$.

Denote by ψ_a the local inverse of ϕ_a mapping 0 to 0. Then for $a \notin \mathcal{C}^0$ the map ψ_a uniquely extends as a biholomorphic map $\psi_a : \mathbb{D} \rightarrow A_a^0$ which we denote a Böttcher parameter.




Figure 1: parameter plane in $[10, 10] \times [10, 10]$ - blue component in the center is \mathcal{C}^0 .




Figure 2: left: dynamical plane for $a = 3.7 + 0.5i$; right: dynamical plane for $a = 16.33 + 1.866i$ in $[10, 10] \times [10, 10]$ - blue component in the center is A_a^0 . Both dynamical planes are generated with parameters from $\mathbb{C} \setminus \overline{\mathcal{C}^0}$.

Let \mathcal{U}^* be the unbounded connected component of $\mathbb{C} \setminus \overline{\mathcal{C}^0}$. Then the Böttcher parameters $\psi_a(z)$ depends holomorphically on $(a, z) \in \mathcal{U}^* \times \mathbb{D}$. Fix $a_0 \in \mathcal{U}^*$ and define a holomorphic motion H of $A_{a_0}^0$, parametrized by \mathcal{U}^* and with base point a_0 via the Böttcher coordinates and parameters as follows:

$$\begin{aligned}
 H : \mathcal{U}^* \times A_{a_0}^0 &\rightarrow \mathbb{C} \\
 H(a, z) &= \psi_a \circ \phi_{a_0}(z),
 \end{aligned}
 \tag{12}$$

so that $H(a, A_{a_0}^0) = A_a^0$. Observe that it follows from (11) that

$$H(a, z) = \frac{a_0}{a}z + O(z^2). \quad (13)$$

Moreover note that ∞ is an isolated boundary point of the motion domain \mathcal{U}^* . Set $\mathcal{U} := \mathcal{U}^* \cup \{\infty\}$.

Our first application of holomorphic explosions is to H . Recall that a simply connected domain $A \subsetneq \mathbb{C}$ is called a K -quasi-disk, if there exists a K -quasi-conformal map $\psi : \mathbb{C} \rightarrow \mathbb{C}$ with $\psi(\mathbb{D}) = A$.

Theorem 3.1. *For $a \in \mathcal{U}^*$, A_a^0 is a K -quasidisk where*

$$K := e^{d_{\mathcal{U}}(\infty, a)} = \frac{1 + |k|}{1 - |k|}, \quad (14)$$

and where $d_{\mathcal{U}}(\cdot, \cdot)$ denotes the hyperbolic distance in \mathcal{U} .

This theorem strengthens a result in [De], where it is shown with a different approach that for all $a_0 \in \mathcal{U}$ the domain $A_{a_0}^0$ is a quasidisk. (observe Figure 2).

The proof uses that the holomorphic motion given by (12) exhibits a holomorphic explosion at ∞ . This is the content of the following Proposition.

Proposition 3.2. *The holomorphic motion $H : \mathcal{U}^* \times A_{a_0}^0 \rightarrow \mathbb{C}$ can be reparametrised in the vertical direction as a holomorphic explosion*

$$\mathcal{H} : \mathcal{U} \times \mathbb{D} \rightarrow \mathbb{C}$$

from $(\infty, 0)$ of order 1. More precisely, there exists a holomorphic motion $\widehat{H} : \mathcal{U} \times \mathbb{D} \rightarrow \mathbb{C}$ with base point ∞ such that

$$H(a, \psi_{a_0}(z)) = \mathcal{H}(a, z) = \frac{2}{a} \cdot \widehat{H}(a, z),$$

where $\widehat{H}_{a_0} = \psi_{a_0}$ is conformal.

Proof. Define $\widehat{H} : \mathcal{U}^* \times \mathbb{D} \rightarrow \mathbb{C}$ by $\widehat{H}(a, z) := \frac{a}{2}H(a, \psi_{a_0}(z)) = \frac{a}{2}\psi_a(z) =: \widehat{\psi}_a(z)$. We shall show that \widehat{H} extends to a holomorphic motion of \mathbb{D} over \mathcal{U} with base point ∞ . Then as $a \mapsto \frac{2}{a}$ has a simple zero at ∞ the theorem will follow.

Notice at first that \widehat{H} is a holomorphic function of the pair of variables (a, z) and that it is vertically univalent with

$$\widehat{H}(a, 0) = 0 \quad \text{and} \quad \frac{\partial}{\partial z} \widehat{H}(a, z)|_{z=0} = \widehat{\psi}'_a(0) = 1.$$

By the Koebe distortion estimates for univalent maps, we have

$$\forall z \in \mathbb{D}, \forall a \in \mathcal{U}^* \quad |\widehat{\psi}_a(z)| \leq \frac{|z|}{(1 - |z|)^2}$$

so that each map \widehat{H}^z is uniformly bounded near ∞ and so extends holomorphically to \mathcal{U} . That is \widehat{H} extends horizontally holomorphically to ∞ . Moreover the univalent functions on \mathbb{D} fixing 0 with derivative 1 form a compact family. It follows that the extension $\psi_\infty := \widehat{H}_\infty : \mathbb{D} \rightarrow \mathbb{C}$ is univalent. Thus the extension is also vertically injective. In order to show that \widehat{H} is a holomorphic motion with base point ∞ we need only show that $\widehat{\psi}_\infty = Id$. Or equivalently that the functions $\widehat{\psi}_a$ converge uniformly to the identity, as $a \rightarrow \infty$. To this end note that $\widehat{\psi}_a$ conjugates $z \mapsto z^2$ to

$$g_a(z) := \frac{a}{2} \cdot f_a\left(\frac{2z}{a}\right) = z^2 + \mathcal{O}(|4z^3/3a|).$$

In fact a standard but tedious computation shows that

$$|g_a(z) - z^2| \leq \left| \frac{4z^3}{3a} \right| e^{|2z/a|}.$$

So g_a converges to $g_\infty(z) := z^2$ locally uniformly in \mathbb{C} . By continuity, $\widehat{\psi}_\infty$ is the Böttcher parameter for g_∞ at 0 and hence $\widehat{\psi}_\infty = Id$. This completes the proof. \square

Proof of Theorem 3.1. The holomorphic motion \widehat{H} extends to a holomorphic motion of $\overline{\mathbb{D}}$ by the λ -Lemma and to a holomorphic motion of the Riemann sphere $\widehat{\mathbb{C}}$ by Ślodkowski's Theorem. We denote by \widehat{H} also such an extension. By the general properties of holomorphic motions over a simply connected domain such as \mathcal{U} , the real dilation of the vertical mapping \widehat{H}_a is bounded by $K_a := e^{d_{\mathcal{U}}(a, \infty)}$, when $\infty \in \mathcal{U}$ is the base point. Thus $A_a^0 = \frac{2}{a} \widehat{H}(a, \mathbb{D})$ is a K_a -quasi disk. \square

Remark 3.3. The holomorphic explosion \mathcal{H} extends uniquely to the holomorphic explosion $\mathcal{H} : \mathcal{U} \times \overline{\mathbb{D}} \rightarrow \mathbb{C}$, by Lemma 2.9.

Let $\{z_i\}_{i \in \mathbb{Z}}$ be the solution set of the equation $f_a(z) = 0$, ordered in increasing order of $\Im(z_i)$ with $z_0 = 0$. Let $A_a^{1,i}$ denote the connected component of $f_a^{-1}(A_a^0)$ which contains z_i . We call the point z_i , the center of $A_a^{1,i}$. Notice that these centers do not depend on the parameter.

Proposition 3.4. *For each $i \in \mathbb{Z} \setminus \{0\}$, there is a holomorphic explosion $\mathcal{H}^i : \mathcal{U} \times \overline{\mathbb{D}} \rightarrow \mathbb{C}$ from (∞, z_i) of order 2 with*

$$\mathcal{H}^i(a, \overline{\mathbb{D}}) = \overline{A}_a^{1,i}.$$

In particular all of the components $A_a^{1,i}$ of the basin of 0 for f_a are also K -quasi-disks with

$$K = e^{d_{\mathcal{U}}(\infty, a)}.$$

Proof. For each $a \in \mathcal{U}^*$ and $i \in \mathbb{Z} \setminus \{0\}$ let k_a^i be the inverse branch of f_a defined on \overline{A}_a^0 , which satisfies $k_a^i(0) = z_i$. And define

$$\begin{aligned} \mathcal{H}^i : \mathcal{U} \times \mathbb{D} &\rightarrow \mathbb{C} \\ \mathcal{U}^* \times \mathbb{D} \ni (a, z) &\mapsto k_a^i \circ \mathcal{H}(a, z) \\ \{\infty\} \times \mathbb{D} \ni (\infty, z) &\mapsto z_i. \end{aligned}$$

This gives the desired holomorphic explosions. From the series form of \mathcal{H}^i near 0

$$\mathcal{H}^i(a, z) = k_a^i(\mathcal{H}(a, z)) = z_i + \frac{2}{a^2 e^{z_i}} \widehat{H}(a, z) + O\left(\frac{1}{a^4}\right),$$

we see that the order is 2. Here \widehat{H} is as in Proposition 3.2. \square

4 Application II

We are interested in the moduli space \mathcal{M}_2 of quadratic rational maps modulo Möbius-conjugacy. Every quadratic rational map R has counting multiplicity three fixed points with multipliers which we generically denote by λ, μ and γ . Denote by $\sigma_1(R) = \lambda + \mu + \gamma$, $\sigma_2(R) = \lambda\mu + \lambda\gamma + \mu\gamma$ and $\sigma_3 = \lambda\mu\gamma$ the three elementary symmetric functions of λ, μ and γ . Following Milnor [Mi], we endow \mathcal{M}_2 with the affine structure given by (σ_1, σ_2) . In more detail $\sigma_3(R) = \sigma_1(R) - 2$ and any pair of numbers $(\sigma_1, \sigma_2) \in \mathbb{C}^2$ defines a unique equivalence class in \mathcal{M}_2 , so that $(\sigma_1, \sigma_2) : \mathcal{M}_2 \rightarrow \mathbb{C}^2$ is biholomorphic [Mi, Lemma 3.1]. For each $\lambda \in \mathbb{C}$ the curve

$$\text{Per}_1(\lambda) = \{ [R] \mid R \text{ has a fixed point of multiplier } \lambda \}.$$

is a straight line in this affine structure, [Mi, Lemma 3.4]. And moreover $\sigma(\lambda, R) := \mu(R)\gamma(R)$ defines an isomorphism between $\text{Per}_1(\lambda)$ and \mathbb{C} , [Mi, Remark 6.9].

For $\lambda \in \mathbb{D}$ we consider the connectedness locus M_λ in $\text{Per}_1(\lambda)$

$$M_\lambda = \{ [R] \in \text{Per}_1(\lambda) \mid J_R \text{ is connected} \},$$

where J_R denotes the Julia set of R .

The special line $\text{Per}_1(0)$ also equals the moduli space for quadratic polynomials

$$\text{Per}_1(0) = \{ [Q_c] \mid Q_c(z) = z^2 + c, c \in \mathbb{C} \}$$

for which $\sigma(Q_c) = 4c$, so that $\sigma(M_0) = 4 \cdot M$ where M is the Mandelbrot set. The external class of each R with $[R] \in \text{Per}_1(\lambda)$ is given by the Blaschke product $B_\lambda(z) = z \frac{z+\bar{\lambda}}{1+\lambda z}$.

Theorem 4.1. *There is a natural dynamically defined holomorphic motion $H : \mathbb{D} \times (4 \cdot M) \rightarrow \mathbb{C}$ over \mathbb{D} with base point 0 of the Mandelbrot set scaled by a factor 4 such that:*

1. For each $\lambda \in \mathbb{D} : H(\lambda, \sigma(0, M_0)) = \sigma(\lambda, M_\lambda)$.
2. For each $c \in M$ and $\lambda \in \mathbb{D}$, any quadratic rational map $R \in \sigma_\lambda^{-1}(H(\lambda, 4c))$ has a polynomial like restriction, hybridly equivalent to a restriction of Q_c .

For $\lambda \in \mathbb{C}^*$ and $B \in \mathbb{C}$ the quadratic rational map $z \mapsto \frac{1}{\lambda}(z + B + 1/z)$ admits ∞ as a fixed point with multiplier λ , and thus represents an element of $\text{Per}_1(\lambda)$. Moreover the maps with the same λ but with $+B$ and $-B$ are conjugate by $z \mapsto -z$, and thus represents the same element of $\text{Per}_1(\lambda)$. Therefore for each $\lambda \in \mathbb{C}^*$ and $A \in \mathbb{C}$ such that $B = \sqrt{A}$, the map

$$(\lambda, A) \mapsto [G_{\lambda, A}], \quad G_{\lambda, A}(z) := \frac{1}{\lambda}(z + \sqrt{A} + 1/z)$$

is well defined without specifying which square root is being used. When a specific choice of square root is relevant, it will be clear from the context, how it is chosen. The maps $G_{\lambda, A}$ admits ± 1 as critical points and a simple computation shows that the product of the two remaining fixed point eigenvalues is given by

$$\sigma_\lambda(A) = \sigma(\lambda, A) = \sigma(\lambda, [G_{\lambda, A}]) = \sigma_\lambda([G_{\lambda, A}]) = \frac{(\lambda - 2)^2 - A}{\lambda^2} \quad (15)$$

[Mi, page 73, column 1](Milnor uses the symbol τ for σ , a^2 for A/λ^2 and μ for λ .) It follows that for each fixed $\lambda \in \mathbb{C}^*$ the map

$$A \mapsto [G_{\lambda,A}] : \mathbb{C} \longrightarrow \text{Per}_1(\lambda)$$

is an isomorphism. We denote by \mathbf{A}_λ the inverse of this isomorphism and for $\lambda \in \mathbb{D}^*$ we write

$$\widehat{M}_\lambda := \mathbf{A}_\lambda(M_\lambda)$$

for the connectedness locus of the family $G_{\lambda,A}$, $A \in \mathbb{C}$. Equivalently \widehat{M}_λ is the set of parameters A for which, either 1, or -1 has bounded orbit under $G_{\lambda,A}$. Note that for any λ the map $G_{\lambda,0}$ is conjugate to itself under $z \mapsto -z$ and thus $0 \notin \widehat{M}_\lambda$ for any $\lambda \in \mathbb{D}^*$. In Figure 3, connectedness loci are shown with different λ values.




Figure 3: Connectedness loci in A -plane with λ parameter values top left: $0.4 - 0.35i$; top right $0.4 + 0.35i$; bottom left e^{-1} ; bottom right $0.2 + 0.2i$, in $[1, 5] \times [-2, 2]$.

Since the external class of the maps R with $[R] \in \text{Per}_1(\lambda)$ is B_λ , it follows from Yoccoz inequality for polynomial-like maps, [Pe][Cor. D2] that for any $r < 1$ the map σ_λ is bounded on

$$\bigcup_{\lambda, |\lambda| < r} M_\lambda.$$

It thus follows immediately from (15) that \mathbf{A}_λ converge to 4 uniformly on M_λ , when $\lambda \rightarrow 0$.

Holomorphic motion of the connectedness loci \widehat{M}_λ with the parameter λ was the subject of the master's thesis of Uhre. We adopt the following result from her thesis. The interested reader can see [Uh, Chapter 8.1-8.3] for a proof.

Theorem 4.2. *There exists a dynamically defined holomorphic motion of the connectedness locus $\widehat{M}_{e^{-1}}$ over \mathbb{D}^* with base point e^{-1} :*

$$\mathbb{A} : \mathbb{D}^* \times \widehat{M}_{e^{-1}} \rightarrow \mathbb{C} \quad \text{with} \quad \mathbb{A}_\lambda(\widehat{M}_{e^{-1}}) = \widehat{M}_\lambda.$$

More precisely for each fixed pair (λ, A) with $\lambda \in \mathbb{D}^*$ and $A \in \widehat{M}_{e^{-1}}$, there exists a global quasi-conformal mapping $h_{\lambda, A} : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ conjugating $G_{e^{-1}, A}$ to $G_{\lambda, \mathbb{A}(\lambda, A)}$ and which restricts to a hybrid conjugacy between quadratic like restrictions of the two maps.

In [Uh], the proof was made for the connectedness locus for the family given by the formula

$$R_{\lambda, \mu}(z) = z \frac{z + \mu}{1 + \lambda z}$$

which is conjugate to $G_{\lambda, A}$, where

$$A = (\lambda - 2)^2 - \lambda^2 \mu \frac{2 - \lambda - \mu}{1 - \lambda \mu} \tag{16}$$

by the affine map

$$\phi_{\lambda, \mu}(z) = \frac{\lambda z + 1}{\sqrt{1 - \lambda \mu}}.$$

Note that

$$\mu \frac{2 - \lambda - \mu}{1 - \lambda \mu} = \sigma \tag{17}$$

i.e., the product of the two other (than λ) fixed point eigenvalues.

Proof of Theorem 4.1. This is now a straight forward application of Corollary 2.3 to the result of Theorem 4.2. This approach provides an alternative to [Uh, Chapter 8.4]. Notice that the extension to 0 corresponds to the case where ∞ is a superattracting fixed point.

Since $\sigma_\lambda(A) = \sigma(\lambda, A) = \sigma(\lambda, [G_{\lambda, A}]) = \sigma(\lambda, \mathbf{A}_\lambda^{-1}(A)) = \sigma_\lambda(\mathbf{A}_\lambda^{-1}(A))$ is holomorphic as a function of the two variables and injective, in fact affine

in the second variable, (see (15)) the holomorphic motion \mathbb{A} is equivalent to the holomorphic motion of $\sigma_{e^{-1}}(M_{e^{-1}})$ over \mathbb{D}^* with base point e^{-1} :

$$\tilde{H} : \mathbb{D}^* \times \sigma_{e^{-1}}(M_{e^{-1}}) \longrightarrow \mathbb{C} \quad \tilde{H}(\lambda, \tau) = \sigma_\lambda \circ \mathbb{A}(\lambda, \sigma_{e^{-1}}^{-1}(\tau)) \quad (18)$$

This relation is illustrated by the following diagram:

$$\begin{array}{ccc}
 \widehat{M}_{e^{-1}} & \xrightarrow{\mathbb{A}} & \widehat{M}_\lambda \\
 \uparrow \mathbf{A}_{e^{-1}} & & \uparrow \mathbf{A}_\lambda \\
 M_{e^{-1}} & & M_\lambda \\
 \downarrow \sigma_{e^{-1}} & & \downarrow \sigma_\lambda \\
 \sigma_{e^{-1}}(M_{e^{-1}}) & \xrightarrow{\tilde{H}} & \sigma_\lambda(M_\lambda)
 \end{array}$$

$\sigma_{e^{-1}}$ (left curved arrow) σ_λ (right curved arrow)

The holomorphic motion \tilde{H} has two special points, whose motions are easy to follow. The center, for which one of the fixed points is persistently super attracting, i.e., it has multiplier 0 so that this trajectory is the constant 0. The other is the root corresponding to the two remaining fixed points persistently coalesce to form a parabolic fixed point of multiplier 1 so that this trajectory is the constant 1. That is, the motions of the points 0 and 1 under \tilde{H} are constants. Thus by Corollary 2.3, \tilde{H} extends to a holomorphic motion over \mathbb{D} . We denote this extension also by \tilde{H} . We write $E = \tilde{H}(0, \sigma_{e^{-1}}(M_{e^{-1}}))$ and denote by $H : \mathbb{D} \times E \longrightarrow \mathbb{C}$ its reparametrization from 0:

$$H(\lambda, \tau) = \tilde{H}(\lambda, \tilde{H}_0^{-1}(\tau)). \quad (19)$$

To prove Theorem 4.1 we just need to show that in fact $E = 4 \cdot M$. To see this note that $[G_{\lambda, A}]$ contains both of the maps $R_{\lambda, \mu}$ for which μ is a solution of the equation (16), which is quadratic in μ . When $\lambda \rightarrow 0$ the maps $R_{\lambda, \mu}$ converge to quadratic polynomials of the form $\mu z + z^2$ uniformly on the sphere. For any fixed $\tau = \sigma_{e^{-1}}(A') \in \sigma(e^{-1}, M_{e^{-1}})$ and any $\lambda \in \mathbb{D}^*$, let $\mu(\lambda, \tau)$ be a solution of the quadratic equation (16) with $A = \mathbb{A}(\lambda, A')$. Then the class $[G_{\lambda, \mathbb{A}(\lambda, A')}]$ contains $R_{\lambda, \mu(\lambda, \tau)}$. Thus for a continuous choice of $\mu(\lambda, \tau)$ this function converge to μ_0 such that for the quadratic polynomial $\mu_0 z + z^2$ the product of the two finite multipliers is $\tau_0 = \tilde{H}(0, \tau)$. This polynomial is affinely conjugate to $z^2 + c$ with $4c = \tau_0$ and has connected filled-in Julia set. Thus $E \subset 4 \cdot M$. Also this polynomial is the unique quadratic polynomial for which $R_{\lambda, \mu(\lambda, \tau)}$ and $G_{\lambda, \mathbb{A}(\lambda, A')}$ are hybridly equivalent according to [D-H, Prop. 14 page 313].

To complete the proof of Theorem 4.1 we need to show surjectivity, i.e $E = 4 \cdot M$. For this we appeal to the Proposition 4.3 below. It follows from the Proposition that for every $\tau_0 = 4c \in 4 \cdot M$ there exists $A' \in \widehat{M}_{e^{-1}}$ such that is $G_{e^{-1}, A'}$ is hybridly equivalent to Q_c . Thus $\tau_0 \in E$, i.e., $4 \cdot M \subset E$. \square

Proposition 4.3. *For every $c \in M$ there exists $A \in \widehat{M}_{e^{-1}}$ such that Q_c and $G_{e^{-1}, A}$ are hybridly equivalent.*

Proof. This is a particular case of [D-H, Prop. 5 page 301] applied to a quadratic like restriction of Q_c and the external class $B_{e^{-1}}$. \square

Then reapplying (15) again or applying Corollary 2.3 directly to \mathbb{A} we find

Corollary 4.4. The holomorphic motion \mathbb{A} of $\widehat{M}_{e^{-1}}$ over \mathbb{D}^* is a quasi-conformal reparametrization of a holomorphic explosion over \mathbb{D} from $(0, 4)$. Indeed writing $A(\tau) = \sigma_{e^{-1}}^{-1} \circ H(e^{-1}, \tau)$, then for any $\tau \in M_0$ we have

$$\mathbb{A}(\lambda, A(\tau)) = (\lambda - 2)^2 - \lambda^2 H(\lambda, \tau), \quad (20)$$

where H is the holomorphic motion in Theorem 4.1 as given by (19).

References

- [De] A. Deniz, Ph.D. thesis, *Entire Transcendental Maps with Two Singular Values*, Roskilde University, IMFUFA Text, 2014.
- [D-H] A. Douady and J. H. Hubbard, *On the Dynamics of Polynomial-like Mappings*, Ann. Sci. Ec. Norm. Sup. (4), vol. 18, no:2, 1985 287–343.
- [M-S-S] R. Mañé, P. Sad and D. Sullivan, *On the Dynamics of Rational Maps*, Ann. Sci. École Norm. Sup. (4), vol. 16, no:2, 1983, 193–217.
- [Mi] J. Milnor, *Geometry and Dynamics of Quadratic Rational maps*, Experiment. Math., vol. 2, no:1, 1993, 37–83.
- [Pe] C. L. Petersen, *On the Pommerenke-Levin-Yoccoz inequality*, Ergodic Theory Dynam. Systems, vol. 13, no: 4 1993, 785–806
- [Sł] Z. Ślodkowski, *Holomorphic motions and polynomial hulls*, Proc. Amer. Math. Soc., vol. 111, no: 2, 1991, 347–355.

- [Uh] E. Uhre, Master's thesis, *Construction of a Holomorphic Motion in Part of the Parameter Space for a Family of Quadratic Rational Maps*, Roskilde University, IMFUFA Text, 2004.