

LYAPUNOV SPECTRUM PROPERTIES AND CONTINUITY OF THE LOWER JOINT SPECTRAL RADIUS

REZA MOHAMMADPOUR

ABSTRACT. In this paper, we study ergodic optimization and multifractal behavior of Lyapunov exponents for matrix cocycles. We show the continuity of entropy spectrum at boundary of Lyapunov spectrum in the sense that $h_{top}(E(\alpha_t)) \rightarrow h_{top}(E(\beta(\mathcal{A})))$ for generic cocycles (in the sense of [BGMV]). We also show that for such cocycles over subshifts of finite type, the Lyapunov spectrum is equal to the closure of the set positive entropy spectrum. Moreover, we prove the restricted variational principle to level sets for such cocycles.

We prove the continuity of the lower joint spectral radius for general cocycles under the assumption that the linear cocycles admit a dominated splitting of index 1.

We show that the singular value pressure is continuous on Hölder and fiber-bunched $GL(2, \mathbb{R})$ -valued cocycles over subshifts of finite type. We also show that the Lyapunov spectrum is a closed and convex set for such cocycles over subshifts of finite type.

CONTENTS

1. Introduction and statement of the results	2
2. Preliminaries	6
2.1. Symbolic dynamics	6
2.2. Multilinear algebra	8
2.3. Fiber bunched cocycles	8
2.4. Typical cocycles	10
2.5. The continuity of Lyapunov exponents	12
3. Thermodynamic Formalism	14
3.1. Convex functions	14
3.2. Legendre transform	14
3.3. Topological entropy	15
3.4. Thermodynamic formalism of subadditive potentials	16
3.5. Thermodynamic formalism of linear cocycles	18
4. Multifractal Formalism of typical cocycles	21

Date: January 14, 2020.

2010 Mathematics Subject Classification. 26A18, 37D35, 37D30, 37H15, 93D30.

Key words and phrases. Lyapunov spectrum, Typical cocycles, Thermodynamic formalism, Random dynamics, Linear cocycles.

5.	Continuity of entropy spectrum of Lyapunov exponents	23
6.	Continuity of the lower joint spectral radius	26
7.	Relation between topological spectrum and Lyapunov spectrum for typical cocycles	28
8.	The continuity of topological pressure	29
8.1.	Application of Theorem 8.1	30
	References	31

1. INTRODUCTION AND STATEMENT OF THE RESULTS

Suppose that $T : X \rightarrow X$ is a continuous function on compact metric space X . Let $\mathcal{A} : X \rightarrow GL(d, \mathbb{R})$ be a measurable function. We define a *linear cocycle* $F : X \times \mathbb{R}^d \rightarrow X \times \mathbb{R}^d$ as

$$F(x, v) = (T(x), \mathcal{A}(x)v).$$

We say that F is generated by T and \mathcal{A} , we will also denote it by (T, \mathcal{A}) . Observe that $F^n(x, v) = (T^n(x), \mathcal{A}^n(x)v)$ for each $n \geq 1$, where

$$\mathcal{A}^n(x) := \mathcal{A}(T^{n-1}(x))\mathcal{A}(T^{n-2}(x))\dots\mathcal{A}(x).$$

If T is invertible then so is F . Moreover, $F^{-n}(x) = (T^{-n}(x), \mathcal{A}^{-n}(x)v)$ for each $n \geq 1$, where

$$\mathcal{A}^{-n}(x) := \mathcal{A}(T^{-n}(x))^{-1}\mathcal{A}(T^{-n+1}(x))^{-1}\dots\mathcal{A}(T^{-1}(x))^{-1}.$$

A simple class of linear cocycles are *locally constant cocycles* which are defined as follows. Assume that $X = \{1, \dots, k\}^{\mathbb{Z}}$ is a symbolic space. Suppose that $T : X \rightarrow X$ is a shift map, i.e. $T(x)_l = (x_{l+1})_l$. We say that $\mathcal{A} : X \rightarrow GL(d, \mathbb{R})$ is locally constant, if there exists $t \in \mathbb{N}$ such that $\mathcal{A}(x)$ depends only on the word $x_{-t}\dots x_t \in \mathcal{L}(2t+1)$ for every $x = (x_i)_{i \in \mathbb{Z}} \in X$. A locally constant cocycle (T, \mathcal{A}) is a cocycle whose generator \mathcal{A} is locally constant. We always assume that all locally constant functions are functions that depend only on the 0-th entry.

We define the *maximal Lyapunov exponent* as follows

$$\beta(\mathcal{A}) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \sup_{x \in X} \|\mathcal{A}^n(x)\|.$$

We introduce for any $x \in X$

$$\chi(x, \mathcal{A}) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \|\mathcal{A}^n(x)\|$$

whenever the limit exists, for definiteness, $\|A\|$ the Euclidean operator norm of a matrix A (i.e. the largest singular value of A). We call it *pointwise Lyapunov exponent* of \mathcal{A} at x .

Assume that μ is a T -invariant measure and $\log^+ \|\mathcal{A}^\pm\| \in L^1(\mu)$. By Kingman's subadditive theorem [V], the *Lyapunov exponent of measure μ*

$$\chi(\mu, \mathcal{A}) := \lim_{n \rightarrow \infty} \frac{1}{n} \int \log \|\mathcal{A}^n(x)\| d\mu(x)$$

exists. If μ is ergodic then $\chi(x, \mathcal{A}) = \chi(\mu, \mathcal{A})$ for a.e. points. The existence of pointwise Lyapunov exponent follows from Kingman's subadditive theorem, but Furstenberg and Kesten [FK1] show that the existence of pointwise Lyapunov exponent for a.e point at first.

It also follows from Kingman subdditive Ergodic Theorem that

$$\chi(\mu, \mathcal{A}^{-1}) := \lim_{n \rightarrow \infty} \frac{1}{n} \int \log \|(\mathcal{A}^n(x))^{-1}\|^{-1} d\mu(x)$$

exists.

In this paper, we investigate the *level sets* of Lyapunov exponents. That is,

$$E_{\mathcal{A}}(\alpha) := \{x \in X, \chi(x, \mathcal{A}) = \alpha\}.$$

We are going to study the continuity of entropy spectrum of Lyapunov exponents, that is, the topological entropy of level sets of points with a common given Lyapunov exponent.

We say that $\Phi := \{\log \phi_n\}_{n=1}^\infty$ is a *subadditive potential* if each ϕ_n is a continuous positive-valued function on X such that

$$0 < \phi_{n+m}(x) \leq \phi_n(x)\phi_m(T^n(x)) \quad \forall x \in X, m, n \in \mathbb{N}.$$

Furthermore, $\Phi = \{\log \phi_n\}_{n=1}^\infty$ is said to be an *almost additive potential* if there exists a constant $C \geq 0$ such that for any $m, n \in \mathbb{N}$, $x \in X$, we have

$$C^{-1}\phi_n(x)\phi_m(T^n(x)) \leq \phi_{n+m}(x) \leq C\phi_n(x)\phi_m(T^n(x)).$$

We denote by $P_\Phi(T)$ the topological pressure of a subadditive potential Φ . We denote by $P_\Phi(q)$ the topological pressure for a subadditive potential $q\Phi$.

The linear cocycles generated by a diffeomorphism map $T : X \rightarrow X$ on a closed Riemannian manifold X and a family of linear maps $\mathcal{A}(x) := D_x T : \mathcal{V}_x \rightarrow \mathcal{V}_{T(x)}$, where \mathcal{V} is a vector bundle on X^1 , are called *derivative cocycles*. Moreover, when $T : X \rightarrow X$ is an Anosov diffeomorphism (or expanding map), Bowen [B] showed that there exists a symbolic coding of T by a subshift of finite type. The smaller size of results for derivative cocycles show that the behaviors of derivative cocycles is significantly less tractable than that of locally constant cocycles. The principal aim of this article is to investigate properties of linear cocycles in a very general setting.

We denote by \mathcal{L} the set of admissible words Σ . We define for $\mathcal{A} : \Sigma \rightarrow GL(d, \mathbb{R})$ and $I \in \mathcal{L}$

$$\|\mathcal{A}(I)\| := \max_{x \in I} \|\mathcal{A}^{|I|}(x)\|.$$

¹It is not necessary trivial bundle.

We define a positive continuous function $\{\varphi_{\mathcal{A},n}\}_{n \in \mathbb{N}}$ on Σ such that

$$\varphi_{\mathcal{A},n}(x) = \|\mathcal{A}^n(x)\|.$$

We denote by $\Phi_{\mathcal{A}}$ the subadditive potential $\{\log \varphi_{\mathcal{A},n}\}_{n=1}^{\infty}$.

We say that $\varphi_{\mathcal{A},\cdot}(x)$ is *quasi-multiplicativite* if there exist $C > 0$ and $m \in \mathbb{N}$ such that for every $I, J \in \mathcal{L}$, there exists $K \in \mathcal{L}$ with $|K| \leq m$ such that

$$\varphi_{\mathcal{A},|IKJ|}(x) \geq C\varphi_{\mathcal{A},|I|}(y)\varphi_{\mathcal{A},|J|}(z),$$

for any $x \in [IKL], y \in [I]$ and $z \in [J]$.

In this paper, we always assume that $T : \Sigma \rightarrow \Sigma$ is topologically mixing subshift of finite type. We denote by $H^r(\Sigma, GL(d, \mathbb{R}))$ the space of all r -Hölder continuous functions. We also denote by $H_b^r(\Sigma, GL(d, \mathbb{R}))$ the space of all r -Hölder continuous and fiber bunched functions, which says that the cocycles are nearly conformal. We define the *typical cocycles* among $H^r(\Sigma, GL(d, \mathbb{R}))$ (see section 2). That is

$$\mathcal{W} := \{\mathcal{A} \in H_b^r(\Sigma, GL(d, \mathbb{R})) : \mathcal{A} \text{ is pinching and twisting}\}.$$

Our main results are Theorems 1.1, 1.2, 1.3, 1.4, and 1.5 formulated as follows:

Theorem 1.1. *Let $\mathcal{A} \in \mathcal{W}$. If $\alpha_t = P'_{\Phi_{\mathcal{A}}}(t)$ for $t > 0$. Then,*

$$h_{top}(E(\alpha_t)) \rightarrow h_{top}(E(\beta(\mathcal{A}))) \text{ when } t \rightarrow \infty.$$

Park [P] shows that every $\mathcal{A} \in \mathcal{W}$ is quasi-multiplicativite. That implies $P_{\Phi_{\mathcal{A}}}(q)$ is convex on $q \in \mathbb{R}$. We use this fact to prove the *restricted variational principle* to level sets as follows.

Theorem 1.2. *Assume that $T : X \rightarrow X$ is topologically mixing subshift of finite type on compact metric space X . Suppose that $\mathcal{A} : X \rightarrow GL(d, \mathbb{R})$ belongs to typical functions \mathcal{W} . Then,*

$$\sup\{h_{\mu}(T) : \mu \in \mathcal{M}(X, T), \chi(\mu, \mathcal{A}) = \alpha\} = \inf\{P_{\Phi_{\mathcal{A}}}(q) - \alpha \cdot q : q \in \mathbb{R}\}.$$

We say that $\mathcal{A} : \Sigma \rightarrow GL(d, \mathbb{R})$ is *almost additive* if there exists a constant κ such that $\|\mathcal{A}(I)\mathcal{A}(J)\| \geq \kappa\|\mathcal{A}(I)\|\|\mathcal{A}(J)\|$ for all $I, J \in \mathcal{L}(n)$, where $\mathcal{L}(n)$ is set of admissible words with length n . Moreover, it is easy to see that $\|\mathcal{A}(I)\mathcal{A}(J)\| \leq \|\mathcal{A}(I)\|\|\mathcal{A}(J)\|$ for all $I, J \in \mathcal{L}(n)$. Bochi and Morris [BM] show that \mathcal{A} is almost additive if \mathcal{A} is 1-domination, which says that the ratio between first and second-singular values of the matrices of the n -th iterate increase uniformly exponentially when n goes to ∞ . So, the condition almost additive for $\mathcal{A} : \Sigma \rightarrow GL(d, \mathbb{R})$ is equivalent that $\{\log \|\mathcal{A}(I)\|\}$ is an almost additive potential for all $I \in \mathcal{L}(n)$. One needs to be careful that quasi-multiplicativity is not equivalence of almost additivity. For instant, consider $T : \{0, 1\}^{\mathbb{Z}} \rightarrow \{0, 1\}^{\mathbb{Z}}$ a shift map. We define a linear cocycle (T, A) such that

$$A_0 = \begin{bmatrix} 2 & 0 \\ 0 & \frac{1}{2} \end{bmatrix}, \quad A_1 = R_{\theta},$$

where θ is irrational angle.

One can define the *lower joint spectral radius* of $\mathcal{A} : \Sigma \rightarrow GL(d, \mathbb{R})$ as follows

$$\check{\rho}(\mathcal{A}) := \liminf_{n \rightarrow \infty} \{ \|\mathcal{A}(I)\|^{\frac{1}{n}} : I \in \mathcal{L}(n) \}.$$

In locally constant cocycles case, Bochi and Morris [BM] investigated the continuity properties of the lower joint spectral radius. They show that $\check{\rho}(\mathcal{A})$ is Lipschitz continuous at \mathcal{A} under 1–domination assumption. Breuillard and Sert [BS] extended Bochi and Morris’s result to the joint spectrum under domination condition. In this case the $\chi(\mu, \mathcal{A})$ depends continuously on the measure μ .

For $\vec{\alpha} \in \mathbb{R}^d$, the *Lyapunov spectrum* of the cocycle is defined as:

$$\vec{L} := \{ \vec{\alpha}, \exists x \in \Sigma \text{ such that } \lim_{n \rightarrow \infty} \frac{1}{n} \log \phi^k(\mathcal{A}^n(x)) = \alpha_k \},$$

where $\phi^k(\mathcal{A}) = \|\mathcal{A}^{\wedge k}\|$ for $1 \leq k \leq d$.

We have

$$\log \check{\rho}(\mathcal{A}) = \min \{ \alpha_1, \vec{\alpha} = (\alpha_1, \alpha_2, \dots, \alpha_d) \in \vec{L} \} = \alpha(\mathcal{A}).$$

In general setting, we prove that the continuity of the minimal Lyapunov exponent under 1–domination assumption. Obviously, our result implies continuity of the lower joint spectral radius.

Theorem 1.3. *Assume that \mathcal{A}_k and \mathcal{A} satisfy 1–domination. Then, $\alpha(\mathcal{A}_k) \rightarrow \alpha(\mathcal{A})$, when $\mathcal{A}_k \rightarrow \mathcal{A}$. Consequently, $\check{\rho}(\mathcal{A}_k) \rightarrow \check{\rho}(\mathcal{A})$ when $\mathcal{A}_k \rightarrow \mathcal{A}$.*

We also show that the Lyapunov spectrum coincides with the closure of the set of positive topological spectrum $h_{top}(E_{\mathcal{A}}(\cdot))$ of Lyapunov exponents for typical cocycles \mathcal{W} .

Theorem 1.4. *Let $\mathcal{A} \in \mathcal{W}$. Then,*

$$L = \overline{\{ \alpha, h_{top}(E_{\mathcal{A}}(\alpha)) > 0 \}}.$$

Furthermore, $\alpha \mapsto h_{top}(E_{\mathcal{A}}(\alpha))$ is concave for $\alpha \in \overset{\circ}{L}$.

In this paper, we also obtain the high dimensional versions of Theorems 1.1, 1.2, 1.3, and 1.4 (see Theorems 4.2, 5.4, 6.4 and 7.4).

Finally, we will investigate on continuity of topological pressure for linear cocycles.

We define the singular value function $\varphi^s : GL(d, \mathbb{R}) \rightarrow [0, \infty)$ with the parameter $s \geq 0$ as follows.

$$\varphi^s(\mathcal{A}) = \sigma_1(\mathcal{A}) \dots \sigma_m(\mathcal{A}) \sigma_{m+1}^{s-m},$$

where $m = [s]^2$. We make the convention $O^0 = 1$. For completeness, if $s > d$, the we also define

$$\varphi^s(\mathcal{A}) = (\det(\mathcal{A}))^{\frac{s}{d}}.$$

² $[s] = s - 1$.

It is well known that φ^s is submultiplicative for all $s \geq 0$. That means, for any $A, B \in GL(d, \mathbb{R})$

$$\varphi^s(AB) \leq \varphi^s(A)\varphi^s(B).$$

The function $(s, A) \mapsto \varphi^s(A)$ is continuous in both A and s where $A \in GL(d, \mathbb{R})$. We define $\tilde{\varphi}_{\mathcal{A}}^s$ on \mathcal{L} as follows, for any $n \in \mathbb{N}$ and $I \in \mathcal{L}(n)$,

$$\tilde{\varphi}_{\mathcal{A}}^s(I) := \max_{x \in I} \varphi_{\mathcal{A}, n}^s(x).$$

$\tilde{\varphi}_{\mathcal{A}}^s$ is also submultiplicative.

Quasi-multiplicativity of $\varphi_{\mathcal{A}, n}$ implies quasi-multiplicativity $\tilde{\varphi}_{\mathcal{A}}$. That is, if there if there exist $C > 0$ and $m \in \mathbb{N}$ such that for every $I, J \in L$, there exists $K \in L$ with $|K| \leq m$ such that $IKJ \in L$ and

$$\tilde{\varphi}_{\mathcal{A}}(IKJ) \geq C\tilde{\varphi}_{\mathcal{A}}(I)\tilde{\varphi}_{\mathcal{A}}(J).$$

We denote by $\tilde{\Phi}_{\mathcal{A}} = \{\log \tilde{\varphi}_{\mathcal{A}}\}$.

Feng and Shmerkin [FS] show that continuity of topological pressure for locally constant cocycles. In fact, the proof of Theorem 1.5 is based on ideas from those work. Moreover, this kind of result is known for typical cocycles by Park.

We show that one can prove the continuity of topological pressure for $H_b^r(\Sigma, GL(2, \mathbb{R}))$ without assumption pinching and twisting.

Theorem 1.5. *The map $(s, \mathcal{A}) \mapsto P_{\tilde{\Phi}_{\mathcal{A}}}(s)$ is continuous on $H_b^r(\Sigma, GL(2, \mathbb{R}))$.*

This paper is organized as follows. In Section 2, we recall some preliminary material regarding symbolic dynamics as well as some results in typical cocycles. In Section 3, we discuss relevant results in thermodynamics formalism of subadditive potentials. In section 4, we discuss multifractal formalism as well as proof of Theorem 1.2. In section 5, we prove Theorem 1.1 and the proof of Theorem 1.3 is given in section 6. In section 7, we prove Theorem 1.4. Finally, we prove Theorem 1.5 in section 8, and also we give an application of the theorem.

Acknowledgements. The author thanks his supervisor, Michał Rams, for his careful reading of an earlier version of this paper and many helpful discussions. He would also like to thank Aaron Brown and Cagri Sert for introducing papers [P] and [BS]. The author was partially supported by the National Science Center grant 2014/13/B/ST1/01033 (Poland).

2. PRELIMINARIES

In this section, we recall some basic facts and definition that we need to prove for main theorems.

2.1. Symbolic dynamics. In this subsection, we recall some definitions and basic facts related to subshift of finite type. For more information see [LM].

Definition 2.1. Let $Q = (q_{ij})$ be a $k \times k$ with $q_{ij} \in \{0, 1\}$. The *one side subshift of finite type* associated to the matrix Q is a left shift map $T : \Sigma_Q^+ \rightarrow \Sigma_Q^+$, where Σ_Q^+ is a set of sequences

$$\Sigma_Q^+ := \{x = (x_i)_{i \in \mathbb{N}_0} : x_i \in \{1, \dots, k\} \text{ and } Q_{x_i, x_{i+1}} = 1 \text{ for all } n \in \mathbb{N}_0\},$$

Similarly, one can define *two sided subshift of finite type* $T : \Sigma_Q \rightarrow \Sigma_Q$, where

$$\Sigma_Q := \{x = (x_i)_{i \in \mathbb{Z}} : x_i \in \{1, \dots, k\} \text{ and } Q_{x_i, x_{i+1}} = 1 \text{ for all } n \in \mathbb{Z}\}.$$

For simplicity, we assume that $\Sigma_Q^+ = \Sigma^+$ and $\Sigma_Q = \Sigma$.

Definition 2.2. The Q is called *primitive*, for every $i, j \in \{1, \dots, k\}$ there is $n_{i,j} > 0$ such that $(Q^{n_{i,j}})_{ij} > 0$.

It is well known that every subshift finite type associated to a primitive matrix Q is equivalent property of *topologically mixing* T . That is, for every $U, V \subset \Sigma$, there is N such that for every $n \geq N$, $T^n(U) \cap V \neq \emptyset$.

We fix $\omega \in (0, 1)$ and consider the space Σ is endowed with the metric d which is defined as follows: For $x = (x_i)_{i \in \mathbb{Z}}, y = (y_i)_{i \in \mathbb{Z}} \in \Sigma$, we have

$$d(x, y) = \omega^k \text{ for all } |i| < k,$$

where k is the largest integer such that $x_i = y_i$.

The subshift of finite type $T : \Sigma \rightarrow \Sigma$ becomes a hyperbolic homeomorphism, if Σ is equipped by the metric d .

We denote by \mathcal{L} the set of collection of admissible words. We denote by $|I|$ the length of $I \in \mathcal{L}$. Denote by $\mathcal{L}(n)$ the *admissible word of length n* . That is, a word i_0, \dots, i_{n-1} with $i_j \in \{1, \dots, q\}$ such that $Q_{x_i, x_{i+1}} = 1$. One can define *the cylinder $[I]$* as follows:

$$[I] = [i_0 \dots i_{n-1}] := \{x \in \Sigma : x_i = i_j \ \forall 0 \leq j \leq n-1\},$$

for any $I = i_0 \dots i_{n-1} \in \mathcal{L}(n)$.

We define *local stable set*

$$W_{loc}^s(x) = \{(y_n)_{n \in \mathbb{Z}} : x_n = y_n \text{ for all } n \geq 1\}$$

and the *local unstable set*

$$W_{loc}^u(x) = \{(y_n)_{n \in \mathbb{Z}} : x_n = y_n \text{ for all } n < 0\}.$$

Furthermore, the global stable and unstable manifolds of x ,

$$W^s(x) := \cup_{n=0}^{\infty} T^{-n}(W_{loc}^s(T^n(x))) \text{ and } W^u(x) := \cup_{n=0}^{\infty} T^n(W_{loc}^u(T^{-n}(x)))$$

are smoothly immersed submanifolds of X and they are characterized by

$$W^s(x) = \{y \in X : \lim_{n \rightarrow \infty} d(T^n(x), T^n(y)) = 0\},$$

$$W^u(x) = \{y \in X : \lim_{n \rightarrow -\infty} d(T^n(x), T^n(y)) = 0\}.$$

2.2. Multilinear algebra. We recall some basic facts about the exterior algebra. We use it for studying the singular value function.

We denote by $\sigma_1, \dots, \sigma_d$ the singular values of the matrix \mathcal{A} , which are the square roots of the eigenvalues of the positive semi definite matrix $\mathcal{A}^* \mathcal{A}$ listed in decreasing order according to multiplicity.

Assume that $\{e_1, \dots, e_d\}$ is the standard orthogonal basis of \mathbb{R}^d and define

$$\wedge^k \mathbb{R}^d := \{e_{i_1} \wedge e_{i_2} \wedge \dots \wedge e_{i_k} : 1 \leq i_1 \leq i_2 \leq \dots \leq i_k \leq d\}$$

for all $k \in \{1, \dots, d\}$ with the convention that $\wedge^0 \mathbb{R}^d = \mathbb{R}$. It is called the k -th exterior power of \mathbb{R}^d .

We are interested in the invertible matrices $GL(d, \mathbb{R})$. We consider induced topology \mathbb{R}^{d^2} for it. For $A \in GL(d, \mathbb{R})$, we define an invertible linear map $A^{\wedge k} : \wedge^k \mathbb{R}^d \rightarrow \wedge^k \mathbb{R}^d$ as follows

$$(A^{\wedge k}(e_{i_1} \wedge e_{i_2} \wedge \dots \wedge e_{i_k})) = Ae_{i_1} \wedge Ae_{i_2} \wedge \dots \wedge Ae_{i_k}.$$

$\wedge^k \mathbb{R}^d$ is represented by a $\binom{n}{k} \times \binom{n}{k}$ whose entries are the $k \times k$ minors of A . It can also show that

$$(AB)^{\wedge k} = A^{\wedge k} B^{\wedge k}.$$

The singular values of $A^{\wedge k}$ are the product $\sigma_{i_1}(A) \dots \sigma_{i_k}(A)$. In addition,

$$\|A^{\wedge k}\| = \sigma_1(A) \dots \sigma_k(A).$$

According to above observation, where A is k -dominated if and only if $A^{\wedge k}$ is 1-dominated.

2.3. Fiber bunched cocycles. We remind that we always assume that $T : \Sigma \rightarrow \Sigma$ is topologically mixing subshift of finite type and $\mathcal{A} : \Sigma \rightarrow GL(d, \mathbb{R})$ is a Hölder continuous function.

We say that $\mathcal{A} : X \rightarrow GL(d, \mathbb{R})$ is a r -Hölder continuous function, if there exists $C > 0$ such that

$$(2.1) \quad \|\mathcal{A}(x) - \mathcal{A}(y)\| \leq Cd(x, y)^r \quad \forall x, y \in \Sigma.$$

We denote by $H^r(\Sigma, GL(d, \mathbb{R}))$ the set of r -Hölder continuous functions. We also show that by $H^r(\Sigma)$, when there is no confusion about $GL(d, \mathbb{R})$.

We denote by $h_r(\mathcal{A})$ the smallest constant C in (2.1). We equip the $H^r(\Sigma, GL(d, \mathbb{R}))$ with the distance

$$D_r(A, B) = \sup_X \|A - B\| + h_r(A - B).$$

It is clear the locally constant functions are ∞ -Hölder i.e, they are r -Hölder for every $r > 0$, with bounded $h_r(A)$.

Definition 2.3. A local stable holonomy for the linear cocycles (\mathcal{A}, T) is a family of matrices $H_{y \leftarrow x}^s \in GL(d, \mathbb{R})$ defined for all $x \in \Sigma$ with $y \in W_{loc}^s(x)$ such that

$$a) \quad H_{x \leftarrow x}^s = Id \text{ and } H_{z \leftarrow y}^s \circ H_{y \leftarrow x}^s = H_{z \leftarrow x}^s \text{ for any } z, y \in W_{loc}^s(x).$$

- b) $\mathcal{A}(x) \circ H_{x \leftarrow y}^s = H_{T(x) \leftarrow T(y)}^s \circ \mathcal{A}(y)$.
 c) $(x, y, v) \mapsto H_{y \leftarrow x}(v)$ is continuous.

Moreover, if $y \in W_{loc}^u(x)$, then there are analogous properties for $H_{x \leftarrow y}^u$.

According (b) in the above definition, one can extend the definition to the global stable holonomy $H_{y \leftarrow x}^s$ for $y \in W^s(x)$ not necessarily in $W_{loc}^s(x)$:

$$H_{y \leftarrow x}^s = \mathcal{A}^n(y)^{-1} \circ H_{T^n(y) \leftarrow T^n(x)}^s \circ \mathcal{A}^n(x),$$

where $n \in \mathbb{N}$ is large enough such that $T^n(y) \in W_{loc}^s(T^n(x))$. One can extend the definition the global unstable holonomy similarly.

Definition 2.4. A r -Hölder continuous function \mathcal{A} is called *fiber bunched* if for any $x \in \Sigma$,

$$(2.2) \quad \|\mathcal{A}(x)\| \|\mathcal{A}(x)^{-1}\| \omega^r < 1,$$

where ω is the hyperbolicity constant defining the metric on the base Σ .

We say that the linear cocycle (\mathcal{A}, T) is fiber-bunched if its generator \mathcal{A} is fiber-bunched. We denoted by $H_b^r(\Sigma, GL(d, \mathbb{R}))$ the family of r -Hölder-continuous and fiber bunched functions.

The geometric interpretation of the fiber bunching condition is as follows. Let $\mathcal{A} \in H_b^r(\Sigma, GL(d, \mathbb{R}))$. The *projection* cocycle associated to \mathcal{A} and T is the map $\mathbb{P}F : \Sigma \times \mathbb{P}\mathbb{R}^d \rightarrow \Sigma \times \mathbb{P}\mathbb{R}^d$ given by

$$\mathbb{P}F(x, v) := (T(x), \frac{\mathcal{A}(x)v}{\|\mathcal{A}(x)v\|}).$$

We denote by $D\mathcal{A}_v$ the derivative of the action $\mathbb{P}\mathbb{R}^d \rightarrow \mathbb{P}\mathbb{R}^d$ on projective space at all points $v \in \mathbb{P}\mathbb{R}^d$. Taking derivative

$$\|D\mathcal{A}_v\| \leq \|\mathcal{A}\| \|\mathcal{A}^{-1}\| \quad \text{and} \quad \|D\mathcal{A}_v^{-1}\| \leq \|\mathcal{A}\| \|\mathcal{A}^{-1}\|$$

for all $v \in \mathbb{P}\mathbb{R}^d$. So, the fiber bunching condition implies that rate of expansion (respectively, contraction) the projective cocycle $\mathbb{P}F$ at every point $x \in \Sigma$ is bounded above by $(\frac{1}{\omega})^r$ (respectively, below by ω^r).

The Hölder continuity and the fiber bunched assumption $\mathcal{A} \in H_b^r(\Sigma, GL(d, \mathbb{R}))$ implies that the convergence of the *canonical holonomy* $H^{s/u}$ (see [BGMV], [KS]). That means, for any $y \in W_{loc}^{s/u}$,

$$H_{y \leftarrow x}^s := \lim_{n \rightarrow \infty} \mathcal{A}^n(y)^{-1} \mathcal{A}^n(x) \quad \text{and} \quad H_{y \leftarrow x}^u := \lim_{n \rightarrow -\infty} \mathcal{A}^n(y)^{-1} \mathcal{A}^n(x).$$

In addition, when the linear cocycle is fiber bunched, the canonical holonomies vary r -Hölder continuously (see [KS]), i.e., there exists $C > 0$ such that for $y \in W_{loc}^{s/u}(x)$,

$$\|H_{x \leftarrow y}^{s/u} - I\| \leq Cd(x, y)^r.$$

In this paper, we will always work with the canonical holonomies for fiber bunched cocycles.

Remark 1. *Even though the locally constant cocycles are not necessary fiber bunched, the canonical holonomies always exist. Indeed, for every $y \in W^s(x)$ there exist m such that $x_n = y_n$ for all $n \geq m$. Then,*

$$H_{x \leftarrow y}^s = \mathcal{A}^{-1}(x) \cdots \mathcal{A}^{m-1}(x)^{-1} \mathcal{A}^{m-1}(y) \cdots \mathcal{A}(y).$$

In particular $H_{x \leftarrow y}^s = Id$, for all $x \in W_{loc}^s(y)$. Similarly, we get the existence of the unstable holonomy.

Remark 2. *If a linear cocycle is not fiber bunched, then it might admit multiple holonomies (see [KS1]).*

2.4. Typical cocycles. We are going to discuss typical cocycles. For details, one is referred to [AV], [BV1] and [V].

Suppose that $p \in \Sigma$ is a periodic point of T , we say the $p \neq z \in \Sigma$ is a *homoclinic point* associated to p if it is the intersection of the stable and unstable manifold of p . That is, $z \in W^s(p) \cap W^u(p)$ (see figure 1). The set of homoclinic points of any periodic point is dense in Σ for hyperbolic systems.




FIGURE 1. Homoclinic point.


We define the *holonomy loop*

$$\psi_p^z := H_{z \leftarrow p}^s \circ H_{p \leftarrow z}^u.$$

Definition 2.5. Suppose that $\mathcal{A} : \Sigma \rightarrow GL(d, \mathbb{R})$ belongs $H_b^r(\Sigma, GL(d, \mathbb{R}))$. We say that \mathcal{A} is *1-typical* if there exists a periodic point p and a homoclinic point z associated to p such that:

- (i) The eigenvalues of $\mathcal{A}^{per(p)}(p)$ have multiplicity 1 and distinct norms. Let $\{v_i\}_{i=1}^d$ be the eigenvectors of $P := \mathcal{A}^{per(p)}(p)$.
- (ii) $\psi_p^z(v_i) \cap W_j = \emptyset$, where W_j spanned by all eigenvectors of P other than v_i for any $1 \leq i, j \leq d$.

For $d = 2$ this second condition means that $\psi_p^z(v_i)$ does not intersect other lines. See Figure 2 for a 1-typical cocycle in dimension 2.


 FIGURE 2. $\psi_p^z(v_1) \neq v_2$

We refer to (i) as the (*pinching*) properties and to (ii) as the (*twisting*) properties.

The cocycles generated by \mathcal{A}^t , $1 \leq t \leq d$ also admit stable and unstable holonomies, namely $(H^{s/u})^t$.

Definition 2.6. Assume that \mathcal{A} is 1-typical. We say \mathcal{A} is *t-typical* for $2 \leq t \leq d-1$, if the points $p, z \in \Sigma$ from Definition 2.5 satisfy

- (I) P^t satisfies the analogous statement to (i) from Definition (2.5) for all t .
Let $\{v_{i_1} \wedge \dots \wedge v_{i_t}\}_{1 \leq i_1 < \dots < i_t \leq d}$ be the eigenvectors of P^t .
- (II) The induced map $(\psi_p^z)^t$ on $(\mathbb{R}^d)^t$ satisfies the analogous statement to (ii) from Definition 2.5.

We say that A is *typical* if A is t -typical for all $1 \leq t \leq d-1$. We denote by $\mathcal{W} \subset H_b^r(\Sigma, GL(d, \mathbb{R}))$ the set of all typical functions. \mathcal{W} is open in $H_b^r(\Sigma)$, and Bonatti, and Viana [BV1] show that \mathcal{W} is dense in $H_b^r(\Sigma, GL(d, \mathbb{R}))$, and its complement has infinite codimension.

Remark 3. Above definition for typical cocycles comes from Park's paper [P] that is a slightly weaker form typical cocycles which was introduced by Bonatti and Viana's paper [BV1].

2.5. The continuity of Lyapunov exponents. Throught, $\mathbb{P}F : \Sigma^+ \times \mathbb{P}\mathbb{R}^d \rightarrow \Sigma^+ \times \mathbb{P}\mathbb{R}^d$ is the projective cocycle associated with linear cocycle $F : \Sigma^+ \times \mathbb{R}^d \rightarrow \Sigma^+ \times \mathbb{R}^d$, which is generated by (\mathcal{A}, T) .

Furstenberg [V, Theorem 6.8] showed that the Lyapunov exponent $\chi(\mu, \mathcal{A})$ of F coincide with the integral of the function $\psi : \Sigma^+ \times \mathbb{P}\mathbb{R}^d \rightarrow \mathbb{R}$,

$$\psi(x, v) = \log \frac{\|A(x)v\|}{\|v\|}$$

for locally constant cocycles under *strongly irreducible* assumption, which there is no finite family of proper subspaces invariant by $\mathcal{A}(x)$ for μ -almost every x . In other words, he showed that

$$\chi(\mu, \mathcal{A}) = \int \psi d(\mu \times \eta),$$

for any stationary measure η of the associated projective cocycle $\mathbb{P}F$. So, one can easily show that we have the continuity of Lyapunov exponents with respect to (\mathcal{A}, μ) ([V, Corollary 6.10]) under strongly irreducibility assumption.

Remark 4. *Bonatti and Viana [BV1], [BGMV] extended Furstenberg's formula to typical cocycles. So, we have the continuity of Lyapunov exponents with respect to (\mathcal{A}, μ) , as well.*

Even though discontinuity of Lyapunov exponents are a common features (see [Bo], [Boc1]), there are some results for the continuity of Lyapunov exponents. For instance, Bocker and Viana [BV] proved continuity of Lyapunov exponents of 2-dimensional locally constant cocycles under a certain assumption. Avila, Eskin and Viana [AEV] announced recently that Bocker and Viana's result remains true in arbitrary dimension.

It was conjectured by Viana [V] that Lyapunov exponents are always continuous among $H_b^\alpha(X, GL(2, \mathbb{R}))$ -cocycles, and that has been proved by Backes, Brown and Butler [BBB]. In fact, they prove Lyapunov exponents vary continuously on any family of $GL(2, \mathbb{R})$ -cocycles with continuous invariant holonomies i.e.,

$$\chi(x, \mathcal{A}_k) \rightarrow \chi(x, \mathcal{A}),$$

when $(\mathcal{A}_k, H^{s,k}, H^{u,k}) \rightarrow (\mathcal{A}, H^s, H^u)$.

We state the main result of Backes, Brown, and Butler as follows.

Theorem 2.7. [BBB, Theorem 2.8] *Suppose that $\{\mathcal{A}_n\}_{n \in \mathbb{N}}$ is a sequence of 2-dimensional linear cocycles over T converging uniformly with holonomies to a cocycle \mathcal{A} and $\{\mu_n\}_{n \in \mathbb{N}}$ a sequence of fully supported, ergodic, T -invariant probability measures converging to an ergodic, T -invariant measure μ with local product structure and full support. Then*

$$\chi(\mu_n, \mathcal{A}_n) \rightarrow \chi(\mu, \mathcal{A}),$$

and,

$$\chi(\mu_n, \mathcal{A}_n^{-1}) \rightarrow \chi(\mu, \mathcal{A}^{-1}).$$

That improves Bocker and Viana's result [BV]. Furthermore, the Butler [Bu] showed in the following example that the fiber-bunching condition is sharp.

Example 2.8. Assume that $T : \{0, 1\}^{\mathbb{Z}} \rightarrow \{0, 1\}^{\mathbb{Z}}$ is a shift map. We define a locally constant cocycle (T, A) such that

$$A_0 = \begin{bmatrix} \sigma & 0 \\ 0 & \sigma^{-1} \end{bmatrix}, \quad A_1 = \begin{bmatrix} \sigma^{-1} & 0 \\ 0 & \sigma \end{bmatrix},$$

where σ is a positive constant greater than 1. We define probability measure ν_p in order to $\nu_p([0]) = p, \nu_p([1]) = 1 - p$, and then Bernoulli measure $\mu_p = \nu_p^{\mathbb{Z}}$. By definition the cocycle (T, A) is fiber bunched if and only if $\sigma^2 < 2^\alpha$ ³.

Butler[Bu] shows that for above example if $\sigma^{4p-2} \geq 2^\alpha$ for $p \in (\frac{1}{2}, 1)$, then for each neighborhood $\mathcal{U} \subset H^\alpha(\{0, 1\}^{\mathbb{Z}}, SL(2, \mathbb{R}))$ of A and every $\kappa \in (0, (2p-1) \log \sigma)$, there is a locally constant cocycle $\mathcal{B} \in \mathcal{U}$ such that $\chi(x, \mathcal{B}) = \kappa$. In particular, A is a discontinuity point for Lyapunov exponents in $H^\alpha(\{0, 1\}^{\mathbb{Z}}, SL(2, \mathbb{R}))$. So, this example shows that we have discontinuity of Lyapunov exponents near fiber bunched cocycles.

The inequality $\sigma^{4p-2} \geq 2^\alpha$ comes from the following observation

$$\lim_{n \rightarrow \infty} \log(\|A^n(x)\| \|A^n(x)^{-1}\|) = (\chi(\mu_p, A) - \chi(\mu_p, A^{-1})) = (4p - 2) \log \sigma,$$

for μ_p almost every $x \in \{0, 1\}^{\mathbb{Z}}$.

Lyapunov exponents are T -invariant maps, thus when μ is ergodic they are constant μ -almost everywhere. In that case, we denote them as $\chi_i(\mathcal{A})$ for $i = 1, \dots, d$.

Theorem 2.9. [P, Theorem D] *Let $\mathcal{A} \in \mathcal{W}$. Then \vec{L} is a convex and closed subset of \mathbb{R}^2 .*

We can use Theorem 2.7 to show that the Lyapunov spectrum on fiber bunched cocycles is a closed and convex set.

Corollary 2.10. *Let $\mathcal{A} \in H_b^r(X, GL(2, \mathbb{R}))$. Then \vec{L} is a convex and closed subset of \mathbb{R}^2 .*

Proof. Since \mathcal{W} is open and dense, for every $\mathcal{A} \in H_b^r(X, GL(2, \mathbb{R}))$ there is a $\mathcal{A}_k \in \mathcal{W}$ such that $\mathcal{A}_k \rightarrow \mathcal{A}$.

By Theorem 2.7,

$$\chi_i(\mathcal{A}_k) \rightarrow \chi_i(\mathcal{A})$$

for $i = 1, 2$. By Theorem 2.9, \vec{L} is a closed and convex subset of \mathbb{R}^2 . \square

³ Σ is equipped by a norm d that is, for all $x \neq y$, $d(x, y) = 2^{-N(x, y)}$, where $N(x, y) = \min\{n, x_n \neq y_n\}$.

3. THERMODYNAMIC FORMALISM

3.1. Convex functions. Let U be an open convex subset of \mathbb{R}^n and f be a real continuous convex function on U . We say a vector $a \in \mathbb{R}^n$ is a *subgradient* of f at x if for all $z \in U$,

$$f(z) \geq f(x) + a^T(z - x),$$

where the right hand side is the scalar product.

For each $x \in \mathbb{R}^n$ set the *subdifferential* of f at point x

$$\partial f(x) := \{a : a \text{ is a subgradient for } f \text{ at } x\}.$$

For $x \in U$, the subdifferential $\partial f(x)$ is always a nonempty convex compact set. Define $\partial^e f(x) := \text{ext}\{\partial f(x)\}$. We say that f is *differentiable* at x when $\partial^e f(x) = \{a\}$.

We define

$$(3.1) \quad \partial f(U) = \cup_{x \in U} \partial f(x) \quad \text{and} \quad \partial^e f(U) = \cup_{x \in U} \partial^e f(x).$$

3.2. Legendre transform. Assume that $f : \mathbb{R}^d \rightarrow \mathbb{R} \cup \{+\infty\}$ is a convex function that is not identically equal to $-\infty$. The Legendre transform of f is the function f^* of a new variable t , defined by

$$t \mapsto f^*(t) := \sup\{t \cdot x - f(x) : x \in \mathbb{R}^d\},$$

where right hand side is scalar product.

It is easy to show that f^* is a convex function and not identically equal to $-\infty$. Let f^{**} be the Legendre transform of f^* . The following result is well known (cf. [R, Theorem 12.2]).

Theorem 3.1. *Assume that $f : \mathbb{R}^k \rightarrow \mathbb{R} \cup \{\infty\}$ is convex and not identically equal to $-\infty$. Let $x \in \mathbb{R}^k$. Suppose that f is lower semi continuous at x , i.e., $\liminf_{y \rightarrow x} f(y) \geq f(x)$. Then $f^{**}(x) = f(x)$.*

Feng and Huang [FH, Corollary 2.5] prove the following corollary as an application the above theorem.

Corollary 3.2. *Assume that A is a non-empty, closed and convex set in \mathbb{R}^k and $g : A \rightarrow \mathbb{R}$ be a concave function. Set*

$$W(x) = \sup\{g(a) + ax : a \in A\}, \quad x \in \mathbb{R}^k$$

and

$$G(a) = \inf\{W(x) - ax : x \in \mathbb{R}^k\}, \quad a \in A.$$

Finally, if g is upper semi continuous at $a \in A$, then $G(a) = g(a)$.

3.3. Topological entropy. Let (X, τ, μ) be a Borel probability space, and $T : X \rightarrow X$ be a measure preserving transformation.

A *partition* of (X, τ, μ) is a subfamily of τ consisting of mutually disjoint elements whose union is X . We denote by α and β the countable partition of X .

Let $\alpha = \{A_i, i \geq 1\}$, where $A_i \in \tau$. We define

$$H_\mu(\alpha) = - \sum_{A \in \alpha} \mu(A) \log \mu(A)$$

to be the *entropy* of α (with the convention $0 \log 0 = 0$).

We denote by $\alpha \vee \beta$ the joint partition $\{A \cap B \mid A \in \alpha, B \in \beta\}$.

Let $T^{-1}(\alpha) = \{T^{-1}(A) \mid A \in \alpha\}$. We define

$$h(\mu, \alpha) = \lim_{n \rightarrow \infty} \frac{1}{n} H_\mu\left(\bigvee_{j=0}^{n-1} T^{-j}(\alpha)\right)$$

to be the entropy of T relative to α ⁴.

Then the metric entropy of μ is defined as

$$h_\mu(T) = \sup h(\mu, \alpha),$$

where the supremum is taken over all countable partitions α with $H_\mu(\alpha) < \infty$.

Assume that (X, d) is a compact metric space and $T : X \rightarrow X$ is a continuous transformation. For any $n \in \mathbb{N}$ we define a new metric d_n on X as follows

$$d_n(x, y) = \max\{d(T^k(x), T^k(y)) : k = 0, \dots, n\},$$

and for any $\varepsilon > 0$, one can define *Bowen ball* $B_n(x, \varepsilon)$ that is an open ball of radius $\varepsilon > 0$ in the metric d_n around x . That is,

$$B_n(x, \varepsilon) = \{y \in X : d_n(x, y) < \varepsilon\}.$$

Given a set $Y \subset X$. If $Y \subset \cup_i B_{n_i}(x_i, \varepsilon)$, then one says that most countable collection of balls set $\mathcal{Y} = \{B_{n_i}(x_i, \varepsilon)\}_i$ covers Y . Consider $N(\mathcal{Y}) = \min_i n_i$. Let $s \geq 0$ and

$$S(Y, s, N, \varepsilon) = \inf \sum_i e^{-sn_i},$$

where the infimum is taken over all collections $\mathcal{Y} = \{B_{n_i}(x_i, \varepsilon)\}_i$ covering Y such that $n(\mathcal{Y}) \geq N$. The quantity $S(Y, s, N, \varepsilon)$ does not decrease with N , consequently

$$S(Y, s, \varepsilon) = \lim_{N \rightarrow \infty} S(Y, s, N, \varepsilon).$$

There is a critical value of the parameter s , which we denote by $h_{top}(T, Y, \varepsilon)$ such that

$$S(Y, s, \varepsilon) = \begin{cases} 0, & s > h_{top}(T, Y, \varepsilon), \\ \infty, & s < h_{top}(T, Y, \varepsilon). \end{cases}$$

⁴Limits exist by subadditivity.

Since $h_{top}(T, Y, \varepsilon)$ does not decrease with ε , the following limit exists,

$$h_{top}(T, Y) = \lim_{\varepsilon \rightarrow 0} (T, Y, \varepsilon).$$

We call $h_{top}(T, Y)$ the *topological entropy* of T restricted to Y or the topological entropy of Y (we denote $h_{top}(Y)$), as there is no confusion about T . We denote $h_{top}(X, T) = h_{top}(T)$.

3.4. Thermodynamic formalism of subadditive potentials. We call (X, T) a *dynamical system*, if $T : X \rightarrow X$ is a continuous map on compact metric space X . We say that $\Phi := \{\log \phi_n\}_{n=1}^{\infty}$ is a *subadditive potential* if each ϕ_n is a continuous positive-valued function on X such that

$$0 < \phi_{n+m}(x) \leq \phi_n(x)\phi_m(T^n(x)) \quad \forall x \in X, m, n \in \mathbb{N}.$$

Furthermore, $\Phi = \{\log \phi_n\}_{n=1}^{\infty}$ is said to be an *almost additive potential* if there exists a constant $C \geq 0$ such that for any $m, n \in \mathbb{N}$, $x \in X$, we have

$$C^{-1}\phi_n(x)\phi_m(T^n(x)) \leq \phi_{n+m}(x) \leq C\phi_n(x)\phi_m(T^n(x)).$$

We introduce the *topological pressure* of Φ as follows. The space X is endowed with a metric d . For any $n \in \mathbb{N}$, one can define a new metric d_n on X by

$$d_n(x, y) = \max\{d(T^k(x), T^k(y)) : k = 0, \dots, n\}.$$

For any $\varepsilon > 0$ a set $E \subset X$ is said to be a (n, ε) -*separated subset* of X if $d_n(x, y) > \varepsilon$ for any two different points $x, y \in E$. We define for a subadditive Φ

$$P_n(T, \Phi, \varepsilon) = \sup\left\{\sum_{x \in E} \phi_n(x) : E \text{ is } (n, \varepsilon)\text{-separated subset of } X\right\}.$$

Since $P_n(T, \Phi, \varepsilon)$ is a decreasing function of ε , We define

$$P(T, \Phi, \varepsilon) = \limsup_{n \rightarrow \infty} \frac{1}{n} \log P_n(T, \Phi, \varepsilon),$$

and

$$P_{\Phi}(T) = \lim_{\varepsilon \rightarrow 0} P(T, \Phi, \varepsilon).$$

We call $P_{\Phi}(T)$ the *topological pressure* of Φ . We denote by $P_{\Phi}(q)$ the topological pressure for subadditive potential $q\Phi$.

Bowen [B2] shows that for any Hölder continuous $\psi : X \rightarrow \mathbb{R}$ on transitive hyperbolic set (X, T) there exists a unique equilibrium measure μ_t (which is also a Gibbs state) for the additive potential $t \log \psi$ for $t \in \mathbb{R}$.

Feng and Käenmäki [FK] extend the Bowen's result for subadditive potential $t\Phi$ of locally constant cocycle under the assumption that the matrices ϕ_n do not preserve a common proper subspace of \mathbb{R}^d (i.e. (T, ϕ) is irreducible).

Recently, Park [P] shows that continuity of topological pressure, and uniqueness of equilibrium measure for typical cocycles. We show that one can prove the continuity of topological pressure on $GL(2, \mathbb{R})$ -fiber bunched cocycles which is weaker than Park's assumptions (see section 8).

Let $\vec{q} = (q_1, \dots, q_d) \in \mathbb{R}_+^d$, and $\vec{\Phi} = (\Phi_1, \dots, \Phi_d) = (\{\log \phi_{n,1}\}_{n=1}^\infty, \dots, \{\log \phi_{n,d}\}_{n=1}^\infty)$. Assume that $\vec{q} \cdot \vec{\Phi} = \sum_{i=1}^d q_i \Phi_i$ is a subadditive potential $\{q_i \log \phi_{n,i}\}_{n=1}^\infty$. We write topological pressure, maximal Lyapunov exponent, and minimal Lyapunov exponent of $\vec{\Phi}$, respectively

$$P_{\vec{q}}(q) = P(T, \vec{q}, \vec{\Phi}), \quad \vec{\beta}(\Phi) = \beta\left(\sum_{i=1}^d \Phi_i\right), \quad \vec{\alpha}(\Phi) = \alpha\left(\sum_{i=1}^d \Phi_i\right).$$

For $\mu \in \mathcal{M}(X, T)$, we write

$$\chi(\mu, \vec{\Phi}) = (\chi(\mu, \Phi_1), \dots, \chi(\mu, \Phi_d)),$$

where $\chi(\mu, \Phi_i) = \lim_{n \rightarrow \infty} \frac{1}{n} \int \log \phi_{n,i}(x) d\mu(x)$ for $i = 1, \dots, d$.

We can define topological pressure by the following variational principle. It was proved by Cao, Feng and Huang [CFH].

Theorem 3.3. [CFH, Theorem 1.1] *Let (X, T) be a dynamical system such that $h_{top}(T) < \infty$. For $\vec{t} \in \mathbb{R}_+^d$, suppose that $\vec{\Phi}$ is a subadditive potential on compact metric space (X, T) and $h_{top}(T) < \infty$. Then*

$$P_{\vec{\Phi}}(T, \vec{t}) = \sup\{h_\mu(T) + \vec{t} \cdot \chi(\mu, \vec{\Phi}) : \mu \in \mathcal{M}(X, T), \chi(\mu, \vec{\Phi}) \neq \infty\}.$$

We denote $P_{\vec{\Phi}}(T, \vec{t}) = P_{\vec{\Phi}}(\vec{t})$. Let $\vec{t} \in \mathbb{R}_+^d$, we denote by $E_{\vec{\Phi}}(\vec{t})$ the collection of invariant measures μ such that

$$h_\mu(T) + \vec{t} \cdot \chi(\mu, \vec{\Phi}) = P_{\vec{\Phi}}(\vec{t}).$$

If $E_{\vec{\Phi}}(\vec{t}) \neq \emptyset$, then each element $E_{\vec{\Phi}}(\vec{t})$ is called an *equilibrium state* for $\vec{t} \cdot \vec{\Phi}$.

In the remaining part of this section, we recall some theorems about multifractal formalism for subadditive potentials.

Theorem 3.4. [FH, Proposition 3.2] *Assume that $h_{top}(T) < \infty$, $\vec{\beta}(\Phi) > -\infty$. Then, $P_{\vec{\Phi}}(\cdot)$ is a real continuous convex function on \mathbb{R}_+^d and*

$$\partial P(\mathbb{R}_+^d) \subset (-\infty, \beta(\Phi_1)] \times \dots \times (-\infty, \beta(\Phi_d)].$$

Furthermore, $P'_{\vec{\Phi}}(\infty) := \vec{\beta}(\Phi)$.

Theorem 3.5. [FH, Theorem 1.1] *Let (X, T) be a dynamical system such that the topological entropy $h_{top}(T)$ is finite. Then $E(\beta(\Phi)) \neq \emptyset$. And*

$$\begin{aligned} h_{top}(T, E(\beta(\Phi))) &= \sup\{h_\mu(T) : \mu \in \mathcal{M}(X, T), \chi(\mu, \Phi) = \beta(\Phi)\} \\ &= \sup\{h_\mu(T) : \mu \in \mathcal{E}(X, T), \chi(\mu, \Phi) = \beta(\Phi)\}. \end{aligned}$$

The topological pressure is related to Lyapunov exponents in the following way.

Proposition 3.6. [FH, Theorem 3.3] *Let (X, T) be a dynamical system such that the entropy map $\mu \mapsto h_\mu(T)$ is upper semi-continuous and $h_{\text{top}}(T) < \infty$. For $t \in \mathbb{R}_+^d$, suppose that $\vec{t} \cdot \vec{\Phi}$ is a subadditive potential on compact metric space X which satisfies $\vec{\beta}(\Phi) > \infty$. Then,*

$$(3.2) \quad \partial P_{\vec{\Phi}}(\vec{t}) = \{\chi(\mu_{\vec{t}}, \vec{\Phi}) : \mu \in \text{Eq}(\vec{\Phi}, \vec{t})\}.$$

Moreover, $\text{Eq}(\vec{\Phi}, \vec{t})$ is a non-empty compact convex subset of $\mathcal{M}(X, T)$, for any $t \in \mathbb{R}_+^d$. Furthermore, the above results hold for $t \in \mathbb{R}^d$ when $\vec{\Phi}$ is almost additive.

The following lemma show that we can always approximate Lyapunov exponent of equilibrium measures by Lyapunov exponent of ergodic measures.

Lemma 3.7. [FH, Lemma 4.7] *Suppose that $h_{\text{top}}(T) < \infty$, $\vec{\beta}(\Phi) > -\infty$, and $q \in \mathbb{R}_+^d$. Let $\vec{\alpha} \in \partial P_{\vec{\Phi}}^e(q)$. Then for any $\varepsilon > 0$, there is a $\nu \in \mathcal{E}(X, T)$ such that*

$$|\chi(\nu, \vec{\Phi}) - \vec{\alpha}| < \varepsilon.$$

Theorem 3.8. [FH, Theorem 4.8] *Keep the assumption of Theorem (3.5), we also assume that the entropy map $\mu \mapsto h_\mu(T)$ is upper semi-continuous on $\mathcal{M}(X, T)$. Then*

- (i) *For any $t \in \mathbb{R}_+^d$, if $\alpha \in \partial P_{\vec{\Phi}}^e(t_*)$, where $t_* = t_-, t_+$, then $E(\vec{\alpha}) \neq \emptyset$. Moreover,*

$$\begin{aligned} h_{\text{top}}(T, E(\vec{\alpha})) &= \inf_{q \in \mathbb{R}_+^d} \{P_{\vec{\Phi}}(q) - \vec{\alpha} \cdot q\} \\ &= \max\{h_\mu(T) : \mu \in \mathcal{M}(X, T), \chi(\mu, \vec{\Phi}) = \vec{\alpha}\}. \end{aligned}$$

- (ii) *If $t \in \mathbb{R}_+^d$ such that $\vec{t} \cdot \vec{\Phi}$ has a unique equilibrium state $\mu_{\vec{t}} \in \mathcal{M}(X, T)$, then $\mu_{\vec{t}}$ is ergodic, $P'_{\vec{\Phi}}(t) = \chi(\mu_{\vec{t}}, \vec{\Phi})$, $E(P'_{\vec{\Phi}}(t)) \neq \emptyset$ and $h_{\text{top}}(T, E(P'_{\vec{\Phi}}(t))) = h_{\mu_{\vec{t}}}(T)$.*

Given an almost additive potential $\Phi = \{\log \phi_n\}_{n=1}^\infty$. Feng and Huang [FH, Lemma A.4.] proved the following lemma:

Lemma 3.9. *Let $\mu \in \mathcal{M}(X, T)$. Then, the map $\mu \mapsto \chi(\mu, \Phi)$ is continuous on $\mathcal{M}(X, T)$.*

3.5. Thermodynamic formalism of linear cocycles. Our motivation for studying thermodynamic formalism for subadditive potential are to deal with linear cocycles. Therefore, it has been proved a lot of results for linear cocycles. In this subsection, we recall some definitions and results that just prove for linear cocycles.

We recall that $T : \Sigma \rightarrow \Sigma$ is topologically mixing subshift of finite type and $\mathcal{A} : \Sigma \rightarrow GL(d, \mathbb{R})$ is a Hölder continuous function.

We study *ergodic optimization* of Lyapunov exponents. Ergodic optimization of Lyapunov exponents is concerned invariant measures in maximizing (or minimizing) the Lyapunov exponents. They were first considered by Rota and Strang [RS] and by Gurvits [GU], respectively. The associated growth rates are called upper joint spectral radius and lower joint spectral radius, respectively; they play an important role in Control Theory (see [J], [Bo]). We will explain in section 5 that ergodic optimization is related to thermodynamic formalism. In particular, we use thermodynamic formalism tools to prove continuity of lower joint spectral radius in section 6.

We define the *maximal Lyapunov exponent* as follows

$$\beta(\mathcal{A}) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \sup_{x \in X} \|\mathcal{A}^n(x)\|.$$

We denote by $\mathcal{M}(X, T)$ the space of all T -invariant Borel probability measure on X . We also denote by $\mathcal{E}(X, T)$ the space of all ergodic T -invariant Borel probability measure on X . Morris [M2] shows that the maximal Lyapunov exponent is equal to the supremum Lyapunov exponent of measure over invariant measures. That means,

$$(3.3) \quad \beta(\mathcal{A}) = \sup_{\mu \in \mathcal{M}(X, T)} \chi(\mu, \mathcal{A}).$$

Feng and Huang [FH] give different proof for this result.

Let us define the set of *maximizing measures* of \mathcal{A} to be the set of measures on X given by

$$\mathcal{M}_{max}(\mathcal{A}) := \{\mu \in \mathcal{M}(X, T), \beta(\mathcal{A}) = \chi(\mu, \mathcal{A})\}.$$

We also define the *minimal Lyapunov exponents* as follows

$$\alpha(\mathcal{A}) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \inf_{x \in X} \|\mathcal{A}^n(x)\|.$$

Similarly, the set of *minimizing measures* is defined as follows

$$\mathcal{M}_{min}(\mathcal{A}) := \{\mu \in \mathcal{M}(X, T), \alpha(\mathcal{A}) = \chi(\mu, \mathcal{A})\}.$$

We remark that supremum (3.3) is attained, so \mathcal{M}_{max} is non-empty set. But, \mathcal{M}_{min} is not necessarily a non-empty.

For $s_1 \geq s_2 \geq \dots \geq s_d$, one can define *generalized singular value function* $\phi^{s_1, \dots, s_d} : GL(d, \mathbb{R}) \rightarrow [0, \infty)$ as

$$\phi^{s_1, \dots, s_d}(\mathcal{A}) = \sigma_1^{s_1}(\mathcal{A}) \sigma_2^{s_2}(\mathcal{A}) \dots \sigma_d^{s_d}(\mathcal{A}),$$

when $s_i \in [0, d]$.

Similarly, we consider $\tilde{\phi}_A^{s_1, \dots, s_d}$ on \mathcal{L} and we define as follows, for any $n \in \mathbb{N}$ and $I \in \mathcal{L}(n)$,

$$\tilde{\phi}_A^{s_1, \dots, s_d}(I) := \max_{x \in I} \phi^{s_1, \dots, s_d}(\mathcal{A}^n(x)).$$

$\tilde{\phi}_{\mathcal{A}}^{s_1, \dots, s_d}$ is also submultiplicative. Moreover, for any $k > 0$ we can define

$$\tilde{\phi}_{\mathcal{A}}^k(I) := \max_{x \in I} \phi^k(\mathcal{A}^n(x)),$$

where $\phi^k(\mathcal{A}) = \|\mathcal{A}^{\wedge k}\|$. Denote

$$\vec{\Phi}_{\mathcal{A}} := (\{\log \tilde{\phi}_{\mathcal{A}}^1\}, \{\log \tilde{\phi}_{\mathcal{A}}^2\}, \dots, \{\log \tilde{\phi}_{\mathcal{A}}^d\}).$$

We define sum of top k Lyapunov exponents for any $n \in \mathbb{N}$ and $I \in \mathcal{L}(n)$ as follows

$$\chi_k(\mathcal{A}) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \tilde{\phi}_{\mathcal{A}}^k(I),$$

of the limit exists.

Similarly, we define sum of top k Lyapunov exponents of measure for any $n \in \mathbb{N}$ and $I \in \mathcal{L}(n)$ as follows

$$\chi_k(\mathcal{A}, \mu) := \lim_{n \rightarrow \infty} \frac{1}{n} \int \log \tilde{\phi}_{\mathcal{A}}^k(I) d\mu,$$

for $\mu \in \mathcal{M}(X, T)$.

We are mainly concerned with the distribution of the Lyapunov exponents of \mathcal{A} . More precisely, for any $\alpha \in \mathbb{R}$, define

$$E_{\mathcal{A}}(\alpha) = \{x \in X, \chi_1(\mathcal{A}) = \alpha\},$$

which is called the α -level set of $\chi(\mathcal{A})$.

We also define higher dimensional of level set of all of Lyapunov exponents as follows

$$E_{\mathcal{A}}(\vec{\alpha}) = \{x \in X, \chi_k(\mathcal{A}) = \alpha_k\},$$

for $1 \leq k \leq d$.

Park [P] uses quasi-multiplicative property $\mathcal{A} \in \mathcal{W}$ to show that continuity of topological pressure which it states in the following theorem. We remark that we prove that for $\mathcal{A} \in H_b^\alpha(\Sigma, GL(2, \mathbb{R}))$ in section 7.

Theorem 3.10. [P, Theorem B] *The map $(s, \mathcal{A}) \rightarrow P_{\tilde{\phi}_{\mathcal{A}}}(s)$ is continuous on $\Lambda_+^d \times \mathcal{W}$, where $\Lambda_+^d = \{(s_1, s_2, \dots, s_d) : s_1 \geq \dots \geq s_d \geq 0\}$.*

In locally constant setting, Feng and Käenmäki [FK] show that there exists a unique equilibrium measure μ_t for potential $t\Phi_{\mathcal{A}}$ under irreducible assumption. Moreover, μ_t is a Gibbs measure and $P_{\Phi_{\mathcal{A}}}(\cdot)$ is differentiable on \mathbb{R}_+ .

Theorem 3.11. [P, Theorem E] *Assume that $\mathcal{A} \in \mathcal{W}$. Then singular value functions $\tilde{\varphi}_{\mathcal{A}}^s$ with $s \in [0, d]$ are quasi-multiplicative.*

Consider the above theorem, the following theorem shows that we have the Feng and Käenmäki's result for typical cocycles.

Theorem 3.12. [F11, Theorem 5.5] *Let $\vec{\Phi}_{\mathcal{A}}$ be quasi-multiplicative and $\vec{t} \in \mathbb{R}_+^d$. $P_{\vec{\Phi}_{\mathcal{A}}}(\vec{t})$ has a unique equilibrium state $\mu_{\vec{t}}$ for the potential $\vec{t} \cdot \vec{\Phi}_{\mathcal{A}}$. Furthermore, $\mu_{\vec{t}}$ has the following Gibbs property: There exists $C \geq 1$ such that*

$$C^{-1} \leq \frac{\mu_{\vec{t}}([J])}{e^{-nP_{\vec{\Phi}_{\mathcal{A}}}(\vec{t}) + \vec{t} \cdot \log \vec{\Phi}_{\mathcal{A}}}} \leq C,$$

for any $n \in \mathbb{N}$ and $x \in [J]$. Furthermore, $P_{\vec{\Phi}_{\mathcal{A}}}(\cdot)$ is differentiable on \mathbb{R}_+^d and $P'_{\vec{\Phi}_{\mathcal{A}}}(\vec{t}) = \chi(\mu_{\vec{t}}, \mathcal{A})$.

Theorem 3.13. *Assume that $h_{top}(T) < \infty$, $\beta(\mathcal{A}) > -\infty$ and $\alpha(\mathcal{A}) < \infty$. If $\mathcal{A} \in \mathcal{W}$, then $P_{\vec{\Phi}_{\mathcal{A}}}(\cdot)$ is a real continuous convex function on \mathbb{R} . Moreover, $\alpha(\mathcal{A})$ exists and it is equal $P'_{\vec{\Phi}_{\mathcal{A}}}(-\infty) := \lim_{t \rightarrow -\infty} \frac{P_{\vec{\Phi}_{\mathcal{A}}}(t)}{t}$. Similarly, $P_{\vec{\Phi}_{\mathcal{A}}}$ is a real continuous convex function on \mathbb{R}^d . Furthermore,*

$$\begin{aligned} \vec{\alpha}(\mathcal{A}) &:= \min\{\alpha_i, \vec{\alpha} \in \vec{L}\} \\ &= \lim_{\vec{t} \rightarrow -\infty} \frac{P_{\vec{\Phi}_{\mathcal{A}}}(\vec{t})}{\vec{t}}. \end{aligned}$$

Proof. See [F, Lemmas 2.2 and 2.3]. We remark that although [F, Lemmas 2.2 and 2.3] only deal with locally constant cocycles, the proof given there works for our theorem under slightly modification. Indeed, Feng uses quasi-multiplicative property to prove the lemmas. Since $\mathcal{A} \in \mathcal{W}$, $\vec{\Phi}_{\mathcal{A}}$ is quasi-multiplicativity by Theorem 3.11. \square

4. MULTIFRACTAL FORMALISM OF TYPICAL COCYCLES

In this section we discuss multifractal formalism of typical cocycles. Our motivation for studying of the multifractal formalism associated to certain iterated function systems with overlaps. For instance, the Hausdorff dimension of level sets has been calculated for 2-dimension-planar affine iterated function systems satisfying strong irreducibility and the strong open set condition by B. Bárány, T. Jordan, A. Käenmäki, and M. Rams [BJKR]. In additive potential setting, the Lyapunov exponents are equal the Birkhoff averages. In this case, the restricted variational principle, topological entropy, and Hausdorff dimension level set has been studied by a lot of authors (see [C]). It is well known that for a dynamical system (X, T) , which is transitive, and Φ is an additive potential, then

$$(4.1) \quad h_{top}(T, E(\alpha)) = \inf\{P_{\Phi}(q) - \alpha q : q \in \mathbb{R}\},$$

and,

$$(4.2) \quad \inf\{P_{\Phi}(q) - \alpha q : q \in \mathbb{R}\} = \sup\{h_{\mu}(T) : \mu \in \mathcal{M}(X, T) \text{ with } \chi(\mu, \Phi) = \alpha\}.$$

We are interested to study (4.1) and (4.2) for typical cocycles. We will see (4.1) in Theorem 4.1 under certain assumptions. Moreover, we will prove (4.2) for typical cocycles in Theorem 4.2. We remark that Feng and Huang [FH, Theorem 5.2] prove the theorem for almost additive potentials. In fact, the proof of Theorem 4.2 is based on ideas from that work.

Theorem 4.1. *Let $\mathcal{A} \in \mathcal{W}$. Suppose that $P_{\vec{\Phi}_{\mathcal{A}}}(\vec{q}) \in \mathbb{R}$ for each $\vec{q} \in \mathbb{R}^d$. Then for $\vec{\alpha} \in \vec{L}$,*

$$(4.3) \quad h_{top}(T, E(\vec{\alpha})) = \inf\{P_{\vec{\Phi}_{\mathcal{A}}}(\vec{q}) - \vec{\alpha} \cdot \vec{q} : \vec{q} \in \mathbb{R}^d\}.$$

Proof. One can find the proof in [FH, Theorem 4.10] and [F, Theorem 1.1]. \square

Theorem 4.2. *Assume that $T : X \rightarrow X$ is a topologically mixing subshift of finite type on compact metric space X . Suppose that $\mathcal{A} : X \rightarrow GL(d, \mathbb{R})$ belongs to typical functions \mathcal{W} . Assume that ω is the range of the map from $\mathcal{M}(X, T)$ to \mathbb{R}^d*

$$\mu \mapsto (\chi_1(\mu, \mathcal{A}), \chi_2(\mu, \mathcal{A}), \dots, \chi_d(\mu, \mathcal{A})).$$

We define

$$\mathbf{h}(\vec{\alpha}) := \sup\{h_{\mu}(T) : \mu \in \mathcal{M}(X, T), \chi_i(\mu, \mathcal{A}) = \alpha_i\},$$

where $\vec{\alpha} \in \omega$. Then,

$$\mathbf{h}(\vec{\alpha}) = \inf\{P_{\vec{\Phi}_{\mathcal{A}}}(\vec{q}) - \vec{\alpha} \cdot \vec{q} : \vec{q} \in \mathbb{R}^d\}.$$

Proof. Fix $\vec{\alpha} = (\alpha_1, \dots, \alpha_d) \in \omega$. For $\mu \in \mathcal{M}(X, T)$, we define

$$V_{\vec{\alpha}}(\mu) := (\chi_1(\mu, \mathcal{A}) - \alpha_1, \dots, \chi_d(\mu, \mathcal{A}) - \alpha_d).$$

It is easy to see that there is $\mu' \in \mathcal{M}(X, T)$ such that $V_{\vec{\alpha}}(\mu') = \vec{0}$.

We write

$$A = \{V_{\vec{\alpha}}(\mu) : \mu \in \mathcal{M}(X, T)\}.$$

$V_{\vec{\alpha}}(\cdot)$ is a continuous affine function on $\mathcal{M}(X, T)$ (see remark 4). Therefore, A is convex compact set in \mathbb{R}^d .

We define $g : A \rightarrow \mathbb{R}$ by

$$g(\vec{t}) = \sup\{h_{\mu}(T) : \mu \in \mathcal{M}(X, T), V_{\vec{\alpha}}(\mu) = \vec{t}\}.$$

It is easy to see that g is a concave and upper semi continuous function on A . We have $\mathbf{h}(\vec{\alpha}) = g(\vec{0})$. We define

$$W(\vec{q}) := \sup\{g(\vec{t}) + \vec{q} \cdot \vec{t} : \vec{t} \in \mathbb{R}^d\},$$

for all $\vec{q} \in A$. Then, we have

$$g(\vec{t}) = \inf\{W(\vec{q}) - \vec{q} \cdot \vec{t} : \vec{q} \in \mathbb{R}^d\}$$

for all $\vec{t} \in A$, by Corollary 3.2. So, we have

$$\mathbf{h}(\vec{\alpha}) = g(\vec{0}) = \inf\{W(\vec{q}) : \vec{q} \in \mathbb{R}^d\}.$$

By Theorem 3.13, $P_{\vec{\Phi}_{\mathcal{A}}}(\vec{q})$ is convex of \mathbb{R}^d . Then, by variational principle

$$W(\vec{q}) = P_{\vec{\Phi}_{\mathcal{A}}}(\vec{q}) - \vec{\alpha} \cdot \vec{q}.$$

□




FIGURE 3. $P_\Phi(\cdot)$ is a convex function for $q \in \mathbb{R}$. The blue line is tangent to $P_\Phi(\cdot)$ at q with slope $\alpha = P'_\Phi(q)$.

Remark 5. *In the locally constant cocycles case, Theorem 4.2 is true under strongly irreducible assumption, which means we do not need pinching assumption in this case.*

5. CONTINUITY OF ENTROPY SPECTRUM OF LYAPUNOV EXPONENTS

In this section, we will discuss continuity of entropy spectrum of Lyapunov exponents, which is the topological entropy of level sets of points with a common given Lyapunov exponent. In the locally constant cocycles case, Lemma 5.1 follows from Feng and Shmerkin's paper [FS]. We show the continuity of entropy spectrum at boundary of Lyapunov exponents for typical cocycles \mathcal{W} in the sense,

$$h_{top}(E(\alpha_t)) \rightarrow h_{top}(E(\beta(\mathcal{A}))) \text{ when } t \rightarrow \infty.$$

We define

$$\lambda_e(\mathcal{A}, (s_1, \dots, s_d)) := \lim_{n \rightarrow \infty} \frac{1}{n} \int \log \phi^{s_1, \dots, s_d}(\mathcal{A}^n(x)) d\mu(x),$$

where $\mu \in Eq(\mathcal{A}, (s_1, \dots, s_d))$.

Lemma 5.1. *Assume that $\mathcal{A}_k, \mathcal{A} \in \mathcal{W}$ with $\mathcal{A}_k \rightarrow \mathcal{A}$. For $t_k, t > 0$, assume that $t_k \rightarrow t$. Let α_k and α be the derivatives of $P_{\Phi_{\mathcal{A}_k}}()$ and $P_{\Phi_{\mathcal{A}}}()$ at t_k and t , respectively. Then,*

$$\lim_{k \rightarrow \infty} h_{top}(E(\alpha_k)) = h_{top}(E(\alpha)).$$

Proof. According to Theorem 3.12 $P_{\Phi_{\mathcal{A}}}()$ is differentiable for any $t > 0$ and there is a unique equilibrium measure μ_t for potential $t\Phi_{\mathcal{A}}$. So, we have

$$h_{top}(E(\alpha)) = h_{\mu_t}(T),$$

where $P'_{\Phi_{\mathcal{A}}}(t) = \alpha$, by Theorem 3.8.

Taking into account the observation above, to prove the theorem it is enough to show that $h_{\mu_{t_k}}(T) \rightarrow h_{\mu_t}(T)$ for proving the theorem.

By definition of $Eq(\mathcal{A}_k, t_k)$,

$$P_{\Phi_{\mathcal{A}_k}}(t_k) = h_{\mu_{t_k}}(T) + \lambda_e(\mathcal{A}_k, t_k).$$

Notice that the Lyapunov exponents are upper semi-continuous. Moreover, mixing subshift of finite type map $T : X \rightarrow X$ implies upper semi-continuity of the entropy map $\mu \mapsto h_{\mu}(T)$. Now, we conclude from above observations and Theorem 3.10,

$$\begin{aligned} P_{\Phi_{\mathcal{A}}}(t) &= \lim_{k \rightarrow \infty} P_{\Phi_{\mathcal{A}_k}}(t_k) \\ &= \lim_{k \rightarrow \infty} h_{\mu_{t_k}}(T) + \lambda_e(\mathcal{A}_k, t_k) \\ &\leq h_{\mu}(T) + \lambda_e(\mathcal{A}, t). \end{aligned}$$

This shows $\mu \in Eq(\mathcal{A}, t)$ and $\mu_k \rightarrow \mu$ ⁵. Moreover, we have equality in the above, that gives the claim. Furthermore, it shows that the continuity of Lyapunov exponent of equilibrium measures. \square

Assume that (μ_t) is a sequence of equilibrium measures for a subadditive potential $t \log \Phi$ for $t > 0$. The following theorem consider the behavior of the (μ_t) as $t \rightarrow \infty$. In the thermodynamic interpretation of the parameter t is the *inverse temperature*, limit $t \rightarrow \infty$ is called *zero temperature limit*, and the accumulation points of the measure (μ_t) as $t \rightarrow \infty$ are called *ground states*.

Zero temperature limit is also related to ergodic optimization, as for $t \rightarrow \infty$ any accumulation points of equilibrium measure μ_t is a maximizing measure Φ .

⁵By weak-* compactness μ_k has a accumulation point, let us call μ . According the above observation μ is an equilibrium measure for $t\Phi_{\mathcal{A}}$. Then uniqueness of equilibrium measure implies the limit.

Theorem 5.2. [M, Theorem 1.1] *Let (X, T) be a dynamical system for which the entropy map $\mu \mapsto h_\mu(T)$ is upper semi-continuous and topological entropy $h_{top}(T) < \infty$. Assume that $\Phi = \{\log \phi_n\}_{n=1}^\infty$ is a subadditive potential on compact metric X . Then a family of equilibrium measures (μ_t) for potential $t\Phi$, where $t > 0$, has a weak* accumulation point as $t \rightarrow \infty$. Any such accumulation point is a Lyapunov maximizing measure for Φ . Furthermore,*

- (i) $\chi(\Phi, \mu) = \lim_{t \rightarrow \infty} \chi(\Phi, \mu_t)$,
- (ii) $h_\mu(T) = \lim_{t \rightarrow \infty} h_{\mu_t}(T) = \max\{h_\nu(T), \nu \in \mathcal{M}_{\max}(\Phi)\}$.

Moreover, above result works for higher dimensional case.

We use the above theorem to prove the following theorem.

Theorem 5.3. *Suppose that $\mathcal{A} \in \mathcal{W}$. If $\alpha_t = P'_{\Phi, \mathcal{A}}(t)$ ⁶ for $t > 0$. Then,*

$$h_{top}(E(\alpha_t)) \rightarrow h_{top}(E(\beta(\mathcal{A}))) \text{ when } t \rightarrow \infty.$$

Proof. Since $\mathcal{A} \in \mathcal{W}$, Theorem 3.12 implies that for $t > 0$, there is a unique equilibrium state μ_t for the potential $t\Phi_{\mathcal{A}}$ such that

$$\chi(\mu_t, \mathcal{A}) = \alpha_t = P'_{\Phi, \mathcal{A}}(t).$$

By Theorem (3.8),

$$h_{top}(E(\alpha_t)) = h_{\mu_t}(T).$$

We know that

$$h_{top}(E(\beta(\mathcal{A}))) = \sup\{h_\mu(T), \mu \in \mathcal{M}(X, T), \chi(\mu, \mathcal{A}) = \beta(\mathcal{A})\}$$

by Theorem 3.5. So, we only need to show that

$$h_{\mu_t}(T) \rightarrow \sup\{h_\mu(T), \mu \in \mathcal{M}(X, T), \chi(\mu, \mathcal{A}) = \beta(\mathcal{A})\}.$$

That follows from Theorem 5.2. □

We denote

$$\vec{\beta}(\mathcal{A}) := (\beta_1(\mathcal{A}), \beta_2(\mathcal{A}^{\wedge 2}), \dots, \beta_d(\mathcal{A}^{\wedge d})),$$

where $\beta_i(\mathcal{A}^{\wedge i}) = \lim_{n \rightarrow \infty} \frac{1}{n} \log \sup_{x \in \Sigma} \|(\mathcal{A}^n)^{\wedge i}(x)\|$ for $i = 1, \dots, d$.

Theorem 5.4. *Suppose that $\mathcal{A}_l, \mathcal{A} \in \mathcal{W}$ with $\mathcal{A}_l \rightarrow \mathcal{A}$, and $\vec{t}_l, \vec{t} \in \mathbb{R}_+^d$ such that $t_l \rightarrow t$. Let $\vec{\alpha}_l$ and $\vec{\alpha}$ be derivative of $P_{\vec{\Phi}, \mathcal{A}_l}(\cdot)$ and $P_{\vec{\Phi}, \mathcal{A}}(\cdot)$ at t_l and t , respectively. Then,*

$$\lim_{l \rightarrow \infty} h_{top}(E(\vec{\alpha}_l)) = h_{top}(E(\vec{\alpha})).$$

Moreover,

$$h_{top}(E(\vec{\alpha}_l)) \rightarrow h_{top}(E(\vec{\beta}(\mathcal{A}))) \text{ when } l \rightarrow \infty.$$

Proof. The proof is similar to Theorem 5.3 and is omitted. □

⁶We remind the reader that $P(\cdot)$ is differentiable for $\mathcal{A} \in \mathcal{W}$ according to Theorem 3.12

6. CONTINUITY OF THE LOWER JOINT SPECTRAL RADIUS

In this section, we are going to prove the continuity of the lower joint spectral radius. This kind of result is known by Bochi and Morris [BM] under 1-domination assumption for locally constant cocycles. Breuillard and Sert [BS] extend their result to the joint spectrum of locally constant cocycles. Moreover, they gave a counter example [BS, Example 4.13] that shows that we have discontinuity the lower joint spectral for typical cocycles. Even though, we have a lot of results for the upper spectral radius, we have few result about the lower spectral radius, which shows that working on the later case is much harder than the former case.

Under t -domination assumption for all $1 \leq t \leq d - 1$, we prove continuity of the vector of lower joint spectral for linear cocycles, which is generated by a topologically mixing subshift of finite type $T : \Sigma \rightarrow \Sigma$ and a Hölder continuous function $\mathcal{A} : \Sigma \rightarrow GL(d, \mathbb{R})$.

Assume that $T : X \rightarrow X$ a diffeomorphism on compact invariant set X . Let $V \oplus W$ be a splitting of the tangent bundle over X that is invariant by the tangent map DT . In this case, if vectors in V are uniformly contracted by DT and vectors in W are uniformly expanded, then the splitting is called *hyperbolic*. Moreover, the splitting is called *dominated*, if each point all vectors in V are more contracted than all vectors in W . Domination could also be called uniform projective hyperbolicity. Indeed, domination is equivalent to V being hyperbolic repeller and W being hyperbolic attractor in the projective bundle.

In the linear cocycles, we are interested in bundles of the form $X \times \mathbb{R}^d$, where the linear cocycle is generated by (T, \mathcal{A}) . Bochi and Gourmelon [BGO] show that a cocycle admits a dominated splitting $V \oplus W$ with $\dim V = k$ if and only if when $n \rightarrow \infty$, the ratio between the $k - th$ and $(k + 1) - th$ singular values of the matrices of the $n - th$ iterate increase uniformly exponentially. In fact, they extended Yoccoz's result [Y] that was proved it for 2-dimensional vector bundles.

Definition 6.1. We say that \mathcal{A} is k -dominated, or that k is an index of domination for \mathcal{A} , if and only if there are constants $C > 0$ and $0 < \tau < 1$ such that

$$\frac{\sigma_{k+1}(\mathcal{A}^n(x))}{\sigma_k(\mathcal{A}^n(x))} \leq Ce^{\tau n} \quad \forall x \in X, \forall n \in \mathbb{N}.$$

We define the *upper joint spectral radius* of $\mathcal{A} : \Sigma \rightarrow GL(d, \mathbb{R})$ as follows

$$\hat{\rho}(\mathcal{A}) := \limsup_{n \rightarrow \infty} \{ \|\mathcal{A}(I)\|^{\frac{1}{n}} : I \in \mathcal{L}(n) \}.$$

It is easy to see that $\beta(\mathcal{A}) = \log \hat{\rho}(\mathcal{A})$. Similarly, we define the *lower joint spectral radius* of $\mathcal{A} : \Sigma \rightarrow GL(d, \mathbb{R})$ as follows

$$\check{\rho}(\mathcal{A}) := \liminf_{n \rightarrow \infty} \{ \|\mathcal{A}(I)\|^{\frac{1}{n}} : I \in \mathcal{L}(n) \}.$$

The *minimal Lyapunov exponents* $\vec{\alpha}(\mathcal{A})_d = (\alpha_1(\mathcal{A}), \dots, \alpha_d(\mathcal{A}))$ is the quantity

$$\lim_{n \rightarrow \infty} \frac{1}{n} \inf_{x \in X} \log \sigma_i(\mathcal{A}^n(x)) = \alpha_i(\mathcal{A}),$$

where $1 \leq i \leq d$. We remark that the limit exists under domination assumption.

The interior and topological closure of a set A is denoted by $\overset{\circ}{A}$ and \overline{A} .

Assume that $f : X \rightarrow X$ is a convex continuous function on compact metric space X . We have $\overline{\partial f(\mathbb{R})} = \partial f(\mathbb{R}) \cup \{f'(\infty)\} \cup \{f'(-\infty)\}$, where $\partial f(\mathbb{R})$ is defined as in (7.4).

Theorem 6.2. *Assume that \mathcal{A} is 1-domination. Then $\alpha(\mathcal{A})$ can be approximated by Lyapunov exponents of equilibrium measures for a subadditive potential $t\Phi_{\mathcal{A}}$ on compact metric space X for $t \in \mathbb{R}$. Moreover, minimizing measure for \mathcal{A} exists.*

Proof. Let $\alpha = \alpha(\mathcal{A})$. By Theorem 3.13, $\alpha = P'_{\Phi_{\mathcal{A}}}(-\infty)$.

According to convexity of $P_{\Phi_{\mathcal{A}}}$, there exists a sequence (t_j) such that $P'_{\Phi_{\mathcal{A}}}(t_j) =: \alpha_j$ exists for every $j \in \mathbb{N}$ and $\alpha_j \rightarrow \alpha$ as $j \rightarrow \infty$. There exists $\mu_j \in \text{Eq}(\mathcal{A}, t_j)$ such that $\chi(\mu_j, \Phi) = \alpha_j$ for all j , by Proposition 3.6. Let μ be an accumulation⁷ point of sequence (μ_j) as $j \rightarrow \infty$. By Lemma 3.9, we have

$$\chi(\mu_j, \mathcal{A}) \rightarrow \chi(\mu, \mathcal{A}) = \alpha.$$

Furthermore, our proof shows that minimizing measure exists. □

Now, we can show that the continuity of lower joint spectral radius.

Theorem 6.3. *Assume that $\mathcal{A}, \mathcal{A}_k$ are 1-domination. Then, $\alpha(\mathcal{A}_k) \rightarrow \alpha(\mathcal{A})$ when $\mathcal{A}_k \rightarrow \mathcal{A}$.*

Proof. According to Theorem 6.2, $\alpha(\mathcal{A})$ can be approximated by Lyapunov exponents of equilibrium measures for a subadditive potential $t\Phi_{\mathcal{A}}$ on compact metric space X for $t \in \mathbb{R}$. So, it is enough to show that

$$(6.1) \quad \chi(\mu_k, \mathcal{A}_k) \rightarrow \chi(\mu, \mathcal{A}),$$

where μ, μ_k are equilibrium measures.

1-domination assumption implies that \mathcal{A} is almost additive. Hence, there exist a unique equilibrium measure for potential $t\Phi_{\mathcal{A}}$ where $t \in \mathbb{R}$. So, (6.1) follows from the proof of Lemma 5.1. □

It is possible to obtain the following generalization of Theorem 6.2 to the joint spectrum of all Lyapunov exponents. One can also obtain continuity lower joint spectral radius for all Lyapunov exponents.

⁷Eq(\mathcal{A}, t) is compact in weak-* topology.

Theorem 6.4. *Assume that \mathcal{A}_k and \mathcal{A} satisfy t -domination for all $1 \leq t \leq d - 1$. Then*

$$\vec{\alpha}(\mathcal{A}_k)_d \rightarrow \vec{\alpha}(\mathcal{A})_d,$$

when $\mathcal{A}_k \rightarrow \mathcal{A}$.

The proof is similar of Theorem 6.3 and is omitted.

7. RELATION BETWEEN TOPOLOGICAL SPECTRUM AND LYAPUNOV SPECTRUM FOR TYPICAL COCYCLES

In this section, we are going to show that the closure of positive topological spectrum is equal the Lyapunov spectrum for typical cocycles. This result is first attempt to characterize Lyapunov spectrum as a set of positive entropy spectrum. The main input of our argument will be this fact that topological pressure is convex for typical cocycles. Then, we can show the concavity of entropy spectrum of Lyapunov exponents.

We recall that $T : X \rightarrow X$ is topologically mixing subshift of finite type and $\mathcal{A} : X \rightarrow GL(d, \mathbb{R})$ is a Hölder continuous function. We always assume that we have more than a matrix, otherwise it might be topological entropy equal zero. Let us recall that the definition of maximal Lyapunov spectrum as follows

$$L = \{\alpha \in \mathbb{R}, \exists x \in \Sigma, \chi_1(\mathcal{A}) = \alpha\}.$$

We also recall the definition the Lyapunov spectrum for all Lyapunov exponents as follows

$$\vec{L} = \{\vec{\alpha} \in \mathbb{R}^d, \exists x \in X, \chi_i(\mathcal{A}) = \alpha_i\},$$

where $i = 1, 2, \dots, d$.

Lemma 7.1. *Let $\mathcal{A} \in W$. Then, $h_{top}(E(\alpha))$ is concave on convex set L .*

Proof. The topological pressure $P_{\Phi_{\mathcal{A}}}(\cdot)$ is convex by Theorem 3.13. Moreover, according to the Theorems 4.1 and 4.2 we have

$$\begin{aligned} h_{top}(E(\alpha)) &= \inf_{t \in \mathbb{R}} \{P_{\Phi_{\mathcal{A}}}(t) - \alpha t\} \\ &= \sup \{h_{\mu}(T) : \mu \in \mathcal{M}(X, T), \chi(\mu, \mathcal{A}) = \alpha\}. \end{aligned}$$


Hence, it is concave. □

Theorem 7.2. *For $\alpha \in \overset{\circ}{L}$, $h_{top}(E(\alpha)) > 0$.*

Proof. By Theorem 7.1 $h_{top}(E(\alpha))$ is concave. So, $h_{top}(E(\alpha)) > 0$. Indeed, a concave function with a maximum in the interior of the domain is non-decreasing to the left of the maximum and non-increasing to the right of the maximum. □

Remark 6. *Entropy spectrum at boundary of Lyapunov spectrum is not necessarily positive. In fact, there is a conjecture, which is known as Meta conjecture, that says that under generic assumptions the entropy spectrum at boundary of Lyapunov spectrum is zero (which would mean that $h_{top}(E(\beta(\mathcal{A})) = h_{top}(E(\alpha(\mathcal{A})) = 0)$); this*

phenomenon is often referred to as ergodic optimization of Lyapunov exponents, see for example [Bo]. In the additive potential case, instead, this phenomenon is often referred to as ergodic optimization of Birkhoff averages, see for example [J].



Theorem 7.3. $\overline{\{\alpha \in \mathbb{R}, h_{top}(E(\alpha)) > 0\}} = L$.

Proof. That is direct consequence Theorem 7.2. □

In fact, Park [P] proves Theorem 2.9 for higher dimensional case. That means, \vec{L} is closed and convex. So, we can obtain the following generalization of Theorem 7.3 to the Lyapunov spectrum of of all Lyapunov exponents.

Theorem 7.4. $\overline{\{\vec{\alpha} \in \mathbb{R}^d, h_{top}(E(\vec{\alpha})) > 0\}} = \vec{L}$.

We remark that the concavity of a function defined on a convex set implies the continuity of the function in the interior, and that the continuity of the entropy under the change of the Lyapunov exponents implies the continuity of the Lyapunov spectrum.

8. THE CONTINUITY OF TOPOLOGICAL PRESSURE

In this section, we are going to prove the continuity topological pressure for $H_b^r(\Sigma, GL(2, \mathbb{R}))$. In the locally constant setting, Feng and Shmerkin [FS] prove that we have the continuity of topological pressure. Recently, Park [P] prove that we have the continuity of topological pressure for typical cocycles. We recall that typical means that a linear cocycle is pinching, twisting and fiber bunching. The techniques we use in the proof are inspired from result [FS]. Our result shows that one can prove the continuity of topological under weaker assumption that Park assumed. The main input our argument is the continuity of Lyapunov expoents that was proved by Backes, Brown, and Butler [BBB] for $H_b^r(\Sigma, GL(2, \mathbb{R}))$.

Moreover, we use our result to show that set of $\varphi_{\mathcal{A},n}$ -equilibrium states for upper triangular matrices that belongs to $H_b^r(\Sigma, GL(2, \mathbb{R}))$ is equal set of equilibrium states its diagonal.

We write for $\mathcal{A} \in H_b^r(X, GL(2, \mathbb{R}))$ and $s \geq 0$.

Theorem 8.1. *The map $(\mathcal{A}, s) \mapsto P_{\tilde{\Phi}_{\mathcal{A}}}(s)$ is continuous on $[0, \infty) \times H_b^r(\Sigma, GL(2, \mathbb{R}))$.*

Proof. Since topological pressure is subadditive, it is upper semi continuous. So, it is enough to show that it is lower semi-continuous.

Assume that $\mathcal{A}_k \in H_b^r(\Sigma, GL(2, \mathbb{R}))$ and $s_k \in (1, 2)$. We can assume that there is an ergodic measure $\mu \in Eq(\mathcal{A}, s)$ by Lemma 3.7. Then, by variational principle

$$(8.1) \quad \begin{aligned} P_{\tilde{\Phi}_{\mathcal{A}_k}}(s_k) &\geq h_\mu(T) + \lambda_e(\mathcal{A}_k, s_k) \\ &= h_\mu(T) + (2 - s_k)\lambda_e(\mathcal{A}_k, 1) + (s_k - 1)\lambda_e(\mathcal{A}_k, 2). \end{aligned}$$

Notice that

$$\lambda_e(\mathcal{A}_k, 2) = \lim_{n \rightarrow \infty} \frac{1}{n} \int \log \|\det(\mathcal{A}_k(x))\| d\mu(x).$$

So, when $\mathcal{A}_k \rightarrow \mathcal{A}$, we have

$$\lambda_e(\mathcal{A}_k, 2) \rightarrow \lambda_e(\mathcal{A}, 2),$$

and

$$\lambda_e(\mathcal{A}_k, 1) \rightarrow \lambda_e(\mathcal{A}, 1).$$

by Theorem 2.7. Then, by (8.1),

$$\liminf_{(\mathcal{A}_k, s_k) \rightarrow (\mathcal{A}, s)} P_{\tilde{\Phi}_{\mathcal{A}_k}}(s_k) \geq h_\mu(T) + \lambda_e(\mathcal{A}, s) = P_{\tilde{\Phi}_{\mathcal{A}}}(s).$$

This proves the continuity of $P_{\tilde{\Phi}_{\mathcal{A}}}(\cdot)$ at (\mathcal{A}, s) . \square

Remark 7. *Recently, C. Freijo and K. Marin [FK2] extend the Backes, Brown, and Butler's result to non-uniformly fiber-bunched cocycles. According to their result, one can prove the above theorem for non-uniformly fiber-bunched cocycles.*

Remark 8. *Recently, Cao, Pesin, and Zhao [CPZ] show that $(\mathcal{A}, s) \mapsto P_{\tilde{\varphi}_{\mathcal{A}}}(s)$ is continuous on $[0, \infty) \times H^\alpha(\Sigma, GL(d, \mathbb{R}))$ with using a different approach.*

8.1. Application of Theorem 8.1. In the locally constant cocycles case, Falconer and Miao [FM, Theorem 2.5] show that the set of φ^s -equilibrium states of upper triangular matrices is precisely the set of φ^s -equilibrium states its diagonal. Käenmäki and Morris [KM, Proposition 6.2] extend Falconer and Miao's result for higher dimensional case. One can use Käenmäki and Morris's proof and Theorem 8.1 to obtain the following result :

Corollary 8.2. *Let $\mathcal{A} \in H^r(\Sigma, GL(2, \mathbb{R}))$ be an upper triangular matrices :*

$$\mathcal{A}(x) := \begin{bmatrix} a(x) & b(x) \\ 0 & c(x) \end{bmatrix}.$$

Then, the set of $\tilde{\varphi}_A^s$ -equilibrium states of upper triangular matrices \mathcal{A} is precisely the set of $\tilde{\varphi}_A^s$ -equilibrium states its diagonal :

$$\mathcal{A}'(x) := \begin{bmatrix} a(x) & 0 \\ 0 & c(x) \end{bmatrix}.$$

Remark 9. Recently, Butler and Park [BP] proved some results in this direction for 2-dimensional cocycles.

REFERENCES

- [AEV] A. Avila, A. Eskin and M. Viana, Continuity of Lyapunov exponents of random matrix-products, In preparation.
- [AV] A. Avila and M. Viana, Simplicity of lyapunov spectra: a sufficient criterion, *Portugaliae Mathematica* **64** (2007), 311-376.
- [BBB] L. Backes, A. Brown, and C. Butler, Continuity of Lyapunov exponents for cocycles with invariant holonomies. *J. Mod. Dyn.* **12**, , 223–260. 2018.
- [Boc] J. Bochi, Discontinuity of the Lyapunov exponents for non-hyperbolic cocycles. <http://www.mat.uc.cl/~jairo.bochi/>.
- [Boc1] J. Bochi, Genericity of zero Lyapunov exponents, *Ergodic Theory and Dynamical Systems*, **22** (2002), 1667-1696.
- [Bo] J. Bochi, Ergodic optimization of Birkhoff averages and Lyapunov exponents . *Proceedings of the International Congress of Mathematicians 2018, Rio de Janeiro*, vol. 2, pp. 1821-1842.
- [BG] J. Bochi and E. Garibaldi, Extremal norms for fiber bunched cocycles, *Journal de l'École polytechnique - Mathématiques*. **6** (2019), 947-1004.
- [BGO] J. Bochi and N.Gourmelon, Some characterizations of domination, *Math.Z.* **263** (1)(2009), 221-231.
- [BM] J. Bochi, I.D. Morris, Continuity properties of the lower spectral radius, *Proc. Lond. Math. Soc.*, **110** 477–509. 2015.
- [BGMV] C. Bonatti, X. Gómez-Mont, and M. Viana. Généricité déxposants de lyapunov non-nuls pour des produits déterministes de matrices. *Annales de Unstitut Henri Poincaré (C)Non Linear Analysis*, vol. 20, Elsevier, 2003, pp. 579-624.
- [BJKR] B. Bárány, T. Jordan, A. Käenmäki, and M. Rams, Birkhoff and Lyapunov spectra on planar self-affine sets, <https://arxiv.org/abs/1805.08004>.
- [BS] E. Breuillard and C. Sert. The joint spectrum, <https://arxiv.org/abs/1809.02404>.
- [B] R. Bowen. Markov partitions for axiom a diffeomorphisms. *American Journal of Mathematics*, **92**(3): 725-747, 1970
- [B1] ———. Topological entropy for noncompact sets. *American Journal of Mathematics*, **184**, 125-136, 1973.
- [B2] ———. Some systems with unique equilibrium states, *Mathematical Systems Theory*, **8**(93): 193-202, 1974.
- [BV] C. Bocker-Neto, and M. Viana. Continuity of Lyapunov exponents for random two-dimensional matrices. *Ergod. Th. & Dynam. Sys.*, **5**, 1413-1442, 2017.
- [BV1] C. Bonatti, and M. Viana, Lyapunov exponents with multiplicity 1 for deterministic products of matrices. *Ergod. Th. & Dynam. Sys.*, **24** (5): 1295-1330, 2004.
- [Bu] C. Butler, discontinuity of Lyapunov exponents near fiber bunched cocycles, *Ergodic Theory and Dynamical Systems*, **38**(2018), 523–539.
- [BP] C. Butler and K. Park, Thermodynamic formalism for $GL_2(\mathbb{R})$ -cocycles with canonical holonomies, <https://arxiv.org/abs/1909.11548>.

- [C] V. Climenhaga, The thermodynamic approach to multifractal analysis. *Ergod. Th. & Dynam. Sys.* **34**(5)(2014), 1409–1450.
- [CFH] Y. Cao, D. Feng, and W. Huang. The thermodynamic formalism for sub-additive potentials. *Discrete Contin. Dyn. Syst.*, **20**(3): 639-657, 2008.
- [CPZ] Y. Cao, Yakov Pesin and Y. Zhao, Dimension estimates for non-conformal repellers and-continuity of sub-additive topological pressure, <https://arxiv.org/abs/1906.07323>.
- [FM] K. Falconer and J. Miao. Dimensions of self-affine fractals and multifractals generated by upper-triangular matrices, *Fractals*, **15**(3):(2007), 289-299.
- [F] D. Feng. Lyapunov exponents for products of matrices and multifractal analysis. Part II. General matrices. *J. Math.*, **170**: 355-394, 2009.
- [F11] F. Feng, Equilibrium states for factor maps between subshifts, *Advances in Mathematics* **226** (2011), no. 3, 2470-2502.
- [FH] D. Feng and W. Huang. Lyapunov spectrum of asymptotically sub-additive potentials. *Comm. Math. Phys.*, **297**(1): 1-43, 2010.
- [FK] D. Feng and A. Käenmäki. Equilibrium states of the pressure function for products of matrices. *Discrete Contin. Dyn. Syst.*, **30**(3): 699-708, 2011.
- [FS] D. Feng and P. Shmerkin. Non-conformal repellers and the continuity of pressure formatrix cocycles, *Geometric and Functional Analysis*, **24**(4): 1101-1128, 2014.
- [FK1] H. Furstenberg, and H. Kesten, Products of random matrices, *the Annals of Mathematical Statistics*, **31**(2) (1960), 457-469.
- [FK2] C. Freijo and K. Marin, Continuity of Lyapunov exponents for non-uniformly fiber bunched cocycles, <https://arxiv.org/abs/1910.14102>.
- [GU] L. Gurvits, Stability of discrete linear inclusion, *Linear Algebra Appl.*, **231** (1995), 47-85.
- [J] O. Jenkinson. Ergodic optimization in dynamical systems. *Ergod. Th.& Dynam. Sys.*, **39**(10): 2593-2618, 2019.
- [KH] A. Katok and B. Hasselblatt. *Introduction to the modern theory of dynamical systems*. Cambridge University Press, London-New York, 1995.
- [KS] B. Kalinin, and V. Sadovskaya, Cocycles with one exponent over partially hyperbolic systems, *Geometriae Dedicata*, **167** (2013), no. 1, 167-188.
- [KS1] B. Kalinin, and V. Sadovskaya, Holonomies and cohomology for cocycles over partially hyperbolic diffeomorphisms, *Discrete and Continuous Dynamical Systems* **36**, **1** (2016), 245-259.
- [KM] A. Käenmäki and I. Morris, Structure of equilibrium states on self-affine sets and strict monotonicity of affinity dimension. *Proc. Lond.Math. Soc.* (**3**) **116** (2018), no. 4, 929-956.
- [LM] L. Douglas and M. Brian, *An introduction to symbolic dynamics and coding*, Cambridge University Press, 1995
- [M] R. Mohammadpour, Zero temperature limits of equilibrium states for sub-additive potentials and approximation of maximal Lyapunov exponent, <https://arxiv.org/abs/1910.07279>.
- [M2] I.D. Morris, A rapidly-converging lower bound for the joint spectral radius via multiplicative ergodic theory. *Adv. Math.* , **225**(6): 3425-3445, 2010.
- [P] K. Park. Quasi-Multiplicativity of typical cocycles. <https://arxiv.org/abs/1903.03928>.
- [R] R. T. Rockafellar, *Convex analysis*. Princeton University Press, Princeton, N.J. 1970.
- [RS] G.-C. Rota and G. Strang, A note on the joint spectral radius, *Indag. Math.*, **22** (1960), 379-381.
- [V] M. Viana, *Lectures on Lyapunov exponents*, Cambridge University Press, 2014.
- [VI] M. Viana, (Dis)continuity of Lyapunov exponents, *Ergodic Theory and Dynamical Systems*, 1-35. doi:10.1017/etds.2018.50.

- [Y] J. Yoccoz, Some questions and remarks about $SL(2, \mathbb{R})$ cocycles, In: *Modern Dynamical Systems and Applications*, Cambridge University Press, Cambridge, (2004), 447-458.

DEPARTMENT OF DYNAMICAL SYSTEMS, INSTITUTE OF MATHEMATICS OF THE POLISH
ACADEMY OF SCIENCES, UL. ŚNIADECKICH 8, 00-656 WARSAW, POLAND

E-mail address: rmohammadpour@impan.pl