

A NEW APPROACH TO A THEOREM OF ENG

HASAN ARSLAN*

*Department of Mathematics, Faculty of Science, Erciyes University, 38039, Kayseri, Turkey

E-mail: hasanarslan@erciyes.edu.tr

ABSTRACT. The main aim of this work is to give a case-free algebraic proof for a theorem of Eng on the Poincaré polynomial of parabolic quotients of finite Coxeter groups evaluated at -1 .

Mathematics Subject Classification (2010): 20F55.

Key words: Coxeter groups, Descent algebras, Poincaré polynomial, the longest element, $q = -1$ phenomenon.

1 Introduction

Let (W, S) be a finite Coxeter system. For any subset J of S , if the simple reflections in J generate W_J , then W_J is called a *standard parabolic subgroup* of W . Denote by X_J the set of distinguished left coset representatives W_J in W and let w_0 be the longest element of W . We use the same notations P_J and N_J as in [10] for the number of elements in X_J of even and odd length, respectively and let $D_J = P_J - N_J$. In [4], Theorem 1], O. Eng proved the following theorem about -1 phenomenon.

Theorem 1.1. *For any $J \subseteq S$, we have*

$$\sum_{w \in X_J} (-1)^{l(w)} = |\{w \in X_J : w_0 w W_J = w W_J\}|, \quad (1.1)$$

where l denotes the length function on W .

In [10], L. Tan showed the relation $D_J = \sum_{w \in X_J} (-1)^{l(w)}$. O. Eng proved this theorem by using case by case techniques. In [7], V. Reiner gave a case-free geometric proof of Theorem 1.1. Then in [8], V. Reiner, D. Stanton, D. White presented the first case-free algebraic proof of the theorem above. To give another proof of Theorem 1.1 algebraically, we use a different method than that of V. Reiner, D. Stanton, D. White. Our approach depends more on the structure of

the descent algebra of a finite Coxeter group introduced by L. Solomon in [9]. Our proof is new and avoids case by case considerations. Set $x_J = \sum_{w \in X_J} w$ for any subset J of S . Then $\{x_J \mid J \subseteq S\}$ forms a basis for a subalgebra of the group algebra $\mathbb{Q}W$ called *the descent algebra* of W . We denote by $\sum(W)$ the descent algebra corresponding to W . The ascent set of w is defined by

$$\mathcal{R}(w) = \{s \in S : l(ws) > l(w)\}.$$

For any subset I of S , put $Y_I = \{w \in W : \mathcal{R}(w) = I\}$. We consider the element $y_I = \sum_{w \in Y_I} w$ in $\mathbb{Q}W$. Then

$$x_I = \sum_{I \subseteq J} y_J$$

and by Möbius inversion formula

$$y_I = \sum_{I \subseteq J} (-1)^{|J-I|} x_J.$$

Thus the set $\{y_I : I \subseteq S\}$ is a basis of $\sum(W)$, see [9]. In [9], L. Solomon also defined an algebra map from $\sum(W)$ to $\mathbb{Q}\text{Irr}W$ as follows:

$$\Phi : \sum(W) \rightarrow \mathbb{Q}\text{Irr}(W), x_J \mapsto \text{Ind}_{W_J}^W 1_{W_J},$$

where $\mathbb{Q}\text{Irr}W$, 1_{W_J} and $\text{Ind}_{W_J}^W 1_{W_J}$ denote the algebra generated by irreducible characters of W , the trivial character of W_J and the permutation character of W_J in W , respectively. Taking into account the sign character of W , which is defined as $\varepsilon : W \rightarrow \mathbb{N}$, $\varepsilon(w) = (-1)^{l(w)}$ (see [6]), it is well-known from [9] that $y_\emptyset = w_0 \in \sum(W)$ and

$$\Phi(w_0) = \varepsilon. \tag{1.2}$$

The sign character ε of W is actually irreducible and equals to the Steinberg character of W , which is given by the formula $St_W = \sum_{J \subseteq S} (-1)^{|J|} \Phi(x_J)$ due to [3]. In [[2], Main Theorem], D. Blessenohl, C. Hohlweg, M. Schocker showed the symmetry property for the descent algebra $\sum(W)$, that is,

$$\Phi(x)(y) = \Phi(y)(x) \tag{1.3}$$

for all $x, y \in \sum(W)$.

Let $I, J \subseteq S$. We write $I \sim_W J$ if I and J are W -conjugate, that is, there is a $w \in W$ such that $I = wJw^{-1}$.

2 Proof of Theorem 1.1

When we extend linearly the sign character ε of W to the group algebra $\mathbb{Q}W$ and use the equation (1.2), (1.3), then we conclude that

$$\begin{aligned} \sum_{w \in X_J} (-1)^{l(w)} &= \sum_{w \in X_J} \varepsilon(w) = \varepsilon\left(\sum_{w \in X_J} w\right) \\ &= \varepsilon(x_J) = \Phi(w_0)(x_J) = \Phi(x_J)(w_0) \\ &= \text{Ind}_{W_J}^W 1_{W_J}(w_0) = |\{w \in X_J : w_0 w W_J = w W_J\}|. \end{aligned}$$

Therefore, we obtain the equation (1.1) and so we complete a case-free algebraic proof of the Theorem 1.1. Consequently, we can write $D_J = \Phi(x_J)(w_0)$.

Corollary 2.1. *Let $I, J \subseteq S$. If $I \sim_W J$, then we have $D_I = D_J$.*

Proof. The radical of $\sum(W)$ is $\text{Ker } \Phi = \langle x_I - x_J : I \sim_W J \rangle$ from [9]. Since Φ is an algebra morphism, then we get $\Phi(x_I) = \Phi(x_J)$. By considering the proof of Theorem 1.1 given above, one has

$$D_I = \Phi(x_I)(w_0) = \Phi(x_J)(w_0) = D_J.$$

□

Another perspective for the proof of Corollary 2.1 is that when I conjugate to J under W , the corresponding sets of the distinguished coset representatives X_I and X_J are W -pointwise conjugate to each other due to [5]. Therefore, we have $D_I = D_J$.

Corollary 2.2. *For any finite Coxeter system (W, S) , we have*

$$\sum_{J \subseteq S} (-1)^{|J|} D_J = (-1)^{|T|},$$

where T denotes the set of reflections of W , that is, $T = \bigcup_{w \in W} wSw^{-1}$.

Proof. To prove Corollary 2.2 observe that image of the longest element w_0 under the sign character ε of W is $\varepsilon(w_0) = (-1)^{l(w_0)} = (-1)^{|T|}$. Since $w_0 = \sum_{J \subseteq S} (-1)^{|J|} x_J$, we have

$$\varepsilon(w_0) = \sum_{J \subseteq S} (-1)^{|J|} \varepsilon(x_J) = \sum_{J \subseteq S} (-1)^{|J|} \Phi(x_J)(w_0) = \sum_{J \subseteq S} (-1)^{|J|} D_J,$$

using the fact that $\Phi(x_J)(w_0) = D_J$.

□

One can observe that Corollary 2.2 is actually a special case of Proposition 1.11 given in [6].

Example. We consider the Coxeter system (W_n, S_n) of type B_n . In [10], Theorem B], L. Tan proved that $D_J = 0$ for every proper parabolic subgroup of W_n . This statement is also seen from the fact that there is no $w \in X_J$ such that $ww_0w^{-1} \in W_J$ for every proper subset J of S_n . Therefore,

$$\sum_{J \subseteq S_n} (-1)^{|J|} D_J = (-1)^n.$$

The following result due to L. Tan [10] is easily derived from Lemma 2.1 given in [1] and Theorem 1.1.

Lemma 2.3 (Tan [10]). *Suppose $I \subseteq J \subseteq S$. Then $W_J = X_I^J W_I$ and $D_{SI} = D_{SJ} D_{JI}$, where $X_I^J = X_I \cap W_J$.*

It is well-known from Lemma 2.1 in [1] that for $I \subseteq J \subseteq S$, we have $x_I = x_J x_I^J$. Since Φ is an algebra morphism, then we get

$$\Phi(x_I) = \Phi(x_J) \Phi(x_I^J).$$

Taking Theorem 1.1 into consideration we conclude that the relation $D_{SI} = D_{SJ} D_{JI}$ in the sense of [10].

Corollary 2.4. *For any finite Coxeter group W , we have*

$$\sum_{w \in W} (-1)^{l(w)} = 0.$$

Proof. Since the image of x_\emptyset under Φ gives the regular character of W and the equation (1.3), then we conclude

$$\sum_{w \in W} (-1)^{l(w)} = \varepsilon\left(\sum_{w \in W} w\right) = \varepsilon(x_\emptyset) = \phi(w_0)(x_\emptyset) = \phi(x_\emptyset)(w_0) = 0.$$

□

References

- [1] F. Bergeron, N. Bergeron, R. B. Howlett, D. E. Taylor, *A Decomposition of the Descent Algebra of a Finite Coxeter Group*, Journal of Algebraic Combinatorics, 1(1) (1992), 23–44.

- [2] D. Blessenohl, C. Hohlweg, M. Schocker, *A symmetry of the descent algebra of a finite Coxeter group*, Adv. Math., 193(2) (2005), 416–437.
- [3] Curtis, C.W., Reiner, I., *Methods of Representation Theory with Applications to Finite Groups and Orders*, Vol. II, John Wiley and Sons, 1987.
- [4] O. Eng, *Quotients of Poincaré polynomials evaluated at -1*, J. Algebraic Combin., 13(1) (2001), 29–40.
- [5] P. Fleischmann, *On pointwise conjugacy of distinguished coset representatives in Coxeter groups*, J. Group Theory, 5(3) (2002), 269–283.
- [6] J. E. Humphreys, *Reflection Groups and Coxeter Groups*, Cambridge Studies in Advanced Math., vol. 29, Cambridge University Press, 1990.
- [7] V. Reiner, *Note on a theorem of Eng*, Ann. Comb., 6(1) (2002), 117–118.
- [8] V. Reiner, D. Stanton, D. White, *The cyclic sieving phenomenon*, J. Combin. Theory Ser. A, 108(1) (2004), 17–50.
- [9] L. Solomon, *A Mackey formula in the group ring of a Coxeter group*, J. Algebra, 41(2) (1976), 255–264.
- [10] L. Tan, *On the distinguished coset representatives of the parabolic subgroups in finite Coxeter groups*, Comm. Algebra, 22(3) (1994), 1049–1061.