

Curvature-Dimension Condition Meets Gromov's n -Volumic Scalar Curvature

Jialong DENG

Mathematisches Institut, Georg-August-Universität, Göttingen, Germany

E-mail: jialong.deng@mathematik.uni-goettingen.de

Received July 29, 2020, in final form January 23, 2021; Published online February 05, 2021

<https://doi.org/10.3842/SIGMA.2021.013>

Abstract. We study the properties of the n -volumic scalar curvature in this note. Lott–Sturm–Villani's curvature-dimension condition $\text{CD}(\kappa, n)$ was showed to imply Gromov's n -volumic scalar curvature $\geq n\kappa$ under an additional n -dimensional condition and we show the stability of n -volumic scalar curvature $\geq \kappa$ with respect to smGH-convergence. Then we propose a new weighted scalar curvature on the weighted Riemannian manifold and show its properties.

Key words: curvature-dimension condition; n -volumic scalar curvature; stability; weighted scalar curvature $\text{Sc}_{\alpha, \beta}$

2020 Mathematics Subject Classification: 53C23

1 Introduction

The concept of lower bounded curvature on the metric space or the metric measure space has evolved to a rich theory due to Alexandrov's insight. The stability of Riemannian manifolds with curvature bounded below is another deriving force to extend the definition of the curvature bounded below to a broader space. However, the scalar curvature (of Riemannian metrics) bounded below was yet absent from this picture. Gromov proposed a synthetic treatment of scalar curvature bounded below, which was called the n -volumic scalar curvature bounded below, and offered some pertinent conjectures in [18, Section 26]. Motivated by the $\text{CD}(\kappa, n)$ condition, we add an n -dimension condition to the Gromov's definition and introduce the definition of $\text{Sc}_{\alpha, \beta}$ on the smooth metric measure space. Details will be given later.

Theorem 1.1. *Assume that the metric measure space (X^n, d, μ) satisfies n -dimensional condition and the curvature-dimension condition $\text{CD}(\kappa, n)$ for $\kappa \geq 0$ and $n \geq 2$, then (X^n, d, μ) satisfies $\text{Sc}^{\text{vol}_n}(X^n) \geq n\kappa$.*

Theorem 1.2. *If compact metric measure spaces (X_i^n, d_i, μ_i) with $\text{Sc}^{\text{vol}_n}(X_i^n) \geq \kappa \geq 0$ and SC -radius $r_{X_i^n} \geq R > 0$ and (X_i^n, d_i, μ_i) strongly measured Gromov–Hausdorff converge to the compact metric measure space (X^n, d, μ) with n -dimensional condition, then X^n also satisfies $\text{Sc}^{\text{vol}_n}(X^n) \geq \kappa$ and the SC -radius $r_{X^n} \geq R$.*

Theorem 1.3. *Let $(M^n, g, e^{-f} d\text{Vol}_g)$ be the closed smooth metric measure space with $\text{Sc}_{\alpha, \beta} > 0$, then we have the following conclusions:*

1. If M^n is a spin manifold, $\alpha \in \mathbb{R}$ and $\beta \geq \frac{|\alpha|^2}{4}$, then the harmonic spinors of M^n vanish.
2. If the dimension $n \geq 3$, $\alpha \in \mathbb{R}$ and $\beta \geq \frac{(n-2)|\alpha|^2}{4(n-1)}$, then there is a metric \tilde{g} conformal to g with positive scalar curvature.

3. If the dimension $n \geq 3$, $\alpha = 2$, $\beta \geq \frac{n-2}{n-1}$ and (N^{n-1}, \bar{g}) is the compact L_f -stable minimal hypersurface of $(M^n, g, e^{-f} d\text{Vol}_g)$, then there exists a PSC-metric conformal to \bar{g} on N^{n-1} , where \bar{g} is the induced metric of g on N^{n-1} .
4. Assume M^n is a spin manifold and there exists a smooth 1-contracting map $h: (M^n, g) \rightarrow (S^n, g_{\text{st}})$ of non-zero degree. If $\alpha \in \mathbb{R}$, $\beta \geq \frac{|\alpha|^2}{4}$ and $\text{Sc}_{\alpha, \beta} \geq n(n-1)$, then h is an isometry between the metrics g and g_{st} .

The paper is organized as follows. In Section 2, we introduce the notions and show that $\text{CD}(\kappa, n)$ implies $\text{Sc}^{\text{vol}_n} \geq (n-1)\kappa$. In Section 3, we show the stability of spaces with $\text{Sc}^{\text{vol}_n} \geq \kappa$. In Section 4, we present the properties of the smooth metric measure space with $\text{Sc}_{\alpha, \beta} > 0$.

2 CD meets n -volumic scalar curvature

The n -dimensional Aleksandrov space with curvature $\geq \kappa$ equipped with the volume-measure satisfies Lott–Villani–Sturm’s weak curvature-dimension condition for dimension n and curvature $(n-1)\kappa$, i.e., $\text{CD}((n-1)\kappa, n)$, was shown by Petrunin for $\kappa = 0$ (and said that for general curvature $\geq \kappa$ the result followed in a similar way) [32] and then Zhang–Zhu investigated the general case [43]. We will modify Gromov’s definition of n -volumic scalar curvature bounded below in [18, Section 26] to fill the picture, which means Lott–Sturm–Villani’s Ricci curvature ≥ 0 implies Gromov’s scalar curvature ≥ 0 .

The metric measure space (mm-space) $X = (X, d, \mu)$ means that d is the complete separable length metric on X and μ is the locally finite full support Borel measure on X equipped with its Borel σ -algebra. Say that an mm-space $X = (X, d, \mu)$ is locally volume-wise smaller (or not greater) than another such space $X' = (X', d', \mu')$ and write $X <_{\text{vol}} X'$ ($X \leq_{\text{vol}} X'$), if all ϵ -balls in X are smaller (or not greater) than the ϵ -balls in X' , $\mu(B_\epsilon(x)) < \mu'(B_\epsilon(x'))$ ($\mu(B_\epsilon(x)) \leq \mu'(B_\epsilon(x'))$), for all $x \in X, x' \in X'$ and the uniformly small ϵ which depends on X and X' .

From now on, the Riemannian 2-sphere $(S^2(\gamma), d_S, \text{vol}_S)$ is endowed with round metric such that the scalar curvature equal to $2\gamma^{-2}$, $(\mathbf{R}^{n-2}, d_E, \text{vol}_E)$ is endowed with Euclidean metric with flat scalar curvature and the product manifold $S^2(\gamma) \times \mathbf{R}^{n-2}$ is endowed with the Pythagorean product metrics $d_{S \times E} := \sqrt{d_S^2 + d_E^2}$ and the volume $\text{vol}_{S \times E} := \text{vol}_S \otimes \text{vol}_E$.

Thus, we have $S^2(\gamma) <_{\text{vol}} \mathbf{R}^2$. If $0 < \gamma_1 < \gamma_2$, then $S^2(\gamma_1) <_{\text{vol}} S^2(\gamma_2)$. Furthermore, $S^2(\gamma) \times \mathbf{R}^{n-2} <_{\text{vol}} \mathbf{R}^n$. If $0 < \gamma_1 < \gamma_2$, then $S^2(\gamma_1) \times \mathbf{R}^{n-2} <_{\text{vol}} S^2(\gamma_2) \times \mathbf{R}^{n-2}$.

Definition 2.1 (Gromov’s n -volumic scalar curvature). Gromov’s n -volumic scalar curvature of X is bounded below by 0 for $X = (X, d, \mu)$ if X is locally volume-wise not greater than \mathbf{R}^n .

Gromov’s n -volumic scalar curvature of X bounds from below by $\kappa > 0$ for $X = (X, d, \mu)$ if X is locally volume-wise smaller than $S^2(\gamma) \times \mathbf{R}^{n-2}$ for all $\gamma > \sqrt{\frac{2}{\kappa}}$, i.e., $X <_{\text{vol}} S^2(\gamma) \times \mathbf{R}^{n-2}$ and $\gamma > \sqrt{\frac{2}{\kappa}}$, where $S^2(\gamma) \times \mathbf{R}^{n-2} = (S^2(\gamma) \times \mathbf{R}^{n-2}, d_{S \times E}, \text{vol}_{S \times E})$.

The n -volumic scalar curvature is sensitive to the scaling of the measure, but the curvature condition $\text{CD}(\kappa, n)$ of Lott–Villani–Sturm [38, Definition 1.3] is invariant up to scalars of the measure only [38, Proposition 1.4(ii)]. Therefore, the n -dimensional condition needs to be put into the definition of Gromov’s n -volumic scalar curvature. In fact, the n -dimensional condition is the special case of Young’s point-wise dimension in dynamical systems [41, Theorem 4.4].

Definition 2.2 (n -dimensional condition). For given positive natural number n , the mm-space $X = (X, d, \mu)$ satisfies the n -dimensional condition if

$$\lim_{r \rightarrow 0} \frac{\mu(B_r(x))}{\text{vol}_E(B_r(\mathbf{R}^n))} = 1$$

for every $x \in X$, where $B_r(\mathbf{R}^n)$ is the closed r -ball in the Euclidean space \mathbf{R}^n and the $B_r(x)$ is the closed r -ball with the center $x \in X$.

From now on, the superscript of n in the space X^n means the mm-space (X^n, d, μ) satisfies n -dimensional condition.

Note that a closed smooth n -manifold M^n ($n \geq 3$) admits a Riemannian metric with constant negative scalar curvature and a Riemannian metric of non-negative scalar curvature which is not identically zero, then by a conformal change of the metric we get a metric of positive scalar curvature according to Kazdan–Warner theorem [25]. Furthermore, if there is a scalar-flat Riemannian metric g on M^n , but g is not Ricci-flat metric, then g can be deformed to a metric with positive scalar curvature according to Kazdan theorem [24, Theorem B] or by using Ricci-flow with an easy argument. Hence we will focus more on promoting the positive scalar curvature to positive n -volumic scalar curvature.

Definition 2.3 (n -volumic scalar curvature). Assume $X^n = (X^n, d, \mu)$ is the compact mm-space and satisfies the n -dimensional condition, we call

1. the n -volumic scalar curvature of X^n is positive, i.e., $\text{Sc}^{\text{vol}_n}(X^n) > 0$, if there exists $r_{X^n} > 0$ such that the measures of ϵ -balls in X^n are smaller than the volumes of ϵ -balls in \mathbf{R}^n for $0 < \epsilon \leq r_{X^n}$.
2. the n -volumic scalar curvature of X^n is bounded below by 0, i.e., $\text{Sc}^{\text{vol}_n}(X^n) \geq 0$, if there exists $r_{X^n} > 0$ such that the measures of ϵ -balls in X^n are not greater than the volumes of ϵ -balls in \mathbf{R}^n for $0 < \epsilon \leq r_{X^n}$.

The r_{X^n} is called scalar curvature radius (SC-radius) of X^n for $\text{Sc}^{\text{vol}_n}(X^n) \geq 0$.

3. the n -volumic scalar curvature of X^n is bounded below by $\kappa > 0$, i.e., $\text{Sc}^{\text{vol}_n}(X^n) \geq \kappa > 0$, if, for any γ with $\gamma > \sqrt{\frac{2}{\kappa}}$, there exists $r_{X^n, \gamma} > 0$ such that the measures of ϵ -balls in X^n are smaller than the volumes of ϵ -balls in $S^2(\gamma) \times \mathbf{R}^{n-2}$ for $0 < \epsilon \leq r_{X^n, \gamma}$.

We call $r_{X^n} := \inf_{\gamma > \sqrt{\frac{2}{\kappa}}} r_{X^n, \gamma}$ is the SC-radius of X^n for $\text{Sc}^{\text{vol}_n}(X^n) \geq \kappa > 0$.

In particular, we will focus on the case of $\inf_{\gamma > \sqrt{\frac{2}{\kappa}}} r_{X^n, \gamma} \neq 0$ for stability in Section 3.

If the mm-space X^n is locally compact, then the definition of the n -volumic scalar curvature bounded below only modifies the definition of the $r_{X^n, \gamma} > 0$ to a positive continuous function of X^n .

Two mm-spaces (X^n, d, μ) and (X_1^n, d_1, μ_1) are isometric if there exists a one-to-one map $f: X^n \rightarrow X_1^n$ such that $d_1(f(a), f(b)) = d(a, b)$ for a and b are in X^n and $f_*\mu = \mu_1$, where $f_*\mu$ is the push-forward measure, i.e., $f_*\mu(U) = \mu(f^{-1}(U))$ for a measurable subset $U \subset X_1^n$. If X^n satisfies $\text{Sc}^{\text{vol}_n}(X^n) \geq \kappa \geq 0$, then each mm-space (X_1^n, d_1, μ_1) that is isometric to (X^n, d, μ) also satisfies $\text{Sc}^{\text{vol}_n}(X_1^n) \geq \kappa \geq 0$.

Proposition 2.4. *Let g be a C^2 -smooth Riemannian metric on a closed oriented n -manifold M^n with induced metric measure space $(M^n, d_g, \text{dVol}_g)$, then the scalar curvature of g is positive, $\text{Sc}_g > 0$, if and only if $\text{Sc}^{\text{vol}_n}(M^n) > 0$, and $\text{Sc}_g \geq \kappa > 0$ if and only if $\text{Sc}^{\text{vol}_n}(M^n) \geq \kappa > 0$.*

Proof. For a C^2 -smooth Riemannian metric g , one has

$$\text{dVol}_g(B_r(x)) = \text{vol}_E(B_r(\mathbf{R}^n)) \left[1 - \frac{\text{Sc}_g(x)}{6(n+2)} r^2 + O(r^4) \right]$$

for $B_r(x) \subset M^n$ as $r \rightarrow 0$. Hence $(M^n, d_g, \text{dVol}_g)$ satisfies the n -dimensional condition.

If we have $\text{Sc}_g > 0$, then, since M^n is compact, there exists $r_{M^n} > 0$, so that $\text{dVol}_g(B_r(x)) < \text{vol}_E(B_r(\mathbf{R}^n))$ for all $0 < r \leq r_{M^n}$. On the other hand, if there exists $r_{M^n} > 0$ such that $\text{dVol}_g(B_r(x)) < \text{vol}_E(B_r(\mathbf{R}^n))$ for all $0 < r \leq r_{M^n}$, then Sc_g must be greater than 0.

If $\text{Sc}^{\text{vol}_n}(M^n) \geq \kappa > 0$, then $\text{Sc}_g \geq \kappa > 0$. Otherwise, assume there exist small $\epsilon > 0$ such that $\text{Sc}_g \geq \kappa - \epsilon > 0$. That means that there exists a point x_0 in M^n such that $\text{Sc}_g(x_0) = \kappa - \epsilon$, as M^n is compact and the scalar curvature is a continuous function on M^n . Thus, we can find a small r -ball $B_r(x_0)$ such that the volume of $B_r(x_0)$ is greater than the volume of the r -ball in the $S^2(\gamma) \times \mathbf{R}^{n-2}$ for $\gamma = \sqrt{\frac{2}{\kappa - \epsilon}}$, which is a contradiction.

On the other hand, $\text{Sc}_g \geq \kappa > 0$ implies $\text{Sc}^{\text{vol}_n}(M^n) \geq \kappa > 0$. Assume $\text{Sc}_g(x_1) = \kappa$ for some $x_1 \in M^n$, then there exists r_1 such that $\text{dVol}_g(B_{r_1}(x)) \leq \text{dVol}_g(B_{r_1}(x_1))$ for r_1 -balls in M^n and

$$\text{dVol}_g(B_r(x_1)) = \text{vol}_E(B_r(\mathbf{R}^n)) \left[1 - \frac{\kappa}{6(n+2)} r^2 + O(r^4) \right]$$

as $r \rightarrow 0$. Thus, for any γ with $\gamma > \sqrt{\frac{2}{\kappa}}$, there exists $r_{M^n, \gamma} > 0$ such that the measures of ϵ -balls in M^n are smaller than the volumes of ϵ -balls in $S^2(\gamma) \times \mathbf{R}^{n-2}$ for $0 < \epsilon \leq r_{M^n, \gamma}$, i.e., $\text{Sc}^{\text{vol}_n}(M^n) \geq \kappa > 0$. ■

Therefore, we have $S_{\frac{\kappa}{n-1}}^n <_{\text{vol}} S^2(\gamma) \times \mathbf{R}^{n-2}$ for all $\gamma > \sqrt{\frac{2}{n\kappa}}$. Here $S_{\frac{\kappa}{n-1}}^n$ is the Riemannian manifold S^n with constant sectional curvature $\frac{\kappa}{n-1}$.

Remark 2.5. For a closed smooth Riemannian manifold (M^n, g) , $\text{Sc}^{\text{vol}_n}(M^n) \geq 0$ implies $\text{Sc}_g \geq 0$. Otherwise, there exists a point in M^n such that the scalar curvature is negative, then the volume of small ball will be greater than the volume of the small ball in Euclidean space, which is a contradiction.

On the other hand, one can consider the case of the scalar-flat metric, i.e., $\text{Sc}_g \equiv 0$. If g is a strongly scalar-flat metric, meaning a metric with scalar curvature zero such that M^n has no metric with positive scalar curvature, then g is also Ricci flat according to Kazdan theorem above. Thus, we have

$$\text{dVol}_g(B_r(x)) = \text{vol}_E(B_r(\mathbf{R}^n)) \left[1 - \frac{\|\text{Rie}(x)\|_g^2}{120(n+2)(n+4)} r^4 + O(r^6) \right]$$

for $B_r(x) \subset M^n$ as $r \rightarrow 0$ [14, Theorem 3.3]. Here Rie is the Riemannian tensor. Therefore, if g is a not flat metric, then $M^n <_{\text{vol}} \mathbf{R}^n$. If g is a flat metric, then $M^n \leq_{\text{vol}} \mathbf{R}^n$. Thus $\text{Sc}_g \geq 0$ implies $\text{Sc}^{\text{vol}_n}(M^n) \geq 0$ for a strongly scalar-flat metric g .

However, $\text{Sc}_g \geq 0$ may not imply $\text{Sc}^{\text{vol}_n}(M^n) \geq 0$. There are a lot of scalar-flat metrics but not strongly scalar flat metrics, i.e., $\text{Sc}_g \equiv 0$ but not $\text{Ric}_g \neq 0$. For instance, the product metric on $S^2(1) \times \Sigma$, where Σ is a closed hyperbolic surface, is the scalar-flat metric, but not the Ricci-flat metric. For those metrics, we have

$$\text{dVol}_g(B_r(x)) = \text{vol}_E(B_r(\mathbf{R}^n)) \left[1 + \frac{-3\|\text{Rie}(x)\|_g^2 + 8\|\text{Ric}(x)\|_g^2}{360(n+2)(n+4)} r^4 + O(r^6) \right]$$

for $B_r(x) \subset M^n$ as $r \rightarrow 0$ [14, Theorem 3.3]. If $8\|\text{Ric}(x)\|_g^2 > -3\|\text{Rie}(x)\|_g^2$ for some point, then $\text{Sc}_g \geq 0$ does not imply $\text{Sc}^{\text{vol}_n}(M^n) \geq 0$.

Theorem 2.6. Assume that the mm-space (X^n, d, μ) satisfies n -dimensional condition and the curvature-dimension condition $\text{CD}(\kappa, n)$ for $\kappa \geq 0$ and $n \geq 2$, then (X^n, d, μ) satisfies $\text{Sc}^{\text{vol}_n}(X^n) \geq n\kappa$.

Proof. In fact, one only needs the generalized Bishop–Gromov volume growth inequality, which is implied by the curvature-dimension of X^n [38, Theorem 2.3].

(i) If $\kappa = 0$, then

$$\frac{\mu(B_r(x))}{\mu(B_R(x))} \geq \left(\frac{r}{R}\right)^n$$

for all $0 < r < R$. That is

$$\frac{\mu(B_r(x))}{\text{vol}_E(B_r(\mathbf{R}^n))} = \frac{\mu(B_r(x))}{\alpha(n)r^n} \geq \frac{\mu(B_R(x))}{\alpha(n)R^n} = \frac{\mu(B_R(x))}{\text{vol}_E(B_R(\mathbf{R}^n))},$$

where $\alpha(n) = \frac{\text{vol}_E(B_r(\mathbf{R}^n))}{r^n}$. Combining the n -dimensional condition,

$$\lim_{r \rightarrow 0} \frac{\mu(B_r(x))}{\text{vol}_E(B_r(\mathbf{R}^n))} = 1,$$

that implies $\text{Sc}^{\text{vol}_n}(X) \geq 0$.

(ii) If $\kappa > 0$, then

$$\frac{\mu(B_r(x))}{\mu(B_R(x))} \geq \frac{\int_0^r [\sin(t\sqrt{\frac{\kappa}{(n-1)}})]^{n-1} dt}{\int_0^R [\sin(t\sqrt{\frac{\kappa}{(n-1)}})]^{n-1} dt}$$

for all $0 < r \leq R \leq \pi\sqrt{\frac{(n-1)}{\kappa}}$.

Since the scalar curvature of the product manifold $S^2(\gamma) \times \mathbf{R}^{n-2}$ is $n\kappa$, where $\gamma = \sqrt{\frac{2}{n\kappa}}$, then there exists $C_1, C_2 > 0$ such that

$$1 - \frac{n\kappa}{6(n+2)}r_1^2 - C_2r_1^4 \leq \text{vol}_{S \times E}(\widetilde{B_{r_1}(y)}) := \frac{\text{vol}_{S \times E}(B_{r_1}(y))}{\text{vol}_E(B_{r_1}(\mathbf{R}^n))} \leq 1 - \frac{n\kappa}{6(n+2)}r_1^2 + C_2r_1^4,$$

for $y \in S^2(\gamma) \times \mathbf{R}^{n-2}$ and $r_1 \leq C_1$, where C_1, C_2 are decided by the product manifold $S^2(\gamma) \times \mathbf{R}^{n-2}$.

Let

$$\mu(\widetilde{B_r(x)}) := \frac{\mu(B_r(x))}{\text{vol}_E(B_r(\mathbf{R}^n))}$$

and

$$f(r) := \frac{\int_0^r [\sin(t\sqrt{\frac{\kappa}{(n-1)}})]^{n-1} dt}{\text{vol}_E(B_r(\mathbf{R}^n))},$$

then the generalized Bishop–Gromov inequality can be re-formulated as

$$\mu(\widetilde{B_R(x)}) \leq \mu(\widetilde{B_r(x)}) \frac{f(R)}{f(r)}$$

for all $0 < r < R \leq \pi\sqrt{\frac{(n-1)}{\kappa}}$. The asymptotic expansion of $f(r)$ is

$$f(r) = \frac{\frac{1}{n}r^n \left[\frac{\kappa}{(n-1)}\right]^{\frac{(n-1)}{2}} - \frac{(n-1)}{6(n+2)}r^{n+2} \left[\frac{\kappa}{(n-1)}\right]^{\frac{n+1}{2}} + O(r^{n+4})}{\text{vol}_E(B_r(\mathbf{R}^n))}$$

as $r \rightarrow 0$. Thus, the asymptotic expansion of $\frac{f(R)}{f(r)}$ is

$$\frac{f(R)}{f(r)} = \frac{1 - \frac{n\kappa}{6(n+2)}R^2 + O(R^4)}{1 - \frac{n\kappa}{6(n+2)}r^2 + O(r^4)}$$

as $R \rightarrow 0, r \rightarrow 0$. The n -dimensional condition, $\lim_{r \rightarrow 0} \mu(\widetilde{B_r(x)}) = 1$, implies that

$$\mu(\widetilde{B_R(x)}) \leq 1 - \frac{n\kappa}{6(n+2)}R^2 + O(R^4)$$

as $R \rightarrow 0$. Therefore, for any κ' with $0 < \kappa' < \kappa$, there exists $\epsilon_{\kappa'} > 0$ such that for any $0 < R \leq \epsilon_{\kappa'}$, we have

$$\mu(\widetilde{B_R(x)}) < \text{vol}_{S \times E}(\widetilde{B_R(y)}),$$

where $\text{vol}_{S \times E}(\widetilde{B_R(y)}) = \frac{\text{vol}_{S \times E}(B_R(y))}{\text{vol}_E(B_R(\mathbf{R}^n))}$ is defined as before, the balls $B_R(y)$ are in $S^2(\gamma) \times \mathbf{R}^{n-2}$ and $\gamma = \sqrt{\frac{2}{n\kappa'}}$. That is $X^n <_{\text{vol}} S^2(\gamma) \times \mathbf{R}^{n-2}$, for all $\gamma > \sqrt{\frac{2}{n\kappa}}$, i.e., $\text{Sc}^{\text{vol}_n}(X^n) \geq n\kappa$.

In fact, one has the classical Bishop inequality by adding the n -dimensional condition to the generalized Bishop–Gromov volume growth inequality. It means that

- if $\kappa = 0$, $\mu(B_R(x)) \leq \text{vol}_E(B_R(\mathbf{R}^n))$ for all $R > 0$,
- if $\kappa > 0$, $\mu(B_R(x)) \leq \text{vol}_{S^n}(B_R(S_{\frac{\kappa}{n-1}}^n))$ for $0 < R \leq \pi\sqrt{\frac{(n-1)}{\kappa}}$.

In other words, if $\kappa = 0$, $X^n \leq_{\text{vol}} \mathbf{R}^n$. If $\kappa > 0$, $X^n \leq_{\text{vol}} S_{\frac{\kappa}{n-1}}^n$. We have $S_{\frac{\kappa}{n-1}}^n <_{\text{vol}} S^2(\gamma) \times \mathbf{R}^{n-2}$ for all $\gamma > \sqrt{\frac{2}{n\kappa}}$. Then $X^n <_{\text{vol}} S^2(\gamma) \times \mathbf{R}^{n-2}$ for all $\gamma > \sqrt{\frac{2}{n\kappa}}$.

Thus, we also get $\text{Sc}^{\text{vol}_n}(X^n) \geq n\kappa$. ■

Remark 2.7. Hence the mm-space (X^n, d, μ) with $\text{Sc}^{\text{vol}_n}(X^n) \geq n\kappa$ includes the mm-spaces that satisfies n -dimensional condition and the generalized Bishop–Gromov volume growth inequality as stated in the proof, e.g., the mm-spaces with the Riemannian curvature condition $\text{RCD}(\kappa, n)$ [2] or with the measure concentration property $\text{MCP}(\kappa, n)$ [29].

Question 2.8. Let $\text{Al}^n(1)$ be an orientable compact n -dimensional Aleksandrov space with curvature ≥ 1 , then do all continuous maps ϕ from $\text{Al}^n(1)$ to the sphere S^n with standard metric of non-zero degree satisfy $\text{Lip}(\phi) \geq C(n)$? Here $\text{Lip}(\phi)$ is the Lipschitz constant of ϕ , ϕ maps the boundary of $\text{Al}^n(1)$ to a point in S^n and $C(n)$ is a constant depending only on the dimension n .

Question 2.9. Assume the compact mm-space (X, d, μ) satisfies the curvature-dimension condition $\text{CD}(n-1, n)$, n -dimensional condition and the covering dimension is also n , then do all continuous maps ϕ from (X, d, μ) to the sphere S^n with standard metric, where ϕ is non-trivial in the homotopy class of maps, satisfy $\text{Lip}(\phi) \geq C_1(n)$, where $C_1(n)$ is a constant depending only on n ?

Remark 2.10. The questions above are inspired by Gromov's spherical Lipschitz bounded theorem in [19, Section 3] and the results above. The finite covering dimension is equal to the cohomological dimension over integer ring \mathbb{Z} for the compact metric space according to the Alexandrov theorem. The best constant of $C(n)$ and $C_1(n)$ would be 1 if both questions have positive answers.

Proposition 2.11 (quadratic scaling). *Assume the compact mm-space (X^n, d, μ) satisfies $\text{Sc}^{\text{vol}_n}(X^n) \geq \kappa > 0$, then $\text{Sc}^{\text{vol}_n}(\lambda X^n) \geq \lambda^{-2}\kappa > 0$ and $r_{\lambda X^n} = \lambda r_{X^n}$ for all $\lambda > 0$, where $\lambda X^n := (X^n, \lambda \cdot d, \lambda^n \cdot \mu)$.*

Proof. First, we will show that the n -dimensional condition is stable under scaling. Let $d' := \lambda \cdot d$, $\mu' := \lambda^n \cdot \mu$, $B'_r(x)$ be an r -ball in the (X^n, d') , and $B_r(x)$ be an r -ball in the (X^n, d) , then $B'_r(x) = B_{\frac{r}{\lambda}}(x)$ as the subset in the X^n . One has

$$\lim_{r \rightarrow 0} \frac{\mu'(B'_r(x))}{\text{vol}_E(B_r(\mathbf{R}^n))} = \lim_{r \rightarrow 0} \frac{\mu'(B_{\frac{r}{\lambda}}(x))}{\text{vol}_E(B_r(\mathbf{R}^n))} = \lim_{r \rightarrow 0} \frac{\lambda^n \cdot \mu(B_{\frac{r}{\lambda}}(x))}{\lambda^n \cdot \text{vol}_E(B_{\frac{r}{\lambda}}(\mathbf{R}^n))} = 1,$$

then λX^n satisfies the n -dimensional condition.

Since $\lambda \cdot (S^2(\gamma) \times \mathbf{R}^{n-2}) = \lambda \cdot S^2(\gamma) \times \lambda \cdot \mathbf{R}^{n-2} = S^2(\lambda\gamma) \times \lambda \cdot \mathbf{R}^{n-2}$, we have

$$\lambda \cdot X^n <_{\text{vol}} \lambda \cdot (S^2(\gamma) \times \mathbf{R}^{n-2}) = S^2(\lambda\gamma) \times \lambda \cdot \mathbf{R}^{n-2}$$

for all $\lambda\gamma > \sqrt{\frac{2}{\lambda\kappa}}$ and $0 < \epsilon \leq \lambda r_{X^n}$. That means $\text{Sc}^{\text{vol}_n}(\lambda X^n) \geq \lambda^{-2}\kappa > 0$ and $r_{\lambda X^n} = \lambda r_{X^n}$. ■

We also have $\text{Sc}^{\text{vol}_n}(\lambda X^n) \geq 0$ (> 0), if $\text{Sc}^{\text{vol}_n}(X^n) \geq 0$ (> 0).

Remark 2.12. Since the n -dimensional condition and definition of n -volumic scalar curvature is locally defined, we have the following construction.

- Global to local: Let the locally compact mm-space (X^n, d, μ) satisfy $\text{Sc}^{\text{vol}_n}(X^n) \geq \kappa \geq 0$ and $Y^n \subset X$ be an open subset. Then, if (Y^n, d_Y) is a complete length space, $(Y^n, d_Y, \mu_{\lfloor Y})$ satisfies $\text{Sc}^{\text{vol}_n}(Y^n) \geq \kappa \geq 0$ and $r_{Y^n} = r_{X^n}$. Where d_Y is the induced metric of d and $\mu_{\lfloor Y}$ is the restriction operator, namely, $\mu_{\lfloor Y}(A) := \mu(Y^n \cap A)$ for $A \subset X^n$.
- Local to global: Let $\{Y_i^n\}_{i \in I}$ be a finite open cover of a locally compact mm-space (X^n, d, μ) . Assume that (Y_i^n, d_{Y_i}) is a complete length space and $(Y_i^n, d_{Y_i}, \mu_{\lfloor Y_i})$ satisfies $\text{Sc}^{\text{vol}_n}(Y_i^n) \geq \kappa \geq 0$, then (X^n, d, μ) satisfies $\text{Sc}^{\text{vol}_n}(X^n) \geq \kappa \geq 0$ and r_{X^n} can be chosen as a partition of unity of the functions $\{r_{Y_i^n}\}_{i \in I}$.

Question 2.13. Assume that $\text{Sc}^{\text{vol}_{n_1}}(X_1^{n_1}) \geq \kappa_1 (\geq 0)$ for the compact mm-space $(X_1^{n_1}, d_1, \mu_1)$ and $\text{Sc}^{\text{vol}_{n_2}}(X_2^{n_2}) \geq \kappa_2 (\geq 0)$ for the compact mm-space $(X_2^{n_2}, d_2, \mu_2)$, then do we have

$$\text{Sc}^{\text{vol}_{n_1+n_2}}(X_1^{n_1} \times X_2^{n_2}) \geq \kappa_1 + \kappa_2, \quad r_{X_1^{n_1} \times X_2^{n_2}} = \min\{r_{X_1^{n_1}}, r_{X_2^{n_2}}\}$$

for $(X_1^{n_1} \times X_2^{n_2}, d_3, \mu_3)$? Here $X_1^{n_1} \times X_2^{n_2}$ is endowed with the measure $\mu_3 := \mu_1 \otimes \mu_2$ and with the Pythagorean product metric $d_3 := \sqrt{d_1^2 + d_2^2}$.

3 smGH-convergence

Let $\{\mu_n\}_{n \in \mathbf{N}}$ and μ be Borel measures on the space X , then the sequence $\{\mu_n\}_{n \in \mathbf{N}}$ is said to converge *strongly* (also called setwise convergence in other literature) to a limit μ if $\lim_{n \rightarrow \infty} \mu_n(\mathcal{A}) = \mu(\mathcal{A})$ for every \mathcal{A} in the Borel σ -algebra.

A map $f: X \rightarrow Y$ is called an ϵ -isometry between compact metric spaces X and Y , if $|d_X(a, b) - d_Y(f(a), f(b))| \leq \epsilon$ for all $a, b \in X$ and it is almost surjective, i.e., for every $y \in Y$, there exists an $x \in X$ such that $d_Y(f(x), y) \leq \epsilon$.

In fact, if f is an ϵ -isometry $X \rightarrow Y$, then there is a (4ϵ) -isometry $f': Y \rightarrow X$ such that for all $x \in X$, $y \in Y$, $d_X(f' \circ f(x), x) \leq 3\epsilon$, $d_Y(f \circ f'(y), y) \leq \epsilon$.

Definition 3.1 (smGH-convergence). Let $(X_i, d_i, \mu_i)_{i \in \mathbf{N}}$ and (X, d, μ) be compact mm-spaces. X_i converges to X in the strongly measured Gromov–Hausdorff topology (smGH-convergence) if there are measurable ϵ_i -isometries $f_i: X_i \rightarrow X$ such that $\epsilon_i \rightarrow 0$ and $f_{i*}\mu_i \rightarrow \mu$ in the strong topology of measures as $i \rightarrow \infty$.

If the spaces $(X_i, d_i, \mu_i, p_i)_{i \in \mathbf{N}}$ and (X, d, μ, p) are locally compact pointed mm-spaces, it is said that X_i converges to X in the pointed strongly measured Gromov–Hausdorff topology (psmGH-convergence) if there are sequences $r_i \rightarrow \infty$, $\epsilon_i \rightarrow 0$, and measurable pointed ϵ_i -isometries $f_i: B_{r_i}(p_i) \rightarrow B_{r_i}(p)$, such that $f_{i*}\mu_i \rightarrow \mu$, where the convergence is strong convergence.

Remark 3.2. Let $(X_i, d_i, \mu_i)_{i \in \mathbf{N}}$ converge to (X, d, μ) in the measured Gromov–Hausdorff topology, then there are measurable ϵ_i -isometries $f_i: X_i \rightarrow X$ such that $f_{i*}\mu_i$ weakly converges to μ . If there is a Borel measure ν on X such that $\sup_i f_{i*}\mu_i \leq \nu$, i.e., $\sup_i f_{i*}\mu_i(\mathcal{A}) \leq \nu(\mathcal{A})$ for every \mathcal{A} in the Borel σ -algebra on X , then X_i smGH-converges to X (see [26, Lemma 4.1]).

Remark 3.3. The n -dimensional condition is not preserved by the measured Gromov–Hausdorff convergence as the following example shows. Let $\{a_i S^2 := (S^2, a_i d_S)\}$ ($a_i \in (0, 1)$) be a sequence of space, then the limit of $a_i S^2$ under the measured Gromov–Hausdorff convergence is a point when a_i goes to 0. The limit exists as the Ricci curvature of $a_i S^2$ is bounded below by 1.

Remark 3.4. The n -dimensional condition is not preserved by the smGH-convergence since the limits of $\lim_{r \rightarrow 0} \lim_{i \rightarrow \infty} \frac{\mu_i(B_r(x))}{\text{vol}_E(B_r(\mathbf{R}^n))}$ may not be commutative for some mm-spaces (X, d, μ_i) . Assume the total variation distance of the measures goes to 0 as $i \rightarrow \infty$, i.e.,

$$d_{TV}(\mu_i, \mu) := \sup_{\mathcal{A}} |\mu_i(\mathcal{A}) - \mu(\mathcal{A})| \rightarrow 0,$$

where \mathcal{A} runs over the Borel σ -algebra of X , then the limits are commutative.

One can also define the *total variation Gromov–Hausdorff convergence* (tvGH-convergence) for mm-spaces by replacing the strong topology with the topology induced by the total variation distance in definition of smGH-convergence. Then tvGH-convergence implies smGH-convergence and the n -dimensional condition is preserved by tvGH-convergence.

Theorem 3.5 (stability). *If compact mm-spaces (X_i^n, d_i, μ_i) with $\text{Sc}^{\text{vol}_n}(X_i^n) \geq \kappa \geq 0$, SC-radius $r_{X_i^n} \geq R > 0$, and (X_i^n, d_i, μ_i) smGH-converge to the compact mm-space (X^n, d, μ) with n -dimensional condition, then X^n also satisfies $\text{Sc}^{\text{vol}_n}(X^n) \geq \kappa$ and the SC-radius $r_{X^n} \geq R$.*

Proof. Fix an $x \in X^n$ and let $B_r(x)$ be the small r -ball on X^n where $r < R$, then there exists $x_i \in X_i^n$ such that $f_i^{-1}(B_r(x)) \subset B_{r+4\epsilon_i}(x_i)$ where $B_{r+4\epsilon_i}(x_i) \subset X_i^n$ and $r + 4\epsilon_i \leq R$. Thus, $f_{i*}\mu_i(B_r(x)) \leq \mu_i(B_{r+4\epsilon_i}(x_i))$.

- For $\kappa = 0$, since $\text{Sc}^{\text{vol}_n}(X_i^n) \geq 0$ and SC-radius $\geq R > 0$, then $\mu_i(B_r(x_i)) \leq \text{vol}_E(B_r(\mathbf{R}^n))$ for all $0 < r \leq R$ and all i . Therefore, $f_{i*}\mu_i(B_r(x)) < \mu_i(B_{r+4\epsilon_i}(x_i)) \leq \text{vol}_E(B_{r+4\epsilon_i}(\mathbf{R}^n))$ for $r + 4\epsilon_i \leq R$. Since ϵ_i that is not related to r can be arbitrarily small, then $\mu(B_r(x)) \leq \text{vol}_E(B_r(\mathbf{R}^n))$.
- For $\kappa > 0$, we have $\mu_i(B_r(x_i)) < \text{vol}_{S \times E}(B_r(S^2(\gamma) \times \mathbf{R}^{n-2}))$ for all $0 < r \leq R$, all i , and $\gamma > \sqrt{\frac{2}{\kappa}}$. Thus, $f_{i*}\mu_i(B_r(x)) < \mu_i(B_{r+4\epsilon_i}(x_i)) < \text{vol}_{S \times E}(B_{r+4\epsilon_i}(S^2(\gamma) \times \mathbf{R}^{n-2}))$ for $r + 4\epsilon_i \leq R$. Since ϵ_i that is not related to r can be arbitrarily small, then $\mu(B_r(x)) \leq \text{vol}_{S \times E}(B_r(S^2(\gamma) \times \mathbf{R}^{n-2}))$ for $\gamma > \sqrt{\frac{2}{\kappa}}$. Thus, $\mu(B_r(x)) < \text{vol}_{S \times E}(B_r(S^2(\sqrt{\frac{2}{\kappa+\epsilon'}}) \times \mathbf{R}^{n-2}))$, where $0 < \epsilon'$ is independence on r and ϵ' can as small as we want. Therefore, we have $\text{Sc}^{\text{vol}_n}(X^n) \geq \kappa$. ■

Definition 3.6 (tangent space). The mm-space (Y, d_Y, μ_Y, o) is a tangent space of (X^n, d, μ) at $p \in X^n$ if there exists a sequence $\lambda_i \rightarrow \infty$ such that $(X^n, \lambda_i \cdot d, \lambda_i^n \cdot \mu, p)$ psmGH-converges to (Y, d_Y, μ_Y, o) as $\lambda_i \rightarrow \infty$.

Therefore, (Y, d_Y, μ_Y, o) also satisfies the n -dimensional condition and can be written as Y^n .

Corollary 3.7. Assume the compact mm-space (X^n, d, μ) with $\text{Sc}^{\text{vol}_n}(X^n) \geq \kappa \geq 0$ and the tangent space (Y^n, d_Y, μ_Y, o) of X^n exists at the point p , then (Y^n, d_Y, μ_Y, o) satisfies $\text{Sc}^{\text{vol}_n}(Y^n) \geq 0$ and the SC-radius $\geq r_{X^n}$.

Proof. Since the n -volumic scalar curvature has the quadratic scaling property, i.e., $\text{Sc}^{\text{vol}_n}(\lambda X^n) \geq \lambda^{-2}\kappa \geq 0$ and $r_{\lambda X^n} = \lambda r_{X^n}$ for all $\lambda > 0$, where $\lambda X^n := (X^n, \lambda \cdot d, \lambda^n \cdot \mu)$, then $\text{Sc}^{\text{vol}_n}(Y^n) \geq 0$ is implied by the stability theorem. ■

The mm-spaces with $\text{Sc}^{\text{vol}_n} \geq 0$ includes some of the Finsler manifolds, for instance, \mathbf{R}^n equipped with any norm and with the Lebesgue measure satisfies $\text{Sc}^{\text{vol}_n} \geq 0$ and any smooth compact Finsler manifold is a $\text{CD}(\kappa, n)$ space for appropriate finite κ and n [30]. It is well-known that Gigli's infinitesimally Hilbertian [12] can be seen as the Riemannian condition in $\text{RCD}(\kappa, n)$ space. Thus, infinitesimally Hilbertian can also be used as a Riemannian condition in the mm-spaces with $\text{Sc}^{\text{vol}_n} \geq 0$.

Definition 3.8 ($\text{RSC}(\kappa, n)$ space). The compact mm-space (X^n, d, μ) with the n -dimensional condition is a Riemannian n -volumic scalar curvature $\geq \kappa$ space ($\text{RSC}(\kappa, n)$ space) if it is infinitesimally Hilbertian and satisfies the $\text{Sc}^{\text{vol}_n}(X^n) \geq \kappa \geq 0$.

Note that any finite-dimensional Alexandrov spaces with curvature bounded below are infinitesimally Hilbertian. Then

$$\text{Al}^n(\kappa) \Rightarrow \text{RCD}((n-1)\kappa, n) \Rightarrow \text{RSC}((n(n-1)\kappa, n)$$

on (X^n, d, \mathcal{H}^n) , where the measure \mathcal{H}^n is the n -dimensional Hausdorff measure that satisfies the n -dimensional condition.

Question 3.9. Are $\text{RSC}(\kappa, n)$ spaces stable under tvGH-convergence?

Remark 3.10 (convergence of compact mm-spaces). For the compact metric measure spaces with probability measures, one can consider mGH-convergence, Gromov–Prokhorov convergence, Gromov–Hausdorff–Prokhorov convergence, Gromov–Wasserstein convergence, Gromov–Hausdorff–Wasserstein convergence, Gromov's \square -convergence, Sturm's \mathbb{D} -convergence [39, Section 27], and Gromov–Hausdorff-vague convergence [3]. smGH-convergence implies those convergences for compact metric measure spaces with probability measures, since the measures converge strongly in smGH-convergence and converge weakly in other situations.

Note that mm-spaces with infinitesimally Hilbertian are not stable under mGH-convergence [12]. It is not clear if the infinitesimally Hilbertian are preserved under smGH-convergence or tvGH-convergence.

4 Smooth mm-space with $\text{Sc}_{\alpha, \beta} > 0$

Let the smooth metric measure space $(M^n, g, e^{-f} d\text{Vol}_g)$ (also known as the weighted Riemannian manifold in some references), where f is a C^2 -function on M^n , g is a C^2 -Riemannian metric and $n \geq 2$, satisfy the curvature-dimension condition $\text{CD}(\kappa, n)$ for $\kappa \geq 0$, then M^n also satisfies $\text{Sc}^{\text{vol}_n}(M^n) \geq n\kappa$.

Motivated by the importance of the Ricci Bakry-Emery curvature, i.e.,

$$\text{Ric}_f^M = \text{Ricc} + \text{Hess}(f),$$

the weighted sectional curvature of smooth mm-space was proposed and discussed in [40]. On the other hand, Perelman defined and used the P-scalar curvature in his \mathcal{F} -functional in [31, Section 1]. Inspired by the P-scalar curvature, i.e., $\text{Sc}_g + 2 \Delta_g f - \|\nabla_g f\|_g^2$, we propose another scalar curvature on the smooth mm-space.

Definition 4.1 (weighted scalar curvature $\text{Sc}_{\alpha,\beta}$). The weighted scalar curvature $\text{Sc}_{\alpha,\beta}$ on the smooth mm-space $(M^n, g, e^{-f} \text{dVol}_g)$ is defined by

$$\text{Sc}_{\alpha,\beta} := \text{Sc}_g + \alpha \Delta_g f - \beta \|\nabla_g f\|_g^2.$$

Note that the Laplacian Δ_g here is the trace of the Hessian and $\text{Sc}^{\text{vol}_n}(M^n) \geq \kappa \geq 0$ is equivalent to $\text{Sc}_{\alpha,\beta} \geq \kappa \geq 0$ for $\alpha = 3$ and $\beta = 3$ (see [33, Theorem 8] or the proof of Corollary 4.10 below).

Example 4.2.

1. For $\alpha = \frac{2(n-1)}{n}$ and $\beta = \frac{(n-1)(n-2)}{n^2}$, the $\text{Sc}_{\frac{2(n-1)}{n}, \frac{(n-1)(n-2)}{n^2}}$ is the Chang–Gursky–Yang’s conformally invariant scalar curvature for the smooth mm-space [7]. That means for a C^2 -smooth function w on M^n , one has

$$\text{Sc}_{\frac{2(n-1)}{n}, \frac{(n-1)(n-2)}{n^2}}(e^{2w}g) = e^{-2w} \text{Sc}_{\frac{2(n-1)}{n}, \frac{(n-1)(n-2)}{n^2}}(g).$$

2. For $\alpha = 2$ and $\beta = \frac{m+1}{m}$, where $m \in \mathbb{N} \cup \{0, \infty\}$, the $\text{Sc}_{2, \frac{m+1}{m}}$ is Case’s weighted scalar curvature and Case also defined and studied the weighted Yamabe constants in [6]. Case’s weighted scalar curvature is the classical scalar curvature if $m = 0$. If $m = \infty$, then it is Perelman’s P -scalar curvature.

Note that the results in this paper are new for those examples.

4.1 Spin manifold and $\text{Sc}_{\alpha,\beta} > 0$

For an orientable closed surface with density $(\Sigma, g, e^{-f} \text{dVol}_g)$ with $\text{Sc}_{\alpha,\beta} > 0$ and $\beta \geq 0$, then the inequality,

$$0 < \int_{\Sigma} \text{Sc}_{\alpha,\beta} \text{dVol}_g = \int_{\Sigma} (\text{Sc}_g + \alpha \Delta_g f - \beta \|\nabla_g f\|_g^2) \text{dVol}_g = 4\pi\chi(\Sigma) - \beta \int_{\Sigma} \|\nabla_g f\|_g^2 \text{dVol}_g,$$

implies that $\chi(\Sigma) > 0$. Thus, Σ is a 2-sphere.

The following proposition of vanishing harmonic spinors is owed to Perelman essentially and the proof is borrowed from [1, Proposition 1].

Proposition 4.3 (vanishing harmonic spinors). *Assume the smooth mm-space $(M^n, g, e^{-f} \text{dVol}_g)$ is closed and spin. If $\alpha \in \mathbb{R}$, $\beta \geq \frac{|\alpha|^2}{4}$ and $\text{Sc}_{\alpha,\beta} > 0$, then the harmonic spinor of M^n vanishes.*

Proof. Let ψ be a harmonic spinor, though the Schrödinger–Lichnerowicz–Weitzenboeck formula

$$\mathbb{D}^2 = \nabla^* \nabla + \frac{1}{4} \text{Sc}_g,$$

one has

$$\begin{aligned} 0 &= \int_M \left[\|\nabla_g \psi\|_g^2 + \frac{1}{4} (\text{Sc}_{\alpha,\beta} - \alpha \Delta_g f + \beta \|\nabla_g f\|_g^2) \|\psi\|_g^2 \right] d\text{Vol}_g \\ &= \int_M \left[\|\nabla_g \psi\|_g^2 + \left(\frac{1}{4} \text{Sc}_{\alpha,\beta} + \frac{\beta}{4} \|\nabla_g f\|_g^2 \right) \|\psi\|_g^2 + \frac{\alpha}{4} \langle \nabla_g f, \nabla_g \|\psi\|_g^2 \rangle_g \right] d\text{Vol}_g. \end{aligned}$$

Then one gets

$$\begin{aligned} \frac{|\alpha|}{4} |\langle \nabla_g f, \nabla_g \|\psi\|_g^2 \rangle_g| &\leq \frac{|\alpha|}{4} (c \|\nabla_g f\|_g \|\psi\|_g \times 2c^{-1} \|\nabla_g \psi\|_g) \\ &\leq \frac{|\alpha|c^2}{8} \|\nabla_g f\|_g^2 \|\psi\|_g^2 + \frac{c^{-2}|\alpha|}{2} \|\nabla_g \psi\|_g^2. \end{aligned}$$

Therefore,

$$0 \geq \int_M \left[\left(1 - \frac{c^{-2}|\alpha|}{2} \right) \|\nabla_g \psi\|_g^2 + \frac{2\beta - c^2|\alpha|}{8} \|\nabla_g f\|_g^2 \|\psi\|_g^2 + \frac{1}{4} \text{Sc}_{\alpha,\beta} \|\psi\|_g^2 \right] d\text{Vol}_g,$$

where $c \neq 0$. If $c^{-2}|\alpha| \leq 2$, $\beta \geq \frac{c^2|\alpha|}{2}$ and $\text{Sc}_{\alpha,\beta} > 0$, then $\psi = 0$. So the conditions $\alpha \in \mathbb{R}$ and $\beta \geq \frac{|\alpha|^2}{4}$ are needed

$$\begin{aligned} \frac{|\alpha|}{4} |\langle \nabla_g f, \nabla_g \|\psi\|_g^2 \rangle_g| &\leq \frac{|\alpha|}{4} (\|\nabla_g f\|_g \|\psi\|_g \times 2\|\nabla_g \psi\|_g) \\ &= \frac{|\alpha|}{2} (c_1 \|\nabla_g f\|_g \|\psi\|_g \times c_1^{-1} \|\nabla_g \psi\|_g) \\ &\leq \frac{|\alpha|}{4} (c_1^2 \|\nabla_g f\|_g^2 \|\psi\|_g^2 + c_1^{-2} \|\nabla_g \psi\|_g^2). \end{aligned}$$

Thus,

$$0 \geq \int_M \left[\left(1 - \frac{c_1^{-2}|\alpha|}{4} \right) \|\nabla_g \psi\|_g^2 + \frac{\beta - c_1^2|\alpha|}{4} \|\nabla_g f\|_g^2 \|\psi\|_g^2 + \frac{1}{4} \text{Sc}_{\alpha,\beta} \|\psi\|_g^2 \right] d\text{Vol}_g,$$

where $c_1 \neq 0$. If $c_1^{-2}|\alpha| \leq 4$, $\beta \geq c_1^2|\alpha|$ and $\text{Sc}_{\alpha,\beta} > 0$, then $\psi = 0$. Also the conditions $\alpha \in \mathbb{R}$ and $\beta \geq \frac{|\alpha|^2}{4}$ are needed. ■

The following 3 corollaries come from the proposition of vanishing of harmonic spinors.

Corollary 4.4. *Assume the smooth mm-space $(M^n, g, e^{-f} d\text{Vol}_g)$ is closed and spin. If $\alpha \in \mathbb{R}$, $\beta \geq \frac{|\alpha|^2}{4}$ and $\text{Sc}_{\alpha,\beta} > 0$, then the \widehat{A} -genus and the Rosenberg index of M^n vanish.*

Proof. Since the $C^*(\pi_1(M^n))$ -bundle in the construction of the Rosenberg index [35] is flat, there are no correction terms due to curvature of the bundle. Then the Schrödinger–Lichnerowicz–Weitzenböck formula and the argument in the proof of vanishing harmonic spinors can be applied without change. ■

Corollary 4.5. *Assume that M^n is a closed spin n -manifold and f is a smooth function on M^n . If one of the following conditions is met,*

- (1) $N \subset M^n$ is a codimension one closed connected submanifold with trivial normal bundle, the inclusion of fundamental groups $\pi_1(N^{n-1}) \rightarrow \pi_1(M^n)$ is injective and the Rosenberg index of N does not vanish, or

- (2) $N \subset M^n$ is a codimension two closed connected submanifold with trivial normal bundle, $\pi_2(M^n) = 0$, the inclusion of fundamental groups $\pi_1(N^{n-1}) \rightarrow \pi_1(M^n)$ is injective and the Rosenberg index of N does not vanish, or
- (3) $N = N_1 \cap \cdots \cap N_k$, where $N_1 \cdots N_k \subset M$ are closed submanifolds that intersect mutually transversely and have trivial normal bundles. Suppose that the codimension of N_i is at most two for all $i \in \{1, \dots, k\}$ and $\pi_2(N) \rightarrow \pi_2(M)$ is surjective and $\hat{A}(N) \neq 0$,

then M^n does not admit a Riemannian metric g such that the smooth mm-space $(M^n, g, e^{-f} \text{dvol}_g)$ satisfies $\text{Sc}_{\alpha, \beta} > 0$ for the dimension $n \geq 3$, $\alpha \in \mathbb{R}$ and $\beta \geq \frac{|\alpha|^2}{4}$.

Proof. The results in the [22, Theorem 1.1] and [42, Theorem 1.9] can be applied to show that the Rosenberg index of M^n does not vanish and Corollary 4.4 implies the theorem. ■

Let $\mathcal{R}_f(M^n) := \{(g, f)\}$ be the space of densities, where g is a smooth Riemannian metric on M^n and f is a smooth function on M^n and $\mathcal{R}_f^+(M^n) \subset \mathcal{R}_f(M^n)$ is the subspace of densities such that the smooth mm-space $(M^n, g, e^{-f} \text{dvol}_g)$ satisfies $\text{Sc}_{\alpha, \beta} > 0$. Furthermore, let $\mathcal{R}_f^+(M^n)$ be endowed with the smooth topology.

Corollary 4.6. Assume M^n is a closed spin n -manifold, $n \geq 3$, $\alpha \in \mathbb{R}$ and $\beta \geq \frac{|\alpha|^2}{4}$ and $\mathcal{R}_f^+(M^n) \neq \emptyset$, then there exists a homomorphism

$$A_{m-1}: \pi_{m-1}(\mathcal{R}_f^+(M^n)) \rightarrow KO_{n+m}$$

such that

- $A_0 \neq 0$, if $n \equiv 0, 1 \pmod{8}$,
- $A_1 \neq 0$, if $n \equiv -1, 0 \pmod{8}$,
- $A_{8j+1-n} \neq 0$, if $n \geq 7$ and $8j - n \geq 0$.

Proof. Since the results in the [23, Section 4.4] and [9] depend on the existence of exotic spheres with non-vanishing α -invariant. Let $\phi: M^n \rightarrow M^n$ be a diffeomorphism of M^n and $(g, f) \in \mathcal{R}_f^+(M^n)$, then $(\phi^*g, f \circ \phi)$ is also in $\mathcal{R}_f^+(M^n)$. Combining it with Proposition 4.3 shows that Hitchin's construction of the map A [23, Proposition 4.6] can be applied to the case of $\mathcal{R}_f^+(M^n)$ and then we can finish the proof with the arguments in [23, Section 4.4] and [9, Section 2.5]. ■

4.2 Conformal to PSC-metrics

Proposition 4.7 (conformal to PSC-metrics). Let $(M^n, g, e^{-f} \text{dVol}_g)$ be a closed smooth mm-space with $\text{Sc}_{\alpha, \beta} > 0$. If the dimension $n \geq 3$, $\alpha \in \mathbb{R}$ and $\beta \geq \frac{(n-2)|\alpha|^2}{4(n-1)}$, then there is a metric \tilde{g} conformal to g with positive scalar curvature (PSC-metric).

Proof. One only needs to show for all nontrivial u , $\int_M -u L_g u \text{dVol}_g > 0$ as in the Yamabe problem [36], where

$$L_g := \Delta_g - \frac{n-2}{4(n-1)} \text{Sc}_g$$

is conformal Laplacian operator. To see this,

$$\begin{aligned}
\int_M -u L_g u \, d\text{Vol}_g &= \int_M \left[\|\nabla_g u\|_g^2 + \frac{n-2}{4(n-1)} \text{Sc}_g u^2 \right] d\text{Vol}_g \\
&= \int_M \left[\|\nabla_g u\|_g^2 + \frac{n-2}{4(n-1)} (\text{Sc}_{\alpha,\beta} - \alpha \triangle_g f + \beta \|\nabla_g f\|_g^2) u^2 \right] d\text{Vol}_g \\
&= \int_M \left[\|\nabla_g u\|_g^2 + \frac{n-2}{4(n-1)} (\text{Sc}_{\alpha,\beta} + \beta \|\nabla_g f\|_g^2) u^2 \right. \\
&\quad \left. + \frac{\alpha(n-2)}{2(n-1)} \langle \nabla_g f, \nabla_g u \rangle_g u \right] d\text{Vol}_g.
\end{aligned}$$

Through the inequality

$$\langle \nabla_g f, \nabla_g u \rangle_g u \leq c_2 \|\nabla_g f\|_g u \times c_2^{-1} \|\nabla_g u\|_g \leq \frac{c_2^2 \|\nabla_g f\|_g^2 u^2 + c_2^{-2} \|\nabla_g u\|_g^2}{2},$$

one gets

$$\begin{aligned}
\int_M -u L_g u \, d\text{Vol}_g &\geq \int_M \left[\left(1 - \frac{|\alpha| c_2^{-2} (n-2)}{4(n-1)} \right) \|\nabla_g u\|_g^2 \right. \\
&\quad \left. + \frac{(\beta - |\alpha| c_2^{-2}) (n-2)}{4(n-1)} \|\nabla_g f\|_g^2 u^2 + \frac{n-2}{4(n-1)} \text{Sc}_{\alpha,\beta} u^2 \right] d\text{Vol}_g,
\end{aligned}$$

where $c_2 \neq 0$.

If $|\alpha| c_2^{-2} \leq \frac{4(n-1)}{n-2}$, $\beta \geq c_2^2 |\alpha|$ and $\text{Sc}_{\alpha,\beta} > 0$, then

$$\int_M -u L_g u \, d\text{Vol}_g > 0.$$

So the conditions $n > 2$, $\alpha \in \mathbb{R}$ and $\beta \geq \frac{(n-2)\alpha^2}{4(n-1)}$ are needed. ■

Remark 4.8. The proof was borrowed from [1, Proposition 2]. The two propositions above offer a geometric reason why the condition of the vanishing of \hat{A} -genus (without simply connected condition) does not imply that M^n can admit a PSC-metric for the closed spin manifold M^n .

The proposition of conformal to PSC-metrics has following 3 corollaries.

Corollary 4.9 (weighted spherical Lipschitz bounded). *Let $(M^n, g, e^{-f} d\text{Vol}_g)$ be a closed orientable smooth mm-space with $\text{Sc}_{\alpha,\beta} \geq \kappa > 0$, $3 \leq n \leq 8$, $\alpha \in \mathbb{R}$ and $\beta \geq \frac{(n-2)|\alpha|^2}{4(n-1)}$, then the Lipschitz constant of the continuous map ϕ from $(M^n, g, e^{-f} d\text{Vol}_g)$ to the sphere S^n with standard metric of non-zero degrees has uniformly non-zero lower bounded.*

Proof. There is a metric \tilde{g} conformal to g with scalar curvature $\geq n(n-1)$ by the proposition of conformal PSC-metrics. For the continuous map ϕ from (M^n, \tilde{g}) to S^n of non-zero degrees, the Lipschitz constant of ϕ is greater than a constant that depends only on the dimensions n by Gromov's spherical Lipschitz bounded theorem [19, Section 3]. Since the conformal function has the positive upper bound by the compactness of the manifold, then the Lipschitz constant has uniformly non-zero lower bounded. ■

Corollary 4.10. *For the closed smooth mm-space $(M^n, g, e^{-f} d\text{Vol}_g)$ ($n \geq 3$) with $\text{Sc}^{\text{vol}_n}(M^n) > 0$, there is a metric \hat{g} conformal to g with PSC-metric. In particular, the \hat{A} -genus and Rosenberg index vanish with additional spin condition.*

For the closed orientable smooth mm-space $(M^n, g, e^{-f} d\text{Vol}_g)$ ($3 \leq n \leq 8$) with $\text{Sc}^{\text{vol}_n}(M^n) \geq \kappa > 0$, then the Lipschitz constant of the continuous map ϕ from $(M^n, g, e^{-f} d\text{Vol}_g)$ to the sphere S^n with standard metric of non-zero degrees has uniformly non-zero lower bounded.

Proof. The volume of the small disk of $(M^n, g, e^{-f} d\text{Vol}_g)$ was computed in [33, Theorem 8],

$$\mu(B_r(x)) = \text{vol}_E(B_r(\mathbf{R}^n)) \left[1 - \frac{\text{Sc}_g + 3 \Delta_g f - 3 \|\nabla_g f\|_g^2}{6(n+2)} r^2 + O(r^4) \right]$$

as $r \rightarrow 0$. Since $\text{Sc}^{\text{vol}_n}(M^n) > 0$, i.e., $\mu(B_r(x)) < \text{vol}_E(B_r)$ as $r \rightarrow 0$, then

$$\text{Sc}_g + 3 \Delta_g f - 3 \|\nabla_g f\|_g^2 > 0.$$

Therefore, the propositions of vanishing harmonic spinors and of conformal PSC-metrics and Corollary 4.4 imply it. \blacksquare

Remark 4.11. Since any weighted Riemannian manifold (with non-trivial Borel measure) is infinitesimally Hilbertian (see [28]), Corollary 4.10 also works for $(M^n, g, e^{-f} d\text{Vol}_g)$ with RSC(κ, n) condition.

Enlargeability as an obstruction to the existence of a PSC-metric on a closed manifold was introduced by Gromov–Lawson. We call a manifold enlargeable as Gromov–Lawson’s definition in [21, Definition 5.5]

Corollary 4.12. *Assume M^n ($n \geq 3$) is a closed spin smooth enlargeable manifold, then $\mathcal{R}_f^+(M^n)$ is an empty set for $\alpha \in \mathbb{R}$ and $\beta \geq \frac{(n-2)|\alpha|^2}{4(n-1)}$.*

In particular, $(\mathbb{T}^n, g, e^{-f} d\text{Vol}_g)$ does not satisfy $\text{Sc}^{\text{vol}_n}(\mathbb{T}^n) > 0$ for any C^2 -smooth Riemannian metrics g and C^2 -smooth functions f on the torus \mathbb{T}^n .

Proof. Since a closed enlargeable manifold cannot carry a PSC-metric [21, Theorem 5.8], Proposition 4.7 implies $\mathcal{R}_f^+(M^n) = \emptyset$ for $\alpha \in \mathbb{R}$ and $\beta \geq \frac{(n-2)|\alpha|^2}{4(n-1)}$.

\mathbb{T}^n is an important example of enlargeable manifolds and then Corollary 4.10 implies that $(\mathbb{T}^n, g, e^{-f} d\text{Vol}_g)$ does not satisfy $\text{Sc}^{\text{vol}_n}(\mathbb{T}^n) > 0$ for $n \geq 3$. For dimension 2, the conditions of $\text{Sc}_{\alpha, \beta} > 0$ and $\beta \geq 0$ imply that the oriented surface is 2-sphere. \blacksquare

4.3 f -minimal hypersurface and $\text{Sc}_{\alpha, \beta} > 0$

In addition to using the Dirac operator method, Schoen–Yau’s minimal hypersurface method [37] is another main idea. For an immersed orientable hypersurface $N^{n-1} \subset M^n$, the weighted mean curvature vector H_f of N^{n-1} is defined by Gromov in [16, Section 9.4.E],

$$H_f = H + (\nabla_g f)^\perp,$$

where H is the mean curvature vector field of the immersion, $(\cdot)^\perp$ is the projection on the normal bundle of N^{n-1} . The first and second variational formulae for the weighted volume functional of N^{n-1} were derived in Bayle’s thesis (also see [34]). We take the detailed presentation of such derivation for [8]. The (N^{n-1}, \bar{g}) with the induced metric is called f -minimal hypersurface if the weighted mean curvature vector H_f vanishes identically.

In fact, the definition of f -minimal hypersurface can also be derived from the first variational formula. Furthermore, an f -minimal hypersurface is a minimal hypersurface of (M^n, \tilde{g}) , where \tilde{g} is the conformal metric of g , $\tilde{g} = e^{-\frac{2f}{n-1}} g$.

The connection between the geometry of the ambient smooth mm-space and the f -minimal hypersurfaces occurs via the second variation of the weighted volume functional. For a hypersurface (N^{n-1}, \bar{g}) , the L_f operator is defined by

$$L_f := \Delta_f + |A|^2 + \text{Ric}_f^M(\nu, \nu),$$

where ν is the unit normal vector, $|A|^2$ denotes the square of the norm of the second fundamental form A of N^{n-1} and

$$\Delta_f := \Delta_{\bar{g}} - \langle \nabla_{\bar{g}} f, \nabla_{\bar{g}} \cdot \rangle$$

is the weighted Laplacian. Through the second variational formula, a two-sided f -minimal hypersurface N^{n-1} is stable (called L_f -stable) if for any compactly supported smooth function $u \in C_c^\infty(N^{n-1})$, it holds that

$$-\int_N u L_f u e^{-f} d\text{Vol}_{\bar{g}} \geq 0.$$

Proposition 4.13. *Let $(M^n, g, e^{-f} d\text{Vol}_g)$ be a closed orientable smooth mm-space with $\text{Sc}_{\alpha, \beta} > 0$ and (N^{n-1}, \bar{g}) be the compact L_f -stable minimal hypersurface of $(M^n, g, e^{-f} d\text{Vol}_g)$. If the dimension $n \geq 3$, $\alpha = 2$, and $\beta \geq \frac{n-2}{n-1}$, then there exists a PSC-metric conformal to \bar{g} on N^{n-1} .*

Proof. The f -minimal hypersurface (N^{n-1}, \bar{g}) is L_f -stable if and only if (N^{n-1}, \tilde{g}) is stable as a minimal hypersurface on (M^n, \tilde{g}) , where $\tilde{g} := e^{-\frac{2f}{n-1}} g$ and \tilde{g} is the induced metric of \tilde{g} (see [8, Appendix]). On the other hand, the scalar curvature of (M^n, \tilde{g}) is

$$\text{Sc}_{\tilde{g}} = e^{\frac{f}{n-1}} \left(\text{Sc}_g + 2 \Delta_g f - \frac{n-2}{n-1} \|\nabla_g f\|_g^2 \right).$$

Thus, $\text{Sc}_{\alpha, \beta} > 0$ with $n \geq 3$, $\alpha = 2$, and $\beta \geq \frac{n-2}{n-1}$ imply $\text{Sc}_{\tilde{g}} > 0$. Then the standard Schoen–Yau's argument can be applied to show that \tilde{g} conformal to a PSC-metric on N^{n-1} . ■

Remark 4.14. The minimal hypersurface method poses a stricter condition to the valid range of α, β than that of the Dirac operator method.

Since the oriented closed manifolds with a PSC-metric in 2 and 3 dimensions are classified by Gauess–Bonnet theorem and Perelman–Thurston geometrization theorem, then Proposition 4.13 can give the following elementary applications:

Corollary 4.15. *Let $(M^n, g, e^{-f} d\text{Vol}_g)$ be a closed orientable smooth mm-space with $\text{Sc}_{\alpha, \beta} > 0$.*

1. *If $n = 3$, $\alpha = 2$, and $\beta \geq \frac{1}{2}$, then there is no closed immersed L_f -stable minimal 2-dimensional surface with positive genus.*
2. *If $n = 4$, $\alpha = 2$, and $\beta \geq \frac{2}{3}$, then the closed immersed L_f -stable minimal 3-dimensional submanifold must be spherical 3-manifolds, $S^2 \times S^1$ or the connected sum of spherical 3-manifolds and copies of $S^2 \times S^1$.*

Remark 4.16 (historical remark). The prototype of Corollary 4.15(1) is the Schoen–Yau's classic result, which said that the oriented closed 3-manifold with a PSC-metric has no compact immersed stable minimal surface of positive genus [37]. The Schoen–Yau result had been generalized to Perelman's P-scalar curvature > 0 by Fan [10]. Note that one can also consider the noncompact immersed L_f -stable minimal 2-dimensional surface under the condition of Corollary 4.15(1) since an oriented complete stable minimal surface in a complete oriented 3-manifold with a PSC-metric is conformally equivalent to the complex plane \mathbb{C} showed by Fischer–Colbrie–Schoen [11].

The smooth mm-space with $\text{Sc}_{\alpha, \beta} > 0$ under suit ranges of α and β implies the manifold admits PSC-metrics, but the manifold (itself) that can admit PSC-metrics does not necessarily imply there exists $\text{Sc}_{\alpha, \beta} > 0$.

Question 4.17. Does the smooth mm-space with $\text{Sc}_{\alpha,\beta} > 0$ under suitable ranges of α and β give more topological restriction on the manifold than the PSC-metric on the manifold?

Question 4.18. Let M be a closed smooth manifold, f be a smooth function on M and h be a smooth function that is negative for some point p on M . What is the range of α and β such that there exists a smooth Riemannian metric g on M satisfying

$$\text{Sc}_g + \alpha \Delta_g f - \beta \|\nabla_g f\|_g^2 = h,$$

i.e., $\text{Sc}_{\alpha,\beta}(g) = h$?

Let (M, g_i) be smooth Riemannian manifolds and $\{g_i\}_{i \in \mathbb{N}}$ C^0 -converges to g , then $\{g_i\}_{i \in \mathbb{N}}$ also smGH-converges to g . Gromov showed that the scalar curvature $\geq \kappa$ is stable under C^0 -convergence in [17, Section 1.8].

Question 4.19. Assume smooth mm-spaces $(M^n, g_i, e^{-f} d\text{Vol}_{g_i})$ all satisfy $\text{Sc}^{\text{vol}_n}(M^n) \geq 0$ such that $\{g_i\}_{i \in \mathbb{N}}$ C^2 -converges to the smooth Riemannian metric g on M^n , then does $(M^n, g, e^{-f} d\text{Vol}_g)$ also satisfy $\text{Sc}^{\text{vol}_n}(M^n) \geq 0$?

Question 4.20. Let mm-spaces $(M^n, g, e^{-f} d\text{Vol}_g)$ with $\text{Sc}^{\text{vol}_n}(M^n) \geq \kappa > 0$, where M^n is a closed smooth manifold, g and f are C^0 -smooth, then does there exist a PSC-metric on M^n ?

Since the role of $\text{Sc}_{\alpha,\beta} > 0$ on the smooth mm-space is similar to the role of $\text{Sc} > 0$ on the manifold, one can try to extend the knowledge about $\text{Sc} > 0$ to $\text{Sc}_{\alpha,\beta} > 0$.

4.4 Weighted rigidity

Gromov's conjecture that said if a smooth Riemannian metric g satisfies $g \geq g_{\text{st}}$ and $\text{Sc}(g) \geq \text{Sc}(g_{\text{st}}) = n(n-1)$ on S^n then $g = g_{\text{st}}$, was proved by Llarull [27] and called Llarull rigidity theorem. A map $h: (M^n, g_M) \rightarrow (N^n, g_N)$ is said to be ϵ -contracting if $\|h_* v\|_{g_N} \leq \epsilon \|v\|_{g_M}$ for all tangent vectors v on M^n .

Proposition 4.21 (weighted rigidity). Assume the smooth mm-space $(M^n, g, e^{-f} d\text{Vol}_g)$ is closed and spin and there exists a smooth 1-contracting map $h: (M^n, g) \rightarrow (S^n, g_{\text{st}})$ of non-zero degree. If $\alpha \in \mathbb{R}$, $\beta \geq \frac{|\alpha|^2}{4}$ and $\text{Sc}_{\alpha,\beta} \geq n(n-1)$, then h is an isometry between the metrics g and g_{st} . Furthermore, if $\alpha > 0$, then f is a constant function.

Proof. One just need to insert the tricks in the proof of Proposition 4.3 to the proof in [27, Theorem 4.1]. Following the setup of Llarull, we only prove the even-dimensional ($2n$) case without loss of generality.

First, we will show that h is an isometry. Fix $p \in M^{2n}$. Let $\{e_1, \dots, e_{2n}\}$ be a g -orthonormal tangent frame near p such that $(\nabla_g e_k)_p = 0$ for each k . Let $\{\epsilon_1, \dots, \epsilon_{2n}\}$ be a g_{st} -orthonormal tangent frame near $h(p) \in S^{2n}$ such that $(\nabla_{g_{\text{st}}} \epsilon_k)_{h(p)} = 0$ for each k . Moreover, the bases $\{e_1, \dots, e_{2n}\}$ and $\{\epsilon_1, \dots, \epsilon_{2n}\}$ can be chosen so that $\epsilon_j = \lambda_j h_* e_j$ for appropriate $\{\lambda_j\}_{j=1}^{2n}$. This is possible since h_* is symmetric. Since h is 1-contracting map, $\lambda_k \geq 1$ for each k .

Then one constructs the twisted vector bundles $S \otimes E$ over M^{2n} as Llarull did. Let R^E be the curvature tensor of E and ψ be a twisted spinor, then one gets

$$\langle R^E \psi, \psi \rangle_g \geq -\frac{1}{4} \sum_{i \neq j} \frac{1}{\lambda_i \lambda_j} \|\psi\|_g.$$

For the twisted Dirac operator \mathbb{D}_E , one has $\mathbb{D}_E^2 = \nabla^* \nabla + \frac{1}{4} \text{Sc}_g + R^E$ and

$$\begin{aligned} \int_M \langle \mathbb{D}_E^2 \psi, \psi \rangle_g d\text{Vol}_g &= \int_M \left[\|\nabla_g \psi\|_g^2 + \frac{1}{4} (\text{Sc}_{\alpha, \beta} - \alpha \Delta_g f + \beta \|\nabla_g f\|_g^2) \|\psi\|_g^2 \right. \\ &\quad \left. + \langle R^E \psi, \psi \rangle_g \right] d\text{Vol}_g \\ &= \int_M \left[\|\nabla_g \psi\|_g^2 + \left(\frac{1}{4} \text{Sc}_{\alpha, \beta} + \frac{\beta}{4} \|\nabla_g f\|_g^2 \right) \|\psi\|_g^2 \right. \\ &\quad \left. + \frac{\alpha}{4} \langle \nabla_g f, \nabla_g \|\psi\|_g^2 \rangle_g + \langle R^E \psi, \psi \rangle_g \right] d\text{Vol}_g. \end{aligned}$$

Because $\lambda_k \geq 1$ for each k , one gets

$$\langle R^E \psi, \psi \rangle_g \geq \frac{-2n(2n-1)}{4} \|\psi\|_g^2$$

and then

$$\begin{aligned} \frac{|\alpha|}{4} |\langle \nabla_g f, \nabla_g \|\psi\|_g^2 \rangle_g| &\leq \frac{|\alpha|}{4} (\|\nabla_g f\|_g \|\psi\|_g \times 2 \|\nabla_g \psi\|_g) \\ &= \frac{|\alpha|}{2} (c_1 \|\nabla_g f\|_g \|\psi\|_g \times c_1^{-1} \|\nabla_g \psi\|_g) \\ &\leq \frac{|\alpha|}{4} (c_1^2 \|\nabla_g f\|_g^2 \|\psi\|_g^2 + c_1^{-2} \|\nabla_g \psi\|_g^2), \end{aligned}$$

where $c_1 \neq 0$. Therefore,

$$\begin{aligned} \int_M \langle \mathbb{D}_E^2 \psi, \psi \rangle_g d\text{Vol}_g &\geq \int_M \left[\left(1 - \frac{c_1^{-2} |\alpha|}{4} \right) \|\nabla_g \psi\|_g^2 + \frac{\beta - c_1^2 |\alpha|}{4} \|\nabla_g f\|_g^2 \|\psi\|_g^2 \right. \\ &\quad \left. + \frac{1}{4} (\text{Sc}_{\alpha, \beta} - 2n(2n-1)) \|\psi\|_g^2 \right] d\text{Vol}_g. \end{aligned}$$

Furthermore, since $\alpha \in \mathbb{R}$, $\beta \geq \frac{|\alpha|^2}{4}$ and $\text{Sc}_{\alpha, \beta} \geq 2n(2n-1)$, one can choose c_1 such that $c_1^{-2} |\alpha| \leq 4$, then $\beta - c_1^2 |\alpha| \geq 0$. Thus,

$$\int_M \langle \mathbb{D}_E^2 \psi, \psi \rangle_g d\text{Vol}_g \geq \int_M \frac{1}{4} [\text{Sc}_{\alpha, \beta} - 2n(2n-1)] \|\psi\|_g^2 d\text{Vol}_g \geq 0.$$

The fact $\text{Index}(\mathbb{D}_{E+}) \neq 0$ implies $\ker(\mathbb{D}_E) \neq 0$ and then $\text{Sc}_{\alpha, \beta} = 2n(2n-1)$. Then using the inequality $\langle R^E \psi, \psi \rangle_g \geq -\frac{1}{4} \sum_{i \neq j} \frac{1}{\lambda_i \lambda_j} \|\psi\|_g^2$, one gets

$$\int_M \langle \mathbb{D}_E^2 \psi, \psi \rangle_g d\text{Vol}_g \geq \int_M \frac{1}{4} \left[\sum_{i \neq j} \left(1 - \frac{1}{\lambda_i \lambda_j} \right) \right] \|\psi\|_g^2 d\text{Vol}_g \geq 0.$$

Choosing $\psi \neq 0$ such that $\mathbb{D}_E \psi = 0$, one has

$$0 \leq 1 - \frac{1}{\lambda_i \lambda_j} \leq 0$$

for $i \neq j$. Thus, $\lambda_k = 0$ for all $1 \leq k \leq 2n$ and h is an isometry.

Second, we will show that f is a constant function. Since $\text{Sc}_{\alpha, \beta} = 2n(2n-1)$, $\text{Sc}_g = 2n(2n-1)$, $\alpha > 0$ and $\beta \geq \frac{|\alpha|^2}{4}$, then $\Delta_g f \geq 0$. One has

$$\int_M \Delta_g f d\text{Vol}_g = 0$$

for a closed manifold M^n , so one gets $\Delta_g f = 0$. That implies $\nabla_g f = 0$ so that f is a constant function on M^n . ■

Corollary 4.22. *Let the closed and spin smooth mm-space $(M^n, g, e^{-f} d\text{Vol}_g)$ with $\text{Sc}^{\text{vol}_n}(M^n) \geq n(n-1)$ and there exists a smooth 1-contracting map $h: (M^n, g) \rightarrow (S^n, g_{\text{st}})$ of non-zero degree, then h is an isometry between the metrics g and g_{st} .*

Proof. Combining the weighted rigidity theorem and the proof of Corollary 4.10 can imply it. ■

As Llarull rigidity theorem (and the weighted rigidity theorem) still holds if the condition that h is 1-contracting is replaced by the condition that h is area-contracting, Gromov called such metrics area-extremal metrics, asked which manifolds possess area-extremal metrics, and conjectured that Riemannian symmetric spaces should have area-extremal metrics [15], [18, Section 17] and [20, Section 4.2]. Goette–Semmelmann showed that several classes of symmetric spaces with non-constant curvatures are area-extremal [13].

Question 4.23. *Can Goette–Semmelmann’s results [13] be generalized to the smooth mm-space with $\text{Sc}_{\alpha, \beta} > 0$ under other suitable conditions?*

Since Corollary 4.15(1) showed that the closed orientable immersed L_f -stable minimal 2-dimensional surface in the closed orientable smooth mm-space $(M^n, g, e^{-f} d\text{Vol}_g)$ with $\text{Sc}_{2, \beta} > 0$ ($\beta \geq \frac{1}{2}$) is 2-sphere, then one can consider rigidity of area-minimizing 2-sphere in 3-dimensional smooth mm-space. Bray’s volume comparison theorem [4, Chapter 3, Theorem 18] is another rigidity theorem that needs the conditions of Ricci curvature and scalar curvature bounded below. There are other rigidity phenomena involving scalar curvature, see [5].

Question 4.24. *Can Bray’s volume comparison theorem be extended to the smooth mm-space?*

Question 4.25. *What is the correct Einstein field equation on the smooth mm-space?*

If one replaces the Ricci and scalar curvature on the left hand side of Einstein field question by Ric_f^M and $\text{Sc}_{\alpha, \beta}$ for the smooth mm-space $(M^n, g, e^{-f} d\text{Vol}_g)$, then what is the stress–energy tensor on the right hand side in this case?

Acknowledgements

I am grateful to Thomas Schick for his help, the referees for their useful comments, and the funding from China Scholarship Council.

References

- [1] Abedin F., Corvino J., On the P -scalar curvature, *J. Geom. Anal.* **27** (2017), 1589–1623.
- [2] Ambrosio L., Gigli N., Savaré G., Metric measure spaces with Riemannian Ricci curvature bounded from below, *Duke Math. J.* **163** (2014), 1405–1490, [arXiv:1109.0222](#).
- [3] Athreya S., Löhr W., Winter A., The gap between Gromov-vague and Gromov–Hausdorff-vague topology, *Stochastic Process. Appl.* **126** (2016), 2527–2553, [arXiv:1407.6309](#).
- [4] Bray H.L., The Penrose inequality in general relativity and volume comparison theorems involving scalar curvature, Ph.D. Thesis, Stanford University, 1997, [arXiv:0902.3241](#).
- [5] Brendle S., Rigidity phenomena involving scalar curvature, in Surveys in Differential Geometry, *Surv. Differ. Geom.*, Vol. 17, *Int. Press*, Boston, MA, 2012, 179–202, [arXiv:1008.3097](#).
- [6] Case J.S., Conformal invariants measuring the best constants for Gagliardo–Nirenberg–Sobolev inequalities, *Calc. Var. Partial Differential Equations* **48** (2013), 507–526, [arXiv:1112.3977](#).
- [7] Chang S.-Y.A., Gursky M.J., Yang P., Conformal invariants associated to a measure, *Proc. Natl. Acad. Sci. USA* **103** (2006), 2535–2540.
- [8] Cheng X., Mejia T., Zhou D., Stability and compactness for complete f -minimal surfaces, *Trans. Amer. Math. Soc.* **367** (2015), 4041–4059, [arXiv:1210.8076](#).

- [9] Crowley D., Schick T., The Gromoll filtration, KO -characteristic classes and metrics of positive scalar curvature, *Geom. Topol.* **17** (2013), 1773–1789, [arXiv:1204.6474](#).
- [10] Fan E.M., Topology of three-manifolds with positive P -scalar curvature, *Proc. Amer. Math. Soc.* **136** (2008), 3255–3261.
- [11] Fischer-Colbrie D., Schoen R., The structure of complete stable minimal surfaces in 3-manifolds of nonnegative scalar curvature, *Comm. Pure Appl. Math.* **33** (1980), 199–211.
- [12] Gigli N., On the differential structure of metric measure spaces and applications, *Mem. Amer. Math. Soc.* **236** (2015), vi+91 pages, [arXiv:1205.6622](#).
- [13] Goette S., Semmelmann U., Scalar curvature estimates for compact symmetric spaces, *Differential Geom. Appl.* **16** (2002), 65–78, [arXiv:math.DG/0010199](#).
- [14] Gray A., Vanhecke L., Riemannian geometry as determined by the volumes of small geodesic balls, *Acta Math.* **142** (1979), 157–198.
- [15] Gromov M., Positive curvature, macroscopic dimension, spectral gaps and higher signatures, in Functional Analysis on the Eve of the 21st Century, Vol. II (New Brunswick, NJ, 1993), *Progr. Math.*, Vol. 132, Birkhäuser Boston, Boston, MA, 1996, 1–213.
- [16] Gromov M., Isoperimetry of waists and concentration of maps, *Geom. Funct. Anal.* **13** (2003), 178–215.
- [17] Gromov M., Dirac and Plateau billiards in domains with corners, *Cent. Eur. J. Math.* **12** (2014), 1109–1156, [arXiv:1811.04318](#).
- [18] Gromov M., 101 questions, problems and conjectures around scalar curvature, 2017, available at <https://www.ihes.fr/~gromov/wp-content/uploads/2018/08/101-problemsOct1-2017.pdf>.
- [19] Gromov M., Metric inequalities with scalar curvature, *Geom. Funct. Anal.* **28** (2018), 645–726, [arXiv:1710.04655](#).
- [20] Gromov M., Four lectures on scalar curvature, [arXiv:1908.10612](#).
- [21] Gromov M., Lawson Jr. H.B., Positive scalar curvature and the Dirac operator on complete Riemannian manifolds, *Inst. Hautes Études Sci. Publ. Math.* **58** (1983), 83–196.
- [22] Hanke B., Pape D., Schick T., Codimension two index obstructions to positive scalar curvature, *Ann. Inst. Fourier (Grenoble)* **65** (2015), 2681–2710, [arXiv:1402.4094](#).
- [23] Hitchin N., Harmonic spinors, *Adv. Math.* **14** (1974), 1–55.
- [24] Kazdan J.L., Deformation to positive scalar curvature on complete manifolds, *Math. Ann.* **261** (1982), 227–234.
- [25] Kazdan J.L., Warner F.W., Scalar curvature and conformal deformation of Riemannian structure, *J. Differential Geometry* **10** (1975), 113–134.
- [26] Lasserre J.B., On the setwise convergence of sequences of measures, *J. Appl. Math. Stochastic Anal.* **10** (1997), 131–136.
- [27] Llarull M., Sharp estimates and the Dirac operator, *Math. Ann.* **310** (1998), 55–71.
- [28] Lučić D., Pasqualetto E., Infinitesimal Hilbertianity of weighted Riemannian manifolds, *Canad. Math. Bull.* **63** (2020), 118–140, [arXiv:1809.05919](#).
- [29] Ohta S., On the measure contraction property of metric measure spaces, *Comment. Math. Helv.* **82** (2007), 805–828.
- [30] Ohta S., Finsler interpolation inequalities, *Calc. Var. Partial Differential Equations* **36** (2009), 211–249.
- [31] Perelman G., The entropy formula for the Ricci flow and its geometric applications, [arXiv:math.DG/0211159](#).
- [32] Petrunin A., Alexandrov meets Lott–Villani–Sturm, *Münster J. Math.* **4** (2011), 53–64, [arXiv:1003.5948](#).
- [33] Qian Z., Estimates for weighted volumes and applications, *Quart. J. Math. Oxford Ser. (2)* **48** (1997), 235–242.
- [34] Rosales C., Cañete A., Bayle V., Morgan F., On the isoperimetric problem in Euclidean space with density, *Calc. Var. Partial Differential Equations* **31** (2008), 27–46, [arXiv:math.DG/0602135](#).
- [35] Rosenberg J., C^* -algebras, positive scalar curvature, and the Novikov conjecture, *Inst. Hautes Études Sci. Publ. Math.* **58** (1983), 197–212.
- [36] Schoen R., Conformal deformation of a Riemannian metric to constant scalar curvature, *J. Differential Geom.* **20** (1984), 479–495.

-
- [37] Schoen R., Yau S.T., Existence of incompressible minimal surfaces and the topology of three-dimensional manifolds with nonnegative scalar curvature, *Ann. of Math.* **110** (1979), 127–142.
 - [38] Sturm K.-T., On the geometry of metric measure spaces. II, *Acta Math.* **196** (2006), 133–177.
 - [39] Villani C., Optimal transport: old and new, *Grundlehren der Mathematischen Wissenschaften*, Vol. 338, Springer-Verlag, Berlin, 2009.
 - [40] Wylie W., Sectional curvature for Riemannian manifolds with density, *Geom. Dedicata* **178** (2015), 151–169, [arXiv:1311.0267](#).
 - [41] Young L.S., Dimension, entropy and Lyapunov exponents, *Ergodic Theory Dynam. Systems* **2** (1982), 109–124.
 - [42] Zeidler R., An index obstruction to positive scalar curvature on fiber bundles over aspherical manifolds, *Algebr. Geom. Topol.* **17** (2017), 3081–3094, [arXiv:1512.06781](#).
 - [43] Zhang H.-C., Zhu X.-P., Ricci curvature on Alexandrov spaces and rigidity theorems, *Comm. Anal. Geom.* **18** (2010), 503–553, [arXiv:0912.3190](#).