# Randomized extended block Kaczmarz for solving least squares

Kui Du<sup>\*</sup>, Wu-Tao Si<sup>†</sup>, Xiao-Hui Sun<sup>‡</sup>

#### Abstract

Randomized iterative algorithms have recently been proposed to solve large-scale linear systems. In this paper, we present a simple randomized extended block Kaczmarz algorithm that exponentially converges in the mean square to the unique minimum  $\ell_2$ -norm least squares solution of a given linear system of equations. The proposed algorithm is pseudoinverse-free and therefore different from the projection-based randomized double block Kaczmarz algorithm of Needell, Zhao, and Zouzias. We emphasize that our method works for all types of linear systems (consistent or inconsistent, overdetermined or underdetermined, full-rank or rank-deficient). Moreover, our approach can utilize efficient implementations on distributed computing units, yielding remarkable improvements in computational time. Numerical examples are given to show the efficiency of the new algorithm.

**Keywords**. general linear systems, minimum  $\ell_2$ -norm least squares solution, randomized extended (block) Kaczmarz, exponential convergence

AMS subject classifications: 65F10, 65F20

#### **1** Introduction

The Kaczmarz method [27] is a simple iterative method for solving a linear systems of equations

$$\mathbf{A}\mathbf{x} = \mathbf{b}, \quad \mathbf{A} \in \mathbb{R}^{m \times n}, \quad \mathbf{b} \in \mathbb{R}^{m}.$$

Due to its simplicity and numerical performance, the Kaczmarz method has found many applications in many fields, such as computer tomography [34, 28, 24], image reconstruction [45, 25], digital signal processing [9, 32], etc. At each step, the method projects the current iterate onto one hyperplane defined by a row of the system. More precisely, assuming that the *i*th row  $\mathbf{A}_{i,:}$ has been selected at the *k*th iteration, then the *k*th estimate vector  $\mathbf{x}^k$  is obtained by

$$\mathbf{x}^{k} = \mathbf{x}^{k-1} - \alpha_{k} \frac{\mathbf{A}_{i,:} \mathbf{x}^{k-1} - \mathbf{b}_{i}}{\mathbf{A}_{i,:} (\mathbf{A}_{i,:})^{\mathrm{T}}} (\mathbf{A}_{i,:})^{\mathrm{T}},$$

where  $(\mathbf{A}_{i,:})^{\mathrm{T}}$  denotes the transpose of  $\mathbf{A}_{i,:}$ ,  $\mathbf{b}_i$  is the *i*th component of  $\mathbf{b}$ , and  $\alpha_k$  is a stepsize. Numerical experiments show that using the rows of the coefficient matrix in the Kaczmarz method in random order, rather than in their given order, can often greatly improve the convergence [26, 34]. In a seminal paper [48], Strohmer and Vershynin proposed a randomized Kaczmarz (RK) algorithm which exponentially converges in expectation to the solutions of consistent, overdetermined, full-rank linear systems. The convergence result was extended and refined in various directions including inconsistent [29, 37, 51, 17, 41, 38, 21], underdetermined

<sup>\*</sup>School of Mathematical Sciences, Xiamen University, Xiamen 361005, China (kuidu@xmu.edu.cn). <sup>†</sup>School of Mathematical Sciences, Xiamen University, Xiamen 361005, China (19020171152496@stu.xmu.edu.cn).

<sup>&</sup>lt;sup>‡</sup>School of Mathematical Sciences, Xiamen University, Xiamen 361005, China (19020190154621@stu.xmu.edu.cn).

or rank-deficient linear systems [33, 20, 46, 15], ridge regression problems [23, 31], linear feasibility problems [11], convex feasibility problems [36], block variants [39, 40, 35], acceleration strategies [30, 47, 3, 4, 5, 6, 50], and many others [2, 44, 13, 14, 22].

Let  $\mathbf{A}^{\dagger}$  denote the Moore-Penrose pseudoinverse<sup>1</sup> [7] of  $\mathbf{A}$ . In this paper, we are interested in the vector  $\mathbf{A}^{\dagger}\mathbf{b}$ . Here we would like to make clear what  $\mathbf{A}^{\dagger}\mathbf{b}$  stands for different types of linear systems (see [7, 19]):

- (1) If  $\mathbf{A}\mathbf{x} = \mathbf{b}$  is consistent with full-column rank  $\mathbf{A}$ , i.e., rank $(\mathbf{A}) = n$ , then  $\mathbf{A}^{\dagger}\mathbf{b}$  is the unique solution. In this case, we have  $m \ge n$  and the linear system is overdetermined when m > n.
- (2) If  $\mathbf{A}\mathbf{x} = \mathbf{b}$  is consistent with rank $(\mathbf{A}) < n$ , then  $\mathbf{A}^{\dagger}\mathbf{b}$  is the unique minimum  $\ell_2$ -norm solution. In this case, we have  $m \ge n$  or m < n, and the linear system is overdetermined (resp. underdetermined) when m > n (resp. m < n). The matrix  $\mathbf{A}$  can be of full-row rank, i.e., rank $(\mathbf{A}) = m$ , or rank-deficient, i.e., rank $(\mathbf{A}) < m$ .
- (3) If  $\mathbf{A}\mathbf{x} = \mathbf{b}$  is inconsistent with rank $(\mathbf{A}) = n$ , then  $\mathbf{A}^{\dagger}\mathbf{b}$  is the unique least squares solution. In this case, we have  $m \ge n$  and the linear system is overdetermined when m > n.
- (4) If  $\mathbf{A}\mathbf{x} = \mathbf{b}$  is inconsistent with rank $(\mathbf{A}) < n$ , then  $\mathbf{A}^{\dagger}\mathbf{b}$  is the unique minimum  $\ell_2$ -norm least squares solution. In this case, we have  $m \ge n$  or m < n, and the linear system is overdetermined (resp. underdetermined) when m > n (resp. m < n). The matrix  $\mathbf{A}$  can be of full-row rank, i.e., rank $(\mathbf{A}) = m$ , or rank-deficient, i.e., rank $(\mathbf{A}) < m$ .

If  $\mathbf{A}\mathbf{x} = \mathbf{b}$  is inconsistent, Needell [37] showed that RK does not converge to  $\mathbf{A}^{\dagger}\mathbf{b}$ . To resolve this problem, Zouzias and Freris [51] proposed a randomized extended Kaczmarz (REK) algorithm, which uses RK twice [30, 13] at each iteration and exponentially converges in the mean square to  $\mathbf{A}^{\dagger}\mathbf{b}$ . More precisely, assuming that the *j*th column  $\mathbf{A}_{:,j}$  and the *i*th row  $\mathbf{A}_{i,:}$  have been selected at the *k*th iteration, REK generates two vectors  $\mathbf{z}^k$  and  $\mathbf{x}^k$  via two RK updates (one for  $\mathbf{A}^{\mathrm{T}}\mathbf{z} = \mathbf{0}$  from  $\mathbf{z}^{k-1}$  and the other for  $\mathbf{A}\mathbf{x} = \mathbf{b} - \mathbf{z}^k$  from  $\mathbf{x}^{k-1}$ ):

$$\begin{aligned} \mathbf{z}^{k} &= \mathbf{z}^{k-1} - \frac{(\mathbf{A}_{:,j})^{\mathrm{T}} \mathbf{z}^{k-1}}{(\mathbf{A}_{:,j})^{\mathrm{T}} \mathbf{A}_{:,j}} \mathbf{A}_{:,j}, \\ \mathbf{x}^{k} &= \mathbf{x}^{k-1} - \frac{\mathbf{A}_{i,:} \mathbf{x}^{k-1} - \mathbf{b}_{i} + \mathbf{z}_{i}^{k}}{\mathbf{A}_{i,:} (\mathbf{A}_{i,:})^{\mathrm{T}}} (\mathbf{A}_{i,:})^{\mathrm{T}}. \end{aligned}$$

For general linear systems (consistent or inconsistent, full-rank or rank-deficient), the vector  $\mathbf{x}^k$  generated by REK exponentially converges to  $\mathbf{A}^{\dagger}\mathbf{b}$  if  $\mathbf{z}^0 \in \mathbf{b} + \operatorname{range}(\mathbf{A})$  and  $\mathbf{x}^0 \in \operatorname{range}(\mathbf{A}^T)$ [30, 13]. To accelerate the convergence, the following projection-based block variants [39, 40] of RK and REK were developed. For a subset  $\mathcal{I} \subset \{1, 2, \ldots, m\}$  and a subset  $\mathcal{J} \subset \{1, 2, \ldots, n\}$ , denote by  $\mathbf{A}_{\mathcal{I},:}$  and  $\mathbf{A}_{:,\mathcal{J}}$  the row submatrix of  $\mathbf{A}$  indexed by  $\mathcal{I}$  and the column submatrix of  $\mathbf{A}$  indexed by  $\mathcal{J}$ , respectively. Assuming that the subset  $\mathcal{I}_i$  has been selected at the *k*th iteration, the randomized block Kaczmarz (RBK) algorithm [39] generates the *k*th estimate  $\mathbf{x}^k$  via

$$\mathbf{x}^{k} = \mathbf{x}^{k-1} - (\mathbf{A}_{\mathcal{I}_{i},:})^{\dagger} (\mathbf{A}_{\mathcal{I}_{i},:} \mathbf{x}^{k-1} - \mathbf{b}_{\mathcal{I}_{i}}).$$

Assuming that the subsets  $\mathcal{J}_j$  and  $\mathcal{I}_i$  have been selected at the k iteration, the randomized double block Kaczmarz (RDBK) algorithm [40] generates the kth estimate  $\mathbf{x}^k$  via

$$\begin{aligned} \mathbf{z}^{k} &= \mathbf{z}^{k-1} - \mathbf{A}_{:,\mathcal{J}_{j}}(\mathbf{A}_{:,\mathcal{J}_{j}})^{\dagger} \mathbf{z}^{k-1}, \\ \mathbf{x}^{k} &= \mathbf{x}^{k-1} - (\mathbf{A}_{\mathcal{I}_{i},:})^{\dagger} (\mathbf{A}_{\mathcal{I}_{i},:} \mathbf{x}^{k-1} - \mathbf{b}_{\mathcal{I}_{i}} + \mathbf{z}_{\mathcal{I}_{i}}^{k}). \end{aligned}$$

<sup>&</sup>lt;sup>1</sup>Every  $m \times n$  matrix **A** has a unique Moore-Penrose pseudoinverse. In particular, in this paper we will use the following property of the pseudoinverse:  $\mathbf{A}^{\mathrm{T}} = \mathbf{A}^{\mathrm{T}} \mathbf{A} \mathbf{A}^{\dagger}$ .

Numerical experiments demonstrate that the convergence can be significantly accelerated if appropriate blocks of the coefficient matrix are used. The main drawback of projection-based block methods is that they are not adequate for distributed implementations.

Recently, Necoara [35] proposed a randomized average block Kaczmarz (RABK) algorithm for consistent linear systems, which takes a convex combination of several RK updates (i.e., the projections of the current iterate onto several hyperplanes) as a new direction with some stepsize. Assuming that the subset  $\mathcal{I}$  has been selected at the *k*th iteration, RABK generates the *k*th estimate  $\mathbf{x}^k$  via

$$\mathbf{x}^{k} = \mathbf{x}^{k-1} - \alpha_{k} \left( \sum_{i \in \mathcal{I}} \omega_{i}^{k} \frac{\mathbf{A}_{i,:} \mathbf{x}^{k-1} - \mathbf{b}_{i}}{\mathbf{A}_{i,:} (\mathbf{A}_{i,:})^{\mathrm{T}}} (\mathbf{A}_{i,:})^{\mathrm{T}} \right),$$
(1)

where the weights  $\omega_i^k \in [0,1]$  such that  $\sum_{i \in \mathcal{I}} \omega_i^k = 1$ , and the stepsize  $\alpha_k \in (0,2)$ . The convergence analysis reveals that RABK is extremely effective when it is given a good sampling of the rows into well-conditioned blocks. A block version of RABK (i.e., parallel randomized block Kaczmarz), which takes a convex combination of the RBK updates, was proposed and studied by Richtárik and Takáč [47]. Shortly afterwards, Du and Sun [15] proposed a doubly stochastic block Gauss-Seidel (DSBGS) algorithm, which randomly chooses a submatrix of the coefficient matrix at each iteration. Assuming that the subsets  $\mathcal{I}$  and  $\mathcal{J}$  have been selected at the *k*th iteration, DSBGS generates the *k*th estimate  $\mathbf{x}^k$  via

$$\mathbf{x}^{k} = \mathbf{x}^{k-1} - \alpha_{k} \frac{\mathbf{I}_{:,\mathcal{J}} (\mathbf{A}_{\mathcal{I},\mathcal{J}})^{\mathrm{T}} (\mathbf{I}_{:,\mathcal{I}})^{\mathrm{T}}}{\|\mathbf{A}_{\mathcal{I},\mathcal{J}}\|_{\mathrm{F}}^{2}} (\mathbf{A} \mathbf{x}^{k-1} - \mathbf{b}),$$

where **I** denotes the identity matrix,  $\mathbf{A}_{\mathcal{I},\mathcal{J}}$  denotes the submatrix that lies in the rows indexed by  $\mathcal{I}$  and the columns indexed by  $\mathcal{J}$ , and  $\|\cdot\|_{\mathrm{F}}$  is the Frobenius norm. Exponential convergence of DSBGS for consistent linear systems was proved. By setting  $\mathcal{I} \subset \{1, 2, \ldots, m\}$  and  $\mathcal{J} = \{1, 2, \ldots, n\}$ , DSBGS recovers a special case of RABK, i.e., RABK with weight

$$\omega_i^k = \frac{\mathbf{A}_{i,:}(\mathbf{A}_{i,:})^{\mathrm{T}}}{\|\mathbf{A}_{\mathcal{I},:}\|_{\mathrm{F}}^2}, \qquad i \in \mathcal{I}.$$

Note that both RABK and DSBGS are very easy to implement on distributed computing units, yielding remarkable improvements in computational time. We emphasize that convergence results in the mean square of RABK and DSBGS are obtained only for consistent linear systems.

In this paper, based on the REK algorithm and the RABK algorithm, we present a simple randomized extended block Kaczmarz (REBK) algorithm that exponentially converges in the mean square to the unique minimum  $\ell_2$ -norm (least squares) solution of a given general linear system (full-rank or rank-deficient, overdetermined or underdetermined, consistent or inconsistent). Our method is different from those projection-based block methods, for example, those in [18, 1, 8, 43, 39, 40, 16]. At each step, REBK, as a direct extension of REK, uses two special RABK (which also can be viewed as special DSBGS) updates (one for  $\mathbf{A}^{\mathrm{T}}\mathbf{z} = \mathbf{0}$  from  $\mathbf{z}^{k-1}$ and the other for  $\mathbf{A}\mathbf{x} = \mathbf{b} - \mathbf{z}^k$  from  $\mathbf{x}^{k-1}$ ; see Section 2 for details). Compared with REK, REBK usually has a better convergence rate and can exploit the high-level basic linear algebra subroutine (BLAS2), even fast matrix-vector multiplies (for example, if submatrices of  $\mathbf{A}$  have circulant or Toeplitz structures, then the Fast Fourier Transform technique can be used), and therefore could be more efficient. Compared with RDBK, REBK can be implemented on distributed computing units. We refer the reader to [39, 35] for more advantages of block methods. Numerical examples are given to illustrate the efficiency of REBK.

Organization of the paper. In the rest of this section, we give some notation. In Section 2 we describe the randomized extended block Kaczmarz algorithm and prove its convergence

theory. Both the exponential convergence of the norm of the expected error and the exponential convergence of the expected norm of the error are discussed. In Section 3 we report the numerical results. Finally, we present brief concluding remarks in Section 4.

Notation. For any random variable  $\boldsymbol{\xi}$ , let  $\mathbb{E} [\boldsymbol{\xi}]$  denote its expectation. For an integer  $m \geq 1$ , let  $[m] := \{1, 2, 3, \ldots, m\}$ . Lowercase (upper-case) boldface letters are reserved for column vectors (matrices). For any vector  $\mathbf{u} \in \mathbb{R}^m$ , we use  $\mathbf{u}_i$ ,  $\mathbf{u}^T$ , and  $\|\mathbf{u}\|_2$  to denote the *i*th element, the transpose, and the  $\ell_2$ -norm of  $\mathbf{u}$ , respectively. We use  $\mathbf{I}$  to denote the identity matrix whose order is clear from the context. For any matrix  $\mathbf{A} \in \mathbb{R}^{m \times n}$ , we use  $\mathbf{A}^T$ ,  $\mathbf{A}^{\dagger}$ ,  $\|\mathbf{A}\|_{\mathrm{F}}$ , range( $\mathbf{A}$ ),  $\sigma_1(\mathbf{A}) \geq \sigma_2(\mathbf{A}) \geq \cdots \geq \sigma_r(\mathbf{A}) > 0$  to denote the transpose, the Moore-Penrose pseudoinverse, the Frobenius norm, the column space, and all the nonzero singular values of  $\mathbf{A}$ , respectively. Obviously, r is the rank of  $\mathbf{A}$ . For index sets  $\mathcal{I} \subseteq [m]$  and  $\mathcal{J} \subseteq [n]$ , let  $\mathbf{A}_{\mathcal{I},:}$ ,  $\mathbf{A}_{:,\mathcal{J}}$ , and  $\mathbf{A}_{\mathcal{I},\mathcal{J}}$  denote the row submatrix indexed by  $\mathcal{I}$ , the column submatrix indexed by  $\mathcal{J}$ , respectively. We call  $\{\mathcal{I}_1, \mathcal{I}_2, \ldots, \mathcal{I}_s\}$  a partition of [m] if  $\mathcal{I}_i \cap \mathcal{I}_j = \emptyset$  for  $i \neq j$  and  $\cup_{i=1}^s \mathcal{I}_i = [m]$ . Similarly,  $\{\mathcal{J}_1, \mathcal{J}_2, \ldots, \mathcal{J}_t\}$  denotes a partition of [n] if  $\mathcal{J}_i \cap \mathcal{J}_j = \emptyset$  for  $i \neq j$  and  $\cup_{j=1}^s \mathcal{J}_j = [n]$ . We use  $|\mathcal{I}|$  to denote the cardinality of a set  $\mathcal{I} \subseteq [m]$ .

# 2 The randomized extended block Kaczmarz algorithm

In this section, based on given partitions of [m] and [n], we propose the following randomized extended block Kaczmarz algorithm (see Algorithm 1) for solving consistent or inconsistent linear systems. We emphasize that this algorithm can be implemented on distributed computing units.

 $\begin{array}{l} \textbf{Algorithm 1: Randomized extended block Kaczmarz (REBK)} \\ \hline \textbf{Let } \{\mathcal{I}_1, \mathcal{I}_2, \dots, \mathcal{I}_s\} \text{ and } \{\mathcal{J}_1, \mathcal{J}_2, \dots, \mathcal{J}_t\} \text{ be partitions of } [m] \text{ and } [n], \text{ respectively.} \\ \hline \textbf{Let } \alpha > 0. \text{ Initialize } \mathbf{z}^0 \in \mathbb{R}^m \text{ and } \mathbf{x}^0 \in \mathbb{R}^n. \\ \textbf{for } k = 1, 2, \dots, \textbf{do} \\ & \text{Pick } j \in [t] \text{ with probability } \|\mathbf{A}_{:,\mathcal{J}_j}\|_{\mathrm{F}}^2 / \|\mathbf{A}\|_{\mathrm{F}}^2 \\ & \text{Set } \mathbf{z}^k = \mathbf{z}^{k-1} - \frac{\alpha}{\|\mathbf{A}_{:,\mathcal{J}_j}\|_{\mathrm{F}}^2} \mathbf{A}_{:,\mathcal{J}_j} (\mathbf{A}_{:,\mathcal{J}_j})^{\mathrm{T}} \mathbf{z}^{k-1} \\ & \text{Pick } i \in [s] \text{ with probability } \|\mathbf{A}_{\mathcal{I}_i,:}\|_{\mathrm{F}}^2 / \|\mathbf{A}\|_{\mathrm{F}}^2 \\ & \text{Set } \mathbf{x}^k = \mathbf{x}^{k-1} - \frac{\alpha}{\|\mathbf{A}_{\mathcal{I}_i,:}\|_{\mathrm{F}}^2} (\mathbf{A}_{\mathcal{I}_i,:})^{\mathrm{T}} (\mathbf{A}_{\mathcal{I}_i,:}\mathbf{x}^{k-1} - \mathbf{b}_{\mathcal{I}_i} + \mathbf{z}_{\mathcal{I}_i}^k) \end{array}$ 

Here we only consider constant stepsize for simplicity. By choosing the row partition parameter s = m, the column partition parameter t = n, and the stepsize  $\alpha = 1$ , we recover the well-known randomized extended Kaczmarz algorithm of Zouzias and Freris [51]. REBK uses two RABK updates (see (1)) at each step:

• RABK update for  $\mathbf{A}^{\mathrm{T}}\mathbf{z} = \mathbf{0}$  from  $\mathbf{z}^{k-1}$ 

$$\mathbf{z}^{k} = \mathbf{z}^{k-1} - \alpha \left( \sum_{l \in \mathcal{J}_{j}} \omega_{l}^{k} \frac{(\mathbf{A}_{:,l})^{\mathrm{T}} \mathbf{z}^{k-1}}{(\mathbf{A}_{:,l})^{\mathrm{T}} \mathbf{A}_{:,l}} \mathbf{A}_{:,l} \right), \qquad \omega_{l}^{k} = \frac{(\mathbf{A}_{:,l})^{\mathrm{T}} \mathbf{A}_{:,l}}{\|\mathbf{A}_{:,\mathcal{J}_{j}}\|_{\mathrm{F}}^{2}};$$

• RABK update for  $\mathbf{A}\mathbf{x} = \mathbf{b} - \mathbf{z}^k$  from  $\mathbf{x}^{k-1}$ 

$$\mathbf{x}^{k} = \mathbf{x}^{k-1} - \alpha \left( \sum_{l \in \mathcal{I}_{i}} \omega_{l}^{k} \frac{\mathbf{A}_{l,:} \mathbf{x}^{k-1} - \mathbf{b}_{l} + \mathbf{z}_{l}^{k}}{\mathbf{A}_{l,:} (\mathbf{A}_{l,:})^{\mathrm{T}}} (\mathbf{A}_{l,:})^{\mathrm{T}} \right), \qquad \omega_{l}^{k} = \frac{\mathbf{A}_{l,:} (\mathbf{A}_{l,:})^{\mathrm{T}}}{\|\mathbf{A}_{\mathcal{I}_{i},:}\|_{\mathrm{F}}^{2}}$$

We note that if  $\mathbf{z}^0 = \mathbf{0}$  in REBK, then all  $\mathbf{z}^k \equiv \mathbf{0}$ , which yields the update of  $\mathbf{x}^k$  is exactly the same as that of RABK.

Before proving the convergence theory of REBK for general linear systems, we give the following notation. Let  $\mathbb{E}_{k-1}[\cdot]$  denote the conditional expectation conditioned on the first k-1 iterations of REBK. That is,

$$\mathbb{E}_{k-1}\left[\cdot\right] = \mathbb{E}\left[\cdot|j_1, i_1, j_2, i_2, \dots, j_{k-1}, i_{k-1}\right],$$

where  $j_l$  is the *l*th column block chosen and  $i_l$  is the *l*th row block chosen. We denote the conditional expectation conditioned on the first k - 1 iterations and the *k*th column block chosen as

$$\mathbb{E}_{k-1}^{i}\left[\cdot\right] = \mathbb{E}\left[\cdot|j_1, i_1, j_2, i_2, \dots, j_{k-1}, i_{k-1}, j_k\right].$$

Then by the law of total expectation we have

$$\mathbb{E}_{k-1}\left[\cdot\right] = \mathbb{E}_{k-1}\left[\mathbb{E}_{k-1}^{i}\left[\cdot\right]\right].$$

#### 2.1 The exponential convergence of the norm of the expected error

In this subsection we show the exponential convergence of the norm of the expected error, i.e.,

$$\|\mathbb{E}\left[\mathbf{x}^{k}-\mathbf{A}^{\dagger}\mathbf{b}\right]\|_{2}.$$

The convergence of the norm of the expected error depends on the positive number  $\delta$  defined as

$$\delta := \max_{1 \le i \le r} \left| 1 - \frac{\alpha \sigma_i^2(\mathbf{A})}{\|\mathbf{A}\|_{\mathrm{F}}^2} \right|.$$

The following lemma will be used and its proof is straightforward (e.g., via the singular value decomposition).

**Lemma 1.** Let  $\alpha > 0$  and  $\mathbf{A} \in \mathbb{R}^{m \times n}$  be any nonzero real matrix with rank $(\mathbf{A}) = r$ . For every  $\mathbf{u} \in \text{range}(\mathbf{A}^{T})$ , it holds

$$\left\| \left( \mathbf{I} - \alpha \frac{\mathbf{A}^{\mathrm{T}} \mathbf{A}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}} \right)^{k} \mathbf{u} \right\|_{2} \leq \delta^{k} \|\mathbf{u}\|_{2}.$$

We give the convergence of the norm of the expected error of REBK in the following theorem.

**Theorem 2.** For any given consistent or inconsistent linear system  $\mathbf{A}\mathbf{x} = \mathbf{b}$ , let  $\mathbf{x}^k$  be the kth iterate of REBK with  $\mathbf{z}^0 \in \mathbb{R}^m$  and  $\mathbf{x}^0 \in \text{range}(\mathbf{A}^T)$ . It holds

$$\|\mathbb{E}\left[\mathbf{x}^{k}-\mathbf{A}^{\dagger}\mathbf{b}\right]\|_{2} \leq \delta^{k}\left(\|\mathbf{x}^{0}-\mathbf{A}^{\dagger}\mathbf{b}\|_{2}+\frac{\alpha k\|\mathbf{A}^{\mathrm{T}}\mathbf{z}^{0}\|_{2}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\right).$$

Proof. Note that

$$\mathbb{E}_{k-1}\left[\mathbf{z}^{k}\right] = \mathbf{z}^{k-1} - \mathbb{E}_{k-1}\left[\frac{\alpha}{\|\mathbf{A}_{:,\mathcal{J}_{j}}\|_{\mathrm{F}}^{2}}\mathbf{A}_{:,\mathcal{J}_{j}}(\mathbf{A}_{:,\mathcal{J}_{j}})^{\mathrm{T}}\right]\mathbf{z}^{k-1}$$
$$= \left(\mathbf{I} - \alpha \frac{\mathbf{A}\mathbf{A}^{\mathrm{T}}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\right)\mathbf{z}^{k-1},$$

and therefore

$$\mathbb{E}\left[\mathbf{z}^{k}\right] = \mathbb{E}\left[\mathbb{E}_{k-1}\left[\mathbf{z}^{k}\right]\right] = \left(\mathbf{I} - \alpha \frac{\mathbf{A}\mathbf{A}^{\mathrm{T}}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\right) \mathbb{E}\left[\mathbf{z}^{k-1}\right] = \left(\mathbf{I} - \alpha \frac{\mathbf{A}\mathbf{A}^{\mathrm{T}}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\right)^{k} \mathbf{z}^{0}.$$

By  $\mathbf{A}^{\mathrm{T}}\mathbf{b} = \mathbf{A}^{\mathrm{T}}\mathbf{A}\mathbf{A}^{\dagger}\mathbf{b}$ , we have

$$\begin{split} \mathbb{E}_{k-1} \left[ \mathbf{x}^{k} - \mathbf{A}^{\dagger} \mathbf{b} \right] &= \mathbb{E}_{k-1} \left[ \mathbb{E}_{k-1}^{i} \left[ \mathbf{x}^{k} - \mathbf{A}^{\dagger} \mathbf{b} \right] \right] \\ &= \mathbb{E}_{k-1} \left[ \mathbb{E}_{k-1}^{i} \left[ \mathbf{x}^{k-1} - \mathbf{A}^{\dagger} \mathbf{b} - \frac{\alpha}{\|\mathbf{A}_{\mathcal{I}_{i},:}\|_{\mathrm{F}}^{2}} (\mathbf{A}_{\mathcal{I}_{i},:})^{\mathrm{T}} (\mathbf{A}_{\mathcal{I}_{i},:} \mathbf{x}^{k-1} - \mathbf{b}_{\mathcal{I}_{i}} + \mathbf{z}_{\mathcal{I}_{i}}^{k}) \right] \right] \\ &= \mathbb{E}_{k-1} \left[ \mathbf{x}^{k-1} - \mathbf{A}^{\dagger} \mathbf{b} - \alpha \frac{\mathbf{A}^{\mathrm{T}} (\mathbf{A} \mathbf{x}^{k-1} - \mathbf{b} + \mathbf{z}^{k})}{\|\mathbf{A}\|_{\mathrm{F}}^{2}} \right] \\ &= \mathbf{x}^{k-1} - \mathbf{A}^{\dagger} \mathbf{b} - \alpha \frac{\mathbf{A}^{\mathrm{T}} \mathbf{A} \mathbf{x}^{k-1} - \mathbf{A}^{\mathrm{T}} \mathbf{b}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}} - \alpha \frac{\mathbf{A}^{\mathrm{T}} }{\|\mathbf{A}\|_{\mathrm{F}}^{2}} \mathbb{E}_{k-1} \left[ \mathbf{z}^{k} \right] \\ &= \mathbf{x}^{k-1} - \mathbf{A}^{\dagger} \mathbf{b} - \alpha \frac{\mathbf{A}^{\mathrm{T}} \mathbf{A} \mathbf{x}^{k-1} - \mathbf{A}^{\mathrm{T}} \mathbf{A} \mathbf{A}^{\dagger} \mathbf{b}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}} - \alpha \frac{\mathbf{A}^{\mathrm{T}} }{\|\mathbf{A}\|_{\mathrm{F}}^{2}} \mathbb{E}_{k-1} \left[ \mathbf{z}^{k} \right] \\ &= \left( \mathbf{I} - \alpha \frac{\mathbf{A}^{\mathrm{T}} \mathbf{A}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}} \right) (\mathbf{x}^{k-1} - \mathbf{A}^{\dagger} \mathbf{b}) - \alpha \frac{\mathbf{A}^{\mathrm{T}} }{\|\mathbf{A}\|_{\mathrm{F}}^{2}} \left( \mathbf{I} - \alpha \frac{\mathbf{A} \mathbf{A}^{\mathrm{T}}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}} \right) \mathbf{z}^{k-1}. \end{split}$$

Taking expectation gives

$$\begin{split} \mathbb{E}\left[\mathbf{x}^{k}-\mathbf{A}^{\dagger}\mathbf{b}\right] &= \mathbb{E}\left[\mathbb{E}_{k-1}\left[\mathbf{x}^{k}-\mathbf{A}^{\dagger}\mathbf{b}\right]\right] \\ &= \left(\mathbf{I}-\alpha\frac{\mathbf{A}^{\mathrm{T}}\mathbf{A}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\right)\mathbb{E}\left[\mathbf{x}^{k-1}-\mathbf{A}^{\dagger}\mathbf{b}\right] - \alpha\frac{\mathbf{A}^{\mathrm{T}}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\left(\mathbf{I}-\alpha\frac{\mathbf{A}\mathbf{A}^{\mathrm{T}}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\right)\mathbb{E}\left[\mathbf{z}^{k-1}\right] \\ &= \left(\mathbf{I}-\alpha\frac{\mathbf{A}^{\mathrm{T}}\mathbf{A}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\right)\mathbb{E}\left[\mathbf{x}^{k-1}-\mathbf{A}^{\dagger}\mathbf{b}\right] - \alpha\frac{\mathbf{A}^{\mathrm{T}}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\left(\mathbf{I}-\alpha\frac{\mathbf{A}\mathbf{A}^{\mathrm{T}}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\right)^{k}\mathbf{z}^{0} \\ &= \left(\mathbf{I}-\alpha\frac{\mathbf{A}^{\mathrm{T}}\mathbf{A}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\right)\mathbb{E}\left[\mathbf{x}^{k-1}-\mathbf{A}^{\dagger}\mathbf{b}\right] - \alpha\left(\mathbf{I}-\alpha\frac{\mathbf{A}^{\mathrm{T}}\mathbf{A}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\right)^{k}\frac{\mathbf{A}^{\mathrm{T}}\mathbf{z}^{0}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}} \\ &= \left(\mathbf{I}-\alpha\frac{\mathbf{A}^{\mathrm{T}}\mathbf{A}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\right)^{2}\mathbb{E}\left[\mathbf{x}^{k-2}-\mathbf{A}^{\dagger}\mathbf{b}\right] - 2\alpha\left(\mathbf{I}-\alpha\frac{\mathbf{A}^{\mathrm{T}}\mathbf{A}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\right)^{k}\frac{\mathbf{A}^{\mathrm{T}}\mathbf{z}^{0}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}} \\ &= \cdots \\ &= \left(\mathbf{I}-\alpha\frac{\mathbf{A}^{\mathrm{T}}\mathbf{A}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\right)^{k}\left(\mathbf{x}^{0}-\mathbf{A}^{\dagger}\mathbf{b}\right) - \alpha k\left(\mathbf{I}-\alpha\frac{\mathbf{A}^{\mathrm{T}}\mathbf{A}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\right)^{k}\frac{\mathbf{A}^{\mathrm{T}}\mathbf{z}^{0}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}. \end{split}$$

Applying the norms to both sides we obtain

$$\begin{split} \|\mathbb{E}\left[\mathbf{x}^{k}-\mathbf{A}^{\dagger}\mathbf{b}\right]\|_{2} &= \left\|\left(\mathbf{I}-\alpha\frac{\mathbf{A}^{\mathrm{T}}\mathbf{A}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\right)^{k}\left(\mathbf{x}^{0}-\mathbf{A}^{\dagger}\mathbf{b}\right)-\alpha k\left(\mathbf{I}-\alpha\frac{\mathbf{A}^{\mathrm{T}}\mathbf{A}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\right)^{k}\frac{\mathbf{A}^{\mathrm{T}}\mathbf{z}^{0}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\right\|_{2} \\ &\leq \left\|\left(\mathbf{I}-\alpha\frac{\mathbf{A}^{\mathrm{T}}\mathbf{A}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\right)^{k}\left(\mathbf{x}^{0}-\mathbf{A}^{\dagger}\mathbf{b}\right)\right\|_{2}+\left\|\alpha k\left(\mathbf{I}-\alpha\frac{\mathbf{A}^{\mathrm{T}}\mathbf{A}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\right)^{k}\frac{\mathbf{A}^{\mathrm{T}}\mathbf{z}^{0}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\right\|_{2} \\ &\leq \delta^{k}\left(\|\mathbf{x}^{0}-\mathbf{A}^{\dagger}\mathbf{b}\|_{2}+\frac{\alpha k\|\mathbf{A}^{\mathrm{T}}\mathbf{z}^{0}\|_{2}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\right). \end{split}$$

Here the last inequality follows from the fact that  $\mathbf{x}^0 \in \operatorname{range}(\mathbf{A}^T)$ ,  $\mathbf{A}^{\dagger}\mathbf{b} \in \operatorname{range}(\mathbf{A}^T)$ ,  $\mathbf{A}^T\mathbf{z}^0 \in \operatorname{range}(\mathbf{A}^T)$ , and Lemma 1.

Remark 3. To ensure convergence of the expected error, it suffices to have

$$\delta = \max_{1 \le i \le r} \left| 1 - \frac{\alpha \sigma_i^2(\mathbf{A})}{\|\mathbf{A}\|_{\mathrm{F}}^2} \right| < 1,$$

which implies

$$0 < \alpha < \frac{2 \|\mathbf{A}\|_{\mathrm{F}}^2}{\sigma_1^2(\mathbf{A})}.$$

The optimal  $\alpha$  in Theorem 2 is (see [42])

$$\frac{2\|\mathbf{A}\|_{\mathrm{F}}^2}{\sigma_1^2(\mathbf{A}) + \sigma_r^2(\mathbf{A})} = \operatorname*{argmin}_{0 < \alpha < \frac{2\|\mathbf{A}\|_{\mathrm{F}}^2}{\sigma_1^2(\mathbf{A})}} \max_{1 \le i \le r} \left| 1 - \frac{\alpha \sigma_i^2(\mathbf{A})}{\|\mathbf{A}\|_{\mathrm{F}}^2} \right|.$$

and the corresponding convergence rate  $\delta$  is

$$\frac{\sigma_1^2(\mathbf{A}) - \sigma_r^2(\mathbf{A})}{\sigma_1^2(\mathbf{A}) + \sigma_r^2(\mathbf{A})}$$

### 2.2 The exponential convergence of the expected norm of the error

In this subsection we show the exponential convergence of the expected norm of the error, i.e.,

$$\mathbb{E}\left[\|\mathbf{x}^k - \mathbf{A}^\dagger \mathbf{b}\|_2^2\right].$$

The convergence of the expected norm of the error depends on the positive numbers  $\eta$  and  $\rho$  defined as

$$\eta := 1 - \frac{(2\alpha - \alpha^2 \beta_{\max}^{\mathcal{I}}) \sigma_r^2(\mathbf{A})}{\|\mathbf{A}\|_{\mathrm{F}}^2}, \qquad \rho := 1 - \frac{(2\alpha - \alpha^2 \beta_{\max}^{\mathcal{J}}) \sigma_r^2(\mathbf{A})}{\|\mathbf{A}\|_{\mathrm{F}}^2},$$

where

$$\beta_{\max}^{\mathcal{I}} := \max_{i \in [s]} \frac{\|\mathbf{A}_{\mathcal{I}_i,:}\|_2^2}{\|\mathbf{A}_{\mathcal{I}_i,:}\|_{\mathrm{F}}^2}, \qquad \beta_{\max}^{\mathcal{J}} := \max_{j \in [t]} \frac{\|\mathbf{A}_{:,\mathcal{J}_j}\|_2^2}{\|\mathbf{A}_{:,\mathcal{J}_j}\|_{\mathrm{F}}^2}$$

The following lemmas will be used extensively in this paper.

**Lemma 4.** Let  $\mathbf{A} \in \mathbb{R}^{m \times n}$  be any nonzero real matrix with  $\operatorname{rank}(\mathbf{A}) = r$ . For every  $\mathbf{u} \in \operatorname{range}(\mathbf{A})$ , it holds

$$\|\mathbf{A}^{\mathrm{T}}\mathbf{u}\|_2^2 \geq \sigma_r^2(\mathbf{A})\|\mathbf{u}\|_2^2.$$

**Lemma 5.** Let  $\mathbf{A} \in \mathbb{R}^{m \times n}$  be any nonzero real matrix. For every  $\mathbf{u} \in \mathbb{R}^m$ , it holds

$$\mathbf{u}^{\mathrm{T}}(\mathbf{A}\mathbf{A}^{\mathrm{T}})^{2}\mathbf{u} \leq \|\mathbf{A}\|_{2}^{2}\|\mathbf{A}^{\mathrm{T}}\mathbf{u}\|_{2}^{2}.$$

The proof of Lemma 4 is straightforward (e.g., via the singular value decomposition), and Lemma 5 follows from

$$\mathbf{u}^{\mathrm{T}}(\mathbf{A}\mathbf{A}^{\mathrm{T}})^{2}\mathbf{u} = \mathbf{u}^{\mathrm{T}}\mathbf{A}(\mathbf{A}^{\mathrm{T}}\mathbf{A})\mathbf{A}^{\mathrm{T}}\mathbf{u} \leq \|\mathbf{A}^{\mathrm{T}}\mathbf{A}\|_{2}\|\mathbf{A}^{\mathrm{T}}\mathbf{u}\|_{2}^{2} = \|\mathbf{A}\|_{2}^{2}\|\mathbf{A}^{\mathrm{T}}\mathbf{u}\|_{2}^{2}.$$

In the following lemma we show that the vector  $\mathbf{z}^k$  generated in REBK with  $\mathbf{z}^0 \in \mathbf{b} + \text{range}(\mathbf{A})$  converges to

$$\mathbf{b}_{\perp} =: (\mathbf{I} - \mathbf{A}\mathbf{A}^{\dagger})\mathbf{b},$$

which is the orthogonal projection of  $\mathbf{z}^0$  onto the set  $\{\mathbf{z} \mid \mathbf{A}^T \mathbf{z} = \mathbf{0}\}$ .

**Lemma 6.** For any given consistent or inconsistent linear system  $\mathbf{A}\mathbf{x} = \mathbf{b}$ , let  $\mathbf{z}^k$  be the vector generated in REBK with  $\mathbf{z}^0 \in \mathbf{b} + \text{range}(\mathbf{A})$ . Assume  $0 < \alpha < 2/\beta_{\text{max}}^{\mathcal{J}}$ . It holds

$$\mathbb{E}\left[\|\mathbf{z}^k - \mathbf{b}_{\perp}\|_2^2\right] \le \rho^k \|\mathbf{z}^0 - \mathbf{b}_{\perp}\|_2^2.$$

*Proof.* By  $(\mathbf{A}_{:,\mathcal{J}_j})^{\mathrm{T}}\mathbf{b}_{\perp} = \mathbf{0}$ , we have

$$\mathbf{z}^{k} - \mathbf{b}_{\perp} = \mathbf{z}^{k-1} - \mathbf{b}_{\perp} - \frac{\alpha}{\|\mathbf{A}_{:,\mathcal{J}_{j}}\|_{\mathrm{F}}^{2}} \mathbf{A}_{:,\mathcal{J}_{j}} (\mathbf{A}_{:,\mathcal{J}_{j}})^{\mathrm{T}} (\mathbf{z}^{k-1} - \mathbf{b}_{\perp}).$$
(2)

By  $\mathbf{z}^0 - \mathbf{b}_{\perp} = \mathbf{A}\mathbf{A}^{\dagger}\mathbf{z}^0 \in \operatorname{range}(\mathbf{A})$  and  $\mathbf{A}_{:,\mathcal{J}_j}(\mathbf{A}_{:,\mathcal{J}_j})^{\mathrm{T}}(\mathbf{z}^{k-1} - \mathbf{b}_{\perp}) \in \operatorname{range}(\mathbf{A})$ , we can show that  $\mathbf{z}^k - \mathbf{b}_{\perp} \in \operatorname{range}(\mathbf{A})$  by induction. It follows from (2) that

Taking the conditioned expectation on the first k-1 iterations yields

Taking expectation again gives

$$\mathbb{E}\left[\|\mathbf{z}^{k} - \mathbf{b}_{\perp}\|_{2}^{2}\right] = \mathbb{E}\left[\mathbb{E}_{k-1}\left[\|\mathbf{z}^{k} - \mathbf{b}_{\perp}\|_{2}^{2}\right]\right]$$
$$\leq \rho \mathbb{E}\left[\|\mathbf{z}^{k-1} - \mathbf{b}_{\perp}\|_{2}^{2}\right]$$
$$\leq \rho^{k} \|\mathbf{z}^{0} - \mathbf{b}_{\perp}\|_{2}^{2}.$$

This completes the proof.

We give the main convergence result of REBK in the following theorem.

**Theorem 7.** For any given consistent or inconsistent linear system  $\mathbf{A}\mathbf{x} = \mathbf{b}$ , let  $\mathbf{x}^k$  be the kth iterate of REBK with  $\mathbf{z}^0 \in \mathbf{b} + \operatorname{range}(\mathbf{A})$  and  $\mathbf{x}^0 \in \operatorname{range}(\mathbf{A}^T)$ . Assume that  $0 < \alpha < 2/\max(\beta_{\max}^{\mathcal{I}}, \beta_{\max}^{\mathcal{J}})$ . For any  $\varepsilon > 0$ , it holds

$$\mathbb{E}\left[\|\mathbf{x}^{k}-\mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2}\right] \leq (1+\varepsilon)^{k}\eta^{k}\|\mathbf{x}^{0}-\mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2} + \left(1+\frac{1}{\varepsilon}\right)\frac{\alpha^{2}\beta_{\max}^{\mathcal{I}}\|\mathbf{z}^{0}-\mathbf{b}_{\perp}\|_{2}^{2}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\sum_{l=0}^{k-1}\rho^{k-l}(1+\varepsilon)^{l}\eta^{l}.$$

*Proof.* Let

$$\widehat{\mathbf{x}}^{k} = \mathbf{x}^{k-1} - \frac{\alpha}{\|\mathbf{A}_{\mathcal{I}_{i},:}\|_{\mathrm{F}}^{2}} (\mathbf{A}_{\mathcal{I}_{i},:})^{\mathrm{T}} \mathbf{A}_{\mathcal{I}_{i},:} (\mathbf{x}^{k-1} - \mathbf{A}^{\dagger} \mathbf{b}),$$

which is actually one RABK update for the linear system  $\mathbf{A}\mathbf{x} = \mathbf{A}\mathbf{A}^{\dagger}\mathbf{b}$  from  $\mathbf{x}^{k-1}$ . It follows from

$$\mathbf{x}^{k} - \widehat{\mathbf{x}}^{k} = \frac{\alpha}{\|\mathbf{A}_{\mathcal{I}_{i},:}\|_{\mathrm{F}}^{2}} (\mathbf{A}_{\mathcal{I}_{i},:})^{\mathrm{T}} (\mathbf{b}_{\mathcal{I}_{i}} - \mathbf{A}_{\mathcal{I}_{i},:} \mathbf{A}^{\dagger} \mathbf{b} - \mathbf{z}_{\mathcal{I}_{i}}^{k})$$

that

$$\|\mathbf{x}^{k} - \widehat{\mathbf{x}}^{k}\|_{2}^{2} = \frac{\alpha^{2}}{\|\mathbf{A}_{\mathcal{I}_{i}:}\|_{\mathrm{F}}^{4}} \|(\mathbf{A}_{\mathcal{I}_{i}:})^{\mathrm{T}}(\mathbf{b}_{\mathcal{I}_{i}} - \mathbf{A}_{\mathcal{I}_{i}:}\mathbf{A}^{\dagger}\mathbf{b} - \mathbf{z}_{\mathcal{I}_{i}}^{k})\|_{2}^{2}$$

$$\leq \frac{\alpha^{2}}{\|\mathbf{A}_{\mathcal{I}_{i}:}\|_{\mathrm{F}}^{2}} \frac{\|\mathbf{A}_{\mathcal{I}_{i}:}\|_{2}^{2}}{\|\mathbf{A}_{\mathcal{I}_{i}:}\|_{\mathrm{F}}^{2}} \|\mathbf{b}_{\mathcal{I}_{i}} - \mathbf{A}_{\mathcal{I}_{i}:}\mathbf{A}^{\dagger}\mathbf{b} - \mathbf{z}_{\mathcal{I}_{i}}^{k}\|_{2}^{2}$$

$$\leq \frac{\alpha^{2}\beta_{\max}^{\mathcal{I}}}{\|\mathbf{A}_{\mathcal{I}_{i}:}\|_{\mathrm{F}}^{2}} \|\mathbf{b}_{\mathcal{I}_{i}} - \mathbf{A}_{\mathcal{I}_{i}:}\mathbf{A}^{\dagger}\mathbf{b} - \mathbf{z}_{\mathcal{I}_{i}}^{k}\|_{2}^{2}.$$
(3)

It follows from

$$\begin{split} \mathbb{E}_{k-1}\left[\|\mathbf{x}^{k}-\widehat{\mathbf{x}}^{k}\|_{2}^{2}\right] &= \mathbb{E}_{k-1}\left[\mathbb{E}_{k-1}^{i}\left[\|\mathbf{x}^{k}-\widehat{\mathbf{x}}^{k}\|_{2}^{2}\right]\right] \\ &\leq \mathbb{E}_{k-1}\left[\mathbb{E}_{k-1}^{i}\left[\frac{\alpha^{2}\beta_{\max}^{\mathcal{I}}}{\|\mathbf{A}_{\mathcal{I}_{i},:}\|_{\mathrm{F}}^{2}}\|\mathbf{b}_{\mathcal{I}_{i}}-\mathbf{A}_{\mathcal{I}_{i},:}\mathbf{A}^{\dagger}\mathbf{b}-\mathbf{z}_{\mathcal{I}_{i}}^{k}\|_{2}^{2}\right]\right] \qquad (by (3)) \\ &= \mathbb{E}_{k-1}\left[\frac{\alpha^{2}\beta_{\max}^{\mathcal{I}}\|\mathbf{b}-\mathbf{A}\mathbf{A}^{\dagger}\mathbf{b}-\mathbf{z}^{k}\|_{2}^{2}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\right] \end{split}$$

that

$$\mathbb{E}\left[\|\mathbf{x}^{k} - \widehat{\mathbf{x}}^{k}\|_{2}^{2}\right] = \mathbb{E}\left[\mathbb{E}_{k-1}\left[\|\mathbf{x}^{k} - \widehat{\mathbf{x}}^{k}\|_{2}^{2}\right]\right]$$

$$\leq \frac{\alpha^{2}\beta_{\max}^{\mathcal{I}}}{\|\mathbf{A}\|_{F}^{2}} \mathbb{E}\left[\|\mathbf{b} - \mathbf{A}\mathbf{A}^{\dagger}\mathbf{b} - \mathbf{z}^{k}\|_{2}^{2}\right]$$

$$\leq \frac{\alpha^{2}\beta_{\max}^{\mathcal{I}}\rho^{k}}{\|\mathbf{A}\|_{F}^{2}}\|\mathbf{z}^{0} - \mathbf{b}_{\perp}\|_{2}^{2}. \quad (by \text{ Lemma } 6)$$
(4)

By  $\mathbf{x}^0 \in \operatorname{range}(\mathbf{A}^{\mathrm{T}}), \ \mathbf{A}^{\dagger}\mathbf{b} \in \operatorname{range}(\mathbf{A}^{\mathrm{T}}), \ (\mathbf{A}_{\mathcal{I}_i,:})^{\mathrm{T}}(\mathbf{A}_{\mathcal{I}_i,:}\mathbf{x}^{k-1} - \mathbf{b}_{\mathcal{I}_i} + \mathbf{z}_{\mathcal{I}_i}^k) \in \operatorname{range}(\mathbf{A}^{\mathrm{T}}), \text{ and}$ 

$$\mathbf{x}^{k} - \mathbf{A}^{\dagger} \mathbf{b} = \mathbf{x}^{k-1} - \mathbf{A}^{\dagger} \mathbf{b} - \frac{\alpha}{\|\mathbf{A}_{\mathcal{I}_{i},:}\|_{\mathrm{F}}^{2}} (\mathbf{A}_{\mathcal{I}_{i},:})^{\mathrm{T}} (\mathbf{A}_{\mathcal{I}_{i},:} \mathbf{x}^{k-1} - \mathbf{b}_{\mathcal{I}_{i}} + \mathbf{z}_{\mathcal{I}_{i}}^{k}),$$

we can show that  $\mathbf{x}^k - \mathbf{A}^{\dagger} \mathbf{b} \in \text{range}(\mathbf{A}^{\mathrm{T}})$  by induction. By

$$\begin{aligned} \|\widehat{\mathbf{x}}^{k} - \mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2} &= \|\mathbf{x}^{k-1} - \mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2} - \frac{2\alpha \|\mathbf{A}_{\mathcal{I}_{i}::}(\mathbf{x}^{k-1} - \mathbf{A}^{\dagger}\mathbf{b})\|_{2}^{2}}{\|\mathbf{A}_{\mathcal{I}_{i}::}\|_{\mathrm{F}}^{2}} \\ &+ \alpha^{2}(\mathbf{x}^{k-1} - \mathbf{A}^{\dagger}\mathbf{b})^{\mathrm{T}} \left( \left( \frac{\mathbf{A}_{\mathcal{I}_{i}:}}{\|\mathbf{A}_{\mathcal{I}_{i}::}\|_{\mathrm{F}}} \right)^{\mathrm{T}} \frac{\mathbf{A}_{\mathcal{I}_{i}::}}{\|\mathbf{A}_{\mathcal{I}_{i}::}\|_{\mathrm{F}}} \right)^{2} (\mathbf{x}^{k-1} - \mathbf{A}^{\dagger}\mathbf{b}) \\ &\leq \|\mathbf{x}^{k-1} - \mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2} - \frac{(2\alpha - \alpha^{2}\beta_{\max}^{\mathcal{I}})\|\mathbf{A}_{\mathcal{I}_{i}::}(\mathbf{x}^{k-1} - \mathbf{A}^{\dagger}\mathbf{b})\|_{2}^{2}}{\|\mathbf{A}_{\mathcal{I}_{i}::}\|_{\mathrm{F}}^{2}}, \quad (\text{by Lemma 5}) \end{aligned}$$

we have

which yields

$$\mathbb{E}\left[\|\widehat{\mathbf{x}}^{k} - \mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2}\right] \leq \eta \mathbb{E}\left[\|\mathbf{x}^{k-1} - \mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2}\right].$$
(5)

Note that for any  $\varepsilon > 0$ , we have

$$\begin{aligned} \|\mathbf{x}^{k} - \mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2} &= \|\mathbf{x}^{k} - \widehat{\mathbf{x}}^{k} + \widehat{\mathbf{x}}^{k} - \mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2} \\ &\leq (\|\mathbf{x}^{k} - \widehat{\mathbf{x}}^{k}\|_{2} + \|\widehat{\mathbf{x}}^{k} - \mathbf{A}^{\dagger}\mathbf{b}\|_{2})^{2} \\ &\leq \|\mathbf{x}^{k} - \widehat{\mathbf{x}}^{k}\|_{2}^{2} + \|\widehat{\mathbf{x}}^{k} - \mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2} + 2\|\mathbf{x}^{k} - \widehat{\mathbf{x}}^{k}\|_{2}\|\widehat{\mathbf{x}}^{k} - \mathbf{A}^{\dagger}\mathbf{b}\|_{2} \\ &\leq \left(1 + \frac{1}{\varepsilon}\right)\|\mathbf{x}^{k} - \widehat{\mathbf{x}}^{k}\|_{2}^{2} + (1 + \varepsilon)\|\widehat{\mathbf{x}}^{k} - \mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2}. \end{aligned}$$
(6)

Combining (4), (5), and (6) yields

$$\begin{split} \mathbb{E}\left[\|\mathbf{x}^{k}-\mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2}\right] &\leq \left(1+\frac{1}{\varepsilon}\right) \mathbb{E}\left[\|\mathbf{x}^{k}-\widehat{\mathbf{x}}^{k}\|_{2}^{2}\right] + (1+\varepsilon)\mathbb{E}\left[\|\widehat{\mathbf{x}}^{k}-\mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2}\right] \\ &\leq \left(1+\frac{1}{\varepsilon}\right) \frac{\alpha^{2}\beta_{\max}^{\mathcal{I}}\rho^{k}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}} \|\mathbf{z}^{0}-\mathbf{b}_{\perp}\|_{2}^{2} + (1+\varepsilon)\eta\mathbb{E}\left[\|\mathbf{x}^{k-1}-\mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2}\right] \\ &\leq \left(1+\frac{1}{\varepsilon}\right) \frac{\alpha^{2}\beta_{\max}^{\mathcal{I}}\|\mathbf{z}^{0}-\mathbf{b}_{\perp}\|_{2}^{2}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}} (\rho^{k}+\rho^{k-1}(1+\varepsilon)\eta) \\ &+ (1+\varepsilon)^{2}\eta^{2}\mathbb{E}\left[\|\mathbf{x}^{k-2}-\mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2}\right] \\ &\leq \cdots \\ &\leq \left(1+\frac{1}{\varepsilon}\right) \frac{\alpha^{2}\beta_{\max}^{\mathcal{I}}\|\mathbf{z}^{0}-\mathbf{b}_{\perp}\|_{2}^{2}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}} \sum_{l=0}^{k-1} \rho^{k-l}(1+\varepsilon)^{l}\eta^{l} \\ &+ (1+\varepsilon)^{k}\eta^{k}\|\mathbf{x}^{0}-\mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2}. \end{split}$$

This completes the proof.

**Remark 8.** For the case REBK with s = m, t = n and  $\alpha = 1$  (i.e., REK), we have

$$\beta_{\max}^{\mathcal{I}} = \max_{i \in [m]} \frac{\|\mathbf{A}_{i,:}\|_{2}^{2}}{\|\mathbf{A}_{i,:}\|_{F}^{2}} = 1, \quad \beta_{\max}^{\mathcal{J}} = \max_{j \in [n]} \frac{\|\mathbf{A}_{:,j}\|_{2}^{2}}{\|\mathbf{A}_{:,j}\|_{F}^{2}} = 1.$$

Therefore,

$$\eta = 1 - \frac{(2\alpha - \alpha^2 \beta_{\max}^{\mathcal{I}})\sigma_r^2(\mathbf{A})}{\|\mathbf{A}\|_{\mathrm{F}}^2} = 1 - \frac{\sigma_r^2(\mathbf{A})}{\|\mathbf{A}\|_{\mathrm{F}}^2},$$

and

$$\rho = 1 - \frac{(2\alpha - \alpha^2 \beta_{\max}^{\mathcal{J}})\sigma_r^2(\mathbf{A})}{\|\mathbf{A}\|_{\mathrm{F}}^2} = 1 - \frac{\sigma_r^2(\mathbf{A})}{\|\mathbf{A}\|_{\mathrm{F}}^2}.$$

It follows from

$$\widehat{\mathbf{x}}^{k} - \mathbf{A}^{\dagger} \mathbf{b} = \left( \mathbf{I} - \frac{(\mathbf{A}_{i,:})^{\mathrm{T}} \mathbf{A}_{i,:}}{\|\mathbf{A}_{i,:}\|_{2}^{2}} \right) (\mathbf{x}^{k-1} - \mathbf{A}^{\dagger} \mathbf{b})$$

$$\mathbf{x}^k - \widehat{\mathbf{x}}^k = rac{\mathbf{b}_i - \mathbf{A}_{i,:} \mathbf{A}^\dagger \mathbf{b} - \mathbf{z}_i^k}{\|\mathbf{A}_{i,:}\|_2^2} (\mathbf{A}_{i,:})^{\mathrm{T}}$$

that

and

$$(\widehat{\mathbf{x}}^k - \mathbf{A}^{\dagger} \mathbf{b})^{\mathrm{T}} (\mathbf{x}^k - \widehat{\mathbf{x}}^k) = 0.$$

Then we have

$$\|\mathbf{x}^k - \mathbf{A}^{\dagger}\mathbf{b}\|_2^2 = \|\mathbf{x}^k - \widehat{\mathbf{x}}^k\|_2^2 + \|\widehat{\mathbf{x}}^k - \mathbf{A}^{\dagger}\mathbf{b}\|_2^2,$$

which yields the following convergence for REK (see [13]):

$$\begin{split} \mathbb{E}\left[\|\mathbf{x}^{k}-\mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2}\right] &= \mathbb{E}\left[\|\mathbf{x}^{k}-\widehat{\mathbf{x}}^{k}\|_{2}^{2}\right] + \mathbb{E}\left[\|\widehat{\mathbf{x}}^{k}-\mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2}\right] \\ &\leq \frac{\alpha^{2}\rho^{k}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\|\mathbf{z}^{0}-\mathbf{b}_{\perp}\|_{2}^{2} + \rho\mathbb{E}\left[\|\mathbf{x}^{k-1}-\mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2}\right] \\ &\leq \frac{2\alpha^{2}\rho^{k}\|\mathbf{z}^{0}-\mathbf{b}_{\perp}\|_{2}^{2}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}} + \rho^{2}\mathbb{E}\left[\|\mathbf{x}^{k-2}-\mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2}\right] \\ &\leq \cdots \\ &\leq \rho^{k}\left(\frac{k\|\mathbf{z}^{0}-\mathbf{b}_{\perp}\|_{2}^{2}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}} + \|\mathbf{x}^{0}-\mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2}\right). \end{split}$$

Actually our proof is a modification of that of Zouzias and Freris [51]. We reorganize the arguments used by Zouzias and Freris and refine the analysis to get a better convergence estimate. **Remark 9.** Let  $\hat{\rho} := \max(\eta, \rho)$  and  $\beta_{\max} := \max(\beta_{\max}^{\mathcal{I}}, \beta_{\max}^{\mathcal{J}})$ . Then we have

$$\widehat{\rho} = 1 - \frac{(2\alpha - \alpha^2 \beta_{\max})\sigma_r^2(\mathbf{A})}{\|\mathbf{A}\|_{\mathrm{F}}^2}$$

.

By Theorem 7, we have

$$\begin{split} \mathbb{E}\left[\|\mathbf{x}^{k}-\mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2}\right] &\leq (1+\varepsilon)^{k}\eta^{k}\|\mathbf{x}^{0}-\mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2} + \left(1+\frac{1}{\varepsilon}\right)\frac{\alpha^{2}\beta_{\max}^{\mathcal{I}}\|\mathbf{z}^{0}-\mathbf{b}_{\perp}\|_{2}^{2}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\sum_{l=0}^{k-l}\rho^{k-l}(1+\varepsilon)^{l}\eta^{l} \\ &\leq (1+\varepsilon)^{k}\hat{\rho}^{k}\|\mathbf{x}^{0}-\mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2} + \left(1+\frac{1}{\varepsilon}\right)\frac{\alpha^{2}\beta_{\max}^{\mathcal{I}}\|\mathbf{z}^{0}-\mathbf{b}_{\perp}\|_{2}^{2}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\hat{\rho}^{k}\sum_{l=0}^{k-l}(1+\varepsilon)^{l} \\ &\leq (1+\varepsilon)^{k}\hat{\rho}^{k}\|\mathbf{x}^{0}-\mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2} + \left(1+\frac{1}{\varepsilon}\right)\frac{\alpha^{2}\beta_{\max}^{\mathcal{I}}\|\mathbf{z}^{0}-\mathbf{b}_{\perp}\|_{2}^{2}}{\|\mathbf{A}\|_{\mathrm{F}}^{2}}\hat{\rho}^{k}\frac{(1+\varepsilon)^{k}-1}{\varepsilon} \\ &\leq (1+\varepsilon)^{k}\hat{\rho}^{k}\left(\|\mathbf{x}^{0}-\mathbf{A}^{\dagger}\mathbf{b}\|_{2}^{2} + \frac{(1+\varepsilon)\alpha^{2}\beta_{\max}^{\mathcal{I}}\|\mathbf{z}^{0}-\mathbf{b}_{\perp}\|_{2}^{2}}{\varepsilon^{2}}\|\mathbf{A}\|_{\mathrm{F}}^{2}\right), \end{split}$$

which shows that REBK exponentially converges in the mean square to the minimum  $\ell_2$ -norm least squares solution of a given linear system of equations with the rate  $(1 + \varepsilon)\hat{\rho}$  if  $0 < \alpha < 2/\beta_{\text{max}}$ . Setting  $\alpha = 1/\beta_{\text{max}}$  yields

$$\widehat{\rho} = 1 - \frac{\sigma_r^2(\mathbf{A})}{\beta_{\max} \|\mathbf{A}\|_{\mathrm{F}}^2},$$

which is better than the rate of REK (see Remark 8)

$$\rho = 1 - \frac{\sigma_r^2(\mathbf{A})}{\|\mathbf{A}\|_{\mathrm{F}}^2}$$

if  $\beta_{\max} < 1$ . A smaller  $\beta_{\max}$  means a faster convergence in terms of iterations. Recalling that

$$\beta_{\max}^{\mathcal{I}} := \max_{i \in [s]} \frac{\|\mathbf{A}_{\mathcal{I}_i,:}\|_2^2}{\|\mathbf{A}_{\mathcal{I}_i,:}\|_F^2} \quad and \quad \beta_{\max}^{\mathcal{J}} := \max_{j \in [t]} \frac{\|\mathbf{A}_{:,\mathcal{J}_j}\|_2^2}{\|\mathbf{A}_{:,\mathcal{J}_j}\|_F^2},$$

we have

$$\max_{i \in [s]} \frac{1}{|\mathcal{I}_i|} \le \max_{i \in [s]} \frac{1}{\operatorname{rank}(\mathbf{A}_{\mathcal{I}_i,:})} \le \beta_{\max}^{\mathcal{I}} \le 1$$

and

$$\max_{j \in [t]} \frac{1}{|\mathcal{J}_j|} \le \max_{j \in [t]} \frac{1}{\operatorname{rank}(\mathbf{A}_{:,\mathcal{I}_j})} \le \beta_{\max}^{\mathcal{J}} \le 1.$$

Therefore,

$$\max\left(\max_{i\in[s]}\frac{1}{|\mathcal{I}_i|}, \max_{j\in[t]}\frac{1}{|\mathcal{J}_j|}\right) \le \beta_{\max} \le 1,$$

which means that REBK is at least as fast as REK in terms of iterations. The numerical results in Section 3 show that the convergence of REBK with appropriate block sizes and stepsizes is much faster than that of REK both in the numbers of iterations and the computing times.

**Remark 10.** It was shown in [20] that the convergence of  $\mathbf{x}^k$  to  $\mathbf{A}^{\dagger}\mathbf{b}$  under the expected norm of the error (Theorem 7) is a stronger form of convergence than the convergence of the norm of the expected error (Theorem 2), as the former also guarantees that the variance of  $\mathbf{x}_i^k$  (the *i*th element of  $\mathbf{x}^k$ ) converges to zero for i = 1, ..., n. By Remark 3, we know  $0 < \alpha < 2 \|\mathbf{A}\|_{\mathrm{F}}^2 / \sigma_1^2(\mathbf{A})$ guarantees the convergence of the norm of the expected error. By Remark 9, we know  $0 < \alpha < 2/\beta_{\mathrm{max}}$  guarantees the convergence of the expected norm of the error. However, since the convergence estimate in Remark 9 usually is not sharp, the stepsize  $\alpha$  satisfying  $2/\beta_{\mathrm{max}} \leq \alpha < 2\|\mathbf{A}\|_{\mathrm{F}}^2/\sigma_1^2(\mathbf{A})$  is also possible to result in convergence (see Figure 1, Tables 2 and 3 in Section 3).

## 3 Numerical results

In this section, we compare the performance of the randomized extended block Kaczmarz (REBK) algorithm proposed in this paper against the randomized extended Kaczmarz (REK) algorithm [51] and the projection-based randomized double block Kaczmarz (RDBK) algorithm [40] on a variety of test problems. We do not claim optimized implementations of the algorithms, and only run on small or medium-scale problems. The purpose is only to demonstrate that even in these simple examples, REBK offers significant advantages to REK. All experiments are performed using MATLAB (version R2019a) on a laptop with 2.7-GHz Intel Core i7 processor, 16 GB memory, and Mac operating system.

To construct an inconsistent linear system, we set  $\mathbf{b} = \mathbf{A}\mathbf{x} + \mathbf{r}$  where  $\mathbf{x}$  is a vector with entries generated from a standard normal distribution and the residual  $\mathbf{r} \in \text{null}(\mathbf{A}^{T})$ . Note that one can obtain such a vector  $\mathbf{r}$  by the MATLAB function null. For all algorithms, we set  $\mathbf{z}^{0} = \mathbf{b}$  and  $\mathbf{x}^{0} = \mathbf{0}$  and stop if the error  $\|\mathbf{x}^{k} - \mathbf{A}^{\dagger}\mathbf{b}\|_{2} \leq 10^{-5}$ . We report the average number of iterations (denoted as ITER) and the average computing time in seconds (denoted as CPU) of REK, RDBK, and REBK. Note that  $\mathbf{A}\setminus\mathbf{b}$  will usually not be the same as pinv(A)\*b when A is rank-deficient or underdetermined. We use MATLAB's lsqminnorm (which is typically more efficient than pinv) to solve the small least squares problems at each step of RDBK. We refer the reader to [39, 40] for more numerical aspects of RDBK. We also report the speed-up of REBK against REK, which is defined as

$$speed-up = \frac{CPU \text{ of REK}}{CPU \text{ of REBK}}$$

For the block methods, we assume that the subsets  $\{\mathcal{I}_i\}_{i=1}^{s-1}$  and  $\{\mathcal{J}_j\}_{j=1}^{t-1}$  have the same size  $\tau$  (i.e.,  $|\mathcal{I}_i| = |\mathcal{J}_j| = \tau$ ). We consider the row partition  $\{\mathcal{I}_i\}_{i=1}^s$ :

$$\mathcal{I}_i = \{ (i-1)\tau + 1, (i-1)\tau + 2, \dots, i\tau \}, \quad i = 1, 2, \dots, s-1, \\ \mathcal{I}_s = \{ (s-1)\tau + 1, (s-1)\tau + 2, \dots, m \}, \quad |\mathcal{I}_s| \le \tau,$$

and the column partition  $\{\mathcal{J}_j\}_{j=1}^t$ :

$$\mathcal{J}_j = \{ (j-1)\tau + 1, (j-1)\tau + 2, \dots, j\tau \}, \quad j = 1, 2, \dots, t-1, \\ \mathcal{J}_t = \{ (t-1)\tau + 1, (t-1)\tau + 2, \dots, n \}, \quad |\mathcal{J}_t| \le \tau.$$

#### 3.1 Synthetic data

Two types of coefficient matrices are generated as follows.

• Type I: For given  $m, n, r = rank(\mathbf{A})$ , and  $\kappa > 1$ , we construct a matrix  $\mathbf{A}$  by

$$\mathbf{A} = \mathbf{U}\mathbf{D}\mathbf{V}^{\mathrm{T}},$$

where  $\mathbf{U} \in \mathbb{R}^{m \times r}$  and  $\mathbf{V} \in \mathbb{R}^{n \times r}$ . Entries of  $\mathbf{U}$  and  $\mathbf{V}$  are generated from a standard normal distribution, and then, columns are orthonormalized,

$$[\mathtt{U},\sim]=\mathtt{qr}(\mathtt{randn}(\mathtt{m},\mathtt{r}),\mathtt{0});\qquad [\mathtt{V},\sim]=\mathtt{qr}(\mathtt{randn}(\mathtt{n},\mathtt{r}),\mathtt{0});$$

The matrix **D** is an  $r \times r$  diagonal matrix whose diagonal entries are uniformly distributed numbers in  $(1, \kappa)$ ,

$$D = diag(1 + (\kappa - 1) \cdot * rand(r, 1));$$

So the condition number of **A** is upper bounded by  $\kappa$ .

• Type II: For given m, n, entries of A are generated from a standard normal distribution,

$$A = randn(m, n);$$

So **A** is a full-rank matrix almost surely.

In Figure 1, we plot the error  $\|\mathbf{x}^k - \mathbf{A}^{\dagger}\mathbf{b}\|_2$  of REBK with a fixed block size ( $\tau = 10$ ) and different stepsizes ( $\alpha$  from  $0.75/\beta_{\text{max}}$  to  $2.62/\beta_{\text{max}}$ ) for two inconsistent linear systems with coefficient matrices of Types I ( $\mathbf{A} = \mathbf{U}\mathbf{D}\mathbf{V}^{\text{T}}$  with m = 500, n = 250, r = 150,  $\kappa = 2$ ) and II ( $\mathbf{A} = \text{randn}(500, 250)$ ). It is observed that the convergence of REBK becomes faster as the increase of the stepsize, and then slows down after reaching the fastest rate.




Figure 1: The average (10 trials of each case) error  $\|\mathbf{x}^k - \mathbf{A}^{\dagger}\mathbf{b}\|_2$  of REBK with block size  $\tau = 10$ and different stepsizes  $\alpha$  from  $0.75/\beta_{\text{max}}$  to  $2.62/\beta_{\text{max}}$  for two inconsistent linear systems. Left: Type I matrix  $\mathbf{A} = \mathbf{U}\mathbf{D}\mathbf{V}^{\text{T}}$  with m = 500, n = 250, r = 150,  $\kappa = 2$ . Right: Type II matrix  $\mathbf{A} =$ randn(500,250).

In Tables 1 and 2, we report the numbers of iterations and the computing times of the REK, RDBK, and REBK algorithms for solving inconsistent linear systems. For the block algorithms (RDBK and REBK), a fixed block size  $\tau = 10$  is used. For the REBK algorithm, empirical stepsizes  $\alpha = 1.75/\beta_{\text{max}}$  and  $\alpha = 2.25/\beta_{\text{max}}$  are used for Type I and Type II matrices, respectively. From these two tables, we observe: (i) in all cases, the RDBK and REBK algorithms vastly outperform the REK algorithm in terms of both the numbers of iterations and the computing times; (ii) for Type I matrix, the convergence rates of the RDBK and REBK algorithms are almost the same in terms of the numbers of iterations; (iii) for Type II matrix, REBK performs better than RDBK in terms of the numbers of iterations.

Table 1: The average (10 trials of each algorithm) ITER and CPU of REK, RDBK( $\tau = 10$ ), and REBK( $\tau = 10$ ,  $\alpha = 1.75/\beta_{\text{max}}$ ) for inconsistent linear systems with random coefficient matrices **A** of Type I: **A** = **UDV**<sup>T</sup>.

$m \times n$	rank	κ	REK		RDBK		REBK				
			ITER	CPU	ITER	CPU	α	ITER	CPU	speed-up	
$250 \times 500$	150	2	5826	0.26	572	0.21	10.87	586	0.05	4.90	
$250 \times 500$	150	10	65520	2.87	6166	2.19	9.36	7365	0.63	4.59	
$500 \times 1000$	250	2	10068	0.59	1000	0.43	11.82	991	0.13	4.60	
$500 \times 1000$	250	10	114297	6.61	10209	4.29	10.85	10259	1.23	5.36	
$500 \times 250$	150	2	5755	0.25	562	0.19	10.70	578	0.03	7.32	
$500 \times 250$	150	10	63741	2.76	5784	1.90	10.13	6424	0.36	7.60	
$500 \times 250$	250	2	9971	0.43	940	0.31	12.47	961	0.06	7.81	
$500 \times 250$	250	10	119182	5.14	11328	3.73	10.99	10783	0.61	8.43	
$1000 \times 500$	250	2	9959	0.55	974	0.39	12.10	987	0.10	5.53	
$1000 \times 500$	250	10	118134	6.54	11236	4.44	11.20	10349	1.03	6.36	
$1000 \times 500$	500	2	20188	1.11	2007	0.80	13.84	2115	0.21	5.20	
$1000 \times 500$	500	10	254117	14.01	25361	10.00	12.67	20432	2.03	6.92	

Table 2: The average (10 trials of each algorithm) ITER and CPU of REK, RDBK( $\tau = 10$ ), and REBK( $\tau = 10$ ,  $\alpha = 2.25/\beta_{\text{max}}$ ) for inconsistent linear systems with random coefficient matrices **A** of Type II: **A** = randn(m,n).

$m \times n$	rank	$\sigma_1(\mathbf{A})$	RE	REK		RDBK		REBK			
		$\overline{\sigma_r(\mathbf{A})}$	ITER	CPU	ITER	CPU	$\alpha$	ITER	CPU	speed-up	
$250 \times 120$	120	5.25	18060	0.66	1646	0.50	13.48	1337	0.06	10.54	
$500 \times 250$	250	5.73	41016	1.79	3811	1.26	14.50	2885	0.17	10.81	
$750 \times 370$	370	5.80	59660	2.91	5929	2.20	16.23	4115	0.36	8.07	
$1000 \times 500$	500	5.74	83093	4.61	8183	3.25	16.42	5422	0.55	8.41	

In Figure 2, we plot the error  $\|\mathbf{x}^k - \mathbf{A}^{\dagger}\mathbf{b}\|_2$  and the computing times of REBK with block sizes  $\tau = 5, 10, 20, 50, 100, 200$  and stepsize  $\alpha = 1.75/\beta_{\text{max}}$  for two inconsistent linear systems with coefficient matrices of Types I ( $\mathbf{A} = \mathbf{U}\mathbf{D}\mathbf{V}^{\text{T}}$  with  $m = 20000, n = 5000, r = 4500, \kappa = 2$ ) and II ( $\mathbf{A} = \mathbf{randn}(20000, 5000)$ ). The average numbers of required iterations are also reported. We observe: (i) increasing block size and using the empirical stepsize  $\alpha = 1.75/\beta_{\text{max}}$  lead to a better convergence in terms of the numbers of iterations; (ii) with the increase of block size, the computing time first decreases, then increases after reaching the minimum value, and finally tends to be stable. This means that for sufficiently large block size the decrease in iteration complexity cannot compensate for the increase in cost per iteration. On the other hand, if a distributed version of REBK is implemented, a larger  $\tau$  will be better.




Figure 2: The average (10 trials of each case) error  $\|\mathbf{x}^k - \mathbf{A}^{\dagger}\mathbf{b}\|_2$  and CPU of REBK with different block sizes  $\tau = 10, 50, 100, 200$  and stepsize  $\alpha = 1.75/\beta_{\text{max}}$  for inconsistent linear systems. The average numbers of required iterations are also reported. Upper: Type I matrix  $\mathbf{A} = \mathbf{U}\mathbf{D}\mathbf{V}^{\mathrm{T}}$  with m = 20000, n = 5000, r = 4500, and  $\kappa = 2$ . Lower: Type II matrix  $\mathbf{A} = \mathrm{randn}(20000, 5000)$ .

In Figure 3, we plot the computing times of the REK, RDBK, and REBK algorithms for inconsistent linear systems with coefficient matrices of Types I ( $\mathbf{A} = \mathbf{U}\mathbf{D}\mathbf{V}^{\mathrm{T}}$  with m = 2000, 4000, ..., 20000, n = 500, r = 250,  $\kappa = 2$ ) and II ( $\mathbf{A} = \operatorname{randn}(\mathfrak{m},\mathfrak{n})$  with m = 2000, 4000, ..., 20000, n = 500). For all cases, the block size  $\tau = 10$  and the stepsize  $\alpha = 1.75/\beta_{\mathrm{max}}$  are used. We observe that both RDBK and REBK are better than REK, and REBK is the best.




Figure 3: The average (10 trials of each algorithm) CPU of REK, RDBK( $\tau = 10$ ), and REBK( $\tau = 10, \alpha = 1.75/\beta_{\text{max}}$ ) for inconsistent linear systems. Left: Type I matrix  $\mathbf{A} = \mathbf{U}\mathbf{D}\mathbf{V}^{\text{T}}$  with  $m = 2000, 4000, \ldots, 20000, n = 500, r = 250$ , and  $\kappa = 2$ . Right: Type II matrix  $\mathbf{A} = \text{randn}(\textbf{m,n})$  with  $m = 2000, 4000, \ldots, 20000$  and n = 500.

#### 3.2 Real-world data

Finally, we test REK and REBK using eight inconsistent linear systems with coefficient matrices from the University of Florida sparse matrix collection [10]. The eight matrices are abtaha1, flower\_5\_1, football, lp\_nug15, relat6, relat7, Sandi\_authors, and WorldCities. In Table 3, we report the numbers of iterations and the computing times for the REK and REBK algorithms. For each matrix, we tested two stepsizes of REBK, the first is  $1/\beta_{max}$  and the second is empirical. We observe that REBK based on good choices of block size and stepsize significantly outperforms REK. Moreover, good stepsize and block size are problem dependent.

## 4 Concluding remarks

We have proposed a randomized extended block Kaczmarz (REBK) algorithm for solving general linear systems and prove its convergence theory. At each step, REBK uses two RABK (with special choice of weights) updates. The new algorithm can utilize efficient implementations on distributed computing units. Numerical experiments show that the crucial point for guaranteeing fast convergence is to obtain good block size and stepsize. Finding appropriate variable stepsize by the adaptive extrapolation [35] and proposing more effective partitions based on the techniques of [39, 12, 49, 35] should be valuable topics. We also note that RABK allows the flexibility that the distributions from which blocks are selected do not require the blocks to form a partition of the columns, or rows. Designing variants of REBK based on RABK with random samplings that do not depend on the partitions is straightforward. We believe the technique used in the proof of Theorem 7 still works for these variants. Although the analysis will be more complicated. Besides, developing parallel and accelerated variants of REBK based on the approach used by Richtárik and Takáč [47] is also worth exploring. We will work on these topics in the future.

Matrix	m  imes n	rank	$\sigma_1(\mathbf{A})$	REK		REBK				
			$\overline{\sigma_r(\mathbf{A})}$	ITER	CPU	$\tau$	$\alpha$	ITER	CPU	speed-up
abtaha1	$14596 \times 209$	209	12.23	276946	89.38	10	1.82	151395	68.30	1.31
	14550 × 205						5	56064	25.34	3.53
flower_5_1	$211 \times 201$	179	13.70	135117	5.16	5	1	136037	6.15	0.84
110wer_0_1							4	34381	1.55	3.34
football	$35 \times 35$	19	166.47	810792	21.99	5	1	858215	30.64	0.72
IUUUDAII							2	409995	14.63	1.50
lp_nug15	$6330 \times 22275$	5698	2.73	216924	220.64	20	3.53	40539	199.67	1.10
							5	31039	158.29	1.39
relat6	$2340 \times 157$	137	7.74	34536	2.43	10	1	34273	3.81	0.64
101400							2.5	13971	1.56	1.56
relat7	$21924\times1045$	1012	10.85	550810	283.69	10	1	542100	466.89	0.61
							2.5	218287	188.81	1.50
Sandi authors	$86 \times 86$	72	189.58	2525141	73.28	5	1	2533343	99.36	0.74
banar_authors							2.5	999294	39.15	1.87
WorldCities	$315 \times 100$	100	66.00	120699	4.32	5	1.13	105647	4.52	0.96
							2.5	47372	2.02	2.14

Table 3: The average (10 trials of each algorithm) ITER and CPU of REK and REBK( $\tau, \alpha$ ) for inconsistent linear systems with coefficient matrices from [10]. For each matrix, two stepsizes of REBK are tested: the first is  $1/\beta_{\text{max}}$ , and the second is empirical.

# Acknowledgments

The authors are thankful to the referees for their detailed comments and valuable suggestions that have led to remarkable improvements. The research of the first author was supported by the National Natural Science Foundation of China (No.11771364) and the Fundamental Research Funds for the Central Universities (No.20720180008).

# References

- M. Arioli, I. Duff, J. Noailles, and D. Ruiz. A block projection method for sparse matrices. SIAM J. Sci. Statist. Comput., 13(1):47–70, 1992.
- [2] Z.-Z. Bai and W.-T. Wu. On convergence rate of the randomized Kaczmarz method. *Linear Algebra Appl.*, 553:252–269, 2018.
- [3] Z.-Z. Bai and W.-T. Wu. On greedy randomized Kaczmarz method for solving large sparse linear systems. SIAM J. Sci. Comput., 40(1):A592–A606, 2018.
- [4] Z.-Z. Bai and W.-T. Wu. On relaxed greedy randomized Kaczmarz methods for solving large sparse linear systems. Appl. Math. Lett., 83:21–26, 2018.
- [5] Z.-Z. Bai and W.-T. Wu. On greedy randomized coordinate descent methods for solving large linear least-squares problems. *Numer. Linear Algebra Appl.*, 26(4):e2237, 15, 2019.
- [6] Z.-Z. Bai and W.-T. Wu. On partially randomized extended Kaczmarz method for solving large sparse overdetermined inconsistent linear systems. *Linear Algebra Appl.*, 578:225–250, 2019.

- [7] A. Ben-Israel and T. N. E. Greville. Generalized Inverses: Theory and Applications., volume 15 of CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC. Springer-Verlag, New York, second edition, 2003.
- [8] R. Bramley and A. Sameh. Row projection methods for large nonsymmetric linear systems. SIAM J. Sci. Statist. Comput., 13(1):168–193, 1992.
- C. Byrne. A unified treatment of some iterative algorithms in signal processing and image reconstruction. *Inverse Problems*, 20(1):103–120, 2004.
- [10] T. A. Davis and Y. Hu. The University of Florida sparse matrix collection. ACM Trans. Math. Software, 38(1):Art. 1, 25, 2011.
- [11] J. A. De Loera, J. Haddock, and D. Needell. A sampling Kaczmarz-Motzkin algorithm for linear feasibility. SIAM J. Sci. Comput., 39(5):S66–S87, 2017.
- [12] L. A. Drummond, I. S. Duff, R. Guivarch, D. Ruiz, and M. Zenadi. Partitioning strategies for the block Cimmino algorithm. J. Engrg. Math., 93:21–39, 2015.
- [13] K. Du. Tight upper bounds for the convergence of the randomized extended Kaczmarz and Gauss-Seidel algorithms. Numer. Linear Algebra Appl., 26(3):e2233, 14, 2019.
- [14] K. Du and H. Gao. A new theoretical estimate for the convergence rate of the maximal weighted residual Kaczmarz algorithm. Numer. Math. Theory Methods Appl., 12(2):627– 639, 2019.
- [15] K. Du and X. Sun. A doubly stochastic block Gauss-Seidel algorithm for solving linear equations. arXiv preprint arXiv:1912.13291, 2019.
- [16] I. S. Duff, R. Guivarch, D. Ruiz, and M. Zenadi. The augmented block Cimmino distributed method. SIAM J. Sci. Comput., 37(3):A1248–A1269, 2015.
- [17] B. Dumitrescu. On the relation between the randomized extended Kaczmarz algorithm and coordinate descent. BIT, 55(4):1005–1015, 2015.
- [18] T. Elfving. Block-iterative methods for consistent and inconsistent linear equations. Numer. Math., 35(1):1–12, 1980.
- [19] G. H. Golub and C. F. Van Loan. *Matrix computations*. Johns Hopkins Studies in the Mathematical Sciences. Johns Hopkins University Press, Baltimore, MD, fourth edition, 2013.
- [20] R. M. Gower and P. Richtárik. Randomized iterative methods for linear systems. SIAM J. Matrix Anal. Appl., 36(4):1660–1690, 2015.
- [21] J. Haddock and D. Needell. On Motzkin's method for inconsistent linear systems. BIT, 59(2):387–401, 2019.
- [22] J. Haddock and D. Needell. Randomized projection methods for linear systems with arbitrarily large sparse corruptions. SIAM J. Sci. Comput., 41(5):S19–S36, 2019.
- [23] A. Hefny, D. Needell, and A. Ramdas. Rows versus columns: randomized Kaczmarz or Gauss-Seidel for ridge regression. SIAM J. Sci. Comput., 39(5):S528–S542, 2017.
- [24] G. T. Herman. Fundamentals of computerized tomography. Advances in Pattern Recognition. Springer, Dordrecht, second edition, 2009. Image reconstruction from projections.

- [25] G. T. Herman and R. Davidi. Image reconstruction from a small number of projections. *Inverse Problems*, 24(4):045011, 17, 2008.
- [26] G. T. Herman and L. B. Meyer. Algebraic reconstruction techniques can be made computationally efficient. *IEEE Trans. Medical Imaging*, 12(3):600–9, 1993.
- [27] S. Kaczmarz. Angenäherte auflösung von systemen linearer gleichungen. Bull. Intern. Acad. Polonaise Sci. Lett., Cl. Sci. Math. Nat. A, 35:355–357, 1937.
- [28] A. C. Kak and M. Slaney. Principles of computerized tomographic imaging, volume 33 of Classics in Applied Mathematics. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2001. Reprint of the 1988 original.
- [29] D. Leventhal and A. S. Lewis. Randomized methods for linear constraints: convergence rates and conditioning. *Math. Oper. Res.*, 35(3):641–654, 2010.
- [30] J. Liu and S. J. Wright. An accelerated randomized Kaczmarz algorithm. Math. Comp., 85(297):153–178, 2016.
- [31] Y. Liu and C.-Q. Gu. Variant of greedy randomized Kaczmarz for ridge regression. Appl. Numer. Math., 143:223–246, 2019.
- [32] D. A. Lorenz, S. Wenger, F. Schöpfer, and M. Magnor. A sparse kaczmarz solver and a linearized bregman method for online compressed sensing. In *IEEE International Conference* on *Image Processing*, pages 1347 – 1351, 2015.
- [33] A. Ma, D. Needell, and A. Ramdas. Convergence properties of the randomized extended Gauss-Seidel and Kaczmarz methods. SIAM J. Matrix Anal. Appl., 36(4):1590–1604, 2015.
- [34] F. Natterer. The mathematics of computerized tomography, volume 32 of Classics in Applied Mathematics. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2001. Reprint of the 1986 original.
- [35] I. Necoara. Faster randomized block kaczmarz algorithms. SIAM J. Matrix Anal. Appl., 40(4):1425–1452, 2019.
- [36] I. Necoara, P. Richtárik, and A. Patrascu. Randomized projection methods for convex feasibility: conditioning and convergence rates. *SIAM J. Optim.*, 29(4):2814–2852, 2019.
- [37] D. Needell. Randomized Kaczmarz solver for noisy linear systems. BIT, 50(2):395–403, 2010.
- [38] D. Needell, N. Srebro, and R. Ward. Stochastic gradient descent, weighted sampling, and the randomized Kaczmarz algorithm. *Math. Program.*, 155(1-2, Ser. A):549–573, 2016.
- [39] D. Needell and J. A. Tropp. Paved with good intentions: analysis of a randomized block Kaczmarz method. *Linear Algebra Appl.*, 441:199–221, 2014.
- [40] D. Needell, R. Zhao, and A. Zouzias. Randomized block Kaczmarz method with projection for solving least squares. *Linear Algebra Appl.*, 484:322–343, 2015.
- [41] S. Petra and C. Popa. Single projection Kaczmarz extended algorithms. Numer. Algorithms, 73(3):791–806, 2016.
- [42] B. T. Poljak. Gradient methods for minimizing functionals. Ž. Vyčisl. Mat i Mat. Fiz., 3:643–653, 1963.

- [43] C. Popa. Extensions of block-projections methods with relaxation parameters to inconsistent and rank-deficient least-squares problems. BIT, 38(1):151–176, 1998.
- [44] C. Popa. Convergence rates for Kaczmarz-type algorithms. Numer. Algorithms, 79(1):1–17, 2018.
- [45] C. Popa and R. Zdunek. Kaczmarz extended algorithm for tomographic image reconstruction from limited data. *Math. Comput. Simulation*, 65(6):579–598, 2004.
- [46] M. Razaviyayn, M. Hong, N. Reyhanian, and Z.-Q. Luo. A linearly convergent doubly stochastic Gauss-Seidel algorithm for solving linear equations and a certain class of overparameterized optimization problems. *Math. Program.*, 176(1-2, Ser. B):465–496, 2019.
- [47] P. Richtárik and M. Takáč. Stochastic reformulations of linear systems algorithms and convergence theory. arXiv preprint arXiv:1706.01108, 2017.
- [48] T. Strohmer and R. Vershynin. A randomized Kaczmarz algorithm with exponential convergence. J. Fourier Anal. Appl., 15(2):262–278, 2009.
- [49] F. S. Torun, M. Manguoglu, and C. Aykanat. A novel partitioning method for accelerating the block Cimmino algorithm. SIAM J. Sci. Comput., 40(6):C827–C850, 2018.
- [50] J.-J. Zhang. A new greedy Kaczmarz algorithm for the solution of very large linear systems. Appl. Math. Lett., 91:207–212, 2019.
- [51] A. Zouzias and N. M. Freris. Randomized extended Kaczmarz for solving least squares. SIAM J. Matrix Anal. Appl., 34(2):773–793, 2013.