Canonical extensions of lattices are more than perfect

Andrew P. K. Craig, Maria J. Gouveia and Miroslav Haviar

In memory of Bjarni Jónsson

Abstract. In [2] we introduced TiRS graphs and TiRS frames to create a new natural setting for duals of canonical extensions of lattices. In this continuation of [2] we answer Problem 2 from there by characterising the perfect lattices that are dual to TiRS frames (and hence TiRS graphs). We introduce a new subclass of perfect lattices called PTi lattices and show that the canonical extensions of lattices are PTi lattices, and so are 'more' than just perfect lattices. We introduce morphisms of TiRS structures and put our correspondence between TiRS graphs and TiRS frames from [2] into a full categorical framework. We illustrate our correspondences between classes of perfects lattices and classes of TiRS graphs by examples.

Mathematics Subject Classification. 06B23, 06D50, 06B15.

Keywords. bounded lattice, canonical extension, perfect lattice, RS frame, TiRS frame, TiRS graph, PTi lattice.

1. Introduction

An important aspect of the study of lattice-based algebras in recent decades has been the theory of canonical extensions. This has its origins in the 1951–52 papers of Jónsson and Tarski [13]. We refer to Gehrke and Vosmaer [10] for a survey of the theory of canonical extensions for lattice-based algebras and, for further background, to recent papers by Gehrke [8] and Goldblatt [11] and the references there, in particular to the first section of [11] called "A biography of canonical extension".

The canonical extensions of general (bounded) lattices were first introduced by Gehrke and Harding [9] as the complete lattices of Galois-closed sets associated with a polarity between the filter lattice and the ideal lattice of the given lattice. (The same polarity was also used in the lattice representation of Hartonas and Dunn [12].) A new construction of the canonical extension

of a general lattice was provided in [3] where it was based on a topological representation of lattices by Ploščica [16]. The Ploščica representation presented a well-known representation of general lattices due to Urquhart [18] in the spirit of the theory of natural dualities of Clark and Davey [1]. It used maximal partial maps into the two-element set to represent elements of the first and second duals of a given lattice.

An another construction of the canonical extensions of general lattices was presented in [2] where Ploščica's topological representation was used in tandem with Gehrke's representation of perfect lattices via RS frames. (For the latter we refer to papers [6] by Dunn, Gehrke and Palmigiano and [7] by Gehrke.) In [2] we also demonstrated a one-to-one correspondence between TiRS frames forming a subclass of the RS frames and TiRS graphs which we introduced as an abstraction of the duals of general lattices in the Ploščica representation. This has led to a new dual representation of the class of all finite lattices via finite TiRS frames, or equivalently finite TiRS graphs, which generalises the well-known Birkhoff dual representation between finite distributive lattices and finite posets from the 1930s. (Here we remark that every poset is a TiRS graph.) We use a common concept of *TiRS structures* when we refer to both TiRS graphs and TiRS frames without distinguishing between the two classes.

This paper has two goals:

- (1) To describe the additional properties that perfect lattices dual to TiRS structures possess. This was listed as "Problem 2" in [2].
- (2) To describe the appropriate morphisms of TiRS structures and hence to extend the one-to-one correspondence between the TiRS structures from [2] into a full categorical framework.

We also show that the canonical extensions of lattices are PTi lattices, which follows from their construction in [2] using Ploščica's and Gehrke's representations in tandem. We present an example of a perfect but not PTi lattice together with its dual TiRS graph and an example of a PTi lattice that is not the canonical extension of any lattice together with its dual TiRS frame.

2. Preliminaries

For a bounded lattice \mathbf{L} , a completion of \mathbf{L} is defined to be a pair (e, \mathbf{C}) where \mathbf{C} is a complete lattice and $e \colon \mathbf{L} \hookrightarrow \mathbf{C}$ is an embedding. By a filter element (ideal element) of a completion (e, \mathbf{C}) of a bounded lattice \mathbf{L} we mean an element of \mathbf{C} which is a meet (join) of elements from $e(\mathbf{L})$. By $\mathbb{F}(\mathbf{C})$ and $\mathbb{I}(\mathbf{C})$ are denoted the sets of all filter and ideal elements of \mathbf{C} , respectively. (We remark that in the older literature the filter (ideal) elements had been called closed (open) elements.) A completion (e, \mathbf{C}) of a bounded lattice \mathbf{L} is called dense if every element of \mathbf{C} can be expressed as both a join of meets and a meet of joins of elements from $e(\mathbf{L})$. A completion (e, \mathbf{C}) of \mathbf{L} is called compact if, for any sets $A \subseteq \mathbb{F}(\mathbf{C})$ and $B \subseteq \mathbb{I}(\mathbf{C})$ with $\bigwedge A \leqslant \bigvee B$, there exist finite subsets $A' \subseteq A$ and $B' \subseteq B$ such that $\bigwedge A' \leqslant \bigvee B'$. (We remark

that the sets A, B in the definition of compactness above can alternatively be taken as arbitrary subsets of L.)

Gehrke and Harding [9] defined abstractly the canonical extension \mathbf{L}^{δ} of a general bounded lattice \mathbf{L} as a dense and compact completion of \mathbf{L} . They proved that every bounded lattice \mathbf{L} has a canonical extension and that it is unique up to an isomorphism that fixes the elements of \mathbf{L} . Concretely, they constructed \mathbf{L}^{δ} as the complete lattice of Galois-stable sets of the polarity R between the filter lattice $\mathrm{Filt}(\mathbf{L})$ and the ideal lattice $\mathrm{Idl}(\mathbf{L})$ of \mathbf{L} where the polarity is given by $(F, I) \in R$ if $F \cap I \neq \emptyset$.

A filter-ideal pair (F, I) will be called *maximal* if F and I are maximal with respect to being disjoint from one another. In our final section we shall use the following result from [9]:

Lemma 2.1 ([9, Lemma 3.4]). Let (e, \mathbf{C}) be a canonical extension of \mathbf{L} .

- (1) $x \in J^{\infty}(\mathbf{C})$ if and only if $x = \bigwedge e[F]$ for some maximal pair (F, I) of \mathbf{L} :
- (2) $x \in M^{\infty}(\mathbf{C})$ if and only if $x = \bigvee e[I]$ for some maximal pair (F, I) of \mathbf{L} .

Further, each element of \mathbf{C} is a join of completely join irreducibles and a meet of completely meet irreducibles.

Ploščica's dual [16, Section 1] of a bounded lattice **L** is a graph with topology, $D(\mathbf{L}) = (\mathcal{L}^{mp}(\mathbf{L}, \underline{\mathbf{2}}), E, \mathcal{T})$, where $\mathcal{L}^{mp}(\mathbf{L}, \underline{\mathbf{2}})$ is the set of maximal partial homomorphisms from **L** into $\underline{\mathbf{2}}$. The graph relation E is defined by

$$(f,g) \in E$$
 if $(\forall a \in dom \ f \cap dom \ g) \ f(a) \leqslant g(a)$,

or equivalently,

$$(f,g) \in E$$
 if $f^{-1}(1) \cap g^{-1}(0) = \emptyset$.

The topology \mathcal{T} has as a subbasis of closed sets the set $\{V_a, W_a \mid a \in L\}$, with $V_a = \{f \in \mathcal{L}^{\mathrm{mp}}(\mathbf{L}, \underline{\mathbf{2}}) \mid f(a) = 0\}$ and $W_a = \{f \in \mathcal{L}^{\mathrm{mp}}(\mathbf{L}, \underline{\mathbf{2}}) \mid f(a) = 1\}$.

TiRS graphs were defined by the present authors in [2] as an abstraction of the graphs $D^{\flat}(\mathbf{L}) = (\mathcal{L}^{mp}(\mathbf{L}, \mathbf{2}), E)$ obtained from Ploščica's duals of bounded lattices \mathbf{L} by forgetting the topology.

For a graph $\mathbf{X} = (X, E)$ and $x \in X$, the sets $\{y \in X \mid (x, y) \in E\}$ and $\{y \in X \mid (y, x) \in E\}$ were denoted in [2] by xE and Ex respectively. We defined the conditions (S), (R) and (Ti) for any graph $\mathbf{X} = (X, E)$ as follows:

- (S) for every $x, y \in X$, if $x \neq y$ then $xE \neq yE$ or $Ex \neq Ey$;
- (R) (i) for all $x, z \in X$, if $zE \subseteq xE$ then $(z, x) \notin E$;
 - (ii) for all $y, z \in X$, if $Ez \subsetneq Ey$ then $(y, z) \notin E$;
- (Ti) for all $x, y \in X$, if $(x, y) \in E$, then there exists $z \in X$ such that $zE \subseteq xE$ and $Ez \subseteq Ey$.

A $TiRS\ graph$ was in [2] defined as a graph $\mathbf{X} = (X, E)$ with a reflexive relation E and satisfying the conditions (R), (S) and (Ti). For any bounded lattice \mathbf{L} , its dual graph $\mathbf{X} = \mathrm{D}^{\flat}(\mathbf{L})$ is a TiRS graph [2, Proposition 2.3].

We further recall that a *frame* is a structure (X_1, X_2, R) , where X_1 and X_2 are non-empty sets and $R \subseteq X_1 \times X_2$. For an arbitrary frame $\mathbf{F} = (X_1, X_2, R)$ the conditions (S) and (R) are defined as follows:

- (S) for all $x_1, x_2 \in X_1$ and $y_1, y_2 \in X_2$,
 - (i) $x_1 \neq x_2$ implies $x_1 R \neq x_2 R$;
 - (ii) $y_1 \neq y_2$ implies $Ry_1 \neq Ry_2$.
- (R) (i) for every $x \in X_1$ there exists $y \in X_2$ such that $\neg(xRy)$ and $\forall w \in X_1 \ ((w \neq x \& xR \subseteq wR) \Rightarrow wRy);$
 - (ii) for every $y \in X_2$ there exists $x \in X_1$ such that $\neg(xRy)$ and $\forall z \in X_2$ $((z \neq y \& Ry \subseteq Rz) \Rightarrow xRz)$.

The frames that satisfy the conditions (R) and (S) are called reduced separated frames, or RS frames for short, and were introduced by Gehrke [7] as a two-sorted generalisation of Kripke frames to be used for relational semantics of substructural logics.

The (Ti) condition introduced in [2] for frames (X_1, X_2, R) was motivated by the (Ti) condition on graphs:

- (Ti) for every $x \in X_1$ and for every $y \in X_2$, if $\neg(xRy)$ then there exist $w \in X_1$ and $z \in X_2$ such that
 - (i) $\neg (wRz)$;
 - (ii) $xR \subseteq wR$ and $Ry \subseteq Rz$;
 - (iii) for every $u \in X_1$, if $u \neq w$ and $wR \subseteq uR$ then uRz;
 - (iv) for every $v \in X_2$, if $v \neq z$ and $Rz \subseteq Rv$ then wRv.

A $TiRS\ frame\ was\ in\ [2]\ defined\ as\ a\ frame\ (X_1,X_2,R)\ that\ satisfies\ conditions\ (R),\ (S)\ and\ (Ti),\ i.e.\ it\ is\ an\ RS\ frame\ that\ satisfies\ condition\ (Ti).$ A one-to-one correspondence between TiRS graphs and TiRS frames was then shown in [2]. We recall here some facts of this correspondence that will be needed in the next section.

Definition 2.2 ([2, Definition 2.5]). Let $\mathbf{X} = (X, E)$ be a graph. The associated frame $\rho(\mathbf{X})$ is the frame $(X_1, X_2, R_{\rho(\mathbf{X})})$ where

(i) $X_1 = X/\sim_1$ for the equivalence relation \sim_1 on X given by

$$x \sim_1 y \text{ if } xE = yE;$$

(ii) $X_2 = X/\sim_2$ for the equivalence relation \sim_2 on X given by

$$x \sim_2 y \text{ if } Ex = Ey;$$

(iii) $R_{\rho(\mathbf{X})}$ is the relation given by

$$[x]_1 R_{\rho(\mathbf{X})}[y]_2 \iff (x,y) \notin E,$$

where $[x]_1$ and $[y]_2$ are, respectively, the \sim_1 -equivalence class of x and the \sim_2 -equivalence class of y.

We omit the subscript $\rho(\mathbf{X})$ in $R_{\rho(\mathbf{X})}$ whenever it is clear to which relation R refers.

If $\mathbf{X}=(X,E)$ is a TiRS graph, then the associated frame $\rho(\mathbf{X})=(X_1,X_2,R_{\rho(\mathbf{X})})$ is a TiRS frame [2, Proposition 2.6]. Then it follows that if

L is a bounded lattice, $\mathbf{X} = D^{\flat}(\mathbf{L})$ is its dual TiRS graph and $\rho(D^{\flat}(\mathbf{L}))$ is the associated frame, then $\rho(D^{\flat}(\mathbf{L}))$ is a TiRS frame (cf. [2, Corollary 2.7]).

Definition 2.3 ([2, Definition 2.8]). Let $\mathbf{F} = (X_1, X_2, R)$ be a TiRS frame. The associated graph $gr(\mathbf{F})$ is $(H_{\mathbf{F}}, K_{\mathbf{F}})$ where the vertex set $H_{\mathbf{F}}$ is the subset of $X_1 \times X_2$ of all pairs (x, y) that satisfy the following conditions:

- (i) $\neg (xRy)$,
- (ii) for every $u \in X_1$, if $u \neq x$ and $xR \subseteq uR$ then uRy,
- (iii) for every $v \in X_2$, if $v \neq y$ and $Ry \subseteq Rv$ then xRv.

and the edge set $K_{\mathbf{F}}$ is formed by the pairs ((x,y),(w,z)) such that $\neg(xRz)$.

We omit the subscript \mathbf{F} in $H_{\mathbf{F}}$ and in $K_{\mathbf{F}}$ whenever it is clear which vertex set and edge set we refer to.

In [2, Proposition 2.10] we showed that if $\mathbf{F} = (X_1, X_2, R)$ is a TiRS frame, then its associated graph $gr(\mathbf{F})$ is a TiRS graph.

Definition 2.4 ([2, Definition 2.11]). Two graphs $\mathbf{X} = (X, E_X)$ and $\mathbf{Y} = (Y, E_Y)$ are *isomorphic* (denoted $\mathbf{X} \simeq \mathbf{Y}$) if there exists a bijective map $\alpha \colon X \to Y$ such that

$$\forall x_1, x_2 \in X \quad (x_1, x_2) \in E_X \iff (\alpha(x_1), \alpha(x_2)) \in E_Y$$

and we refer to such a map as the graph-isomorphism $\alpha \colon \mathbf{X} \to \mathbf{Y}$.

Two frames $\mathbf{F} = (X_1, X_2, R_F)$ and $\mathbf{G} = (Y_1, Y_2, R_G)$ are isomorphic (denoted $\mathbf{F} \simeq \mathbf{G}$) if there exists a pair (β_1, β_2) of bijective maps $\beta_i \colon X_i \to Y_i$ (i = 1, 2) with

$$\forall x_1 \in X_1 \ \forall x_2 \in X_2 \quad \left(x_1 R_F x_2 \iff \beta_1(x_1) R_G \beta_2(x_2) \right)$$

and we refer to such a pair as the frame-isomorphism $(\beta_1, \beta_2) \colon \mathbf{F} \to \mathbf{G}$.

Now for a TiRS graph $\mathbf{X}=(X,E)$, a map $\alpha_X\colon X\to gr(\rho(\mathbf{X}))$ is defined by $\alpha_X(x)=([x]_1,[x]_2)$. The next result shows that α_X is a graph isomorphism and that the correspondence between TiRS graphs and TiRS frames is one-to-one.

Theorem 2.5 ([2, Theorem 2.13]). Let $\mathbf{X} = (X, E)$ be a TiRS graph and $\mathbf{F} = (X_1, X_2, R)$ be a TiRS frame. Then

- (a) the graphs **X** and $gr(\rho(\mathbf{X}))$ are isomorphic;
- (b) the frames ${\bf F}$ and $\rho(\operatorname{gr}({\bf F}))$ are isomorphic.

3. TiRS graph and TiRS frame morphisms

In this section we extend the one-to-one correspondence between TiRS graphs and TiRS frames from [2] into the full categorical framework. We start by defining the concepts of TiRS graph and TiRS frame morphisms.

Definition 3.1. Let $\mathbf{X} = (X, E_X)$ and $\mathbf{Y} = (Y, E_Y)$ be TiRS graphs. A *TiRS graph morphism* is a map $\varphi \colon X \to Y$ that satisfies the following conditions:

(i) for $x_1, x_2 \in X$, if $(x_1, x_2) \in E_X$ then $(\varphi(x_1), \varphi(x_2)) \in E_Y$;

- (ii) for $x_1, x_2 \in X$, if $x_1 E_X \subseteq x_2 E_X$ then $\varphi(x_1) E_Y \subseteq \varphi(x_2) E_Y$;
- (iii) for $x_1, x_2 \in X$, if $E_X x_1 \subseteq E_X x_2$ then $E_Y \varphi(x_1) \subseteq E_Y \varphi(x_2)$.

We note that every graph isomorphism and its inverse are TiRS graph morphisms.

Definition 3.2. Let $\mathbf{F} = (X_1, X_2, R_F)$ and $\mathbf{G} = (Y_1, Y_2, R_G)$ be TiRS frames. A *TiRS frame morphism* $\psi \colon \mathbf{F} \to \mathbf{G}$ is a pair $\psi = (\psi_1, \psi_2)$ of maps $\psi_1 \colon X_1 \to Y_1$ and $\psi_2 \colon X_2 \to Y_2$ that satisfies the following conditions:

- (i) for $x \in X_1$ and $y \in X_2$, if $\psi_1(x)R_G\psi_2(y)$ then xR_Fy ;
- (ii) for $x, w \in X_1$, if $xR_F \subseteq wR_F$ then $\psi_1(x)R_G \subseteq \psi_1(w)R_G$;
- (iii) for $y, z \in X_2$, if $R_F y \subseteq R_F z$ then $R_G \psi_2(y) \subseteq R_G \psi_2(z)$;
- (iv) for $x \in X_1$ and $y \in X_2$, if $(x, y) \in H_{\mathbf{F}}$ then $(\psi_1(x), \psi_2(y)) \in H_{\mathbf{G}}$.

We note that a frame isomorphism is a TiRS morphism.

Henceforth we shall refer to TiRS graph morphisms and to TiRS frame morphisms simply as graph morphisms and frame morphisms respectively.

Our main result in this section puts our one-to-one correspondence between TiRS graphs and TiRS frames into a full categorical framework. The last two statements are illustrated by the diagrams in Fig. 1.

Theorem 3.3. Let $\mathbf{X} = (X, E_X)$ and $\mathbf{Y} = (Y, E_Y)$ be TiRS graphs and let $\mathbf{F} = (X_1, X_2, R_F)$ and $\mathbf{G} = (Y_1, Y_2, R_G)$ be TiRS frames.

- (1) If $\varphi \colon \mathbf{X} \to \mathbf{Y}$ is a TiRS graph morphism, then, for $\rho(\varphi)_1 \colon X/\sim_1 \to Y/\sim_1$ and $\rho(\varphi)_2 \colon X/\sim_2 \to Y/\sim_2$ the maps defined by $\rho(\varphi)_1([x]_1) = [\varphi(x)]_1$ and $\rho(\varphi)_2([x]_2) = [\varphi(x)]_2$, for all $x \in X$, the pair $\rho(\varphi) = (\rho(\varphi)_1, \rho(\varphi)_2)$ is a TiRS frame morphism from $\rho(\mathbf{X})$ to $\rho(\mathbf{Y})$.
- (2) If the pair $\psi = (\psi_1, \psi_2) \colon \mathbf{F} \to \mathbf{G}$ is a TiRS frame morphism, then the map $\operatorname{gr}(\psi) \colon \operatorname{gr}(\mathbf{F}) \to \operatorname{gr}(\mathbf{G})$ defined by $\operatorname{gr}(\psi)(x,y) = (\psi_1(x), \psi_2(y))$, for $(x,y) \in H_F$, is a TiRS graph morphism.
- (3) If $\varphi \colon \mathbf{X} \to \mathbf{Y}$ is a TiRS graph morphism, then $\operatorname{gr}(\rho(\varphi)) \circ \alpha_X = \alpha_Y \circ \varphi$.
- (4) If $\psi \colon \mathbf{F} \to \mathbf{G}$ is a TiRS frame morphism, then $\rho(\operatorname{gr}(\psi)) \circ \beta_F = \beta_G \circ \psi$.

Proof. (1) First we show that $\rho(\varphi)_1$ is well defined. Let $x,y\in X$. If $[x]_1=[y]_1$ then $xE_X=yE_X$ which implies $\varphi(x)E_Y=\varphi(y)E_Y$ and so $[\varphi(x)]_1=[\varphi(y)]_1$, by the definition of a TiRS graph morphism. Similarly we prove that $[\varphi(x)]_2=[\varphi(y)]_2$ whenever $[x]_2=[y]_2$. Next we prove that conditions (i) to (iv) of the definition of a TiRS frame morphism are satisfied by $\rho(\varphi)$. For (i), let $x,y\in X$ and assume $\rho(\varphi)_1([x]_1)R_{\rho(\mathbf{Y})}\rho(\varphi)_2([y]_2)$. Then $(\varphi(x),\varphi(y))\notin E_Y$ yielding that $(x,y)\notin E_X$ and so $[x]_1R_{\rho(\mathbf{X})}[y]_2$. For (ii), let $x,w\in X$. Then the following holds:

$$[x]_1 R_{\rho(\mathbf{X})} \subseteq [w]_1 R_{\rho(\mathbf{X})} \iff w E_X \subseteq x E_X \Rightarrow \varphi(w) E_Y \subseteq \varphi(x) E_Y$$
$$\iff [\varphi(x)]_1 R_{\varrho(\mathbf{Y})} \subseteq [\varphi(w)]_1 R_{\varrho(\mathbf{Y})}.$$

Hence (ii) is satisfied. Similarly we conclude that (iii) holds. Finally (iv) follows from [2, Lemma 2.12] where we showed that for a TiRS graph $\mathbf{X} = (X, E)$, the elements of $H_{\rho(\mathbf{X})}$ are exactly the pairs $([x]_1, [x]_2)$, with $x \in X$.




FIGURE 1. TiRS graph morphisms and TiRS frame morphisms

(2) First we note that condition (iv) of the definition of a TiRS frame morphism satisfied by ψ guarantees that the map $\operatorname{gr}(\psi)$ is well defined. Next we prove that conditions (i) to (iii) of the definition of a TiRS graph morphism are satisfied by $\operatorname{gr}(\psi)$. Let $(x,y),(w,z) \in H_{\mathbf{F}}$. If $((x,y),(w,z)) \in K_{\mathbf{F}}$ then $\neg(xR_{\mathbf{F}}z)$ which implies $\neg(\psi_1(x)R_{\mathbf{G}}\psi_2(z))$ and consequently

$$(\operatorname{gr}(\psi)(x,y),\operatorname{gr}(\psi)(w,z)) = ((\psi_1(x),\psi_2(y)),(\psi_1(w),\psi_2(z))) \in K_{\mathbf{G}}.$$

Hence (x, y), (w, z) satisfies (i). For (ii), we observe that,

$$(x,y)K_{\mathbf{F}} \subseteq (w,z)K_{\mathbf{F}} \iff wR_F \subseteq xR_F$$

and

$$\operatorname{gr}(\psi)(x,y)K_{\mathbf{G}} \subseteq \operatorname{gr}(\psi)(w,z)K_{\mathbf{G}} \iff \psi_1(w)R_G \subseteq \psi_1(x)R_G,$$

which follows from (iii) of [2, Lemma 3.9]. As ψ is a TiRS morphism, we also have

$$wR_F \subseteq xR_F \Rightarrow \psi_1(w)R_G \subseteq \psi_1(x)R_G.$$

Hence (x, y), (w, z) satisfies (ii). Similarly we conclude that (iii) also holds.

(3) Let $x \in X$. We have that

$$(\operatorname{gr}(\rho(\varphi)) \circ \alpha_X)(x) = \operatorname{gr}(\rho(\varphi))([x]_1, [x]_2)$$

$$= (\rho(\varphi)_1([x]_1), \rho(\varphi)_2([x]_2))$$

$$= ([\varphi(x)]_1, [\varphi(x)]_2)$$

$$= (\alpha_Y \circ \varphi)(x).$$

(4) Let $x \in X_1$. There exist $y \in X_2$ such that $(x, y) \in H_{\mathbf{F}}$. We have that

$$(\rho(gr(\psi)) \circ \beta_F)(x) = \rho(gr(\psi)([(x,y)]_1) = [gr(\psi)(x,y)]_1 = [(\psi_1(x), \psi_2(y))]_1,$$

where $\psi_1: X_1 \to Y_1$ and $\psi_2: X_2 \to Y_2$ satisfy $\psi = (\psi_1, \psi_2)$. Since $(x, y) \in H_{\mathbf{F}}$ and ψ is a TiRS morphism, we also have $(\psi_1(x), \psi_2(y)) \in H_{\mathbf{G}}$ and so

$$[(\psi_1(x), \psi_2(y))]_1 = \beta_G(\psi(x)) = (\beta_G \circ \psi)(x).$$

Corollary 3.4. The category of TiRS graphs with TiRS graph morphisms is equivalent to the category of TiRS frames with TiRS frame morphisms via the functors given by ρ and gr as described above.

There are other definitions of morphisms between two frames (or contexts) $\mathbf{F} = (X_1, X_2, R_F)$ and $\mathbf{G} = (Y_1, Y_2, R_G)$ that are used in the literature. Deiters and Erné [5] use a pair of maps (α, β) where $\alpha \colon X_1 \to Y_1$ and $\beta \colon X_2 \to Y_2$ as we do above. Gehrke [7, Section 3] uses a pair of relations (R, S) where $R \subseteq X_2 \times Y_1$ and $S \subseteq X_1 \times Y_2$. More recently, Moshier [15] (see also Jipsen [14]) defined a context morphism to be a single relation $S \subseteq X_1 \times Y_2$.

4. Perfect lattices dual to TiRS structures

Consider a complete lattice \mathbf{C} and let $\mathbf{F}(\mathbf{C}) = (J^{\infty}(\mathbf{C}), M^{\infty}(\mathbf{C}), \leqslant)$ where $J^{\infty}(\mathbf{C})$ and $M^{\infty}(\mathbf{C})$ denote the sets of completely join-irreducible and completely meet-irreducible elements of \mathbf{C} , respectively. We will refer to $\mathbf{F}(\mathbf{C})$ as the frame coming from \mathbf{C} . For the opposite direction, consider an RS frame $\mathbb{F} = (X, Y, R)$. For $A \subseteq X$ and $B \subseteq Y$, let

$$R_{\triangleright}(A) = \{ y \in Y \mid (\forall a \in A)(aRy) \} \text{ and } R_{\triangleleft}(B) = \{ x \in X \mid (\forall b \in B)(xRb) \}.$$

Now consider the complete lattice of Galois-closed sets (ordered by inclusion):

$$\mathcal{G}(\mathbb{F}) = \{ A \subset X \mid A = (R_{\triangleleft} \circ R_{\triangleright})(A) \}.$$

By results from Gehrke [7, Section 2] we know that the completely join-irreducible elements and completely meet-irreducible elements of $\mathcal{G}(\mathbb{F})$ are identified as follows:

$$J^{\infty}(\mathcal{G}(\mathbb{F})) = \{ (R_{\triangleleft} \circ R_{\triangleright})(\{x\}) \mid x \in X \} \text{ and } M^{\infty}(\mathcal{G}(\mathbb{F})) = \{ Ry \mid y \in Y \}.$$

Below we introduce a condition that refines the class of perfect lattices. At the end of this section we will conclude that every perfect lattice that is the canonical extension of some bounded lattice will have this property.

Definition 4.1. A perfect lattice satisfies the condition (PTi) if for all $x \in J^{\infty}(\mathbf{C})$ and for all $y \in M^{\infty}(\mathbf{C})$, if $x \nleq y$ then there exist $w \in J^{\infty}(\mathbf{C})$, $z \in M^{\infty}(\mathbf{C})$ such that

- (i) $w \leqslant x$ and $y \leqslant z$
- (ii) $w \nleq z$
- (iii) $(\forall u \in J^{\infty}(\mathbf{C}))(u < w \Rightarrow u \leqslant z)$
- (iv) $(\forall v \in M^{\infty}(\mathbf{C}))(y < v \Rightarrow w \leqslant v)$

In Fig. 2 we give a pictorial depiction of the (PTi) condition. We have indicated the sets $\uparrow x$, $\uparrow w$, $\downarrow y$ and $\downarrow z$. We see that the (PTi) condition for **C** essentially starts with an arbitrary disjoint filter-ideal pair $(\uparrow x, \downarrow y)$ generated by elements $x \in J^{\infty}(\mathbf{C})$ and $y \in M^{\infty}(\mathbf{C})$. It says that every such




Figure 2. The (PTi) condition illustrated.

disjoint filter-ideal pair is contained in a maximal disjoint filter-ideal pair $(\uparrow w, \downarrow z)$ where again $w \in J^{\infty}(\mathbf{C})$ and $z \in M^{\infty}(\mathbf{C})$ (here maximality is understood such that neither of $\uparrow w$ and $\downarrow z$ can be enlarged without breaking their disjointness).

Lemma 4.2. Let \mathbf{C} be a perfect lattice. If \mathbf{C} satisfies (PTi) then the RS frame $\mathbf{F}(\mathbf{C}) = (J^{\infty}(\mathbf{C}), M^{\infty}(\mathbf{C}), \leqslant)$ satisfies (Ti).

Proof. First observe that when translating the condition (Ti) from a general RS frame to $\mathbb{F}(\mathbf{C})$ we have that $xR = \uparrow x$ and $Ry = \downarrow y$. The fact that $\mathbb{F}(\mathbf{C})$ satisfies (Ti) follows then from the fact that u < w implies $u \neq w$ and $\uparrow w \subseteq \uparrow u$.

We want to characterise the condition (PTi) on the Galois closed sets arising from an RS frame $\mathbb{F} = (X, Y, R)$. The following lemma will assist us in this task.

Lemma 4.3. Consider the RS frame $\mathbb{F} = (X, Y, R)$. Then

- (i) $w \in (R_{\triangleleft} \circ R_{\triangleright})(\{x\})$ if and only if $xR \subseteq wR$;
- (ii) $(R_{\triangleleft} \circ R_{\triangleright})(\{w\}) \subseteq (R_{\triangleleft} \circ R_{\triangleright})(\{x\})$ if and only if $xR \subseteq wR$;
- (iii) $(R_{\triangleleft} \circ R_{\triangleright})(\{x\}) \subseteq Ry \text{ if and only if } xRy.$

Proof. For (i) we have

$$w \in (R_{\triangleleft} \circ R_{\triangleright})(\{x\})$$
 \Leftrightarrow $(\forall z \in R_{\triangleright}(\{x\}))(wRz)$
 \Leftrightarrow $(\forall z \in Y)(xRz \Rightarrow wRz)$
 \Leftrightarrow $xR \subseteq wR.$

To assist with the proof of (ii), note that $R_{\triangleright}(\{x\}) = Rx$ and $R_{\triangleright}(\{w\}) = Rw$. If we assume that $xR \subseteq wR$ then the fact that $R_{\triangleleft} : \mathcal{P}(Y) \to \mathcal{P}(X)$ is order-reversing gives us that $(R_{\triangleleft} \circ R_{\triangleright})(\{w\}) \subseteq (R_{\triangleleft} \circ R_{\triangleright})(\{x\})$. For the converse, if $(R_{\triangleleft} \circ R_{\triangleright})(\{w\}) \subseteq (R_{\triangleleft} \circ R_{\triangleright})(\{x\})$ then since $R_{\triangleright} : \mathcal{P}(X) \to \mathcal{P}(Y)$ is order-reversing and since $(R_{\triangleright} \circ R_{\triangleleft} \circ R_{\triangleright})(\{w\}) = R_{\triangleright}(\{w\})$, we get $xR \subseteq wR$. The statement (iii) is exactly [7, Proposition 2.6].

We want to prove that when an RS frame $\mathbb{F} = (X, Y, R)$ satisfies the (Ti) condition, the perfect lattice of Galois-closed sets $\mathcal{G}(\mathbb{F})$ satisfies (PTi). In order to make the proof easier to follow, it will be useful to translate the condition (PTi) from the setting of a general perfect lattice to the setting of $\mathcal{G}(\mathbb{F})$.

Lemma 4.4. Let $\mathbb{F} = (X, Y, R)$ be an RS frame. Assume that the following set of conditions is satisfied by $\mathcal{G}(\mathbb{F})$:

For all $x \in X$ and all $y \in Y$, if $\neg(xRy)$ then there exist $p \in X$, $q \in Y$ such that

- (i) $xR \subseteq pR$ and $Ry \subseteq Rq$
- (ii) $\neg (pRq)$
- (iii) $(\forall u \in X)(pR \subseteq uR \Rightarrow uRq)$
- (iv) $(\forall v \in Y)(Rq \subsetneq Rv \Rightarrow pRv)$

Then the lattice $\mathcal{G}(\mathbb{F})$ satisfies (PTi).

Proof. This follows using Lemma 4.3 to translate (PTi) conditions to the complete lattice $\mathcal{G}(\mathbb{F})$.

Lemma 4.5. Let $\mathbb{F} = (X, Y, R)$ be an RS frame. If \mathbb{F} satisfies (Ti) then $\mathcal{G}(\mathbb{F})$ satisfies (PTi).

Proof. Let $\mathbb{F} = (X, Y, R)$ be an RS frame satisfying (Ti) (i.e. a TiRS frame). Take arbitrary $x \in X$ and $y \in Y$ and assume that $\neg(xRy)$. In the perfect lattice $\mathcal{G}(\mathbb{F})$ coming from \mathbb{F} consider the sets $A = (R_{\triangleleft} \circ R_{\triangleright})(\{x\})$ and B = Ry. Then $A \in J^{\infty}(\mathcal{G}(\mathbb{F}))$, $B \in M^{\infty}(\mathcal{G}(\mathbb{F}))$ and $A \nsubseteq B$ using Lemma 4.3(iii).

We have that

$$(R_{\triangleleft} \circ R_{\triangleright})(\{x\}) \nsubseteq Ry \quad \Rightarrow \quad (\exists w \in X)(xR \subseteq wR \& \neg (wRy))$$

$$\Rightarrow \quad (\exists w \in X) \Big[xR \subseteq wR \& \Big]$$

$$(\exists p \in X)(\exists q \in Y) \Big(\neg (pRq) \& wR \subseteq pR \& Ry \subseteq Rq \\ \& (\forall u \in X)(pR \subsetneq uR \Rightarrow uRq) \Big]$$

$$\& (\forall v \in Y)(Rq \subsetneq Rv \Rightarrow pRv) \Big) \Big]$$

The only part of the (PTi) condition for $\mathcal{G}(\mathbb{F})$ that is not now immediate is the fact that we need $xR \subseteq pR$. This follows from the $xR \subseteq wR \subseteq pR$ and the transitivity of set containment.

Now we are ready to show that the canonical extensions of lattices are PTi lattices and so they indeed are 'more' than just perfect lattices. For this we cite our final result from [2]:

Proposition 4.6 ([2, Corollary 3.11]). Let **L** be a bounded lattice and **X** = $D^{\flat}(\mathbf{L})$ be its dual TiRS graph. Let $\rho(\mathbf{X})$ be the frame associated to **X** and $G(\rho(\mathbf{X}))$ be its corresponding perfect lattice of Galois-closed sets.

The lattice $G(\rho(\mathbf{X}))$ is the canonical extension of \mathbf{L} .

The result can be illustrated by the diagram in Fig. 3. The given bounded lattice \mathbf{L} is firstly assigned its Ploščica dual space $\mathrm{D}(\mathbf{L}) = (\mathcal{L}^{\mathrm{mp}}(\mathbf{L}, \underline{\mathbf{2}}), E, \mathcal{T})$, and then the Ploščica dual graph $\mathbf{X} = \mathrm{D}^{\flat}(\mathbf{L}) = (\mathcal{L}^{\mathrm{mp}}(\mathbf{L}, \underline{\mathbf{2}}), E)$ is obtained by forgetting the topology. This is a TiRS graph and so the frame $\rho(\mathbf{X})$ associated to \mathbf{X} in our one-to-one correspondence developed in [2] between TiRS graphs and TiRS frames is a TiRS frame. Hence by Lemma 4.5 above, the perfect lattice $\mathrm{G}(\rho(\mathbf{X}))$ of Galois-closed sets corresponding in Gehrke's representation to the frame $\rho(\mathbf{X})$ is a PTi lattice. By Proposition 4.6, the lattice $\mathrm{G}(\rho(\mathbf{X}))$ is the canonical extension of the given lattice \mathbf{L} .




FIGURE 3. Ploščica and Gehrke in tandem.

Hence we have our final result of this section:

Theorem 4.7. The canonical extension of a bounded lattice is a PTi lattice.

Gehrke and Vosmaer [10] showed that the canonical extension of a lattice need not be meet-continuous, and hence need not always be algebraic. Theorem 4.7 gives us further information about the structure of canonical extensions of bounded lattices.

5. Examples

Our goal in this section is to illustrate that the PTi condition adds to the current description of the canonical extension of a bounded lattice. We focus on non-distributive examples. Canonical extensions of distributive lattices are known to be completely distributive complete lattices. To show that our new condition does indeed add to the current description, we give an example of a perfect lattice that is not PTi. Giving an example of a PTi lattice that is not the canonical extension of a lattice would be the same as giving an example of a TiRS graph that is not of the form $(\mathcal{L}^{mp}(\mathbf{L}, \mathbf{2}), E)$ for some bounded lattice \mathbf{L} . Hence this is the same as the representable TiRS graph (representable poset) problem.

Our goals are:

- (1) Give an example of a complete non-distributive lattice which is a PTi lattice but is not the canonical extension of any bounded lattice.
- (2) Give an example of a perfect non-distributive lattice that is not a PTi lattice.




FIGURE 4. A TiRS graph that is not the graph of MPH's of any bounded lattice (left) and its dual PTi lattice A_L that is not a canonical extension (right). The double-headed arrows on the graph emphasize that transitivity holds amongst the vertical edges.

Example 5.1. Consider the complete lattice A_L depicted on the right in Fig. 4. We will denote by m, the middle element of the infinite chain $\omega \oplus \mathbf{1} \oplus \omega^{\partial}$ and y is above the bottom and below the top but incomparable with all other elements.

The TiRS graph dual to A_L is $X = \{p_i \mid i \in \omega\} \cup \{q_j \mid j \in \omega\} \cup \{k\}$ with the relation E given by

$$p_0 < p_1 < p_2 < \ldots < p_n < p_{n+1} < \ldots < q_{n+1} < q_n < q_{n-1} \ldots < q_1 < q_0$$

 $\cup \{(k, p_i) \mid i \in \omega\} \cup \{(k, q_j) \mid j \geqslant 1\}$

(it is depicted on the left in Fig. 4). To be clear, the p_i 's and q_j 's form a poset (it is transitive) that is order-isomorphic to $\omega \oplus \omega^{\partial}$ while the element k is related to everything except the top of the chain.

Recall that MPE's are ordered by: $\varphi \leq \psi$ if and only if $\varphi^{-1}(1) \subseteq \psi^{-1}(1)$. The MPE's φ_1 and φ_0 are defined by $\varphi_1(x) = 1$ and $\varphi_0(x) = 0$ for all $x \in X$. All but one of the other MPE's have $k \mapsto 0$ and then they split the chain at some point. When φ splits the chain by sending the p_i 's to 0 and the q_j 's to 1 then you get the limit point in the middle of A_L . The interesting MPE is the map does the following for $a \in X$:

$$\varphi(a) = \begin{cases} 1 & \text{if } a = k \\ 0 & \text{if } a = q_0 \\ - & \text{otherwise} \end{cases}$$

This interesting MPE is the incomparable point that makes A_L non-distributive.




FIGURE 5. An RS frame that is not Ti (left) and its dual perfect lattice that is not PTi (right).

It is quite easy to show that the lattice A_L is a PTi lattice; we indicated on the right in Fig. 4 what the elements $w \in J^{\infty}(A_L)$ and $z \in M^{\infty}(A_L)$ are for the chosen elements $x \in J^{\infty}(A_L)$ and $y \in M^{\infty}(A_L)$. The fact that the lattice A_L is not the canonical extension of any bounded lattice is harder to show and it follows from Proposition 5.2 below.

Proposition 5.2. There is no bounded lattice L and lattice embedding $e: L \to A_L$ such that (e, A_L) is the canonical extension of L.

Proof. Suppose there are no bounded lattice L and an embedding $e : L \to A_L$ such that (e, A_L) is the canonical extension of L. Clearly the top and bottom element of A_L are, respectively e(1) and e(0) where 1 and 0 are the top and bottom element of L. Now consider the set of elements $(A_L) \setminus \{e(0), e(1), m\}$. It is easy to see that each of these elements is completely join-irreducible in A_L and hence we have, by Lemma 2.1, that each of these elements is the meet of the embedding of a filter of L. Hence each of the elements of $(A_L) \setminus \{e(0), e(1), m\}$ is a filter element. Dually, it is easy to see that each element of $(A_L) \setminus \{e(0), e(1), m\}$ is completely meet-irreducible and again by Lemma 2.1 they are all the join of the embedding of an ideal of L and hence are all ideal elements. Thus every element of $(A_L) \setminus \{e(0), e(1), m\}$ is both ideal and filter and hence must be of the form e(a) for some $a \in L$. Now consider the element m. Since $m = \bigwedge \omega^{\partial}$, and since every element of ω^{∂} is the image of an element of L under e, we have that m is a filter element of A_L . Also, $m = \bigvee \omega$ and every element of ω is the image of an element of L under e. Therefore m is also an ideal element of A_L . Hence m must be of the form e(b) for some $b \in L$. Thus we have that $\mathbf{L} \cong A_L$ and that the embedding e is a bijection.

Now we show that (e, A_L) cannot be the canonical extension of **L**. Observe that since $m = \bigwedge \omega^{\partial} = \bigvee \omega$ we have that $\bigwedge \omega^{\partial} \leqslant \bigvee \omega$. However, for any finite subset $A' \subseteq \omega^{\partial}$ and any finite subset $B' \subseteq \omega$ we will have $\bigvee B' < \bigwedge A'$. Hence (e, A_L) is not a compact completion of **L**.

Example 5.3. We consider the complete lattice M_L depicted on the right in Fig. 5. The order is given by the poset $\mathbf{1} \oplus \omega$ with an additional element y incomparable to all elements except the top and the bottom. It can easily be seen that M_L is a perfect lattice $(J^{\infty}(M_L) = M^{\infty}(M_L) = \{y\} \cup \{x_i \mid i \geqslant 1\})$. It is not PTi since there are no w and z for the pair $x_j \nleq y \ (j \geqslant 1)$.

The RS frame corresponding to it was already mentioned in [2, page 128] as an example of an RS frame which is not TiRS (it is indicated on the left in Fig. 5): Let $X_1 = \{a_i\}_{i \in \omega}$, $X_2 = \{b_i\}_{i \in \omega}$ and let

$$R = \{(a_1, b_0), (a_0, b_1)\} \cup \{(a_i, b_i) \mid 2 \leq i, 1 \leq j \leq i\}.$$

By considering $\neg(a_0Rb_0)$ it is rather straightforward to show that (X_1, X_2, R) does not satisfy (Ti).




FIGURE 6. The correspondences between classes of PTi lattices and classes of TiRS graphs.

Our final picture Fig. 6 describes the correspondence between PTi lattices and TiRS graphs and between their important subclasses: (i) the canonical extensions of bounded lattices inside the PTi lattices and representable graphs (as dual graphs of bounded lattices) inside the TiRS graphs; (ii) the canonical extensions of bounded distributive lattices inside the canonical extensions of bounded lattices and representable posets (as dual graphs of bounded distributive lattices) inside the representable graphs.

A natural question that we asked already in [2, pages 126–127] was which TiRS graphs arise as duals of bounded lattices. In the case of bounded distributive lattices (denoted as BDL's in Fig. 6) this question reduces to the question of which posets are representable posets which seems to be extremely hard. Examples of non-representable posets are also examples of non-representable graphs as any poset is automatically a TiRS graph. We mention an example of a non-representable poset due to Tan [17] from the 1970s: $T := \omega \oplus \omega^{\delta}$. The perfect lattice corresponding to this TiRS graph is the PTi lattice $T_L := \omega \oplus \mathbf{1} \oplus \omega^{\delta}$.

Acknowledgements

The first author gratefully acknowledges the hospitality of Matej Bel University during his research visit in September 2017. The third author acknowledges support from Slovak grant VEGA 1/0337/16 and the hospitality of the University of Lisbon during his visit in September 2019.

References

- [1] Clark D.M., Davey, B.A.: Natural Dualities for the Working Algebraist, Cambridge University Press (1998)
- [2] Craig, A.P.K, Gouveia, M.J., Haviar, M.: TiRS graphs and TiRS frames: a new setting for duals of canonical extensions, Algebra Universalis 74, 123–138 (2015)
- [3] Craig, A.P.K., Haviar, M., Priestley, H.A.: A fresh perspective on canonical extensions for bounded lattices, Appl. Categ. Structures 20, 725–749 (2013)
- [4] Craig, A.P.K., Haviar, M.: Reconciliation of approaches to the construction of canonical extensions of bounded lattices, Math. Slovaca 64, 1335–1356 (2014)
- [5] Deiters, K., Erné, M.: Sums, products and negations of contexts and complete lattices, Algebra Universalis **60**, 469–496 (2009)
- [6] Dunn, J.M., Gehrke, M., Palmigiano, A.: Canonical extensions and relational completeness of some substructural logics, J. Symbolic Logic 70, 713-740 (2005)
- [7] Gehrke, M.: Generalized Kripke frames, Studia Logica 84, 241–275 (2006)
- [8] Gehrke, M.: Canonical extensions: an algebraic approach to Stone duality, Algebra Universalis 79: 63 (2018)
- [9] Gehrke, M., Harding, J.: Bounded lattice expansions, J. Algebra 238, 345–371 (2001)
- [10] Gehrke, M., Vosmaer, J.: A view of canonical extension. In: Proceedings of the Eighth International Tbilisi Symposium, TbiLLC, 2009. Lecture Notes in Computer Science vol. 6618, pp. 77–100. Logic, Language and Computation (Tbilisi, 2011)
- [11] Goldblatt, R.: Canonical extensions and ultraproducts of polarities, Algebra Universalis 79: 80 (2018)
- [12] Hartonas, C., Dunn, J.M.: Stone duality for lattices, Algebra Universalis 37, 391–401 (1997)
- [13] Jónsson, B., Tarski, A.: Boolean algebras with operators, I & II, Amer. J. Math. 73, 891–939 (1951) & 74, 127–162 (1952)
- [14] Jipsen, P., Categories of algebraic contexts equivalent to idempotent semirings and domain semirings, RAMiCS 2012, LNCS 7560, 195–206 (2012)
- [15] Moshier, M.A.,: A relational category of formal contexts, Pre-print.
- [16] Ploščica, M.: A natural representation of bounded lattices, Tatra Mountains Math. Publ. 5, 75–88 (1995)
- [17] Tan, T.: On representable posets. PhD Thesis, University of Manitoba (1974)
- [18] Urquhart, A.: A topological representation theory for lattices, Algebra Universalis $\bf 8$, $\bf 45-58$ (1978)

Andrew P. K. Craig
Department of Mathematics and Applied Mathematics
University of Johannesburg
PO Box 524, Auckland Park, 2006
South Africa

e-mail: acraig@uj.ac.za

Maria J. Gouveia

e-mail: mjgouveia@fc.ul.pt

Faculdade de Ciências da Universidade de Lisboa

P-1749-016 Lisboa

Portugal

Miroslav Haviar Department of Mathematics Faculty of Natural Sciences, M. Bel University Tajovského 40, 974 01 Banská Bystrica Slovakia e-mail: miroslav.haviar@umb.sk