

CONGRUENCES FOR COEFFICIENTS OF MODULAR FUNCTIONS IN LEVELS 3, 5, AND 7 WITH POLES AT 0

PAUL JENKINS AND RYAN KECK

ABSTRACT. We give congruences modulo powers of $p \in \{3, 5, 7\}$ for the Fourier coefficients of certain modular functions in level p with poles only at 0, answering a question posed by Andersen and the first author and continuing work done by the authors and Moss. The congruences involve a modulus that depends on the base p expansion of the modular form's order of vanishing at ∞ .

1. INTRODUCTION

A modular form $f(z)$ of level N and weight k is a complex valued function which is holomorphic on the upper half plane, satisfies the equation

$$f\left(\frac{az+b}{cz+d}\right) = (cz+d)^k f(z) \text{ for all } \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(N),$$

and is holomorphic at the cusps of $\Gamma_0(N)$. Letting $q = e^{2\pi iz}$, modular forms have a Fourier expansion $f(z) = \sum_{n \geq 0} a(n)q^n$ with Fourier coefficients $a(n)$. A *weakly holomorphic* modular form is a modular form that is allowed to be meromorphic at the cusps; we define $M_k^\sharp(N)$ to be the space of weakly holomorphic modular forms of weight k and level N that are holomorphic away from the cusp at ∞ , and in the same notation as [7], we use $M_k^b(N)$ to denote forms holomorphic away from the cusp at 0. For prime N , these are the only cusps. Modular forms in both of these spaces also have Fourier expansions at infinity, where the constraint $n \geq 0$ is relaxed to $n \gg -\infty$ if the form has a pole at infinity.

The Fourier coefficients of modular forms often satisfy interesting congruences. For the j -invariant $j(z) = q^{-1} + 744 + \sum_{n=1}^{\infty} c(n)q^n \in M_0^\sharp(1)$, Lehner [11, 12] proved that the $c(n)$ satisfy the congruence

$$c(2^a 3^b 5^c 7^d n) \equiv 0 \pmod{2^{3a+8} 3^{2b+3} 5^{c+1} 7^d} \text{ if } a, b, c, d \geq 1.$$

Kolberg [9, 10], Aas [1], and Allatt and Slater [2] strengthened Lehner's congruences for $j(z)$. Furthermore, Griffin [6] gave a canonical basis for $M_0^\sharp(1)$ and extended Kolberg's and Aas's results to the basis elements. Similarly, the first author, Andersen, and Thornton [3, 8] proved congruences for the Fourier coefficients of elements of canonical bases for $M_0^\sharp(p)$ with $p = 2, 3, 5, 7$. The authors and Moss [7] proved congruences for the Fourier coefficients of elements of a canonical basis for $M_0^b(2)$. It is natural to wonder whether a similar result holds for bases of $M_0^b(p)$ for the other genus zero primes, mirroring the results of $M_0^\sharp(p)$.

Date: March 13, 2024.

2010 Mathematics Subject Classification. Primary 11F30, 11F37.

This work was partially supported by a grant from the Simons Foundation (#281876 to Paul Jenkins).

Let $p \in \{2, 3, 5, 7, 13\}$. Taking $\eta(z) = q^{\frac{1}{24}} \prod_{n=1}^{\infty} (1 - q^n)$ to be the Dedekind eta function, a Hauptmodul for $\Gamma_0(p)$ is

$$\phi^{(p)}(z) = \left(\frac{\eta(pz)}{\eta(z)} \right)^{24/(p-1)} = q + \frac{24}{p-1}q^2 + \cdots,$$

which vanishes at ∞ and has a pole only at 0. The functions $(\phi^{(p)}(z))^m$ for $m \geq 0$ are a basis for $M_0^b(p)$. Andersen and the first author used powers of $\phi^{(p)}(z)$ to prove congruences involving $\psi^{(p)} = \frac{1}{\phi^{(p)}} = q^{-1} - \frac{24}{p-1} + \cdots \in M_0^{\sharp}(p)$ in [3], and made the following remark: “Additionally, it appears that powers of the function $[\phi^{(p)}(z)]$ have Fourier coefficients with slightly weaker divisibility properties... It would be interesting to more fully understand these congruences.” The authors with Moss in [7] proved congruences for the Fourier coefficients of $\phi^{(2)}(z)$ and its powers. In this paper, we use similar techniques to obtain congruences for $\phi^{(p)}(z)$ and its powers for $p = 3, 5, 7$.

Write $\phi^{(p)}(z)^m = \sum_{n=m}^{\infty} a^{(p)}(m, n)q^n$. Let χ_S be the characteristic function on S , which outputs 1 when the input is an element of S and 0 otherwise. The main result of this paper is the following theorem.

Theorem 1. *Let $p \in \{3, 5, 7\}$. Let $n = p^\alpha n'$ where $p \nmid n'$. Express the base p expansion of m as $a = \sum_{i=1}^{\infty} a_i p^{i-1}$, where $a_i = 0$ for all sufficiently large i . Consider the rightmost α digits $a_\alpha \dots a_2 a_1$. Let i' be the index of the rightmost nonzero digit, or $i' = -1$ if $a_1 = a_2 = \dots = a_\alpha = 0$. Let*

$$\gamma_3(m, \alpha) = \begin{cases} 3 - a_{i'} + 2\#\{i \mid a_i = 0, i > i'\} + \#\{i \mid a_i = 1, i > i'\} & \text{if } i' \geq 0, \\ 0 & \text{otherwise.} \end{cases}$$

$$\gamma_5(m, \alpha) = \gamma_7(m, \alpha) = \begin{cases} \chi_{\{1,2\}}(a_{i'}) + \#\{i \mid a_i \in \{0, 1\}, i > i'\} & \text{if } i' \geq 0, \\ 0 & \text{otherwise.} \end{cases}$$

Then

$$a^{(p)}(m, p^\alpha n') \equiv 0 \pmod{p^{\gamma_p(m, \alpha)}}.$$

The power of p in the congruence includes a count of the number of digits in the base p expansion of m that are 0, 1, or 2. This result is similar to the one found for $\phi^{(2)}$, but it is more complicated to state because there are more digits in bases 3, 5, 7. We note that this congruence is not sharp. For $m = 1$, Allatt and Slater in [2] proved a stronger result that provides an exact congruence for many n .

As an example, the base 3 expansion for $m = 102$ is $m = \dots 00010210$. Table 1 gives values of γ_3 . Notice that once α surpasses 5—the leftmost nonzero digit in the base 3 expansion of m occurs in the 5th place— γ_3 always increases by 2 as α increases by 1. This illustrates that $\gamma_3(m, \alpha)$ is unbounded for a fixed m . Similar examples can be constructed for γ_5 and γ_7 .

α	0	1	2	3	4	5	6	7	8	9	...	α	...
$\gamma(102, \alpha)$	0	0	2	2	4	5	7	9	11	13	...	$2(\alpha - 5) + 5$...

TABLE 1. Values of $\gamma_3(m, \alpha)$ for $m = 102$

Section 2 contains the machinery and definitions we use for proving Theorem 1 and the proof itself is in Section 3. In Section 4 we discuss the $p = 13$ case; although 13 is also a genus zero prime, we do not obtain congruences modulo 13.

We thank the referee for helpful comments which helped to improve this paper.

2. PRELIMINARY LEMMAS

The operator U_p on a function $f(z)$ is given by

$$U_p f(z) = \frac{1}{p} \sum_{j=0}^{p-1} f\left(\frac{z+j}{p}\right).$$

If $M_k^!(N)$ is the space of weakly holomorphic modular forms of weight k and level N , meaning we allow poles at any cusp, then we have $U_p : M_k^!(N) \rightarrow M_k^!(N)$ if p divides N . If $f(z)$ has the Fourier expansion $\sum_{n=n_0}^{\infty} a(n)q^n$, then the effect of U_p on the Fourier expansion is given by $U_p f(z) = \sum_{n=n_0}^{\infty} a(pn)q^n$.

We prove Theorem 1 by first proving several facts about $\phi^{(p)}$ and its image under the U_p operator. Most importantly, $U_p(\phi^{(p)})$ is a polynomial with integer coefficients in $\phi^{(p)}$. If all of the coefficients are divisible by some number, then every p th coefficient of $\phi^{(p)}$ is divisible by that number. The same holds true when applying U_p to any power of $\phi^{(p)}$ as well.

The following result describes how U_p applied to a modular function behaves under the Fricke involution. This will help us in Lemma 5 to write $U_p(\phi^{(p)})^m$ as a polynomial in $\phi^{(p)}$.

Lemma 2 ([4, Theorem 4.6]). *Let p be prime and let $f(z)$ be a level p modular function. Then*

$$p(U_p f)\left(\frac{-1}{pz}\right) = p(U_p f)(pz) + f\left(\frac{-1}{p^2 z}\right) - f(z).$$

The Fricke involution $\begin{pmatrix} 0 & -1 \\ p & 0 \end{pmatrix}$ swaps the cusps of $\Gamma_0(p)$, which are 0 and ∞ . We will use this fact in the proof of Lemma 5, and the following relations between $\phi^{(p)}(z)$ and $\psi^{(p)}(z)$ will help us compute this involution.

Lemma 3 ([3, Lemma 3]). *The functions $\phi(z)$ and $\psi(z)$ satisfy the relations*

$$\begin{aligned} \phi^{(p)}\left(\frac{-1}{pz}\right) &= p^{-12/(p-1)}\psi^{(p)}(z), \\ \psi^{(p)}\left(\frac{-1}{pz}\right) &= p^{12/(p-1)}\phi^{(p)}(z). \end{aligned}$$

The following lemma is a special case of a result of Lehner [12]. It provides a polynomial with functions as its coefficients whose roots are modular forms used in the proof of Theorem 7.

Lemma 4 ([12, Theorem 2]). *There exist integers $b_j^{(p)}$ such that*

$$U_p \phi^{(p)}(z) = p \sum_{j=1}^p b_j^{(p)} \phi^{(p)}(z)^j.$$

Furthermore, let $h^{(p)}(z) = p^{12/(p-1)}\phi^{(p)}(z/p)$. Then

$$(h^{(p)}(z))^p + \sum_{j=1}^p (-1)^j g_j(z) (h^{(p)}(z))^{p-j} = 0$$

where

$$g_j(z) = (-1)^{j+1} p^{12/(p-1)+2} \sum_{\ell=j}^p b_\ell^{(p)} \phi^{(p)}(z)^{\ell-j+1}.$$

In the following lemma, we extend the result from the first part of Lemma 4, writing $U_p(\phi^{(p)})^m$ as an integer polynomial in $\phi^{(p)}$. In particular, we give the degree of that polynomial. An alternative approach can be seen in [5, Lemma 4.1.1].

Lemma 5. *For all $m \geq 1$, $U_p(\phi^{(p)})^m \in \mathbb{Z}[\phi^{(p)}]$. In particular,*

$$U_p(\phi^{(p)})^m = \sum_{j=\lceil m/p \rceil}^{pm} d(m, j) (\phi^{(p)})^j$$

where $d(m, j) \in \mathbb{Z}$, and $d(m, pm)$ is not 0.

Proof. We proceed as in [7, Lemma 5]; this is a straightforward generalization from 2 to p . Using Lemmas 2 and 3, we have that

$$\begin{aligned} U_p \phi^{(p)}(-1/pz)^m &= U_p \phi^{(p)}(pz)^m + p^{-1} \phi^{(p)}(-1/p^2z)^m - p^{-1} \phi^{(p)}(z)^m \\ &= U_p \phi^{(p)}(pz)^m + p^{-1-12m/(p-1)} \psi^{(p)}(pz)^m - p^{-1} \phi^{(p)}(z)^m \\ &= p^{-1-12m/(p-1)} q^{-pm} + O(q^{-pm+p}). \end{aligned}$$

Thus,

$$p^{1+12m/(p-1)} U_p \phi^{(p)}(-1/pz)^m = q^{-pm} + O(q^{-pm+p}).$$

Because $\phi^{(p)}(z)^m$ is holomorphic at ∞ , $U_p \phi^{(p)}(z)^m$ is holomorphic at ∞ . So $U_p \phi^{(p)}(-1/pz)^m$ is holomorphic at 0 and, since the Fourier expansion starts with q^{-pm} , it must be a polynomial of degree pm in $\psi^{(p)}(z)$. Let $b(m, j) \in \mathbb{Z}$ such that

$$p^{1+12m/(p-1)} U_p \phi^{(p)}(-1/pz)^m = \sum_{j=0}^{pm} b(m, j) \psi^{(p)}(z)^j,$$

and we note that $b(m, pm)$ is not 0. Now replace z with $-1/pz$ and use Lemma 3 to get

$$p^{1+12m/(p-1)} U_p \phi^{(p)}(z)^m = \sum_{j=0}^{pm} b(m, j) p^{12j/(p-1)} \phi^{(p)}(z)^j,$$

which gives

$$U_p \left(\phi^{(p)}(z)^m \right) = \sum_{j=0}^{pm} b(m, j) p^{12(j-m)/(p-1)-1} \phi^{(p)}(z)^j.$$

Because $(\phi^{(p)}(z))^m = q^m + \dots$, if m is divisible by p , the leading term of the above sum is $q^{m/p}$, and otherwise the smallest power of q present in the polynomial is at least $\lceil m/p \rceil$, so the sum starts with $j = \lceil m/p \rceil$ as desired. Notice that $b(m, j) p^{12(j-m)/(p-1)-1}$ is an integer because the coefficients of $\phi^{(p)}(z)^m$ are integers. \square

We may repeatedly use Lemma 5 to write $U_p^\alpha(\phi^{(p)})^m$ as a polynomial in $\phi^{(p)}$. Let

$$(2.1) \quad f_{(p)}(\ell) = \lceil \ell/p \rceil, \quad f_{(p)}^0(\ell) = \ell, \quad \text{and} \quad f_{(p)}^\alpha(\ell) = f_{(p)}(f_{(p)}^{\alpha-1}(\ell)) \quad \text{for } \alpha \geq 1.$$

Using Lemma 5, the smallest exponent of q appearing in $U_p^\alpha(\phi^{(p)})^m$ is at least $f_{(p)}^\alpha(m)$.

Lemma 6 provides a connection between $\gamma_p(m, \alpha)$ and the integers $f_{(p)}^\alpha(m)$: γ_p is counting the number of 0s and 1s to the left of the first nonzero digit in the base p expansion of m . The key difference between the following lemma and its corresponding lemma in [7] is that there are more digits in bases 3, 5, 7.

Lemma 6. *The number of 0s to the left of the rightmost nonzero digit in the first α digits of the base p expansion of m is equal to the number of integers congruent to 1 modulo p in the list*

$$m, f_{(p)}(m), f_{(p)}^2(m), \dots, f_{(p)}^{\alpha-1}(m),$$

except when the rightmost nonzero digit is 1 (in which case there is exactly one more in the list). Similarly, the number of 1s to the left of the rightmost nonzero digit in the base p expansion of m is equal to the number of integers congruent to 2 modulo p in the above list, again with the exception of when the rightmost nonzero digit is 2.

Proof. Write the base p expansion of m as $a_r \dots a_2 a_1$, and consider its first α digits, $a_\alpha \dots a_2 a_1$, where $a_i = 0$ for $i > r$ if $\alpha > r$. If $a_i = 0$ for $1 \leq i \leq \alpha$, then all of the integers in the list are zero modulo p . Otherwise, suppose that $a_i = 0$ for $1 \leq i < i'$ and $a_{i'} \neq 0$. Apply $f_{(p)}$ repeatedly to m . Each application of $f_{(p)}$ deletes the rightmost 0 from the expansion, until $a_{i'}$ is the rightmost remaining digit; that is, $f_{(p)}^{i'-1}(m) = a_\alpha \dots a_{i'+1} a_{i'}$. In particular, the rightmost digit is nonzero. Having reduced to this case, we now treat only the case where m is not divisible by p .

If m is not divisible by p , and $a_1 \in \{1, 2\}$, then at least one number in the list, namely m , is congruent to either 1 or 2 modulo p . Also, $f_{(p)}(m) = \lceil m/p \rceil = (m+a)/p$ for some $a \neq 0$. Applied to the base p expansion of m , $f_{(p)}$ deletes a_1 and propagates a 1 leftward through the base p expansion, replacing any digit that was previously equal to $p-1$ with zero. This is essentially the operation of carrying in addition. This process then terminates upon encountering the rightmost digit less than $p-1$ (if it exists), which becomes one greater. As in the case where p divides m , we apply f repeatedly to delete the new leading 0s. But if the first nonzero digit to the left was either a 0 or a 1 before we propagated a 1 leftward, it is now a 1 or a 2 respectively. So now when we repeat this process until all digits are accounted for, we notice that any digit that was either a 0 or a 1 becomes a 1 or a 2 respectively (with the exception of the first nonzero digit), which proves the lemma. \square

3. PROOF OF THE MAIN THEOREM

Theorem 1 will follow from the next theorem.

Theorem 7. *Let $f_{(p)}^\alpha(m)$ be as in (2.1). Let $\gamma_p(m, \alpha)$ be as in Theorem 1, and let $\alpha \geq 1$. Define $P^{(p)}(\ell, a)$ to be the set of polynomials in $\phi^{(p)}$ with lowest power ℓ*

having coefficient d_ℓ divisible by p^a and each subsequent coefficient d_k being divisible by at least $p^{\delta_p(k-\ell)+a}$, where $\delta_3 = 4$ and $\delta_5 = \delta_7 = 1$. Then

$$(3.1) \quad U_p^\alpha(\phi^{(p)})^m \in P^{(p)}(f_{(p)}^\alpha(m), \gamma_p(m, \alpha)).$$

Since every coefficient of such a polynomial is divisible by $p^{\gamma_p(m, \alpha)}$, this shows that the coefficient of $q^{p^\alpha n}$ in $(\phi^{(p)})^m$ must vanish modulo $p^{\gamma_p(m, \alpha)}$. Note that these methods do not give meaningful congruences for the case when $\alpha = 0$.

Theorem 7 is an improvement on the following result by Lehner [12] for $p = 3$.

Theorem 8. [12, Equation 3.24] Write $U_3^\alpha(\phi^{(3)})^m$ as $\sum d(m, j, \alpha)(\phi^{(3)})^j \in \mathbb{Z}[\phi^{(3)}]$. For any integer k , let $\nu_3(k)$ be the highest power of 3 dividing k . Then

$$\nu_3(d(m, j, \alpha)) \geq 4(j-1) + \alpha(2 - 4(1-m)).$$

In particular, Lehner's bound sometimes only gives the trivial result that the 3-adic valuation of $d(m, j, \alpha)$ is greater than some negative integer. Lehner also proved congruences for $p = 5, 7$, but they experience similar issues [11].

In proving Theorem 7, we use techniques from Lemmas 5 and 6 of [3]. From the definition of U_p , we have

$$(3.2) \quad U_p \phi^{(p)}(z)^m = p^{-1} \sum_{j=1}^p \phi^{(p)}\left(\frac{z+j}{p}\right)^m.$$

Define $h_\ell^{(p)}(z) = p^{12/(p-1)} \phi^{(p)}\left(\frac{z+\ell}{p}\right)$. Consider the polynomials

$$F^{(p)}(x) = x^p + \sum_{j=1}^{p-1} (-1)^j g_j(z) x^{p-j}$$

with the $g_j(z)$ functions from Lemma 4 as coefficients. We claim that $F^{(p)}(x)$ has the $h_\ell^{(p)}(z)$ as roots. By Lemma 4, we know that $h_0^{(p)}(z)$ is a root; the others are roots because the $g_j(z)$ are fixed under $z \mapsto z+1$, but for this transformation of z , $h_\ell^{(p)}(z)$ gets sent to $h_{\ell+1}^{(p)}(z)$.

Since $F^{(p)}$ has the functions $h_\ell^{(p)}(z)$ as its roots, the coefficients $g_j(z)$ are the symmetric polynomials in the roots. We can now use Newton's identities for the sums of powers of roots of a polynomial. Writing $F^{(p)}(x) = \prod_{i=1}^n (x - x_i)$, let $S_\ell = x_1^\ell + \cdots + x_n^\ell$. It follows that

$$S_\ell = g_1 S_{\ell-1} - g_2 S_{\ell-2} + \cdots + (-1)^{\ell+1} \ell g_\ell.$$

Let $R^{(p)}$ be the set of polynomials of the form $\sum_{n=1}^N d_n \phi^{(p)}(z)^n$, where $d_n \in \mathbb{Z}$ and where for $n \geq 2$, $\nu_p(d_n) \geq \delta_p(n-1)$, where δ_p is as in Theorem 7. We will need the following $R^{(p)}$ product lemma, which implies that when multiplying two polynomials in $R^{(p)}$, we must multiply in δ_p extra copies of p for the product to be in $R^{(p)}$.

Lemma 9. ($R^{(p)}$ product lemma) If $f, g \in R^{(p)}$, then $p^{\delta_p} fg \in R^{(p)}$.

Proof. We only need to prove this for the product

$$p^{\delta_p} \left(p^{\delta_p(i-1)} d_i \phi^{(p)}(z)^i \right) \left(p^{\delta_p(j-1)} d'_j \phi^{(p)}(z)^j \right),$$

since then the lemma will hold for the product of any two polynomials by linearity, as the sum of any two elements in $R^{(p)}$ is clearly also an element of $R^{(p)}$. Observe that

$$\begin{aligned} p^{\delta_p} \left(p^{\delta_p(i-1)} d_i \phi^{(p)}(z)^i \right) \left(p^{\delta_p(j-1)} d'_j \phi^{(p)}(z)^j \right) &= p^{\delta_p(i-1+j-1+1)} d_i d'_j \phi^{(p)}(z)^{i+j} \\ &= p^{\delta_p((i+j)-1)} (d_i d'_j) \phi^{(p)}(z)^{i+j}, \end{aligned}$$

which is clearly an element of $R^{(p)}$. \square

We prove Theorem 7 by first showing the theorem holds when $\alpha = 1$. This is comparable to Lemma 6 from [3], which gives a subring of $\mathbb{Z}[\phi]$ which is closed under the U_p operator. Here, we employ a similar technique to prove divisibility properties of the polynomial coefficients in Lemma 5. We then show that applying U_p to a polynomial in the set $P^{(p)}(f_{(p)}^\alpha(m), \gamma_p(m, \alpha))$ will carry it to the set $P^{(p)}(f_{(p)}^{\alpha+1}(m), \gamma_p(m, \alpha + 1))$, which we refer to as the polynomial step. Implicitly, this proves the result by induction. This structure differs from [7] because it allows us to prove the polynomial step in a much cleaner way. Another approach to proving the base case can be found in [5, Lemma 4.1.1]. In this paper, we will only treat the case $p = 3$, but the computations are similar for $p = 5, 7$. Therefore, we will simply use ϕ in place of $\phi^{(3)}$.

Let $\alpha = 1$. We seek to prove the statement

$$U_p(\phi)^m = \sum_{j=\lceil m/p \rceil}^{pm} d(m, j) (\phi)^j$$

with

$$(3.3) \quad p^{\delta_p(j - \lceil m/p \rceil) + c_m^{(p)}} \mid d(m, j)$$

where

$$c_m^{(3)} = \begin{cases} 2 & m \equiv 1 \pmod{3}, \\ 1 & m \equiv 2 \pmod{3}, \\ 0 & \text{otherwise,} \end{cases}$$

$$c_m^{(5)} = c_m^{(7)} = \begin{cases} 1 & m \equiv 1, 2 \pmod{p}, \\ 0 & \text{otherwise.} \end{cases}$$

We prove (3.3) by induction on m .

Given Lemma 5 and the statement we are trying to prove, we see that proving (3.3) is equivalent to proving the following three statements:

$$\begin{aligned} m \equiv 0 \pmod{3} &: \exists r \in R^{(3)}, U_3 \phi^m = 3^{-4(\lceil m/3 \rceil - 1)} r, \\ m \equiv 1 \pmod{3} &: \exists r \in R^{(3)}, U_3 \phi^m = 3^{-4(\lceil m/3 \rceil - 1) + 2} r, \\ m \equiv 2 \pmod{3} &: \exists r \in R^{(3)}, U_3 \phi^m = 3^{-4(\lceil m/3 \rceil - 1) + 1} r. \end{aligned}$$

Furthermore, by equation (3.2) and considering the functions $h_\ell^{(3)}$, we see that $U_3 \phi^m = 3^{-1-6m} S_m$. Let $c_m = 0, 2, 1$ if m is congruent to 0, 1, 2 modulo 3 respectively. Then for the theorem to be true, we require

$$(3.4) \quad S_m = 3^{6m+5-4\lceil m/3 \rceil + c_m} r$$

for some $r \in R^{(3)}$. We will prove this by induction.

Our base cases are $m = 1, 2, 3$. From Lemma 4,

$$\begin{aligned} g_1(z) &= 10 \cdot 3^9 \phi(z) + 4 \cdot 3^{14} \phi^2(z) + 3^{18} \phi^3(z), \\ g_2(z) &= -4 \cdot 3^{14} \phi(z) - 3^{18} \phi^2(z), \\ g_3(z) &= 3^{18} \phi(z). \end{aligned}$$

We find S_1, S_2 , and S_3 as follows:

$$\begin{aligned} S_1 &= g_1 = 3^9(Q_1(\phi)), \\ S_2 &= g_1 S_1 - 2g_2 = 3^{14}(Q_2(\phi)), \\ S_3 &= g_1 S_2 - S_1 g_2 + 3g_3 = 3^{19}(Q_3(\phi)), \end{aligned}$$

where $Q_1, Q_2, Q_3 \in R^{(3)}$. Note that since g_1 and S_1 each are of the form $3^9 r$ for some $r \in R^{(3)}$, we use the $R^{(p)}$ product lemma to quickly deduce that their product is of the form $3^{18-4} r = 3^{14} r$ for some $r \in R^{(3)}$, from which we easily see $S_2 = 3^{14}(Q_2(\phi))$. Similarly, 3^{23} divides both $g_1 S_2$ and $S_1 g_2$, which means that to keep $g_1 S_2$ and $S_1 g_2$ in $R^{(p)}$, we lose 3^4 and get $S_3 = 3^{19}(Q_3(\phi))$ using the same lemma. Comparing with (3.4), we see that the base case is proved.

Suppose that for some $m \geq 4, i \in \{1, 2, 3\}$ we have

$$S_{m-i} = r_i 3^{6(m-i)+5-4[(m-i)/3]+c_{m-i}}$$

for some $r_i \in R^{(3)}$. We want to show that $S_m = 3^{6m+5-4[m/3]+c_m} r_0$ for some $r_0 \in R^{(3)}$. Recall that for $m \geq 4$,

$$S_m = g_1 S_{m-1} - g_2 S_{m-2} + g_3 S_{m-3}.$$

If each of the terms is of the form $3^{6m+5-4[m/3]+c_m} r_0$ for some $r_0 \in R^{(3)}$, then we are done. We will check each term.

Using the $R^{(p)}$ product lemma, the power of 3 that we get from $g_1 S_{m-1}$ is

$$\begin{aligned} &9 + 6(m-1) + 5 - 4[(m-1)/3] + c_{m-1} - 4 \\ &= 6m + 4 - 4[(m-1)/3] + c_{m-1}, \end{aligned}$$

which we want to be greater than $6m + 5 - 4[m/3] + c_m$. In other words, we want to check whether

$$-1 - 4[(m-1)/3] + c_{m-1} \geq -4[m/3] + c_m,$$

i.e.

$$(3.5) \quad -1 + c_{m-1} \geq -4([\!m/3] - [(m-1)/3]) + c_m.$$

Checking the three cases depending on the value of m modulo 3, we see that (3.5) is true.

The power of 3 that divides $g_2 S_{m-2}$ is

$$\begin{aligned} &14 + 6(m-2) + 5 - 4[(m-2)/3] + c_{m-2} - 4 \\ &= 6m + 3 - 4[(m-2)/3] + c_{m-2}, \end{aligned}$$

which we want to be greater than $6m + 5 - 4[m/3] + c_m$. In other words, we want to check whether

$$-2 + c_{m-2} \geq -4([\!m/3] - [(m-2)/3]) + c_m.$$

As before, we find that this is true.

In the final term, the power of 3 that divides $g_3 S_{m-3}$ is

$$\begin{aligned} & 18 + 6(m-3) + 5 - 4\lceil(m-3)/3\rceil + c_{m-3} - 4 \\ & = 6m + 1 - 4\lceil(m-3)/3\rceil + c_{m-3}, \end{aligned}$$

which we want to be greater than $6m + 5 - 4\lceil m/3\rceil + c_m$. In other words, we want to check whether

$$-4 - 4\lceil(m-3)/3\rceil + c_{m-3} \geq -4\lceil m/3\rceil + c_m,$$

i.e.

$$-4 \geq -4(\lceil m/3\rceil - \lceil(m-3)/3\rceil).$$

As before, this is true, proving the theorem for $\alpha = 1$.

Moving forward to the polynomial step, we have shown that

$$U_3(\phi(z)^m) = \sum_{n=i}^{3m} d(m, n)(\phi(z))^n,$$

where $i = \lceil \frac{m}{3} \rceil$, and $3^2 \mid d(m, i)$ if $m \equiv 1 \pmod{3}$, or $3 \mid d(m, i)$ if $m \equiv 2 \pmod{3}$. Furthermore, $\nu_3(d(m, n)) \geq 4(n-i) + c_m$, where $n \neq i$. To take into account the equivalence class of k modulo 3, we can rewrite this as

$$U_3(\phi(z)^m) = 3^{c_m} \sum_{n=i}^{3m} d'(m, n)(\phi(z))^n,$$

where c_m is as above.

Suppose that

$$\sum_{n=i}^j d_n(\phi(z))^n$$

is a polynomial where $\nu_3(d_n) \geq 4(n-i)$ for $n \geq i$; without loss of generality, we assume that $3 \nmid d_i$. We will show that

$$(3.6) \quad U_3 \left(\sum_{n=i}^j d_n \phi(z)^n \right) = 3^{c_i} \sum_{n=i'}^{3j} d'_n \phi(z)^n,$$

where $d'_n \in \mathbb{Z}$, the quantity 3^{c_i} is 3^2 , 3, or 1 if i is 1, 2, or 0 modulo 3 respectively, and $\nu_3(d'_n) \geq 4(n-i') + c_i$ for $n \geq i'$.

We have already shown that i' is at least $\lceil \frac{i}{3} \rceil$ and that the degree of the resulting polynomial is $3j$, so the only question is whether the 3^{c_i} term appears and whether the d'_n satisfy the appropriate divisibility by increasing powers of 3.

We use the sets $P^{(3)}(\ell, a)$ as defined in the statement of Theorem 7. Note first that $U_3(d_i \phi^i)$ is in the set $P^{(3)}(i', c_i)$. Since U_3 is linear over the sum, we find the set to which each $U_3(d_n \phi^n)$ belongs, and then show that each of those sets is contained in $P^{(3)}(i', c_i)$. It follows that (3.6) holds. We do this by showing the containments

$$\begin{aligned} P^{(3)}(i', c_i) & \supseteq P^{(3)} \left(\left\lceil \frac{i+1}{3} \right\rceil, c_{i+1} + 4(i+1-i) \right) \\ & \supseteq \dots \\ & \supseteq P^{(3)} \left(\left\lceil \frac{j}{3} \right\rceil, c_j + 4(j-i) \right). \end{aligned}$$

In order to show all of these containments, we only need to show consecutive set inclusions, or that for each m , $P^{(3)}(\lceil \frac{m}{3} \rceil, c_m + 4(m-i)) \subseteq P^{(3)}(\lceil \frac{m-1}{3} \rceil, c_{m-1} + 4(m-1-i))$.

Now we move to cases.

Case 1. If $m \equiv 0 \pmod{3}$, then $\lceil \frac{m-1}{3} \rceil = \lceil \frac{m}{3} \rceil$, and we want to show that $P^{(3)}(\lceil \frac{m}{3} \rceil, 4(m-i)) \subseteq P^{(3)}(\lceil \frac{m-1}{3} \rceil, 1 + 4(m-1-i))$. Observe that

$$\begin{aligned} P^{(3)}\left(\left\lceil \frac{m}{3} \right\rceil, 4(m-i)\right) &\subseteq P^{(3)}\left(\left\lceil \frac{m-1}{3} \right\rceil, 4 + 4(m-1-i)\right) \\ &\subseteq P^{(3)}\left(\left\lceil \frac{m-1}{3} \right\rceil, 1 + 4(m-1-i)\right), \end{aligned}$$

as desired.

Case 2. If $m \equiv 1 \pmod{3}$, then $\lceil \frac{m-1}{3} \rceil = \lceil \frac{m}{3} \rceil - 1$, and we want to show that $P^{(3)}(\lceil \frac{m}{3} \rceil, 2 + 4(m-i)) \subseteq P^{(3)}(\lceil \frac{m-1}{3} \rceil, 4(m-1-i))$. Observe that

$$\begin{aligned} P^{(3)}\left(\left\lceil \frac{m}{3} \right\rceil, 2 + 4(m-i)\right) &\subseteq P^{(3)}\left(\left\lceil \frac{m-1}{3} \right\rceil + 1, 4 + 2 + 4(m-1-i)\right) \\ &\subseteq P^{(3)}\left(\left\lceil \frac{m-1}{3} \right\rceil, 4(m-1-i)\right), \end{aligned}$$

as desired.

Case 3. If $m \equiv 2 \pmod{3}$, then $\lceil \frac{m-1}{3} \rceil = \lceil \frac{m}{3} \rceil$, and we want to show that $P^{(3)}(\lceil \frac{m}{3} \rceil, 1 + 4(m-i)) \subseteq P^{(3)}(\lceil \frac{m-1}{3} \rceil, 2 + 4(m-1-i))$. Observe that

$$\begin{aligned} P^{(3)}\left(\left\lceil \frac{m}{3} \right\rceil, 1 + 4(m-i)\right) &\subseteq P^{(3)}\left(\left\lceil \frac{m-1}{3} \right\rceil, 4 + 1 + 4(m-1-i)\right) \\ &\subseteq P^{(3)}\left(\left\lceil \frac{m-1}{3} \right\rceil, 2 + 4(m-1-i)\right), \end{aligned}$$

as desired.

Thus, the result holds for $p = 3$. In similar fashion, the result can also be shown to hold for $p = 5, 7$, completing the proof of Theorem 7. This method of proving the polynomial step could be used to create a similar proof for $p = 2$, simplifying the argument in [7].

Now Theorem 1 follows easily from Theorem 7.

Theorem 1. *Let $p \in \{3, 5, 7\}$. Let $n = p^\alpha n'$ where $p \nmid n'$. Express the base p expansion of m as $a = \sum_{i=1}^{\infty} a_i p^{i-1}$, where $a_i = 0$ for all sufficiently large i . Consider the rightmost α digits $a_\alpha \dots a_2 a_1$. Let i' be the index of the rightmost nonzero digit, or $i' = -1$ if $a_1 = a_2 = \dots = a_\alpha = 0$. Let*

$$\begin{aligned} \gamma_3(m, \alpha) &= \begin{cases} 3 - a_{i'} + 2\#\{i \mid a_i = 0, i > i'\} + \#\{i \mid a_i = 1, i > i'\} & \text{if } i' \geq 0, \\ 0 & \text{otherwise.} \end{cases} \\ \gamma_5(m, \alpha) = \gamma_7(m, \alpha) &= \begin{cases} \chi_{\{1,2\}}(a_{i'}) + \#\{i \mid a_i \in \{0, 1\}, i > i'\} & \text{if } i' \geq 0, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Then

$$a^{(p)}(m, p^\alpha n') \equiv 0 \pmod{p^{\gamma_p(m, \alpha)}}.$$

Proof. The leading coefficient of a polynomial in the set $P^{(p)}(f^\alpha(m), \gamma_p(m, \alpha))$ is divisible by $p^{\gamma_p(m, \alpha)}$. On the other hand, by Theorem 7, $U_{(p)}^\alpha \phi^m$ is an element of that set, so every p^α th coefficient is divisible by $p^{\gamma_p(m, \alpha)}$. \square

4. THE CASE $p = 13$

Since 13 is also a genus zero prime, it is natural to consider whether any congruences hold for $\phi^{(13)}(z)$ or its powers. Computationally, it appears that unless $m \equiv 5 \pmod{13}$, expressing $U_{13}\phi^{(13)}(z)^m$ as a polynomial in $\phi^{(13)}(z)$ gives a coefficient not divisible by 13. Furthermore, in the case of $m \equiv 5 \pmod{13}$, it appears that there are two coefficients of that polynomial that are exactly divisible by 13. If we attempt to use the methods in this paper, this prevents us from chaining properly in the polynomial step. There may be some way around this issue, but it would take a different approach. Computationally, for any prime p less than 1000, it appears that the Fourier coefficient of q^p in $\phi^{(13)}(z)$ is divisible by 13 precisely when $\tau(p)$ is divisible by 13, but no further congruences of the style given in this paper are immediately apparent.

REFERENCES

1. Hans-Fredrik Aas, *Congruences for the coefficients of the modular invariant $j(\tau)$* , Math. Scand. **15** (1964), 64–68.
2. P. Allatt and J. B. Slater, *Congruences on some special modular forms*, J. London Math. Soc. (2) **17** (1978), no. 3, 380–392.
3. Nickolas Andersen and Paul Jenkins, *Divisibility properties of coefficients of level p modular functions for genus zero primes*, Proc. Amer. Math. Soc. **141** (2013), no. 1, 41–53.
4. Tom M. Apostol, *Modular functions and Dirichlet series in number theory*, second ed., Graduate Texts in Mathematics, vol. 41, Springer-Verlag, New York, 1990.
5. Frank Calegari, *Congruences between modular forms*, Arizona Winter School notes, <http://swc.math.arizona.edu/aws/2013/2013CalegariLectureNotes.pdf>, 2013.
6. Michael Griffin, *Divisibility properties of coefficients of weight 0 weakly holomorphic modular forms*, Int. J. Number Theory **7** (2011), no. 4, 933–941.
7. Paul Jenkins, Ryan Keck, and Eric Moss, *Congruences for coefficients of level 2 modular functions with poles at 0*, Archiv der Mathematik **111** (2018), no. 4, 369–378.
8. Paul Jenkins and D. J. Thornton, *Congruences for coefficients of modular functions*, Ramanujan J. **38** (2015), no. 3, 619–628.
9. O. Kolberg, *The coefficients of $j(\tau)$ modulo powers of 3*, Arbok Univ. Bergen Mat.-Natur. Ser. **1962** (1962), no. 16, 7.
10. ———, *Congruences for the coefficients of the modular invariant $j(\tau)$* , Math. Scand. **10** (1962), 173–181.
11. Joseph Lehner, *Divisibility properties of the Fourier coefficients of the modular invariant $j(\tau)$* , Amer. J. Math. **71** (1949), 136–148.
12. ———, *Further congruence properties of the Fourier coefficients of the modular invariant $j(\tau)$* , Amer. J. Math. **71** (1949), 373–386.

DEPARTMENT OF MATHEMATICS, BRIGHAM YOUNG UNIVERSITY, PROVO, UT 84602
E-mail address: `jenkins@math.byu.edu`

E-mail address: `ryank@mathematics.byu.edu`