

Hypoellipticity and the Mori-Zwanzig formulation of stochastic differential equations

Yuanran Zhu^b, Daniele Venturi^{a,*}

^a*Department of Applied Mathematics, University of California Santa Cruz
Santa Cruz (CA) 95064*

^b*Department of Applied Mathematics, University of California Merced
Merced (CA) 95343*

Abstract

We develop a thorough analysis of the Mori-Zwanzig (MZ) formulation for stochastic dynamical systems driven by multiplicative white noise. To this end, we first derive a new type of MZ equation, which we call effective Mori-Zwanzig (EMZ) equation, that governs the temporal dynamics of noise-averaged observables. Such dynamics is generated by a Kolmogorov operator obtained by averaging Itô's representation of the stochastic Liouvillian of the system. Building upon recent work on hypoelliptic operators, we prove that the generator of the EMZ orthogonal dynamics has a spectrum that lies within cusp-shaped region of the complex plane. This allows to rigorously prove that the EMZ memory kernel and fluctuation terms converge exponentially fast in time to a unique equilibrium state. We apply the new theoretical results to the stochastic dynamics of an interacting particle system widely studied in molecular dynamics, and show that such equilibrium state admits an explicit representation.

1. Introduction

The Mori-Zwanzig (MZ) formulation is a technique originally developed in statistical mechanics [31, 57] to formally integrate out phase variables in nonlinear dynamical systems by means of a projection operator. One of the main features of such formulation is that it allows us to systematically derive exact generalized Langevin equations (GLEs) [58, 5, 9, 48] for quantities of interest, e.g., macroscopic observables, based on microscopic equations of motion. Such GLEs can be found in a variety of applications, including particle dynamics [28, 52, 23, 17, 16], fluid dynamics [35, 18, 34], and, more generally, systems described by nonlinear partial differential equations (PDEs) [47, 4, 44, 42, 40, 41, 2, 46, 45, 30, 29]. Computing the solution to the MZ equation is usually a daunting task. One of the main difficulties is the approximation of the memory integral (convolution term) and the fluctuation term, which encode the interaction between the so-called orthogonal dynamics and the dynamics of the quantity of interest. The orthogonal dynamics is essentially a high-dimensional flow that satisfies a hard-to-solve integro-differential equation. The mathematical properties of such flow, and hence the properties of the MZ memory integral and the MZ fluctuation term for a given dynamical system and observable, are not well understood. Recent work in this direction aimed at obtaining existence conditions and operator bounds for the orthogonal dynamics propagator, i.e., the operator semigroup that pushes forward in time the orthogonal dynamics. For example, Kupferman, Givon and Hald [19] proved existence and uniqueness of the orthogonal dynamics for a classical dynamical system with Mori's projection operator. More recently, we were able to prove uniform boundedness of the orthogonal dynamics propagator for Hamiltonian systems using semigroup estimates [53, 15].

The main objective of this paper is to generalize these results to the MZ formulation of stochastic differential equations (SDEs) [16, 22]. In particular, we aim at developing a thorough mathematical analysis of the so-called effective Mori-Zwanzig (EMZ) equation governing the dynamics of noise-averaged observables, i.e., smooth functions of a stochastic flow [26] generated by an SDE. To this end, we build upon recent work

*Corresponding author
Email address: venturi@ucsc.edu (Daniele Venturi)

of Eckmann & Hairer [14, 12, 13], Hérau & Nier [21] and Helffer & Nier [20] on the spectral properties of hypoelliptic operators, and show that the generator of EMZ orthogonal dynamics propagator has a discrete spectrum that lies within cusp-shaped region of the complex plane. This allows us to rigorously prove exponential relaxation to a unique equilibrium state for both the EMZ memory kernel and the EMZ fluctuation term.

This paper is organized as follows. In Section 2, we develop a self-consistent MZ formulation for stochastic differential equations driven by multiplicative Gaussian white noise. This yields a stochastic GLE for smooth functions of the stochastic flow. In Section 3, we average such stochastic GLE over the white noise to obtain an effective Mori Zwanzig (EMZ) equation governing the dynamics of noise-averaged observables. To study the theoretical properties of the EMZ equation, in Section 4, we briefly review Hörmander’s theory of linear hypoelliptic operators, and show how such theory can be used to prove exponential convergence of the EMZ orthogonal dynamics propagator. In Section 5, we apply our theoretical results to the stochastic dynamics of an interacting particle system. The main findings are summarized in Section 6.

2. The Mori-Zwanzig formulation for stochastic differential equations

Let us consider a d -dimensional stochastic differential equation on a smooth manifold \mathcal{M}

$$\frac{d\mathbf{x}(t)}{dt} = \mathbf{F}(\mathbf{x}(t)) + \boldsymbol{\sigma}(\mathbf{x}(t))\boldsymbol{\xi}(t), \quad \mathbf{x}(0) = \mathbf{x}_0 \sim \rho_0(\mathbf{x}), \quad (1)$$

where $\mathbf{F} : \mathcal{M} \mapsto \mathbb{R}^d$ and $\boldsymbol{\sigma} : \mathcal{M} \rightarrow \mathbb{R}^{d \times m}$ are smooth functions, $\boldsymbol{\xi}(t)$ is m -dimensional Gaussian white noise with independent components, and \mathbf{x}_0 is a random initial state characterized in terms of a probability density function¹ $\rho_0(\mathbf{x})$. The solution (1) is a d -dimensional stochastic flow on the manifold \mathcal{M} , which is known as Brownian flow [26]. As is well known, if $\mathbf{F} : \mathcal{M} \mapsto \mathbb{R}^d$ and $\boldsymbol{\sigma} : \mathcal{M} \rightarrow \mathbb{R}^{d \times m}$ are of class \mathcal{C}^{k+1} ($k \geq 0$) with uniformly bounded derivatives, then the solution to (1) is global, and that the corresponding flow is a stochastic flow of diffeomorphisms [3, 51, 50] of class \mathcal{C}^k . This means that such stochastic flow is differentiable k times (with continuous derivative), with respect to the initial condition for all t . Next, we define the vector-valued smooth phase space function

$$\mathbf{u} : \begin{array}{l} \mathcal{M} \rightarrow \mathbb{R}^M \\ \mathbf{x} \mapsto \mathbf{u}(\mathbf{x}) \end{array} \quad (3)$$

Similar to the deterministic case [54, 53], it is possible to define a composition operator $\mathcal{E}(t, s)$, i.e., a stochastic Koopman operator, that pushes forward in time the observable $\mathbf{u}(t) = \mathbf{u}(\mathbf{x}(t))$. Specifically, we have

$$\mathbf{u}(\mathbf{x}(t)) = \mathcal{E}(t, s)\mathbf{u}(\mathbf{x}(s)), \quad (4)$$

where $\mathcal{E}(t, s)$ can be formally written as

$$\mathcal{E}(t, s) = \overrightarrow{\mathcal{T}} e^{\int_s^t \mathcal{L}(\tau) d\tau}. \quad (5)$$

Here $\overrightarrow{\mathcal{T}}$ is the time ordering operator placing later operators to the right, and $\mathcal{L}(\tau)$ is the (random) infinitesimal generator of the composition operator². The generator of the semigroup $\mathcal{E}(t, s)$ can be determined by Dyson’s series expansion [36], i.e.,

$$\mathcal{E}(t, s) = \sum_{n=1}^{\infty} \mathcal{D}_n(t, s), \quad \mathcal{D}_n(t, s) = \overrightarrow{\mathcal{T}} \frac{1}{n!} \int_s^t \int_s^{t_1} \cdots \int_s^{t_{n-1}} dt_1 dt_2 \cdots dt_n \mathcal{L}(t_1) \mathcal{L}(t_2) \cdots \mathcal{L}(t_n). \quad (6)$$

¹If the system (1) evolves from a deterministic initial states then we have

$$\rho_0(\mathbf{x}) = \prod_{i=1}^d \delta(x_i - x_i(0)). \quad (2)$$

² The evolution operator $\mathcal{E}(t, s)$ has a dual construction in the space of probability density functions [11].

Substituting (6) into (4) yields

$$\mathbf{u}(\mathbf{x}(t)) = \sum_{n=1}^{\infty} \mathcal{D}_n(t, s) \mathbf{u}(\mathbf{x}(s)). \quad (7)$$

Next, we match each term of the series expansion (7) with the terms of the stochastic Taylor expansion of $\mathbf{u}(\mathbf{x}(t))$ [25, Chapter 5]. This allows us to construct an explicit expression for the stochastic Liouvillian $\mathcal{L}(\tau)$ that generates $\mathcal{E}(t, s)$. If we interpret the SDE (1) in the Itô sense, then we can use the classical Itô-Taylor expansion to obtain³

$$\mathcal{L}(t) = \sum_{k=1}^d F_k(\mathbf{x}_0) \frac{\partial}{\partial x_{0k}} + \frac{1}{2} \sum_{j=1}^m \sum_{i,k=1}^d \sigma_{ij}(\mathbf{x}_0) \sigma_{kj}(\mathbf{x}_0) \frac{\partial^2}{\partial x_{0i} \partial x_{0k}} + \sum_{j=1}^m \sum_{i=1}^d \sigma_{ij}(\mathbf{x}_0) \xi_j(t) \frac{\partial}{\partial x_{0i}}. \quad (9)$$

With the generator $\mathcal{L}(t)$ available, we can now derive the Mori-Zwanzig equation governing the evolution of the stochastic observable (4). To this end, we introduce a projection operator \mathcal{P} and the complementary projection $\mathcal{Q} = \mathcal{I} - \mathcal{P}$. By following the formal procedure outlined in [55, 54, 11] we obtain the following exact evolution equation

$$\frac{d}{dt} \mathbf{u}(t) = \mathcal{E}(t, 0) \mathcal{P} \mathcal{L}(t) \mathbf{u}(\mathbf{x}_0) + \mathcal{Y}(t, 0) \mathcal{Q} \mathcal{L}(t) \mathbf{u}(0) + \int_0^t \mathcal{E}(s, 0) \mathcal{P} \mathcal{L}(s) \mathcal{Y}(t, s) \mathcal{Q} \mathcal{L}(t) \mathbf{u}(0) ds, \quad (10)$$

where $\mathbf{u}(0) = \mathbf{u}(\mathbf{x}_0)$ and

$$\mathcal{Y}(t, s) = \overrightarrow{\mathcal{T}} e^{\int_s^t \mathcal{Q} \mathcal{L}(\tau) d\tau}. \quad (11)$$

The three terms at the right hand side of (10) are called, respectively, streaming term, fluctuation (or noise) term, and memory term. It is often more convenient (and tractable) to compute the evolution of the observable $\mathbf{u}(t)$ within a closed linear space, e.g., the image of the projection operator \mathcal{P} . To this end, we apply such projection to both sides of equation (10) to obtain⁴

$$\frac{d}{dt} \mathcal{P} \mathbf{u}(t) = \mathcal{P} \mathcal{E}(t, 0) \mathcal{P} \mathcal{L}(t) \mathbf{u}(0) + \int_0^t \mathcal{P} \mathcal{E}(s, 0) \mathcal{P} \mathcal{L}_{\omega}(s) \mathcal{Y}(t, s) \mathcal{Q} \mathcal{L}(t) \mathbf{u}(0) ds. \quad (12)$$

Depending on the choice of the projection operator, the MZ equation (12) can yield evolution equations for different quantities. For example, if we use Chorin's projection [5, 6, 53, 44], then (12) is an evolution equation for the conditional expectation of $\mathbf{u}(t)$. Similarly, if we use Mori's projection [54, 39], then (12) is an evolution equation for the temporal auto-correlation function of $\mathbf{u}(t)$ (see [55] for further details).

3. Effective Mori-Zwanzig (EMZ) equation

The Mori-Zwanzig formulation we discussed in the previous Section provides a general framework to handle dimension reduction in stochastic dynamical systems. However, equations (10) and (12) are not easy to use in practice because the generator $\mathcal{L}(t)$ is time-dependent and it is also a function of the random process $\boldsymbol{\xi}(t)$ (see Eq. (9)). To overcome this difficulty we average the evolution (4) over $\boldsymbol{\xi}(t)$, conditional to the (random) initial state \mathbf{x}_0 . This allows us to define a composition operator $\mathcal{M}(t, 0)$ that pushes forward in time the average of the observable $\mathbf{u}(t)$ over the noise, i.e.,

$$\mathbb{E}_{\boldsymbol{\xi}(t)}[\mathbf{u}(\mathbf{x}(t)) | \mathbf{x}_0] = \mathcal{M}(t, 0) \mathbf{u}(\mathbf{x}_0). \quad (13)$$

³Similarly, by using the Stratonovich-Taylor expansion, we obtain the following infinitesimal generator

$$\mathcal{L}(t) = \sum_{k=1}^d F_k(\mathbf{x}_0) \frac{\partial}{\partial x_{0k}} + \sum_{j=1}^m \sum_{i=1}^d \sigma_{ij}(\mathbf{x}_0) \circ \xi_j(t) \frac{\partial}{\partial x_{0i}}, \quad (8)$$

where $\circ \xi_j(t)$ denotes the stochastic increment in the Stratonovich sense.

⁴The projected fluctuating term $\mathcal{P} \mathcal{Y}(t, 0) \mathcal{Q} \mathcal{L}(t) \mathbf{u}(\mathbf{x}_0)$ is identically zero since $\mathcal{P} \mathcal{Q} = 0$.

$\mathcal{M}(t, 0)$ is a Markovian semigroup [37] generated by the following (backward) Kolmogorov operator [38, 25, 43]

$$\mathcal{K}(\mathbf{x}_0) = \sum_{k=1}^d F_k(\mathbf{x}_0) \frac{\partial}{\partial x_{0k}} + \frac{1}{2} \sum_{j=1}^m \sum_{i,k=1}^d \sigma_{ij}(\mathbf{x}_0) \sigma_{kj}(\mathbf{x}_0) \frac{\partial}{\partial x_{0i} \partial x_{0k}}, \quad (14)$$

i.e.,

$$\mathcal{M}(t, 0) = e^{t\mathcal{K}}. \quad (15)$$

By following the mathematical steps we outlined in Section 2, it is straightforward to obtain the following effective Mori-Zwanzig (EMZ) equation governing the evolution of the noise-averaged observable (13)

$$\frac{\partial}{\partial t} e^{t\mathcal{K}} \mathbf{u}(0) = e^{t\mathcal{K}} \mathcal{P} \mathcal{K} \mathbf{u}(0) + e^{t\mathcal{Q}\mathcal{K}\mathcal{Q}} \mathcal{Q} \mathcal{K} \mathbf{u}(0) + \int_0^t e^{s\mathcal{K}} \mathcal{P} \mathcal{K} e^{(t-s)\mathcal{Q}\mathcal{K}\mathcal{Q}} \mathcal{Q} \mathcal{K} \mathbf{u}(0) ds. \quad (16)$$

Applying the projection operator \mathcal{P} to (16) yields

$$\frac{\partial}{\partial t} \mathcal{P} e^{t\mathcal{K}} \mathbf{u}(0) = \mathcal{P} e^{t\mathcal{K}} \mathcal{P} \mathcal{K} \mathbf{u}(0) + \int_0^t \mathcal{P} e^{s\mathcal{K}} \mathcal{P} \mathcal{K} e^{(t-s)\mathcal{Q}\mathcal{K}\mathcal{Q}} \mathcal{Q} \mathcal{K} \mathbf{u}(0) ds. \quad (17)$$

Note that the projected EMZ equation for stochastic systems has the same structure as the classical projected MZ equation for deterministic (autonomous) systems [55, 53, 54]. However, the Liouville operator \mathcal{L} is now replaced by a Kolmogorov operator \mathcal{K} .

3.1. EMZ with Mori's projection operator

Let us consider the weighted Hilbert space $H = L^2(\mathcal{M}, \rho)$, where ρ is a positive weight function in \mathcal{M} . For instance, ρ can be the probability density function of the random initial state \mathbf{x}_0 . Let

$$\langle h, g \rangle_\rho = \int_{\mathcal{M}} h(\mathbf{x}) g(\mathbf{x}) \rho(\mathbf{x}) d\mathbf{x} \quad h, g \in H \quad (18)$$

be the inner product in H . We introduce the following projection operator

$$\mathcal{P}h = \sum_{i,j=1}^M G_{ij}^{-1} \langle u_i(0), h \rangle_\rho u_j(0), \quad h \in H. \quad (19)$$

where $G_{ij} = \langle u_i(0), u_j(0) \rangle_\rho$ and $u_i(0) = u_i(\mathbf{x})$ ($i = 1, \dots, M$) are M linearly independent functions. With \mathcal{P} defined as in (19), we can write the EMZ equation (16), and its projected version (17), as

$$\frac{d\mathbf{u}(t)}{dt} = \mathbf{\Omega} \mathbf{u}(t) + \int_0^t \mathbf{K}(t-s) \mathbf{u}(s) ds + \mathbf{f}(t), \quad (20)$$

$$\frac{d}{dt} \mathcal{P} \mathbf{u}(t) = \mathbf{\Omega} \mathcal{P} \mathbf{u}(t) + \int_0^t \mathbf{K}(t-s) \mathcal{P} \mathbf{u}(s) ds. \quad (21)$$

where $\mathbf{u}(t) = [u_1(t), \dots, u_M(t)]^T$ and

$$G_{ij} = \langle u_i(0), u_j(0) \rangle_\rho \quad (\text{Gram matrix}), \quad (22a)$$

$$\Omega_{ij} = \sum_{k=1}^M G_{jk}^{-1} \langle u_k(0), \mathcal{K} u_i(0) \rangle_\rho \quad (\text{streaming matrix}), \quad (22b)$$

$$K_{ij}(t) = \sum_{k=1}^M G_{jk}^{-1} \langle u_k(0), \mathcal{K} e^{t\mathcal{Q}\mathcal{K}\mathcal{Q}} \mathcal{Q} \mathcal{K} u_i(0) \rangle_\rho \quad (\text{memory kernel}), \quad (22c)$$

$$f_i(t) = e^{t\mathcal{Q}\mathcal{K}\mathcal{Q}} \mathcal{Q} \mathcal{K} u_i(0) \quad (\text{fluctuation term}). \quad (22d)$$

The Kolmogorov operator \mathcal{K} is not skew-adjoint relative to $\langle \cdot, \cdot \rangle_\rho$, and therefore in general it is not possible to represent the memory kernel $\mathbf{K}(t)$ as a function of the auto-correlation of $\mathbf{f}(t)$ using the second fluctuation-dissipation theorem [55].

3.2. An Example: EMZ for the Ornstein-Uhlenbeck SDE

Let us consider the Ornstein-Uhlenbeck process defined by the solution to the Itô stochastic differential equation

$$\frac{d}{dt}x = \theta(\mu - x) + \sigma\xi(t), \quad (23)$$

where σ , μ , and θ are positive parameters and $\xi(t)$ is Gaussian white noise with correlation function $\langle \xi(t), \xi(s) \rangle = \delta(t - s)$. As is well-known, the Ornstein-Uhlenbeck process is ergodic and it admits a stationary (equilibrium) Gaussian distribution $\rho_{eq} = \mathcal{N}(\mu, \sigma^2/2\theta)$. Let $x(0)$ be a random initial state with probability density function ρ_0 . The formal solution of (23) can be written as

$$x(t) = x(0)e^{-\theta t} + \mu(1 - e^{-\theta t}) + \sigma \int_0^t e^{-\theta(t-s)} \xi(s) ds. \quad (24)$$

We define the phase space function $u(t) = x(t) - \mu$ (centered Ornstein-Uhlenbeck process). By using the formal solution (24) we obtain the following expressions for the conditional mean and conditional autocovariance function

$$\mathbb{E}_{\xi(t)}[x(t) - \mu | x(0)] = (x(0) - \mu)e^{-\theta t}, \quad (25)$$

$$\mathbb{E}_{\xi(t)}[(x(t) - \mu)(x(s) - \mu) | x(0)] = \frac{\sigma^2}{2\theta} \left(e^{-\theta|t-s|} - e^{-\theta(t+s)} \right). \quad (26)$$

Averaging over the random initial state (distributed as ρ_{eq}) yields

$$\mathbb{E}_{x(0)}[\mathbb{E}_{\xi(t)}[x(t) - \mu | x(0)]] = \mathbb{E}_{x(0)}[x(0)e^{-\theta t} - \mu e^{-\theta t}] = 0, \quad (27)$$

$$\mathbb{E}_{x(0)}[\mathbb{E}_{\xi(t)}[(x(t) - \mu)(x(s) - \mu) | x(0)]] = \frac{\sigma^2}{2\theta} e^{-\theta|t-s|}. \quad (28)$$

At this point, we define the projection operators

$$\mathcal{P}_1 h = \frac{\langle h \rangle_{\rho_{eq}}}{\langle u(0) \rangle_{\rho_{eq}}} u(0), \quad \mathcal{P}_2 h = \frac{\langle h, u(0) \rangle_{\rho_{eq}}}{\langle u(0), u(0) \rangle_{\rho_{eq}}} u(0). \quad (29)$$

The Kolmogorov backward operator associated with (23) is

$$\mathcal{K} = -\theta h \frac{\partial}{\partial h} + \frac{\sigma^2}{2} \frac{\partial^2}{\partial h^2}. \quad (30)$$

By using the identity

$$\frac{d}{dt} \mathbb{E}_{x(0)}[\mathbb{E}_{\xi(t)}[u(t)u(0) | x(0)]] = \frac{d}{dt} \mathbb{E}_{x(0)}[\mathbb{E}_{\xi(t)}[u(t) | x(0)]u(0)] = \left\langle \frac{d}{dt} \mathcal{M}(t, 0) u(0), u(0) \right\rangle_{\rho_{eq}}, \quad (31)$$

one can verify that the EMZ equations (20) and (21) derived, respectively, with the projection operators \mathcal{P}_1 and \mathcal{P}_2 defined in (29), can be written as

$$\frac{d}{dt} M(t) = -\theta M(t), \quad \frac{d}{dt} C(t) = -\theta C(t), \quad (32)$$

where $M(t) = \mathbb{E}_{\xi(t)}[x(t) - \mu | x(0)]$ and $C(t) = \mathbb{E}_{x(0)}[\mathbb{E}_{\xi(t)}[(x(t) - \mu)(x(0) - \mu) | x(0)]]$. Clearly, equations (32) are the exact evolution equations governing the (non-stationary) mean and the (stationary) autocorrelation function of process $u(t)$. In fact, their solutions coincide with (25) and (28), respectively.

4. Analysis of the EMZ equation

In this section, we develop an in-depth mathematical analysis of the effective Mori-Zwanzig equation (16) using Hörmander's analysis [21, 20, 49, 33]. In particular, we build upon the result of Hérau and Nier [21], Eckmann and Hairer [14, 12, 13], and Helffer and Nier [20] on linear hypoelliptic operators to prove that the generator of the EMZ orthogonal dynamics, i.e., $\mathcal{Q}\mathcal{K}\mathcal{Q}$, is hypoelliptic. Consequently, the propagator $e^{t\mathcal{Q}\mathcal{K}\mathcal{Q}}$ converges exponentially fast (in time) to statistical equilibrium. This implies that both the EMZ memory kernel (22c) and fluctuation term (22d) converge exponentially fast to an equilibrium state. One of the key results of such analysis is the fact that the spectrum of $\mathcal{Q}\mathcal{K}\mathcal{Q}$ lies within a cusp-shaped region of the complex half-plane. For consistency with the literature on hypoelliptic operators, we will use the negative of \mathcal{K} and $\mathcal{Q}\mathcal{K}\mathcal{Q}$ as semigroup generators⁵ and write the semigroups appearing in EMZ equation (16) as $e^{-t\mathcal{K}}$ and $e^{-t\mathcal{Q}\mathcal{K}\mathcal{Q}}$.

4.1. Analysis of the Kolmogorov operator

In this section we review some of key results related to the Kolmogorov operator (14). Such operator can be written as a Hörmander-type operator of the form

$$\mathcal{K}(\mathbf{x}) = \sum_{i=1}^m \mathcal{X}_i^*(\mathbf{x})\mathcal{X}_i(\mathbf{x}) + \mathcal{X}_0(\mathbf{x}) + f(\mathbf{x}). \quad (33)$$

where $\mathcal{X}_i(\mathbf{x})$ ($0 \leq i \leq m$) denotes a first-order differential partial operator in the variable x_i with space-dependent coefficients, $\mathcal{X}_i^*(\mathbf{x})$ is the formal adjoint of $\mathcal{X}_i(\mathbf{x})$ in $L^2(\mathbb{R}^n)$, and $f(\mathbf{x})$ is a function that has at most polynomial growth at infinity. To derive useful spectral estimates for \mathcal{K} , it is convenient to first provide some definitions

Definition 1. Let N be a real number. Define

$$\text{Pol}_0^N = \left\{ f \in C^\infty(\mathbb{R}^n) \mid \sup_{\mathbf{x} \in \mathbb{R}^n} (1 + \|\mathbf{x}\|)^{-N} |\partial^\alpha f(\mathbf{x})| \leq C_\alpha \right\},$$

where α is a multi-index of arbitrary order. Note that Pol_0^N is the set of infinitely differentiable functions growing at most polynomially as $\|\mathbf{x}\| \rightarrow \infty$. Similarly, we define the space of k -th order differential operators with coefficients growing at most polynomially with \mathbf{x} as

$$\text{Pol}_k^N = \left\{ \mathcal{G} : C^\infty(\mathbb{R}^n) \rightarrow C^\infty(\mathbb{R}^n) \mid \mathcal{G}(\mathbf{x}) = G_0(\mathbf{x}) + \sum_{j=1}^n \sum_{i=1}^k G_j^i(\mathbf{x})\partial_j^i, \quad G_j^i \in \text{Pol}_0^N \right\}.$$

It is straightforward to verify that if $\mathcal{X} \in \text{Pol}_k^N$ and $\mathcal{Y} \in \text{Pol}_l^M$ then the operator commutator $[\mathcal{X}, \mathcal{Y}] = \mathcal{X}\mathcal{Y} - \mathcal{Y}\mathcal{X}$ is in Pol_{k+l-1}^{N+M} .

Definition 2. A family of operators

$$\mathcal{A}_i(\mathbf{x}) = \sum_{j=1}^n A_{ij}(\mathbf{x})\partial_j \quad i = 1, \dots, m \quad (34)$$

is called *non-degenerate* if there are two constants N and C such that

$$\|\mathbf{y}\|^2 \leq C(1 + \|\mathbf{x}\|^2)^N \sum_{i=1}^m \langle \mathcal{A}_i(\mathbf{x}), \mathbf{y} \rangle^2 \quad \forall \mathbf{x}, \mathbf{y} \in \mathbb{R}^n,$$

where $\langle \mathcal{A}_i(\mathbf{x}), \mathbf{y} \rangle = \sum_{j=1}^n A_{ij}(\mathbf{x})y_j$.

⁵If \mathcal{K} and $\mathcal{Q}\mathcal{K}\mathcal{Q}$ are dissipative then $-\mathcal{K}$ and $-\mathcal{Q}\mathcal{K}\mathcal{Q}$ are accretive.

It was recently shown by Eckmann and Hairer in [12, 13] that \mathcal{K} is hypoelliptic if the Lie algebra generated by the operators $\{\mathcal{X}_0, \dots, \mathcal{X}_m\}$ in (33) is non-degenerate. The main result can be summarized as follows

Proposition 1 (Eckmann and Hairer [13]). *If $\{\mathcal{X}_0, \dots, \mathcal{X}_m\}$ and f in (33) satisfy the following conditions:*

1. $\mathcal{X}_j \in \text{Pol}_1^N$ for all $j = 0, \dots, m$, and $f \in \text{Pol}_0^N$;
2. *There exists a finite integer M such that the family of operators consisting of $\{\mathcal{X}_i\}_{i=0}^m$, $\{[\mathcal{X}_i, \mathcal{X}_j]\}_{i,j=1}^m$, $\{[\mathcal{X}_i, [\mathcal{X}_j, \mathcal{X}_k]]\}_{i,j,k=1}^m$ and so on up to the commutators of rank M is non-degenerate;*

Then the operator \mathcal{K} defined in (33) and $\partial_t + \mathcal{K}$ are both hypoelliptic.

Conditions 1 and 2 are called *poly-Hörmander conditions*. The hypoellipticity of the operator $\partial_t + \mathcal{K}$ guarantees smoothness (in time) of the transition probability governed by the Kolmogorov backward equation [12]. Hereafter we review additional important properties of the Kolmogorov operator \mathcal{K} . As a differential operator with C^∞ tempered coefficients (i.e. with all derivatives polynomially bounded), \mathcal{K} and its formal adjoint \mathcal{K}^* are defined in the Schwartz space $\mathcal{S}(\mathbb{R}^n)$, which is dense in $L^p(\mathbb{R}^n)$ ($1 \leq p < \infty$). On the other hand, since \mathcal{K} and \mathcal{K}^* are both closable operators, all estimates we obtain in this Section hold naturally in $\mathcal{S}(\mathbb{R}^n)$, which can be extended to $L^2(\mathbb{R}^n)$. Hence, we do need to distinguish between \mathcal{K} and its closed extension in $L^2(\mathbb{R}^n)$. We now introduce a family of weighted Sobolev spaces

$$\mathcal{S}^{\alpha, \beta} = \{u \in \mathcal{S}'(\mathbb{R}^n) : \Lambda^\alpha \bar{\Lambda}^\beta u \in L^2(\mathbb{R}^n), \quad \alpha, \beta \in \mathbb{R}\}, \quad (35)$$

where $\mathcal{S}'(\mathbb{R}^n)$ the space of tempered distributions in \mathbb{R}^n . The operator $\bar{\Lambda}^\beta$ is product operator defined as $\bar{\Lambda}^\beta := (1 + \|\mathbf{x}\|^2)^{\beta/2}$, while Λ^α is a pseudo-differential operator (see [14, 13, 21]) that reduces to

$$\Lambda^2 = 1 - \Delta \quad (36)$$

for $\alpha = 2$. The weighted Sobolev space (35) is equipped with the scalar product

$$\langle h, g \rangle_{\alpha, \beta} = \langle \Lambda^\alpha \bar{\Lambda}^\beta h, \Lambda^\alpha \bar{\Lambda}^\beta g \rangle_{L^2},$$

which induces the Sobolev norm $\|\cdot\|_{\alpha, \beta}$. Throughout the paper $\|\cdot\|$ denotes the standard L^2 norm. With the above definitions it is possible to prove the following important estimate on spectrum of the Kolmogorov operator \mathcal{K} .

Theorem 1 (Eckmann and Hairer [13]). *Let $\mathcal{K} \in \text{Pol}_2^N$ be an operator of the form (33) satisfying conditions 1. and 2. in Proposition 1. Suppose that the closure of \mathcal{K} is a maximal-accretive operator in $L^2(\mathbb{R}^n)$ and that for every $\epsilon > 0$ there are two constants $\delta > 0$ and $C > 0$ such that*

$$\|u\|_{\delta, \delta} \leq C(\|u\|_{0, \epsilon} + \|\mathcal{K}u\|) \quad (37)$$

for all $u \in \mathcal{S}(\mathbb{R}^n)$. If, in addition, there exist two constants $\delta > 0$ and $D > 0$ such that

$$\|u\|_{0, \epsilon} \leq D(\|u\| + \|\mathcal{K}u\|), \quad (38)$$

then \mathcal{K} has compact resolvent when considered as an operator acting on $L^2(\mathbb{R}^n)$, whose spectrum $\sigma(\mathcal{K})$ is contained in the following cusp-shaped region of the complex plane (see Figure 1):

$$\mathcal{S}_{\mathcal{K}} = \{z \in \mathbb{C} : \text{Re } z \geq 0, \quad |\text{Im } z| < (8C_1)^{M/2}(1 + \text{Re } z)^M\} \quad (39)$$

for some positive constant C_1 and $M \in \mathbb{N}$.

One of the key estimates used by Eckmann and Hairer in the proof of Theorem 1 is

$$\frac{1}{4}|z + 1|^{2/M} \|u\|^2 \leq C_1 ([1 + \text{Re } z]^2 \|u\|^2 + \|(\mathcal{K} - z)u\|^2), \quad \forall \text{Re } z \geq 0. \quad (40)$$

In a series of papers, Hérau, Nier and Helffer [21, 20] proved that the Kolmogorov operator \mathcal{K} corresponding to classical Langevin dynamics generates a semigroup $e^{-t\mathcal{K}}$ that decays exponentially fast to an equilibrium state. Hereafter we show that similar results can be obtained for Kolmogorov operators in the general form (33).




Figure 1: Sketch of the cusp-shaped region of the complex plane enclosing the spectrum of \mathcal{K} and $\mathcal{Q}\mathcal{K}\mathcal{Q}$. The curve γ_{ext} represents the boundary of the cusp. Specifically, we denote by $\partial\mathcal{S}'_{\mathcal{K}}$ the curve defined by the union of γ_{ext} and $\bar{\gamma}$, while $\partial\mathcal{S}'_{\mathcal{K}}$ is defined by the union γ_{ext} (up to intersection with γ_{int}) and γ_{int} .

Theorem 2. *Suppose that \mathcal{K} satisfies all conditions in Theorem 1. If the spectrum of \mathcal{K} in $L^2(\mathbb{R}^n)$ is such that*

$$\sigma(\mathcal{K}) \cap i\mathbb{R} = \{0\}, \quad (41)$$

and the eigenvalue 0 (if any) has finite algebraic multiplicity, then for any $0 < \alpha < \min(\operatorname{Re} \sigma(\mathcal{K})/\{0\})$, there exists a positive constant $C = C(\alpha)$ such that the estimate

$$\|e^{-t\mathcal{K}}u_0 - \pi_0 u_0\| \leq C e^{-\alpha t} \|u_0\| \quad (42)$$

holds for all $u_0 \in L^2(\mathbb{R}^n)$ and for all $t > 0$, where π_0 is the spectral projection onto the kernel of \mathcal{K} .

Proof. The Kolmogorov operator \mathcal{K} is closed, maximal-accretive and densely defined in $L^2(\mathbb{R}^n)$. Hence, by the Lumer-Phillips theorem, the semigroup $e^{-t\mathcal{K}}$ is a contraction in $L^2(\mathbb{R}^n)$. It was shown in [13, 21] that the core of \mathcal{K} is a Schwartz space, and that the hypoelliptic estimate (40) holds for any $u \in L^2(\mathbb{R}^n)$. According to Theorem 1, \mathcal{K} only has a discrete spectrum, i.e., $\sigma(\mathcal{K}) = \sigma_{dis}(\mathcal{K})$. Condition (41) requires that $\lambda = 0$ is the only eigenvalue on the imaginary axis $i\mathbb{R}$, and that such eigenvalue has finite algebraic multiplicity. This condition, together with Theorem 6.1 in [20], allows us to obtain a weakly convergent⁶ Dunford integral representation of the semigroup $e^{-t\mathcal{K}}$ given by

$$e^{-t\mathcal{K}}u_0 - \pi_0 u_0 = \frac{1}{2\pi i} \int_{\partial\mathcal{S}'_{\mathcal{K}}} e^{-tz} (z - \mathcal{K})^{-1} u_0 dz, \quad (44)$$

where $\partial\mathcal{S}'_{\mathcal{K}} = \gamma_{int} \cup \gamma_{ext}$ is the union of the two curves shown in Figure 1, and $(z - \mathcal{K})^{-1}$ is the resolvent of \mathcal{K} . Equation (44) allows us to formulate the semigroup estimation problem as an estimation problem involving an integral in the complex plane. In particular, to derive the upper bound (42), we just need an

⁶Weak convergence is in the sense

$$\langle e^{-t\mathcal{K}}u_0 - \pi_0 u_0, \phi \rangle = \frac{1}{2\pi i} \int_{\partial\mathcal{S}'_{\mathcal{K}}} \langle e^{-tz} (z - \mathcal{K})^{-1} u_0, \phi \rangle dz \quad (43)$$

for $u_0 \in L^2(\mathbb{R}^n)$ and $\phi \in D(\mathcal{K}^*)$

upper bound for the norm of resolvent $(z - \mathcal{K})^{-1}$. To derive such bound, we notice that for all $z \notin \mathcal{S}_{\mathcal{K}}$, where $\mathcal{S}_{\mathcal{K}}$ is the cusp (39), and $\operatorname{Re} z \geq 0$, we have $|z + 1|^{2/M} \geq (8C_1)(1 + \operatorname{Re} z)^2$. A substitution of this inequality into (40) yields, for all $u \in L^2(\mathbb{R}^n)$

$$\frac{1}{8}|z + 1|^{2/M}\|u\|^2 \leq C_1\|(\mathcal{K} - z)u\|^2, \quad \forall \operatorname{Re} z \geq 0, z \notin \mathcal{S}_{\mathcal{K}}.$$

Hence, $\|(\mathcal{K} - z)^{-1}\| \leq \sqrt{8C_1}|z + 1|^{-1/M}$. Next, we rewrite the Dunford integral (44) as

$$\frac{1}{2\pi i} \int_{\partial \mathcal{S}'_{\mathcal{K}}} e^{-tz}(z - \mathcal{K})^{-1}u_0 dz = \frac{1}{2\pi i} \int_{\gamma_{int}} e^{-tz}(z - \mathcal{K})^{-1}u_0 dz + \frac{1}{2\pi i} \int_{\gamma_{ext}} e^{-tz}(z - \mathcal{K})^{-1}u_0 dz. \quad (45)$$

Since $(\mathcal{K} - z)^{-1}$ is a compact linear operator, we have that for any $0 < \alpha < \min(\operatorname{Re} \sigma(\mathcal{K})/\{0\})$ there exists a constant $C_\alpha > 0$ such that $\|(\mathcal{K} - \alpha)u\| \geq C_\alpha\|u\|$. On the other hand, \mathcal{K} is also a real operator, which implies that for all complex numbers $z = (x + iy) \notin \sigma(\mathcal{K})$, we have

$$\|(\mathcal{K} - (x + iy))u\|^2 = \|(\mathcal{K} - x)u\|^2 + y^2\|u\|^2 \geq (C_x^2 + y^2)\|u\|^2,$$

i.e.,

$$\|(\mathcal{K} - (x + iy))^{-1}u\| \leq \frac{1}{\sqrt{C_x^2 + y^2}}\|u\|. \quad (46)$$

This suggests that the resolvent $(\mathcal{K} - z)^{-1}$ is uniformly bounded by $1/C_x$ along the line γ_{int} , i.e.,

$$\left\| \frac{1}{2\pi i} \int_{\gamma_{int}} e^{-tz}(z - \mathcal{K})^{-1}u_0 dz \right\| \leq C e^{-\alpha t}\|u_0\|. \quad (47)$$

The boundary γ_{ext} is defined by all complex numbers $z = x + iy$ such that $|\operatorname{Im} z| = (8C_1)^{M/2}(1 + \operatorname{Re} z)^M$. Also, if $z \notin \mathcal{S}_{\mathcal{K}}$ then we have shown that the norm of the resolvent is bounded by $\|(\mathcal{K} - z)^{-1}\| \leq \sqrt{8C_1}|z + 1|^{-1/M}$. These two inequalities allow us to write, for $M = 1$ and all $M \geq 2$,

$$\begin{aligned} \left\| \frac{1}{2\pi i} \int_{\gamma_{ext}} e^{-tz}(z - \mathcal{K})^{-1}u_0 dz \right\| &\leq C\|u_0\| \int_{(8C_1)^{M/2}(1+x)^M}^{\infty} \exp\{-t[(8C_1)^{-\frac{1}{2}}y^{\frac{1}{M}} - 1]\}y^{-\frac{1}{M}} dy \\ &= C\|u_0\| \int_x^{\infty} e^{-th}(h + 1)^{M-2} dh \\ &\leq C e^{-xt}\|u_0\| \left[\frac{1}{t} + \frac{1}{t^2} + \dots + \frac{1}{t^M} \right] \quad t > 0. \end{aligned} \quad (48)$$

At this point we recall that $e^{-t\mathcal{K}}$ is a dissipative semigroup and that π_0 is a projection operator into the kernel of \mathcal{K} . This allows us to write $\|e^{-t\mathcal{K}}u_0 - \pi_0 u_0\| = \|e^{-t\mathcal{K}}(u_0 - \pi_0 u_0)\| \leq \|u_0 - \pi_0 u_0\|$. By combining this inequality with (44), (45), (47) and (48) we see that there exists a constant $C = C(\alpha)$ such that

$$\|e^{-t\mathcal{K}}u_0 - \pi_0 u_0\| \leq C e^{-\alpha t}\|u_0\|. \quad (49)$$

This completes the proof. \square

Another direct consequence of the estimate (40) is the unconditional convergence of the formal power series expansion (Taylor series) of the semigroup $e^{-t\mathcal{K}}$. This is an important result that yields convergence of operator series expansions – such as the MZ-Dyson expansion [54] – of the EMZ equation for stochastic dynamical systems.

Corollary 2.1. *Assuming that \mathcal{K} satisfies all conditions listed in Theorem 2, then for any $u_0 \in L^2(\mathbb{R}^n)$ the power series expansion*

$$e^{-(t+s)\mathcal{K}}u_0 = \lim_{m \rightarrow \infty} \sum_{n=0}^m \frac{(-s)^n}{n!} e^{-t\mathcal{K}}\mathcal{K}^n u_0 \quad (50)$$

converges in operator norm for any $t > 0$ and $s \geq t$.

Proof. By using the resolvent identity $(z - \mathcal{K})^{-1}\mathcal{K} = z(z - \mathcal{K})^{-1} - \mathcal{I}$, Cauchy's integral theorem, and the Dunford integral representation (44), we obtain

$$\begin{aligned} e^{-t\mathcal{K}}\mathcal{K}u_0 - \pi_0\mathcal{K}u_0 &= \frac{1}{2\pi i} \int_{\partial S'_\mathcal{K}} e^{-tz}(z - \mathcal{K})^{-1}\mathcal{K}u_0 dz, \\ &= \frac{1}{2\pi i} \int_{\partial S'_\mathcal{K}} ze^{-tz}(z - \mathcal{K})^{-1}u_0 dz, \end{aligned} \quad (51)$$

which holds for any $t > 0$. As before, we split the integral along $\partial S'_\mathcal{K}$ into the sum of two integrals (see Eq. (45))

$$e^{-t\mathcal{K}}\mathcal{K}u_0 - \pi_0\mathcal{K}u_0 = \frac{1}{2\pi i} \left(\int_{\gamma_{int}} ze^{-tz}(z - \mathcal{K})^{-1}u_0 dz + \int_{\gamma_{ext}} ze^{-tz}(z - \mathcal{K})^{-1}u_0 dz \right). \quad (52)$$

If $z = x + iy$ is in γ_{int} then we have $|z| \leq (x^2 + (8C_1)^M(1+x)^{2M})^{1/2}$. This inequality, combined with the uniform boundedness of the resolvent (46), allows us to write

$$\left\| \frac{1}{2\pi i} \int_{\gamma_{int}} e^{-tz}z(z - \mathcal{K})^{-1}u_0 dz \right\| \leq Ce^{-xt}\|u_0\|. \quad (53)$$

If $z = x + iy$ is in γ_{ext} then $|z| = \sqrt{[(8C_1)^{-1/2}|y|^{1/M} - 1]^2 + y^2}$. To derive an upper bound for the second integral in (52), we only need to consider the case where y is large enough so that $|y|^{2/M} < |y|^2$, i.e., $|z| \leq C|y|$. A substitution of this estimate into the second integral in (52) yields

$$\begin{aligned} \left\| \frac{1}{2\pi i} \int_{\gamma_{ext}} z^n e^{-tz}(z - \mathcal{K})^{-1}u_0 dz \right\| &\leq C\|u_0\| \int_{(8C_1)^{M/2}(1+x)^M}^{\infty} \exp\{-t[(8C_1)^{-1/2}y^{1/M} - 1]\}y^{1-1/M} dy \\ &\leq C\|u_0\| \int_x^{\infty} e^{-th}(h+1)^{2M-2} dh \\ &\leq \underbrace{Ce^{-\alpha t}\|u_0\| \left[\frac{1}{t} + \frac{1}{t^2} + \dots + \frac{1}{t^{2M-1}} \right]}_{B(t)} \quad t > 0. \end{aligned} \quad (54)$$

Combining (53) and (54) we conclude that the Dunford integral (51) is bounded by $B(t)$. By the triangle inequality, we have that for any fixed $t > 0$ and any $n \in \mathbb{N}$

$$\left\| \left\| e^{-t\mathcal{K}/n}\mathcal{K}u_0 \right\| - \|\pi_0\mathcal{K}u_0\| \right\| \leq \left\| e^{-t\mathcal{K}/n}\mathcal{K}u_0 - \pi_0\mathcal{K}u_0 \right\| \leq B\left(\frac{t}{n}\right). \quad (55)$$

Using the operator identity $e^{-t\mathcal{K}}\mathcal{K}^n = (e^{-t\mathcal{K}/n}\mathcal{K})^n$ we obtain

$$\|e^{-t\mathcal{K}}\mathcal{K}^n u_0\| \leq \left\| e^{-t\mathcal{K}/n}\mathcal{K}u_0 \right\|^n \leq \left(B\left(\frac{t}{n}\right) + \|\pi_0\mathcal{K}u_0\| \right)^n. \quad (56)$$

As easily seen from the the definition of $B(t)$ given in (54), for each fixed $t > 0$ we have

$$\lim_{n \rightarrow \infty} \frac{\|e^{-t\mathcal{K}}\mathcal{K}^n u_0\|}{n!} \leq \lim_{n \rightarrow \infty} \frac{1}{n!} \left(B\left(\frac{t}{n}\right) + \|\pi_0\mathcal{K}u_0\| \right)^n = 0. \quad (57)$$

By using the Lagrangian representation of a truncated power series residual [15, p. 104], we see that (57) implies that (50) converges in operator norm for any $t > 0$ and any $s \geq t$. \square

4.2. Analysis of the projected Kolmogorov operator

In this section we analyze the semigroup $e^{-t\mathcal{Q}\mathcal{K}\mathcal{Q}}$ generated by the operator $\mathcal{Q}\mathcal{K}\mathcal{Q}$, where \mathcal{K} is the Kolmogorov operator (14), \mathcal{P} is a finite-rank projection, and $\mathcal{Q} = \mathcal{I} - \mathcal{P}$ is the complementary projection. Such semigroup appears in the EMZ memory and fluctuation terms (see Eqs. (20), (22c) and (22d)). In principle, the projection operator \mathcal{P} and therefore the complementary projection \mathcal{Q} can be chosen arbitrarily [53, 6]. Here we restrict our analysis to finite-rank, self-adjoint projections in L^2 . Mori's projection (19) is one of such projections.

Theorem 3. *Let $\mathcal{P} : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ be a finite-rank self-adjoint projection operator. If \mathcal{K} satisfies all conditions listed in Theorem 1, then the operator $\mathcal{Q}\mathcal{K}\mathcal{Q}$ is also maximal accretive and has a compact resolvent. Moreover, the spectrum of $\mathcal{Q}\mathcal{K}\mathcal{Q}$ lies within the cusp*

$$\mathcal{S}_{\mathcal{Q}\mathcal{K}\mathcal{Q}} = \{z \in \mathbb{C} \mid \operatorname{Re} z \geq 0, |\operatorname{Im} z| < (8C_{\mathcal{Q}})^{M_{\mathcal{Q}}/2}(1 + \operatorname{Re} z)^{M_{\mathcal{Q}}}\} \quad (58)$$

for some the positive constants $C_{\mathcal{Q}}$ and integer $M_{\mathcal{Q}}$.

Proof. We first show that if \mathcal{K} is closely defined and maximal accretive, and that $\mathcal{Q}\mathcal{K}\mathcal{Q}$ has the same properties. According to Lumer-Phillips theorem [15], the adjoint of a maximal-accretive operator is accretive, and therefore

$$\begin{aligned} \operatorname{Re}\langle \mathcal{K}f, f \rangle &\geq 0, & \forall f \in D(\mathcal{K}), \\ \operatorname{Re}\langle \mathcal{K}^*f, f \rangle &\geq 0, & \forall f \in D(\mathcal{K}^*). \end{aligned}$$

On the other hand, if \mathcal{P} is a self-adjoint operator in $L^2(\mathbb{R}^n)$ then $\mathcal{Q} = \mathcal{I} - \mathcal{P}$ is also a self-adjoint. This implies that

$$\begin{aligned} \operatorname{Re}\langle \mathcal{Q}\mathcal{K}\mathcal{Q}f, f \rangle &= \operatorname{Re}\langle \mathcal{K}\mathcal{Q}f, \mathcal{Q}f \rangle \geq 0, & \forall f \in D(\mathcal{K}), \\ \operatorname{Re}\langle (\mathcal{Q}\mathcal{K}\mathcal{Q})^*f, f \rangle &= \operatorname{Re}\langle \mathcal{K}^*\mathcal{Q}f, \mathcal{Q}f \rangle \geq 0, & \forall f \in D(\mathcal{K}^*), \end{aligned}$$

i.e., $\mathcal{Q}\mathcal{K}\mathcal{Q}$ and its adjoint $\mathcal{Q}\mathcal{K}^*\mathcal{Q}$ are both maximal-accretive. $\mathcal{Q}\mathcal{K}\mathcal{Q}$ is also a closable operator defined in $D(\mathcal{K})$. This can be seen by decomposing it as $\mathcal{Q}\mathcal{K}\mathcal{Q} = \mathcal{K} - \mathcal{K}\mathcal{P} - \mathcal{P}\mathcal{K}\mathcal{Q}$. In fact, if \mathcal{K} is a closed operator then $\mathcal{Q}\mathcal{K}\mathcal{Q}$ is closed since $\mathcal{K}\mathcal{P}$ and $\mathcal{P}\mathcal{K}\mathcal{Q}$ are bounded [24], as we shall see hereafter. By using the Lumer-Phillips theorem again, we conclude that $\mathcal{Q}\mathcal{K}\mathcal{Q}$ is also maximal accretive, and its closure generates a contraction semigroup $e^{-t\mathcal{Q}\mathcal{K}\mathcal{Q}}$ in $L^2(\mathbb{R}^n)$. Next we show that if \mathcal{K} satisfies the hypoelliptic estimate $\|u\|_{\delta,\delta} \leq C(\|u\| + \|\mathcal{K}u\|)$, then so does $\mathcal{Q}\mathcal{K}\mathcal{Q}$, i.e.

$$\|u\|_{\delta,\delta} \leq C(\|u\| + \|\mathcal{Q}\mathcal{K}\mathcal{Q}u\|). \quad (59)$$

By using triangle inequality

$$\|u\|_{\delta,\delta} \leq C(\|u\| + \|\mathcal{K}u\|) \leq C(\|u\| + \|\mathcal{K}\mathcal{P}u\| + \|\mathcal{Q}\mathcal{K}\mathcal{Q}u\| + \|\mathcal{P}\mathcal{K}\mathcal{Q}u\|).$$

To prove (59), it is sufficient to show that $\mathcal{K}\mathcal{P}$ and $\mathcal{P}\mathcal{K}\mathcal{Q}$ are bounded operators in $L^2(\mathbb{R}^n)$. To this end, we recall that any finite-rank projection admits the canonical representation

$$\mathcal{P} = \sum_{i=1}^m \lambda_i \langle \cdot, \phi_i \rangle \varphi_i, \quad (60)$$

where $\{\phi_i\}_{i=1}^m$ and $\{\varphi_i\}_{i=1}^m$ are elements $L^2(\mathbb{R}^n)$. This implies that

$$\|\mathcal{K}\mathcal{P}u\| = \left\| \sum_{i=1}^m \lambda_i \langle u, \phi_i \rangle \mathcal{K}\varphi_i \right\| \leq \sum_{i=1}^m |\lambda_i| \|\mathcal{K}\varphi_i\| \|\phi_i\| \|u\| \leq C\|u\|, \quad (61)$$

$$\|\mathcal{P}\mathcal{K}\mathcal{Q}u\| = \left\| \sum_{i=1}^m \lambda_i \langle \mathcal{K}\mathcal{Q}u, \phi_i \rangle \varphi_i \right\| = \left\| \sum_{i=1}^m \lambda_i \langle u, \mathcal{Q}\mathcal{K}^*\phi_i \rangle \varphi_i \right\| \leq \sum_{i=1}^n |\lambda_i| \|\mathcal{Q}\mathcal{K}^*\phi_i\| \|\varphi_i\| \|u\| \leq C\|u\|. \quad (62)$$

This proves that $\mathcal{K}\mathcal{P}$ and $\mathcal{P}\mathcal{K}\mathcal{Q}$ are bounded operators. At this point we notice that if $\mathcal{Q}\mathcal{K}\mathcal{Q}$ is accretive, then $(\mathcal{Q}\mathcal{K}\mathcal{Q} + \mathcal{I})$ invertible. Moreover,

$$\|u\|_{\delta,\delta} \leq C(\|u\| + \|\mathcal{Q}\mathcal{K}\mathcal{Q}u\|) \leq \sqrt{2}C\|(\mathcal{Q}\mathcal{K}\mathcal{Q} + \mathcal{I})u\| \Rightarrow \|(\mathcal{Q}\mathcal{K}\mathcal{Q} + \mathcal{I})^{-1}u\|_{\delta,\delta} \leq \sqrt{2}C\|u\|,$$

i.e., $(\mathcal{Q}\mathcal{K}\mathcal{Q} + \mathcal{I})^{-1}$ is a bounded operator from L^2 into the Sobolev space $\mathcal{S}^{\delta,\delta}$ (see Eq. (35)). Recall that $\mathcal{S}^{\delta,\delta}$ is compactly embedded into L^2 (Lemma 3.2 [13]). Hence, $(\mathcal{Q}\mathcal{K}\mathcal{Q} + \mathcal{I})^{-1}$ is compact from L^2 into L^2 and therefore $\mathcal{Q}\mathcal{K}\mathcal{Q}$ has a compact resolvent [24]. To prove that the discrete spectrum of $\mathcal{Q}\mathcal{K}\mathcal{Q}$ lies within the cusp $\mathcal{S}_{\mathcal{Q}\mathcal{K}\mathcal{Q}}$ defined in (58), we follow the procedure outlined in [13]. To this end, let $\mathcal{K} \in \text{Pol}_2^N$. Then, for $\delta = \max\{2, N\}$ we have the bound

$$\|(\mathcal{K} + \mathcal{I})u\| \leq C\|u\|_{\delta,\delta}, \quad \forall u \in \mathcal{S}_n,$$

and

$$\|(\mathcal{Q}\mathcal{K}\mathcal{Q} + \mathcal{I})u\| \leq \|\mathcal{Q}\|(\|\mathcal{K}u\| + \|\mathcal{K}\mathcal{P}u\|) + \|u\| \leq C(\|\mathcal{K}u\| + \|u\|) \leq \sqrt{2}C\|(\mathcal{K} + \mathcal{I})u\| \leq C\|u\|_{\delta,\delta}.$$

Recall that $\mathcal{Q}\mathcal{K}\mathcal{Q} : D(\mathcal{Q}\mathcal{K}\mathcal{Q}) \rightarrow L^2(\mathbb{R}^{2d})$ is a maximal accretive operator. Therefore, by Lemma 4.5 in [13], for all $\delta > 0$ we can find an integer $M_{\mathcal{Q}} > 0$ and a constant C such that

$$\langle u, [(\mathcal{Q}\mathcal{K}\mathcal{Q} + \mathcal{I})^*(\mathcal{Q}\mathcal{K}\mathcal{Q} + \mathcal{I})]^{1/M_{\mathcal{Q}}} \rangle \leq C\|u\|_{\delta,\delta}^2. \quad (63)$$

By using the hypoelliptic estimate (63), (59), Proposition B.1 in [20] and the triangle inequality for $z = \text{Re } z + i \text{Im } z$, we obtain

$$\begin{aligned} \frac{1}{2}|z + 1|^{2/M_{\mathcal{Q}}}\|u\|^2 &\leq C\|u\|_{\delta,\delta}^2 + \|(\mathcal{Q}\mathcal{K}\mathcal{Q} - z)\|^2 \\ &\leq C_{\mathcal{Q}}([1 + \text{Re } z]^2\|u\|^2 + \|(\mathcal{Q}\mathcal{K}\mathcal{Q} - z)u\|^2). \end{aligned}$$

This result, together with the compactness of the resolvent of $\mathcal{Q}\mathcal{K}\mathcal{Q}$, implies that if $z \in \sigma(\mathcal{Q}\mathcal{K}\mathcal{Q})$ (discrete spectrum of $\mathcal{Q}\mathcal{K}\mathcal{Q}$) then

$$\frac{1}{4}|\text{Im } z|^{2/M_{\mathcal{Q}}}\|u\|^2 \leq \frac{1}{4}|z + 1|^{2/M_{\mathcal{Q}}}\|u\|^2 \leq C_{\mathcal{Q}}(1 + \text{Re } z)^2\|u\|^2.$$

Therefore the spectrum of $\mathcal{Q}\mathcal{K}\mathcal{Q}$ is contained in the cusp-shaped region $\mathcal{S}_{\mathcal{Q}\mathcal{K}\mathcal{Q}}$ defined in equation (58). If $z \notin \mathcal{S}_{\mathcal{Q}\mathcal{K}\mathcal{Q}}$, then we have resolvent estimate

$$\|(\mathcal{Q}\mathcal{K}\mathcal{Q} - z)^{-1}\| \leq \sqrt{8C_{\mathcal{Q}}}|z + 1|^{-1/M_{\mathcal{Q}}}. \quad (64)$$

□

Remark. The main assumption at the basis of Theorem 3 is that \mathcal{P} is a finite-rank self-adjoint projection operator. Mori's projection (19) is one of such projections. If \mathcal{P} is of finite-rank then both $\mathcal{K}\mathcal{P}$ and $\mathcal{P}\mathcal{K}\mathcal{Q}$ are bounded operators, which yields the hypoelliptic estimate (37). On the other hand, if \mathcal{P} is an infinite-rank projection, e.g., Chorin's projection [53, 7, 8, 58], then $\mathcal{K}\mathcal{P}$ and $\mathcal{P}\mathcal{K}\mathcal{Q}$ may not be bounded. We do not know if Theorem 3 holds for infinite-rank projections.

With the resolvent estimate (64) available, we can now prove the analog of Theorem 2 and Corollary 2.1, where \mathcal{K} replaced by $\mathcal{Q}\mathcal{K}\mathcal{Q}$. These results establish exponential relaxation to equilibrium of $e^{-t\mathcal{Q}\mathcal{K}\mathcal{Q}}$ and convergence of its formal power series expansion (Taylor series).

Theorem 4. *Assume that \mathcal{K} satisfies all conditions listed in Theorem 1. Let $\mathcal{P} : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ is a self-adjoint, finite-rank projection operator. If the spectrum of operator $\mathcal{Q}\mathcal{K}\mathcal{Q}$ in $L^2(\mathbb{R}^n)$ satisfies*

$$\sigma(\mathcal{Q}\mathcal{K}\mathcal{Q}) \cap i\mathbb{R} \subseteq \{0\}, \quad (65)$$

where the eigenvalue 0 (if any) has finite algebraic multiplicity, then for any $0 < \alpha_{\mathcal{Q}} < \min(\text{Re } \sigma(\mathcal{Q}\mathcal{K}\mathcal{Q})/\{0\})$, there exists a positive constant $C = C(\alpha_{\mathcal{Q}})$ such that

$$\|e^{-t\mathcal{Q}\mathcal{K}\mathcal{Q}}u_0 - \pi_0^{\mathcal{Q}}u_0\| \leq Ce^{-\alpha_{\mathcal{Q}}t}\|u_0\|, \quad (66)$$

for all $u_0 \in L^2(\mathbb{R}^n)$ and for all $t > 0$, where $\pi_0^{\mathcal{Q}}$ is the spectral projection onto the kernel of $\mathcal{Q}\mathcal{K}\mathcal{Q}$.

Proof. The proof closely follows the proof of Theorem 2. Therefore it is omitted. \square

Corollary 4.1. *Suppose that \mathcal{K} satisfies all conditions listed in Theorem 2. Then for any $u_0 \in L^2(\mathbb{R}^n)$ the power series expansion*

$$e^{-(t+s)\mathcal{Q}\mathcal{K}\mathcal{Q}}u_0 = \lim_{q \rightarrow \infty} \sum_{n=0}^q \frac{(-s)^n}{n!} e^{-t\mathcal{Q}\mathcal{K}\mathcal{Q}} (\mathcal{Q}\mathcal{K}\mathcal{Q})^n u_0 \quad (67)$$

converges in norm for any $t > 0, s \geq t$.

Proof. The proof closely follows the proof of Corollary 2.1. Therefore it is omitted. \square

Remark. The convergence results in Corollary 2.1 and Corollary 4.1 hold for all $t > 0$. An immediate consequence of such results is that the operator polynomial expansion method we proposed in [54] (see also [53] and the references therein) is convergent when applied to the EMZ equations (20)-(21) corresponding to stochastic systems of the form (1). Similarly, first principle MZ memory calculation method we recently developed in [55] is convergent as well, when applied to the EMZ memory memory (22c) and fluctuation (22d) terms. A very important consequence of the bound (66) is that such terms converge to an equilibrium state exponentially fast (in time). In fact, we have the following

Corollary 4.2. *For any scalar observable $u(t) = u(x(t))$ with initial condition $u(0) = u_0$, the EMZ memory kernel (22c) converges exponentially fast to the equilibrium state $\langle \mathcal{Q}\mathcal{K}^*u_0, \pi_0^{\mathcal{Q}}\mathcal{K}u_0 \rangle$, with rate $\alpha_{\mathcal{Q}}$. In other words, there exists a positive constant C such that*

$$|K(t) - \langle \mathcal{Q}\mathcal{K}^*u_0, \pi_0^{\mathcal{Q}}\mathcal{K}u_0 \rangle| \leq C e^{-\alpha_{\mathcal{Q}}t}. \quad (68)$$

Proof. According to the definition (22c) we have

$$\begin{aligned} |K(t) - \langle \mathcal{Q}\mathcal{K}^*u_0, \pi_0^{\mathcal{Q}}\mathcal{K}u_0 \rangle| &= |\langle u_0, \mathcal{K}e^{t\mathcal{Q}\mathcal{K}\mathcal{Q}}\mathcal{Q}\mathcal{K}u_0 \rangle - \langle \mathcal{Q}\mathcal{K}^*u_0, \pi_0^{\mathcal{Q}}\mathcal{K}u_0 \rangle| \\ &= |\langle \mathcal{Q}\mathcal{K}^*u_0, e^{t\mathcal{Q}\mathcal{K}\mathcal{Q}}\mathcal{K}u_0 \rangle - \langle \mathcal{Q}\mathcal{K}^*u_0, \pi_0^{\mathcal{Q}}\mathcal{K}u_0 \rangle|. \end{aligned} \quad (69)$$

Since $\pi_0^{\mathcal{Q}}$ is a projection we have that $\pi_0^{\mathcal{Q}}\pi_0^{\mathcal{Q}} = \pi_0^{\mathcal{Q}}$, and therefore $\pi_0^{\mathcal{Q}}(\mathcal{K}u_0 - \pi_0^{\mathcal{Q}}\mathcal{K}u_0) = 0$. By using the estimates (66), (69), and the Cauchy-Schwartz inequality we obtain

$$|K(t) - \langle \mathcal{Q}\mathcal{K}^*u_0, \pi_0^{\mathcal{Q}}\mathcal{K}u_0 \rangle| \leq C \|\mathcal{Q}\mathcal{K}^*u_0\| \|\mathcal{K}u_0 - \pi_0^{\mathcal{Q}}\mathcal{K}u_0\| e^{-\alpha_{\mathcal{Q}}t}. \quad (70)$$

\square

Remark. It is straightforward to generalize Corollary 4.2 to matrix-valued memory kernels (22c) and obtain the following exponential convergence result

$$\|\mathbf{K}(t) - \mathbf{G}^{-1}\mathbf{C}^{\mathcal{Q}}\|_M \leq C \|\mathbf{G}^{-1}\mathbf{D}^{\mathcal{Q}}\|_M e^{-\alpha_{\mathcal{Q}}t}, \quad (71)$$

where $\|\cdot\|_M$ denotes any matrix norm and \mathbf{G} is the Gram matrix (22a). Also, the matrix $\mathbf{C}^{\mathcal{Q}}$ has entries $C_{ij}^{\mathcal{Q}} = \langle \mathcal{Q}\mathcal{K}^*u_i(0), \pi_0^{\mathcal{Q}}\mathcal{K}u_j(0) \rangle$, while $D_{ij}^{\mathcal{Q}} = \|\mathcal{Q}\mathcal{K}u_i(0)\| \|\mathcal{K}u_j(0) - \pi_0^{\mathcal{Q}}\mathcal{K}u_j(0)\|$. The proof of (71) follows immediately from the following inequality

$$\begin{aligned} \langle u_i(0), \mathcal{K}e^{t\mathcal{Q}\mathcal{K}\mathcal{Q}}\mathcal{Q}\mathcal{K}u_j(0) \rangle - \langle \mathcal{Q}\mathcal{K}^*u_i(0), \pi_0^{\mathcal{Q}}\mathcal{K}u_j(0) \rangle &= \langle \mathcal{Q}\mathcal{K}^*u_i(0), \mathcal{K}e^{t\mathcal{Q}\mathcal{K}\mathcal{Q}}\mathcal{Q}\mathcal{K}u_j(0) - \pi_0^{\mathcal{Q}}\mathcal{K}u_j(0) \rangle \\ &\leq \|\mathcal{Q}\mathcal{K}u_i(0)\| \|\mathcal{K}u_j(0) - \pi_0^{\mathcal{Q}}\mathcal{K}u_j(0)\|. \end{aligned} \quad (72)$$

In fact, a substitution of (72) into (22c) yields (71).

Remark. Similarly, we can prove that the fluctuation term (22d) reaches the equilibrium state exponentially fast in time. If we choose the initial condition as $\mathbf{u}_0 = \mathcal{Q}\mathcal{K}\mathbf{u}_0$ then for all $j = 1, \dots, m$, we have

$$\|f_j(t) - \pi_0^{\mathcal{Q}}\mathcal{Q}\mathcal{K}u_j(0)\| = \|e^{-t\mathcal{Q}\mathcal{K}\mathcal{Q}}\mathcal{Q}\mathcal{K}u_j(0) - \pi_0^{\mathcal{Q}}\mathcal{Q}\mathcal{K}u_j(0)\| \leq Ce^{-\alpha\varrho t}\|\mathcal{Q}\mathcal{K}u_j(0)\|. \quad (73)$$

Let us now introduce the tensor product space $V = \otimes_{i=1}^m L_i^2(\mathbb{R}^n)$ and the following norm

$$\|\mathbf{u}(t)\|_V := \|(\|u_1(t)\|, \|u_2(t)\|, \dots, \|u_m(t)\|)\|_M, \quad (74)$$

where $\|\cdot\|$ is the standard $L^2(\mathbb{R}^n)$ norm, and $\|\cdot\|_M$ is any matrix norm. Then from (73) it follows that

$$\|\mathbf{f}(t) - \pi_0^{\mathcal{Q}}\mathcal{Q}\mathcal{K}\mathbf{u}_0\|_V \leq Ce^{-\alpha\varrho t} (\|\mathcal{Q}\mathcal{K}u_1(0)\|, \dots, \|\mathcal{Q}\mathcal{K}u_m(0)\|)\|_M. \quad (75)$$

5. Application to molecular dynamics

All results we obtained so far can be applied to arbitrary stochastic dynamical systems of the form (1), provided the MZ projection operator is of finite-rank. In this section, we study in detail a specific system, i.e., an interacting particle system widely used in molecular dynamics simulations, and show that the EMZ memory kernel (22c) and fluctuation term (22d) decay exponentially fast in time to a unique equilibrium state. Such state is defined by the projector $\pi_0^{\mathcal{Q}}$ appearing in Theorem 3 and Corollary 4.2. Hereafter we will determine the exact expression of such projector for a system of interacting identical particles modeled by the following SDE in \mathbb{R}^{2d} (see [27])

$$\begin{cases} \frac{d\mathbf{q}}{dt} = \frac{1}{m}\mathbf{p} \\ \frac{d\mathbf{p}}{dt} = -\nabla V(\mathbf{q}) - \frac{\gamma}{m}\mathbf{p} + \sigma\xi(t) \end{cases}, \quad (76)$$

where m is the mass of each particle, $V(\mathbf{q})$ is the interaction potential and $\xi(t)$ is a d -dimensional Gaussian white noise process modeling physical Brownian motion. The parameters γ and σ represent, respectively, the magnitude of the fluctuations and the dissipation. Such parameters are linked by the fluctuation-dissipation relation $\sigma = (2\gamma/\beta)^{1/2}$, where β is proportional to the inverse of the thermodynamic temperature. The stochastic dynamical system (76) is widely used in statistical mechanics to model mesoscopic dynamics of liquids and gases. Letting the mass m in (76) go to zero, and setting $\gamma = 1$ yields the so-called overdamped Langevin dynamics, i.e., Langevin dynamics where no average acceleration takes place. The (negative) Kolmogorov operator (14) associated with the SDE (76) is given by

$$\mathcal{K} = -\frac{\mathbf{p}}{m} \cdot \nabla_{\mathbf{q}} + \nabla_{\mathbf{q}}V(\mathbf{q}) \cdot \nabla_{\mathbf{p}} + \gamma \left(\frac{\mathbf{p}}{m} \cdot \nabla_{\mathbf{p}} - \frac{1}{\beta}\Delta_{\mathbf{p}} \right), \quad (77)$$

where “ \cdot ” denotes the standard dot product. If the interaction potential $V(\mathbf{q})$ is strictly positive at infinity then the Langevin equation (76) admits an unique invariant Gibbs measure given by

$$\rho_{eq}(\mathbf{p}, \mathbf{q}) = \frac{1}{Z}e^{-\beta H(\mathbf{p}, \mathbf{q})}, \quad (78)$$

where

$$H(\mathbf{p}, \mathbf{q}) = \frac{\|\mathbf{p}\|_2^2}{2m} + V(\mathbf{q}), \quad (79)$$

is the Hamiltonian and Z is the partition function. At this point we introduce the unitary transformation $\mathcal{U} : L^2(\mathbb{R}^{2d}) \rightarrow L^2(\mathbb{R}^{2d}; \rho_{eq})$ defined by

$$(\mathcal{U}g)(\mathbf{p}, \mathbf{q}) = \sqrt{Z}e^{\beta H(\mathbf{p}, \mathbf{q})/2}g(\mathbf{p}, \mathbf{q}), \quad (80)$$

where $L^2(\mathbb{R}^{2d}; \rho_{eq})$ is a weighted Hilbert space endowed with the inner product

$$\langle h, g \rangle_{eq} = \int h(\mathbf{p}, \mathbf{q})g(\mathbf{p}, \mathbf{q})\rho_{eq}(\mathbf{p}, \mathbf{q})d\mathbf{p}d\mathbf{q}. \quad (81)$$

The linear transformation (80) is an isomorphism between $L^2(\mathbb{R}^{2d})$ and $L^2(\mathbb{R}^{2d}; \rho_{eq})$. In fact, for any $\tilde{u} \in L^2(\mathbb{R}^{2d})$, there exists a unique $u \in L^2(\mathbb{R}^{2d}, \rho_{eq})$ such that $\tilde{u} = (e^{-\beta H/2}/\sqrt{Z})u$ and

$$\|\tilde{u}\|_{L^2} = \|u\|_{L^2_{eq}}. \quad (82)$$

By applying (80) to (77) we construct the transformed Kolmogorov operator $\tilde{\mathcal{K}} = \mathcal{U}^{-1}\mathcal{K}\mathcal{U}$, which has the explicit expression

$$\tilde{\mathcal{K}} = -\frac{\mathbf{p}}{m} \cdot \nabla_{\mathbf{q}} + \nabla V(\mathbf{q}) \cdot \nabla_{\mathbf{p}} + \frac{\gamma}{\beta} \left(-\nabla_{\mathbf{p}} + \frac{\beta}{2m} \mathbf{p} \right) \cdot \left(\nabla_{\mathbf{p}} + \frac{\beta}{2m} \mathbf{p} \right). \quad (83)$$

This operator can be written in the canonical form (33) as

$$\tilde{\mathcal{K}} = \sum_{i=1}^d \mathcal{X}_i^* \mathcal{X}_i - \mathcal{X}_0, \quad (84)$$

provided we set

$$\begin{cases} \mathcal{X}_0 = \frac{\mathbf{p}}{m} \cdot \nabla_{\mathbf{q}} - \nabla V(\mathbf{q}) \cdot \nabla_{\mathbf{p}}, \\ \mathcal{X}_i = \sqrt{\frac{\gamma}{\beta}} \left(\partial_{p_i} + \frac{\beta}{2m} p_i \right), \\ \mathcal{X}_i^* = \sqrt{\frac{\gamma}{\beta}} \left(-\partial_{p_i} + \frac{\beta}{2m} p_i \right). \end{cases} \quad (85)$$

Note that \mathcal{X}_0 is skew-adjoint in $L^2(\mathbb{R}^{2d})$. Also, \mathcal{X}_i^* and \mathcal{X}_i can be interpreted as creation and annihilation operator of a quantum harmonic oscillator [56]. The Kolmogorov operator $\tilde{\mathcal{K}}$ and its formal adjoint $\tilde{\mathcal{K}}^*$ are both accretive, closable and with maximally accretive closure in $L^2(\mathbb{R}^{2d})$ (see, e.g., [21, 20, 12]) Similar to the Kolmogorov operator $\tilde{\mathcal{K}} = \mathcal{U}^{-1}\mathcal{K}\mathcal{U}$, we can transform the MZ projection operators \mathcal{P} and \mathcal{Q} into operators in the “flat” Hilbert space $L^2(\mathbb{R}^{2d})$ as $\tilde{\mathcal{P}} = \mathcal{U}^{-1}\mathcal{P}\mathcal{U}$ and $\tilde{\mathcal{Q}} = \mathcal{U}^{-1}\mathcal{Q}\mathcal{U}$. The relationship between $L^2(\mathbb{R}^{2d})$, $L^2(\mathbb{R}^{2d}; \rho_{eq})$ and the operators defined between such spaces can be summarized by the following commutative diagram

$$\begin{array}{ccc} L^2(\mathbb{R}^{2d}) & \xrightarrow{\mathcal{U}} & L^2(\mathbb{R}^{2d}; \rho_{eq}) \\ \tilde{\mathcal{P}}, \tilde{\mathcal{K}}, \tilde{\mathcal{Q}} \downarrow & & \downarrow \mathcal{P}, \mathcal{K}, \mathcal{Q} \\ L^2(\mathbb{R}^{2d}) & \xleftarrow{\mathcal{U}^{-1}} & L^2(\mathbb{R}^{2d}; \rho_{eq}) \end{array}$$

The properties of all operators in $L^2(\mathbb{R}^{2d})$ and $L^2(\mathbb{R}^{2d}; \rho_{eq})$ are essentially the same since \mathcal{U} is an isomorphism. For instance if \mathcal{P} is compact and self-adjoint then $\tilde{\mathcal{P}}$ is also compact and self-adjoint.

Next, we apply the analytical results we obtained in Section 4.1 and Section 4.2 to the particle system described by the SDE (76). To this end, we just need to verify whether $\tilde{\mathcal{K}}$ is a poly-Hörmander operator, i.e., if the operators $\{\mathcal{X}_i\}_{i=0}^d$ appearing in (84)-(85) satisfy the poly-Hörmander conditions in Proposition 1 and the estimate in Theorem 1 (see Section 4.1). This can be achieved by imposing additional conditions on the particle interaction potential $V(\mathbf{q})$ (see [12, Proposition 3.7]). In particular, following Helffer and Nier [20], we assume $V(\mathbf{q})$ satisfies the following weak ellipticity hypothesis

Hypothesis 1. *The particle interaction potential $V(\mathbf{q})$ is of class $C^\infty(\mathbb{R}^d)$, and for all $\mathbf{q} \in \mathbb{R}^d$ it satisfies*

1. $\forall \alpha \in \mathbb{N}^d$ such that $|\alpha| = 1$, $|\partial_{\mathbf{q}}^\alpha V(\mathbf{q})| \leq C_\alpha \sqrt{1 + \|\nabla V(\mathbf{q})\|^2}$ for some positive constant C_α ,
2. There exists $M \in \mathbb{N}$, and $C \geq 1$, such that $C^{-1}(1 + \|\mathbf{q}\|^2)^{1/(2M)} \leq \sqrt{1 + \|\nabla V(\mathbf{q})\|^2} \leq C(1 + \|\mathbf{q}\|^2)^{M/2}$.

Hypothesis 1 holds for any particle interaction potential that grows at most polynomially at infinity, i.e., $V(\mathbf{q}) \simeq \|\mathbf{q}\|^M$ as $\|\mathbf{q}\| \rightarrow \infty$. With this hypothesis, it is possible to prove the following

Proposition 2 (Helffer and Nier [20]). *Consider the Langevin equation (76) with particle interaction potential $V(\mathbf{q})$ satisfying Hypothesis 1. Then the operator $\tilde{\mathcal{K}}$ defined in (83) has a compact resolvent, and a discrete spectrum bounded by the cusp $\mathcal{S}_{\tilde{\mathcal{K}}}$. Moreover, there exists a positive constant C such that the estimate*

$$\|e^{-t\tilde{\mathcal{K}}}u_0 - \tilde{\pi}_0\tilde{u}_0\| \leq Ce^{-\alpha t}\|\tilde{u}_0\| \quad (86)$$

holds for all $\tilde{u}_0 \in L^2(\mathbb{R}^{2d})$ and for all $t > 0$, where $\tilde{\pi}_0$ is the orthogonal projection onto the kernel of $\tilde{\mathcal{K}}$ in $L^2(\mathbb{R}^{2d})$.

By using the isomorphism (80) we can rewrite Proposition 2 in $L^2(\mathbb{R}^{2d}; \rho_{eq})$ as

$$\|e^{-t\mathcal{K}}u_0 - \pi_0u_0\|_{L^2_{eq}} = \|e^{-t\tilde{\mathcal{K}}}\tilde{u}_0 - \tilde{\pi}_0\tilde{u}_0\|_{L^2} \leq Ce^{-\alpha t}\|\tilde{u}_0\|_{L^2} = Ce^{-\alpha t}\|u_0\|_{L^2_{eq}}, \quad (87)$$

where $\pi_0 = \mathcal{U}\tilde{\pi}_0\mathcal{U}^{-1}$ is the orthogonal projection $\pi_0(\cdot) = \mathbb{E}[\cdot]$. The inequality (87) is completely equivalent to the estimate (42). It is also possible to obtain a priori estimate on the convergence rate α by building a connection between the Kolmogorov operator and the Witten Laplacian (see [21, 20] for further details).

Our next task is to derive an estimate for the operator $\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}}$, and for the semigroup $e^{-t\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}}}$ generated by $\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}}$. According to Theorem 3, the spectrum of $\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}}$ is bounded by the cusp $\mathcal{S}_{\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}}}$, provided that \mathcal{P} is a finite-rank projection. On the other hand, Theorem 4 establishes exponential convergence of $e^{-t\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}}}$ to equilibrium if $\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}}$ satisfies condition (65). To determine the exact form of the spectral projection $\tilde{\pi}_0^{\tilde{\mathcal{Q}}}$, i.e., the projection onto the kernel of $\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}}$ (see Theorem 4) we consider a general Mori-type projection $\tilde{\mathcal{P}}$ and its unitarily equivalent version $\tilde{\mathcal{P}} = \mathcal{U}^{-1}\mathcal{P}\mathcal{U}$

$$\mathcal{P}(\cdot) = \sum_{i=1}^m \langle \cdot, g_i(\mathbf{q}) \rangle_{eq} g_i(\mathbf{q}), \quad \tilde{\mathcal{P}}(\cdot) = \sum_{i=1}^m \langle \cdot, g_i(\mathbf{q}) \rangle_{eq/2} g_i(\mathbf{q}) e^{-\beta H/2}, \quad (88)$$

where $\{g_j(\mathbf{q})\}_{j=1}^m$ are zero-mean⁷ orthonormal basis functions. In (88) we used the shorthand notation

$$\langle h \rangle_{eq/2} = \frac{1}{Z} \int g(\mathbf{p}, \mathbf{q}) e^{-\beta H(\mathbf{p}, \mathbf{q})/2} d\mathbf{p} d\mathbf{q}. \quad (89)$$

Lemma 5. *Suppose that the particle interaction potential $V(\mathbf{q})$ in (77) satisfies Hypothesis 1. Then*

$$Ker(\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}}) = Ker(\tilde{\mathcal{K}}) \oplus Id(\tilde{\mathcal{P}}), \quad (90)$$

$$\sigma(\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}}) \cap i\mathbb{R} \subseteq \{0\}, \quad (91)$$

where $\tilde{\mathcal{K}}$ and $\tilde{\mathcal{P}}$ are defined in (83) and (88), respectively, and $Id(\tilde{\mathcal{P}}) = \{u \in L^2(\mathbb{R}^{2d}) : \tilde{\mathcal{P}}u = u\}$ is a finite-dimensional linear space.

Proof. We first prove (90). If $u \in Ker(\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}})$ then $\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}}u = 0$. Therefore we can only have one of the following three cases

1. $\tilde{\mathcal{Q}}u = 0$;
2. $\tilde{\mathcal{Q}}u \neq 0, \tilde{\mathcal{K}}\tilde{\mathcal{Q}}u = 0$;
3. $\tilde{\mathcal{Q}}u \neq 0, \tilde{\mathcal{K}}\tilde{\mathcal{Q}}u \neq 0, \tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}}u = 0$.

Hereafter we discuss each case separately.

Case 1. If $\tilde{\mathcal{Q}}u = 0$ then $\tilde{\mathcal{P}}u = u - \tilde{\mathcal{Q}}u = u$, i.e., $u \in Id(\tilde{\mathcal{P}})$.

⁷Zero-mean here means that $\langle g_j \rangle_{eq} = 0$.

Case 2. If $\tilde{Q}u \neq 0$ and $\tilde{K}\tilde{Q}u = 0$, then $\tilde{Q}u \in Ker(\tilde{K})$. We first prove that $\tilde{Q}u \in Ker(\tilde{K}) \Rightarrow u \in Id(\tilde{P}) \oplus Ker(\tilde{K})$. Since \tilde{P} is an orthogonal, finite-rank projection in $L^2(\mathbb{R}^n)$ (see Eq. (88)), we have that⁸ $Id(\tilde{P}) = R(\tilde{P}) = \text{span}\{g_i(\mathbf{q})e^{-\beta H(\mathbf{p}, \mathbf{q})/2}\}_{i=1}^m$. At this point, we write u as

$$u = \tilde{P}u + \tilde{Q}u = \sum_{i=1}^m c_i g_i(\mathbf{q})e^{-\beta H/2} + c'e^{-\beta H/2} \in Id(\tilde{P}) \oplus Ker(\tilde{K}), \quad c_i, c' \in \mathbb{R}.$$

Let us verify that if $u \in Id(\tilde{P}) \oplus Ker(\tilde{K})$ then $\tilde{K}\tilde{Q}u = 0$, i.e. $\tilde{Q}u \in Ker(\tilde{K})$. Indeed,

$$\tilde{Q}u = u - \tilde{P}u = c'e^{-\beta H/2} - c'\tilde{P}e^{-\beta H/2} = c'e^{-\beta H/2} \in Ker(\tilde{K}) \Rightarrow \tilde{K}\tilde{Q}u = 0.$$

Case 3. Let us assume that $\tilde{Q}u \neq 0$, $\tilde{K}\tilde{Q}u \neq 0$ and $\tilde{Q}\tilde{K}\tilde{Q}u = 0$. If u satisfies these three conditions, then we have

$$\langle \tilde{Q}\tilde{K}\tilde{Q}u, u \rangle = \langle \tilde{K}\tilde{Q}u, \tilde{Q}u \rangle = 0, \quad \tilde{Q}u \neq 0, \quad \tilde{K}\tilde{Q}u \neq 0. \quad (92)$$

Upon definition of $g = \tilde{Q}u$, we can write (92) as $\langle \tilde{K}g, g \rangle = 0, g \neq 0, \tilde{K}g \neq 0$. This suggests that $\text{Re}\langle \tilde{K}g, g \rangle = \sum_{i=1}^d \langle \mathcal{X}_i g, \mathcal{X}_i g \rangle = 0$, which implies that g belongs to the kernel of all annihilation operators \mathcal{X}_i , $1 \leq i \leq d$. Therefore g must be of the form $g = \mu(\mathbf{q})e^{-\beta \|\mathbf{p}\|^2/(4m)}$. On the other hand, since $\tilde{Q}\tilde{K}\tilde{Q}u = \tilde{Q}\tilde{K}g = 0$, we have that $\tilde{P}\tilde{K}g = \tilde{K}g$. A substitution of general form of g into this expression yields

$$\tilde{K}g = \mathcal{X}_0 g = \sum_{i=1}^d p_i \left[\frac{1}{m} \partial_{q_i} \mu(\mathbf{q}) - \frac{\beta}{2m} \mu(\mathbf{q}) \partial_{q_i} V(\mathbf{q}) \right] e^{-\beta \|\mathbf{p}\|^2/(4m)} = \sum_{i=1}^d F_i(\mathbf{q}) p_i e^{-\beta \|\mathbf{p}\|^2/(4m)}. \quad (93)$$

Thus,

$$\tilde{P}\tilde{K}g = \sum_{i=1}^N \langle \tilde{K}g, g_i(\mathbf{q}) \rangle_{eq/2} g_j(\mathbf{q}) e^{-\frac{\beta}{2} V(\mathbf{q})} e^{-\beta \|\mathbf{p}\|^2/(4m)}. \quad (94)$$

The function $\mathbf{F}(\mathbf{q})$ appearing in (93) is a d -dimensional vector field that depends on the position \mathbf{q} . It is clear that $\tilde{P}\tilde{K}g = \tilde{K}g$ if and only if $\tilde{K}g = 0$, which contradicts the assumption $\tilde{K}\tilde{Q}u \neq 0$. Hence we conclude that there is no u that simultaneously satisfies $\tilde{Q}\tilde{K}\tilde{Q}u = 0$, $\tilde{Q}u \neq 0$ and $\tilde{K}\tilde{Q}u \neq 0$. In other words, the kernel of operator $\tilde{K}\tilde{Q}\tilde{K}$ is the finite-dimensional linear space $Ker(\tilde{K}) \oplus Id(\tilde{P})$. Next we prove the spectral condition (91). Such condition states that the only eigenvalue of $\tilde{Q}\tilde{K}\tilde{Q}$ on the imaginary axis $i\mathbb{R}$ is the origin and that such eigenvalue has finite algebraic multiplicity. Equivalently, this means that for all $u \in L^2(\mathbb{R}^{2d})$ such that $\tilde{Q}\tilde{K}\tilde{Q}u = i\lambda u$ ($\lambda \in \mathbb{R}$), we have that $\lambda = 0$. To see this, let us choose $g = \tilde{Q}u$. Then $\tilde{Q}\tilde{K}\tilde{Q}u = i\lambda u$ implies that $\text{Re}\langle \tilde{Q}\tilde{K}\tilde{Q}u, u \rangle = \text{Re}\langle \tilde{K}g, g \rangle = 0$, which again implies $g = \mu(\mathbf{q})e^{-\beta \|\mathbf{p}\|^2/(4m)}$. Since g is a real function and \mathcal{X}_0 a real operator, we have $\text{Im}\langle \tilde{K}g, g \rangle = \text{Im}\langle \mathcal{X}_0 g, g \rangle = 0$. Therefore, if $\tilde{Q}\tilde{K}\tilde{Q}u = i\lambda u$ and $\lambda \in \mathbb{R}$, then we must have $\lambda = 0$. □

Remark. Lemma 5 can be generalized to other projection operators of the form (88). To this end, one needs to check if condition $\tilde{P}\tilde{K}g = \tilde{K}g$ in Case 3 implies $\tilde{K}g = 0$ and vice versa. For most projection operators of the form (88), this is true and therefore $Ker(\tilde{Q}\tilde{K}\tilde{Q}) = Ker(\tilde{K}) \oplus Id(\tilde{P})$. However, there are cases where this is not true. For instance, if $\mathcal{P}, \tilde{\mathcal{P}}$ are defined as

$$\mathcal{P}(\cdot) = \langle (\cdot), p_j \rangle_{eq} p_j \quad \tilde{\mathcal{P}}(\cdot) = \langle (\cdot), p_j \rangle_{eq/2} p_j e^{-\beta \mathcal{H}/2}, \quad (95)$$

then we have

$$\tilde{\mathcal{K}}g = \sum_{i=1}^d [\mathbf{F}(\mathbf{q})]_i p_i e^{-\frac{\beta}{4m} \|\mathbf{p}\|^2}, \quad \tilde{\mathcal{P}}\tilde{\mathcal{K}}g = \langle \tilde{\mathcal{K}}g, p_j \rangle_{eq/2} p_j e^{-\frac{\beta}{2} V(\mathbf{q})} e^{-\frac{\beta}{4m} \|\mathbf{p}\|^2} \quad (96)$$

⁸ $R(\tilde{P})$ denotes the range of the projection \tilde{P} .

where $g = \tilde{Q}u$. The condition $\tilde{\mathcal{P}}\tilde{\mathcal{K}}g = \tilde{\mathcal{K}}g$ in Case 3 is satisfied if the following system of equations

$$\begin{cases} [\mathbf{F}(\mathbf{q})]_i = 0, & i \neq j \\ [\mathbf{F}(\mathbf{q})]_j = \langle [F(q)]_j, p_j \rangle_{e_q/2} e^{-\frac{\beta}{2}V(q)}, & i = j \end{cases} \quad (97)$$

admits a non-zero solution. Hence in this case, $\text{Ker}(\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}}) \neq \text{Ker}(\tilde{\mathcal{K}}) \oplus \text{Id}(\tilde{\mathcal{P}})$, and therefore the EMZ fluctuation term $\mathbf{f}(t)$ and the memory kernel $\mathbf{K}(t)$ converge to different constant matrices.

Lemma 5 allows us to prove the following exponential convergence result for the semigroup $e^{-t\mathcal{Q}\mathcal{K}\mathcal{Q}}$.

Proposition 3. *Suppose that the particle interaction potential $V(\mathbf{q})$ in (77) satisfies Hypothesis 1. Then there exists a positive constant C such that*

$$\|e^{-t\mathcal{Q}\mathcal{K}\mathcal{Q}}u_0 - \pi_0^{\mathcal{Q}}u_0\|_{L_{e_q}^2} \leq Ce^{-\alpha_{\mathcal{Q}}t}\|u_0\|_{L_{e_q}^2} \quad (98)$$

for all $u_0 \in L^2(\mathbb{R}^{2d}, \rho_{e_q})$ and for all $t > 0$. In (98), \mathcal{P} is the finite-rank projection (88), while

$$\pi_0^{\mathcal{Q}}(\cdot) = \pi_0(\cdot) + \mathcal{P}(\cdot) \quad (99)$$

is the orthogonal projection onto the linear space $\text{Ker}(\mathcal{Q}\mathcal{K}\mathcal{Q}) = \text{Ker}(\mathcal{K}) \oplus \text{Id}(\mathcal{P})$.

Proof. We first rewrite (98) as an $L^2(\mathbb{R}^{2d})$ estimation problem as

$$\|e^{-t\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}}}\tilde{u}_0 - \tilde{\pi}_0^{\tilde{\mathcal{Q}}}\tilde{u}_0\|_{L^2} \leq Ce^{-\alpha_{\mathcal{Q}}t}\|\tilde{u}_0\|_{L^2} \quad (100)$$

where $\tilde{\pi}_0^{\tilde{\mathcal{Q}}} = \mathcal{U}^{-1}\pi_0^{\mathcal{Q}}\mathcal{U}$. According to Proposition 2, the transformed Kolmogorov operator $\tilde{\mathcal{K}}$ is of the form (33). We know that such operator has compact resolvent and a spectrum enclosed in cusp-shaped region of the complex plane (see Figure 1). Then, by Theorem 3, the operator $\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}}$ has exactly the same properties, provided $\tilde{\mathcal{P}}$ is a finite-rank projection. To derive the estimate (98) we use Theorem 4. To this end, we need to make sure that the following two conditions are satisfied

Condition 1. The spectral projection appearing in Theorem 4 can be replaced by a projection operator $\tilde{\pi}_0^{\tilde{\mathcal{Q}}}$ in $L^2(\mathbb{R}^{2d})$, which projects any $u \in L^2(\mathbb{R}^{2d})$ onto the linear subspace $\text{Ker}(\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}})$, i.e., $R(\tilde{\pi}_0^{\tilde{\mathcal{Q}}}) = \text{Ker}(\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}}) = \text{Id}(\tilde{\mathcal{P}}) \oplus \text{Ker}(\tilde{\mathcal{K}})$;

Condition 2. $\pi_0^{\mathcal{Q}}$ is an orthogonal projection in $L^2(\mathbb{R}^{2d}; \rho_{e_q})$ given by (99) (unique up to an isomorphism).

Proof of Condition 1. The projection operator $\tilde{\mathcal{P}}$ is self-adjoint, and therefore $\text{Id}(\tilde{\mathcal{P}}) = \text{Id}(\tilde{\mathcal{P}}^*)$. In [20], Helffer and Nier proved that $\text{Ker}(\tilde{\mathcal{K}}) = \text{Ker}(\tilde{\mathcal{K}}^*)$. Hence,

$$\text{Ker}(\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}}) = \text{Ker}(\tilde{\mathcal{Q}}\tilde{\mathcal{K}}^*\tilde{\mathcal{Q}}) = \text{Ker}(\tilde{\mathcal{K}}) \oplus \text{Id}(\tilde{\mathcal{P}}) \quad (101)$$

Next, consider the following orthogonal decomposition of the Hilbert space $L^2(\mathbb{R}^{2d})$

$$L^2(\mathbb{R}^{2d}) = \text{Ker}(\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}}) \oplus^{\perp} \text{Ker}(\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}})^{\perp},$$

If we define a projection operator $\pi_0^{\tilde{\mathcal{Q}}}$ with range $R(\pi_0^{\tilde{\mathcal{Q}}}) = \text{Ker}(\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}})$, then for any $u_0 \in L^2(\mathbb{R}^{2d})$, we have the orthogonal decomposition

$$\tilde{u}_0 = \pi_0^{\tilde{\mathcal{Q}}}\tilde{u}_0 + (\tilde{u}_0 - \pi_0^{\tilde{\mathcal{Q}}}\tilde{u}_0), \quad \text{where } \pi_0^{\tilde{\mathcal{Q}}}\tilde{u}_0 \in \text{Ker}(\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}}), \quad \tilde{u}_0 - \pi_0^{\tilde{\mathcal{Q}}}\tilde{u}_0 \in \text{Ker}(\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}})^{\perp}.$$

We now verify that $\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}}$ maps the linear subspace $\text{Ker}(\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}})^{\perp}$ into itself, i.e., that for any $u \in \text{Ker}(\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}})^{\perp}$ we have that $\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}}u \in \text{Ker}(\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}})^{\perp}$. To this end, we notice that if $w \in \text{Ker}(\tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}})$ then

$$\langle \tilde{\mathcal{Q}}\tilde{\mathcal{K}}\tilde{\mathcal{Q}}u, w \rangle = \langle u, \tilde{\mathcal{Q}}\tilde{\mathcal{K}}^*\tilde{\mathcal{Q}}w \rangle = 0.$$

This follows directly from $Ker(\tilde{Q}\tilde{K}\tilde{Q}) = Ker(\tilde{Q}\tilde{K}^*\tilde{Q})$. On the other hand, if $u \in Ker(\tilde{Q}\tilde{K}\tilde{Q})^\perp$ then $\tilde{Q}\tilde{K}\tilde{Q}u \neq 0$ and therefore we must have $\tilde{Q}\tilde{K}\tilde{Q}u \in Ker(\tilde{Q}\tilde{K}\tilde{Q})^\perp$. The operator $\tilde{Q}\tilde{K}\tilde{Q}$ and its adjoint $\tilde{Q}\tilde{K}^*\tilde{Q}$ can be decomposed as

$$\tilde{Q}\tilde{K}\tilde{Q} = \tilde{Q}\tilde{K}\tilde{Q}|_{Ker(\tilde{Q}\tilde{K}\tilde{Q})} \oplus \tilde{Q}\tilde{K}\tilde{Q}|_{Ker(\tilde{Q}\tilde{K}\tilde{Q})^\perp}, \quad \tilde{Q}\tilde{K}^*\tilde{Q} = \tilde{Q}\tilde{K}^*\tilde{Q}|_{Ker(\tilde{Q}\tilde{K}\tilde{Q})} \oplus \tilde{Q}\tilde{K}^*\tilde{Q}|_{Ker(\tilde{Q}\tilde{K}\tilde{Q})^\perp}.$$

This allows us to deform the domain of the Dunford integral representing $e^{t\tilde{Q}\tilde{K}\tilde{Q}}\tilde{u}_0 - \tilde{\pi}_0^{\tilde{Q}}\tilde{u}_0$ from $[-i\infty, +i\infty]$ to the cusp $S'_{\tilde{Q}\tilde{K}\tilde{Q}}$, as we did in Theorem 2. This yields

$$e^{t\tilde{Q}\tilde{K}\tilde{Q}}\tilde{u}_0 - \tilde{\pi}_0^{\tilde{Q}}\tilde{u}_0 = e^{t\tilde{Q}\tilde{K}\tilde{Q}} \left(\tilde{u}_0 - \tilde{\pi}_0^{\tilde{Q}}\tilde{u}_0 \right) = \frac{1}{2\pi i} \int_{\partial S'_{\tilde{Q}\tilde{K}\tilde{Q}}} e^{-tz} \left(z - \tilde{Q}\tilde{K}\tilde{Q} \right)^{-1} \tilde{u}_0 dz.$$

At this point we can follow the exact same procedure in the proofs of Theorem 2 and Theorem 3, to prove the semigroup estimate (98).

Proof of Condition 2. Here we show that $\tilde{\pi}_0^{\tilde{Q}}$ is an orthogonal projection in $L^2(\mathbb{R}^{2d})$ given by

$$\tilde{\pi}_0^{\tilde{Q}}(\cdot) = \tilde{\pi}_0(\cdot) + \tilde{P}(\cdot) = \langle \cdot \rangle_{eq/2} e^{-\beta H/2} + \sum_{i=1}^m \langle \cdot, g_i(\mathbf{q}) \rangle_{eq/2} g_i(\mathbf{q}) e^{-\beta H/2}. \quad (102)$$

The projection $\tilde{\pi}_0^{\tilde{Q}}$ is unique up to an isomorphism. Since $\tilde{\pi}_0^{\tilde{Q}}$ and its adjoint are both projections, they can be used to decompose $L^2(\mathbb{R}^{2d})$ as

$$L^2(\mathbb{R}^{2d}) = Ker\left(\tilde{\pi}_0^{\tilde{Q}}\right) \oplus R\left(\tilde{\pi}_0^{\tilde{Q}}\right), \quad L^2(\mathbb{R}^{2d}) = Ker\left([\tilde{\pi}_0^{\tilde{Q}}]^*\right) \oplus R\left([\tilde{\pi}_0^{\tilde{Q}}]^*\right)$$

It follows from (101) that $R(\tilde{\pi}_0^{\tilde{Q}}) = R([\tilde{\pi}_0^{\tilde{Q}}]^*) = Ker(\tilde{Q}\tilde{K}\tilde{Q})$. This implies $Ker(\tilde{\pi}_0^{\tilde{Q}}) = Ker([\tilde{\pi}_0^{\tilde{Q}}]^*)$. Since for all $u, w \in L^2(\mathbb{R}^{2d})$, $w - \tilde{\pi}_0^{\tilde{Q}}w \in Ker(\tilde{\pi}_0^{\tilde{Q}})$, we have that

$$\langle \tilde{\pi}_0^{\tilde{Q}}u, w - \tilde{\pi}_0^{\tilde{Q}}w \rangle = \langle u, [\tilde{\pi}_0^{\tilde{Q}}]^*(w - \tilde{\pi}_0^{\tilde{Q}}w) \rangle = 0.$$

Therefore $\tilde{\pi}_0^{\tilde{Q}}$ is an orthogonal projection operator. On the other hand, since $Id(\tilde{P}) = R(\tilde{P})$ and \tilde{P} is an orthogonal projection, we have that $\tilde{\pi}_0^{\tilde{Q}}$ is given by (102) (unique up to isomorphism). \square

5.1. EMZ memory and fluctuation terms

Theorem 3 allows us to prove that the EMZ memory kernel (22c) and the fluctuation term (22d) of the particle system converge exponentially fast to an equilibrium state for any observable (3). Specifically,

Corollary 5.1. *Suppose that the particle interaction potential $V(\mathbf{q})$ in (77) satisfies Hypothesis 1. Let \mathcal{P} be a Mori-type projection operator associated with the scalar observable $u(t) = u(x(t))$ such that $Ker(\mathcal{Q}\mathcal{K}\mathcal{Q}) = Ker(\mathcal{K}) \oplus Id(\mathcal{P})$. Then the memory kernel of (22c) converges exponentially converges to the equilibrium state $\mathbb{E}[\mathcal{K}u_0]\mathbb{E}[\mathcal{Q}\mathcal{K}^*u_0]$ with rate $\alpha_{\mathcal{Q}}$, i.e.*

$$|K(t) - \mathbb{E}[\mathcal{K}u_0]\mathbb{E}[\mathcal{Q}\mathcal{K}^*u_0]| \leq C e^{-\alpha_{\mathcal{Q}}t}. \quad (103)$$

Proof. The Corollary follows immediately from (68), (99) and the fact that $\mathcal{P}\mathcal{Q} = 0$. \square

We emphasize that the equilibrium state $\mathbb{E}[\mathcal{K}u_0]\mathbb{E}[\mathcal{Q}\mathcal{K}^*u_0]$ can be explicitly calculated. It is straightforward to extend (103) to matrix-valued memory kernels (22c). By following the same steps that lead us to (71), we obtain

$$\|\mathbf{K}(t) - \mathbf{G}^{-1}\mathbf{C}^{\mathcal{Q}}\|_M \leq C \|\mathbf{G}^{-1}\mathbf{D}^{\mathcal{Q}}\|_M e^{-\alpha_{\mathcal{Q}}t}, \quad (104)$$

where $\|\cdot\|_M$ denotes any matrix norm, and \mathbf{G} is the Gram matrix (22a). The entries of the matrix $\mathbf{C}^\mathcal{Q}$ are given by $C_{ij}^\mathcal{Q} = \mathbb{E}[\mathcal{Q}\mathcal{K}^*u_i(0)]\mathbb{E}[\mathcal{K}u_j(0)]$, while $D_{ij}^\mathcal{Q} = \mathbb{E}[\mathcal{Q}\mathcal{K}u_i(0)]\mathbb{E}[\mathcal{Q}\mathcal{K}u_j(0) - \mathbb{E}[\mathcal{K}u_j(0)]]$.

The components of the EMZ fluctuation term (22d) decay to an equilibrium state as well, exponentially fast in time. In fact, if we choose the initial condition as $\mathbf{u}_0 = \mathcal{Q}\mathcal{K}\mathbf{u}_0$, then (66) yields the following $L^2(\mathbb{R}^n, \rho_{eq})$ -equivalent estimate

$$\|f_i(t) - \mathbb{E}[\mathcal{Q}\mathcal{K}u_i(0)]\|_M \leq Ce^{-\alpha_\mathcal{Q}t}\mathbb{E}[\mathcal{Q}\mathcal{K}u_i(0)]. \quad (105)$$

The inequality (105) can be written in a vector form as⁹

$$\|\mathbf{f}(t) - \mathbb{E}[\mathcal{Q}\mathcal{K}\mathbf{u}(0)]\|_{V_{eq}} \leq Ce^{-\alpha_\mathcal{Q}t} \left(\sqrt{\mathbb{E}[(\mathcal{Q}\mathcal{K}u_1(0))^2]}, \dots, \sqrt{\mathbb{E}[(\mathcal{Q}\mathcal{K}u_m(0))^2]} \right) \|_M, \quad (106)$$

where $\|\cdot\|_{V_{eq}}$ is a norm in the tensor product space $V_{eq} = \otimes_{i=1}^m L_i^2(\mathbb{R}^n, \rho_{eq})$, defined similarly to (74).

6. Summary

We developed an in-dept analysis of the Mori-Zwanzig formulation for nonlinear dynamical systems driven by multiplicative Gaussian white noise, and evolving from random initial states. To this end, we first derived a new type of MZ equation, which we called effective Mori-Zwanzig (EMZ) equation, that governs the temporal dynamics of noise-averaged observables. Such dynamics is generated by a Kolmogorov operator obtained by averaging the Itô's representation of the system's Liouvillian. Building upon recent work of Eckmann, Hairer, Helffer, Hérau and Nier [13, 20] on the spectral properties of hypoelliptic operators, we proved that the EMZ memory kernel and fluctuation terms converge to a computable equilibrium state exponentially fast in time. This allows us to effectively study the asymptotic dynamics of any smooth quantity of interest depending on the stochastic flow. We applied our theoretical results to the Langevin dynamics of a particle system widely used in molecular dynamics simulations, and proved that for smooth, polynomial bounded particle interaction potentials, the EMZ memory and fluctuation terms decay exponentially fast to statistical equilibrium. We conclude by emphasizing that the mathematical framework we developed in this paper can be generalized and applied to other stochastic dynamical systems, e.g., systems driven by fractional Brownian motion with anomalous long-time behavior [1, 10, 32].

Acknowledgements This research was partially supported by the Air Force Office of Scientific Research (AFOSR) grant FA9550-16-586-1-0092. We would like to thank Prof. F. Hérau, Prof. B. Helffer and Prof. F. Nier for helpful discussions on the spectral analysis of Kolmogorov operators.

References

- [1] A. Bazzani, G. Bassi, and G. Turchetti. Diffusion and memory effects for stochastic processes and fractional Langevin equations. *Physica A*, 324(3-4):530–550, 2003.
- [2] C. Brennan and D. Venturi. Data-driven closures for stochastic dynamical systems. *J. Comp. Phys.*, 372:281–298, 2018.
- [3] A. Budhiraja, P. Dupuis, and V. Maroulas. Large deviations for stochastic flows of diffeomorphisms. *Bernoulli*, (1):234–257, 2010.
- [4] H. Cho, D. Venturi, and G. E. Karniadakis. Statistical analysis and simulation of random shocks in Burgers equation. *Proc. R. Soc. A*, 2171(470):1–21, 2014.
- [5] A. J. Chorin, O. H. Hald, and R. Kupferman. Optimal prediction and the Mori-Zwanzig representation of irreversible processes. *Proc. Natl. Acad. Sci. USA*, 97(7):2968–2973, 2000.

⁹Estimating the convergence constant $\alpha_\mathcal{Q}$ in (104) and (106) is non-trivial. Such constant coincides with the real part of the smallest non-zero eigenvalue of $\mathcal{Q}\mathcal{K}\mathcal{Q}$.

- [6] A. J. Chorin, O. H. Hald, and R. Kupferman. Optimal prediction with memory. *Physica D*, 166(3-4):239–257, 2002.
- [7] A. J. Chorin, R. Kupferman, and D. Levy. Optimal prediction for Hamiltonian partial differential equations. *J. Comput. Phys.*, 162(1):267–297, 2000.
- [8] A. J. Chorin and P. Stinis. Problem reduction, renormalization and memory. *Comm. App. Math. and Comp. Sci.*, 1(1):1–27, 2006.
- [9] E. Darve, J. Solomon, and A. Kia. Computing generalized Langevin equations and generalized Fokker-Planck equations. *Proc. Natl. Acad. Sci. USA*, 106(27):10884–10889, 2009.
- [10] S. Denisov, W. Horsthemke, and P. Hänggi. Generalized Fokker-Planck equation: Derivation and exact solutions. *Eur. Phys. J. B*, 68(4):567–575, 2009.
- [11] J. M. Dominy and D. Venturi. Duality and conditional expectations in the Nakajima-Mori-Zwanzig formulation. *J. Math. Phys.*, 58(8):082701, 2017.
- [12] J. P. Eckmann and M. Hairer. Non-equilibrium statistical mechanics of strongly anharmonic chains of oscillators. *Commun. Math. Phys.*, 212(1):105–164, 2000.
- [13] J. P. Eckmann and M. Hairer. Spectral properties of hypoelliptic operators. *Commun. Math. Phys.*, 235(2):233–253, 2003.
- [14] J. P. Eckmann, C. A. Pillet, and L. Rey-Bellet. Non-equilibrium statistical mechanics of anharmonic chains coupled to two heat baths at different temperatures. *Commun. Math. Phys.*, 201(3):657–697, 1999.
- [15] K.-J. Engel and R. Nagel. *One-parameter semigroups for linear evolution equations*, volume 194. Springer, 1999.
- [16] P. Espanol. Hydrodynamics from dissipative particle dynamics. *Phys. Rev. E*, 52(2):1734, 1995.
- [17] P. Espanol and P. Warren. Statistical mechanics of dissipative particle dynamics. *EPL*, 30(4):191, 1995.
- [18] S. K. J. Falkena, C. Quinn, J. Sieber, J. Frank, and H. A. Dijkstra. Derivation of delay equation climate models using the Mori-Zwanzig formalism. *Proc. R. Soc. A*, 475, 2019.
- [19] D. Givon, R. Kupferman, and O. H. Hald. Existence proof for orthogonal dynamics and the Mori-Zwanzig formalism. *Isr. J. Math.*, 145(1):221–241, 2005.
- [20] B. Helffer and F. Nier. *Hypoelliptic estimates and spectral theory for Fokker-Planck operators and Witten Laplacians*. Springer, 2005.
- [21] F. Hérau and F. Nier. Isotropic hypoellipticity and trend to equilibrium for the Fokker-Planck equation with a high-degree potential. *Arch. Ration. Mech. Anal.*, 171(2):151–218, 2004.
- [22] T. Hudson and X. Li. Coarse-graining of overdamped Langevin dynamics via the Mori-Zwanzig formalism. *arXiv:1810.08175*, 2018.
- [23] N. G. Van Kampen and I. Oppenheim. Brownian motion as a problem of eliminating fast variables. *Physica A*, 138(1-2):231–248, 1986.
- [24] T. Kato. *Perturbation theory for linear operators*, volume 132. Springer Science & Business Media, 2013.
- [25] P. E. Kloeden and E. Platen. *Numerical solution of stochastic differential equations*, volume 23. Springer Science & Business Media, 2013.
- [26] H. Kunita. *Stochastic flows and stochastic differential equations*. Cambridge university press, 1997.

- [27] H. Lei, N.A. Baker, and X. Li. Data-driven parameterization of the generalized Langevin equation. *Proc. Natl. Acad. Sci.*, 113(50):14183–14188, 2016.
- [28] Z. Li, , X. Bian, X. Li, and G. E. Karniadakis. Incorporation of memory effects in coarse-grained modeling via the Mori-Zwanzig formalism. *J. Chem. Phys.*, 143:243128, 2015.
- [29] K. K. Lin and F. Lu. Data-driven model reduction, Wiener projections, and the Mori-Zwanzig formalism. *arXiv*, pages 1908.07725(1–30), 2019.
- [30] F. Lu, K. K. Lin, and A. J. Chorin. Data-based stochastic model reduction for the Kuramoto–Sivashinsky equation. *Physica D*, 340:46–57, 2017.
- [31] H. Mori. Transport, collective motion, and Brownian motion. *Prog. Theor. Phys.*, 33(3):423–455, 1965.
- [32] T. Morita, H. Mori, and K. Mashiyama. Contraction of state variables in non-equilibrium open systems. II. *Prog. Theor. Phys.*, 64(2):500–521, 1980.
- [33] M. Ottobre, G. A. Pavliotis, and K. P. Starov. Exponential return to equilibrium for hypoelliptic quadratic systems. *Journal of Functional Analysis*, 262:4000–4039, 2012.
- [34] E. J. Parish and K. Duraisamy. A dynamic subgrid scale model for large eddy simulations based on the Mori–Zwanzig formalism. *J. Comp. Phys.*, 349:154–175, 2017.
- [35] E. J. Parish and K. Duraisamy. Non-Markovian closure models for large eddy simulations using the Mori-Zwanzig formalism. *Phys. Rev. Fluids*, 2(1):014604, 2017.
- [36] M. E. Peskin. *An introduction to quantum field theory*. CRC Press, 2018.
- [37] G. Da Prato and J. Zabczyk. *Ergodicity for infinite dimensional systems*, volume 229. Cambridge University Press, 1996.
- [38] H. Risken. *The Fokker-Planck equation: methods of solution and applications*. Springer-Verlag, second edition, 1989. Mathematics in science and engineering, vol. 60.
- [39] I. Snook. *The Langevin and generalised Langevin approach to the dynamics of atomic, polymeric and colloidal systems*. Elsevier, 2006.
- [40] P. Stinis. Stochastic optimal prediction for the Kuramoto–Sivashinsky equation. *Multiscale Modeling & Simulation*, 2(4):580–612, 2004.
- [41] P. Stinis. Renormalized reduced models for singular PDEs. *Comm. Appl. Math. and Comput. Sci.*, 8(1):39–66, 2013.
- [42] P. Stinis. Renormalized Mori–Zwanzig-reduced models for systems without scale separation. *Proc. Royal Soc. A*, 471(2176):20140446, 2015.
- [43] D. Venturi. The numerical approximation of nonlinear functionals and functional differential equations. *Physics Reports*, 732:1–102, 2018.
- [44] D. Venturi, H. Cho, and G. E. Karniadakis. The Mori-Zwanzig approach to uncertainty quantification. In R. Ghanem, D. Higdon, and H. Owhadi, editors, *Handbook of uncertainty quantification*. Springer, 2016.
- [45] D. Venturi, M. Choi, and G. E. Karniadakis. Supercritical quasi-conduction states in stochastic Rayleigh-Bénard convection. *Int. J. Heat and Mass Transfer*, 55(13-14):3732–3743, 2012.
- [46] D. Venturi and G. E. Karniadakis. New evolution equations for the joint response-excitation probability density function of stochastic solutions to first-order nonlinear PDEs. *J. Comput. Phys.*, 231:7450–7474, 2012.
- [47] D. Venturi and G. E. Karniadakis. Convolutionless Nakajima-Zwanzig equations for stochastic analysis in nonlinear dynamical systems. *Proc. R. Soc. A*, 470(2166):1–20, 2014.

- [48] D. Venturi, T. P. Sapsis, H. Cho, and G. E. Karniadakis. A computable evolution equation for the joint response-excitation probability density function of stochastic dynamical systems. *Proc. R. Soc. A*, 468(2139):759–783, 2012.
- [49] C. Villani. Hypocoercivity. *Memoirs of the American Mathematical Society*, 202(950), 2009.
- [50] S. Watanabe. *Lectures on Stochastic Differential Equations and Malliavin Calculus*. Lectures delivered at the Indian Institute of Science, Bangalore, 1984.
- [51] V. Wihstutz and MA. Pinsky. *Diffusion processes and related problems in analysis, volume II: Stochastic flows*, volume 27. Springer Science & Business Media, 2012.
- [52] Y. Yoshimoto, I. Kinefuchi, T. Mima, A. Fukushima, T. Tokumasu, and S. Takagi. Bottom-up construction of interaction models of non-Markovian dissipative particle dynamics. *Phys. Rev. E*, 88(4):043305, 2013.
- [53] Y. Zhu, J. M. Dominy, and D. Venturi. On the estimation of the Mori-Zwanzig memory integral. *J. Math. Phys.*, 59(10):103501, 2018.
- [54] Y. Zhu and D. Venturi. Faber approximation of the Mori-Zwanzig equation. *J. Comp. Phys.*, (372):694–718, 2018.
- [55] Y. Zhu and D. Venturi. Generalized Langevin equations for systems with local interactions. *arXiv*, pages 1906.04918(1–27), 2019.
- [56] J. Zinn-Justin. *Quantum field theory and critical phenomena*. Oxford Univ. Press, fourth edition, 2002.
- [57] R. Zwanzig. Memory effects in irreversible thermodynamics. *Phys. Rev.*, 124(4):983, 1961.
- [58] R. Zwanzig. Nonlinear generalized Langevin equations. *J. Stat. Phys.*, 9(3):215–220, 1973.