arXiv:2001.04614v1 [math.RT] 14 Jan 2020

A NOTE ON SINGULAR EQUIVALENCES AND IDEMPOTENTS

DAWEI SHEN

Abstract. Let Λ be an Artin algebra and let e be an idempotent in Λ. We study certain functors which preserve the singularity categories. Suppose $\operatorname{pd}\Lambda e_{e\Lambda e}<\infty$ and $\operatorname{id}{}_{\Lambda}\frac{\Lambda/\langle e\rangle}{\operatorname{rad}\Lambda/\langle e\rangle}<\infty$, we show that there is a singular equivalence between $e\Lambda e$ and Λ.

1. Introduction

Let R be a left noetherian ring. Let $\operatorname{mod} R$ be the category of finitely generated left R-modules and let $\operatorname{proj} R$ be the full subcategory of projective left R-modules. A complex in $\operatorname{mod} R$ is called *perfect* if it is quasi-isomorphic to a bounded complex in $\operatorname{proj} R$. The *singularity category* of R is the Verdier quotient of the bounded derived category of $\operatorname{mod} R$ by the thick subcategory of perfect complexes [2, 10]. If there is a triangle equivalence between the singularity categories of two rings R and S, then such an equivalence is called a *singular equivalence* [4].

Let e be an idempotent in R. There is a recollement

$$\operatorname{mod} R/\langle e\rangle \overset{\longleftarrow}{\longrightarrow}\overset{\longleftarrow}{\longrightarrow} \operatorname{mod} R \overset{\longleftarrow i_\lambda \longrightarrow}{\longleftarrow i \longrightarrow} \operatorname{mod} eRe.$$

Here, i takes any M in $\operatorname{mod} R$ to eM and i_λ takes any N in $\operatorname{mod} eRe$ to $Re\otimes_{eRe}N$. Suppose

(1) $\operatorname{pd}{}_{eRe}eR<\infty$, and
(2) $\operatorname{pd}{}_R M<\infty$ for every $M\in\operatorname{mod} R$ annihilated by e,

then i induces a singular equivalence between R and eRe; see [3].

In the present paper, we investigate the singular equivalence induced by the functor i_λ. We have the following

Theorem I. *Let R be a left noetherian ring and e be an idempotent in R. Suppose*

(1) $\operatorname{pd} Re_{eRe}<\infty$, *and*
(2) *every $M\in\operatorname{mod} R$ admits a projective resolution P such that $P^{-i}\in\operatorname{add} Re$ for every sufficiently large i,*

then i_λ induces singular equivalence between eRe and R.

2010 *Mathematics Subject Classification.* Primary 18E30; Secondary 16G10.
Key words and phrases. Singularity category, triangular matrix ring.

Let Λ be an Artin algebra and e be an idempotent in Λ. The following conditions are studied in [11].

$$(\alpha)\ \mathrm{id}\,{}_{\Lambda}\tfrac{\Lambda/\langle e\rangle}{\mathrm{rad}\,\Lambda/\langle e\rangle} < \infty \qquad (\beta)\ \mathrm{pd}\,{}_{e\Lambda e}e\Lambda < \infty$$
$$(\gamma)\ \mathrm{pd}\,{}_{\Lambda}\tfrac{\Lambda/\langle e\rangle}{\mathrm{rad}\,\Lambda/\langle e\rangle} < \infty \qquad (\delta)\ \mathrm{pd}\,{}_{\Lambda}e\Lambda_{e\Lambda e} < \infty$$

It turns out that

(1) (α) and (β) hold if and only if (γ) and (δ) hold;
(2) i induces a singular equivalence if and only if (β) and (γ) hold.

As a complement, we have the following

Theorem II. *Let Λ be an Artin algebra and e be an idempotent in Λ. Then i_λ induces a singular equivalence if and only if (α) and (δ) hold.*

2. Singularity categories

Let R be a left noetherian ring. Let $\mathbf{K}^{-,\mathrm{b}}(\mathrm{proj}\,R)$ be the homotopy category of bounded above complexes in $\mathrm{proj}\,R$ with bounded cohomologies. It is a triangulated category, whose translation functor Σ is the shift of complexes [12].

A complex P in $\mathbf{K}^{-,\mathrm{b}}(\mathrm{proj}\,R)$ is perfect if there is an integer ℓ such that the $(-i)$-th coboundary of P is projective for every $i \geq \ell$. Let $\mathbf{K}^{\mathrm{b}}(\mathrm{proj}\,R)$ be the full subcategory of perfect complexes.

The singularity category of R is the quotient triangulated category

$$\mathbf{D}_{\mathrm{sg}}(R) \coloneqq \mathbf{K}^{-,\mathrm{b}}(\mathrm{proj}\,R)/\mathbf{K}^{\mathrm{b}}(\mathrm{proj}\,R).$$

Let $\underline{\mathrm{mod}}\,R$ be the projectively stable category of $\mathrm{mod}\,R$. For any M in $\mathrm{mod}\,R$, take a projective precover $\pi\colon P \to M$; the syzygy ΩM of M is the kernel of π. There is a functor

$$\Omega\colon\ \underline{\mathrm{mod}}\,R \to \underline{\mathrm{mod}}\,R$$

sending a module M to the syzygy ΩM. For any M in $\mathrm{mod}\,R$, take a projective resolution pM of M. There is a functor

$$p\colon\ \underline{\mathrm{mod}}\,R \to \mathbf{D}_{\mathrm{sg}}(R)$$

sending a module M to the projective resolution pM. For any $\ell \geq 0$, there is a natural isomorphism

$$p\Omega^{\ell}(M) \cong \Sigma^{-\ell}pM.$$

Lemma 2.1 ([4, Lemma 2.1])**.** *For any $X \in \mathbf{D}_{\mathrm{sg}}(R)$, there is $n \geq 0$ and $M \in \mathrm{mod}\,R$ such that*

$$\Sigma^{-n}X \cong pM.$$

Lemma 2.2 ([9, Exemple 2.3])**.** *For any $M, N \in \mathrm{mod}\,R$, there is an isomorphism*

$$\varinjlim_{\ell\geq 0} \underline{\mathrm{Hom}}_R(\Omega^{\ell}M, \Omega^{\ell}N) \cong \mathrm{Hom}_{\mathbf{D}_{\mathrm{sg}}(R)}(pM, pN).$$

Let e be an idempotent in R. There is a functor

$$i_\lambda\colon\ \mathrm{mod}\,eRe \to \mathrm{mod}\,R$$

such that $i_\lambda(M) = Re \otimes_{eRe} M$ for every $M \in \operatorname{mod} eRe$. It is a full faithful functor which takes projectives to projectives. The functor i_λ restricts to a full faithful functor i'_λ such that

$$i'_\lambda \colon \operatorname{proj} eRe \xrightarrow{\simeq} \operatorname{add} Re \xrightarrow{\subseteq} \operatorname{proj} R.$$

Here, we denote by $\operatorname{add} Re$ the full subcategory of summands of finite direct sums of copies of Re.

We now restate and prove Theorem I.

Theorem 2.3. *Suppose* $\operatorname{pd} Re_{eRe} < \infty$*, then* i_λ *induces a full faithful triangle functor*

$$\mathbf{D}_{\mathrm{sg}}(i_\lambda) \colon \mathbf{D}_{\mathrm{sg}}(eRe) \to \mathbf{D}_{\mathrm{sg}}(R).$$

Moreover, $\mathbf{D}_{\mathrm{sg}}(i_\lambda)$ *is a triangle equivalence if and only if for every* $M \in \operatorname{mod} R$*, there is an integer* ℓ *and a projective resolution* P *of* M *such that* $P^{-i} \in \operatorname{add} Re$ *for every* $i \geq \ell$*.*

Proof. Let X be in $\mathbf{K}^{-,\mathrm{b}}(\operatorname{proj} eRe)$ which is exact at degree $\leq \ell$. Then $i_\lambda(X)$ is exact at degree $\leq \ell - \operatorname{pd} Re_{eRe}$. Therefore i_λ induces a triangle functor

$$\mathbf{K}^{-,\mathrm{b}}(i_\lambda) \colon \mathbf{K}^{-,\mathrm{b}}(\operatorname{proj} eRe) \to \mathbf{K}^{-,\mathrm{b}}(\operatorname{proj} R).$$

Since i_λ is full faithful, $\mathbf{K}^{-,\mathrm{b}}(i_\lambda)$ is also full faithful. Since i_λ preserves perfect complexes, it induces a triangle functor $\mathbf{D}_{\mathrm{sg}}(i_\lambda)$ such that

$$\mathbf{D}_{\mathrm{sg}}(i_\lambda) \colon \mathbf{D}_{\mathrm{sg}}(eRe) \xrightarrow{\simeq} \mathbf{K}^{-,\mathrm{b}}(\operatorname{add} Re)/\mathbf{K}^{\mathrm{b}}(\operatorname{add} Re) \to \mathbf{D}_{\mathrm{sg}}(R).$$

Let $f \colon X \to Y$ be a morphism in $\mathbf{K}^{-,\mathrm{b}}(\operatorname{proj} R)$, where X^i is zero for every $i \leq \ell$ and Y belongs to $\mathbf{K}^{-,\mathrm{b}}(\operatorname{add} Re)$. Let $\sigma_{\geq \ell} Y$ be the stupid truncation of Y at degree $\geq \ell$, then it belongs to $\mathbf{K}^{\mathrm{b}}(\operatorname{add} Re)$. Since f factors through $\sigma_{\geq \ell} Y$, by [8, Proposition 10.2.6] the functor $\mathbf{D}_{\mathrm{sg}}(i_\lambda)$ is full faithful.

By Lemma 2.1 and 2.2, the singularity category of R is triangle equivalent to the stabilization of the stable category $\underline{\operatorname{mod}}\, R$. Then the denseness of $\mathbf{D}_{\mathrm{sg}}(i_\lambda)$ follows from [5, Corollary 2.13]. □

3. Triangular matrix rings

Let T and S be two rings and M be an S-T-bimodule. We consider the triangular matrix ring

$$R = \begin{pmatrix} T & 0 \\ M & S \end{pmatrix}.$$

Following [1, III.2] a left R-module is given by

$$(X, Y, \phi) = \left\{ \begin{pmatrix} x \\ y \end{pmatrix} \mid x \in X, y \in Y \right\},$$

where X is a left T-module, Y is a left S-module and $\phi \colon M \otimes_T X \to Y$ is a left S-module map. The action is given by

$$\begin{pmatrix} t & 0 \\ m & s \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} tx \\ \phi(m \otimes x) + sy \end{pmatrix}$$

for every $t \in T$, $s \in S$ and $m \in M$.

Let $e = \operatorname{diag}(0, 1)$ be in R. Then $Re = eRe$ and

$$\operatorname{add} Re = \{(0, Y, 0) \mid Y \in \operatorname{proj} S\}.$$

Lemma 3.1. *Let X be a left T-module and Y be a left S-module.*
(1) $\operatorname{pd}{}_S Y = \operatorname{pd}{}_R(0, Y, 0)$*;*
(2) $(X, M \otimes_T X, 1)$ *is a projective left R-module if and only if X is a projective left T-module.*

We have the following; compare [3, Theorem 4.1].

Proposition 3.2. *Let T and S be left noetherian rings, and let M be an S-T-bimodule such that ${}_S M$ is finitely generated. Assume that T has finite left global dimension, then there is a triangle equivalence*

$$\mathbf{D}_{\mathrm{sg}}(\begin{pmatrix} T & 0 \\ M & S \end{pmatrix}) \cong \mathbf{D}_{\mathrm{sg}}(S).$$

Proof. One checks that the triangular matrix ring R is left noetherian. Let (X, Y, ϕ) be in $\operatorname{mod} R$. Since $\operatorname{pd}{}_T X$ is finite, by Lemma 3.1 there a projective resolution P of (X, Y, ϕ) such that $P^{-i} \in \operatorname{add} Re$ for every $i > \operatorname{pd}{}_T X$. Then by Theorem I there is a triangle equivalence between $\mathbf{D}_{\mathrm{sg}}(R)$ and $\mathbf{D}_{\mathrm{sg}}(S)$. □

Example 3.3 (see [4], Proposition 4.1)**.** Let k be a field, Λ be a finite dimensional k-algebra and M be a finite dimensional left Λ-module. Then M is a Λ-k-bimodule. The *one-point extension* of Λ by M is the triangular matrix algebra

$$\Lambda[M] = \begin{pmatrix} k & 0 \\ M & \Lambda \end{pmatrix}.$$

By Theorem I there a singular equivalence between Λ and $\Lambda[M]$.

4. Artin algebras

Let Λ be an Artin algebra. We need the following well known

Lemma 4.1. *Let M be in $\operatorname{mod}\Lambda$ and P_M be a minimal projective resolution of M. For any semi-simple Λ-module S and any $i \geq 0$ there is an isomorphism*

$$\operatorname{Ext}^i_\Lambda(M, S) \cong \operatorname{Hom}_\Lambda(P_M^{-i}, S).$$

Proof. For $i = 0$, it is obvious. For $i = 1$, let K be the (-1)-th coboundary of P_M. There is an exact sequence

$$K \rightarrowtail P_M^0 \overset{\pi}{\twoheadrightarrow} M.$$

Applying $\operatorname{Hom}_\Lambda(-, S)$ to it, we obtain an exact sequence

$$\operatorname{Hom}_\Lambda(M, S) \overset{\pi^*}{\rightarrowtail} \operatorname{Hom}_\Lambda(P_M^0, S) \to \operatorname{Hom}_\Lambda(K, S) \twoheadrightarrow \operatorname{Ext}^1_\Lambda(M, S).$$

Since π is a projective cover and S is semi-simple, π^* is surjective. Then

$$\operatorname{Hom}_\Lambda(K, S) \cong \operatorname{Ext}^1_\Lambda(M, S).$$

Since P_M^{-1} is a projective cover of K, there is an isomorphism

$$\operatorname{Ext}^1_\Lambda(M, S) \cong \operatorname{Hom}_\Lambda(P_M^{-1}, S).$$

By shifting one proves the isomorphism for $i \geq 2$. □

For any $P \in \operatorname{proj}\Lambda$, we have the following

$$P \in \operatorname{add}\Lambda e \iff \operatorname{Hom}_\Lambda(P, \tfrac{\Lambda/\langle e\rangle}{\operatorname{rad}\Lambda/\langle e\rangle}) = 0. \tag{4.1}$$

We now restate and prove Theorem II.

Theorem 4.2. *Let Λ be an Artin algebra and e be an idempotent in Λ. Suppose $\operatorname{pd}\Lambda e_{e\Lambda e} < \infty$, then $\mathbf{D}_{\mathrm{sg}}(i_\lambda)$ is a triangle equivalence if and only if $\operatorname{id}_\Lambda \frac{\Lambda/\langle e\rangle}{\operatorname{rad}\Lambda/\langle e\rangle} < \infty$.*

Proof. " $\Longrightarrow$ " Let P be a minimal projective resolution of $\Lambda/\operatorname{rad}\Lambda$. If $\mathbf{D}_{\mathrm{sg}}(i_\lambda)$ is dense, by Theorem I there is $\ell \geq 0$ such that $P^{-i} \in \operatorname{add}\Lambda e$ for every $i > \ell$. By Lemma 4.1 and (4.1) we have

$$\operatorname{Ext}^i_\Lambda(\Lambda/\operatorname{rad}\Lambda, \tfrac{\Lambda/\langle e\rangle}{\operatorname{rad}\Lambda/\langle e\rangle}) = 0.$$

Then $\operatorname{id}_\Lambda \frac{\Lambda/\langle e\rangle}{\operatorname{rad}\Lambda/\langle e\rangle}$ is finite.

" $\Longleftarrow$ " Let M be in $\operatorname{mod}\Lambda$ and P_M be a minimal projective resolution of M. If $\operatorname{id}_\Lambda \frac{\Lambda/\langle e\rangle}{\operatorname{rad}\Lambda/\langle e\rangle}$ is finite, by Lemma 4.1 we have

$$P_M^{-i} \in \operatorname{add}\Lambda e, \forall\, i > \operatorname{id}_\Lambda \tfrac{\Lambda/\langle e\rangle}{\operatorname{rad}\Lambda/\langle e\rangle}.$$

From Theorem I we infer that $\mathbf{D}_{\mathrm{sg}}(i_\lambda)$ is a triangle equivalence. □

We end this section by an example.

Let Λ be an Artin algebra. Let S be a semi-simple left Λ-module with $\operatorname{id}_\Lambda S \leq 1$. Denote by

$${}^\perp S = \{M \in \operatorname{mod}\Lambda \mid \operatorname{Hom}_\Lambda(M, S) = \operatorname{Ext}^1_\Lambda(M, S) = 0\}$$

the perpendicular category of S in $\operatorname{mod}\Lambda$. By Lemma 4.1 a finitely generated left Λ-module M belongs to ${}^\perp S$ if and only if M admits a projective presentation

$$P^{-1} \to P^0 \twoheadrightarrow M$$

such that both P^{-1} and P^0 belong to ${}^\perp S$.

Let e be an idempotent in Λ such that $\operatorname{proj}\Lambda \cap {}^\perp S$ coincides with $\operatorname{add}\Lambda e$. Recall the functor i_λ decomposes into

$$i_\lambda \colon \operatorname{mod} e\Lambda e \xrightarrow{\simeq} {}^\perp S \xrightarrow{\subseteq} \operatorname{mod}\Lambda.$$

We have the following; compare [6, Proposition 2.13].

Proposition 4.3. *Keep the notation as previous.*

(1) $\operatorname{gl.dim} e\Lambda e \leq \operatorname{gl.dim}\Lambda \leq \operatorname{gl.dim} e\Lambda e + 2$;

(2) *there is a singular equivalence between $e\Lambda e$ and Λ.*

Proof. (1) Let $\Omega^2 M$ be the minimal second syzygy for M in $\operatorname{mod}\Lambda$. Since $\operatorname{id}_\Lambda S \leq 1$, $\operatorname{Ext}^i_\Lambda(M, S) = 0$ for every $i \geq 2$. By Lemma 4.1 $\Omega^2 M \in {}^\perp S$. Since i_λ preserves projective resolutions, for any N in $\operatorname{mod} e\Lambda e$ we have

$$\operatorname{pd}_{e\Lambda e} N = \operatorname{pd}_\Lambda i_\lambda(N).$$

Then

$$\operatorname{pd}_\Lambda M \leq \operatorname{pd}_\Lambda \Omega^2 M + 2 = \operatorname{pd}_{e\Lambda e} i(\Omega^2 M) + 2.$$

(2) From [7, Proposition 1.1] we infer that ${}^\perp S$ is an exact subcategory of $\operatorname{mod}\Lambda$. Then the right $e\Lambda e$-module Λe is projective. By Theorem II there is a singular equivalence between $e\Lambda e$ and Λ. □

Acknowledgments

The author is thankful to Professor Xiao-Wu Chen for discussions. This work is supported by National Natural Science Foundation of China (No. 11801141).

References

[1] M. Auslander, I. Reiten, S. O. Smalø, *Representation theory of Artin algebras*, Cambridge Studies in Advanced Mathematics, 36. Cambridge University Press, Cambridge, 1995.

[2] R.-O. Buchweitz, *Maximal Cohen-Macaulay modules and Tate-cohomology over Gorenstein rings*, Unpublished manuscript, Available at: `http://hdl.handle.net/1807/16682`, 1987.

[3] X.-W. Chen, *Singularity categories, Schur functors and triangular matrix rings*, Algebr. Represent. Theory **12** (2009), 181–191.

[4] X.-W. Chen, *The singularity category of an algebra with radical square zero*, Doc. Math. **16** (2011), 921–936.

[5] X.-W. Chen, *The singularity category of a quadratic monomial algebra*, The Quarterly Journal of Mathematics, **69** (2018), 1015–1033.

[6] X.-W. Chen and Y. Ye, *Retractions and Gorenstein homological properties*, Algebr. Represent. Theory **17** (2014), 713–733.

[7] W. Geigle and L. Lenzing, *Perpendicular categories with applications to representations and sheaves*. J. Algebra **144** (1991), 273–343.

[8] M. Kashiwara and P. Schapira, *Categories and sheaves*, Grundlehren der Mathematischen Wissenschaften, 332. Springer-Verlag, Berlin, 2006.

[9] B. Keller and D. Vossieck, *Sous les catégories dérivées*, C. R. Acad. Sci. Paris Sér. I Math. **305** (1987), 225–228.

[10] D. Orlov, *Triangulated categories of singularities and D-branes in Landau-Ginzburg models*, Tr. Mat. Inst. Steklova **246** (2004), 240–262.

[11] C. Psaroudakis, Ø. Skartsæterhagen and Ø. Solberg, *Gorenstein categories, singular equivalences and finite generation of cohomology rings in recollements*, Trans. Amer. Math. Soc. Ser. B **1** (2014), 45–95.

[12] J.-L. Verdier, *Catégories dérivées: quelques résultats (état 0)*, Cohomologie étale, 262–311, Lecture Notes in Math., 569. Springer, Berlin, 1977.

School of Mathematics and Statistics, Henan University, 475004, Kaifeng, Henan, P. R. China

E-mail address: `sdw12345@mail.ustc.edu.cn`