

# CURVATURE BOUNDS FOR REGULARIZED RIEMANNIAN METRICS

DANIEL LUCKHARDT\* AND JAN-BERNHARD KORDASS§

**ABSTRACT.** We investigate regularization of riemannian metrics by mollification. Assuming both-sided bounds on the Ricci tensor and a lower injectivity radius bound we obtain a uniform estimate on the change of the sectional curvature. Actually, our result holds for any metric with a uniform bound on the  $W^{2,p}$ -harmonic radius.

## § 1. Introduction

The goal of this note is to show that regularization by a naive mollification of a riemannian metric satisfying certain geometric conditions can be set up to control the deviation of the sectional curvature.

A riemannian metric on a smooth manifold  $M$  is a section in the symmetric  $(0,2)$ -tensor bundle over  $M$ . The regularity of this section is then referred to as the regularity of the metric. Under certain geometric limit processes, it is a common phenomenon to loose a controlled level of regularity. Gromov-Hausdorff limits of isometry classes of smooth riemannian metrics that satisfy certain curvature bounds, are a prominent example thereof. We will briefly recall fundamental results in this area in § 2.4. In this work we are interested in procedures to regain regularity and at the same time control the deviation of curvature. Various such techniques have been studied with a view towards different goals (e.g. [CG85; Abr88]).

A fundamental tool in deriving and phrasing our results are *chart norms*, that control the regularity properties of the metric tensor and its derivatives in a given chart. Let  $\psi: (B(0,r),0) \rightarrow (M,p_0)$  be a harmonic chart in a smooth, pointed,  $n$ -dimensional riemannian manifold  $(M^n, g, p_0)$ , where we denote by  $B(0,r) \subset \mathbb{R}^n$  the open ball of radius  $r$  with respect to the euclidean norm. Recall that a chart is called *harmonic*, if the coordinate-functions of  $\psi$  are harmonic with respect to the Laplace-Beltrami operator of  $g$ . The harmonic chart norm  $\|\psi\|_{W^{m,p,r}}^{\text{harm}}$  is bounded by  $Q \geq 0$ , if  $Q$  gives control of the derivatives of the metric tensor, its inverse and its first  $m$  derivatives in the  $L^p$ -norm. We refer to § 2.3 for a more detailed explanation.

---

*Date:* October 2019.

\*Supported by the ISF project “Action now: geometry and dynamics of group actions”.

§Supported by the SNSF-Project 200021E-172469 and the DFG-Priority programme Geometry at infinity (SPP 2026).

Unless otherwise stated, we will always denote by  $M$  a smooth manifold of dimension  $n$ . Given a collection of charts  $\{\psi_i: B(0, r) \rightarrow M\}_{i \in I}$ , a partition of unity  $\{\rho_i\}_{i \in I}$ , and a fixed mollifier function  $\varphi_t$  for  $t \in (0, T]$ , we define a mollified riemannian metric  $g^{[t]}$  (cf. Definition 3.2). This metric is defined on  $\bigcup_{i \in I} \psi_i(B(0, r/2))$  and, provided bounds on the harmonic chart norms, has a curvature tensor that can be controlled as follows:

**Theorem A.** *Given  $p > 2n$  and  $r, Q > 0$ . Choose  $\beta \in (0, 1 - 2n/p)$ . Then there is some  $T \in (0, r/2)$  such that for any pointed smooth riemannian manifold  $(M, g, p_0)$  and any finite collection of charts with a corresponding partition of unity*

$$\{\psi_i: (B(0, r), 0) \rightarrow (M, \psi_i(0))\}_{i \in I}, \quad \vec{\varrho} := \{\varrho_i: M \rightarrow [0, 1]\}_{i \in I}$$

*we have that for any section  $\vec{v} \in (T\psi(B(0, r)))^{\times 3} \times T^*\psi(B(0, r))$*

$$\left( \forall i \in I: \|\psi_i\|_{W^{2,p,r}}^{\text{harm}} \leq Q \right) \implies R_{g^{[t]}}(x)(\vec{v}) - \sup_{B(x, e^{Qt})} R_g(\vec{v}) \leq C \|\vec{v}\|_g t^\beta$$

*for any  $t \in (0, T]$ ,  $x \in \bigcap_{i \in I} \psi_i(B(0, r/2))$  and with*

$$C = C(n, p, r, Q, \beta, \{\|\varrho_i \circ \psi_i\|_{C^2}\}_{i \in I}, \#I).$$

In the theorem, we denote by  $\|\vec{v}\|_g$  the semi-norm obtained as the supremum of the point-wise product  $\|v_1(x)\|_{g(x)} \cdots \|v_4(x)\|_{g(x)}$  (cf. §3 for further details).

The new metric  $g^{[t]}$  is obtained by applying a mollification operator  $P_t$  in the charts  $\psi_i$ . We can summarize the ansatz of our proof of this statement as follows:

$$(1.1) \quad \begin{aligned} R_{g^{[t]}}(x)(\vec{v}) &\leq P_t(R_g(\vec{v}))(x) + |R_{g^{[t]}}(x)(\vec{v}) - P_t(R_g(\vec{v}))(x)| \\ &\leq \sup_{y \in \psi(B(0, T))} R_g(y)(\vec{v}) + \|R_{g^{[t]}}(\vec{v}) - P_t R_g(\vec{v})\|_{L^\infty} \end{aligned}$$

where in the last step we used that convolution does not increase the supremum of a function. Hence it suffices to find an  $L^\infty$ -estimate for the “commutator”

$$R_{g^{[t]}} - P_t R_g \quad “= [R, P_t]g”$$

which will be obtained in Lemma 3.5.

Finally, we state our main result on the deviation of the sectional curvature. Define for a riemannian manifold  $(M, g)$

$$\overline{K}(g) := \sup K_g(x)(v, w),$$

$$\underline{K}(g) := \sup K_g(x)(v, w)$$

where the suprema are taken over all  $x \in M$  and  $v, w \in T_x M$  such that  $\langle v, w \rangle_g > 0$ . Moreover, we denote by  $\|(M, g)\|_{W^{2,p,r}}^{\text{harm}}$  the lowest bound  $Q$  such that for each point  $p_0 \in M$  there is a harmonic chart  $\psi: B(0, r) \rightarrow M$  mapping 0 to  $p_0$  such that the metric tensor  $\psi^*g$  on  $B(0, r)$  is controlled by  $Q$  with respect to the  $W^{2,p}$ -norm.

**Theorem B.** *Let  $p > 2n$ ,  $r, Q > 0$ , and  $\beta \in (0, 1 - 2n/p)$ . Then there is some  $T > 0$  such that for any smooth riemannian manifold  $(M, g)$  with  $\|(M, g)\|_{W^{2,p},r}^{\text{harm}} \leq Q$  there is a locally finite cover of charts*

$$\psi_i: (B(0, r), 0) \rightarrow (M, \psi_i(0)),$$

*indexed by  $i \in I$ , such that the mollified metrics  $g^{[t]}$  have at any  $x \in M$  sectional curvature  $K_{g^{[t]}}(x)$  in the interval*

$$(1.2) \quad \left[ \underline{K}(g \cdot |_{B(x,t)}), \bar{K}(g \cdot |_{B(x,t)}) \right] \cdot [1 - Ct, 1 + Ct] + [-Ct^\beta, Ct^\beta]$$

*for all  $t \in (0, T]$ , where  $C = C(n, Q, r, p)$ . Moreover, we have that*

$$(1.3) \quad \|M\|_{C^{2,\beta},r} \leq Q'$$

*for some  $Q' = Q'(n, \beta, Q)$ .*

The advantage of the regularized metric is especially that it admits point-wise control on the curvature tensor that is stable under Gromov-Hausdorff limits, i.e. as apparent from  $(2.K_{C^m,\alpha})$  below, a Gromov-Hausdorff limit of regularized metrics is again a riemannian manifold with  $\|M\|_{C^{2,\beta},r} \leq Q'$ .

**Corollary C.** *Let  $\iota > 0$  and  $\kappa \geq 0$ . Then there exist  $r > 0$  and  $T > 0$  such that for any smooth riemannian manifold  $(M, g)$  with*

$$\text{inj rad}(M) \geq \iota, \quad \|\text{Ric } M\|_{L^\infty} \leq \kappa$$

*there exist charts  $\psi_i: B(0, r) \rightarrow M$  such that the mollified metrics  $g^{[t]}$  have at any  $x \in M$  sectional curvature  $K_{P_t g}(x)$  in the interval (1.2) for all  $t \in (0, T]$ , where  $C = C(n, \iota, \kappa)$ . Moreover, the norm  $\|M\|_{C^{2,\beta},r}$  is bounded in terms of  $n, \beta$  and  $Q$ .*

**Corollary D.** *Let  $\delta \in (0, 1)$  and assume one of the conditions (1.4a) to (1.4d):*

$$\begin{aligned} (1.4a) \quad & \# \pi_2(M) < \infty, \text{ or} \\ (1.4b) \quad & K_g \in [\delta, 1] \text{ and} \\ (1.4c) \quad & \left\{ \begin{array}{l} \dim M \text{ is even, or} \\ \delta \geq 1/4 - \varepsilon', \quad \text{where } \varepsilon' \approx 10^{-6} \end{array} \right. \end{aligned}$$

$$(1.4d) \quad K_g \leq 1, \text{ Ric}_g \geq \delta, \text{ and } \dim M = 3.$$

*Then there exist charts  $\psi_i: B(0, r) \rightarrow M$  such that the mollified metrics  $g^{[t]}$  have at any  $x \in M$  sectional curvature  $K_{P_t g}(x)$  in the interval (1.2) for all  $t \in (0, T]$ , where  $C = C(n, \kappa)$ . Moreover, the norm  $\|M\|_{C^{2,\beta},r}$  is bounded in terms of  $n, \beta$  and  $Q$ .*

As mentioned before, there are other techniques towards a regularization of riemannian metrics. In [Abr88], Abresch constructs a smoothing operator  $S_\varepsilon$ , which satisfies  $\|\nabla^m R_{S_\varepsilon g}\| \leq C \frac{1}{(1+\varepsilon\Lambda)^{m+2}}$ , where  $C = C(n)$  is a constant and  $\Lambda$  is a bound on the sectional curvature. In particular,  $S_\varepsilon$  preserves isometries of the original metric. Similarly, it is known that given a bound on the sectional curvature, the application of Ricci flow amounts to

a regularization of the metric, which gives  $\|\nabla^m R_{g(t)}\| \leq C(n, m, t)$  [BMR84; Shi89; Ron96; Kap05; He16]. In contrast to our results, this again preserves the isometries of the original metric, but just as Abresch's result requires stronger bounds on the curvature. The crucial point is that both methods do not provide a bound on the difference  $\|R_g - R_{S_\varepsilon g}\|$  or  $\|R_g - R_{g(t)}\|$ .

We are motivated by the viewpoint of moduli spaces of riemannian metrics with pinched curvature and regard Theorem B as a result on a controlled perturbation within such a space.

The paper is structured as follows: In §2 we will recall mollification, chart norms, and known uniform bounds on the regularity of the metric tensor under geometric conditions. The subsequent §3 gives a proof of the main technical tool, Theorem A, beginning with a local version of its statement. This is then used in §4 to derive the main result, Theorem B, and its corollaries.

## § 2. Review of mollification and chart norms

We will give a short introduction to Hölder spaces and mollification. In the subsequent two subsections we will explain norm bounds for a riemannian metric that are independent of a distinguished coordinate system, and state the fundamental examples and properties of such systems.

### § 2.1 Hölder spaces

Besides  $L^p$ -classes we will use Hölder spaces of functions  $f: \mathbb{R}^n \rightarrow \mathbb{R}^N$ . For  $m = 0, 1, \dots$  define

$$(2.1a) \quad \nabla^m f: \Omega \rightarrow \mathbb{R}^{N \cdot n^m}$$

to be the function of all derivatives of order  $m$  and

$$(2.1b) \quad \nabla^{\leq m} f: \Omega \rightarrow \mathbb{R}^{N \cdot n^0 + \dots + N \cdot n^m}$$

to be the collection of all derivatives of order 0 to order  $m$ . Further let

$$(2.2a) \quad |(x_1, \dots, x_n)| := \max\{|x_1|, \dots, |x_n|\}$$

for  $x = (x_1, \dots, x_n) \in \mathbb{R}^n$  (which is in contrast to the euclidean norm  $|0x|$ ). Recall that the Hölder norm—for  $\alpha \in [0, 1]$ ,  $m = 0, 1, \dots$ , and a domain  $\Omega \subset \mathbb{R}^n$ —is given by

$$(2.2b) \quad \|f\|_{C^{m,\alpha}} = \|f\|_{C^{m,\alpha}(\Omega)} := \|f\|_{C^m} + \sum_{k=0}^m \|\nabla^k f\|_\alpha$$

where  $\|f\|_\alpha = 0$  in case  $\alpha = 0$  and otherwise

$$(2.2c) \quad \|f\|_\alpha := \sup_{\substack{x, y \in \Omega \\ x \neq y}} \frac{|f(x) - f(y)|}{|x - y|^\alpha}.$$

Denote by  $C^{m,\alpha}(\Omega) = C^{m,\alpha}(\Omega, \mathbb{R})$  the corresponding spaces of real-valued functions on a domain  $\Omega \subset \mathbb{R}^n$ . If  $\Omega$  is a bounded open set with Lipschitz

boundary, Hölder spaces are connected to Sobolev spaces by Sobolev's inequality which states for  $k - n/p = r + \alpha$ ,  $p \in (n, \infty]$ ,  $r < k$ ,  $\alpha \in (0, 1]$  that

$$(2.S) \quad \|f\|_{C^{r,\alpha}} \leq C \|f\|_{W^{k,p}}.$$

with  $C = C(n, p)$ . From  $\|-\|_\alpha \leq \|-\|_1$  and the mean value theorem we get the elementary estimate

$$(2.3) \quad \|f\|_{C^{m,\alpha}} \leq \|f\|_{C^{m+1}}.$$

## § 2.2 Mollification

The tool for regularization will be mollification, i.e. convolution with a smooth function. Convolution can be defined for any compactly supported function  $f: \mathbb{R}^n \rightarrow \mathbb{R}$  and any locally integrable function  $g: \mathbb{R}^n \rightarrow \mathbb{R}$  via

$$(2.4) \quad f * g(x) := \int f(x-h)g(h)dh.$$

If one of the functions  $f, g$  is multi-valued, i.e.  $\mathbb{R}^n \rightarrow \mathbb{R}^N$ , the convolution is defined component-wise. It is elementary to see that if in addition  $f \in C^m$ , then  $f * g \in C^k$  for  $m = 0, 1, \dots$  [Hör83, Theorem 1.3.1]. Moreover, in this case

$$(2.5) \quad \partial^I(f * g) = (\partial^I f) * g$$

where  $I$  is a multi-index of order  $m$ . We will always assume that  $p, q \in [1, \infty]$  satisfy the relation  $1 = 1/p + 1/q$ . The classical key tools will be Hölder's inequality

$$(2.H) \quad \|fg\|_{L^1} \leq \|f\|_{L^p} \|g\|_{L^q}$$

and Young's convolution inequality

$$(2.Y) \quad \|f * g\|_{L^\infty} \leq \|f\|_{L^p} \|g\|_{L^q}.$$

For the definition of mollification, fix as the mollification kernel a smooth function  $\varphi: \mathbb{R}^n \rightarrow [0, 1]$  supported on  $[-1, 1]^n$  with  $\varphi(0) = \int \varphi(x)dx = 1$ . Set for  $t > 0$

$$(2.6a) \quad \varphi_t(x) := t^{-n} \varphi(x/t).$$

By substitution we have that  $\int \varphi_t(x)dx = \int \varphi(x)dx = 1$ . Define the mollification operator by

$$(2.6b) \quad P_t: f \mapsto \varphi_t * f.$$

As a first application of Young's convolution inequality in conjunction with (2.5) note that

$$(2.6c) \quad \|\nabla^m P_t f\|_{L^\infty} \leq \|\nabla^m \varphi_t\|_{L^1} \|f\|_{L^\infty}.$$

### § 2.3 Definition of chart Norms

Hölder classes of riemannian metrics allow to formulate celebrated regularity results in a more concise and little bit stronger fashion. Let  $(M, g, p_0)$  be a pointed  $n$ -dimensional riemannian manifold. We introduce norms on charts, given by pointed maps

$$\psi: (B(0, r), 0) \rightarrow (M, p_0),$$

i.e.  $\psi(0) = p_0$ , where  $B(x, r)$  denotes the open ball of radius  $r$  around  $x$  in the metric space to which  $x$  belongs—here  $0$  belongs to  $\mathbb{R}^n$  with the euclidean metric. In contrast, we will denote a closed ball by  $B[x, r]$ . We will mainly use and adapt definitions from [Pet16, §§ 11.3.1-11.3.5].

**Definition 2.1.** For a chart  $\psi: (B(0, r), 0) \rightarrow (M, g, p_0)$  compatible with the smooth atlas of  $M$  we define  $\|\psi\|_{C^{m,\alpha}}$ , the **chart norm of  $\psi$  on the scale of  $r$**  as the minimal\* quantity  $Q \geq 0$  for which the following conditions are fulfilled

- (1) for the differentials we have the bounds  $|D\psi| \leq e^Q$  on  $B(0, r)$  and  $|D\psi^{-1}| \leq e^Q$  on  $\psi(B(0, r))$ . Equivalently, this condition can be expressed in coordinates on  $\psi$  by

$$(2.N_0) \quad e^{-2Q} \delta_{kl} v^k v^l \leq g_{kl} \leq e^{2Q} \delta_{kl} v^k v^l$$

for every vector  $v \in \mathbb{R}$ .

- (2) for the semi-norm from (2.2c) and any  $k = 0, 1, \dots, m$

$$(2.N_{C^{m,\alpha}}) \quad r^{k+\alpha} \|\nabla^k g_{..}\|_\alpha \leq Q$$

where  $g_{..} := \psi^* g$ .

In this case we write

$$\|\psi\|_{C^{m,\alpha},r} \leq Q.$$

We define  $\|\psi\|_{C^{m,\alpha}}^{\text{harm}}$ , the **harmonic chart norm of  $\psi$  on the scale of  $r$** , by additionally imposing the condition

- (3) the chart  $\psi$  is harmonic, meaning that each coordinate function  $x_k$  ( $k = 1, \dots, n$ ) is harmonic with respect to  $g_{..}$ , i.e. the Laplace-Beltrami operator vanishes

$$(2.N_{\text{harm}}) \quad \Delta_{g_{..}} x_k = 0.$$

In this case we write

$$\|\psi\|_{C^{m,\alpha},r}^{\text{harm}} \leq Q.$$

We can directly extend this definition by

$$\begin{aligned} \|(M, g, p_0)\|_{C^{m,\alpha},r} &= \inf_{\psi: (B(0,r),0) \rightarrow (M,p_0)} \|\psi\|_{C^{m,\alpha}}, \\ \|(M, g)\|_{C^{m,\alpha},r} &= \sup_{p_0 \in M} \|(M, g, p_0)\|_{C^{m,\alpha},r} \end{aligned}$$

---

\*Note that there are only finitely many defining conditions.

(mutatis mutandis for  $\|(M, g, p_0)\|_{C^{m,\alpha},r}^{\text{harm}}$  and  $\|(M, g)\|_{C^{m,\alpha},r}^{\text{harm}}$ ). In the same manner we can introduce the norm bounds on Sobolev scales

$$\|\psi\|_{W^{m,p},r} \leq Q, \quad \|(M, g, p_0)\|_{W^{m,p},r} \leq Q, \quad \|(M, g)\|_{W^{m,p},r} \leq Q$$

$$\left( \|\psi\|_{W^{m,p},r}^{\text{harm}} \leq Q, \quad \|(M, g, p_0)\|_{W^{m,p},r}^{\text{harm}} \leq Q, \quad \|(M, g)\|_{W^{m,p},r}^{\text{harm}} \leq Q, \quad \text{resp.} \right)$$

by retaining condition (2.N<sub>0</sub>) (as well as (2.N<sub>harm</sub>) if appropriate) and replacing condition (2.N<sub>C<sup>m,α</sup></sub>) by

$$(2.N_{W^{m,p}}) \quad r^{k-n/p} \|\nabla^k g\|_{L^p} \leq Q.$$

for  $k = 0, 1, \dots, m$ . Note that  $\|(M, g, p_0)\|_{C^{m,\alpha},r}$  and  $\|(M, g, p_0)\|_{C^{m,\alpha},r}^{\text{harm}}$  are realized by charts by an application of the Arzelà–Ascoli theorem. (The same holds for  $\|(M, g, p_0)\|_{W^{m,p},r}$  and  $\|(M, g, p_0)\|_{W^{m,p},r}^{\text{harm}}$  as a consequence of the Banach–Alaoglu theorem.) Finally, let

$$\mathcal{M}^n(C^{m,\alpha} \leq_r Q), \quad \mathcal{M}^n(W^{m,p} \leq_r Q)$$

$$\left( \mathcal{M}^n(C^{m,\alpha} \leq_r^{\text{harm}} Q), \quad \mathcal{M}^n(W^{m,p} \leq_r^{\text{harm}} Q), \quad \text{resp.} \right)$$

denote the space of isomorphism class of  $n$ -dimensional, pointed, smooth riemannian manifolds  $(M, g, p_0)$  with

$$\|(M, g)\|_{C^{m,\alpha},r} \leq Q, \quad \|(M, g)\|_{W^{m,p},r} \leq Q, \quad \text{resp.}$$

$$\left( \|(M, g)\|_{C^{m,\alpha},r}^{\text{harm}} \leq Q, \quad \|(M, g)\|_{W^{m,p},r}^{\text{harm}} \leq Q, \quad \text{resp.} \right).$$

These spaces are endowed with the Gromov–Hausdorff topology. Note the elementary estimate [Pet16, Proposition 11.3.2 (4)]

$$(2.D) \quad (2.N_0) \implies$$

$$e^{-Q} \min\{|x y|, 2r - |0 x|\} \leq |\psi(x) \psi(y)|_g \leq e^Q |x y| \leq e^Q |x y|$$

for all  $x, y \in B(0, r)$  and  $|\cdot|$  the euclidean norm.

Having introduced spaces with a global bound on the metric tensor in local coordinates, one may be inclined to ask why we did not assume any regularity assumption on changes of coordinates. The answer is found in Schauder estimates, standard estimates on the regularity of solutions of elliptic PDEs. The crucial fact can be stated as follows [GT15, Problem 6.1 (a)]: On a bounded open set  $\Omega$  let  $u: \Omega \rightarrow \mathbb{R}$  be a  $C^{m+2,\alpha}$ -solution ( $m \geq 0$ ) of  $(a^{ij}(x)\partial_i\partial_j + b^i(x)\partial_i + c(x))u = f$  (summation convention) and assume that the coefficients of  $L$  satisfy  $a^{ij}\xi_i\xi_j \geq \lambda|\xi|^2$  and  $\|\nabla^m a\|_\alpha, \|\nabla^m b\|_\alpha, \|\nabla^m c\|_\alpha \leq \Lambda$ . If  $\Omega' \subset \Omega$  with  $\overline{\Omega'} \subsetneq \Omega$ , then

$$\|u\|_{C^{m+2,\alpha}} \leq C(\|u\|_{C^0} + \|f\|_{C^{m,\alpha}})$$

on  $\Omega'$  with  $C = C(n, m, \alpha, \lambda, \Lambda, |\Omega' \partial \Omega|_{\text{H}})$  where  $|\Omega' \partial \Omega|_{\text{H}}$  denotes the Hausdorff distance.

If we apply this statement to a transition function  $\psi_i^{-1} \circ \psi_j$  for two charts  $\psi_i, \psi_j$  with  $\|\psi_i\|_{C^{m,\alpha},r}^{\text{harm}}, \|\psi_j\|_{C^{m,\alpha},r}^{\text{harm}} \leq Q$  we first notice that by harmonicity  $\|\Delta_g \psi_i^{-1} \circ \psi_j\|_{C^{m,\alpha}} = \|0\|_{C^{m,\alpha}} = 0$  and moreover, in harmonic coordinates,

as calculated e.g. in [Pet16, § 11.2.3],  $\Delta_g = g^{ij}\partial_i\partial_j$ . If we restrict to the domain  $\Omega' := \psi_j^{-1}(\psi_i(B(0, r/2)) \cap \psi_j(B(0, r/2)))$  the above estimate becomes

$$(2.Sch) \quad \|\psi_i^{-1} \circ \psi_j\|_{C^{m+2,\alpha}} \leq C \|\psi_i^{-1} \circ \psi_j\|_{C^0} \leq C \cdot r/2$$

where  $C = C(n, m, \alpha, Q, r)$ . For non-harmonic chart norms a similar result with one lower degree of regularity can be found in Taylor [Tay06].

## § 2.4 Precompactness theorems for riemannian manifolds

We review some fundamental results. For every  $n \geq 2$  and  $s, Q > 0$

$$(2.K_{C^{m,\alpha}}) \quad \mathcal{M}^n(C^{m,\alpha} \leq_r Q) \text{ is compact}$$

with respect to the Gromov-Hausdorff topology. This statement is sometimes called Fundamental Theorem of Convergence Theory [Pet16, § 11.3.5].

Let  $\iota > 0$ ,  $\alpha \in (0, 1)$ ,  $p \in (n, \infty)$ , and  $\underline{\kappa} \in \mathbb{R}$ . For all  $Q > 0$  there is  $r > 0$  such that every pointed riemannian manifold  $(M, g, p_0)$  satisfies

$$(2.K_{Ric}) \quad \text{injectivity radius}(M) \geq \iota \text{ and } Ric \geq -\underline{\kappa}^2 \\ \implies (M, g, p_0) \in \mathcal{M}^n(C^\alpha \leq_r^{\text{harm}} Q) \cap \mathcal{M}^n(W^{1,p} \leq_r^{\text{harm}} Q);$$

see [AC92]. Likewise, we obtain a result that is stronger by one derivative for an absolute Ricci bound: For every  $\kappa \geq 0$ ,  $\iota > 0$ ,  $\alpha \in (0, 1)$ ,  $p \in (n, \infty)$ , and  $Q \in (0, \log(2))$  there is an  $r > 0$  such that any smooth manifold  $(M, g, p_0)$

$$(2.K_{|Ric|}) \quad \text{injectivity radius}(M) \geq \iota \text{ and } \|Ric\|_{L^\infty} \leq \kappa \\ \implies (M, g, p_0) \in \mathcal{M}^n(C^{1,\alpha} \leq_r^{\text{harm}} Q) \cap \mathcal{M}^n(W^{2,p} \leq_r^{\text{harm}} Q);$$

the result appeared several times, e.g. [And90] and [HH97, Theorem 11], and found its ways into textbooks [Pet16, § 11.4.1]. Note that the Hölder parts of (2.K<sub>Ric</sub>) and (2.K<sub>|Ric|</sub>) are implied by their Sobolev parts via Sobolev's inequality (2.S).

*Remark 2.2.* Actually, these regularity results can be improved in at least two ways that we do not use but would like to emphasize. The exponent  $\alpha$  can be improved from  $\alpha \in (0, 1)$  to 1 by replacing the Hölder scales with Hölder-Zygmund scales  $C_*^s$ ,  $s > 0$ . The space  $C_*^s(\Omega)$  coincides with  $C^{m,\alpha}$  if  $s = m + \alpha$  and  $\alpha \neq 0, 1$ . Otherwise  $C^m \subset C_*^m(\omega)$ . See [Tay07] for the case of lower Ricci bounds and [And+04] for the case of absolute Ricci bounds. The norm bound in case of an absolute bound on Ricci curvature can be generalized to the case of manifolds with boundary [And+04].

## § 3. Mollified riemannian curvature tensor

In this section we begin by proving a local version of Theorem A. Recall that  $g_\cdot = \psi^*g$  and  $P_t(g_\cdot) = \varphi_t * g_\cdot$ . The results will be formulated using the Sobolev chart norm  $\|-\|_{W^{2,p},r}$  from (2.N <sub>$W^{m,p}$</sub> ).



**Proposition 3.1.** *Let  $p > 2n$ ,  $r, Q > 0$  and choose  $\beta \in (1/2, 1 - n/p)$ . There exists some  $T \in (0, r/2)$  such that for a pointed smooth riemannian manifold  $(M, g, p_0)$ , a chart  $\psi: (B(0, r), 0) \rightarrow (M, p_0)$ , and any section  $\vec{v} \in \Gamma T^{3,1} B(0, r)$  we have*

$$\|\psi\|_{W^{2,p},r} \leq Q \implies R_{P_t(g_{\cdot})}(\psi(x))(\vec{v}) - \sup_{y \in B(x,t)} R_{g_{\cdot}}(y)(\vec{v}) \leq C \|\vec{v}\|_{L^\infty}^4 t^\beta$$

for any  $t \in (0, T]$ ,  $x \in B(0, r/2)$ , a constant  $C = C(n, p, r, Q, \beta)$ , and considering  $\vec{v}$  as a function valued in  $\mathbb{R}^{4n}$  by the canonical euclidean identification.

The following generalization of the above proposition to a convex combination of mollified metrics will be of even greater interest. Recall that the  $g$ -norm of a section  $v$  in a tangent bundle is defined by  $\|v\|_g := \sup_{x \in M} \|v(x)\|_{g(x)}$ . For a cotangent vector  $v_x \in T_x^* M$  the  $g$ -norm  $\|v_x\|_{g(x)}$  is defined as  $\|(w \mapsto \langle w, - \rangle)^{-1}(v_x)\|_{g(x)}$ . The  $g$ -norm of a section  $v \in \Gamma T^* M$  is  $\|v\|_g := \sup_{x \in M} \|v(x)\|_{g(x)}$ . Finally, for a section in a product of bundles  $\vec{v} = (v_1, \dots, v_k, v_{k+1}, \dots, v_{k+l}) \in \Gamma(TM \times \dots \times TM \times T^*M \times \dots \times T^*M)$  we define  $\|\vec{v}\|_g = \sup_{x \in M} \|v_1(x)\|_{g(x)} \cdots \|v_k(x)\|_{g(x)} \cdot \|v_{k+1}(x)\|_{g(x)} \cdots \|v_{k+l}(x)\|_{g(x)}$ .

**Definition 3.2.** Let  $r > 0$  and  $M$  be a smooth manifold, which is covered by a locally finite family of charts  $\{\psi_i\}_I$  with a **corresponding partition of unity**  $\{\varrho_i\}_I$ , i.e.

$$(3.1a) \quad \{\psi_i: B(0, r) \rightarrow M\}_{i \in I}, \quad \{\varrho_i: M \rightarrow [0, 1]\}_{i \in I}$$

such that for each  $i \in I$  the function  $\varrho_i$  is smooth and

$$(3.1b) \quad \text{supp } \varrho_i \subset \psi_i(B(0, 3r/4)), \quad \sum_{j \in I} \varrho_j \Big|_{\bigcup_{j \in I} \psi_j(B(0, r/2))} = 1.$$

Define for  $t \in (0, r/2)$  the **mollified metrics**

$$(3.1c) \quad g^{[t]} := P_{t, \{\psi_i\}_I, \{\varrho_i\}_I}(g) := \sum_{i \in I} \varrho_i g^{[t, \psi_i]}$$

on  $\bigcup_{j \in I} \psi_j(B(0, r/2))$  where

$$(3.1d) \quad g^{[t, \psi_i]} := (\psi_i^{-1})^* P_t(\psi_i^* g).$$

**Theorem A.** *Given  $p > 2n$  and  $r, Q > 0$ . Choose  $\beta \in (0, 1 - 2n/p)$ . Then there is some  $T \in (0, r/2)$  such that for any pointed smooth riemannian manifold  $(M, g, p_0)$  and any finite collection of charts with a corresponding partition of unity*

$$\{\psi_i: (B(0, r), 0) \rightarrow (M, \psi_i(0))\}_{i \in I}, \quad \vec{\varrho} := \{\varrho_i: M \rightarrow [0, 1]\}_{i \in I}$$

we have that for any section  $\vec{v} \in (T\psi(B(0, r)))^{\times 3} \times T^*\psi(B(0, r))$

$$\left( \forall i \in I: \|\psi_i\|_{W^{2,p},r}^{\text{harm}} \leq Q \right) \implies R_{g^{[t]}}(x)(\vec{v}) - \sup_{B(x, e^Q t)} R_g(\vec{v}) \leq C \|\vec{v}\|_g t^\beta$$

for any  $t \in (0, T]$ ,  $x \in \bigcap_{i \in I} \psi_i(B(0, r/2))$  and with

$$C = C\left(n, p, r, Q, \beta, \{\|\varrho_i \circ \psi_i\|_{C^2}\}_{i \in I}, \#I\right).$$

Throughout the proof, whose ansatz was pointed out in (1.1), we will consider a metric tensor  $g_{\cdot\cdot} = \psi^*g$  with

$$(3.2a) \quad \|\psi\|_{W^{2,p},r}, \|\psi\|_{C^{1,\alpha},r} \leq Q$$

such that

$$(3.2b) \quad \frac{1}{p} + \frac{1}{q} = 1, \quad n < p < \infty, \quad \frac{n}{p} < \alpha < 1, \quad \text{and} \quad \beta \leq \alpha - n/p.$$

These conditions imply  $p, q \in (0, \infty)$  and  $\alpha \in (0, 1)$ . To guarantee this condition without loss of generality, choose  $\alpha \in (1 - 2n/p, 1 - n/p)$ , apply Sobolev's inequality (2.S), and replace  $Q$  by  $\max\{Q, CQ\}$ , where  $C$  is the constant from Sobolev's inequality. By  $g^{\cdot\cdot}$  we denote the inverse of  $g_{\cdot\cdot}$ .

**Lemma 3.3.** *In local coordinates the riemannian curvature tensor  $R$  of a smooth manifold  $(M, g)$  can be written using notation (2.1b) as*

$$R^{\cdot\cdot\cdot\cdot} = A(g^{\cdot\cdot}, \nabla^2 g_{\cdot\cdot}) + B(\nabla g_{\cdot\cdot}, \nabla^{\leq 1} g^{\cdot\cdot})$$

where

- $A$  is bilinear;
- $B$  is a polynomial of degree not bigger than 4 and without constant terms; and
- the coefficients of  $A$  and  $B$  depend only on the dimension.

*Proof.* The standard coordinate definition the riemannian curvature tensor has the form [Tay13, § C3]

$$\begin{aligned} \Gamma_{kl}^i &= \frac{1}{2} g^{im} (g_{mk,l} + g_{ml,k} - g_{kl,m}) \\ R_{\sigma\mu\nu}^{\rho} &= \partial_{\mu} \Gamma_{\nu\sigma}^{\rho} - \partial_{\nu} \Gamma_{\mu\sigma}^{\rho} + \Gamma_{\mu\lambda}^{\rho} \Gamma_{\nu\sigma}^{\lambda} - \Gamma_{\nu\lambda}^{\rho} \Gamma_{\mu\sigma}^{\lambda} \end{aligned}$$

where  $f_{i,j} = \partial_j f_i = \frac{\partial}{\partial x_j} f_i$ . From this we read that

$$R^{\cdot\cdot\cdot\cdot} = L_2(g^{\cdot\cdot}, \nabla^2 g_{\cdot\cdot}) + L_1(\nabla g^{\cdot\cdot}, g_{\cdot\cdot}) + Q_1(g^{\cdot\cdot}, \nabla g_{\cdot\cdot})$$

where the coefficients of  $L_2$ ,  $L_1$ , and  $Q_1$  depend only on the dimension and

- $L_2$  is linear in  $\nabla^2 g_{\cdot\cdot}$  and  $g^{\cdot\cdot}$ ,
- $L_1$  is linear in  $\nabla g^{\cdot\cdot}$  as well as in  $\nabla g_{\cdot\cdot}$ ,
- $Q_1$  is quadratic in both  $g^{\cdot\cdot}$  and  $\nabla g_{\cdot\cdot}$ .

Obviously, choose  $A = L_2$  and  $B = L_1 + Q_1$ . □

**Lemma 3.4.** *Let  $f, g \in L^{\infty}(\Omega, \mathbb{R}^N)$  and  $p \in \mathbb{R}[X_1, \dots, X_N]$  a polynomial without constant term. Then*

$$\|p(f) - p(g)\|_{L^{\infty}} \leq C \|f - g\|_{L^{\infty}}$$

where  $C = C(p, \|f\|_{L^{\infty}}, \|g\|_{L^{\infty}})$ .

*Proof.* By the triangle inequality, it is sufficient to prove the claim for a monomial  $p = X_{i_1} \cdots X_{i_d}$ . Then the claim follows from a telescope argument

$$\begin{aligned} & \|f_{i_1} \cdots f_{i_d} - g_{i_1} \cdots g_{i_d}\|_{L^\infty} \\ & \leq \|(f_{i_1} - g_{i_1})f_{i_2} \cdots f_{i_d}\|_{L^\infty} + \|g_{i_1}(f_{i_2} - g_{i_2}) \cdots f_{i_d}\|_{L^\infty} + \cdots \\ & \quad \cdots + \|g_{i_1} \cdots (f_{i_d} - g_{i_d})\|_{L^\infty} \end{aligned}$$

since each term  $g_{i_1} \cdots (f_{i_k} - g_{i_k}) \cdots f_{i_d}$  is  $L^\infty$ -bounded by  $\|f\|_{L^\infty}^{k-1} \cdot \|g\|_{L^\infty}^{n-k-1} \cdot \|f - g\|_{L^\infty}$  (for  $k = 1, \dots, d$ ).  $\square$

We now turn to the crucial  $L^p$ -estimate. It is a consequence of Hölder's inequality (2.H) for indices 1 and  $\infty$  stating that for any  $q \in (1, \infty)$

$$\begin{aligned} \|\varphi_t\|_{L^q} &= \|t^{-nq} \chi_{[-t,t] \times \mathbb{R}^n} (\varphi(h/t))^q\|_{L^1}^{1/q} \\ &\leq \|t^{-nq} \chi_{[-t,t] \times \mathbb{R}^n}\|_{L^1}^{1/q} \|(\varphi(h/t))^q\|_{L^\infty}^{1/q} \\ &= t^{-n} \|\chi_{[-t,t] \times \mathbb{R}^n}\|_{L^1}^{1/p} \|(\varphi(h/t))^q\|_{L^\infty}^{1/q} \\ &= t^{-n} (2t)^{n \cdot 1/q} \cdot 1^{1/q} \\ &= 2^{n/q} t^{-n(1-1/q)} \\ (3.3) \quad &= 2^{n/q} t^{-n/p}. \end{aligned}$$

Recall the definitions of the commutator of two operators  $P, Q$

$$[P, Q]: f \mapsto PQ - QP$$

and the multiplication operator

$$T_g: f \mapsto gf.$$

Finally, note that condition (2.N<sub>0</sub>) implies that the eigenvalues of  $g_{\cdot}$  are in  $[e^{-2Q}, e^{2Q}]$ . This implies

$$(3.4) \quad \det(g_{\cdot}) \in [e^{-2Qn}, e^{2Qn}].$$

**Lemma 3.5** (Commutator-like estimate). *Let  $n, p, q, \alpha$  satisfy (3.2b) and  $\Omega \subset \mathbb{R}^n$  be an open domain. Let  $f \in L^p(\mathbb{R}^n)$  and  $a_{(-)}(-): \Omega \times [0, T] \rightarrow \mathbb{R}$  such that for some constant  $C_a$  we have*

$$\|a_t\|_\alpha \leq C_a \quad \text{and} \quad \|a_t - a_0\|_{C^0} \leq C_a t^\alpha$$

for all  $t \in [0, T]$ . Then

$$\|P_t T_{a_0}(f) - T_{a_t} P_t(f)\|_{L^\infty} \leq 2^{(n+1)/q} C_a \|f\|_{L^p} t^{\alpha-n/p}.$$

*Proof.* First note that by Young's convolution inequality (2.Y) and (3.3)

$$\begin{aligned}
\| [P_t, T_g]f \|_{L^\infty} &= \operatorname{ess\,sup}_x \left| \int (g(x-h) - g(x)) f(x-h) \varphi_t(h) dh \right| \\
&\leq \operatorname{ess\,sup}_x \int |g(x-h) - g(x)| \cdot |f(x-h)| \varphi_t(h) dh \\
&\leq \operatorname{ess\,sup}_x \int \|g\|_\alpha |h|^\alpha |f(x-h)| \varphi_t(h) dh \\
&\leq \|g\|_\alpha t^\alpha \|P_t(|f|)\|_{L^\infty} \\
&= \|g\|_\alpha t^\alpha \|f\|_{L^p} \|\varphi_t\|_{L^q} \\
&\leq 2^{n/q} t^{\alpha-n/p} \|g\|_\alpha \|f\|_{L^p}.
\end{aligned}$$

Using this estimate as well as (3.3) and (2.Y) again we conclude the proof:

$$\begin{aligned}
&\|P_t T_{a_0}(f) - T_{a_t} P_t(f)\|_{L^\infty} \\
&\leq \| [P_t, T_{a_t}]f \|_{L^\infty} + \|P_t T_{a_0}(f) - P_t T_{a_t} f\|_{L^\infty} \\
&= \| [P_t, T_{a_t}]f \|_{L^\infty} + \|P_t T_{a_0-a_t}(f)\|_{L^\infty} \\
&\leq \| [P_t, T_{a_t}]f \|_{L^\infty} + \|\varphi_t\|_{L^q} \|T_{a_0-a_t}(f)\|_{L^p} \\
&\leq 2^{n/q} t^{\alpha-n/p} \|a_t\|_\alpha \|f\|_{L^p} + 2^{n/q} t^{-n/p} \|a_0 - a_t\|_{C^0} \|f\|_{L^p} \\
&\leq C_a t^\alpha \|f\|_{L^p} 2^{n/q} t^{-n/p} + C_a t^\alpha \|f\|_{L^p} 2^{n/q} t^{-n/p} \\
&= 2^{(n+1)/q} C_a \|f\|_{L^p} t^{\alpha-n/p}.
\end{aligned}$$

□

**Lemma 3.6.** *Let  $r > 0$ ,  $m = 0, 1, \dots$ , and  $\alpha \in (0, 1]$ . Given a function  $f \in C^{m,\alpha}(B(0, r), \mathbb{R})$  we have*

$$\|f - P_t f\|_{C^m} \leq t^\alpha \|f\|_{C^{m,\alpha}}$$

on  $B(0, r/2)$  for  $t \in (0, r/2)$ .

*Proof.* Let  $k = 0, \dots, m$ . The lemma follows from

$$\begin{aligned}
\|\nabla^k(P_t(f) - f)\|_{C^0} &= \|P_t(\nabla^k f) - \nabla^k f\|_{C^0} \\
&= \sup_{x \in B(0, r/2)} \left| P_t(\nabla^k f) - \int \varphi_t(h) \nabla^k f(x) dh \right| \\
&= \sup_{x \in B(0, r/2)} \left| \int \varphi_t(h) (\nabla^k f(x-h) - \nabla^k f(x)) dh \right| \\
&\leq \sup_{x \in B(0, r/2)} \int |\varphi_t(h)| \cdot |h|^\alpha \|\nabla^k f\|_\alpha dh \\
&\leq \|\nabla^k f\|_\alpha \int |\varphi_t(h)| t^\alpha dh \\
&= t^\alpha \|\nabla^k f\|_\alpha.
\end{aligned}$$

□

*Proof of Proposition 3.1.* As discussed above at (3.2a) we can assume without loss of generality  $\|\psi\|_{W^{2,p,r}}, \|\psi\|_{C^{1,\alpha,r}} \leq Q$  with properties (3.2b). By assumption (2.N $_{C^{m,\alpha}}$ ) and Lemma 3.6 this implies

$$(3.5a) \quad \|P_t(g_{..}) - g_{..}\|_{C^1} \leq C_{n,r} Q t^\alpha$$

for  $t \in (0, r/2)$  and for a dimension- and radius-depending constant  $C_{n,r}$ . This in conjunction with the bounds (3.4) implies

$$(3.5b) \quad \det(g_{\cdot\cdot}), \det(P_t(g_{\cdot\cdot})) \in [e^{-2Qn-1}, e^{2Qn+1}]$$

for all  $t \in (0, T]$  when  $T$  is chosen sufficiently small. This implies that

$$(3.5c) \quad \|P_t(g_{\cdot\cdot})^{-1}\|_{C^{1,\alpha}}, \|g_{\cdot\cdot}\|_{C^{1,\alpha}} \leq C_{Q,r}$$

for a constant  $C_{Q,r}$  [CDK11, Corollary 16.30]. Let  $M_{kl}$  denote the  $(k, l)$ -minor of  $g_{\cdot\cdot}$ ,  $M_{kl}^t$  the  $(k, l)$ -minor of  $P_t(g_{\cdot\cdot})$ , and  $M_{\cdot\cdot}, M_{\cdot\cdot}^t$  the corresponding matrices of minors. Lastly, we need an estimate on the difference of the inverse of the mollified metric:

$$\begin{aligned} & \| (P_t(g_{\cdot\cdot}))^{-1} - g_{\cdot\cdot}^{-1} \|_{C^1} \\ & \leq \left\| \frac{1}{\det P_t(g_{\cdot\cdot})} M_{\cdot\cdot}^t - \frac{1}{\det(g_{\cdot\cdot})} M_{\cdot\cdot} \right\|_{C^1} \\ & \leq \left| \frac{1}{\det P_t(g_{\cdot\cdot})} - \frac{1}{\det(g_{\cdot\cdot})} \right| \|M_{\cdot\cdot}^t\|_{C^1} + \frac{1}{\det(g_{\cdot\cdot})} \|M_{\cdot\cdot}^t - M_{\cdot\cdot}\|_{C^1} \end{aligned}$$

proceeding by using (3.5b) twice—directly and by the standard estimate

$$\left| \frac{1}{\xi} - \frac{1}{\eta} \right| \leq \left| \left( \frac{d}{dx} \frac{1}{x} \right) (x_0) \right| |\xi - \eta| = |x_0|^{-2} |\xi - \eta| \text{ for } \xi, \eta \geq x_0$$

$$\begin{aligned} & \leq \frac{|\det P_t(g_{\cdot\cdot}) - \det(g_{\cdot\cdot})|}{(e^{-2Qn-1})^2} \|M_{\cdot\cdot}^t\|_{C^1} + e^{2Qn+1} \|M_{\cdot\cdot}^t - M_{\cdot\cdot}\|_{C^1} \\ & \leq e^{4Qn+2} \cdot C_{n,r} \cdot t^\alpha Q \cdot C_n + e^{2Qn+1} \cdot C'_{n,r} \cdot t^\alpha Q \end{aligned}$$

$$(3.5d) \quad \leq C_{n,r,\alpha,Q} t^\alpha.$$

where in the penultimate step we used Lemma 3.6 and (3.5a) as well as standard product estimates [CDK11, Theorem 16.28] providing suitable constants  $C_n$  and  $C'_{n,r}$ .

Finally, we can apply the ansatz proposed in (1.1):

$$(3.6) \quad \begin{aligned} R_{P_t(g_{\cdot\cdot})}(x)(\vec{v}) & \leq P_t(R_{\cdot\cdot\cdot})(x)(\vec{v}) + \left| P_t(R_{\cdot\cdot\cdot})(x)(\vec{v}) - R_{P_t(g_{\cdot\cdot})}(x)(\vec{v}) \right| \\ & \leq \sup_{y \in B(x,t)} R_{g_{\cdot\cdot}}(y)(\vec{v}) + \|P_t(R_{\cdot\cdot\cdot}) - R_{P_t(g_{\cdot\cdot})}\|_{L^\infty} \|\vec{v}\|_{L^\infty}^4. \end{aligned}$$

It remains to show that  $\|P_t(R_{\cdot\cdot\cdot}) - R_{P_t(g_{\cdot\cdot})}\|_{L^\infty}$  vanishes with modulus  $t^\alpha$  as  $t \rightarrow 0$ . Set  $b := B(\nabla g_{\cdot\cdot}, \nabla^{\leq 1} g_{\cdot\cdot})$  and  $b_t := B(\nabla P_t g_{\cdot\cdot}, \nabla^{\leq 1} (P_t g_{\cdot\cdot})^{-1})$ . Observe

$$\begin{aligned} & \|P_t(R_{\cdot\cdot\cdot}) - R_{P_t(g_{\cdot\cdot})}\|_{L^\infty} \\ & \leq \|P_t A(g_{\cdot\cdot}, \nabla^2 g_{\cdot\cdot}) - A((P_t g_{\cdot\cdot})^{-1}, \nabla^2 P_t g_{\cdot\cdot})\|_{L^\infty} + \|P_t(b) - b_t\|_{L^\infty} \\ & \leq \|P_t A(g_{\cdot\cdot}, \nabla^2 g_{\cdot\cdot}) - A((P_t g_{\cdot\cdot})^{-1}, P_t \nabla^2 g_{\cdot\cdot})\|_{L^\infty} \\ & \quad + \|P_t(b) - b\|_{L^\infty} + \|b - b_t\|_{L^\infty} \end{aligned}$$

We will give estimates for each summand:

- For the first summand express  $A$  as  $A(x, y) = \sum_{i,j} a_{ij} x_i y_j$ . By linearity of convolution  $P_t(A(x, y)) = \sum_{i,j} a_{ij} P_t T_{x_i}(y_j)$ . On the other hand  $A(x, P_t y) = \sum_{i,j} a_{ij} T_{x_i} P_t(y_j)$ . This is to say that the difference of

both expressions is  $\sum_{i,j} a_{ij}[P_t, T_{x_i}](y_j)$ . Thereby, taking  $x_{(-)} = g^\cdot$  and  $y_{(\cdot)} = \nabla^2 g_\cdot$ , we are in situation of Lemma 3.5—providing the desired estimate.

- For the second summand Lemma 3.6 gives the desired estimate.
- In case of the third summand Lemma 3.4 is applicable due to (3.5c) and  $(2.N_{C^m, \alpha})$  providing the estimate

$$\|b_{\vec{v}} - b_{\vec{v}, t}\|_{L^\infty} \leq C' \left\| \left( \nabla g_\cdot, \nabla^{\leq 1} g^\cdot \right) - \left( \nabla P_t g_\cdot, \nabla^{\leq 1} (P_t g_\cdot)^{-1} \right) \right\|_{L^\infty}$$

where  $C' = C'(B, \|(\nabla g_\cdot, \nabla^{\leq 1} g^\cdot, \nabla P_t g_\cdot, \nabla^{\leq 1} (P_t g_\cdot)^{-1})\|_{L^\infty})$ . The estimate we seek follows now from (3.5a) and (3.5d).

Thus (3.6) becomes  $R_{P_t(g_\cdot)}(x)(\vec{v}) \leq \sup_{y \in B(x, t)} R_{g_\cdot}(y)(\vec{v}) + Ct^\beta$  for a constant  $C$  and  $\beta \leq \alpha - n/p$  as in the claim of the theorem.  $\square$

*Proof of Theorem A.* For the proof we examine the metric tensor on some chart  $\psi$  with  $\|\psi\|_{W^{2,p,r}}^{\text{harm}} \leq Q$ , e.g.  $\psi = \psi_i$  for one  $i \in I$ . For any metric tensor  $\tilde{g}$  on  $M$  representation with respect to  $\psi$ -coordinates is indicated by  $\tilde{g}_\cdot$ —or  $\tilde{g}^\cdot$  for the inverse, e.g.  $g^{[t]\cdot} = (g^{[t]})^{-1}$ . All sums  $\sum_i$  range over the entire index set  $I$ . By abuse of notation we write  $\varrho_i = \varrho_i \circ \psi$ . We define

$$(3.7a) \quad g^{\Delta t, \psi_i} := g - g^{[t, \psi_i]}.$$

Further we agree on the shorthands

$$(3.7b) \quad A_{\vec{v}}(x, y) := A(x, y)(\vec{v})$$

$$(3.7c) \quad b_{\vec{v}}^{[t]} := B(\nabla g^{[t]\cdot}, \nabla^{\leq 1} g^{[t]\cdot})(\vec{v})$$

$$(3.7d) \quad b_{\vec{v}}^{[t, \psi_i]} := B(\nabla g^{[t, \psi_i]\cdot}, \nabla^{\leq 1} g^{[t, \psi_i]\cdot})(\vec{v})$$

$$(3.7e) \quad C_\varrho := \sum_i \|\varrho_i \circ \psi_i\|_{C^2}$$

$$(3.7f) \quad \bar{v} := \left\| \begin{pmatrix} v^{\psi_{11}} v^{\psi_{21}} v^{\psi_{31}} v^{\psi_{41}} \dots, v^{\psi_{1n}} v^{\psi_{2n}} v^{\psi_{3n}} v^{\psi_{4n}} \\ \dots, v^{\psi_{1n}} v^{\psi_{2n}} v^{\psi_{3n}} v^{\psi_{4n}} \end{pmatrix} \right\|_{L^\infty}.$$

where  $v^\psi := \psi^* v$ .

Observe that due to (2.Sch)  $|\text{supp}(\varrho_i) \cap \partial(\psi^{-1} \circ \psi_i(B(0, r)))|_H \leq e^{-2Qr/4}$  for all  $i \in I$ . By (2.Sch) the estimates (3.5c) and (3.5d) imply

$$(3.8a) \quad \|g^{[t, \psi]\cdot}|_{\text{supp } \varrho_i}\|_{C^{1, \alpha}} \leq C_{n, Q, r, \alpha}, \quad \|g^{[t]\cdot}\|_{C^{1, \alpha}} \leq C_{n, Q, r, \alpha, C_\varrho},$$

$$(3.8b) \quad \|(g^{\Delta t, \psi_i})^\cdot|_{\text{supp } \varrho_i}\|_{C^1} \leq C'_{n, r, \alpha, Q} t^\alpha, \quad \|(g^{\Delta t})^\cdot\|_{C^1} \leq C'_{n, r, \alpha, Q, C_\varrho} t^\alpha$$

for every  $i \in I$  and respective constants. Moreover, by (2.Sch) we have the estimate

$$\begin{aligned}
 \left\| \nabla^2 g^{[t, \psi_i] \cdot \cdot} \right\|_{L^\infty} &= \left\| \nabla^2 (\psi_i^{-1} \circ \psi)^* P_t(\psi_i^* g) \right\|_{L^\infty} \\
 &\leq C_n \|\psi_i^{-1} \circ \psi\|_{C^3}^2 \|\nabla^2 P_t(\psi_i^* g)\|_{L^\infty} \\
 &= C_n \|\psi_i^{-1} \circ \psi\|_{C^3}^2 \|P_t(\nabla^2 \psi_i^* g)\|_{L^\infty} \quad \text{by (2.5)} \\
 &\leq C_{n, \alpha, Q, r} \|\varphi_t\|_{L^q} \|\nabla^2 \psi_i^* g\|_{L^p} \quad \text{by (2.Sch) and (2.Y)} \\
 (3.8c) \quad &\leq C_{n, \alpha, Q, r, p} t^{-n/p} \quad \text{by (3.3)}.
 \end{aligned}$$

Before starting with ansatz (1.1), we decompose  $R_{g^{[t]}}$  into a convex combination of functions. To this end observe

$$\begin{aligned}
 R_{g^{[t]}}(\vec{v}) &= A_{\vec{v}} \left( g^{[t] \cdot \cdot}, \nabla^2 g^{[t] \cdot \cdot} \right) + b_{\vec{v}}^{[t]} \\
 &= A_{\vec{v}} \left( g^{[t] \cdot \cdot}, \nabla^2 \sum_i \varrho_i g^{[t, \psi_i] \cdot \cdot} \right) + b_{\vec{v}}^{[t]} \\
 &= A_{\vec{v}} \left( g^{[t] \cdot \cdot}, \nabla^2 \sum_i \varrho_i g \cdot \cdot - \sum_i \nabla^2 \varrho_i g^{\Delta t, \psi_i} \cdot \cdot \right) + b_{\vec{v}}^{[t]} \\
 &= A_{\vec{v}} \left( g^{[t] \cdot \cdot}, \nabla^2 g \cdot \cdot - \sum_i \nabla^2 \varrho_i g^{\Delta t, \psi_i} \cdot \cdot \right) + b_{\vec{v}}^{[t]} \\
 &= A_{\vec{v}} \left( g^{[t] \cdot \cdot}, \sum_i \varrho_i \nabla^2 g \cdot \cdot - \sum_i \nabla^2 \varrho_i g^{\Delta t, \psi_i} \cdot \cdot \right) + b_{\vec{v}}^{[t]} \\
 &= \sum_i A_{\vec{v}} \left( g^{[t] \cdot \cdot}, \varrho_i \nabla^2 g \cdot \cdot - \nabla^2 \varrho_i g^{\Delta t, \psi_i} \cdot \cdot \right) + b_{\vec{v}}^{[t]} \\
 &= \sum_i A_{\vec{v}} \left( g^{[t] \cdot \cdot}, \varrho_i \nabla^2 \left( g^{[t, \psi_i]} + g^{\Delta t, \psi_i} \right) \cdot \cdot - \nabla^2 \varrho_i g^{\Delta t, \psi_i} \cdot \cdot \right) + b_{\vec{v}}^{[t]} \\
 &= \sum_i A_{\vec{v}} \left( g^{[t] \cdot \cdot}, \varrho_i \nabla^2 g^{[t, \psi_i]} \cdot \cdot - \left( \nabla^2 \varrho_i g^{\Delta t, \psi_i} \cdot \cdot - \varrho_i \nabla^2 g^{\Delta t, \psi_i} \cdot \cdot \right) \right) + b_{\vec{v}}^{[t]}.
 \end{aligned}$$

Thus for a dimension-dependent constant  $C_n$  and using definition (2.2a)

$$\begin{aligned}
 &\left\| R_{g^{[t]}}(\vec{v}) - \sum_i A_{\vec{v}} \left( g^{[t] \cdot \cdot}, \varrho_i \nabla^2 g^{[t, \psi_i]} \cdot \cdot \right) - b_{\vec{v}}^{[t]} \right\|_{L^\infty} \\
 &\leq \sum_i \left\| A_{\vec{v}} \left( g^{[t] \cdot \cdot}, \nabla^2 \varrho_i g^{\Delta t, \psi_i} \cdot \cdot - \varrho_i \nabla^2 g^{\Delta t, \psi_i} \cdot \cdot \right) \right\|_{L^\infty} \\
 &\leq C_n \bar{v} \sum_i \left\| g^{[t] \cdot \cdot} \right\|_{L^\infty} \left\| \nabla^2 \varrho_i g^{\Delta t, \psi_i} \cdot \cdot - \varrho_i \nabla^2 g^{\Delta t, \psi_i} \cdot \cdot \right\|_{L^\infty} \\
 &\leq C'_{n, Q, r, \alpha} \bar{v} \sum_i \left\| \nabla^2 \varrho_i g^{\Delta t, \psi_i} \cdot \cdot - \varrho_i \nabla^2 g^{\Delta t, \psi_i} \cdot \cdot \right\|_{L^\infty} \\
 &\leq C'_{n, Q, r, \alpha} \bar{v} \sum_i \left( \left\| \nabla^2 \varrho_i \right\|_{L^\infty} \left\| g^{\Delta t, \psi_i} \cdot \cdot \right\|_{L^\infty} + 2 \left\| \nabla \varrho_i \right\|_{L^\infty} \left\| \nabla g^{\Delta t, \psi_i} \cdot \cdot \right\|_{L^\infty} \right) \\
 &\leq 2C'_{n, Q, r, \alpha} \bar{v} C_\varrho \sum_i \left\| g^{\Delta t, \psi_i} \cdot \cdot \right\|_{W^{1, \infty}}
 \end{aligned}$$

where we used (3.8a) in the antepenultimate step, and definition (3.7e) in the last step. As  $g^{\Delta t, \psi_i} \cdot \cdot = \psi^*(\psi_i^{-1})^*(\psi_i^* g)^{\Delta t} = (\psi \circ \psi_i^{-1})^*(\psi_i^* g)^{\Delta t}$ , combining estimate (2.Sch) and (3.5a) gives

$$\left\| R_{g^{[t]}}(\vec{v}) - \sum_i A_{\vec{v}} \left( g^{[t] \cdot \cdot}, \varrho_i \nabla^2 g^{[t, \psi_i]} \cdot \cdot \right) - b_{\vec{v}}^{[t]} \right\|_{L^\infty} \leq C' \bar{v} t^\alpha$$

for a constant  $C' = C'(n, Q, r, \{\varrho_i \circ \psi_i\}_{i \in I}, \alpha)$ . Finally, this gives

$$\begin{aligned}
 \mathbf{R}_{g^{[t]}}(x)(\vec{v}) &\leq \sum_i A_{\vec{v}} \left( g^{[t]\cdot\cdot}, \varrho_i \nabla^2 g^{[t, \psi_i]}\cdot\cdot \right) (x) + b_{\vec{v}}^{[t]}(x) + C' t^\alpha \\
 &\leq \sum_i \varrho_i \left( A_{\vec{v}} \left( g^{[t]\cdot\cdot}, \nabla^2 g^{[t, \psi_i]}\cdot\cdot \right) + b_{\vec{v}}^{[t]} \right) (x) + C' t^\alpha \\
 (3.9) \quad &\leq \sup_{\substack{i \in I \\ |x \psi_i(0)|_g < r/2}} \left( A_{\vec{v}} \left( g^{[t]\cdot\cdot}, \nabla^2 g^{[t, \psi_i]}\cdot\cdot \right) + b_{\vec{v}}^{[t]} \right) (x) + C' t^\alpha.
 \end{aligned}$$

This means that it is sufficient to find an estimate

$$\left( A_{\vec{v}} \left( g^{[t]\cdot\cdot}, \nabla^2 g^{[t, \psi_i]}\cdot\cdot \right) + b_{\vec{v}}^{[t]} \right) (x) - \mathbf{R}_g(x)(\vec{v}) \leq C'' \bar{v} t^\alpha$$

for every  $i \in I$ .

We are now in a position to apply an estimate similar to the one from the proof of Proposition 3.1:

$$\begin{aligned}
 (3.10) \quad &\left( A_{\vec{v}} \left( g^{[t]\cdot\cdot}, \nabla^2 g^{[t, \psi_i]}\cdot\cdot \right) + b_{\vec{v}}^{[t]} \right) (x) \leq \mathbf{R}_{g^{[t, \psi_i]}\cdot\cdot}(x)(\vec{v}) + \\
 &+ \left\| A_{\vec{v}} \left( g^{[t]\cdot\cdot} - g^{[t, \psi_i]}\cdot\cdot, \nabla^2 g^{[t, \psi_i]}\cdot\cdot \right) \right\|_{L^\infty} + \left\| b_{\vec{v}}^{[t]} - b_{\vec{v}}^{[t, \psi_i]} \right\|_{L^\infty}
 \end{aligned}$$

The proof is reduced to seeking bounds for the second and third summand.

With regard to the second summand we estimate for a constant  $C_n$  using definition (2.2a)

$$\begin{aligned}
 &\left\| A_{\vec{v}} \left( g^{[t]\cdot\cdot} - g^{[t, \psi_i]}\cdot\cdot, \nabla^2 g^{[t, \psi_i]}\cdot\cdot \right) \right\|_{L^\infty} \\
 &\leq C_n \bar{v} \left\| g^{[t]\cdot\cdot} - g^{[t, \psi_i]}\cdot\cdot \right\|_{L^\infty} \left\| \nabla^2 g^{[t, \psi_i]}\cdot\cdot \right\|_{L^\infty} \\
 &\leq C_n \bar{v} \left( \left\| (g^{\Delta t})\cdot\cdot \right\|_{L^\infty} + \left\| (g^{\Delta t, \psi_i})\cdot\cdot \right\|_{L^\infty} \right) \left\| \nabla^2 g^{[t, \psi_i]}\cdot\cdot \right\|_{L^\infty} \\
 &\leq C_n \bar{v} \left( C'_{n, r, \alpha, Q} t^\alpha + C'_{n, r, \alpha, Q, C_\varrho} t^\alpha \right) C_{n, \alpha, Q, r, p} t^{n/p}
 \end{aligned}$$

where we used (3.8a) and (3.8c) in the last step. To summarize: the second summand vanishes with modulus  $t^{\alpha-n/p}$ .

As for the third summand by Lemma 3.4 we have a constant  $C = C(\|(\nabla g^{[t]}\cdot\cdot, \nabla^{\leq 1} g^{[t]}\cdot\cdot)\|_{L^\infty})$  such that we can estimate

$$\begin{aligned}
 \left\| b_{\vec{v}}^{[t]} - b_{\vec{v}}^{[t, \psi_i]} \right\|_{L^\infty} &\leq C \bar{v} \left\| \left( \nabla g^{[t]}\cdot\cdot, \nabla^{\leq 1} g^{[t]}\cdot\cdot \right) - \left( \nabla g^{[t, \psi_i]}\cdot\cdot, \nabla^{\leq 1} g^{[t, \psi_i]}\cdot\cdot \right) \right\|_{L^\infty} \\
 &\leq C \bar{v} \left( \left\| g^{[t]\cdot\cdot} - g^{[t, \psi_i]}\cdot\cdot \right\|_{W^{1, \infty}} + \left\| \nabla g^{[t]}\cdot\cdot - \nabla g^{[t, \psi_i]}\cdot\cdot \right\|_{L^\infty} \right).
 \end{aligned}$$

Finally, by (3.8b) we have a bound

$$\begin{aligned}
 \left\| g^{[t]\cdot\cdot} - g^{[t, \psi_i]}\cdot\cdot \right\|_{W^{1, \infty}} &\leq \left\| g^{[t]\cdot\cdot} - g^\cdot \right\|_{W^{1, \infty}} + \left\| g^\cdot - g^{[t, \psi_i]}\cdot\cdot \right\|_{W^{1, \infty}} \\
 &\leq \left( C'_{n, r, \alpha, Q} + C'_{n, r, \alpha, Q, C_\varrho} \right) t^\alpha
 \end{aligned}$$

and by (3.5a) and (2.Sch) there is a bound

$$\begin{aligned}
 \left\| \nabla g^{[t]}\cdot\cdot - \nabla g^{[t, \psi_i]}\cdot\cdot \right\|_{L^\infty} &\leq \left\| \nabla g^{[t]}\cdot\cdot - \nabla g^\cdot \right\|_{L^\infty} + \left\| \nabla g^\cdot - \nabla g^{[t, \psi_i]}\cdot\cdot \right\|_{L^\infty} \\
 &\leq \left( C_{n, \alpha, Q, C_\varrho} + C'_{n, r, \alpha, Q} \right) t^\alpha.
 \end{aligned}$$



Thus combining (3.9) with (3.10) we obtain

$$\begin{aligned}
R_{g^{[t]}}(x)(\vec{v}) &\leq \sup_{\substack{i \in I \\ |x \psi_i(0)|_g < r/2}} R_{g^{[t, \psi_i]}}(x)(\vec{v}) + C\bar{v}t^\alpha \\
&\leq \sup \left\{ R_g(y)(\vec{v}) \mid y \in \bigcup_{\substack{i \in I \\ |x \psi_i(0)|_g < r/2}} \psi_i(B(0, t)) \right\} + C\bar{v}t^\beta \\
&\leq \sup_{y \in B(x, e^Q t)} R_g(y)(\vec{v}) + C\bar{v}t^\beta
\end{aligned}$$

where in the penultimate step we used Proposition 3.1 which is applicable since  $R_{g^{[t, \psi_i]}}(\vec{v}) = R_{g^{[t, \psi_i]}}(\vec{v})$  is coordinate independent; and in the last step the distance estimate (2.D) came to hand. By (2.N<sub>0</sub>), we further have  $\bar{v} \leq n^4 \cdot \left(\sqrt{e^{2Q}}\right)^4 \|\vec{v}\|_g$ . This implies the bound we sought.  $\square$

#### § 4. Consequence for sectional curvature

This section is devoted to the main result of the paper, which is again phrased in terms of Sobolev chart norms  $\|\cdot\|_{W^{2,p,r}}^{\text{harm}}$ , see (2.N<sub>W<sup>m,p</sup></sub>).

**Theorem B.** *Let  $p > 2n$ ,  $r, Q > 0$ , and  $\beta \in (0, 1 - 2n/p)$ . Then there is some  $T > 0$  such that for any smooth riemannian manifold  $(M, g)$  with  $\|(M, g)\|_{W^{2,p,r}}^{\text{harm}} \leq Q$  there is a locally finite cover of charts*

$$\psi_i: (B(0, r), 0) \rightarrow (M, \psi_i(0)),$$

*indexed by  $i \in I$ , such that the mollified metrics  $g^{[t]}$  have at any  $x \in M$  sectional curvature  $K_{g^{[t]}}(x)$  in the interval*

$$(1.2) \quad \left[ \underline{K}(g \cdot |_{B(x,t)}), \overline{K}(g \cdot |_{B(x,t)}) \right] \cdot [1 - Ct, 1 + Ct] + [-Ct^\beta, Ct^\beta]$$

*for all  $t \in (0, T]$ , where  $C = C(n, Q, r, p)$ . Moreover, we have that*

$$(1.3) \quad \|M\|_{C^{2,\beta},r} \leq Q'$$

*for some  $Q' = Q'(n, \beta, Q)$ .*

A glance at (2.K<sub>|Ric|</sub>) gives immediately:

**Corollary C.** *Let  $\iota > 0$  and  $\kappa \geq 0$ . Then there exist  $r > 0$  and  $T > 0$  such that for any smooth riemannian manifold  $(M, g)$  with*

$$\text{inj rad}(M) \geq \iota, \quad \|\text{Ric } M\|_{L^\infty} \leq \kappa$$

*there exist charts  $\psi_i: B(0, r) \rightarrow M$  such that the mollified metrics  $g^{[t]}$  have at any  $x \in M$  sectional curvature  $K_{P_t g}(x)$  in the interval (1.2) for all  $t \in (0, T]$ , where  $C = C(n, \iota, \kappa)$ . Moreover, the norm  $\|M\|_{C^{2,\beta},r}$  is bounded in terms of  $n, \beta$  and  $Q$ .*

**Corollary D.** *Let  $\delta \in (0, 1)$  and assume one of the conditions (1.4a) to (1.4d):*

$$\begin{aligned}
 (1.4a) \quad & \left\{ \begin{array}{l} \#\pi_2(M) < \infty, \text{ or} \\ \dim M \text{ is even, or} \\ \delta \geq 1/4 - \varepsilon', \text{ where } \varepsilon' \approx 10^{-6} \end{array} \right. \\
 (1.4b) \quad & K_g \in [\delta, 1] \text{ and} \\
 (1.4c) \quad & \\
 (1.4d) \quad & K_g \leq 1, \text{Ric}_g \geq \delta, \text{ and } \dim M = 3.
 \end{aligned}$$

*Then there exist charts  $\psi_i: B(0, r) \rightarrow M$  such that the mollified metrics  $g^{[t]}$  have at any  $x \in M$  sectional curvature  $K_{P_t g}(x)$  in the interval (1.2) for all  $t \in (0, T]$ , where  $C = C(n, \kappa)$ . Moreover, the norm  $\|M\|_{C^{2,\beta},r}$  is bounded in terms of  $n, \beta$  and  $Q$ .*

*Proof.* Each of (1.4a) to (1.4d) implies a uniform positive lower bound on the injectivity radius [Tus00, p. 6].  $\square$

The first step to prove Theorem B is to introduce the notion of a locally  $N$ -finite cover: a cover  $\{U_i\}_{i \in I}$  of a space  $X$  is **locally  $N$ -finite** if every point of  $x$  is contained in at most  $N$  members of  $\{U_i\}_I$ . By help of this terminology one can reduce Theorem B to the following claim:

**Lemma 4.1.** *Let  $p > 2n$ ,  $r, Q > 0$ ,  $\beta \in (0, 1 - 2n/p)$  and  $N$  be a natural number. Then there is  $T' > 0$  such that for any smooth riemannian manifold  $M$  with  $\|(M, g)\|_{W^{2,p},r}^{\text{harm}} \leq Q$  and any*

$$\{\psi_i: (B(0, r), 0) \rightarrow (M, \psi_i(0))\}_{i \in I}, \quad \vec{\varrho} := \{\varrho_i: M \rightarrow [0, 1]\}_{i \in I}$$

*a cover of  $M$  by charts and a corresponding partition of unity with*

- $\|\psi_i\|_{W^{2,p},r}^{\text{harm}} \leq Q$ ,
- $\{\psi_i(B(0, r)) \mid i \in I\}$  is locally  $N$ -finite,
- $M$  is covered by  $\{\psi_i(B(0, e^{-Q}r/2))\}_{i \in I}$ , and
- $\|\varrho_i \circ \psi_i\|_{C^2} < C_I$  for all  $i \in I$  and a constant  $C_I$

*the mollified metrics  $g^{[t]}$  have at any  $x \in M$  sectional curvature  $K_{P_t g}(x)$  in the interval (1.2) for all  $t \in (0, T']$ , where  $C = C(n, Q, r, p)$ .*

*Proof of Theorem B using Lemma 4.1.* By Gromov [Gro07, Proposition 5.2] every Gromov-Hausdorff compact class  $\mathcal{M}$  of (isometry classes of) pointed metric spaces has the following property: for every space  $(M, d, p_0) \in \mathcal{M}$  the maximal number of disjoint closed balls of radius  $\varepsilon$  that fit into  $B[p_0, R]$  is bounded by a finite number  $N(\varepsilon, R)$ . By Sobolev's inequality (2.S) we have the inclusion  $\mathcal{M}(W^{2,p} \leq_r Q) \subset \mathcal{M}(C^{1,\alpha} \leq_r CQ)$  for  $\alpha = n/p$  and a constant  $C = C(n, p)$ . Hence by (2.KC $^{m,\alpha}$ ) the space  $\mathcal{M}(W^{2,p} \leq_r Q)$  is precompact in the Gromov-Hausdorff topology and Gromov's result is applicable.

Now we apply Gromov's result to  $\varepsilon = re^{-2Q}/5$  and  $R = 2re^Q$  obtaining a bound  $N$ . Let  $\{B[x_i, \varepsilon]\}_{i \in I}$  be some maximal disjoint system of  $\varepsilon$ -balls in  $(M, g, p_0) \in \mathcal{M}(W^{2,p} \leq_r Q)$ . The balls  $\{B(x_i, re^{-2Q}/2)\}_{i \in I}$  cover  $M$  due to maximality of the system. On the other hand for any  $x \in M$  the cardinality of the set  $\{i \in I \mid x \in B(x_i, re^Q)\} \subset \{i \in$

$I \mid B(x_i, re^Q) \subset B[x, R] \}$  is bounded by  $N$ . For each point  $x_i$  choose a chart  $\psi_i: (B(0, r), 0) \rightarrow (M, x_i)$  with  $\|\psi_i\|_{W^{2,p},r} \leq Q$ . By (2.D),  $B(x_i, re^{-2Q}/2) \subset \psi_i(B(0, e^{-Q}r/2))$  and, hence  $\{\psi_i(B(0, e^{-Q}r/2))\}_{i \in I}$  is covering. By the same estimate,  $\psi_i(B(0, r)) \subset B[x_i, re^Q]$  and, thus,  $\{\psi_i(B(0, r))\}_{i \in I}$  is  $N$ -finite.

To find a suitable partition of unity, choose any bump function  $b: \mathbb{R}^d \rightarrow \mathbb{R}$  with  $\text{supp } b \subset B(0, 3r/4)$  and  $b|_{B(0, re^{-2Q}/2)} \equiv 1$ . We define for all  $i \in I$

$$(4.1a) \quad b_i(x) := \begin{cases} b \circ \psi_i^{-1}(x) & \text{if } x \in \psi_i(B(0, r)) \\ 0 & \text{otherwise} \end{cases}$$

$$(4.1b) \quad \varrho_i(x) := \frac{1}{\sum_{j \in I} b_j(x)} b_i(x).$$

Due to (2.Sch) there is a uniform  $C^{3,\beta}$ -bound on the transition maps  $\psi_i^{-1} \circ \psi_j$  for  $i, j \in I$ . Moreover,  $b$  is  $C^{3,\beta'}$ -bounded since its support is compact. Hence there is a uniform  $C^{3,\beta}$ -bound on  $b_i \circ \psi_j = b \circ \psi_i^{-1} \circ \psi_j$  for each  $i \in I$ . Since the denominator in (4.1b) is at least 1 at each  $x \in M$  for at least one  $i$  (namely the  $i$  for which  $|\psi_i(0)x| \leq e^{-Q}r/2$ ),  $\varrho_i$  is  $C^{2,\beta}$ -bounded uniformly in  $i \in I$  as well. This puts us in a situation to apply Lemma 4.1.

Due to (2.6c) we have a uniform  $C^{2,\beta}$ -bound on  $P_T(\psi_i^*g)$  and thus a uniform  $C^{2,\beta}$ -bound on the pullback metrics  $(\psi_i^{-1} \circ \psi_j)^* P_T(\psi_i^*g)$  for all  $i, j \in I$  as well. Together with the bound on the  $\varrho_i$ 's from last paragraph this implies a uniform  $C^{2,\beta}$ -bound on  $\psi_i^*g^{[T]}$  for each  $i \in I$ . Thus the bound (1.3) holds.  $\square$

*Proof of Lemma 4.1.* We apply Theorem A obtaining a constant

$$C_1 = C(n, p, r, Q, \beta, C_I, N).$$

From  $\{\psi_i\}_{i \in I}$ ,  $\{\varrho_i\}_{i \in I}$  we get a mollification of  $g$ :  $g^{[-]}: [0, T] \rightarrow \Gamma(\text{Sym}^{0,2} M)$  with  $g^{[0]} := g$  and the property

$$(4.2) \quad R_{g^{[t]}}(x)(\vec{v}) - \sup_{y \in B(p, e^Q t)} R_g(y)(\vec{v}) \leq C_1 \|\vec{v}\|_g t^\beta$$

for any  $x \in M$  and any section  $\vec{v} \in (TM)^{\times 3} \times T^*M$ . For convenience we abbreviate

$$\begin{aligned} \langle -, - \rangle_t &:= \langle -, - \rangle_{g^{[t]}}, & \|-\|_t &:= \|-\|_{g^{[t]}}, \\ R_t &:= R_{g^{[t]}}, & K_t &:= K_{g^{[t]}}. \end{aligned}$$

We further agree on the following shorthands for intervals:  $[a \pm b] := [a - b, a + b]$ ,  $[\pm b] := [0 \pm b]$ , and

$$[f(x)]_{x \in X} := \left[ \inf_{x \in X} f(x), \sup_{x \in X} f(x) \right].$$

Moreover, from multi-linearity of the curvature tensor (use e.g.  $R(v, w) = -R(-v, w)$ ) we get immediately the reversed version of (4.2)

$$-R_{g^{[t]}}(x)(\vec{v}) + \inf_{y \in B(x, e^Q t)} R_g(y)(\vec{v}) \geq -C_1 \|\vec{v}\|_g t^\beta.$$

We seek a  $T' \in (0, T]$  such that the claim of the theorem holds, i.e.

$$K_t(x)(v, w) = \frac{R_t(\vec{v})}{\|v\|_t \|w\|_t - \langle v, w \rangle_t},$$

where  $R_t(\vec{v}) = \langle R_t(x)(v, w)w, v \rangle_t$ , is in the interval

$$(4.3) \quad [K_0(y)(v, w)]_{\substack{y \in B(x, e^Q) \\ v, w \in T_y M}} + [\pm C t^\beta]$$

for all  $t \in (0, T']$ ,  $x \in M$ , and  $v, w \in T_x M$  linear independent. Since the sectional curvature depends only on the plane spanned by  $v$  and  $w$ , we can assume without loss of generality that

$$\|v\|_0 = \|w\|_0 = 1, \quad \langle v, w \rangle_0 = 0.$$

Fix some chart  $\psi: (B(0, r), 0) \rightarrow (M, p_0)$  with  $\|\psi\|_{W^{2,p},r}^{\text{harm}} \leq Q$ . We extend  $v$  and  $w$  to sections  $v_\psi, w_\psi \in \Gamma T^{3,1} \psi(B(0, r))$  using the euclidean identification, i.e. by pushing forward the constant sections  $v$  and  $w$  along  $\psi$ . Again  $\vec{v}_\psi := (v_\psi, w_\psi, w_\psi, \langle -, v_\psi \rangle_t)$ .

By (2.N<sub>0</sub>) we have  $\|v_\psi\|_{\text{eucl.}}, \|w_\psi\|_{\text{eucl.}} \leq e^{2Q}$ . Hence  $\|v_\psi\|_g, \|w_\psi\|_g \leq e^{4Q}$ . Thus  $\|\vec{v}_\psi\|_g \leq e^{4 \cdot 4Q} = e^{16Q}$ . By Sobolev's inequality the entries of the metric tensor are at least Lipschitz with some bound  $C_{n,p,r,Q}$ . Hence we have

$$\begin{aligned} (\|v_\psi\|_0 \|w_\psi\|_0 - \langle v_\psi, w_\psi \rangle_0)(\psi(y)) &= 1 - \langle v_\psi, w_\psi \rangle_0(\psi(y)) \\ &\in [1 \pm C_{n,p,r,Q} e^{2Q} \cdot |0 y|] \end{aligned}$$

for  $y \in B(0, r)$ . The Lipschitz bound on the metric tensor implies further via Lemma 3.6

$$(\|v_\psi\|_t \|w_\psi\|_t - \langle v_\psi, w_\psi \rangle_t)(\psi(y)) \in [1 \pm C'_2 t]$$

for  $y \in B(0, e^Q t)$  and  $t \in [0, T]$ . Choose  $T'$  so small that  $C'_2 T' \leq 1/2$ . Hence we can choose a constant  $C_2 > 0$  such that

$$\frac{1}{\|v_\psi\|_t \|w_\psi\|_t - \langle v_\psi, w_\psi \rangle_t}(\psi(y)) \in [1 \pm C_2 t]$$

and choose  $T' > 0$  so small that  $1 - C_2 t \geq 1/2$  for all  $t \in (0, T']$ .

Gathering all estimates above we conclude the proof:

$$\begin{aligned} K_t(x)(v, w) &= \frac{R_t(\vec{v}_\psi)}{\|v_\psi\|_t \|w_\psi\|_t - \langle v_\psi, w_\psi \rangle_t}(x) \\ &\in R_t(x)(\vec{v}_\psi) [1 \pm C_2 t] \\ &\subset \left( [R_0(y)(\vec{v}_\psi)]_{y \in B(x, e^Q t)} + [\pm C_1 \|\vec{v}_\psi\|_g t^\beta] \right) [1 \pm C_2 t] \end{aligned}$$

for some new constant  $C'$  this is included in

$$\begin{aligned} &\subset \left( [R_0(y)(\vec{v}_\psi)]_{y \in B(x, e^Q t)} \right) [1 \pm C_2 t] + [\pm C' t^\beta] \\ &\subset \left( [K_0(y)(v_\psi, w_\psi)]_{y \in B(x, e^Q t)} \right) [1 \pm C'_2 t] [1 \pm C_2 t] + [\pm C' t^\beta]. \end{aligned}$$

This is contained in the interval (4.3) for some constant  $C$ . This proves the theorem.  $\square$

## References

- [Abr88] U. Abresch. “Über das Glätten Riemann’scher Metriken”. Habilitation. 1988.
- [AC92] M. T. Anderson and J. Cheeger. “ $C^\alpha$ -compactness for manifolds with Ricci curvature and injectivity radius bounded below”. In: *Journal of Differential Geometry* 35.2 (1992), pp. 265–281.
- [And+04] M. Anderson et al. “Boundary regularity for the Ricci equation, geometric convergence, and Gel’fand’s inverse boundary problem”. In: *Inventiones mathematicae* 158.2 (2004), pp. 261–321.
- [And90] M. T. Anderson. “Convergence and rigidity of manifolds under Ricci curvature bounds”. In: *Inventiones Mathematicae* 102.2 (1990), pp. 429–445.
- [BMR84] J. Bemelmans, Min-Oo, and E. A. Ruh. “Smoothing Riemannian metrics”. In: *Mathematische Zeitschrift* 188.1 (1984), pp. 69–74.
- [CDK11] G. Csató, B. Dacorogna, and O. Kneuss. *The Pullback Equation for Differential Forms*. Progress in Nonlinear Differential Equations and Their Applications 83. Boston: Birkhäuser, 2011.
- [CG85] J. Cheeger and M. Gromov. “On the Characteristic Numbers of Complete Manifolds of Bounded Curvature and Finite Volume”. In: *Differential Geometry and Complex Analysis*. 1985.
- [Gro07] M. Gromov. *Metric Structures for Riemannian and Non-Riemannian Spaces*. Ed. by J. Lafontaine and P. Pansu. Trans. from the French by S. M. Bates. Modern Birkhäuser Classics. Birkhäuser, 2007.
- [GT15] D. Gilbarg and N. Trudinger. *Elliptic Partial Differential Equations of Second Order*. Classics in Mathematics. Berlin Heidelberg: Springer, 2015.
- [He16] F. He. *Existence and applications of Ricci flows via pseudolocality*. Version 2. 2016. arXiv: 1610.01735 [math.DG].
- [HH97] E. Hebey and M. Herzlich. “Harmonic coordinates, harmonic radius and convergence of Riemannian manifolds”. In: *Rendiconti di Matematica e delle sue Applicazioni*. 7th ser. 17.4 (1997), pp. 569–605.
- [Hör83] L. Hörmander. *The Analysis of Linear Partial Differential Operators. Distribution Theory and Fourier Analysis*. Vol. 1. Grundlehren 256. Springer, 1983.
- [Kap05] V. Kapovitch. “Curvature bounds via Ricci smoothing”. In: *Illinois Journal of Mathematics* 49.1 (2005), pp. 259–263.
- [Pet16] P. Petersen. *Riemannian Geometry*. 3rd ed. Graduate Texts in Mathematics 171. Springer, 2016.
- [Ron96] X. Rong. “On the fundamental groups of manifolds of positive sectional curvature”. In: *Annals of Mathematics. Second Series* 143.2 (1996), pp. 397–411.

- [Shi89] W.-X. Shi. “Deforming the metric on complete Riemannian manifolds”. In: *Journal of Differential Geometry* 30.1 (1989), pp. 223–301.
- [Tay06] M. E. Taylor. “Existence and regularity of isometries”. In: *Transactions of the American Mathematical Society* 358.6 (June 2006), pp. 2415–2423.
- [Tay07] M. E. Taylor. *Tools for PDE. Pseudodifferential Operators, Pseudodifferential Operators, and Layer Potentials*. Mathematical Surveys and Monographs 81. American Mathematical Society, 2007.
- [Tay13] M. E. Taylor. *Partial Differential Equations*. Vol. 2. 2nd ed. Applied Mathematical Sciences 116. Springer, 2013.
- [Tus00] W. Tuschmann. “Endlichkeitssätze und positive Krümmung”. Habilitation. Max-Planck-Institut für Mathematik in den Naturwissenschaften, 2000.

(D. LUCKHARDT) DEPARTMENT OF MATHEMATICS, BEN-GURION UNIVERSITY OF THE NEGEV, ISRAEL.

*E-mail address:* luckhard@post.bgu.ac.il

(J.-B. KORDAß) DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DE FRIBOURG, SWITZERLAND.

*E-mail address:* jan-bernhard.kordass@unifr.ch