

On number of pendants in local antimagic chromatic number

Gee-Choon Lau^{a,1}, Wai-Chee Shiu^{b,c}, Ho-Kuen Ng^d

^a*Faculty of Computer & Mathematical Sciences,
Universiti Teknologi MARA (Segamat Campus),
85000, Johor, Malaysia.
geeclau@yahoo.com*

^b*College of Global Talents, Beijing Institute of Technology,
Zhuhai, China.*

^c*Department of Mathematics, The Chinese University of Hong Kong,
Shatin, Hong Kong.
wcshiu@associate.hkbu.edu.hk*

^d*Department of Mathematics, San José State University,
San José CA 95192 USA.
ho-kuen.ng@sjsu.edu*

Abstract

An edge labeling of a connected graph $G = (V, E)$ is said to be local antimagic if it is a bijection $f : E \rightarrow \{1, \dots, |E|\}$ such that for any pair of adjacent vertices x and y , $f^+(x) \neq f^+(y)$, where the induced vertex label $f^+(x) = \sum f(e)$, with e ranging over all the edges incident to x . The local antimagic chromatic number of G , denoted by $\chi_{la}(G)$, is the minimum number of distinct induced vertex labels over all local antimagic labelings of G . Let $\chi(G)$ be the chromatic number of G . In this paper, sharp upper and lower bounds of $\chi_{la}(G)$ for G with pendant vertices, and sufficient conditions for the bounds to equal, are obtained. Consequently, for $k \geq 1$, there are infinitely many graphs with $k \geq \chi(G) - 1$ pendant vertices and $\chi_{la}(G) = k + 1$. We conjecture that every tree T_k , other than certain caterpillars, spiders and lobsters, with $k \geq 1$ pendant vertices has $\chi_{la}(T_k) = k + 1$.

Keywords: Local antimagic labeling, local antimagic chromatic number, cut-vertices, pendants.

2010 AMS Subject Classifications: 05C78, 05C69.

1 Introduction


A connected graph $G = (V, E)$ is said to be *local antimagic* if it admits a *local antimagic edge labeling*, i.e., a bijection $f : E \rightarrow \{1, \dots, |E|\}$ such that the induced vertex labeling $f^+ : V \rightarrow \mathbb{Z}$ given by $f^+(u) = \sum f(e)$ (with e ranging over all the edges incident to u) has the property that any two adjacent vertices have distinct induced vertex labels [1]. Thus, f^+ is a coloring of G . Clearly, the order of G must be at least 3. The vertex label $f^+(u)$ is called the *induced color* of u under f (the *color* of u , for short, if no ambiguous occurs). The number of distinct induced colors under f is denoted by $c(f)$, and is called the *color number* of f . The *local antimagic chromatic number* of G , denoted by $\chi_{la}(G)$, is $\min\{c(f) : f \text{ is a local antimagic labeling of } G\}$. Clearly, $2 \leq \chi_{la}(G) \leq |V(G)|$. Haslegrave [4] proved that every graph is local antimagic. This paper relates the number of pendant vertices of G to $\chi_{la}(G)$. Sharp upper and lower bounds, and sufficient conditions for the bounds to equal, are obtained. Consequently, there exist infinitely many graphs with $k \geq \chi(G) - 1 \geq 1$ pendant vertices and $\chi_{la}(G) = k + 1$. We conjecture that every tree T_k , other than certain caterpillars, with $k \geq 1$ pendant vertices has $\chi_{la}(T_k) = k + 1$. The following two results in [7] are needed.

Lemma 1.1. *Let G be a graph of size q containing k pendants. Let f be a local antimagic labeling of G such that $f(e) = q$. If e is not a pendant edge, then $c(f) \geq k + 2$.*

¹Corresponding author.

Theorem 1.2. *Let G be a graph having k pendants. If G is not K_2 , then $\chi_{la}(G) \geq k + 1$ and the bound is sharp.*

The sharp bound for $k \geq 2$ is given by the star S_k with maximum degree k . The left labeling below is another example for $k = 2$. The right labeling shows that the lower bound is sharp for $k = 1$.



For $1 \leq i \leq t$, $a_i, n_i \geq 1$ and $d = \sum_{i=1}^t n_i \geq 3$, a spider of d legs, denoted $Sp(a_1^{[n_1]}, a_2^{[n_2]}, \dots, a_t^{[n_t]})$, is a tree formed by identifying an end-vertex of n_i path(s) of length a_i . The vertex u of degree d is the core of the spider. Note that $Sp(1^{[n]})$ is the star graph of n pendant vertices with $\chi_{la}(Sp(1^{[n]})) = n + 1$. We first give a family of spiders with k pendant vertices to have $\chi_{la} > k + 1$.

Theorem 1.3. *For $n \geq 3$,*

$$\chi_{la}(Sp(2^{[n]})) = \begin{cases} n + 2 & \text{if } n \geq 4 \\ n + 1 & \text{otherwise.} \end{cases}$$

Proof. Let the neighbors of u be v_1, \dots, v_n . Let w_i be the pendant adjacent to v_i , $1 \leq i \leq n$. By Theorem 1.2, it is easy to conclude that $\chi_{la}(Sp(2^{[3]})) = 4$. Consider $n \geq 4$. Let f be a local antimagic labeling of $Sp(2^{[n]})$ with minimum $c(f)$. By Lemma 1.1 and symmetry, suffice to consider $f(v_1 w_1) = 2n = f^+(w_1)$. By definition, $f^+(u) \neq f^+(v_1)$. Now, $f^+(u) > n(n+1)/2 > 2n \geq f^+(w_i)$ and $f^+(v_1) = 2n > f^+(w_i)$ for $1 \leq i \leq n$. Thus, $c(f) \geq n + 2$ and $\chi_{la}(Sp(2^{[n]})) \geq n + 2$.

Define a bijection $f : E(Sp(2^{[n]})) \rightarrow [1, 2n]$ such that $f(uv_i) = i$ and $f(v_i w_i) = 2n + 1 - i$. We have $f^+(u) = n(n+1)/2 > f^+(v_i) = 2n + 1 > f^+(w_i) = 2n + 1 - i$, $1 \leq i \leq n$. Consequently, $\chi_{la}(Sp(2^{[n]})) = n + 2$ for $n \geq 4$. The theorem holds. \square

2 Adding pendant edges

Suppose G has $e \geq 2$ edges with $\chi_{la}(G) = t \geq 2$ such that the corresponding local antimagic labeling f induces a t -independent set $\{V_1, V_2, \dots, V_t\}$ with $|V_i| = n_i \geq 1$. Moreover, each non-pendant vertex must be in one of V_i for $1 \leq i \leq r \leq t$, and V_i is a singleton consisting of a pendant vertex for $r + 1 \leq i \leq t$ if $r < t$. Let $f^+(v) = c_i$ for each vertex $v \in V_i$. Without loss of generality, we assume that $c_1 < c_2 < \dots < c_r$ and that $c_{r+1} < c_{r+2} < \dots < c_t$. Note that c_{r+1} to c_t do not exist for $r = t$. By Theorem 1.2, G contains at most $t - 1$ pendant vertices.

Let $b \geq 0$ be the number of pendant vertices in $\cup_{i=1}^r V_i$ so that G has exactly $t - r + b$ pendant vertices. Let $f(u'v') = e \leq f^+(u') < f^+(v')$ such that $v' \in V_i$, $1 \leq i \leq r$ and that $u' \in V_j$. Suppose $e = f^+(u')$, then u' is a pendant vertex. We now have (a) $j < i \leq r$ so that $1 \leq b \leq j$ or (b) $i \leq r < j$ so that $0 \leq b \leq i - 1$. Otherwise, if $e < f^+(u')$, then u' is not a pendant vertex so that $j < i \leq r$ and $0 \leq b \leq j - 1$.

Let $G(V_i, s)$ be a graph obtained from G by adding s pendant edges $v_a w_{a,k}$ ($1 \leq k \leq s$) to each vertex in $V_i = \{v_a \mid 1 \leq a \leq n_i\}$.

Theorem 2.1. *Suppose G and $G(V_i, s)$ are as defined above. Let $r \geq 2$ and*

$$e + sn_i \geq \begin{cases} c_r & \text{for } i < r \\ c_{r-1} & \text{for } i = r \end{cases}$$

such that $s \geq 1$ if $n_i = 1$, and $s \geq 2$ is even if $n_i \geq 2$.

- (1) Suppose $e < c_1$. For $1 \leq i \leq r$, $\chi_{la}(G(V_i, s)) = sn_i + t - r + 1$.
- (2) Suppose $c_1 \leq e < c_2$, then $\chi_{la}(G(V_1, s)) = sn_1 + t - r + 1$. For $2 \leq i \leq r$, $sn_i + t - r + b + 1 \leq \chi_{la}(G(V_i, s)) \leq sn_i + t - r + 2$. Moreover, if $c_1 = e$ and $b = 1$, then $\chi_{la}(G(V_i, s)) = sn_i + t - r + 2$.
- (3) Suppose $c_{j-1} \leq e < c_j$ for $3 \leq j \leq r$, then
- (a) $sn_i + t - r + b \leq \chi_{la}(G(V_i, s)) \leq sn_i + t - r + j - 1$ for $1 \leq i \leq j - 1$, if V_i has a pendant vertex;
 - (b) $sn_i + t - r + b + 1 \leq \chi_{la}(G(V_i, s)) \leq sn_i + t - r + j - 1$ for $1 \leq i \leq j - 1$, if V_i has no pendant vertex;
 - (c) $sn_i + t - r + b + 1 \leq \chi_{la}(G(V_i, s)) \leq sn_i + t - r + j$ for $j \leq i \leq r$.
- Moreover, when $c_{j-1} = e$ and $b = j - 1$, $\chi_{la}(G(V_i, s)) = sn_i + t - r + j - 1$ for $1 \leq i \leq j - 1$, and $\chi_{la}(G(V_i, s)) = sn_i + t - r + j$ for $j \leq i \leq r$.

In particular, if $c_{r-1} \leq e < c_r$, then in $G(V_r, s)$, the condition on $e + sn_i$ is simplified to $s \geq 1$ for $n_r = 1$, and $s \geq 2$ is even for $n_r \geq 2$.

Proof. Note that G must contain a non-pendant vertex v such that $f^+(v) > e$. So $c_r > e \geq c_t$. Moreover, $uv \notin E(G)$ implies that $uv \notin E(G(V_i, s))$. For $1 \leq i \leq r$, define a bijection $g_i : E(G(V_i, s)) \rightarrow [1, e + sn_i]$ such that $g_i(e) = f(e)$ if $e \in E(G)$ and that

$$g_i(v_a w_{a,k}) = e + [k + \frac{(-1)^k}{2} - \frac{1}{2}]n_i - (-1)^k(a - \frac{1}{2}) + \frac{1}{2} = \begin{cases} e + (k-1)n_i + a & \text{if } k \text{ is odd,} \\ e + kn_i + 1 - a & \text{if } k \text{ is even.} \end{cases}$$

Therefore $g_i^+(v_a) = c_i + es + \frac{s}{2}(sn_i + 1) > e + sn_i$ when s is even; $g_i^+(v_a) = c_i + es + \frac{s}{2}(s+1) + a - 1 > e + sn_i$ when s is odd (in this case $n_i = 1$ and $a = 1$).

Observe that $g_i^+(v) = f^+(v)$ for each $v \notin V_i$ so that $c_1, \dots, c_{i-1}, c_{i+1}, \dots, c_t$ are vertex colors under g_i . Moreover, for $v_a \in V_i$, $f^+(v_a)$ is replaced by $g_i^+(v_a) > c_t$ for $1 \leq a \leq n_i$ and $\{g_i^+(w_{a,k}) = g_i(v_a w_{a,k}) \mid 1 \leq a \leq n_i, 1 \leq k \leq s\} = [e+1, e+sn_i]$.

- (1) Since $e < c_1$, in $G(V_i, s)$, we have $c_{r+1} < \dots < c_t < e+1 \leq c_1 < \dots < c_r \leq e+sn_i$ when $1 \leq i < r$, and $c_{r+1} < \dots < c_t < e+1 \leq c_1 < \dots < c_{r-1} \leq e+sn_i$ when $i = r$. So $(\{c_j \mid 1 \leq j \leq r\} \setminus \{c_i\}) \subset [e+1, e+sn_i]$ but all the c_{r+1}, \dots, c_t are not in $[e+1, e+sn_i]$. Since $c_i + es + \frac{s}{2}(n_i s + 1) > e + sn_i$, g_i is a local antimagic labeling with induced vertex color set $\{c_{r+1}, c_{r+2}, \dots, c_t\} \cup [e+1, e+sn_i] \cup \{c_i + es + \frac{s}{2}(n_i s + 1)\}$. Therefore, $\chi_{la}(G(V_i, s)) \leq sn_i + t - r + 1$. Since $G(V_i, s)$ contains at least $sn_i + t - r$ pendant vertices, by Theorem 1.2, $\chi_{la}(G(V_i, s)) \geq sn_i + t - r + 1$. Thus, $\chi_{la}(G(V_i, s)) = sn_i + t - r + 1$.
- (2) For $c_1 \leq e < c_2$, in $G(V_i, s)$, similar to the above case, $(\{c_j \mid 1 \leq j \leq r\} \setminus \{c_1, c_i\}) \subset [e+1, e+sn_i]$ but all the c_1, c_{r+1}, \dots, c_t are not in $[e+1, e+sn_i]$.

Suppose $i = 1$, then g_1 is a local antimagic labeling with induced vertex color set $\{c_{r+1}, c_{r+2}, \dots, c_t\} \cup [e+1, e+sn_1] \cup \{c_1 + es + \frac{s}{2}(n_1 s + 1)\}$. Thus, $\chi_{la}(G(V_1, s)) \leq sn_1 + t - r + 1$. By the same argument of (1), we get $\chi_{la}(G(V_1, s)) = sn_1 + t - r + 1$.

Suppose $2 \leq i \leq r$. In this case, $0 \leq b \leq 1$. Similar to the above case, g_i is a local antimagic labeling with induced vertex color set $\{c_{r+1}, c_{r+2}, \dots, c_t, c_1\} \cup [e+1, e+sn_i] \cup \{c_i + es + \frac{s}{2}(sn_i + 1)\}$. Thus, $\chi_{la}(G(V_i, s)) \leq sn_i + t - r + 2$. Combining with Theorem 1.2, we have $sn_i + t - r + b + 1 \leq \chi_{la}(G(V_i, s)) \leq sn_i + t - r + 2$. Moreover, $e = c_1$ implies that $b = 1$ so $\chi_{la}(G(V_i, s)) = sn_i + t - r + 2$.

- (3) For $c_{j-1} \leq e < c_j$, $3 \leq j \leq r$, we have $0 \leq b \leq j - 1$. In $G(V_i, s)$, similar to above case, $\{c_j, \dots, c_r\} \subset [e+1, e+sn_i]$ but all the c_1, \dots, c_{j-1} are not in $[e+1, e+sn_i]$. Similar to the above case, if $1 \leq i \leq j - 1$, then g_i is a local antimagic labeling with induced vertex color set $\{c_k \mid r+1 \leq k \leq t\} \cup (\{c_k \mid 1 \leq k \leq j-1\} \setminus \{c_i\}) \cup [e+1, e+sn_i] \cup \{c_i + es + \frac{s}{2}(n_i s + 1)\}$ so that $\chi_{la}(G(V_i, s)) \leq sn_i + t - r + j - 1$. If $j \leq i \leq r$, then the induced vertex color set is $\{c_k \mid r+1 \leq k \leq t\} \cup \{c_k \mid 1 \leq k \leq j-1\} \cup [e+1, e+sn_i] \cup \{c_i + es + \frac{s}{2}(n_i s + 1)\}$ so that $\chi_{la}(G(V_i, s)) \leq sn_i + t - r + j$.

Note that for $1 \leq i \leq j - 1$, if V_i has a pendant vertex, then $G(V_i, s)$ has $sn_i + t - r + b - 1$ pendant vertices, otherwise $G(V_i, s)$ has $sn_i + t - r + b$ pendant vertices. For $j \leq i \leq t$, $G(V_i, s)$ also has $sn_i + t - r + b$ pendant vertices. Combining with Theorem 1.2, we have the conclusion. Moreover, if $c_{j-1} = e$ and $b = j - 1$, only case (a) exists for $1 \leq i \leq j - 1$ so that $\chi_{la}(G(V_i, s)) = sn_i + t - r + j - 1$, and $\chi_{la}(G(V_i, s)) = sn_i + t - r + j$ for $j \leq i \leq r$.

In particular, if $c_{r-1} \leq e < c_r$, we know the condition $e + sn_r \geq c_{r-1}$ always hold and can be omitted. \square

Example 2.1.

- (1) In [5, Theorem 5], the authors proved that every $G = W_{4k}, k \geq 1$ with $e = 8k$ edges admits a local antimagic labeling with $\chi_{la}(W_{4k}) = 3$ such that for $k \geq 2$, $e < c_1 = 9k + 2 < c_2 = 11k + 1 < c_3 = 2k(12k + 1)$, while W_4 has $c_1 = 11, c_2 = 15, c_3 = 20$. Moreover, $n_1 = n_2 = 2k, n_3 = 1$. Thus, $r = t = 3$. Suppose $k \geq 2$, we can add $s \geq 12k - 3$ (s even) pendant edges to each vertex in $V_i, i = 1$ or 2 , and label them accordingly. We can also add $s \geq 3k + 1$ pendant edges to the vertex in V_3 and label them accordingly. By Theorem 2.1, we get $W_{4k}(V_i, s)$ that has $\chi_{la} = sn_i + 1$ for the respective s and n_i . We can also add edges to W_4 similarly.
- (2) The right graph under Theorem 1.2, say G , has $e = 7$ with $t = r = 2$; $c_1 = 7, c_2 = 14$ and $n_1 = 4, n_2 = 2$. Thus, $\chi_{la}(G(V_1, s)) = 4s + 1$. In particular, since $c_1 = e$ and $b = 1$, we also can get $\chi_{la}(G(V_2, s)) = 2s + 2$.
- (3) The left graph under Theorem 1.2, say H , has $e = 5$ with $t = r = 3$; $c_1 = 4 < c_2 = e < c_3$ and $b = 2$. Thus, $\chi_{la}H(V_3, s) = s + 3$ for $s \geq 1$.


By a similar argument for Theorem 2.1, we can get the following theorem.

Theorem 2.2. Suppose $G \not\cong K_{1,e}$ and $G(V_i, s)$ are as defined above with $r < t$ and $e + s \geq c_r$. Let $r + 1 \leq i \leq t$.

- (1) If $e < c_1$, then $\chi_{la}(G(V_i, s)) = s + t - r$.
- (2) If $c_{j-1} \leq e < c_j$ for $2 \leq j \leq r$, then $s + t - r + b \leq \chi_{la}(G(V_i, s)) \leq s + t - r + j - 1$. Moreover, if $b = j - 1$, then $\chi_{la}(G(V_i, s)) = s + t - r + j - 1$.

Example 2.2.

- (1) It is easy to verify that the graph G below has $\chi_{la}(G) = 9$ with $e = 9 < c_1 = 10, c_2 = 20, c_3 = 25$ and $r = 3 < t = 9$. We may apply Theorem 2.2 (1) accordingly.



- (2) We can add $k \geq 1$ pendant edges to the degree 4 vertex of the left graph under Theorem 1.2 and label the edges by 6 to $5 + k$ bijectively to get a graph, say G , with $k + 2$ pendant vertices. Clearly, $r = 3 < t = k + 3$ with $c_1 = 4, c_2 = 5 < e = k + 5 < c_3 = (k + 3)(k + 8)/2$, $c_i = i + 2$ for $i = 4, \dots, k + 2$, $j = 3$ and $b = j - 1$. By Theorem 2.2 (2), we know $\chi_{la}(G(V_i, s)) = s + k + 2$ for $s \geq (k + 2)(k + 7)/2$.

Suppose G is a graph containing $k \geq 1$ pendant vertices with $\chi_{la}(G) = k + 1$. Keeping the notation defined before Theorem 2.1, we have $t = k + 1$. Then there is only one independent set, which is V_r , containing no pendant vertex. Clearly, $r \geq 1$. So we have $c_1 < \dots < c_{r-1} \leq e < c_r$ and $c_{r+1} < \dots < c_{k+1} \leq e$ for $r \geq 2$, whereas for $r = 1$, $G \cong K_{1,k}, k \geq 2$ with $c_1 = k(k + 1)/2$ and $c_i = i - 1$ for $2 \leq i \leq k + 1$.

Corollary 2.3. Keep all notation defined before Theorem 2.1. Suppose G has k pendant vertices and $\chi_{la}(G) = k + 1$. Let

$$e + sn_i \geq \begin{cases} c_r & \text{for } i \in [1, k + 1] \setminus \{r\} \\ c_{r-1} & \text{for } i = r \end{cases}$$

such that $s \geq 1$ if $n_i = 1$, and $s \geq 2$ is even if $n_i \geq 2$, then $G(V_r, s)$ has $sn_r + k - 1$ pendant vertices with $\chi_{la}(G(V_r, s)) = sn_r + k + 1$, whereas $G(V_i, s)$ has $sn_i + k - 1$ pendant vertices with $\chi_{la}(G(V_i, s)) = sn_i + k$ when $i \in [1, k + 1] \setminus \{r\}$.

Proof. Suppose $r = 1$. Recall that $e = k$ and $c_1 = k(k+1)/2$. Since $G(V_1, s) \cong K_{1,k+s}$, we only consider $G(V_i, s)$, $2 \leq i \leq k+1$. By labeling the s added edges of $G(V_i, s)$ by $k+1$ to $k+s$ bijectively, c_i is now replaced by $i-1+ks+s(s+1)/2 > k+s \geq c_1 > c_{k+1} > \dots > c_2$. Thus, $G(V_i, s)$ now admits a local antimagic labeling with vertex color set $[1, k+s] \setminus \{i-1\} \cup \{i-1+ks+s(s+1)/2\}$. Therefore, $\chi_{la}(G(V_i, s)) \leq s+k$. Since $G(V_i, s)$ has $s+k-1$ pendant vertices, by Theorem 1.2, the equality holds.

Consider $r \geq 2$. Suppose $i = r$, then $G(V_r, s)$ has sn_r+k pendant vertices so that $\chi_{la}(G(V_r, s)) \geq sn_r+k+1$. By a labeling g_i as in the proof of Theorem 2.1, we know g_i is a local antimagic labeling with induced vertex color set $\{c_1, c_2, \dots, c_{r-1}, c_{r+1}, \dots, c_{k+1}\} \cup [e+1, e+sn_r] \cup \{g_i^+(v) \mid v \in V_r\}$ of size sn_r+k+1 . By Theorem 1.2, we have $\chi_{la}(G(V_r, s)) = sn_r+k+1$.

Suppose $i \in [1, k+1] \setminus \{r\}$, then $G(V_i, s)$ has sn_i+k-1 pendant vertices so that $\chi_{la}(G(V_i, s)) \geq sn_i+k$. By a labeling g_r as in the proof of Theorem 2.1, we know g_r is a local antimagic labeling with induced vertex color set $(\{c_1, c_2, \dots, c_{k+1}\} \setminus \{c_i, c_r\}) \cup [e+1, e+sn_i] \cup \{g_r^+(v) \mid v \in V_i\}$ of size sn_r+k . By Theorem 1.2, we have $\chi_{la}(G(V_r, s)) = sn_r+k$. \square

Example 2.3. Consider W_4 under Example 2.1. We have $W_4(V_3, 12)$ with $k = 12$, $\chi_{la}(W_4(V_3, 12)) = 13$ and $c_1 = 11, c_2 = 15 < e = 20 < c_3 = 194$. Moreover, $P_n, n \geq 3$ and $K_{1,k}$, is a tree with $k \geq 2$ pendant vertices and $\chi_{la} = k+1$. We may apply Corollary 2.3 accordingly.

Thus, we get the following.

Theorem 2.4. For $k \geq 1$, there exist infinitely many graphs G with $k \geq \chi(G) - 1 \geq 1$ pendant vertices and $\chi_{la}(G) = k+1$.

Let $G - e$ be the graph G with an edge e deleted. In [6, Lemmas 2.2-2.4], the authors obtained sufficient conditions for $\chi_{la}(G) = \chi_{la}(G - e)$. We note that these lemmas may be applied to $G(V_i, s)$ if all vertices in each $V_j, 1 \leq j \leq t$, are of same degree like $W_{4k}(V_i, s)$ in Example 2.1.

3 Existing results and open problems

In [2], [5, Theorem 7 and Theorem 9] and [7, Theorems 2.4-2.6, 2.9, 3.1, Lemma 2.10 and Section 3], the authors also determined the exact value of $\chi_{la}(G)$ for many families of G with pendant vertices. Particularly, they showed that there are infinitely many caterpillars of k -pendant vertices with $\chi_{la} = k+1$ or $\chi_{la} \geq k+2$. Note that [7, Theorem 2.6] corrected Theorem 2.2 in [8]. A lobster is a tree such that the removal of its pendant vertices resulted in a caterpillar. Note that the graph $Sp(2^{[n]})$ is also a lobster. We end this paper with the followings.

Conjecture 3.1. Every tree T_k , other than certain caterpillars, spiders and lobsters, with $k \geq 2$ pendant vertices has $\chi_{la}(T_k) = k+1$.

Problem 3.1. Characterize all graphs G with $k \geq \chi(G) - 1 \geq 1$ pendant vertex(s) and $\chi_{la}(G) = k+1$.

References

- [1] S. Arumugam, K. Premalatha, M. Bača and A. Semaničová-Feňovčíková, Local antimagic vertex coloring of a graph, *Graphs and Combin.*, **33** (2017) 275–285.
- [2] S. Arumugam, Y.C. Lee, K. Premalatha, T.M. Wang, On local antimagic vertex coloring for corona products of graphs, (2018) arXiv:1808.04956v1.
- [3] R. Frucht and F. Harary, On the corona of two graphs, *Aequationes Math.* **4** (1970), 322-325.
- [4] J. Haslegrave, Proof of a local antimagic conjecture, *Discrete Math. Theor. Comput. Sci.* **20**(1) (2018). <https://doi.org/10.23638/DMTCS-20-1-18>
- [5] G.C. Lau, H.K. Ng and W.C. Shiu, Affirmative solutions on local antimagic chromatic number, (2019) arXiv:1805.02886.

- [6] G.C. Lau, W.C. Shiu and H.K. Ng, On local antimagic chromatic number of cycle-related join graphs, (2018) Discuss. Math. Graph Theory, accepted.
- [7] G.C. Lau, W.C. Shiu and H.K. Ng, On local antimagic chromatic number of graphs with cut-vertices, (2019) arXiv:1805.04801.
- [8] Nuris Hisan Nazula, Slamin and Dafik, Local antimagic vertex coloring of unicyclic graphs, *Indonesian Journal of Combinatorics* **2(1)** (2018), 30-34.