

Connectivity for an unlabelled bridge-addable graph class

Colin McDiarmid
Oxford University

October 29, 2021

Call a class \mathcal{A} of graphs *bridge-addable* if, whenever a graph G in \mathcal{A} has vertices u and v in distinct components, then the graph $G + uv$ (obtained by adding the edge uv) is also in \mathcal{A} . Thus for example the class \mathcal{F} of forests is bridge-addable, as is the class of planar graphs.

Let \mathcal{A}_n denote the set of graphs in \mathcal{A} on vertex set $\{1, \dots, n\}$ (so these are labelled graphs), and similarly for other graph classes. Also, let us use the notation $R_n \in_u \mathcal{A}$ to indicate that the random graph R_n is sampled uniformly from \mathcal{A}_n (and assume implicitly that \mathcal{A}_n is non-empty). Thus

$$\mathbb{P}(R_n \text{ is connected}) = |\mathcal{C}_n| / |\mathcal{A}_n|$$

where \mathcal{C} is the class of connected graphs in \mathcal{A} . The following elementary but useful inequality appeared in McDiarmid, Steger and Welsh [12, 13]. If \mathcal{A} is bridge-addable and $R_n \in_u \mathcal{A}$ then

$$\mathbb{P}(R_n \text{ is connected}) \geq e^{-1} \approx 0.3679 \quad \text{for each } n. \quad (1)$$

If \mathcal{T} denotes the class of trees then $|\mathcal{T}_n|/|\mathcal{F}_n| \rightarrow e^{-\frac{1}{2}}$ as $n \rightarrow \infty$, by a result of Rényi [17] in 1959. It was conjectured in [13] in 2006 that the class of forests is the ‘least connected’ bridge-addable class; and in particular that, if \mathcal{A} is bridge-addable and $R_n \in_u \mathcal{A}$, then

$$\liminf_n \mathbb{P}(R_n \text{ is connected}) \geq e^{-\frac{1}{2}} \approx 0.6065. \quad (2)$$

After several partial results towards proving this conjecture, see [1, 2, 3, 7, 16], the full result was proved by Guillaume Chapuy and Guillem Perarnau [4]. See also [9, 10, 14, 15].

Here we are interested in **unlabelled** graphs, where by contrast little seems to be known. Let $\tilde{\mathcal{A}}_n$ be the set of unlabelled graphs in \mathcal{A}_n , and similarly for other graph classes. We may identify an unlabelled graph as an isomorphism class of labelled graphs. We use the notation $\tilde{R}_n \in_u \tilde{\mathcal{A}}$ to indicate that the random unlabelled graph \tilde{R}_n is sampled uniformly from $\tilde{\mathcal{A}}_n$. Of course, as before

$$\mathbb{P}(\tilde{R}_n \text{ is connected}) = |\tilde{\mathcal{C}}_n| / |\tilde{\mathcal{A}}_n|.$$

We make two conjectures about the probability of being connected, corresponding to (1) and (2) above. The first seems to ask for rather little (though see Theorem 5 below).

Conjecture 1. *There is a $\delta > 0$ such that, if the graph class \mathcal{A} is bridge-addable and $\tilde{R}_n \in_u \tilde{\mathcal{A}}$, then*

$$\mathbb{P}(\tilde{R}_n \text{ is connected}) \geq \delta \quad \text{for each } n. \quad (3)$$

For trees and forests we have

$$|\tilde{\mathcal{T}}_n|/|\tilde{\mathcal{F}}_n| \rightarrow \tau \approx e^{-0.5226} \approx 0.5930 \quad \text{as } n \rightarrow \infty,$$

see for example the first line of table 3 in [6]. The second conjecture is more speculative. Call the class \mathcal{A} of graphs *decomposable* when a graph is in the class if and only if each component is. Examples include the class \mathcal{F} of forests and the class of planar graphs.

Conjecture 2. *If the graph class \mathcal{A} is bridge-addable and decomposable, and $\tilde{R}_n \in_u \tilde{\mathcal{A}}$, then*

$$\liminf_n \mathbb{P}(\tilde{R}_n \text{ is connected}) \geq \tau. \quad (4)$$

The *fragment size* $\text{frag}(G)$ of a graph G is the number of vertices of G less the maximum number of vertices in a component. In the labelled case, if the graph class \mathcal{A} is bridge-addable and $R_n \in_u \mathcal{A}$ then ([9], see [8] for an earlier version)

$$\mathbb{E}[\text{frag}(R_n)] < 2.$$

Is there a corresponding result in the unlabelled case? Here is an awkward example.

Fix an integer $k \geq 3$, and let \mathcal{A} consist of all graphs G such that deleting any bridges from G yields a disjoint union of cycles each of length at least k .

Clearly \mathcal{A} is bridge-addable and decomposable. Let $n = 2k$, and let $\tilde{R}_n \in_u \tilde{\mathcal{A}}$. Then $\mathbb{P}(\tilde{R}_n \text{ is connected}) = \frac{1}{2}$, which is satisfactory (though less than τ); but $\mathbb{E}[\text{frag}(\tilde{R}_n)] = n/4$, which is bad!

Conjecture 3. *For each graph class \mathcal{A} which is bridge-addable and decomposable, there is a constant $c_{\mathcal{A}}$ such that, for each positive integer n , if $\tilde{R}_n \in_u \tilde{\mathcal{A}}$ then*

$$\mathbb{E}[\text{frag}(\tilde{R}_n)] \leq c_{\mathcal{A}}.$$

The fourth and final conjecture is a little speculative (as was the second).

Conjecture 4. *There is a constant c such that, for each graph class \mathcal{A} which is bridge-addable and decomposable, for $\tilde{R}_n \in_u \tilde{\mathcal{A}}$*

$$\limsup_{n \rightarrow \infty} \mathbb{E}[\text{frag}(\tilde{R}_n)] \leq c.$$

Let us return to Conjecture 1, but lower our sights. We can quickly prove a weakened version of the conjecture (with δ replaced by $1/2n$) which is still strong enough to be useful (see the proof of Theorem 4 in [11]).

Theorem 5. *Let \mathcal{A} be a bridge-addable class of graphs, let \mathcal{C} be the class of connected graphs in \mathcal{A} , and let \mathcal{C}^\bullet be the set of vertex-rooted graphs in \mathcal{C} . Then*

$$|\tilde{\mathcal{A}}_n| \leq 2 |\tilde{\mathcal{C}}_n^\bullet| \leq 2n |\tilde{\mathcal{C}}_n| \quad \text{for each } n \in \mathbb{N}. \quad (5)$$

Proof. From each graph $G \in \tilde{\mathcal{A}}_n$ we shall construct a vertex-rooted connected graph $H^v \in \tilde{\mathcal{C}}_n^\bullet$ and $b \in \{0, 1\}$, such that from H^v and b we can determine G . It will follow that $|\tilde{\mathcal{A}}_n| \leq 2 |\tilde{\mathcal{C}}_n^\bullet|$, and since $|\tilde{\mathcal{C}}_n^\bullet| \leq n |\tilde{\mathcal{C}}_n|$ the proof will be complete.

Now for the construction. Observe first that every n -vertex graph has at most $n - 1$ bridges, so there is a vertex incident to at most one bridge. Let $G \in \tilde{\mathcal{A}}_n$. Let C be a smallest component in G , and let v be a vertex in C incident to $b \in \{0, 1\}$ bridges. Add an edge between v and each component of G other than C , forming the graph H . Finally root H at v , and output the rooted graph H^v together with b .

Let B be the set of bridges in H incident to v . Then B consists of b edges from the original graph G and any edges added when forming H . For each edge $e \in B$ let $n(e)$ be the number of vertices in the component of $H \setminus e$ not

containing v . If $b = 1$ and e is the edge in B from G then $n(e) \leq v(C) - 1$. Also $n(e') \geq v(C)$ for any other edge $e' \in B$. Thus, given H^v and b , we can identify these added edges and delete them to recover G , as required. \square

This note grew from an open problem presented at Oberwolfach in May 2014. It was updated and presented as an open problem at the First Armenian Workshop on Graphs, Combinatorics, Probability, in Dzoraget, Armenia, in June 2019. Now it has been amplified by adding Theorem 5 (and some further references).

Barbados?

References

- [1] L. Addario-Berry, C. McDiarmid and B. Reed, Connectivity for bridge-addable monotone graph classes, *Combinatorics, Probability and Computing* **21** (2012) 803 – 815.
- [2] P. Balister, B. Bollobás and S. Gerke, Connectivity of addable graph classes. *J. Combin. Th. B* **98** (2008) 577 – 584.
- [3] P. Balister, B. Bollobás and S. Gerke, Connectivity of random addable graphs, *Proc. ICDM 2008* No 13 (2010) 127 – 134.
- [4] G. Chapuy and G. Perarnau, Connectivity in bridge-addable graph classes: the McDiarmid-Steger-Welsh conjecture, *Journal of Combinatorial Theory B* **136** (2019) 44 – 71.
- [5] G. Chapuy and G. Perarnau, Local convergence and stability of tight bridge-addable graph classes, *Canadian Journal of Mathematics*, to appear. arXiv:1609.03974v1
- [6] M. Kang and C. McDiarmid, Random unlabelled graphs containing few disjoint cycles, *Random Structures and Algorithms* **38** (2011) 174 – 204.
- [7] M. Kang and K. Panagiotou, On the connectivity of random graphs from addable classes, *J. Combinatorial Theory B* **103** (2013) 306 – 312.
- [8] C. McDiarmid, Random graphs from a minor-closed class, *Combinatorics, Probability and Computing* **18** (2009) 583-599.

- [9] C. McDiarmid, Connectivity for random graphs from a weighted bridge-addable class, *Electronic J Combinatorics* **19**(4) (2012) P53.
- [10] C. McDiarmid, Connectivity for bridge-alterable graph classes, *European J. Comb.* **56** (2016) 33 – 39.
- [11] C. McDiarmid and S. Saller, Classes of graphs embeddable in non-constant surfaces, in preparation, 2020.
- [12] C. McDiarmid, A. Steger and D. Welsh, Random planar graphs, *J. Combinatorial Theory B* **93** (2005) 187 – 206.
- [13] C. McDiarmid, A. Steger and D. Welsh, Random graphs from planar and other addable classes, *Topics in Discrete Mathematics* (M. Klazar, J. Kratochvil, M. Loeb, J. Matousek, R. Thomas, P. Valtr, Eds.), Algorithms and Combinatorics 26, Springer, 2006, 231 – 246.
- [14] C. McDiarmid and K. Weller, Relatively bridge-addable classes of graphs, in LATIN 2014 : Theoretical Informatics, Proceedings of the 11th Latin American Symposium, Montevideo, Uruguay, 2014 (editors Alberto Pardo and Alfredo Viola), Springer *LNCS* **8392** (2014) 391 – 398.
- [15] C. McDiarmid and K. Weller, Bridge-addability, edge-expansion and connectivity, *Combinatorics, Probability and Computing* **26** (2017) 697 – 719.
- [16] S. Norin, Connectivity of addable classes of forests, private communication, 2013.
- [17] A. Rényi, Some remarks on the theory of trees, *Publications of the Mathematical Institute of the Hungarian Academy of Sciences* **4** (1959) 73 – 85.