On the basic properties of GC_n sets

Hakop Hakopian, Navasard Vardanyan

Abstract

In this paper the simplest *n*-correct sets in the plane - GC_n sets are studied. An *n*-correct node set \mathcal{X} is called GC_n set if the fundamental polynomial of each node is a product of *n* linear factors. We say that a node uses a line if the line is a factor of the fundamental polynomial of this node. A line is called *k*-node line if it passes through exactly *k* nodes of \mathcal{X} . At most n + 1 nodes can be collinear in any GC_n set and an (n + 1)-node line is called a maximal line. The Gasca-Maeztu conjecture (1982) states that for every GC_n set there exists at least one maximal line. Until now the conjecture has been proved only for the cases $n \leq 5$.

Here, for a line ℓ we introduce and study the concept of ℓ -lowering of the set \mathcal{X} and define so called proper lines. We also provide refinements of several basic properties of GC_n sets regarding the maximal lines, *n*-node lines, the used lines, as well as the subset of nodes that use a given line.

Key words: Polynomial interpolation, the Gasca-Maeztu conjecture, n-correct set, GC_n set, maximal line, proper line.

Mathematics Subject Classification (2010): 41A05, 41A63.

1 Introduction

An *n*-correct set \mathcal{X} in the plane is a node set for which the Lagrange interpolation problem with bivariate polynomials of total degree not exceeding n is unisolvent. A line is called *k*-node line if it passes through exactly k nodes of \mathcal{X} . It is a simple fact that at most n + 1 nodes can be collinear in an *n*-correct set. An (n + 1)-node line is called a maximal line. In *n*-correct node sets with geometric characterization: GC_n sets, introduced by Chung and Yao [10], the fundamental polynomial of each node is a product of n linear factors. The presence of simple fundamental polynomials, in view of the Lagrange formula, means a simple formula for interpolation polynomials. Thus, from both theoretical and practical points of view, the GC_n sets are the simplest *n*-correct sets. Consequently, the study of the sets GC_n is a topical problem in the theory of bivariate polynomial interpolation.

The conjecture of M. Gasca and J. I. Maeztu [11] (GM conjecture) states that every GC_n set has at least one maximal line. Until now the conjecture has been proved only for the cases $n \leq 5$ (see [2] and [12]). The GM conjecture is very important because if it proves to be true then a nice classification of GC_n sets takes place (see the forthcoming Sections 3-5).

We say that a node uses a line if the line is a factor of the fundamental polynomial of this node. The problem of finding the fundamental polynomial of a node in the GC_n sets means finding all the lines used by the node. To solve this important problem, we can look at it more generally. Namely, given a line and a node, find out whether the line is used by the node. Therefore, the problem under consideration is equivalent to the problem of finding all nodes that use a given line.

Denote the subset of nodes of \mathcal{X} that use the line ℓ by \mathcal{X}^{ℓ} . Let us mention that

$$\mathcal{X}^{\ell} = \mathcal{X} \setminus \ell \text{ if } \ell \text{ is a maximal line.}$$
(1.1)

Next, let us formulate a recent result regarding the set \mathcal{X}^{ℓ} for the general ℓ (see also [1], Conjecture 3.7).

Theorem 1.1 ([15], Theorem 3.1). Let \mathcal{X} be a GC_n set, and ℓ be a k-node line, $k \geq 2$. Assume that GM Conjecture holds for all degrees up to n. Then we have that

$$\mathcal{X}^{\ell} = \emptyset, \text{ or}$$

 $\mathcal{X}^{\ell} \text{ is a } GC_{s-2} \text{ subset of } \mathcal{X}, \text{ hence } |\mathcal{X}_{\ell}| = \binom{s}{2},$ (1.2)

where $k - \delta \leq s \leq k$ and $\delta = n + 1 - k$.

Moreover, [15] established the following algorithm for retrieving \mathcal{X}^{ℓ} . At the first step one obtains a node set \mathcal{X}_1 by removing the nodes in a maximal line of \mathcal{X} , which does not intersect the line ℓ at a node of \mathcal{X} or by removing the nodes in a pair of maximal lines of \mathcal{X} which are concurrent together with ℓ , i.e., the two maximal lines and ℓ intersect at a single point. Thus, in view of the forthcoming Proposition 2.6, \mathcal{X}_1 is a GC_{n-1} or GC_{n-2} set. Second step is similar to the first one. The only difference is that instead of the maximal lines of \mathcal{X} now the maximal lines of \mathcal{X}_1 are used. By continuing in the same way, at the last step ω , the line ℓ becomes a maximal line in a GC_{s-1} set \mathcal{X}_{ω} , and hence, according to (1.1), one gets

$$\mathcal{X}^{\ell} = \mathcal{X}_{\omega} \setminus \ell.$$

In this paper, by assuming that GM conjecture is true, we characterize the lines ℓ for which the above-described algorithm can be simplified and done by using only the original maximal lines of \mathcal{X} .

More specifically, we introduce the concept of ℓ -lowering of a GC_n set \mathcal{X} , and denote it by $\hat{\mathcal{X}} := \hat{\mathcal{X}}(\ell)$. This set plays an important role here. We get $\hat{\mathcal{X}}$ by removing, at a single step, the nodes in all maximal lines of \mathcal{X} which do not intersect the line ℓ at a node of \mathcal{X} and by removing the nodes in all pairs of maximal lines of \mathcal{X} which are concurrent together with ℓ . Then we specify cases when ℓ is a maximal line in the obtained GC set $\hat{\mathcal{X}}$ and hence

$$\mathcal{X}^{\ell} = \hat{\mathcal{X}} \setminus \ell.$$

We call such lines *proper* lines. We prove in the forthcoming Theorem 7.1 that all lines ℓ used by more than 3 nodes are either maximal or proper, provided that the number of maximal lines of \mathcal{X} differs from three. Then we prove that in the case of arbitrary non-proper line ℓ one needs to complete only one or two steps of the maximal lines removal in $\hat{\mathcal{X}}$ till ℓ becomes a maximal line. It means that in the general case the above-described algorithm can be carried out by using at most only two steps of the maximal lines removal where the lines are not original maximal lines of \mathcal{X} .

It is worth mentioning that if the number of maximal lines of \mathcal{X} equals three, in which case \mathcal{X} is called *generalized principal lattice*, we have a simple description of the set \mathcal{X}^{ℓ} (see the forthcoming formula (6.4)).

Then, in contrast to the ambiguity present in the formula (1.2), we determine the exact number of the nodes that use a given line, provided that GM conjecture is true. Let us call a node a 2_m -node in \mathcal{X} if it is a point of intersection of two maximal lines. In the forthcoming Theorem 7.4 we prove that a k-node line ℓ either is not used at all, or is used by exactly $\binom{k-r-\hat{r}}{2}$ nodes of \mathcal{X} , where r is the number of 2_m -nodes in $\ell \cap \mathcal{X}$ and \hat{r} is the number of 2_m -nodes in $\ell \cap \mathcal{X}$. Moreover, we have that always $\hat{r} \leq 2$. On other hand we have that $\hat{r} = 0$ if $\#\mathcal{X}^{\ell} > 3$. Furthermore, for each 2_m -node in $\ell \cap \hat{\mathcal{X}}$ we prove that one of the two maximal lines in $\hat{\mathcal{X}}$ to which it belongs is a maximal line in \mathcal{X} and another is a proper line in \mathcal{X} .

Next, we study the set of all *n*-node lines of GC_n set \mathcal{X} denoted by $N(\mathcal{X})$.

Also, in this paper we improve several known properties of GC_n sets.

Let us mention that Carnicer and Gasca started the investigation of the set \mathcal{X}^{ℓ} and proved that a k-node line ℓ can be used by at most $\binom{k}{2}$ nodes of a GC_n set \mathcal{X} and in addition there are no k collinear nodes that use ℓ , provided that GM conjecture is true (see [5], Theorem 4.5). Note that these statements follow readily from Theorem 1.1.

2 GC_n sets and the Gasca-Maeztu conjecture

Let Π_n be the space of bivariate polynomials of total degree at most n :

$$p(x,y) = \sum_{i+j \le n} c_{ij} x^i y^j.$$

We have that $N := \dim \prod_n = \binom{n+2}{2}$. Let \mathcal{X} be a set of N distinct nodes: $\mathcal{X} = \{(x_1, y_1), (x_2, y_2), \dots, (x_N, y_N)\}.$ **Definition 2.1.** A set of nodes \mathcal{X} , $\#\mathcal{X} = N$, is called *n*-correct if for any given numbers $c_1c_2 \ldots c_N$ there exists a unique polynomial $p \in \prod_n$ satisfying the following conditions

$$p(x_i, y_i) = c_i, \quad i = 1, 2, \dots, N.$$
 (2.1)

A polynomial $p \in \Pi_n$ is called an *n*-fundamental polynomial of a node $A \in \mathcal{X}$, if p(A) = 1 and $p(B) = 0 \forall B \in \mathcal{X} \setminus \{A\}$. We denote this polynomial by $p_{A,\mathcal{X}}^*$.

Definition 2.2. Let \mathcal{X} be an *n*-correct set. We say that a node $A \in \mathcal{X}$ uses a line $\ell \in \Pi_1$, if $p_{A,\mathcal{X}}^* = \ell q$, where $q \in \Pi_{n-1}$.

The following proposition is well-known (see, e.g., [12] Proposition 1.5):

Proposition 2.3. Suppose that a polynomial $p \in \Pi_n$ vanishes at n+1 points of a line ℓ . Then we have that $p = \ell r$, where $r \in \Pi_{n-1}$.

Thus at most n + 1 nodes of an *n*-correct set \mathcal{X} can be collinear. A line λ passing through n + 1 nodes of the set \mathcal{X} is called a *maximal line*. Note that from Proposition 2.3 we readily get (1.1).

Below the basic properties of maximal lines are given:

Proposition 2.4 ([3], Prop. 2.1). Let \mathcal{X} be an *n*-correct set. Then

- (i) any two maximal lines of \mathcal{X} intersect necessarily at a node of \mathcal{X} ;
- (ii) any three maximal lines of \mathcal{X} cannot be concurrent;
- (iii) \mathcal{X} can have at most n+2 maximal lines.

Now let us consider a special type of n-correct sets satisfying a geometric characterization (GC) property introduced by K.C. Chung and T.H. Yao:

Definition 2.5 ([10]). An *n*-correct set \mathcal{X} is called a GC_n set (or GC set) if the *n*-fundamental polynomial of each node $A \in \mathcal{X}$ is a product of *n* linear factors.

Thus, each node of a GC_n set uses exactly n lines.

Proposition 2.6 ([5], Prop. 2.3). Let λ be a maximal line of a GC_n set \mathcal{X} . Then the set $\mathcal{X} \setminus \lambda$ is a GC_{n-1} set. Also any maximal line of \mathcal{X} different from λ is a maximal line of $\mathcal{X} \setminus \lambda$.

Next we present the Gasca-Maeztu (GM) conjecture:

Conjecture 2.7 ([11], Sect. 5). For any GC_n set there exists at least one maximal line.

Till now, this conjecture has been confirmed for the degrees $n \leq 5$ (see [2], [12]). For a generalization of the Gasca-Maeztu conjecture to maximal curves see [13].

Denote by $M(\mathcal{X})$ the set of maximal lines of the node set \mathcal{X} .

The concept of the defect introduced by Carnicer and Gasca in [3] is an important characteristic of GC_n sets.

Definition 2.8 ([3]). The *defect* of an *n*-correct set \mathcal{X} is the number $def(\mathcal{X}) := n + 2 - \#M(\mathcal{X}).$

In view of Proposition 2.4 we have that $0 \leq \operatorname{def}(\mathcal{X}) \leq n+2$.

Proposition 2.9 ([5], Crl. 3.5). Let λ be a maximal line of a GC_n set \mathcal{X} such that $\#M(\mathcal{X} \setminus \lambda) \geq 3$. Then we have that

 $def(\mathcal{X} \setminus \lambda) = def(\mathcal{X}) \quad or \quad def(\mathcal{X}) - 1.$

This equality means that $\#M(\mathcal{X} \setminus \lambda) = \#M(\mathcal{X}) - 1$ or $\#M(\mathcal{X})$. In view of Proposition 2.6 all $\#M(\mathcal{X}) - 1$ maximal lines of \mathcal{X} different from λ belong to $M(\mathcal{X} \setminus \lambda)$. Thus there can be at most one *newly emerged* maximal line of $\mathcal{X} \setminus \lambda$.

In the sequel we will use the following

Lemma 2.10 ([7], Lemma 3.4). Suppose that the Gasca-Maeztu conjecture is true for all degrees up to n. Suppose also that \mathcal{X} is a GC_n set with exactly three maximal lines and λ is a maximal line. Then, the GC_{n-1} set $\mathcal{X} \setminus \lambda$ also possesses exactly three maximal lines, hence $defect(\mathcal{X} \setminus \lambda) = defect(\mathcal{X}) - 1$.

Definition 2.11 ([4]). Given an *n*-correct set \mathcal{X} and a line ℓ , \mathcal{X}^{ℓ} is the subset of nodes of \mathcal{X} which use the line ℓ .

Let \mathcal{X} be an *n*-correct set, and ℓ be a line. Then

(i) a maximal line λ is called ℓ -disjoint if $\lambda \cap \ell \cap \mathcal{X} = \emptyset$;

(ii) two maximal lines λ', λ'' are called ℓ -adjoint if $\lambda' \cap \lambda'' \cap \ell \in \mathcal{X}$.

In view of Proposition 2.4 a maximal line ℓ has no ℓ -disjoint or ℓ -adjoint maximal lines.

The following two lemmas of Carnicer and Gasca play an important role in the sequel.

Lemma 2.12 ([5], Lemma 4.4). Let \mathcal{X} be an *n*-correct set and ℓ be a line with $\#(\ell \cap \mathcal{X}) \leq n$. Suppose also that a maximal line λ is ℓ -disjoint. Then we have that $\mathcal{X}^{\ell} = (\mathcal{X} \setminus \lambda)^{\ell}$.

The set $\mathcal{X} \setminus \lambda$ is called ℓ -disjoint reduction of \mathcal{X} .

Lemma 2.13 ([5], Proof of Thm. 4.5). Let \mathcal{X} be an *n*-correct set and ℓ be a line with $\#(\ell \cap \mathcal{X}) \leq n$. Suppose also that two maximal lines λ', λ'' are ℓ -adjoint. Then we have that $\mathcal{X}^{\ell} = [\mathcal{X} \setminus (\lambda' \cup \lambda'')]^{\ell}$. The set $\mathcal{X} \setminus (\lambda' \cup \lambda'')$ is called ℓ -adjoint reduction of \mathcal{X} .

Next, by the motivation of the above two lemmas, we introduce the concept of ℓ -lowering of a GC_n set.

Definition 2.14. Let \mathcal{X} be a GC_n set, ℓ be a k-node line, $2 \leq k \leq n+1$. We say that the set $\hat{\mathcal{X}} = \hat{\mathcal{X}}(\ell)$ is the ℓ -lowering of \mathcal{X} , and briefly denote this by $\mathcal{X} \downarrow_{\ell} \hat{\mathcal{X}}$, if

$$\hat{\mathcal{X}} = \mathcal{X} \setminus (\mathcal{U}_1 \cup \mathcal{U}_2),$$

where \mathcal{U}_1 is the union of the ℓ -disjoint maximal lines of \mathcal{X} , and \mathcal{U}_2 is the union of the (pairs of) ℓ -adjoint maximal lines of \mathcal{X} .

If ℓ is a maximal line then $\mathcal{U}_1 = \mathcal{U}_2 = \emptyset$ and hence $\hat{\mathcal{X}} = \mathcal{X}$. From Lemmas 2.12 and 2.13 we immediately get that

$$\mathcal{X}\downarrow_{\ell} \hat{\mathcal{X}} \Rightarrow \mathcal{X}^{\ell} = \hat{\mathcal{X}}^{\ell}.$$

Definition 2.15. A node $A \in \mathcal{X}$ is called k_m -node if it belongs to exactly k maximal lines.

In view of Proposition 2.4, (ii), there are 0_m , 1_m , and 2_m -nodes only. It is easily seen that $\ell \cap \hat{\mathcal{X}}(\ell)$ contains no 2_m -node of \mathcal{X} .

The following proposition will be used frequently in the sequel.

Proposition 2.16. Suppose that a line $\ell \notin M(\mathcal{X})$ passes through exactly $k \, 1_m$ -nodes of a GC_n set \mathcal{X} . Then the set $\hat{\mathcal{X}}(\ell)$ is a GC_s set with $s = def(\mathcal{X}) + k - 2$.

Proof. Suppose that a line ℓ passes through exactly $k \ 1_m$ -nodes. Then there are exactly $\#M(\mathcal{X})-k$ maximal lines which are either ℓ -disjoint or ℓ -adjoint. Hence, in view of Proposition 2.6, we readily get that the set $\hat{\mathcal{X}}(\ell)$ is a GC_s set with $s = n - [\#M(\mathcal{X}) - k] = def(\mathcal{X}) + k - 2$.

Definition 2.17. Let \mathcal{X} be a GC_n set, ℓ be a k-node line, $2 \leq k \leq n$ and $\mathcal{X} \downarrow_{\ell} \hat{\mathcal{X}}$. Then the line ℓ is called *proper* if it is a maximal line in the set $\hat{\mathcal{X}}$. The line ℓ is called *proper* (-r) if it becomes a maximal line after r steps of application of ℓ -disjoint or ℓ -adjoint reductions, in all, starting with $\hat{\mathcal{X}}$.

We call the ℓ -disjoint maximal line, or the union of the pair of ℓ -adjoint maximal lines, used at the above-mentioned kth step, the (-k) d/a (disjoint/adjoint) item, $k = 1, \ldots, r$.

In the forthcoming Theorem 7.1 we show that if $def(\mathcal{X}) \neq n-1$ then any used line, which is not maximal, is either proper, proper (-1), or (-2). Denote by $Pr(\mathcal{X})$ the set of proper lines of \mathcal{X} .

From (1.1) we immediately get that $\mathcal{X}^{\ell} = \hat{\mathcal{X}} \setminus \ell$ if $\ell \in Pr(\mathcal{X})$.

In Proposition 8.3 we show that $\#Pr(\mathcal{X}) \in \{0,3\}$ if $def(\mathcal{X}) \neq 1$, provided that the Gasca-Maeztu conjecture is true.



Figure 3.1: A Chang-Yao lattice, n = 3.

In the next three sections we will consider the results of Carnicer, Gasca, and Godés, concerning the classification of GC_n sets according to the defects of the sets. Let us present here the following

Theorem 2.18 ([9]). Let \mathcal{X} be a GC_n set. Assume that the GM Conjecture holds for all degrees up to n. Then $def(\mathcal{X}) \in \{0, 1, 2, 3, n - 1\}$.

Of course this implies that $\#M(\mathcal{X}) \in \{3, n-1, n, n+1, n+2\}$.

3 Sets of defect 0 and 1

Let us start with defect 0 sets, i.e., the Chung-Yao lattices. Let a set \mathcal{M} of n+2 lines be in general position, i.e., no two lines are parallel and no three lines are concurrent. Then the Chung-Yao lattice is defined as the set \mathcal{X} of all $\binom{n+2}{2}$ intersection points of these lines (see Fig. 3.1). We have that the n+2 lines of \mathcal{M} are maximal lines for \mathcal{X} . Any particular node here is 2_m -node, i.e., lies in exactly 2 maximal lines. Observe that the product of the remaining n maximal lines gives the fundamental polynomial of the particular node. Thus \mathcal{X} is a GC_n set. Let us mention that any n-correct set \mathcal{X} , with def(\mathcal{X}) = 0, i.e., $\#M(\mathcal{X}) = n+2$, in view of Proposition 2.4, (i) and (ii), forms a Chung-Yao lattice. Recall that there are no n-correct sets with more maximal lines (Proposition 2.4, (iii)).

Evidently the set of used lines in the Chung-Yao lattice coincides with the set of n + 2 maximal lines. For each maximal line ℓ , in view of (1.1), we have that $\mathcal{X}^{\ell} = \mathcal{X} \setminus \ell$, hence $\# \mathcal{X}^{\ell} = \binom{n+1}{2}$.

Thus the total number of line-usages equals:

 $(n+2)\binom{n+1}{2} = \frac{1}{2}(n+2)(n+1)n = n\binom{n+2}{2}.$

Next let us discuss defect 1 sets, i.e., the Carnicer-Gasca lattices. Let a set \mathcal{M} of n + 1 lines be in general position, $n \geq 2$. Then the Carnicer-Gasca lattice \mathcal{X} is defined as $\mathcal{X} := \mathcal{X}^{(2)} \cup \mathcal{X}^{(1)}$, where $\mathcal{X}^{(2)}$ is the set of all intersection nodes of these n + 1 lines, and $\mathcal{X}^{(1)}$ is a set of other n + 1non-collinear nodes, one in each line, to make the line maximal (see Fig. 3.2). We have that $\#\mathcal{X} = \binom{n+1}{2} + (n+1) = \binom{n+2}{2}$. It is easily seen that \mathcal{X} is a GC_n set and has exactly n + 1 maximal lines, i.e., the lines of \mathcal{M} . The set $\mathcal{X}^{(2)}$ consists of 2_m -nodes and the set $\mathcal{X}^{(1)}$ consists of 1_m -nodes. Let us



Figure 3.2: A Carnicer-Gasca lattice, n = 3.

mention that any *n*-correct set \mathcal{X} , with def(\mathcal{X}) = 1, i.e., $\#M(\mathcal{X}) = n + 1$, forms a Carnicer-Gasca lattice (see [3], Proposition 2.4).

It is easily seen that the set of used lines in the Carnicer-Gasca lattice consists of two classes:

1) The set of n + 1 maximal lines;

2) The set of lines passing through at least two 1_m -nodes.

Next we show that the lines of the class 2) are proper:

Proposition 3.1. Suppose that \mathcal{X} is a Carnicer-Gasca lattice and ℓ is a line passing through exactly $k \mid_m$ -nodes, where $k \geq 2$. Suppose also that $\mathcal{X} \downarrow_{\ell} \hat{\mathcal{X}}$. Then $\hat{\mathcal{X}}$ is a Chung-Yao lattice of degree k - 1, where ℓ is a maximal line. Hence $\mathcal{X}_{\ell} = \hat{\mathcal{X}} \setminus \ell$ and $\# \mathcal{X}^{\ell} = \binom{k}{2}$.

Proof. In view of Proposition 2.16 we have that $\hat{\mathcal{X}}$ is a GC_{k-1} set. On the other hand we have that the line ℓ , with $k \ 1_m$ -nodes, as well as the k maximal lines of \mathcal{X} intersecting the line ℓ at these nodes, in all k+1 lines, are maximal lines in $\hat{\mathcal{X}}$.

Denote by n_i the number of line usages in the class i), i = 1, 2. Then we have that the total number of line-usages equals:

 $n_1 + n_2 = (n+1)\binom{n+1}{2} + \binom{n+1}{2} = \frac{1}{2}(n+1)(n+1)n + \frac{1}{2}(n+1)n = n\binom{n+2}{2}.$

4 Defect 2 sets

Let a set \mathcal{M} of n lines be in general position, $n \geq 3$. Then consider the lattice \mathcal{X} defined as

$$\mathcal{X} := \mathcal{X}^{(2)} \cup \mathcal{X}^{(1)} \cup \mathcal{X}^{(0)}, \tag{4.1}$$

where $\mathcal{X}^{(2)}$ is the set of all intersection nodes of these *n* lines, $\mathcal{X}^{(1)}$ is a set of other 2*n* nodes, two in each line, to make the line maximal and $\mathcal{X}^{(0)}$ consists of a single node, denoted by *O*, which does not belong to any line from \mathcal{M} (see Fig. 4.1). Correspondingly, we have that $\#\mathcal{X} = \binom{n}{2} + 2n + 1 = \binom{n+2}{2}$.

Note that all the nodes of $\mathcal{X}^{(k)}$ belong to exactly k maximal lines and thus are k_m -nodes, k = 0, 1, 2.

In the sequel we will need the following characterization of GC_n set \mathcal{X} , with $def(\mathcal{X}) = 2$, due to Carnicer and Gasca:



Figure 4.1: A defect 2 set, n = 4.

Proposition 4.1 ([3], Prop. 2.5). A set \mathcal{X} is a GC_n set of defect 2, i.e., has exactly n maximal lines: $\lambda_1, \ldots, \lambda_n$, where $n \geq 3$, if and only if, (4.1) holds with the following additional properties (see Fig. 4.1):

- (i) There are 3 lines $\ell_1^o, \ell_2^o, \ell_3^o$, called O-lines, concurrent at $O: O = \ell_1^o \cap \ell_2^o \cap \ell_3^o$, such that $\mathcal{X}^{(1)} \subset \ell_1^o \cup \ell_2^o \cup \ell_3^o$;
- (ii) No line ℓ_i^o contains n+1 nodes of \mathcal{X} , i=1,2,3.

Note that each above line ℓ_i^o contains at least two 1_m -nodes. They may contain also 2_m -nodes.

4.1 The used lines in defect 2 sets

Suppose that $M(\mathcal{X}) = \{\lambda_1, \ldots, \lambda_n\}$. Consider a pair of maximal lines $\lambda_i, \lambda_j, 1 \leq i < j \leq n$. We have that the node $A_{ij} := \lambda_i \cap \lambda_j$ uses n-2 maximal lines, i.e., all maximal lines except λ_i, λ_j . Next let us identify the remaining two used lines. We have two 1_m -nodes in each of the two maximal lines, which, in view of the above item (i), lie also in the three O-lines. Denote by n_{ij} the number of O-lines containing at least one of these four 1_m -nodes. Evidently we have that $n_{ij} = 2$ or 3. In the case $n_{ij} = 2$ (the case of the node A_{34} in Fig. 4.1) the four 1_m -nodes belong to two O-lines, which are the two remaining used lines. In the case $n_{ij} = 3$ (the case of the node A_{12} in Fig. 4.1) two of four 1_m -nodes belong to the same O-line, denoted by $\ell_{k(i,j)}^o$, where $1 \leq k(i,j) \leq 3$. Denote also by ℓ_{ij} the line passing through the other two 1_m -nodes. It is easily seen that in this case the lines $\ell_{k(i,j)}^o$ and ℓ_{ij} are the remaining two lines used by the node A_{ij} .

The set of used lines in \mathcal{X} consists of the following three classes:

- 1) The set of n maximal lines;
- 2) The set of three O-lines $\{\ell_1^o, \ell_2^o, \ell_3^o\};$
- 3) The set of lines ℓ_{ij} , $1 \le i < j \le n$, with $n_{ij} = 3$.

Let us verify that these classes are disjoint, i.e., lines from different classes cannot coincide. It suffices to show that the classes 2) and 3) are disjoint. Indeed, any line ℓ_{ij} passes through two 1_m -nodes which belong to two different *O*-lines. Hence ℓ_{ij} cannot coincide with an *O*-line.

Note also that $\ell_{ij} = \ell_{i'j'}$ implies that (i, j) = (i', j'). Indeed, there is an *O*-line which intersects both lines at nodes of \mathcal{X} . Hence without loss of generality we may assume that $\lambda_i = \lambda'_i$, i.e., i = i'. Now suppose by way of contradiction that $j \neq j'$. Then the other two *O*-lines do not intersect the maximal line λ_i at nodes of \mathcal{X} , which contradicts Proposition 4.1, (i).

Next let us show that the lines in the class 2) are proper:

Proposition 4.2. Suppose that \mathcal{X} is a GC_n set of defect 2 and the line $\ell := \ell_i^o$, $i \in \{1, 2, 3\}$, passes through exactly $k \mid_m$ -nodes, where $k \geq 2$. Suppose also that $\mathcal{X} \downarrow_{\ell} \hat{\mathcal{X}}$. Then $\hat{\mathcal{X}}$ is a GC_k set of defect 1, where ℓ is a maximal line. The set $\mathcal{X}^{\ell} = \hat{\mathcal{X}} \setminus \ell$ is a GC_{k-1} set of defect 1.

Proof. In view of Proposition 2.6, we have that $\hat{\mathcal{X}}$ is a GC_k set. Then note that the line ℓ is a (k+1)-node line in $\hat{\mathcal{X}}$, with the node O and $k \ 1_m$ -nodes. Hence the line ℓ , as well as the k maximal lines of \mathcal{X} intersecting the line ℓ at $k \ 1_m$ -nodes are maximal lines in $\hat{\mathcal{X}}$. Now assume by way of contradiction that there is one more maximal line in $\hat{\mathcal{X}}$. Then clearly it coincides with an O line $\ell_j^o, j \neq i$. Observe that each disjoint or adjoint reduction in getting $\hat{\mathcal{X}}$ from \mathcal{X} takes a node or two nodes from ℓ_j^o , respectively. Therefore we obtain that ℓ_j^o is a maximal line in \mathcal{X} , which contradicts Proposition 4.1, (ii).

Definition 4.3. Suppose ℓ is the line which is not maximal or proper. A node $A \in \ell$ is called a $\hat{2}_m$ -node if it is a 2_m -node in the set $\hat{\mathcal{X}} = \hat{\mathcal{X}}(\ell)$.

It can be readily verified that in Fig. 4.2 S is a $\hat{2}_m$ -node in the line ℓ_{12} . In Proposition 7.1 we will prove that if $S \in \ell$ is a $\hat{2}_m$ -node then

 $S = \ell^o \cap \lambda$, where $\ell^o \in Pr(\mathcal{X}) \cap M(\mathcal{X})$ and $\lambda \in M(\mathcal{X})$.

Thus, any $\hat{2}_m$ -node is necessarily an 1_m -node for \mathcal{X} . We also will verify that each used line ℓ may have at most two $\hat{2}_m$ -nodes.

Next let us show that the lines in the class 3) are proper (-1) and are used by only one node.

Proposition 4.4. Suppose that \mathcal{X} is a GC_n set of defect 2, $\ell := \ell_{ij}$, with $n_{ij} = 3, \ 1 \leq i < j \leq n$, and $\mathcal{X} \downarrow_{\ell} \hat{\mathcal{X}}$. Then the point $S := \ell^o_{k(i,j)} \cap \ell$ is the only candidate for a $\hat{2}_m$ -node in ℓ and

 $S \in \hat{\mathcal{X}} \Leftrightarrow S \text{ is an } 1_m \text{-node in } \mathcal{X} \Leftrightarrow S \text{ is a } \hat{2}_m \text{-node } \Leftrightarrow S = \ell^o \cap \lambda^*,$

where $\ell^o := \ell^o_{k(i,j)} \in \Pr(\mathcal{X}) \cap M(\hat{\mathcal{X}})$ and $\lambda^* \in M(\mathcal{X})$. Next we have that $def(\hat{\mathcal{X}}) = 1$, $\#\mathcal{X}^{\ell} = 1$, and the line ℓ is proper (-1), i.e.,

 $\mathcal{X}^{\ell} = \hat{\mathcal{X}} \setminus (\mathcal{C} \cup \ell), \text{ where } \mathcal{C} \text{ is the } d/a \text{ item.}$

Moreover if $S \notin \hat{\mathcal{X}}$ then $\hat{\mathcal{X}}$ is a GC_2 set, $\mathcal{C} = \ell^o_{k(i,j)} \in M(\hat{\mathcal{X}})$. While if $S \in \hat{\mathcal{X}}$, then $\hat{\mathcal{X}}$ is a GC_3 set, $\mathcal{C} = \ell^o_{k(i,j)} \cup \lambda^*$, where $\lambda^* \in M(\mathcal{X})$ and $S \in \lambda^*$.



Figure 4.2: S is a $\hat{2}_m$ -node in ℓ_{12} .

Proof. Note that, in view of Proposition 4.1, (i), all 1_m -nodes of \mathcal{X} belong to the three O-lines. Assume first that $S \notin \hat{\mathcal{X}}$ (see the case of $\ell = \ell_{12}$, and k(i, j) = 3, in Fig. 4.1). Then the line ℓ intersects only two Olines in the nodes of \mathcal{X} . Hence it has just two 1_m -nodes. Thus, in view of Proposition 2.16, we have that $\hat{\mathcal{X}}$ is a GC_2 set. It is easily seen that $M(\hat{\mathcal{X}}) = \{\lambda_i, \lambda_j, \ell^o_{k(i,j)}\}$, hence $def(\hat{\mathcal{X}}) = 1$. Also, ℓ is a 2-node maximal line in $\hat{\mathcal{X}} \setminus \ell^o_{k(i,j)}$ and $\#\mathcal{X}^{\ell} = 1$. Clearly ℓ has no $\hat{2}_m$ -node.

Now assume that $S \in \hat{\mathcal{X}}$ (see the case of $\ell = \ell_{12}, \lambda^* = \lambda_3$, and k(i, j) = 3, in Fig. 4.2). Then, since $S \neq O$ and S is not 2_m -node in \mathcal{X} , we conclude that S is an 1_m -node in \mathcal{X} . Thus $S \in \lambda^*$, where $\lambda^* \in M(\mathcal{X})$. Therefore the line ℓ has exactly three 1_m -nodes. Next, in view of Proposition 2.16, we have that $\hat{\mathcal{X}}$ is a GC_3 set. In this case we have $M(\hat{\mathcal{X}}) = \{\lambda_i, \lambda_j, \lambda^*, \ell^o_{k(i,j)}\}$ and hence $def(\hat{\mathcal{X}}) = 1$. It is easily seen that ℓ is a 3-node line in $\hat{\mathcal{X}}$ with only one $\hat{2}_m$ -node: $S = \ell^o_{k(i,j)} \cap \lambda^*$. Moreover, we have that ℓ is a maximal line in $\hat{\mathcal{X}} \setminus (\ell^o_{k(i,j)} \cup \lambda^*)$ and $\#\mathcal{X}^\ell = 1$.

Denote by n_i the number of line usages in the class i), i = 1, 2, 3. Then we have that the total number of line-usages equals:

 $n_1 + [n_2 + n_3] = n\binom{n+1}{2} + [2\binom{n}{2} + 2n] = \frac{1}{2}n(n+1)n + n(n-1) + 2n = n\binom{n+2}{2}.$ Here we take into account the fact that each 2_m -node uses exactly two

lines from the classes 2) and 3), while each of $2n \ 1_m$ -nodes uses only one line from there, namely from the class 3).

5 Defect 3 sets

Let a set $\mathcal{M} = \{\lambda_1, \dots, \lambda_{n-1}\}$ of n-1 lines be in general position, $n \ge 4$. Then consider the lattice \mathcal{X} defined as

$$\mathcal{X} := \mathcal{X}^{(2)} \cup \mathcal{X}^{(1)} \cup \mathcal{X}^{(0)}, \tag{5.1}$$

where $\mathcal{X}^{(2)}$ is the set of all intersection nodes of these n-1 lines, $\mathcal{X}^{(1)}$ is a set of other 3(n-1) nodes, three in each line, to make the line maximal and $\mathcal{X}^{(0)}$ consists of exactly three non-collinear nodes, denoted by O_1, O_2, O_3 ,



Figure 5.1: A defect 3 set, n = 5.

which do not belong to any line from \mathcal{M} (see Fig. 5.1) Correspondingly, we have that $\#\mathcal{X} = \binom{n-1}{2} + 3(n-1) + 3 = \binom{n+2}{2}$.

Note that in the end all the nodes of $\mathcal{X}^{(k)}$ will belong to exactly k maximal lines and thus become k_m -nodes, k = 0, 1, 2.

Denote by ℓ_i^{oo} , $1 \leq i \leq 3$, the line passing through the two 0_m -nodes in $\{O_1, O_2, O_3\} \setminus \{O_i\}$. We call these lines *OO*-lines. Suppose that $\mathcal{X}^{(1)} = \{A_i^1, A_i^2, A_i^3 \in \lambda_i : 1 \leq i \leq n-1\}$.

In the sequel we will need the following characterization of GC_n set \mathcal{X} , with $def(\mathcal{X}) = 3$, due to Carnicer and Godés:

Proposition 5.1 ([8], Thm. 3.2). A set \mathcal{X} is a GC_n set of defect 3, i.e., has exactly n-1 maximal lines $\lambda_1, \ldots, \lambda_{n-1}$, where $n \ge 4$, if and only if, with some permutation of the indices of the maximal lines and 1_m -nodes, (5.1) holds with the following additional properties (see Fig. 5.1):

- (i) $\mathcal{X}^{(1)} \setminus (\ell_1^{oo} \cup \ell_2^{oo} \cup \ell_3^{oo}) = \{D_1, D_2, D_3\}, \text{ where } D_i := A_i^i;$
- (ii) Each line ℓ_i^{oo} , i = 1, 2, 3, passes through exactly n nodes: $n-2 \ 1_m$ -nodes and two 0_m -nodes. Moreover, $\ell_i^{oo} \cap \lambda_i \notin \mathcal{X}, \ i = 1, 2, 3$;
- (iii) The triples $\{O_1, D_2, D_3\}, \{O_2, D_1, D_3\}, \{O_3, D_1, D_2\}$ are collinear.

Let us denote by ℓ_1^{dd} , ℓ_2^{dd} , ℓ_3^{dd} , the lines passing through the latter triples, respectively, and call them *DD-lines*. Also, the nodes D_i are called *D-nodes*.

5.1 The used lines in defect 3 sets

Let $M(\mathcal{X}) = \{\lambda_1, \ldots, \lambda_{n-1}\}$. Consider a pair of maximal lines λ_i, λ_j , where $1 \leq i < j \leq 3$. Assume that $\{i, j, k\} = \{1, 2, 3\}$. We have that the node $A_{ij} := \lambda_i \cap \lambda_j$ uses n-3 maximal lines, i.e., all the maximal lines except λ_i and λ_j (see the case of the node $A_{12}, k = 3$, in Fig. 5.1). Next let us identify the remaining three used lines. We have three 1_m -nodes in each of two maximal lines, six in all. Two of these nodes belong to the first used line: ℓ_k^{oo} . Two nodes belong to the second used line: ℓ_k^{dd} . Finally, the last used

line is denoted by ℓ_{ij} , which passes through the remaining two 1_m -nodes in the lines λ_i and λ_j .

Now consider a pair of maximal lines λ_i, λ_j , where $1 \leq i \leq 3$ and $4 \leq j \leq n-1$. Assume that $\{i, k_1, k_2\} = \{1, 2, 3\}$. As above the node $A_{ij} := \lambda_i \cap \lambda_j$ uses n-3 maximal lines, i.e., all maximal lines except λ_i and λ_j (see the case of the node A_{34} , i.e., (i, j) = (3, 4) and $(k_1, k_2) = (1, 2)$ in Fig. 5.1). Next we identify the remaining three used lines. Again we have three 1_m -nodes in each of two maximal lines, six in all. Now four of these six nodes lie in the two used OO-lines $\ell_{k_1}^{oo}, \ell_{k_2}^{oo}$. Finally, denote by ℓ_i^j the third used line passing through D_i and the node $\lambda_j \cap \ell_i^{oo}$.

The set of used lines in \mathcal{X} consists of the following five classes:

- 1) The set of n-1 maximal lines;
- 2) The set of three OO-lines $\{\ell_i^{oo}, 1 \leq i \leq 3\};$
- 3) The set of three *DD*-lines $\{\ell_i^{dd}, 1 \le i \le 3\};$
- 4) The set of three lines $\{\ell_{ij}, 1 \le i < j \le 3\};$
- 5) The three sets of n 4 lines $\{\ell_i^j, 4 \le j \le n 1\}, i = 1, 2, 3.$

Proposition 5.2. The classes of lines 1)-5) are disjoint.

Proof. It is enough to show that the classes 2) -5) are disjoint.

Let us start with a line $\ell_2 := \ell_m^{oo}$ from the class 2).

This line cannot coincide with a line ℓ_i^{dd} from the class 3). Indeed, the latter line passes through two *D*-nodes, which, according to Proposition 5.1, (i), do not belong to any line from the class 2).

Then ℓ_2 cannot coincide with a line ℓ_{ij} from the class 4), since the latter line passes through two 1_m -nodes which belong to two different *OO*-lines.

Next ℓ_2 cannot coincide with a line ℓ_i^j from the class 5). Indeed, the latter line passes through a *D*-node, which, as was mentioned above, does not belong to any line from the class 2).

Now consider a line $\ell_3 := \ell_m^{dd}$ from the class 3). This line cannot coincide with a line ℓ_{ij} from the class 4), since the latter line certainly does not pass through two *D*-nodes: D_i and D_j .

Next let us show that ℓ_3 cannot coincide with a line ℓ_i^j from the class 5). Set $\{m, k_1, k_2\} = \{1, 2, 3\}$. We have that the line ℓ_m^{dd} intersects the lines $\ell_{k_1}^{oo}$ and $\ell_{k_2}^{oo}$ at their intersection node O_m . On the other hand the line ℓ_i^j intersects ℓ_i^{oo} at an 1_m -node. Thus, if the given two lines coincide, then we readily conclude that $i \neq k_1, k_2$, hence i = m. Now the line ℓ_m^j passes through the node D_m , while ℓ_3 does not pass through D_m . Hence they cannot coincide.

Finally, let us show that a line ℓ_{ij} from the class 4) cannot coincide with a line $\ell_{i'}^{j'}$ from the class 5). Set $\{i, j, k\} = \{1, 2, 3\}$. We have that the line ℓ_{ij} intersects the maximal lines λ_i and λ_j at nodes belonging to *OO*-lines and the line $\ell_{i'}^{j'}$ passes through the node $D_{i'}$. Thus, if the given two lines coincide, then we conclude that $i' \neq i, j$, hence i' = k. Thus it suffices to show that line ℓ_{ij} does not pass through the node D_k :

Lemma 5.3. Let \mathcal{X} be a GC_n set of defect 3. Then the line ℓ_{ij} , $1 \leq i < j \leq 3$, does not pass through the node D_k , where (i, j, k) = (1, 2, 3).



Figure 5.2: May the node D_1 belong to the line ℓ_{23} ?

Suppose, by way of contradiction, that the line $\ell_{ij} = \ell_{23}$, say, passes through the node D_1 (see Fig. 5.2). Let us apply the Pappus theorem for the pair of triple collinear nodes here: $\{D_1, B, C\}$; and $\{O_1, D_3, D_2\}$. Denote by $\ell(P,Q)$ the line passing through the points P and Q. Observe that $\ell(D_1, D_2) \cap \ell(C, O_1) = O_3$, $\ell(D_1, D_3) \cap \ell(B, O_1) = O_2$, $\ell(C, D_3) \cap \ell(B, D_2) =$ A_{23} . Thus, according to the Pappus theorem we get that the triple of nodes $\{O_2, O_3, A_{23}\}$ is collinear, i.e., the OO-line ℓ_1^{oo} passes through the point of intersection of maximal lines λ_1 and λ_2 , thus contradicting Proposition 5.1, (ii) (cf. the last part of Proposition 3.8's proof in [15]).

Then, let us verify that within each class the differently denoted lines are different. This is evident for the classes 1), 2), 3) and 5). Thus assume that two lines in class 4) coincide: $\ell_{ij} = \ell_{i'j'}$. We have that each of these lines passes through two 1_m -nodes belonging to different *OO*-lines. Hence there is an *O*-line which intersects both lines at nodes of \mathcal{X} . Hence, without loss of generality, we may assume that i = i'. Now suppose by way of contradiction that $j \neq j'$. Then we have that two *OO*-lines do not intersect the maximal line λ_i at nodes of \mathcal{X} , which contradicts Proposition 5.1, (i).

Next we show that the lines in the class 2) are proper.

Proposition 5.4. Suppose that \mathcal{X} is a GC_n set of defect 3 and $\ell = \ell_i^{oo}$, where $i \in \{1, 2, 3\}$. Suppose also that $\mathcal{X} \downarrow_{\ell} \hat{\mathcal{X}}$. Then $\hat{\mathcal{X}}$ is a GC_{n-1} set of defect 2, where ℓ is a maximal line. The set $\mathcal{X}^{\ell} = \hat{\mathcal{X}} \setminus \ell$ is a GC_{n-2} set of defect 2.

Proof. We have that $\hat{\mathcal{X}} = \mathcal{X} \setminus \lambda_i$. It is easily seen that ℓ is a newly emerged maximal line of $\hat{\mathcal{X}}$. In view of Proposition 2.9 we conclude that $def(\hat{\mathcal{X}}) = 2$. It remains to note that the set \mathcal{X}^{ℓ} is a GC_{n-2} set satisfying the conditions of Proposition 4.1, where the O-lines are ℓ_j^{oo} , ℓ_k^{oo} , ℓ_i^{dd} , and $\{i, j, k\} = \{1, 2, 3\}$.

Next we show that the lines in the classes 3), 4), and 5), are proper (-1), (-2), and (-2), and are used by 3, 1, and 1 node, respectively. We also identify the $\hat{2}_m$ -nodes in these lines.

Proposition 5.5. Suppose that \mathcal{X} is a GC_n set of defect 3. Set $\ell_1 := \ell_k^{dd}, \ \ell_2 := \ell_{ij}, \ and \ \ell_3 := l_k^m, \ where \ 1 \le k \le 3, \ 4 \le m \le n-1, \ and \{i, j, k\} = \{1, 2, 3\}.$ Set also $\{k, k_3, k_4\} = \{1, 2, 3\}.$ Then the following points are the only candidates for $\hat{2}_m$ - nodes in the respective lines $\ell_r, r = 1, 2, 3$: $S_1 := \ell_k^{oo} \cap \ell_1, \ S_2 := \ell_k^{oo} \cap \ell_2, \ S_3 := \ell_{k_3}^{oo} \cap \ell_3, \ S_4 := \ell_{k_4}^{oo} \cap \ell_3.$ Also, for each $S = S_q$ and the respective set $\hat{\mathcal{X}} = \hat{\mathcal{X}}(\ell_r)$ we have that

 $S \in \hat{\mathcal{X}} \Leftrightarrow S \text{ is an } 1_m \text{-node in } \mathcal{X} \Leftrightarrow S \text{ is a } \hat{2}_m \text{-node } \Leftrightarrow S = \ell^{oo} \cap \lambda^*,$

where $\ell^{oo} \in \Pr(\mathcal{X}) \cap M(\hat{\mathcal{X}})$ and $\lambda^* \in M(\mathcal{X})$. Next, we have that $def[\hat{\mathcal{X}}(\ell_1)] = def[\hat{\mathcal{X}}(\ell_2)] = 2$, $def[\hat{\mathcal{X}}(\ell_3)] = 1$, and $\#\mathcal{X}^{\ell_1} = 3$, $\#\mathcal{X}^{\ell_2} = \#\mathcal{X}^{\ell_3} = 1$. Further, the lines ℓ_1, ℓ_2 , and ℓ_3 , are proper (-1), (-2), and (-2), respectively:

$$\mathcal{X}^{\ell_1} = \hat{\mathcal{X}} \setminus (\mathcal{C}_1 \cup \ell_1), \ \mathcal{X}^{\ell_2} = \hat{\mathcal{X}} \setminus (\mathcal{C}_2 \cup \ell_1 \cup \ell_2), \ \mathcal{X}^{\ell_3} = \hat{\mathcal{X}} \setminus (\mathcal{C}_3 \cup \mathcal{C}_4 \cup \ell_3), \ (5.2)$$

where C_i is the d/a item.

Moreover, the following assertions hold for the lines ℓ_1 and ℓ_2 : If $S_i \notin \hat{\mathcal{X}}(\ell_i)$, then $\hat{\mathcal{X}}(\ell_i)$ is a GC₃ set, and $\mathcal{C}_i = \ell_k^{oo}$, i = 1, 2. If $S_i \in \hat{\mathcal{X}}(\ell_i)$, then $\hat{\mathcal{X}}(\ell_i)$ is a GC₄ set, $\mathcal{C}_i = \ell_k^{oo} \cup \lambda_i^*$, where $\lambda_i^* \in M(\mathcal{X})$ and $S_i \in \lambda_i^*$, i = 1, 2. The following assertions hold for the line ℓ_3 : If $S_3, S_4 \notin \ell_3$ then $\hat{\mathcal{X}}(\ell_3)$ is GC₃ set, and $\mathcal{C}_i = \ell_{k_i}^{oo}$, i = 3, 4. If $S_3 \in \ell$, $S_4 \notin \ell$ then $\hat{\mathcal{X}}(\ell_3)$ is GC₄ set, $\mathcal{C}_3 = \ell_{k_3}^{oo} \cup \lambda^*$, $\mathcal{C}_4 = \ell_{k_4}^{oo}$. If $S_3, S_4 \in \ell_3$ then $\hat{\mathcal{X}}(\ell_3)$ is GC₅ set, $\mathcal{C}_i = \ell_{k_i}^{oo} \cup \lambda_i^*$, where $\lambda_i^* \in M(\mathcal{X})$ and $S_i \in \lambda_i^*$, i = 3, 4.

Proof. Note that, in view of Proposition 5.1, (i), all 1_m -nodes of \mathcal{X} , except the three *D*-nodes, belong to the three *OO*-lines. Note also that no 2_m -node of a line ℓ belongs to $\hat{\mathcal{X}}(\ell)$.

The line ℓ_1 passes through two *D*-nodes and the point of intersection of two *OO*-lines: the node O_k . Thus $S_1 = \ell_1 \cap \ell_k^{oo}$ is the only candidate for a third 1_m -node in ℓ_1 .

In view of Lemma 5.3 the line ℓ_2 does not pass through *D*-nodes. It passes through two 1_m -nodes which are intersection points with *OO*-lines. Thus $S_2 = \ell_2 \cap \ell_k^{oo}$ is the only candidate for a third 1_m -node in ℓ_2 .

Suppose first that the lines ℓ_1 and ℓ_2 do not have a third 1_m -nodes (see the case of $\ell_1 = \ell_3^{dd}$ and $\ell_2 = \ell_{12}$, i.e., (i, j, k) = (1, 2, 3) in Fig. 5.1). The following holds for both lines $\ell = \ell_1$ and $\ell = \ell_2$.

Observe that ℓ has exactly two 1_m -nodes coinciding with the intersection points with the maximal lines λ_i and λ_j . Thus, in view of Proposition

2.16, we have that \mathcal{X} is a GC_3 set. It consists of the node A_{ij} , six 1_m -nodes: three and three lying in the lines λ_i and λ_j , and the three 0_m -nodes: O_1, O_2, O_3 . Here the line ℓ_k^{oo} with 4 nodes, including two O nodes, is an ℓ -disjoint maximal.

Next, the line ℓ_1 is a maximal line in the set $\hat{\mathcal{X}} \setminus \ell_k^{oo}$ and the line ℓ_2 is a maximal line in the set $\hat{\mathcal{X}} \setminus (\ell_k^{oo} \cup \ell_1)$. Hence we readily get that the first two relations of (5.2) hold with $\mathcal{C}_1 = \mathcal{C}_2 = \ell_k^{oo}$.

Now note that $M(\hat{\mathcal{X}}) = \{\lambda_i, \lambda_j, \ell_k^{oo}\}$ and hence $def(\mathcal{X}) = 2$. Indeed, if there is another maximal line then it is easily seen that it has to pass through two O nodes and 2 other nodes in λ_i and λ_j . But clearly the two OO-lines: ℓ_i^{oo} and ℓ_j^{oo} , pass through only 3 nodes of $\hat{\mathcal{X}}$. Note also that ℓ_1 is a 3-node line in $\hat{\mathcal{X}}$ and ℓ_2 is a 2-node line in $\hat{\mathcal{X}}$, without any $\hat{2}_m$ -node.

Next, suppose that each of the lines $\ell_q, q = 1, 2$, has a third 1_m -node. Then this node is $S_q = \lambda_q^* \cap \ell_k^{oo}$, where $\lambda_q^* \in M(\mathcal{X})$. Note that S_q is the only $\hat{2}_m$ -node in ℓ_q . Now the number of 1_m -nodes in each of the lines equals to three. Thus, in view of Proposition 2.16, we have that $\hat{\mathcal{X}}(\ell_q)$ is a GC_4 set. Here ℓ_k^{oo} and λ_q^* form a pair of ℓ -adjoint maximal lines and we readily get that the first two relations of (5.2) hold with $\mathcal{C}_q = \ell_k^{oo} \cup \lambda_q^*$, q = 1, 2.

Note that $M(\hat{\mathcal{X}}(\ell_q)) = \{\lambda_i, \lambda_j, \ell_k^{oo}, \lambda_q^*\}$ and hence $\operatorname{def}[\hat{\mathcal{X}}(\ell_q)] = 2, q = 1, 2$. Indeed, if there is more than four maximal lines then it is easily seen that $\hat{\mathcal{X}} \setminus \lambda_q^*$ would have more than three maximal lines, which contradicts the consideration in the previous case.

Now consider the line ℓ_3 . It passes through two 1_m -nodes, namely D_k and the point of intersection of λ_m with one *OO*-line ℓ_k^{oo} . Thus the points of intersections with the other two *OO*-lines, i.e., S_3 and S_4 , are the only candidates for a third and forth 1_m -nodes in ℓ_3 .

Suppose first that the line ℓ_3 does not have a third and forth 1_m -nodes (see the case of $\ell_3 = \ell_3^4$, i.e., $(k, m) = (3, 4), (k_3, k_4) = (1, 2)$ in Fig. 5.1). Then ℓ_3 has exactly two 1_m -nodes. Thus, in view of Proposition 2.16, we have that $\hat{\mathcal{X}}$ is a GC_3 set. It consists of the node A_{ij} , six 1_m -nodes: three and three lying in the lines λ_i and λ_j , and the three 0_m -nodes: O_1, O_2, O_3 . Here the lines $\ell_{k_q}^{oo}$, q = 3, 4, with 4 nodes, including two O nodes, are ℓ -disjoint maximal lines. Next, the line ℓ_3 is a maximal line in the set $\hat{\mathcal{X}} \setminus (\ell_{k_3}^{oo} \cup \ell_{k_4}^{oo})$. Hence we readily get that the third relation of (5.2) holds with $\mathcal{C}_q = \ell_{k_q}^{oo}, q = 3, 4$.

Now we have that $M(\hat{\mathcal{X}}) = \{\lambda_i, \lambda_j, \ell_{k_3}^{oo}, \ell_{k_4}^{oo}\}$ and hence $def(\mathcal{X}) = 1$. Note that ℓ_3 is a 2-node line in $\hat{\mathcal{X}}$, without any $\hat{2}_m$ -node.

Next, suppose that the line ℓ_3 has a third 1_m -node, say, S_3 . Then we have that $S_3 = \lambda_3^* \cap \ell_{k_3}^{oo}$, where $\lambda_3^* \in M(\mathcal{X})$. Now the number of 1_m -nodes in ℓ_3 equals to three. Thus, in view of Proposition 2.16, we have that $\hat{\mathcal{X}}(\ell_q)$ is a GC_4 set. Here $\ell_{k_3}^{oo}$ and λ_3^* form a pair of ℓ_3 -adjoint maximal lines and we readily get that the third relation of (5.2) holds with $\mathcal{C}_3 = \ell_{k_3}^{oo} \cup \lambda_3^*$ and

 $\mathcal{C}_4 = \ell_{k_4}^{oo}$. Note that S_3 is the only $\hat{2}_m$ -node in ℓ_3 .

Then we have that $M(\hat{\mathcal{X}}) = \{\lambda_i, \lambda_j, \ell_{k_3}^{oo}, \ell_{k_4}^{oo}, \lambda_3^*\}$ and hence $def(\mathcal{X}) = 1$. Indeed, if there is more than five maximal lines then it is easily seen that $\hat{\mathcal{X}} \setminus \lambda_3^*$ would have more than four maximal lines, which contradicts the consideration in the previous case.

Finally, suppose that the line ℓ_3 has four 1_m -nodes. Then we have that $S_q = \lambda_q^* \cap \ell_{k_q}^{oo}$, where $\lambda_q^* \in M(\mathcal{X})$, q = 3, 4. Now the number of 1_m -nodes in ℓ_3 equals to four. Thus, in view of Proposition 2.16, we have that $\hat{\mathcal{X}}(\ell_q)$ is a GC_5 set. Here $\ell_{k_q}^{oo}$ and λ_q^* , q = 3, 4, form two pairs of ℓ -adjoint maximal lines and we readily get that the third relation of (5.2) holds with $\mathcal{C}_q = \ell_{k_q}^{oo} \cup \lambda_q^*$, q = 3, 4. Note that in this case S_3 and S_4 are the only $\hat{2}_m$ -nodes in ℓ_3 . Note also that $M(\hat{\mathcal{X}}) = \{\lambda_i, \lambda_j, \ell_{k_3}^{oo}, \ell_{k_4}^{oo}, \lambda_3^*, \lambda_4^*\}$ and hence def $(\mathcal{X}) = 1$.

Denote by n_i the number of line usages in the class i), i = 1, ..., 5. Then we have that the total number of line-usages equals:

 $n_1 + \dots + n_5 = (n-1)\binom{n+1}{2} + 3\binom{n}{2} + 9 + 3 + 3(n-4) = n\binom{n+2}{2}.$

6 Generalized principal lattices: defect n-1 sets

A principal lattice is defined as an affine image of the set (see Fig. 6.1)

$$PL_n := \left\{ (i, j) \in \mathbb{N}_0^2 : \quad i + j \le n \right\}.$$

Let us set $I = \{0, 1, ..., n\}$. Observe that the following 3 sets of n + 1 lines, namely $\{x = i : i \in I\}$, $\{y = j : j \in I\}$, and $\{x + y = n - k : k \in I\}$, intersect at PL_n . We have that PL_n is a GC_n set. Moreover, the following is the fundamental polynomial of the node $(i_0, j_0) \in PL_n$:

$$p_{i_0 j_0}^{\star}(x, y) = \prod_{0 \le i < i_0, \ 0 \le j < j_0, \ 0 \le k < k_0} (x - i)(y - j)(x + y - n + k), \quad (6.1)$$

where $k_0 = n - i_0 - j_0$. Next let us bring the definition of the generalized



Figure 6.1: A principal lattice PL_5 .

principal lattice due to Carnicer, Gasca and Godés (see [6], [7]):

Definition 6.1 ([7]). A node set \mathcal{X} is called a generalized principal lattice, briefly GPL_n , if there are 3 sets of lines each containing n + 1 lines

$$\ell_i^j := \ell_i^j(\mathcal{X}), \ i = 0, 1, \dots, n, \ j = 0, 1, 2, \tag{6.2}$$

such that the 3n + 3 lines are distinct,

$$\ell_i^0 \cap \ell_j^1 \cap \ell_k^2 \cap \mathcal{X} \neq \emptyset \iff i+j+k=n,$$

and $\mathcal{X} = \left\{ x_{ijk} \mid x_{ijk} := \ell_i^0 \cap \ell_j^1 \cap \ell_k^2, 0 \le i, j, k \le n, i+j+k=n \right\}.$

Observe that if $0 \leq i, j, k \leq n$, i + j + k = n then the three lines $\ell_i^0, \ell_j^1, \ell_k^2$, intersect at a node $x_{ijk} \in \mathcal{X}$. This implies that each node of \mathcal{X} belongs to only one line of each of the three sets of n + 1 lines. Therefore $\#\mathcal{X} = (n+1)(n+2)/2$.

One can find readily, as in the case of PL_n , the fundamental polynomial of each node $A = x_{i_0 j_0 k_0} \in \mathcal{X}, \ i_0 + j_0 + k_0 = n$:

$$p_A^{\star} = \prod_{0 \le i < i_0, \ 0 \le j < j_0, \ 0 \le k < k_0} \ell_i^0 \ell_j^1 \ell_k^2.$$
(6.3)

Thus \mathcal{X} is a GC_n set.

Let us bring a characterization for GPL_n set due to Carnicer and Godés:

Theorem 6.2 ([7], Thm. 3.6). Assume that GM Conjecture holds for all degrees up to n - 3. Then the following statements are equivalent:

- (i) \mathcal{X} is generalized principal lattice of degree n;
- (ii) \mathcal{X} is a GC_n set with $\#M(\mathcal{X}) = 3$.

From (6.3) we have that the only used lines in the generalized principal lattice \mathcal{X} are

1) The three sets of n lines $\ell_s^r(\mathcal{X})$, where $0 \le s \le n-1$, r = 0, 1, 2. Let $\ell := \ell_{n-k+1}^r(\mathcal{X})$ be a k-node line, $k \ge 2$. Then it is easily seen that

$$\mathcal{X}^{\ell} = \mathcal{X} \setminus (\ell_0^r \cup \ell_1^r \cup \dots \cup \ell_{n-k+1}^r).$$
(6.4)

Hence

$$\mathcal{X}^{\ell}$$
 is a GC_{k-2} subset of \mathcal{X} and $\#\mathcal{X}^{\ell} = \binom{k}{2}$. (6.5)

Next suppose that $\ell = \ell_i^r$ is any used line, which is not a maximal line, i.e., $1 \leq k \leq n-1$. It is easily seen that the maximal line $\lambda_0 := \ell_0^r$ does not intersect ℓ while the remaining two maximal lines intersect ℓ at two different nodes. Thus for ℓ there is only one reduction, which is disjoint one. Therefore we get

$$\mathcal{X}(\ell) = \mathcal{X} \setminus \lambda_0. \tag{6.6}$$

From here we readily get that the 3 *n*-node lines ℓ_1^r , r = 0, 1, 2, are the proper lines of the generalized principal lattice \mathcal{X} . On the other hand, according to Corollary 4.4 [14], there are exactly three *n*-node lines, provided that $n \ge 4$. Therefore we have that

$$\Pr(\mathcal{X}) = \{\ell_1^0, \ell_1^1, \ell_1^2\} = N(\mathcal{X}), \tag{6.7}$$

where the last equality holds if $n \ge 4$.

Now, in view of Lemma 2.10, we conclude that

$$\operatorname{defect} \mathcal{X}(\ell) = \operatorname{defect} \mathcal{X} - 1, \tag{6.8}$$

for any used line $\ell \notin M(\mathcal{X})$ in defect n-1 set \mathcal{X} .

Note that for the maximal line λ_0 , in view of Lemma 2.12, we get that $\#(\lambda_0 \cap \mathcal{X}^{\ell}) = 0$ and for each remaining maximal line λ , in view of the relation 6.4, we get that $\#(\lambda \cap \mathcal{X}^{\ell}) = k - 1$.

Next we have that the total number of line-usages here equals: $3\left[\binom{n+1}{2} + \binom{n}{2} + \cdots + \binom{2}{2}\right] = 3\binom{n+2}{3} = n\binom{n+2}{2}.$

7 On the properties of the set \mathcal{X}^{ℓ}

For GC_n sets of defect n-1 there is a simple formula for the set \mathcal{X}^{ℓ} , namely (6.4), which yields its basic properties given in (6.5). For other GC_n sets we have the following

Theorem 7.1. Let \mathcal{X} be a GC_n set, $\ell \notin M(\mathcal{X})$ be a used line and $\mathcal{X} \downarrow_{\ell} \mathcal{X}$. Assume that GM Conjecture holds for all degrees up to n. Then we have that

$$def(\mathcal{X}) = def(\mathcal{X}) - 1 \quad or \quad def(\mathcal{X}) - 2.$$
(7.1)

Next, by assuming that $def(\mathcal{X}) \neq n-1$ if $n \geq 5$, and the line ℓ is not proper, we get that ℓ is proper (-r) and has at most $r \ \hat{2}_m$ -nodes, where r = 1 or 2. Also we have that the set \mathcal{X}^{ℓ} is GC_0 or GC_1 set and the set $\hat{\mathcal{X}}$ is a GC_k set with $k \leq 5$.

Moreover, if $S \in \ell$ is a $\hat{2}_m$ -node, then $S = \ell^o \cap \lambda$, where $\ell^o \in Pr(\mathcal{X}) \cap M(\hat{\mathcal{X}})$ and $\lambda \in M(\mathcal{X})$.

Proof. The proof is divided into five cases.

Case 1. $def(\mathcal{X}) = 0.$

In this case, all used lines are maximal.

Case 2. $def(\mathcal{X}) = 1$.

In this case we have two classes of used lines and all lines in the class 1) are maximal. Then, by Proposition 3.1, all lines in the class 2) are proper and $def(\hat{\mathcal{X}}) = 0$.

Case 3. $def(\mathcal{X}) = 2$.

In this case we have three classes of used lines and all lines in the class 1) are maximal. Then, by Proposition 4.2, all the lines in the class 2) are proper and $def(\hat{\mathcal{X}}) = 1$.

Thus it remains to consider a line $\ell := \ell_{ij}$ belonging to the class 3). According to Proposition 4.4 the line ℓ is proper (-1), $def(\hat{\mathcal{X}}) = 1$ and $\#\mathcal{X}^{\ell} = 1$. Also, we have that $\hat{\mathcal{X}}$ is a GC_2 or GC_3 set if ℓ contains no $\hat{2}_m$ -node or contains a $\hat{2}_m$ -node, respectively. Case 4. $def(\mathcal{X}) = 3$.

In this case we have five classes of used lines and all lines in the class 1) are maximal. Then, by Proposition 5.4, all the lines in the class 2) are proper and $def(\hat{\mathcal{X}}) = 2$.

Now consider lines $\ell_1 := \ell_k^{dd}$ and $\ell_2 := \ell_{ij}$ belonging to the classes 3) and 4), respectively, where $\{i, j, k\} = \{1, 2, 3\}$. According to Proposition 5.5 the line ℓ_i is proper (-i) and def $[\hat{\mathcal{X}}(\ell_i)] = 2$, i = 1, 2. We also have that \mathcal{X}^{ℓ_1} and \mathcal{X}^{ℓ_2} are GC_1 and GC_0 sets, respectively.

Then $\hat{\mathcal{X}}$ is a GC_3 or GC_4 set if ℓ_i contains no $\hat{2}_m$ -node or contains a $\hat{2}_m$ -node, respectively i = 1, 2.

Next consider a line $\ell := \ell_i^j$ belonging to the class 5), where $1 \leq i \leq 3$, $4 \leq j \leq n-1$. According to Proposition 5.5 the line ℓ is proper (-2), $def[\hat{\mathcal{X}}(\ell)] = 1$ and $\# \mathcal{X}^{\ell} = 1$.

Then $\hat{\mathcal{X}}$ is a GC_3 , GC_4 , and a GC_5 set if ℓ contains zero, one, two $\hat{2}_m$ -nodes, respectively.

Case 5. $def(\mathcal{X}) = n-1$. This case follows simply from the relation (6.8). Finally note that the "Moreover" statement follows from Propositions 4.4 and 5.5.

Note that the equality

$$\operatorname{def}(\hat{\mathcal{X}}) = \operatorname{def}(\mathcal{X}) - 2$$

holds only for the above lines ℓ_i^j belonging to the class 5) of the used lines of defect 3 sets.

Similarly to the proof of Theorem 7.1, by using Propositions 4.4 and 5.5, one can verify the following

Remark 7.2. Assume that the hypotheses of Theorem 7.1 hold, $def(\mathcal{X}) \neq n-1$ if $n \geq 5$, and in addition ℓ is a proper (-r) line, where r = 1 or 2. Then all (-1), (-2) disjoint and adjoint items are proper lines and pairs of proper lines with maximal lines of \mathcal{X} , respectively. The only exception is for the class 4) of the used lines of defect 3 sets, where the (-2) item is necessarily disjoint and proper (-1) line.

Next let us mention a result from [15] (Corollary 4.1), which states that if $n \ge 6$ then for any k-node line in a GC_n set $\mathcal{X}, 2 \le k \le n$, there is either ℓ -disjoint maximal line or a pair of ℓ -adjoint maximal lines. Below we strengthen this result in the case of used lines.

Corollary 7.3. Let \mathcal{X} be a GC_n set, ℓ be a k-node used line, $2 \leq k \leq n$. Assume that GM Conjecture holds for all degrees up to n. Then there is either ℓ -disjoint maximal line or a pair of ℓ -adjoint maximal lines.

Proof. It suffices to verify that for any used line ℓ we have that $\mathcal{X} \neq \hat{\mathcal{X}}(\ell)$. This follows from the statement $\operatorname{def}(\hat{\mathcal{X}}) \leq \operatorname{def}(\mathcal{X}) - 1$ of Theorem 7.1. \Box

Theorem 7.4. Let \mathcal{X} be a GC_n set and ℓ be a k-node used line. Assume that ℓ contains exactly $r \ 2_m$ -nodes and $\hat{r} \ \hat{2}_m$ -nodes. Assume that GM Conjecture holds for all degrees up to n. Then \mathcal{X}^{ℓ} is a GC_{s-2} set and hence $\#\mathcal{X}^{\ell} = {s \choose 2}$, where $s = k - r - \hat{r}$. Moreover, for any used line ℓ we have that $\hat{r} \leq 2$. Furthermore, $\hat{r} = 0$ if $\#\mathcal{X}^{\ell} > 3$.

Proof. First note that Theorem 7.1 implies the statements in the parts "Moreover" and "Furthermore", if defect $\mathcal{X} \neq n-1$. Then, in view of relation (1.1) and Proposition 2.4, we get that Theorem holds with $r = \hat{r} = 0$, if the line ℓ is a maximal line.

Next, in view of Lemmas 2.12 and 2.13, we get that Theorem holds with $\hat{r} = 0$, if the line ℓ is a proper line. Thus Theorem holds if \mathcal{X} is a Chung-Yao or Carnicer-Gasca lattice, since then, in view of Proposition 3.1, all used lines are maximal or proper lines.

If \mathcal{X} is a defect 2 set then, in view of Proposition 4.2, all lines in the classes 1) and 2) are maximal and proper, respectively. Thus consider a line $\ell := \ell_{ij}$ belonging to the class 3). According to Proposition 4.4 the line ℓ , not counting a $\hat{2}_m$ -node, is 2-node line in $\hat{\mathcal{X}}$ and $\#\mathcal{X}^{\ell} = 1$. Also we have that $s = k - r - \hat{r} = 2$.

If \mathcal{X} is a defect 3 set then, in view of Proposition 5.4, all lines in the classes 1) and 2) are maximal and proper, respectively. Thus consider lines $\ell_1 := \ell_k^{dd}$ and $\ell_2 := \ell_{ij}$ belonging to the classes 3) and 4), respectively, where $\{i, j, k\} = \{1, 2, 3\}$. According to Proposition 5.5 the line ℓ_1 , not counting a possible $\hat{2}_m$ -node, is 3-node line in $\hat{\mathcal{X}}$ and $\#\mathcal{X}^{\ell_1} = 3$. Also we have that $s = k - r - \hat{r} = 3$. While the line ℓ_2 , not counting a possible $\hat{2}_m$ -node, is 2-node line in $\hat{\mathcal{X}}$ and $\#\mathcal{X}^{\ell_2} = 1$. Also we have that $s = k - r - \hat{r} = 2$.

Next, consider a line $\ell := \ell_i^j$ belonging to the class 5), where $1 \leq i \leq 3$, $4 \leq j \leq n-1$. According to Proposition 5.5 the line ℓ , not counting possible two $\hat{2}_m$ -nodes, is a 2-node line in $\hat{\mathcal{X}}$ and $\#\mathcal{X}^\ell = 1$. Also we have that $s = k - r - \hat{r} = 2$.

Finally, let \mathcal{X} be a generalized principal lattice, i.e., def $\mathcal{X} = n - 1$. Consider a line ℓ , which is not maximal or proper. Then we have that $r = \hat{r} = 0$. On other hand, in view of the formula (6.5), we have that $\#\mathcal{X}^{\ell} = \binom{k}{2}$.

From Theorem 7.4 we readily get

Corollary 7.5. Let \mathcal{X} be a GC_n set and ℓ be a k-node used line. Assume that ℓ contains exactly $r \ 2_m$ -nodes. Assume that GM Conjecture holds for all degrees up to n. Then $\#\mathcal{X}^{\ell} \in \left\{1, 3, \binom{k-r}{2}\right\}$.

Indeed, if $\hat{r} = 0$ then $\#\mathcal{X}^{\ell} = \binom{k-r}{2}$. While if $\hat{r} \neq 0$ then $\operatorname{def}(\mathcal{X}) \neq n-1$, and, in view of Theorem 7.1, we get that $\#\mathcal{X}^{\ell} \leq 3$, i.e., $\#\mathcal{X}^{\ell} = 1$ or 3.

Below, we restate a result from [15] (Theorem 3.1). In the "Moreover" and "Furthermore" parts we complement it.

Let us call the maximal line of \mathcal{X} passing through an $\hat{2}_m$ -node in $\ell : \ell$ -special maximal.

Theorem 7.6. Let \mathcal{X} be a GC_n set, ℓ be a line with $\#\mathcal{X}^{\ell} = {s \choose 2}$, $s \geq 2$. Assume that GM Conjecture holds for all degrees up to n. Then for any maximal line λ we have that $\#(\lambda \cap \mathcal{X}^{\ell}) = s - 1$, or 0. Moreover, the latter case:

$$\#(\lambda \cap \mathcal{X}^{\ell}) = 0, \tag{7.2}$$

holds, if and only if

- (i) λ is an ℓ -disjoint maximal line, or
- (ii) λ is one of ℓ -adjoint maximal lines, or
- (iii) λ is an ℓ -special maximal line.

Furthermore, for any line ℓ , (iii) may take place for at most two maximal lines λ .

Proof. First suppose that ℓ is a maximal line. Then, in view of the relation (1.1) and Proposition 2.4, (ii), we have that $\#(\lambda \cap \mathcal{X}^{\ell}) = n$ for any other maximal line λ .

Now from Lemma 2.12 and 2.13 applied to both \mathcal{X} and $\hat{\mathcal{X}}$ the direction "if" of the "Moreover" part follows. For the direction "only if" note first that if \mathcal{X} is a defect n-1 set then Theorem follows from the comment in the paragraph after the relation (6.8). Next assume that $def(\mathcal{X}) \neq n-1$. Consider a maximal line λ and assume that it is not ℓ -disjoint, ℓ -adjoint and ℓ -special. Now, according to Definition 2.14 and Proposition 2.6, λ is a maximal line in $\hat{\mathcal{X}}$. Then, in view of the Remark 7.2, λ is not present in any (-1) or (-2) d/a item. Consequently, Proposition 2.6 implies that λ is a maximal line in the GC_{s-2} set \mathcal{X}^{ℓ} and hence $\#(\lambda \cap \mathcal{X}^{\ell}) = s - 1$.

Finally, note that the "Furthermore" part follows from Theorem 7.1. \Box

8 On proper, maximal, and *n*-node lines

Recall that $N(\mathcal{X})$ denotes the set of all *n*-node lines of GC_n set \mathcal{X} . The following Proposition presents four unexpected properties of the set $N(\mathcal{X})$.

Proposition 8.1. Let \mathcal{X} be a GC_n set, $n \geq 4$. Assume that GM Conjecture holds for all degrees up to n. Then $N(\mathcal{X}) \subset Pr(\mathcal{X})$ and $\#N(\mathcal{X}) \in \{0, 1, 2, 3\}$. Moreover, any n-node line intersects each maximal line, except possibly one, at a node of \mathcal{X} . Furthermore, any two n-node lines intersect at a node of \mathcal{X} .

Proof. Assume that ℓ is an *n*-node line. Assume also that ℓ is type (i, j, k), meaning that it passes through $i \ 0_m$ -nodes, $j \ 1_m$ -nodes, and $k \ 2_m$ -nodes. Note that to prove the "Moreover" part it suffices to verify the inequality $j + 2k \ge \#M(\mathcal{X}) - 1$. Consider the following cases:

Case 1. $\#M(\mathcal{X}) = n + 2$.

In this case we have only 2_m -nodes and there is no *n*-node line. Indeed, suppose by way of contradiction that ℓ is an *n*-node line. Then for the number of maximal lines intersecting ℓ we have $2n \leq n+2$, i.e., $n \leq 2$.

Case 2. $\#M(\mathcal{X}) = n + 1$.

In this case we have only 1_m and 2_m -nodes. Thus ℓ is type (0, j, k) = (0, n - k, k). Then we have that $n - k + 2k \le n + 1$. Hence k = 0 or 1.

Recall that there are n + 1 1_m -nodes. Thus only one *n*-node line is possible if $n \ge 5$: type (0, n, 0) or type (0, n - 1, 1). If n = 4 then either one type (0, 4, 0) line is possible or two type (0, 3, 1) lines. In view of Proposition 3.1 all above *n*-node lines are proper lines.

Case 3. $\#M(\mathcal{X}) = n$.

In this case we have one 0_m -node: O.

First suppose that the line ℓ does not pass through O, i.e., it is type (0, j, k) = (0, n - k, k). Then we have that $n - k + 2k \leq n$, i.e., k = 0. In this case, in view of Proposition 4.1, (i), the number of 1_m -nodes in ℓ is less than or equal to 3. Hence $n \leq 3$.

Now suppose that the line ℓ passes through O, i.e., is type (1, j, k) = (1, n - 1 - k, k). Also suppose that ℓ is different from O-lines. In this case, by Proposition 4.1, (i), the number of 1_m -nodes in ℓ equals to 0. Hence k = n - 1 and we have that $2(n - 1) \leq n$. Hence $n \leq 2$.

Next suppose that the line ℓ is an *O*-line. For the number of 1_m -nodes in the three *O*-lines we have $n_1 + n_2 + n_3 = 2n$, where $n_i \ge 2$ is the number of 1_m -nodes in *i*th *O*-line. Suppose all three lines are *n*-node lines. Observe that since $(n-1-k) + 2k \le n$ then each of them has at most one 2_m -node. Therefore we have that $3(n-2) \le 2n$ hence $n \le 6$. Thus for $n \ge 7$ two *n*-node lines are possible in all: of types (1, n - 1, 0) or (1, n - 2, 1). It is easily seen that in the cases n = 4, 5, 6, all three *O*-lines can be *n*-node lines of types (1, n - 1, 0) or (1, n - 2, 1). In view of Proposition 4.2 each *O*-line is a proper line.

Case 4. $\#M(\mathcal{X}) = n - 1$.

In this case we have three non-collinear 0_m -nodes: O_1, O_2, O_3 .

First suppose that the line ℓ does not pass through any O node, i.e., it is type (0, j, k) = (0, n - k, k). Then we get $n - k + 2k \le n - 1$, i.e., $k \le -1$.

Now suppose that the line ℓ passes through one O node, i.e., is type (1, j, k) = (1, n - 1 - k, k). Also suppose that ℓ is different from OO-lines. Then we have that $n - 1 - k + 2k \leq n - 1$. Hence k = 0. In this case, by Proposition 5.1, (iii), the number of 1_m -nodes in ℓ is less than or equal to 3 (the case of DD-line with one $\hat{2}_m$ -node). Thus $n - 1 \leq 3$ and $n \leq 4$. It is

easily seen that in the case of n = 4 a *DD*-line can not have a $\hat{2}_m$ -node and be a 4-node line.

Next suppose that the line ℓ is an *OO*-line. By Proposition 5.1, (ii), these three lines are *n*-node lines of type (2, n-2, 0). In view of Proposition 5.4 each *OO*-line is a proper line.

Case 5. $\#M(\mathcal{X}) = 3$. This case follows from the relation (6.7).

Remark 8.2. Note that all above *n*-node lines are either type (i, j, 0) with $i+j=n, j+2\cdot 0 = \#M(\mathcal{X})-1$, or type (i, j, 1) with $i+j+1=n, j+2\cdot 1 = \#M(\mathcal{X})$. Note that in the first case *n*-node line intersects all but one maximal line of \mathcal{X} , and in the second case it intersects all maximal lines of \mathcal{X} .

Recall that in defect 1 sets the proper lines are the lines passing through at least two of n + 1 1_m-nodes. For other GC sets we have the following

Proposition 8.3. Let \mathcal{X} be a GC_n set, $def(\mathcal{X}) \neq 1$, $n \geq 4$. Assume that GM Conjecture holds for all degrees up to n. Then $\#Pr(\mathcal{X}) \in \{0,3\}$.

Proof. Indeed, there are no proper lines in Chung-Yao lattice. In the defect 2 and defect 3 sets the only proper lines are the three O-lines and three OO-lines, respectively. Finally, the case of defect n-1 sets follows from the relation (6.7).

Next, we complement Proposition 2.9. From Proposition 8.1 and Remark 8.2 we readily get the following

Proposition 8.4. Let \mathcal{X} be a GC_n set, $n \ge 4$. Assume that GM Conjecture holds for all degrees up to n. Then the relation

$$def(\mathcal{X} \setminus \lambda) = def(\mathcal{X}) - 1 \tag{8.1}$$

holds for $\lambda \in M(\mathcal{X})$ if and only if there is a type (i, j, 0) n-node line ℓ such that $\ell \cap \lambda \notin \mathcal{X}$. Hence (8.1) holds for at most three maximal lines λ .

Note that for all other maximal lines λ , in view of Proposition 2.9, we have that $\operatorname{def}(\mathcal{X} \setminus \lambda) = \operatorname{def}(\mathcal{X})$.

For a defect n-1 set there is a simple formula for the fundamental polynomials of nodes $A \in \mathcal{X}$, namely (6.4). This formula gives the *n* used lines of the node *A*. For other GC_n sets we have the following

Proposition 8.5. Let \mathcal{X} be a GC_n set and $def(\mathcal{X}) \neq n-1$ if $n \geq 5$. Then we have that among n lines used by a node $A \in \mathcal{X}$ at most three lines are proper, at most one line is proper (-1), at most one line is proper (-2), and at least n-3 lines are maximal.

Proof. Let us list the following possibilities of n lines used by A:

1) All n lines are maximal lines;

2) n-1 lines are maximal and one line is proper;

3) n-2 lines are maximal and two lines are proper;

4) n-2 lines are maximal, one is proper and one is proper (-1);

5) n-3 lines are maximal and three lines are proper;

6) n-3 lines are maximal, two are proper, and one is proper (-2);

7) n-3 lines are maximal, one is proper, one is proper (-1), and one is proper (-2).

In view of the results in Section 3, one can readily verify the following:

If \mathcal{X} is a Chung-Yao lattice then for all nodes the item 1) holds.

If \mathcal{X} is a Carnicer-Gasca lattice then for all 1_m -nodes the item 1) holds. While for all 2_m -nodes the item 2) holds.

If \mathcal{X} is defect 2 set then for the node O the item 1) holds. For all 1_m -nodes the item 2) holds. For all 2_m -nodes the item 3) or 4) holds.

If \mathcal{X} is defect 3 set then for the three *O*-nodes the item 2) holds. For the three *D*-nodes the item 3) holds. For the remaining 1_m -nodes the item 3) or 4) holds. Finally, for all 2_m -nodes one of the items 5), 6) or 7) holds. \Box

References

- [1] V. Bayramyan and H. Hakopian, On a new property of n-correct and GC_n sets, Adv. Comput. Math., **43**, (2017) 607–626.
- [2] J. R. Busch, A note on Lagrange interpolation in ℝ², Rev. Un. Mat. Argentina, **36** (1990) 33–38.
- [3] J. M. Carnicer and M. Gasca, Planar configurations with simple Lagrange formula, in *Mathematical Methods in CAGD*, eds. T. Lyche and L. L. Schumaker (Vanderbilt University Press, Nashville, Oslo 2000, TN, 2001) 55—62.
- [4] J. M. Carnicer and M. Gasca, A conjecture on multivariate polynomial interpolation, Rev. R. Acad. Cienc. Exactas Fís. Nat. (Esp.), Ser. A Mat. 95 (2001) 145–153.
- [5] J. M. Carnicer and M. Gasca, On Chung and Yao's geometric characterization for bivariate polynomial interpolation, in *Curve and Surface Design*, eds. T. Lyche, M.-L. Mazure, and L. L. Schumaker (Nashboro Press, Saint Malo 2002, Brentwood, 2003) 21–30.
- [6] J. M. Carnicer and M. Gasca, Generation of lattices of points for bivariate interpolation, Numer. Algorithms, **39** (2005) 69–79.
- [7] J. M. Carnicer and C. Godés, Geometric characterization and generalized principal lattices, J. Approx. Theory, 143 (2006) 2–14.
- [8] J. M. Carnicer and C. Godés, Geometric characterization of configurations with defect three, in *Curve and Surface Fitting*, eds. Albert Cohen,

Jean Louis Merrien, Larry L. Schumaker (Nashboro Press, Brentwood, Avignon 2006, TN, 2007) 61–70.

- [9] J. M. Carnicer and C. Godés, Configurations of nodes with defects greater than three. J. Comput. Appl. Math., 233 (2010) 1640–1648.
- [10] K. C. Chung and T. H. Yao, On lattices admitting unique Lagrange interpolations, SIAM J. Numer. Anal., 14 (1977) 735–743.
- [11] M. Gasca and J. I. Maeztu, On Lagrange and Hermite interpolation in \mathbb{R}^k , Numer. Math., **39** (1982) 1–14.
- [12] H. Hakopian, K. Jetter, and G. Zimmermann, The Gasca-Maeztu conjecture for n = 5, Numer. Math., **127** (2014) 685–713.
- [13] H. Hakopian and L. Rafayelyan, On a generalization of Gasca-Maeztu conjecture, New York J. Math., 21 (2015) 351–367.
- [14] H. Hakopian, V. Vardanyan, On a correction of a property of GC_n sets, Adv. Comput. Math., **45**, (2019) 311–325.
- [15] H. Hakopian, V. Vardanyan, On the usage of lines in GC_n sets, Adv. Comput. Math., **45**, (2019) 2721–2743.