

Biorthogonal greedy algorithms in convex optimization

A. V. Dereventsov^{a,*}, V. N. Temlyakov^b

^a*Oak Ridge National Laboratory*

^b*University of South Carolina, Steklov Institute of Mathematics, and Lomonosov Moscow State University*

Abstract

The study of greedy approximation in the context of convex optimization is becoming a promising research direction as greedy algorithms are actively being employed to construct sparse minimizers for convex functions with respect to given sets of elements. In this paper we propose a unified way of analyzing a certain kind of greedy-type algorithms for the minimization of convex functions on Banach spaces. Specifically, we define the class of Weak Biorthogonal Greedy Algorithms for convex optimization that contains a wide range of greedy algorithms. We analyze the introduced class of algorithms and establish the properties of convergence, rate of convergence, and numerical stability, which is understood in the sense that the steps of the algorithm are allowed to be performed not precisely but with controlled computational inaccuracies. We show that the following well-known algorithms for convex optimization — the Weak Chebyshev Greedy Algorithm (co) and the Weak Greedy Algorithm with Free Relaxation (co) — belong to this class, and introduce a new algorithm — the Rescaled Weak Relaxed Greedy Algorithm (co). Presented numerical experiments demonstrate the practical performance of the aforementioned greedy algorithms in the setting of convex minimization as compared to optimization with regularization, which is the conventional approach of constructing sparse minimizers.

Keywords: greedy algorithm, convex optimization, sparsity, Biorthogonal Greedy Algorithm, Banach space

1. Introduction

This paper is devoted to the question of building an approximate sparse solution of a given convex optimization problem. A typical setting is to find an approximate minimum of a real-valued convex function E defined on the

*Corresponding author

Email addresses: dereventsov@gmail.com (A. V. Dereventsov), temlyakovv@gmail.com (V. N. Temlyakov)

Banach space X , i.e.

$$\text{find } x^* = \underset{x \in X}{\operatorname{argmin}} E(x). \quad (1.1)$$

When optimization is performed over the whole space X , it is called an *unconstrained optimization problem*. Usually in practice it is desirable to obtain a minimizer that possesses a certain structure or belongs to a given domain $S \subset X$, in which case problem (1.1) becomes a *constrained optimization problem*.

In particular, it is often preferable that the constructed solution x^* is sparse with respect to a given set of elements $\mathcal{D} \subset X$. A conventional approach to such a task is to impose an additional ℓ_1 -regularization on the original problem (see e.g. [14]) and, instead of (1.1), solve the problem

$$\text{find } x^* = \underset{x \in X}{\operatorname{argmin}} (E(x) + \lambda \|x\|_{\mathcal{D}}), \quad (1.2)$$

where $\lambda > 0$ is an appropriate regularization parameter and $\|\cdot\|_{\mathcal{D}}$ is the atomic norm with respect to the set \mathcal{D} (see e.g. [6]), i.e.

$$\|x\|_{\mathcal{D}} := \inf \left\{ \sum_{g \in \mathcal{D}} |c_g| : x = \sum_{g \in \mathcal{D}} c_g g \right\}. \quad (1.3)$$

While such an approach is quite popular and has its uses, it might not always result in the most appropriate solution since it essentially changes the target function in order to promote sparsity of the solution.

Another way of obtaining a sparse minimizer (without changing the optimization problem) is to procedurally construct a sequence of minimizers with an increasing support, or, more generally, to design an algorithm that after m iterations provides a point x_m such that $E(x_m)$ is close to the $\inf_{x \in S} E(x)$ and that x_m is m -sparse with respect to \mathcal{D} , i.e.

$$x_m = \sum_{j=1}^m c_j g_j \quad \text{with } g_1, \dots, g_m \in \mathcal{D} \quad \text{and } c_1, \dots, c_m \in \mathbb{R}.$$

A wide class of algorithms that fit such requirements is the greedy algorithms in approximation theory, see e.g. [11], [26]. A typical problem of greedy approximation is the following. Let X be a Banach space with the norm $\|\cdot\|$ and let \mathcal{D} be a dictionary, i.e. a dense set of semi-normalized elements of X . The goal of a greedy algorithm is to obtain a sparse (with respect to the dictionary \mathcal{D}) approximation of a given element $f \in X$. Greedy algorithms are iterative by design and generally after m iterations a greedy algorithm constructs an m -term linear combination with respect to \mathcal{D} that approximates the element f .

It is easy to reframe a greedy approximation problem as a convex optimization problem. Indeed, for a given dictionary \mathcal{D} consider the set of all m -term linear combinations with respect to \mathcal{D} (m -sparse with respect to \mathcal{D} elements):

$$\Sigma_m(\mathcal{D}) := \left\{ x \in X : x = \sum_{j=1}^m c_j g_j, \quad g_1, \dots, g_m \in \mathcal{D} \right\}.$$

Greedy algorithms in approximation theory are designed to provide a simple way to build good approximants of f from $\Sigma_m(\mathcal{D})$, hence the problem of greedy approximation is the following:

$$\text{find } x_m = \underset{x \in \Sigma_m}{\operatorname{argmin}} \|f - x\|. \quad (1.4)$$

Clearly, problem (1.4) is a constrained optimization problem of the real-valued convex function $E(x) := \|f - x\|$ over the manifold $\Sigma_m(\mathcal{D}) \subset X$.

At first glance the settings of approximation and optimization problems appear to be very different since in approximation theory our task is to find a sparse approximation of a given element $f \in X$, while in optimization theory we want to find an approximate sparse minimizer of a given target function $E : X \rightarrow \mathbb{R}$ (for instance, energy function or loss function). However it is now well understood that similar techniques can be used for solving both problems. Namely, it was shown in [27] and in follow up papers (see, for instance, [12], [16], [21], [28], and [29]) how methods developed in nonlinear approximation theory (greedy approximation techniques in particular) can be adjusted to find an approximate sparse (with respect to a given dictionary \mathcal{D}) solution to the optimization problem (1.1). Moreover, there is an increasing interest in building such sparse approximate solutions using different greedy-type algorithms, for example, [2], [3], [6], [7], [18], [19], [23], [30], and [31].

With an established framework it is straightforward to adjust a greedy strategy to a context of convex optimization; however, each of these modified techniques requires an individual analysis to guarantee a desirable performance. On the other hand, it is known that the behavior of a greedy method is largely determined by the underlying geometry of the problem setting. In particular, in [10] we present a unified way of analyzing different greedy-type algorithms in Banach spaces. Specifically, we define the class of Weak Biorthogonal Greedy Algorithms ($\mathcal{WBG\mathcal{A}}$) and prove convergence and rate of convergence results for algorithms from this class. Such an approach allows for a simultaneous analysis of a wide range of seemingly different greedy algorithms based on the smoothness characteristic of the problem.

In this paper we adopt the approach of unified analysis for the setting of convex minimization. In Section 2 we adjust the class $\mathcal{WBG\mathcal{A}}$ of algorithms designed for greedy approximation in Banach spaces and derive the class of Weak Biorthogonal Greedy Algorithms for convex optimization ($\mathcal{WBG\mathcal{A}}(\text{co})$), which consists of greedy algorithms designed for convex optimization. We prove convergence and rate of convergence results for algorithms from the class $\mathcal{WBG\mathcal{A}}(\text{co})$ in Theorems 2.1 and 2.2 respectively. Thus, results in Section 2 address two important characteristics of an algorithm — convergence and rate of convergence.

The rate of convergence is an essential characteristic of an algorithm, though in certain practical applications resistance to various perturbations might be of equal importance. A systematic study of the stability of greedy algorithms in Banach spaces was started in [25] and further advanced in [8], where necessary and sufficient conditions for the convergence of a certain algorithm were ob-

tained. A transition to the optimization setting was performed in [12] and [29], where stability results for greedy-type algorithms for convex optimization were obtained. In Section 3 we discuss the stability of the algorithms from the $\mathcal{WBG}\mathcal{A}(\text{co})$ by analyzing convergence properties of the algorithms from $\mathcal{WBG}\mathcal{A}(\text{co})$ under the assumption of imprecise calculations in the steps of the algorithms. We call such algorithms *approximate greedy algorithms* or *algorithms with errors*. We prove convergence and rate of convergence results for the Weak Biorthogonal Greedy Algorithms with errors, which describes the stability of the algorithms from the class $\mathcal{WBG}\mathcal{A}(\text{co})$ — an important characteristic that is crucial for practical implementation.

Since theoretical analysis cannot always predict the practical behavior of an algorithm, it is of interest to observe its actual implementation for particular problems. In Section 4 we demonstrate the performance of some algorithms from the class $\mathcal{WBG}\mathcal{A}(\text{co})$ by employing them to solve various minimization problems. Additionally, we compare these algorithms with a conventional method of obtaining sparse minimizers — optimization with ℓ_1 -regularization (1.2). Lastly, in Sections 5 and 6 we prove the results stated in Sections 2 and 3 respectively.

2. Weak Biorthogonal Greedy Algorithms for Convex Optimization

In this section we introduce and discuss the class of Weak Biorthogonal Greedy Algorithms for convex optimization, denoted as $\mathcal{WBG}\mathcal{A}(\text{co})$. We begin by recalling the relevant terminology.

2.1. Preliminaries

Let X be a real Banach space with the norm $\|\cdot\|$. We say that a set of elements \mathcal{D} from X is a dictionary if each $g \in \mathcal{D}$ has the norm bounded by one and \mathcal{D} is dense in X , that is

$$\|g\| \leq 1 \text{ for any } g \in \mathcal{D}, \text{ and } \overline{\text{span } \mathcal{D}} = X.$$

For notational convenience in this paper we consider *symmetric dictionaries*, i.e. such that

$$g \in \mathcal{D} \text{ implies } -g \in \mathcal{D}.$$

We denote the closure (in X) of the convex hull of \mathcal{D} by $\mathcal{A}_1(\mathcal{D})$:

$$\mathcal{A}_1(\mathcal{D}) := \overline{\text{conv } \mathcal{D}}, \tag{2.1}$$

which is the standard notation in relevant greedy approximation literature.

The modulus of smoothness $\rho(E, S, u)$ of a function $E : X \rightarrow \mathbb{R}$ on a set $S \subset X$ is defined as

$$\rho(E, S, u) := \frac{1}{2} \sup_{x \in S, \|y\|=1} \left| E(x + uy) + E(x - uy) - 2E(x) \right|. \tag{2.2}$$

We note that, in comparison to the modulus of smoothness of a norm (see, for instance, [1, Part 3]), the modulus of smoothness of a function additionally

depends on the chosen set $S \subset X$. That is because a norm is a positive homogeneous function, thus its smoothness on the whole space is determined by its smoothness on the unit sphere, which is not the case for a general function on a Banach space.

The function E is uniformly smooth on $S \subset X$ if $\rho(E, S, u) = o(u)$ as $u \rightarrow 0$. We say that the modulus of smoothness $\rho(E, S, u)$ is of power type $1 \leq q \leq 2$ if $\rho(E, S, u) \leq \gamma u^q$ for some $\gamma > 0$. Note that the class of functions with the modulus of smoothness of a nontrivial power type is completely different from the class of uniformly smooth Banach spaces with the norms of a nontrivial power type since any uniformly smooth norm is not uniformly smooth as a function on any set containing 0. However, it is shown in [4] that if a norm $\|\cdot\|$ has the modulus of smoothness of power type $q \in [1, 2]$, then the function $E(\cdot) := \|\cdot\|^q$ has the modulus of smoothness $\rho(E, S, u)$ of power type q for any set $S \subset X$. In particular, it implies (see e.g. [13, Lemma B.1]) that for any $1 \leq p < \infty$ the function $E : L_p \rightarrow \mathbb{R}$ defined as

$$E_p(x) = \|x\|_{L_p}^p$$

has the modulus of smoothness that satisfies

$$\rho_p(E, X, u) \leq \begin{cases} \frac{1}{p} u^p & 1 \leq p \leq 2, \\ \frac{p-1}{2} u^2 & 2 \leq p < \infty, \end{cases}$$

i.e. $\rho_p(E, X, u)$ is of power type $\min\{p, 2\}$.

A typical smoothness assumption in convex optimization is of the form

$$|E(x + uy) - E(x) - \langle E'(x), uy \rangle| \leq Cu^2$$

with some constant $C > 0$ and any $u \in \mathbb{R}$, $x \in X$, $\|y\| = 1$. In terms of the modulus of smoothness (2.2) such an assumption corresponds to the case $\rho(E, X, u) \leq Cu^2/2$, i.e. that the modulus of smoothness of E is of power type 2.

Throughout the paper we assume that the target function E is Fréchet-differentiable, i.e. that at any $x \in X$ there is a bounded linear functional $E'(x) : X \rightarrow \mathbb{R}$ such that

$$\sup_{\|y\|=1} \left(\lim_{u \rightarrow 0} \frac{E(x + uy) - E(x)}{u} - \langle E'(x), y \rangle \right) = 0.$$

Then the convexity of E implies that for any $x, y \in D$

$$E(y) \geq E(x) + \langle E'(x), y - x \rangle, \quad (2.3)$$

or, equivalently,

$$E(x) - E(y) \leq \langle E'(x), x - y \rangle = \langle -E'(x), y - x \rangle. \quad (2.4)$$

Remark. *The condition of Fréchet-differentiability is not necessary and can be relaxed by considering support functionals in place of the derivative of E , as is done in [9, Chapter 5]. Although the existence of support functionals is guaranteed by the convexity of the target function, we additionally impose the assumption of differentiability for the convenience of presentation.*

2.2. Weak Biorthogonal Greedy Algorithms

Typically in greedy approximation one has to perform a greedy selection from a given dictionary \mathcal{D} , which might not always be possible. In order to guarantee the feasibility of algorithms, it is conventional to perform a *weak* greedy step where the greedy search is relaxed. Such relaxations are represented by a given sequence $\tau := \{t_m\}_{m=1}^\infty$, referred to as a *weakness sequence*.

For a convex Fréchet-differentiable target function $E : X \rightarrow \mathbb{R}$ we define the following class of greedy algorithms.

Weak Biorthogonal Greedy Algorithms ($\mathcal{WBG}\mathcal{A}(\text{co})$).

We say that an algorithm belongs to the class $\mathcal{WBG}\mathcal{A}(\text{co})$ with a weakness sequence $\tau = \{t_m\}_{m=1}^\infty$, $t_m \in [0, 1]$, if sequences of approximators $\{G_m\}_{m=0}^\infty$ and selected elements $\{\varphi_m\}_{m=1}^\infty$ of the dictionary \mathcal{D} satisfy the following conditions at every iteration $m \geq 1$:

- (1) Greedy selection: $\langle -E'(G_{m-1}), \varphi_m \rangle \geq t_m \sup_{\varphi \in \mathcal{D}} \langle -E'(G_{m-1}), \varphi \rangle$;
- (2) Error reduction: $E(G_m) \leq \inf_{\lambda \geq 0} E(G_{m-1} + \lambda \varphi_m)$;
- (3) Biorthogonality: $\langle E'(G_m), G_m \rangle = 0$.

We assume that for a given target function $E : X \rightarrow \mathbb{R}$ the set

$$D = D(E) := \{x \in X : E(x) \leq E(0)\} \subset X$$

is bounded. Coupled with the assumption that E is convex and Fréchet-differentiable, boundedness of D guarantees that

$$\inf_{x \in X} E(x) = \inf_{x \in D} E(x) > -\infty \quad \text{and} \quad \operatorname{argmin}_{x \in X} E(x) = \operatorname{argmin}_{x \in D} E(x) \in D,$$

i.e. there is a nontrivial and attainable minimum of E . Then by condition (2) the sequence of m -sparse approximants $\{G_m\}_{m=0}^\infty$ constructed by an algorithm from the $\mathcal{WBG}\mathcal{A}(\text{co})$ satisfies the relation

$$E(0) = E(G_0) \geq E(G_1) \geq E(G_2) \geq \dots,$$

which guarantees that $G_m \in D$ for all $m \geq 0$.

Remark. In the case $E(x) := \|f - x\|^q$ with any $f \in X$ and $q \geq 1$, the class $\mathcal{WBG}\mathcal{A}(\text{co})$ coincides with the class $\mathcal{WBG}\mathcal{A}$ from the approximation theory, which is introduced and analyzed in [10].

2.3. Examples of algorithms from the $\mathcal{WBG}\mathcal{A}(\text{co})$

In this section we briefly overview a few particular algorithms from the class $\mathcal{WBG}\mathcal{A}(\text{co})$ that will be utilized in the numerical experiments presented in Section 4. By $\tau := \{t_m\}_{m=1}^\infty$ we denote a weakness sequence, i.e. a given sequence of non-negative numbers $t_m \leq 1$, $m = 1, 2, 3, \dots$

We first define the Weak Chebyshev Greedy Algorithm for convex optimization that is introduced and studied in [27].

Weak Chebyshev Greedy Algorithm (WCGA(co)).

Set $G_0^c = 0$ and for each $m \geq 1$ perform the following steps:

1. Take any $\varphi_m^c \in \mathcal{D}$ satisfying

$$\langle -E'(G_{m-1}^c), \varphi_m^c \rangle \geq t_m \sup_{\varphi \in \mathcal{D}} \langle -E'(G_{m-1}^c), \varphi \rangle;$$

2. Denote $\Phi_m^c = \text{span}\{\varphi_k^c\}_{k=1}^m$ and find $G_m^c \in \Phi_m^c$ such that

$$E(G_m^c) = \inf_{G \in \Phi_m^c} E(G).$$

Another algorithm, which utilizes a simpler approach to updating the approximant is the Weak Greedy Algorithm with Free Relaxation for convex optimization (see [27]).

Weak Greedy Algorithm with Free Relaxation (WGAFR(co)).

Set $G_0^f = 0$ and for each $m \geq 1$ perform the following steps:

- (1) Take any $\varphi_m^f \in \mathcal{D}$ satisfying

$$\langle -E'(G_{m-1}^f), \varphi_m^f \rangle \geq t_m \sup_{\varphi \in \mathcal{D}} \langle -E'(G_{m-1}^f), \varphi \rangle;$$

- (2) Find $\omega_m \in \mathbb{R}$ and $\lambda_m \in \mathbb{R}$ such that

$$E((1 - \omega_m)G_{m-1}^f + \lambda_m \varphi_m^f) = \inf_{\lambda, \omega \in \mathbb{R}} E((1 - \omega)G_{m-1}^f + \lambda \varphi_m^f)$$

$$\text{and define } G_m^f = (1 - \omega_m)G_{m-1}^f + \lambda_m \varphi_m^f.$$

The next algorithm — the Rescaled Weak Relaxed Greedy Algorithm for convex optimization — is an adaptation of its counterpart from the approximation theory (see [10]) that can be viewed as a generalization of the Rescaled Pure Greedy Algorithm, introduced in [22] and adapted for convex optimization in [16].

Rescaled Weak Relaxed Greedy Algorithm (RWRGA(co)).

Set $G_0^r = 0$ and for each $m \geq 1$ perform the following steps:

1. Take any $\varphi_m^r \in \mathcal{D}$ satisfying

$$\langle -E'(G_{m-1}^r), \varphi_m^r \rangle \geq t_m \sup_{\varphi \in \mathcal{D}} \langle -E'(G_{m-1}^r), \varphi \rangle;$$

2. Find $\lambda_m \geq 0$ such that

$$E(G_{m-1}^r + \lambda_m \varphi_m^r) = \inf_{\lambda \geq 0} E(G_{m-1}^r + \lambda \varphi_m^r);$$

3. Find $\mu_m \in \mathbb{R}$ such that

$$E(\mu_m(G_{m-1}^r + \lambda_m \varphi_m^r)) = \inf_{\mu \in \mathbb{R}} E(\mu(G_{m-1}^r + \lambda_m \varphi_m^r))$$

and define $G_m^r = \mu_m(G_{m-1}^r + \lambda_m \varphi_m^r)$.

Proposition 2.1. *The WCGA(co), the WGAFR(co), and the RWRGA(co) belong to the class $\mathcal{WBGA}(co)$.*

2.4. Convergence results for the $\mathcal{WBGA}(co)$

In this section we state the results related to convergence and the rate of convergence for algorithms from the class $\mathcal{WBGA}(co)$.

Our setting of an infinite dimensional Banach space makes the formulation of convergence results nontrivial, and thus we require a special sequence which is defined for a given modulus of smoothness $\rho(u) := \rho(E, D, u)$ and a given weakness sequence $\tau = \{t_m\}_{m=1}^\infty$.

Let $E : X \rightarrow \mathbb{R}$ be a convex uniformly smooth function, then $\rho(u) := \rho(E, D, u) : \mathbb{R} \rightarrow \mathbb{R}_+$ is an even convex function. Assume that $\rho(u)$ has the property $\rho(1/\theta_0) \geq 1$ for some $\theta_0 \in (0, 1]$ and

$$\lim_{u \rightarrow 0} \rho(u)/u = 0.$$

Note that assumptions on uniform smoothness of E and boundedness of domain $D \subset X$ guarantee the above properties. Then for a given $0 < \theta \leq \theta_0$ define $\xi_m := \xi_m(\rho, \tau, \theta)$ as the solution of the equation

$$\rho(u) = \theta t_m u. \tag{2.5}$$

Note that conditions on $\rho(u)$ imply that the function

$$s(u) := \begin{cases} \rho(u)/u, & u \neq 0 \\ 0, & u = 0 \end{cases}$$

is continuous and increasing on $[0, \infty)$ with $s(1/\theta_0) \geq \theta_0$. Thus equation (2.5) has the unique solution $\xi_m = s^{-1}(\theta t_m)$ such that $0 < \xi_m \leq 1/\theta_0$.

We now formulate our main convergence result for the $\mathcal{WBGA}(co)$.

Theorem 2.1. *Let E be a uniformly smooth on $D \subset X$ convex function with the modulus of smoothness $\rho(E, D, u)$. Assume that a sequence $\tau := \{t_m\}_{m=1}^\infty$ satisfies the condition that for any $\theta \in (0, \theta_0]$ we have*

$$\sum_{m=1}^{\infty} t_m \xi_m(\rho, \tau, \theta) = \infty.$$

Then for any algorithm from the class $\mathcal{WBGA}(co)$ we have

$$\lim_{m \rightarrow \infty} E(G_m) = \inf_{x \in D} E(x).$$

Here are two simple corollaries of Theorem 2.1.

Corollary 2.1. *Let E be a uniformly smooth on $D \subset X$ convex function. Then any algorithm from the class $\mathcal{WBG}\mathcal{A}(co)$ with a constant weakness sequence $\tau = t \in (0, 1]$ converges, i.e.*

$$\lim_{m \rightarrow \infty} E(G_m) = \inf_{x \in D} E(x).$$

Corollary 2.2. *Let E be a convex function with the modulus of smoothness of power type $1 < q \leq 2$, that is, $\rho(E, D, u) \leq \gamma u^q$. Let a sequence $\tau := \{t_m\}_{m=1}^{\infty}$, $t_m \in (0, 1]$ for $m = 1, 2, 3, \dots$ be such that*

$$\sum_{m=1}^{\infty} t_m^p = \infty, \quad p = \frac{q}{q-1}.$$

Then any algorithm from the class $\mathcal{WBG}\mathcal{A}(co)$ with the weakness sequence τ converges, i.e.

$$\lim_{m \rightarrow \infty} E(G_m) = \inf_{x \in D} E(x).$$

We now proceed to the rate of convergence estimates, which are of interest in both finite dimensional and infinite dimensional settings. A typical assumption in this regard is formulated in terms of the convex hull $\mathcal{A}_1(\mathcal{D})$ of the dictionary \mathcal{D} , defined by (2.1).

Theorem 2.2. *Let E be a convex function with the modulus of smoothness of power type $1 < q \leq 2$, that is, $\rho(E, D, u) \leq \gamma u^q$. Take an element $f^\epsilon \in D$ and a number $\epsilon \geq 0$ such that*

$$E(f^\epsilon) \leq \inf_{x \in D} E(x) + \epsilon, \quad f^\epsilon / A(\epsilon) \in \mathcal{A}_1(\mathcal{D})$$

with some number $A(\epsilon) \geq 1$. Then for any algorithm from the class $\mathcal{WBG}\mathcal{A}(co)$ we have

$$E(G_m) - \inf_{x \in D} E(x) \leq \max \left\{ 2\epsilon, C(q, \gamma) A(\epsilon)^q \left(C(E, q, \gamma) + \sum_{k=1}^m t_k^p \right)^{1-q} \right\},$$

where $p = q/(q-1)$.

Corollary 2.3. *Let E be a convex function with the modulus of smoothness of power type $1 < q \leq 2$, that is, $\rho(E, D, u) \leq \gamma u^q$. If $\operatorname{argmin}_{x \in D} E(x) \in \mathcal{A}_1(\mathcal{D})$ then for any algorithm from the class $\mathcal{WBG}\mathcal{A}(co)$ we have*

$$E(G_m) - \inf_{x \in D} E(x) \leq C(q, \gamma) \left(C(E, q, \gamma) + \sum_{k=1}^m t_k^p \right)^{1-q},$$

where $p = q/(q-1)$.

Remark. *While the results stated in this section are known for the $WCGA(co)$ and the $WGAFR(co)$ (see [27]), they are novel for the $RWRGA(co)$.*

3. Weak Biorthogonal Greedy Algorithms with errors for Convex Optimization

In this section we address the question of the stability of algorithms from the class $\mathcal{WBG}\mathcal{A}(\text{co})$ by introducing the wider class $\mathcal{WBG}\mathcal{A}(\Delta, \text{co})$, which allows for imprecise calculations in the realization of algorithms. Such an approach is of a practical interest since computational inaccuracies often occur naturally in applications. To account for imprecise computations we introduce a sequence $\Delta := \{\delta_m, \epsilon_m\}_{m=1}^{\infty}$, where $\delta_m \in [0, 1]$ and $\epsilon_m \geq 0$ for $m = 1, 2, 3, \dots$, that represents the allowed inaccuracies in the steps of the algorithms. In accordance with the conventional notation (see e.g. [17], [15]), we refer to a given sequence $\Delta := \{\delta_m, \epsilon_m\}_{m=1}^{\infty}$ as an *error sequence*.

For a convex Fréchet-differentiable target function $E : X \rightarrow \mathbb{R}$ we define the following class of greedy algorithms with errors.

Weak Biorthogonal Greedy Algorithms with errors ($\mathcal{WBG}\mathcal{A}(\Delta, \text{co})$).

We say that an algorithm belongs to the class $\mathcal{WBG}\mathcal{A}(\Delta, \text{co})$ with a weakness sequence $\tau = \{t_m\}_{m=1}^{\infty}$, $t_m \in [0, 1]$ and an error sequence $\Delta = \{\delta_m, \epsilon_m\}_{m=1}^{\infty}$, $\delta_m \in [0, 1]$, $\epsilon_m \geq 0$, if sequences of approximators $\{G_m\}_{m=0}^{\infty}$ and selected elements $\{\varphi_m\}_{m=1}^{\infty}$ of the dictionary \mathcal{D} satisfy the following conditions at every iteration $m \geq 1$:

- (1) Greedy selection: $\langle -E'(G_{m-1}), \varphi_m \rangle \geq t_m \sup_{\varphi \in \mathcal{D}} \langle -E'(G_{m-1}), \varphi \rangle$;
- (2) Error reduction: $E(G_m) \leq \inf_{\lambda \geq 0} E(G_{m-1} + \lambda \varphi_m) + \delta_m$;
- (3) Biorthogonality: $|\langle E'(G_m), G_m \rangle| \leq \epsilon_m$;
- (4) Boundedness: $E(G_m) \leq E(0) + C_0$.

Note that in addition to conditions (1)–(3) from the definition of the class $\mathcal{WBG}\mathcal{A}(\text{co})$, for the $\mathcal{WBG}\mathcal{A}(\Delta, \text{co})$ we require the boundedness condition (4) to account for the magnitude of allowed errors Δ . In particular, if the error sequence Δ is summable, i.e. $\sum_{m=1}^{\infty} \delta_m < \infty$, then condition (4) follows directly from (2) with $C_0 = \sum_{m=1}^{\infty} \delta_m$.

Moreover, we assume that the set

$$D \subset D_1 := \{x \in X : E(x) \leq E(0) + C_0\} \subset X,$$

where C_0 is the constant from condition (4), is bounded. Then condition (4) guarantees that $G_m \in D_1$ for all $m \geq 0$ for any algorithm from the $\mathcal{WBG}\mathcal{A}(\Delta, \text{co})$.

Remark. In the error reduction condition (2) from the definition of the class $\mathcal{WBG}\mathcal{A}(\Delta, \text{co})$ the infimum is taken over all $\lambda \geq 0$. In order to simplify this problem, one can consider a wider than the $\mathcal{WBG}\mathcal{A}(\Delta, \text{co})$ class — the class $\mathcal{WBG}\mathcal{A}(\Delta, [0, 1], \text{co})$ of algorithms satisfying conditions (1), (3), (4), and the following condition instead of (2):

- (2') Restricted error reduction: $E(G_m) \leq \inf_{\lambda \in [0, 1]} E(G_{m-1} + \lambda \varphi_m) + \delta_m$.

Then finding such $\lambda \in [0, 1]$ is a line search problem, which is known to be a simple one-dimensional convex optimization problem (see e.g. [5], [20]).

3.1. Examples of algorithms from the $\mathcal{WBG}\mathcal{A}(\Delta, \text{co})$

In this section we briefly overview particular algorithms from the class $\mathcal{WBG}\mathcal{A}(\Delta, \text{co})$ that correspond to the approximate versions of the algorithms considered in Section 2.3. Denote by $\tau := \{t_m\}_{m=1}^\infty$ and $\Delta := \{\delta_m, \epsilon_m\}_{m=1}^\infty$ a weakness sequence and an error sequence respectively, i.e. given sequences of numbers $t_m \in [0, 1]$, $\delta_m \in [0, 1]$, and $\epsilon_m \geq 0$ for $m = 1, 2, 3, \dots$

We begin with the Weak Chebyshev Greedy Algorithm with errors for convex optimization.

Weak Chebyshev Greedy Algorithm with errors (WCGA(Δ, co)).

Set $G_0^c = 0$ and for each $m \geq 1$ perform the following steps:

1. Take any $\varphi_m^c \in \mathcal{D}$ satisfying

$$\langle -E'(G_{m-1}^c), \varphi_m^c \rangle \geq t_m \sup_{\varphi \in \mathcal{D}} \langle -E'(G_{m-1}^c), \varphi \rangle;$$

2. Denote $\Phi_m^c = \text{span}\{\varphi_k^c\}_{k=1}^m$ and find $G_m^c \in \Phi_m^c$ such that

$$E(G_m^c) \leq \inf_{G \in \Phi_m^c} E(G) + \delta_m.$$

Next, we state the Weak Greedy Algorithm with Free Relaxation and errors for convex optimization, introduced and studied in [12].

Weak Greedy Algorithm with Free Relaxation and errors (WGAFR(Δ, co)).

Set $G_0^f = 0$ and for each $m \geq 1$ perform the following steps:

1. Take any $\varphi_m^f \in \mathcal{D}$ satisfying

$$\langle -E'(G_{m-1}^f), \varphi_m^f \rangle \geq t_m \sup_{\varphi \in \mathcal{D}} \langle -E'(G_{m-1}^f), \varphi \rangle;$$

2. Find $\omega_m \in \mathbb{R}$ and $\lambda_m \in \mathbb{R}$ such that

$$E((1 - \omega_m)G_{m-1}^f + \lambda_m \varphi_m^f) \leq \inf_{\lambda, \omega \in \mathbb{R}} E((1 - \omega)G_{m-1}^f + \lambda \varphi_m^f) + \delta_m$$

$$\text{and define } G_m^f = (1 - \omega_m)G_{m-1}^f + \lambda_m \varphi_m^f.$$

Lastly, we introduce a new algorithm — the Rescaled Weak Relaxed Greedy Algorithm with errors for convex optimization.

Rescaled Weak Relaxed Greedy Algorithm with errors (RWRGA(Δ, co)).

Set $G_0^r = 0$ and for each $m \geq 1$ perform the following steps:

1. Take any $\varphi_m^r \in \mathcal{D}$ satisfying

$$\langle -E'(G_{m-1}^r), \varphi_m^r \rangle \geq t_m \sup_{\varphi \in \mathcal{D}} \langle -E'(G_{m-1}^r), \varphi \rangle;$$

2. Find $\lambda_m \geq 0$ such that

$$E(G_{m-1}^r + \lambda_m \varphi_m^r) \leq \inf_{\lambda \geq 0} E(G_{m-1}^r + \lambda \varphi_m^r) + \delta_m/2;$$

3. Find $\mu_m \in \mathbb{R}$ such that

$$E(\mu_m(G_{m-1}^r + \lambda_m \varphi_m^r)) \leq \inf_{\mu \in \mathbb{R}} E(\mu(G_{m-1}^r + \lambda_m \varphi_m^r)) + \delta_m/2$$

and define $G_m^r = \mu_m(G_{m-1}^r + \lambda_m \varphi_m^r)$.

Proposition 3.1. *The WCGA(Δ, co), the WGAFR(Δ, co), and the RWRGA(Δ, co) belong to the class $\mathcal{WBGA}(\Delta, co)$ with*

$$\epsilon_m = \inf_{u > 0} \frac{\delta_m + 2\rho(E, D_1, u \|G_m\|)}{u}.$$

3.2. *Convergence results for the $\mathcal{WBGA}(\Delta, co)$*

In this section we discuss the convergence and rate of convergence results for algorithms from the class $\mathcal{WBGA}(\Delta, co)$.

First, we state the convergence result.

Theorem 3.1. *Let E be a uniformly smooth on $D_1 \subset X$ convex function. Assume that an error sequence $\Delta := \{\delta_m, \epsilon_m\}_{m=1}^\infty$ is such that $\delta_m \rightarrow 0$ and $\epsilon_m \rightarrow 0$ as $m \rightarrow \infty$. Then any algorithm from the class $\mathcal{WBGA}(\Delta, co)$ with a constant weakness sequence $\tau = t \in (0, 1]$ converges, i.e.*

$$\lim_{m \rightarrow \infty} E(G_m) = \inf_{x \in D_1} E(x).$$

Second, we provide the rate of convergence estimate.

Theorem 3.2. *Let E be a convex function with the modulus of smoothness of power type $1 < q \leq 2$, that is, $\rho(E, D_1, u) \leq \gamma u^q$. Take an element $f^\epsilon \in D_1$ and a number $\epsilon \geq 0$ such that*

$$E(f^\epsilon) \leq \inf_{x \in D_1} E(x) + \epsilon, \quad f^\epsilon/A \in \mathcal{A}_1(\mathcal{D}),$$

with some number $A := A(\epsilon) \geq 1$. Then for any algorithm from the class $\mathcal{WBGA}(\Delta, co)$ with a constant weakness sequence $\tau = t \in (0, 1]$ and an error sequence $\Delta = \{\delta_m, \epsilon_m\}_{m=1}^\infty$ with $\delta_m + \epsilon_m \leq cm^{-q}$, $m = 1, 2, 3, \dots$ we have

$$E(G_m) - \inf_{x \in D_1} E(x) \leq \epsilon + C(E, q, \gamma, t, c)A(\epsilon)^q m^{1-q}.$$

Corollary 3.1. *Under the conditions of Theorem 3.2, specifying*

$$A(\epsilon) := \inf \left\{ A > 0 : \exists f \in D_1 : f/A \in \mathcal{A}_1(\mathcal{D}), \quad E(f) \leq \inf_{x \in D_1} E(x) + \epsilon \right\}$$

and denoting

$$\eta_m := \inf \{ \epsilon > 0 : A(\epsilon)^q m^{1-q} \leq \epsilon \},$$

we obtain for any algorithm from the class $\mathcal{WBGA}(\Delta, co)$

$$E(G_m) - \inf_{x \in D_1} E(x) \leq C(E, q, \gamma, t) \eta_m.$$

Remark. *It follows from the proofs of Theorems 3.1 and 3.2, given in Section 6, that the results stated in this section also hold for the class $\mathcal{WBGA}(\Delta, [0, 1], co)$.*

4. Numerical experiments

In this section we demonstrate the performance of the algorithms from the class $\mathcal{WBGA}(co)$ that are discussed in Section 2.3: the Weak Chebyshev Greedy Algorithm ($WCGA(co)$), the Weak Greedy Algorithm with Free Relaxation ($WGAFR(co)$), and the Rescaled Weak Relaxed Greedy Algorithm ($RWRGA(co)$).

For each of the numerical experiments presented below we consider the Banach space $X = \ell_1^{(\dim)}$ of dimensionality \dim , a target function $E : X \rightarrow \mathbb{R}$, and a dictionary $\mathcal{D} \in X$. We then employ the aforementioned algorithms to solve the optimization problem (1.1), i.e. to find a sparse (with respect to the dictionary \mathcal{D}) minimizer

$$x^* = \operatorname{argmin}_{x \in X} E(x).$$

Since greedy algorithms are iterative by design, in Examples 1–2 we obtain and present the trade-off between the sparsity of the solution x^* and the value of $E(x^*)$. In Examples 3–4, we additionally compare the greedy algorithms for convex optimization with a conventional method of finding sparse solutions — the optimization with ℓ_1 -regularization, see (1.2). Specifically, we solve the problem

$$\text{find } x^* = \operatorname{argmin}_{x \in X} \left(E(x) + \lambda \|x\|_{\mathcal{D}} \right),$$

where $\|\cdot\|_{\mathcal{D}}$ is the atomic norm with respect to the dictionary \mathcal{D} , defined by (1.3). To obtain minimizers of different sparsities, the values of the regularization parameter λ are taken from the sequence $\{0.1 \times (0.9)^k\}_{k=0}^{49}$, i.e. 50 regularized optimization problems are solved in every setting.

To avoid an unintentional bias in the selection of dictionary \mathcal{D} and target function E , we generate those randomly, based on certain parameters that are described in the setting of each example. In order to provide a reliable demonstration that is independent of a particular random generation, we compute 100 simulations for each presented example and provide the distribution of the optimization results (shown in Figures 1–4). In the presented pictures the solid lines represent the mean minimization values for each algorithm and the filled areas represent the minimization distribution across all 100 simulations. Finally, to make the results consistent across simulations, we rescale the optimization results to be in the interval $[0, 1]$, i.e. instead of reporting the value of $E(x^*)$ we report

$$\frac{E(x^*) - \inf_{x \in X} E(x)}{E(0) - \inf_{x \in X} E(x)} \in [0, 1].$$

Numerical experiments presented in this section are performed in Python 3.6 with the use of NumPy and SciPy libraries. The source code is available at https://github.com/sukiboo/wbga_co_2020.




Figure 1: Distribution of optimization results for Example 1.

4.1. Example 1

In this example we consider the space $X = \ell_1^{(500)}$, and construct a dictionary \mathcal{D} of size 1000 as linear combinations of the canonical basis $\{e_i\}_{i=1}^{500}$ of X with uniformly distributed coefficients, i.e.

$$\mathcal{D} = \{\varphi_j\}_{j=1}^{1000}, \text{ where } \varphi_j = \sum_{i=1}^{500} c_j^i e_i \text{ with } c_j^i \sim \mathcal{U}(0, 1).$$

The target function $E : X \rightarrow \mathbb{R}$ is chosen as

$$E(x) = \|x - f\|_p^p,$$

where $p = 1.2$ and $f \in X$ is randomly generated as a linear combination of 60 randomly selected elements of \mathcal{D} with normally distributed coefficients, i.e.

$$f = \sum_{k=1}^{60} a_k \varphi_{\sigma(k)}, \text{ where } a_k \sim \mathcal{N}(0, 1) \text{ and } \sigma \text{ is a permutation of } \{1, \dots, 1000\}.$$

Performance of the greedy algorithms in this setting is presented in Figure 1. The average number of iterations required to obtain a minimizer of sparsity 50 is 185 for the RWRGA(co), 93 for the WGAFR(co), and 50 for the WCGA(co).

4.2. Example 2

In this example we once again consider the space $X = \ell_1^{(500)}$ and a dictionary \mathcal{D} of size 1000, constructed as linear combinations of the canonical basis $\{e_i\}_{i=1}^{500}$ of X with uniformly distributed coefficients, i.e.

$$\mathcal{D} = \{\varphi_j\}_{j=1}^{1000}, \text{ where } \varphi_j = \sum_{i=1}^{500} c_j^i e_i \text{ with } c_j^i \sim \mathcal{U}(0, 1).$$




Figure 2: Distribution of optimization results for Example 2.

The target function $E : X \rightarrow \mathbb{R}$ is chosen as

$$E(x) = \|x - f\|_p^p \|g\|_q^q + \|x - g\|_q^q \|f\|_p^p,$$

where $p = 3$, $q = 1.2$, and the elements $f, g \in X$ are each randomly generated as linear combinations of 30 randomly selected elements of \mathcal{D} with normally distributed coefficients, i.e.

$$f = \sum_{k=1}^{30} a_k^1 \varphi_{\sigma_1(k)} \quad \text{and} \quad g = \sum_{k=1}^{30} a_k^2 \varphi_{\sigma_2(k)},$$

where $a_k^1, a_k^2 \sim \mathcal{N}(0, 1)$ and σ_1, σ_2 are permutations of $\{1, \dots, 1000\}$.

Performance of the greedy algorithms in this setting is presented in Figure 2. The average number of iterations required to obtain a minimizer of sparsity 50 is 125 for the RWRGA(co), 78 for the WGAFR(co), and 50 for the WCGA(co).

4.3. Example 3

In this example we additionally compare the greedy algorithms with conventional optimization with ℓ_1 -regularization, see (1.2). Since obtaining the minimization-sparsity trade-off with ℓ_1 -regularization is more expensive computationally than it is for the greedy algorithms, we restrict ourselves to work in a space of smaller dimensionality. Namely, we consider the space $X = \ell_1^{(100)}$, and construct a dictionary \mathcal{D} of size 200 as linear combinations of the canonical basis $\{e_i\}_{i=1}^{100}$ of X with uniformly distributed coefficients, i.e.

$$\mathcal{D} = \{\varphi_j\}_{j=1}^{200}, \quad \text{where} \quad \varphi_j = \sum_{i=1}^{100} c_j^i e_i \quad \text{with} \quad c_j^i \sim \mathcal{U}(0, 1).$$




Figure 3: Distribution of optimization results for Example 3.

The target function $E : X \rightarrow \mathbb{R}$ is chosen as

$$E(x) = \|x - f\|_p^p \|g\|_q^q + \|x - g\|_q^q \|f\|_p^p,$$

where $p = 4$, $q = 1.5$, and the elements $f, g \in X$ are each randomly generated as linear combinations of 30 randomly selected elements of \mathcal{D} with normally distributed coefficients, i.e.

$$f = \sum_{k=1}^{30} a_k^1 \varphi_{\sigma_1(k)} \quad \text{and} \quad g = \sum_{k=1}^{30} a_k^2 \varphi_{\sigma_2(k)},$$

where $a_k^1, a_k^2 \sim \mathcal{N}(0, 1)$ and σ_1, σ_2 are permutations of $\{1, \dots, 200\}$.

Performance of the greedy algorithms and optimization with ℓ_1 -regularization in this setting is presented in Figure 3. The average number of iterations required to obtain a minimizer of sparsity 50 is 105 for the RWRGA(co), 74 for the WGAFR(co), and 50 for the WCGA(co).

4.4. Example 4

In this example we compare the greedy algorithms with conventional optimization with ℓ_1 -regularization in a classical setting of canonical basis instead of a randomly-generated dictionary. Namely, we consider the space $X = \ell_1^{(200)}$, and set a dictionary \mathcal{D} to be the canonical basis $\{e_i\}_{i=1}^{200}$ of X , i.e.

$$\mathcal{D} = \{e_j\}_{j=1}^{200}, \quad \text{where} \quad e_j = (\underbrace{0, \dots, 0}_{j-1}, 1, \underbrace{0, \dots, 0}_{200-j}).$$

The target function $E : X \rightarrow \mathbb{R}$ is chosen as

$$E(x) = \|x - f\|_p^p \|g\|_q^q + \|x - g\|_q^q \|f\|_p^p,$$




Figure 4: Distribution of optimization results for Example 4.

where $p = 7$, $q = 3$, and f, g are randomly generated as elements of X with normally distributed coefficients, i.e.

$$f = \sum_{k=1}^{200} a_k^1 e_k \quad \text{and} \quad g = \sum_{k=1}^{200} a_k^2 e_k, \quad \text{where} \quad a_k^1, a_k^2 \sim \mathcal{N}(0, 1).$$

Performance of the greedy algorithms and optimization with ℓ_1 -regularization in this setting is presented in Figure 4. Note that in this case all greedy algorithms — the RWRGA(co), the WGAFR(co), and the WCGA(co) — coincide due to the fact that elements of the dictionary \mathcal{D} are mutually disjoint. Hence the number of iterations required to obtain a minimizer of sparsity 100 is exactly 100 for all three greedy algorithms.

5. Proofs for Section 2

In this section we provide the proofs of the results from Section 2. We begin with a known lemma.

Lemma 5.1 ([27, Lemma 6.1]). *Let E be a uniformly smooth Fréchet-differentiable convex function on a Banach space X and L be a finite-dimensional subspace of X . Let x_L denote the point from L at which E attains the minimum, i.e.*

$$x_L = \operatorname{argmin}_{x \in L} E(x) \in L.$$

Then for any $\phi \in L$ we have

$$\langle E'(x_L), \phi \rangle = 0.$$

We now prove that the algorithms stated in Section 2.3 belong to the class $\mathcal{WBGA}(\text{co})$.

Proof of Proposition 2.1. It is easy to see that conditions **(1)** and **(2)** from the definition of the class $\mathcal{WBG}\mathcal{A}(\text{co})$ are satisfied for all three algorithms. Condition **(3)** for any $m \geq 1$ follows directly from Lemma 5.1 with $x_L = \phi = G_m$ and

$$L = \Phi_m^c = \text{span}\{\varphi_1^c, \dots, \varphi_m^c\}$$

for the WCGA(co), and

$$L = \text{span}\{G_{m-1}^f, \varphi_m^f\} \quad \text{or} \quad L = \text{span}\{G_{m-1}^r, \varphi_m^r\}$$

for the WGAFR(co) / RWRGA(co) respectively. \square

We proceed by listing the lemmas that will be utilized later in the proofs of the main results. The following simple lemma is well-known (see, for instance, [27]). For the reader's convenience we present its proof here.

Lemma 5.2 ([27, Lemma 6.3]). *Let E be a Fréchet-differentiable convex function. Then the following inequality holds for any $x \in S \subset X$, $y \in X$, and $u \in \mathbb{R}$*

$$0 \leq E(x + uy) - E(x) - u \langle E'(x), y \rangle \leq 2\rho(E, S, u\|y\|).$$

Proof. The left inequality follows directly from (2.3). Next, from the definition of modulus of smoothness (2.2) it follows that

$$E(x + uy) + E(x - uy) \leq 2(E(x) + \rho(E, S, u\|y\|)).$$

From inequality (2.3) we get

$$E(x - uy) \geq E(x) - u \langle E'(x), y \rangle.$$

Combining the above two estimates, we obtain

$$E(x + uy) \leq E(x) + u \langle E'(x), y \rangle + 2\rho(E, S, u\|y\|),$$

which proves the second inequality. \square

Lemma 5.3 ([26, Lemma 6.10]). *For any bounded linear functional F and any dictionary \mathcal{D} , we have*

$$\sup_{g \in \mathcal{D}} \langle F, g \rangle = \sup_{f \in \mathcal{A}_1(\mathcal{D})} \langle F, f \rangle.$$

The following lemma is similar to the result from [24]. For the reader's convenience we present a brief proof of this lemma here.

Lemma 5.4. *Suppose that a sequence $y_1 \geq y_2 \geq y_3 \geq \dots > 0$ satisfies inequalities*

$$y_k \leq y_{k-1}(1 - w_k y_{k-1}), \quad w_k \geq 0$$

for any $k > n$. Then for any $m > n$ we have

$$\frac{1}{y_m} \geq \frac{1}{y_n} + \sum_{k=n+1}^m w_k.$$

Proof. The proof follows directly from the chain of inequalities

$$\frac{1}{y_k} \geq \frac{1}{y_{k-1}}(1 - w_k y_{k-1})^{-1} \geq \frac{1}{y_{k-1}}(1 + w_k y_{k-1}) = \frac{1}{y_{k-1}} + w_k.$$

□

The following lemma is our key tool for establishing convergence and rate of convergence of algorithms from the class $\mathcal{WBG}\mathcal{A}(\text{co})$.

Lemma 5.5 (Error Reduction Lemma). *Let E be a uniformly smooth on $D \subset X$ convex function with the modulus of smoothness $\rho(E, D, u)$. Take a number $\epsilon \geq 0$ and an element $f^\epsilon \in D$ such that*

$$E(f^\epsilon) \leq \inf_{x \in X} E(x) + \epsilon, \quad f^\epsilon/A \in \mathcal{A}_1(\mathcal{D}),$$

with some number $A := A(\epsilon) \geq 1$. Then for any algorithm from the class $\mathcal{WBG}\mathcal{A}(\text{co})$ we have for any $m \geq 1$

$$\begin{aligned} E(G_m) - E(f^\epsilon) &\leq E(G_{m-1}) - E(f^\epsilon) \\ &\quad + \inf_{\lambda \geq 0} \left(-\lambda t_m A^{-1} (E(G_{m-1}) - E(f^\epsilon)) + 2\rho(E, D, \lambda) \right). \end{aligned}$$

Proof. The main idea of the proof is the same as in the proof of the corresponding one-step improvement inequality for the WCGA (see, for instance, [26, Lemma 6.11]). It follows from **(2)** of the definition of the class $\mathcal{WBG}\mathcal{A}(\text{co})$ that

$$E(0) \geq E(G_1) \geq E(G_2) \dots$$

Thus if $E(G_{m-1}) - E(f^\epsilon) \leq 0$ then the claim of Lemma 5.5 is trivial. Assuming $E(G_{m-1}) - E(f^\epsilon) > 0$, Lemma 5.2 provides for any $\lambda \geq 0$

$$E(G_{m-1} + \lambda \varphi_m) \leq E(G_{m-1}) - \lambda \langle -E'(G_{m-1}), \varphi_m \rangle + 2\rho(E, D, \lambda)$$

and by **(1)** from the definition of the class $\mathcal{WBG}\mathcal{A}(\text{co})$ and Lemma 5.3 we get

$$\begin{aligned} \langle -E'(G_{m-1}), \varphi_m \rangle &\geq t_m \sup_{g \in \mathcal{D}} \langle -E'(G_{m-1}), g \rangle \\ &= t_m \sup_{\phi \in \mathcal{A}_1(\mathcal{D})} \langle -E'(G_{m-1}), \phi \rangle \geq t_m A^{-1} \langle -E'(G_{m-1}), f^\epsilon \rangle. \end{aligned}$$

By **(3)** from the definition of the class $\mathcal{WBG}\mathcal{A}(\text{co})$ and by convexity (2.4) we obtain

$$\langle -E'(G_{m-1}), f^\epsilon \rangle = \langle -E'(G_{m-1}), f^\epsilon - G_{m-1} \rangle \geq E(G_{m-1}) - E(f^\epsilon).$$

Thus, by **(2)** from the definition of the $\mathcal{WBG}\mathcal{A}(\text{co})$ we deduce

$$\begin{aligned} E(G_m) &\leq \inf_{\lambda \geq 0} E(G_{m-1} + \lambda \varphi_m) \\ &\leq E(G_{m-1}) + \inf_{\lambda \geq 0} \left(-\lambda t_m A^{-1} (E(G_{m-1}) - E(f^\epsilon)) + 2\rho(E, D, \lambda) \right), \end{aligned}$$

which proves the lemma. □

Proof of Theorem 2.1. The error reduction property **(2)** of the class $\mathcal{WBG}\mathcal{A}(\text{co})$ implies that the sequence of minimizers $\{G_m\}_{m=0}^\infty$ is in D and the sequence $\{E(G_m)\}_{m=0}^\infty$ is non-increasing. Therefore, we have

$$\lim_{m \rightarrow \infty} E(G_m) = a \geq \inf_{x \in D} E(x).$$

Denote

$$b := \inf_{x \in D} E(x) \quad \text{and} \quad \alpha := a - b.$$

We prove that $\alpha = 0$ by contradiction. Indeed, assume that $\alpha > 0$. Then for any $m \geq 0$ we have

$$E(G_m) - b \geq \alpha.$$

We set $\epsilon = \alpha/2$ and find $f^\epsilon \in D$ such that

$$E(f^\epsilon) \leq b + \epsilon \quad \text{and} \quad f^\epsilon/A \in \mathcal{A}_1(D)$$

with some $A := A(\epsilon) \geq 1$. Then by Lemma 5.5 we get

$$E(G_m) - E(f^\epsilon) \leq E(G_{m-1}) - E(f^\epsilon) + \inf_{\lambda \geq 0} (-\lambda t_m A^{-1} \alpha/2 + 2\rho(E, D, \lambda)).$$

Specify $\theta := \min\{\theta_0, \frac{\alpha}{8A}\}$ and take $\lambda = \xi_m(\rho, \tau, \theta)$ given by (2.5). Then we obtain

$$E(G_m) \leq E(G_{m-1}) - 2\theta t_m \xi_m.$$

The assumption

$$\sum_{m=1}^{\infty} t_m \xi_m = \infty$$

implies a contradiction, which proves the theorem. \square

Proof of Theorem 2.2. Denote

$$a_n := E(G_n) - E(f^\epsilon),$$

then the sequence $\{a_n\}_{n=0}^\infty$ is non-increasing. If for some $n \leq m$ we have $a_n \leq 0$ then $E(G_m) - E(f^\epsilon) \leq 0$, which implies

$$E(G_m) - \inf_{x \in D} E(x) \leq \epsilon,$$

and hence the statement of the theorem holds. Thus we assume that $a_n > 0$ for $n \leq m$. By Lemma 5.5 we have

$$a_m \leq a_{m-1} + \inf_{\lambda \geq 0} \left(-\frac{\lambda t_m a_{m-1}}{B} + 2\gamma \lambda^q \right). \quad (5.1)$$

Choose λ from the equation

$$\frac{\lambda t_m a_{m-1}}{A} = 4\gamma \lambda^q,$$

which implies that

$$\lambda = \left(\frac{t_m a_{m-1}}{4\gamma A} \right)^{\frac{1}{q-1}}.$$

Let

$$A_q := 2(4\gamma)^{\frac{1}{q-1}}.$$

Using the notation $p := q/(q-1)$ we get from (5.1)

$$a_m \leq a_{m-1} \left(1 - \frac{\lambda t_m}{2A} \right) = a_{m-1} \left(1 - \frac{t_m^p}{A_q A^p} a_{m-1}^{\frac{1}{q-1}} \right).$$

Raising both sides of this inequality to the power $1/(q-1)$ and taking into account the inequality $x^r \leq x$ for $r \geq 1$, $0 \leq x \leq 1$, we obtain

$$a_m^{\frac{1}{q-1}} \leq a_{m-1}^{\frac{1}{q-1}} \left(1 - \frac{t_m^p}{A_q A^p} a_{m-1}^{\frac{1}{q-1}} \right).$$

Then Lemma 5.4 with $y_k := a_k^{\frac{1}{q-1}}$, $n = 0$, $w_k = t_m^p/(A_q A^p)$, which provides

$$a_m^{\frac{1}{q-1}} \leq C(q, \gamma) A^p \left(C(E, q, \gamma) + \sum_{k=1}^m t_k^p \right)^{-1},$$

that implies

$$a_m \leq C(q, \gamma) A^q \left(C(E, q, \gamma) + \sum_{k=1}^m t_k^p \right)^{1-q},$$

which proves the theorem. \square

6. Proofs for Section 3

In this section we state the proofs for the results from Section 3. We begin with the proof that the algorithms stated in Section 3.1 belong to the class $\mathcal{WBG}\mathcal{A}(\Delta, \text{co})$.

Proof of Proposition 3.1. It is easy to see that conditions **(1)** and **(2)** from the definition of the class $\mathcal{WBG}\mathcal{A}(\Delta, \text{co})$ are satisfied for all three algorithms. Condition **(4)** holds with $C_0 = 1$ since for all three algorithms we have for any $m \geq 1$

$$E(G_m) \leq E(0) + \delta_m \leq E(0) + 1.$$

To guarantee condition **(3)**, first note that for any $m \geq 1$ and any $u > 0$ the definition of modulus of smoothness (2.2) provides

$$E((1+u)G_m) + E((1-u)G_m) \leq 2E(G_m) + 2\rho(E, D_1, u\|G_m\|).$$

Assume that $\langle E'(G_m), G_m \rangle \geq 0$ (the case $\langle E'(G_m), G_m \rangle < 0$ is handled similarly). Then from convexity (2.3) we get

$$E((1+u)G_m) \geq E(G_m) + u \langle E'(G_m), G_m \rangle$$

and from the definitions of the corresponding algorithms we obtain

$$E((1-u)G_m) \geq E(G_m) - \delta_m.$$

Combining the above estimates we deduce

$$\langle E'(G_m), G_m \rangle \leq \frac{\delta_m + 2\rho(E, D_1, u\|G_m\|)}{u}.$$

Taking infimum over $u > 0$ completes the proof. \square

Next, we state necessary technical lemmas that will be utilized in the proof of main results.

Lemma 6.1 ([29, Lemma 3.2]). *Let $\rho(u)$ be a non-negative convex on $[0, 1]$ function with the property $\rho(u)/u \rightarrow 0$ as $u \rightarrow 0$. Assume that a nonnegative sequence $\{\alpha_k\}_{k=1}^\infty$ is such that $\alpha_k \rightarrow 0$ as $k \rightarrow \infty$. Suppose that a nonnegative sequence $\{a_k\}_{k=0}^\infty$ satisfies the inequalities*

$$a_m \leq a_{m-1} + \inf_{0 \leq \lambda \leq 1} (-\lambda v a_{m-1} + B\rho(\lambda)) + \alpha_m, \quad m = 1, 2, 3, \dots$$

with positive numbers v and B . Then

$$\lim_{m \rightarrow \infty} a_m = 0.$$

Lemma 6.2 ([29, Lemma 3.3]). *Suppose a nonnegative sequence a_0, a_1, \dots satisfies the inequalities for $m = 1, 2, 3, \dots$*

$$a_m \leq a_{m-1} + \inf_{0 \leq \lambda \leq 1} (-\lambda v a_{m-1} + B\lambda^q) + \alpha_m, \quad \alpha_m \leq c m^{-q},$$

where $q \in (1, 2]$, $v \in (0, 1]$, and $B > 0$. Then

$$a_m \leq C(q, v, B, a_0, c) m^{1-q} \leq C'(q, B, a_0, c) v^{-q} m^{1-q}.$$

Lastly, we establish a generalized version of Lemma 5.5.

Lemma 6.3 (General Error Reduction Lemma). *Let E be a uniformly smooth on $S \subset X$ convex function with the modulus of smoothness $\rho(E, S, u)$. Take a number $\epsilon \geq 0$ and an element $f^\epsilon \in S$ such that*

$$E(f^\epsilon) \leq \inf_{x \in X} E(x) + \epsilon, \quad f^\epsilon/B \in \mathcal{A}_1(\mathcal{D}),$$

with some number $B \geq 1$. Suppose that $G \in S$ and $\varphi \in \mathcal{D}$ satisfy the following conditions

$$\langle -E'(G), \varphi \rangle \geq \theta \sup_{g \in \mathcal{D}} \langle -E'(G), g \rangle, \quad \theta \in (0, 1]; \quad (6.1)$$

$$|\langle E'(G), G \rangle| \leq \delta, \quad \delta \in [0, 1]. \quad (6.2)$$

Then we have

$$\begin{aligned} \inf_{0 \leq \lambda \leq 1} E(G + \lambda\varphi) &\leq E(G) \\ &+ \inf_{0 \leq \lambda \leq 1} (-\lambda\theta B^{-1}(E(G) - E(f^\epsilon)) + 2\rho(E, S, \lambda)) + \delta. \end{aligned}$$

Proof. If $E(G_{m-1}) - E(f^\epsilon) \leq 0$ then the claim of Lemma 6.3 is trivial. Assuming $E(G_{m-1}) - E(f^\epsilon) > 0$, Lemma 5.2 provides for any $\lambda \geq 0$

$$E(G + \lambda\varphi) \leq E(G) - \lambda \langle -E'(G), \varphi \rangle + 2\rho(E, S, \lambda).$$

By (6.1) and Lemma 5.3 we get

$$\begin{aligned} \langle -E'(G), \varphi \rangle &\geq \theta \sup_{g \in \mathcal{D}} \langle -E'(G), g \rangle \\ &= \theta \sup_{\phi \in \mathcal{A}_1(\mathcal{D})} \langle -E'(G), \phi \rangle \geq \theta B^{-1} \langle -E'(G), f^\epsilon \rangle. \end{aligned}$$

By (6.2) and by convexity (2.4) we obtain

$$\langle -E'(G), f^\epsilon \rangle = \langle -E'(G), f^\epsilon - G \rangle + \langle -E'(G), G \rangle \geq E(G) - E(f^\epsilon) - \delta.$$

Thus

$$E(G + \lambda\varphi) \leq E(G) - \lambda\theta B^{-1}(E(G) - E(f^\epsilon)) + 2\rho(E, S, \lambda) + \delta,$$

which proves the lemma. \square

Proof of Theorem 3.1. Assumption **(4)** from the definition of the class $\mathcal{WBG}\mathcal{A}(\Delta, \text{co})$ implies that for any $m \geq 0$

$$E(G_m) \leq E(0) + C_0 \quad \text{and} \quad G_m \in D_1.$$

Then from Lemma 6.3 with $S = D_1$, $G = G_{m-1}$, $\varphi = \varphi_m$, $\delta = \epsilon_m$, $\theta = t$, $B = A(\epsilon)$ and property **(2)** from the definition of the class $\mathcal{WBG}\mathcal{A}(\Delta, \text{co})$ we obtain

$$\begin{aligned} E(G_m) &\leq \inf_{0 \leq \lambda \leq 1} E(G_{m-1} + \lambda\varphi_m) + \delta_m \\ &\leq E(G_{m-1}) + \inf_{0 \leq \lambda \leq 1} \left(-\lambda t A^{-1}(E(G_{m-1}) - E(f^\epsilon)) \right. \\ &\quad \left. + 2\rho(E, D_1, \lambda) \right) + \delta_m + \epsilon_m. \end{aligned} \quad (6.3)$$

Denote

$$a_n := \max \{E(G_n) - E(f^\epsilon), 0\}.$$

Note that under our assumptions $t \in (0, 1]$ and $A := A(\epsilon) \geq 1$ we always have

$$a_{m-1} + \inf_{0 \leq \lambda \leq 1} (-\lambda t A^{-1} a_{m-1} + 2\rho(E, D_1, \lambda)) \geq 0.$$

Therefore estimate (6.3) implies

$$a_m \leq a_{m-1} + \inf_{0 \leq \lambda \leq 1} (-\lambda t A^{-1} a_{m-1} + 2\rho(E, D_1, \lambda)) + \delta_m + \epsilon_m. \quad (6.4)$$

We apply Lemma 6.1 with $v = tA^{-1}$, $B = 2$, $\rho(u) = \rho(E, D_1, \lambda)$, and $\alpha_m = \delta_m + \epsilon_m$ to obtain

$$\lim_{m \rightarrow \infty} a_m = 0,$$

which implies

$$\limsup_{m \rightarrow \infty} E(G_m) \leq \epsilon + \inf_{x \in D_1} E(x)$$

and, due to the arbitrary nature of choice of $\epsilon > 0$,

$$\lim_{m \rightarrow \infty} E(G_m) = \inf_{x \in D_1} E(x),$$

which completes the proof. \square

Proof of Theorem 3.2. From estimate (6.4) we get

$$\begin{aligned} a_m &\leq a_{m-1} + \inf_{0 \leq \lambda \leq 1} (-\lambda t A^{-1} a_{m-1} + 2\rho(E, D_1, \lambda)) + \delta_m + \epsilon_m \\ &\leq a_{m-1} + \inf_{0 \leq \lambda \leq 1} (-\lambda t A^{-1} a_{m-1} + 2\gamma \lambda^q) + \delta_m + \epsilon_m. \end{aligned}$$

Applying Lemma 6.2 with $v = tA^{-1}$, $B = 2\gamma$, and $\alpha_m = \delta_m + \epsilon_m$ completes the proof. \square

Acknowledgments

The first author acknowledges support given by the Oak Ridge National Laboratory, which is operated by UT-Battelle, LLC., for the U.S. Department of Energy under Contract DE-AC05-00OR22725.

The work was supported by the Russian Federation Government Grant N^o14.W03.31.0031. The paper contains results obtained in frames of the program "Center for the storage and analysis of big data", supported by the Ministry of Science and High Education of Russian Federation (contract 11.12.2018 N^o13/1251/2018 between the Lomonosov Moscow State University and the Fond of support of the National technological initiative projects).

References

References

- [1] B. Beauzamy, Introduction to Banach spaces and their geometry, North-Holland (1982).
- [2] T. Blumensath and M.E. Davies, Gradient Pursuits, IEEE Transactions in Signal Processing, **56**, 2370–2382 (2008).
- [3] T. Blumensath and M.E. Davies, Stagewise Weak Gradient Pursuits, IEEE Transactions in Signal Processing, **57**, 4333–4346 (2009).
- [4] J. Borwein, A. Guirao, P. Hájek, and J. Vanderwerff, Uniformly convex functions on Banach spaces, Proceedings of the American Mathematical Society, **137**(3), 1081–1091 (2009).

- [5] S. Boyd and L. Vandenberghe, *Convex optimization*, Cambridge university press (2004).
- [6] V. Chandrasekaran, B. Recht, P. Parrilo, and A. Willsky, The convex geometry of linear inverse problems, *Foundations of Computational mathematics*, **12**(6), 805–849 (2012).
- [7] K.L. Clarkson, Coresets, Sparse Greedy Approximation, and the Frank-Wolfe Algorithm, *ACM Transactions on Algorithms*, **6**(4), 63 (2010).
- [8] A.V. Dereventsov, On the Approximate Weak Chebyshev Greedy Algorithm in uniformly smooth Banach spaces, *Journal of Mathematical Analysis and Applications*, **436**(1), 288–304 (2016).
- [9] A. Dereventsov, Convergence and rate of convergence of approximate greedy-type algorithms, *Doctoral dissertation* (2017).
- [10] A.V. Dereventsov and V.N. Temlyakov, A unified way of analyzing some greedy algorithms, *Journal of Functional Analysis*, **277**(12) (2019).
- [11] R.A. DeVore, Nonlinear approximation, *Acta Numerica*, **7**, 51–150 (1998).
- [12] R.A. DeVore and V.N. Temlyakov, Convex optimization on Banach spaces, *Foundations of Computational Mathematics*, **16**, 369–394 (2016).
- [13] M.J. Donahue, C. Darken, L. Gurvits, and E. Sontag, Rates of convex approximation in non-Hilbert spaces, *Constructive Approximation*, **13**(2), 187–220 (1997).
- [14] M.A. Figueiredo, R.D. Nowak, and S.J. Wright, Gradient projection for Sparse Reconstruction: Application to Compressed Sensing and Other Inverse Problems, *IEEE Selected Topics in Signal Processing*, **1**(4), 586–597 (2007).
- [15] V.V. Galatenko and E.D. Livshitz, On the convergence of approximate weak greedy algorithms, *East journal on approximations*, **9**(1), 43–50 (2003).
- [16] Z. Gao and G. Petrova, Rescaled Pure Greedy Algorithm for convex optimization, *Calcolo*, **56** (2019).
- [17] R. Gribonval and M. Nielsen, Approximate weak greedy algorithms, *Advances in Computational Mathematics*, **14**, 361–378 (2001).
- [18] M. Jaggi, Revisiting Frank-Wolfe: Projection-free sparse convex optimization, *International Conference on Machine Learning*, **1**, 427–435 (2013).
- [19] M. Jaggi and M. Sulovský, A Simple Algorithm for Nuclear Norm Regularized Problems, *International Conference on Machine Learning* (2010).

- [20] Yu. Nesterov, *Introductory Lectures on Convex Optimization: A Basic Course*, Kluwer Academic Publishers, Boston (2004).
- [21] H. Nguyen and G. Petrova, Greedy strategies for convex optimization, *Calcolo*, 1–18 (2016).
- [22] G. Petrova, Rescaled Pure Greedy Algorithm for Hilbert and Banach spaces, *Applied and Computational Harmonic Analysis*, **41**, 852–866 (2016).
- [23] S. Shalev-Shwartz, N. Srebro, and T. Zhang, Trading accuracy for sparsity in optimization problems with sparsity constraints, *SIAM Journal on Optimization*, **20**(6), 2807–2832 (2010).
- [24] V.N. Temlyakov, Weak Greedy Algorithms, *Advances in Computational Mathematics*, **12**(2–3), 213–227 (2000).
- [25] V.N. Temlyakov, Greedy type algorithms in Banach spaces and applications, *Constructive Approximation*, **21**, 257–292 (2005).
- [26] V.N. Temlyakov, *Greedy approximation*, Cambridge University Press (2011).
- [27] V.N. Temlyakov, Greedy approximation in convex optimization, *Constructive Approximation*, **41**, 269–296 (2015).
- [28] V.N. Temlyakov, Greedy expansions in convex optimization, *Proceedings of the Steklov Institute of Mathematics*, **284**, 244–262 (2014).
- [29] V.N. Temlyakov, Convergence and Rate of Convergence of Some Greedy Algorithms in Convex Optimization, *Trudy MIAN*, **293**, 333–345 (2016); English transl. *Proceedings of Steklov Institute*, **293**, 325–337 (2016).
- [30] A. Tewari, P.K. Ravikumar, and I.S. Dhillon, Greedy algorithms for structurally constrained high dimensional problems, *Advances in Neural Information Processing Systems*, 882–890 (2011).
- [31] T. Zhang, Sequential greedy approximation for certain convex optimization problems, *IEEE Transactions on Information Theory*, **49**(3), 682–691 (2003).