

NJL model estimation of anisotropic electrical conductivity for quark matter in presence of magnetic field

Jayanta Dey* and Sabyasachi Ghosh
*Indian Institute of Technology Bhilai, GEC Campus,
 Sejbahar, Raipur 492015, Chhattisgarh, India*

Aritra Bandyopadhyay and Ricardo L. S. Farias
Departamento de Física, Universidade Federal de Santa Maria, Santa Maria, RS, 97105-900, Brazil

Gastão Krein
Instituto de Física Teórica, Universidade Estadual Paulista, 01140-070 São Paulo, SP, Brazil

Present work has gone through the microscopic calculation of electrical conductivity of quark matter in presence of magnetic field, where NJL model is considered for mapping the interaction picture of the medium. Let us start with Ohm's law'

$$J_D^i = \sigma^{ij} E_j \quad (1)$$

where J_D^i is dissipative current, σ^{ij} electrical conductivity tensor and E_j is electric field. Now for a fluid of quark having spin-color degeneracy g and electric charge q_f dissipative current from kinetic theory framework can be written as,

$$J_D^j = q_f g \int \frac{d^3 p}{(2\pi)^3} \vec{v} \delta f \quad (2)$$

Where δf is small deviation of quark distribution function from the equilibrium Fermi-Dirac distribution of quark $f_0 = \frac{1}{e^{\beta\omega} + 1}$. In terms of 3-momentum (\vec{p}) and energy (ω) particle velocity can be written as $\vec{v} = \frac{\vec{p}}{\omega}$. Now to find δf in presence of electric field \vec{E} and magnetic field \vec{B} we use RTA in Boltzmann's equation, where we can assume the a general force term

$$\vec{F} = \alpha \vec{e} + \beta \vec{b} + \gamma \vec{e} \times \vec{b} \quad (3)$$

where \vec{e} , \vec{b} are unit vector along \vec{E} and \vec{B} . With suitable connection between δf and \vec{F} [1, 2], the coefficients α , β and γ can be found as

$$\begin{aligned} \alpha &= q \left(\frac{\tau_c}{\omega} \right) \frac{1}{1 + (\tau_c/\tau_B)^2} E, \\ \beta &= q \left(\frac{\tau_c}{\omega} \right) \frac{(\tau_c/\tau_B)^2}{1 + (\tau_c/\tau_B)^2} (\vec{e} \cdot \vec{h}) E, \\ \gamma &= -q \left(\frac{\tau_c}{\omega} \right) \frac{(\tau_c/\tau_B)}{1 + (\tau_c/\tau_B)^2} E, \end{aligned} \quad (4)$$

where τ_B and τ_c as magnetic and thermal relaxation time. After taking care of all degeneracy factors of u and d quarks, we get 3 components of electrical conductivity, whose general expressions can be written as

$$\sigma_n = e^2 \beta \frac{20}{9} \int \frac{d^3 p}{(2\pi)^3} \frac{p^2}{\omega^2} \frac{\tau_c (\tau_c/\tau_B)^n}{1 + (\tau_c/\tau_B)^2} f_0 (1 - f_0). \quad (5)$$

with $n = 0, 1, 2$. We will get temperature (T) and magnetic field (B) dependent quark mass from NJL model, briefly discussed below, which will be used in above Eq. (5) to estimate σ_n of quark matter.

The Lagrangian density for the isospin-symmetric two-flavor version of NJL model in presence of electromagnetic field (A^μ) is given by

$$\begin{aligned} \mathcal{L}_{NJL} &= -\frac{1}{4} F^{\mu\nu} F_{\mu\nu} + \bar{\psi} (\not{D} - m) \psi \\ &\quad + G [(\bar{\psi}\psi)^2 + (\bar{\psi}i\gamma_5\vec{\tau}\psi)^2], \end{aligned} \quad (6)$$

*Electronic address: jayantad@iitbhilai.ac.in




FIG. 1: Temperature dependence of three components of electrical conductivities ($\sigma_{0,1,2}$) with $eB = 0.2 \text{ GeV}^2$ and σ (without B) for $\tau_c = 5 \text{ fm}$ (b) and 0.2 fm (d). $\Delta\sigma_n/\sigma = (\sigma_n - \sigma)/\sigma$ for $\tau_c = 5 \text{ fm}$ (a) and 0.2 fm (c).

In quasi-particle approximation, the gap equation for the constituent quark mass M at finite T and B is given by

$$M = m - 2G \sum_{f=u,d} \langle \bar{\psi}_f \psi_f \rangle, \quad (7)$$

where $\langle \bar{\psi}_f \psi_f \rangle$ represents the quark condensate of flavor f , and a thermo-magnetic NJL coupling constant $G(B, T)$ has been considered [3, 4].

Using the governing NJL model Eq. (7) and then plug in to Eq. (5), we have estimated σ_n . In Figs. 1(b) and (d), we present the temperature dependence of the different components of electrical conductivities σ_n (scaled with T) in presence of an external magnetic field as

well as without field case,

$$\sigma = e^2 \beta \frac{20}{9} \int \frac{d^3 p}{(2\pi)^3} \frac{\vec{p}^2}{\omega^2} \tau_c f_0 (1 - f_0). \quad (8)$$

Their difference $\frac{\Delta\sigma_n}{\sigma} = \frac{\sigma_n - \sigma}{\sigma}$ are plotted in Figs. 1(a) and (c). The results are presented for $\tau_c = 5 \text{ fm}$ (a, b) and $\tau_c = 0.2 \text{ fm}$ (c, d) but in same magnetic field ($eB = 0.2 \text{ GeV}^2$), which can be assigned with the zones $\tau_c > \tau_B$ and $\tau_c < \tau_B$ respectively. It means that $eB = 0.2 \text{ GeV}^2$ may be considered as stronger magnetic field for $\tau_c = 5 \text{ fm}$ and weaker magnetic field for $\tau_c = 0.2 \text{ fm}$. Therefore, former case is showing $\sigma_2 > \sigma_0$ and latter case is showing $\sigma_2 < \sigma_0$. It is controlling by the anisotropic function $\frac{(\tau_c/\tau_B)^n}{1+(\tau_c/\tau_B)^2}$. In terms of anisotropy, the above outcomes can be briefly sketched as

$$\text{for } \tau_c = 5 \text{ fm} \\ \sigma^{xx} = \sigma^{yy} < \sigma^{zz} \Rightarrow \text{larger anisotropy} \quad (9)$$

$$\text{for } \tau_c = 0.2 \text{ fm}, \\ \sigma^{xx} = \sigma^{yy} \approx \sigma^{zz} \Rightarrow \text{smaller anisotropy} \quad (10)$$

when external magnetic field $eB = 0.2 \text{ GeV}^2$ is in the z-direction. As seen from Fig. 1, all the components of the electrical conductivities increase with temperature with a kink near the quark-hadron phase transition temperature T_c and rate of increments are also different for two different phases.

References

- [1] A. Harutyunyan, A. Sedrakian, Phys. Rev. **C 94** (2016) 025805.
- [2] J. Dey, S. Satapathy, A. Mishra, S. Paul, S. Ghosh, arXiv:1908.04335 [hep-ph].
- [3] R.L.S. Farias, K.P. Gomes, G. Krein, and M.B. Pinto, Phys. Rev. **C 90**, 025203 (2014).
- [4] R.L.S. Farias, V.S. Timteo, S.S. Avancini, M.B. Pinto, and G. Krein, Eur. Phys. J. **A 53** (2017) 101.