

TRANSVERSE HILBERT SCHEMES, BI-HAMILTONIAN SYSTEMS, AND HYPERKÄHLER GEOMETRY

ROGER BIELAWSKI

Dedicated to the memory of Sir Michael Francis Atiyah (1929-2019)

ABSTRACT. We give a characterisation of Atiyah's and Hitchin's transverse Hilbert schemes of points on a symplectic surface in terms of bi-Poisson structures. Furthermore, we describe the geometry of hyperkähler manifolds arising from the transverse Hilbert scheme construction, with particular attention paid to the monopole moduli spaces.

1. INTRODUCTION

In chapter 6 of the monograph [1], Atiyah and Hitchin consider the following construction. Let Y be a complex symplectic surface with a holomorphic submersion π onto a 1-dimensional complex manifold X . They associate to it an open subset of the Hilbert scheme of n points on Y consisting of 0-dimensional complex subspaces D of length n such that $\pi|_D$ is an isomorphism onto its scheme-theoretic image. They observe that this *transverse Hilbert scheme* $Y_\pi^{[n]}$ is a symplectic manifold equipped with holomorphic submersion $\pi^{[n]}$ onto $S^n X$, the fibres of which are Lagrangian submanifolds. In particular, if X is a domain in \mathbb{C} , then the components of $\pi^{[n]}$ define n functionally independent and Poisson-commuting Hamiltonians on $Y_\pi^{[n]}$, i.e. a completely integrable system. Atiyah and Hitchin observe further that sometimes one can perform this construction on the fibres of the twistor space of a 4-dimensional hyperkähler manifold and obtain a new twistor space which then might lead to a new hyperkähler manifold. Their main example of this construction is $Y = \mathbb{C}^* \times \mathbb{C}$ with π the projection onto the second factor. The corresponding transverse Hilbert scheme is the space of based rational maps of degree n and the hyperkähler metric resulting from applying the construction to the twistor space of $S^1 \times \mathbb{R}^3$ is the L^2 -metric on the moduli space of Euclidean monopoles of charge n . Further examples of this construction are given in [20, 5].

The purpose of this article is to characterise both symplectic and hyperkähler manifolds arising from this construction. Partial results in this direction have been obtained in [6] and in [17, 18]. They rely on the existence of a certain endomorphism of the tangent bundle of $Y_\pi^{[n]}$. In the present work our point of view is different. We observe that $Y_\pi^{[n]}$ is equipped with a second Poisson structure, compatible with the symplectic form. Thus $Y_\pi^{[n]}$ is a completely integrable *bi-Hamiltonian system*. We then show that a nondegenerate bi-Poisson manifold M^{2n} arises as an (open subset) of a transverse Hilbert scheme on a symplectic surface Y with a submersion $Y \rightarrow \mathbb{C}$ essentially exactly then, when the coefficients of the minimal polynomial of

The author is a member of the DFG Priority Programme 2026 "Geometry at infinity".

the corresponding recursion operation (see §2.3 for a definition) form a submersion to \mathbb{C}^n .

We then turn our attention to hyperkähler manifolds arising from the transverse Hilbert scheme construction on the fibres of the twistor space a 4-dimensional hyperkähler manifold with a tri-Hamiltonian vector field. We show that the essential feature of the geometry of a manifold M arising from this construction is the existence of a bivector Π on M which lies in Salamon's component $\Lambda^2 E \otimes S^2 H$ of $\Lambda^2 T^{\mathbb{C}} M$ and satisfies $D\Pi = 0$, where D is the Penrose-Ward-Salamon differential operator on $\Lambda^2 E \otimes S^2 H$ [21]. The bivector Π is not Poisson, but its $(2, 0)$ -component with respect to each complex structure is a (generically log-symplectic) holomorphic Poisson bivector. Moreover, this holomorphic Poisson bivector is compatible with the parallel holomorphic symplectic form arising from the hyperkähler structure.

In the last section we identify the bivector Π on moduli spaces of $SU(2)$ -monopoles, i.e. hyperkähler transverse Hilbert schemes on $S^1 \times \mathbb{R}^3$, in terms of solutions to Nahm's equations.

2. TRANSVERSE HILBERT SCHEMES AND BI-HAMILTONIAN SYSTEMS

2.1. Transverse Hilbert schemes. Let X be a complex manifold, C a complex manifold of dimension 1, and $\pi : X \rightarrow C$ a holomorphic map. The *transverse Hilbert scheme* $X_\pi^{[n]}$ of n points in X is an open subset of the full Hilbert scheme $X^{[n]}$ consisting of those $D \in X^{[n]}$ such that $\pi|_D$ is an isomorphism onto its scheme-theoretic image [1]. Since $C^{[n]} = S^n C$, this simply means that $\pi(D)$ consists of n points (with multiplicities). First of all, observe that $X_\pi^{[n]}$ is always smooth, unlike the full Hilbert scheme $X^{[n]}$:

Proposition 2.1. *Let $\pi : X \rightarrow C$ be a holomorphic map from a complex manifold X to a 1-dimensional complex manifold C . Then the transverse Hilbert scheme $X_\pi^{[n]}$ is smooth.*

Proof. Since $D \in X_\pi^{[n]}$ satisfies $D \simeq \pi(D) \in S^n C$, such a D is a local complete intersection (l.c.i.). Now the claim follows from general results of deformations theory (see, e.g., [13, Theorem 1.1.(c)]). \square

The transverse Hilbert scheme comes equipped with a canonical map $\pi^{[n]} : X_\pi^{[n]} \rightarrow S^n C$. If π is a submersion, then so is $\pi^{[n]}$. In this case, points of $X_\pi^{[n]}$ such that $\pi(D) = n_1 p_1 + \dots + n_k p_k$, with p_1, \dots, p_k distinct points of C , correspond to a choice of a section s_i of π in a neighbourhood of each p_i , truncated to order n_i (in other words s_i is an $(n_i - 1)$ -jet of sections at p_i). Let us remark that Atiyah and Hitchin consider only the case when π is a submersion (they say π is a “complex fibration”, and the proof of the smoothness of $X_\pi^{[n]}$, given on p. 53 in [1], makes clear that π must be a submersion).

Suppose now that X has a symplectic structure. If $\dim X = 2$, then a theorem of Beauville [2] implies that $X^{[n]}$ (which is smooth, owing to a well-known result of Fogarty), and hence $X_\pi^{[n]}$, carries an induced symplectic structure. For higher dimensional X , there is no induced symplectic structure on $X^{[n]}$, not even on its smooth locus.

2.2. Log-symplectic Poisson structures. A Poisson structure on a (smooth or complex) manifold M^{2n} is given by a bivector $\Pi \in \Gamma(\Lambda^2 T M)$ such that the Schouten

bracket $[\Pi, \Pi]$ vanishes. The *symplectic locus* of the Poisson structure is the set of points m where the induced map $\#_\Pi : T_m^*M \rightarrow T_mM$ is an isomorphism. Its complement is called the *degeneracy locus*. A Poisson structure is called *log-symplectic* if $\Pi^n \in \Gamma(\Lambda^{2n}TM)$ meets the zero section of the $\Lambda^{2n}TM$ transversely. These structures were studied by Goto [10] in the holomorphic case, and by Guillemin, Miranda and Pires in the smooth category [12] (see also [11, 8]). The name is justified by the fact that the dual 2-form $\omega = \Pi^{-1}$ has a logarithmic singularity along the degeneracy locus. The degeneracy locus Δ of a log-symplectic Poisson structure is a smooth Poisson hypersurface with codimension one symplectic leaves and $M \setminus \Delta$ is a union of open symplectic leaves.

We recall from [12] that if f is a local defining function for Δ , then ω can be decomposed as

$$(2.1) \quad \omega = \alpha \wedge \frac{df}{f} + \beta,$$

for a 1-form α and a 2-form β . Moreover, the restrictions of α and β to Δ are closed, $\alpha|_\Delta$ is intrinsically defined and its kernel is the tangent space to the symplectic leaf of Π .

2.3. Bi-Poisson structures . A *bi-Poisson* structure on a (real or complex) manifold M is a pair (Π_1, Π_2) of linearly independent bivectors such that every linear combination of Π_1 and Π_2 is a Poisson structure. In other words Π_1 and Π_2 satisfy $[\Pi_1, \Pi_1] = 0$, $[\Pi_2, \Pi_2] = 0$, $[\Pi_1, \Pi_2] = 0$, where $[\cdot, \cdot]$ is the Schouten bracket.

A bi-Poisson structure is called *nondegenerate*, if the pencil $t_1\Pi_1 + t_2\Pi_2$ contains a symplectic structure. In what follows, we shall consider only nondegenerate bi-Poisson structures and assume that Π_1 is symplectic. Following Magri and Morosi [19] (see also [9]) we can define the *recursion operator* $R = \#_{\Pi_2} \circ \#_{\Pi_1}^{-1}$. It is an endomorphism of TM and Magri and Morosi show that 1) its Nijenhuis tensor vanishes; and 2) the eigenvalues of R form a commuting family with respect to both Poisson brackets.

Furthermore, $\det R = (\mu_R)^2$ for a well defined function μ_R on M (μ_R is the quotient of the Pfaffians of Π_2 and of Π_1). Thus Π_2 is log-symplectic if and only if 0 is a regular value of μ_R . Since $(\Pi_1, \Pi_2 - \lambda\Pi_1)$ is a nondegenerate bi-Poisson structure for each scalar λ , the characteristic polynomial of R is of the form $\chi_R(\lambda) = \mu_R(\lambda)^2$. We shall refer to $\mu_R(\lambda)$ as the *Pfaffian polynomial* of R . We observe:

Proposition 2.2. *Let (Π_1, Π_2) be a real (resp. holomorphic) bi-Poisson structure on a smooth (resp. complex) manifold M^{2n} with Π_1 symplectic. If the coefficients of the Pfaffian polynomial of the recursion operator R define a submersion $p : M \rightarrow \mathbb{R}^n$ (resp. $p : M \rightarrow \mathbb{C}^n$), then the Poisson structure $\Pi_2 - \lambda\Pi_1$ is log-symplectic for every λ .* \square

We also recall the following property of bi-Poisson structures, proved by Magri and Morosi in [19]:

Proposition 2.3 (Magri-Morosi). *Let (M, Π_1, Π_2) be a bi-Poisson manifold with Π_1 symplectic. Then, for any polynomial $\rho(z)$, the bivector Π_ρ defined by*

$$\Pi_\rho(\alpha, \cdot) = \rho(R)\Pi_1(\alpha, \cdot), \quad \alpha \in \Omega^1(M)$$

defines a Poisson structure on M , compatible with Π_1 . \square

2.4. Transverse Hilbert schemes on symplectic surfaces. Beauville's construction [2] of a symplectic form on the Hilbert scheme $S^{[n]}$ of n points in a symplectic surface S has been extended by Bottacin [7] to Poisson surfaces: any Poisson structure on a complex surface S induces a Poisson structure on $S^{[n]}$. Therefore, if (S, ω) is a symplectic surface and $\pi : S \rightarrow \mathbb{C}$ a holomorphic map, we obtain two Poisson bivectors on $S^{[n]}$: Π_1 induced by ω^{-1} and Π_2 induced by $\pi \cdot \omega^{-1}$, where $\omega^{-1} = \#_\omega \omega$ (i.e. the bivector dual to ω) and π is viewed as a function on S . Since the Poisson structures on S are compatible, Π_1 and Π_2 are compatible (compatibility is trivial on the open dense subset where D consists of distinct points, and hence $[\Pi_1, \Pi_2]$ vanishes everywhere). Observe that the corresponding recursion operator R (cf. §2.3) is the endomorphism of $TS^{[n]}$ given by the multiplication by π on each $T_D S^{[n]} \simeq H^0(D, \mathcal{N}_{D/S})$ where $\mathcal{N}_{D/S}$ denotes the normal sheaf of D in S . This is the endomorphism considered in [6, 17, 18]. The coefficients of its Pfaffian polynomial define a map $S^{[n]} \rightarrow S^n \mathbb{C} \simeq \mathbb{C}^n$. Its restriction to the transverse Hilbert scheme $S_\pi^{[n]}$ coincides with the canonical map $\pi^{[n]}$ introduced in §2.1. Let us prove the following properties of $S_\pi^{[n]}$ and $\pi^{[n]}$.

Proposition 2.4. *Let S be a complex symplectic surface with a holomorphic map $\pi : S \rightarrow \mathbb{C}$. Then the transverse Hilbert scheme $S_\pi^{[n]}$ is a nondegenerate bi-Poisson manifold with the following properties:*

- (i) *the coefficients of the minimal polynomial of the corresponding recursion operator R coincide with the canonical map $\pi^{[n]} : S_\pi^{[n]} \rightarrow \mathbb{C}^n$;*
- (ii) *at any point of its degeneracy locus, the Poisson structure $\Pi_2 - \lambda \Pi_1$ has rank $2n - 2$ ($\lambda \in \mathbb{C}$);*
- (iii) *on the subset of $\pi^{[n]}$ -regular points, the Poisson structure $\Pi_2 - \lambda \Pi_1$ is log-symplectic for every $\lambda \in \mathbb{C}$.*

Proof. We already know that $S_\pi^{[n]}$ is a nondegenerate bi-Poisson manifold. Owing to the definition of the recursion operator, we know that the geometric multiplicity of each eigenvalue is even. Now observe that the multiplication by π defines also an endomorphism \bar{R} of $T_{\pi(D)} \mathbb{C}^{[n]}$. The geometric multiplicity of every eigenvalue of \bar{R} is equal to 1 (since $\pi(D) \in S^n \mathbb{C}$ has length n). We also know that the characteristic polynomial of \bar{R} is equal to the Pfaffian polynomial $\mu_R(\lambda)$ of R , and that the characteristic polynomial of R is $\mu_R(\lambda)^2$. Putting this together, we conclude that the geometric multiplicity of every eigenvalue of R is equal to 2 and that the minimal polynomial of R is equal to $\mu_R(\lambda)$. This proves statements (i) and (ii). The third statement follows from Proposition 2.2. \square

Remark 2.5. Statement (i) has been shown in [17, Remark 2.4] under the assumption that π is a submersion.

Remark 2.6. Since, owing to the above mentioned result of Bottacin, any Poisson structure on S induces a Poisson structure on $S^{[n]}$, we can conclude that if S is a Poisson surface with a holomorphic map $\pi : S \rightarrow \mathbb{C}$, then $S^{[n]}$ is a bi-Poisson manifold. The bi-Poisson structure will, however, be degenerate if S is not symplectic.

Remark 2.7. Suppose that (z, u) are Darboux coordinates for the symplectic form ω on an open subset U of S , i.e. $\omega = dz \wedge du$ on U . Suppose also that $\pi(z, u) = z$ (which implies that π is a submersion on U). Then the corresponding open subset $U_\pi^{[n]}$ can be described as an open subset of $\{(q(z), p(z))\}$, where $q(z)$ is a monic

polynomial of degree n and $p(z)$ is a polynomial of degree at most $n-1$, such that, for every root z_i of q , $(z_i, p(z_i)) \in U$. On the open dense subset of $U_\pi^{[n]}$, where the roots are distinct, the two Poisson structures are given by:

$$\Pi_1 = \sum_{i=1}^n \frac{\partial}{\partial z_i} \wedge \frac{\partial}{\partial u_i}, \quad \Pi_2 = \sum_{i=1}^n z_i \frac{\partial}{\partial z_i} \wedge \frac{\partial}{\partial u_i}, \quad \text{where } u_i = p(z_i).$$

We can now characterise transverse Hilbert schemes on symplectic surfaces, in the case when π is a submersion (i.e. the case originally considered by Atiyah and Hitchin):

Theorem 2.8. *Let (M^{2n}, Π_1, Π_2) be a holomorphic bi-Poisson manifold with Π_1 symplectic. Assume that the coefficients of the Pfaffian polynomial of the corresponding recursion operator R define a submersion $p : M \rightarrow \mathbb{C}^n$ and that, for each $\lambda \in \mathbb{C}$, if the degeneracy locus D_λ of $\Pi_2 - \lambda\Pi_1$ is nonempty, then its symplectic foliation is simple.*

Then there exists a symplectic surface S with a holomorphic submersion $\pi : S \rightarrow \mathbb{C}$ and a local bi-Poisson biholomorphism $\Phi : (M, \Pi_1, \Pi_2) \rightarrow S_\pi^{[n]}$.

Proof. Let $\mu_m(\lambda)$ denote the Pfaffian polynomial of R_m . We consider the following incidence variety (cf. [1, pp. 40-43], [18]):

$$T = \{(\lambda, m) \in \mathbb{C} \times M; m \in D_\lambda\} = \{(\lambda, m) \in \mathbb{C} \times M; \mu_m(\lambda) = 0\}.$$

Due to the assumptions and to Proposition 2.2, T is smooth and the symplectic foliation on each D_λ is simple with codimension one leaves. We thus obtain an integrable simple foliation \mathcal{F} of T , the leaf space of which is a 2-dimensional complex manifold S with a canonical holomorphic submersion $\pi : S \rightarrow \mathbb{C}$.

On each D_λ there is a canonically defined (closed) 1-form α_λ (cf. (2.1) and the following lines), the kernel of which corresponds to the symplectic foliation of D_λ . Thus $\alpha_\lambda \wedge d\lambda$ defines a nondegenerate, hence symplectic, 2-form on S .

The scheme-theoretic inverse image of a point $m \in M$ defines a 0-dimensional subspace Z_m of T with structure sheaf isomorphic to $\mathbb{C}[\lambda]/(\mu_m(\lambda))$. The projection $T \rightarrow \mathbb{C}$ maps Z_m isomorphically onto a 0-dimensional subspace of \mathbb{C} of length n . Thus Z_m descends to an element of $S_\pi^{[n]}$, and we obtain a holomorphic map $\Phi : M \rightarrow S_\pi^{[n]}$. It remains to show that Φ is a local diffeomorphism. Since the coefficients of μ_m define a submersion, the corresponding Hamiltonian vector fields do not vanish anywhere. Near any point $p \in M$ we have therefore the “action-angle” coordinates on a neighbourhood U (given by coefficients of μ and the local free action of \mathbb{C}^n). Let S_p be the symplectic surface obtained from U by the above procedure. On $(S_p)_\pi^{[n]}$ there are analogous “action-angle” coordinate and therefore we obtain a holomorphic map $\psi : (S_p)_\pi^{[n]} \rightarrow U$. Fernandes [9] shows that, on the subset where the eigenvalues are distinct, there exist local coordinates z_i, u_i such that

$$\Pi_1 = \sum_{i=1}^n \frac{\partial}{\partial z_i} \wedge \frac{\partial}{\partial u_i}, \quad \Pi_2 = \sum_{i=1}^n z_i \frac{\partial}{\partial z_i} \wedge \frac{\partial}{\partial u_i}.$$

Thus, owing to Remark 2.7, ψ is the inverse of Φ on the open dense subset where the roots of μ_m are distinct (and a bi-Poisson isomorphism) and, hence, ψ is the inverse of $\Phi|_U$. \square

Remark 2.9. Presumably the result remains true without the assumption that the symplectic foliations of D_λ are simple, provided we replace “symplectic surface” with “2-dimensional symplectic stack”.

3. HYPERKÄHLER GEOMETRY OF TRANSVERSE HILBERT SCHEMES

3.1. Ward transform. Let us briefly recall the essential features of the Ward transform [22, 21] in the case of hypercomplex manifolds. Let Z be complex manifold with a surjective holomorphic submersion $\pi : Z \rightarrow \mathbb{P}^1$, and let $M^\mathbb{C}$ be the Kodaira moduli space of sections with normal bundle isomorphic to $\mathcal{O}(1)^{\oplus n}$. The twistor double fibration in this case is simply

$$(3.1) \quad M^\mathbb{C} \xleftarrow{\tau} M^\mathbb{C} \times \mathbb{P}^1 \xrightarrow{\nu} Z.$$

If F is an $M^\mathbb{C}$ -uniform (i.e. $h^0(\nu(\tau^{-1}(m)))$ is constant on $M^\mathbb{C}$) holomorphic vector bundle on Z , then we obtain an induced holomorphic vector bundle $\hat{F} = \tau_* \nu^* F$ on $M^\mathbb{C}$. In particular, if we denote by E the vector bundle induced from $T_\pi Z \otimes \pi^* \mathcal{O}(-1)$ (where $T_\pi Z = \ker d\pi$) and by H the trivial vector bundle with fibre \mathbb{C}^2 , we have $TM \simeq E \otimes H$. Furthermore, the vector bundle induced from $\pi^* \mathcal{O}(k)$, $k \geq 0$, is simply $S^k H$, and if F is $M^\mathbb{C}$ -trivial (i.e. trivial on each line $\nu(\tau^{-1}(m))$), then the bundle induced from $F \otimes \pi^* \mathcal{O}(k)$ is $\hat{F} \otimes S^k H$, for any $k \geq 0$.

Recall also that an induced vector bundle comes equipped with a first-order differential operator, which arises as the pushforward of a partial connection on $\nu^* F$, which is basically the exterior derivative in the fibre directions. If F is $M^\mathbb{C}$ -trivial, then this operator is a linear connection on \hat{F} . We can identify this operator for bundles of the form $\hat{F} \otimes S^k H$, which are induced from $F \otimes \pi^* \mathcal{O}(k)$, where F is $M^\mathbb{C}$ -trivial. We fix an isomorphism $H \simeq H^*$ (which corresponds to a choice of isomorphism $H^1(\mathbb{P}^1, \mathcal{O}(-2)) \simeq \mathbb{C}$). We denote by α the natural projection $S^k H \otimes H \rightarrow S^{k+1} H$ (which corresponds to multiplication of sections of $\mathcal{O}(k)$ and of $\mathcal{O}(1)$), and use the same letter for the corresponding map on $W \otimes S^k H \otimes H \rightarrow W \otimes S^{k+1} H$ for any holomorphic vector bundle W . The induced differential operators D are then:

1. on $S^k H$, $D = \alpha \circ d$;
2. on $\hat{F} \otimes S^k H$, $D = \alpha \circ (\nabla \otimes \text{Id}) + \text{Id} \otimes (\alpha \circ d)$, where F is $M^\mathbb{C}$ -trivial and ∇ denotes the induced connection on \hat{F} . The principal symbol of D is α .

The construction of D as the push-forward of a partial connection shows, in particular, that $Ds = 0$ if and only if $s = \tau_* \eta^* \tilde{s}$ for a holomorphic section \tilde{s} of $F(k)$ on Z .

3.2. Hyper-Poisson bivectors. Let M be a hypercomplex manifold with twistor space Z . As discussed above, the vector bundle $T_\pi Z \otimes \pi^* \mathcal{O}(-1)$ is M -trivial, and hence the operator D on the induced vector bundle E is a linear connection ∇ . Recall that the tensor product of ∇ and the flat connection on H is a torsion-free linear connection on TM known as the *Obata connection*. The induced operator D on TM is therefore the composition of the Obata connection and the projection $H \otimes H \rightarrow S^2 H$. Similarly, the vector bundle $\Lambda^r(T_\pi Z \otimes \pi^* \mathcal{O}(-1))$ is M -trivial, and hence the vector bundle on M induced from $\Lambda^r T_\pi Z$ is $\Lambda^r E \otimes S^r H$. This is a direct summand of $\Lambda^r T^\mathbb{C} M$, which Salamon [21, Prop. 4.2] identifies with the subspace of finite linear combinations of multivectors of type $(r, 0)$ for different

complex structures. Salamon also shows that its $Sp(1)$ -invariant complement is

$$(3.2) \quad B^r = \bigcap_{\zeta} (T_{\zeta}^{1,r-1}M \oplus T_{\zeta}^{2,r-2}M \oplus \cdots \oplus T_{\zeta}^{r-1,1}M),$$

where $\zeta \in \mathbb{P}^1$ labels different complex structures of the hypercomplex structure. As explained in the previous subsection, a multivector field $\Pi \in \Gamma(\Lambda^r E \otimes S^r H)$ arises from a holomorphic section of $\Lambda^r T_{\pi} Z$ precisely if it satisfies the equation $D\Pi = 0$. In that case, for any complex structure I_{ζ} , the $(r, 0)$ -component of Π is the corresponding holomorphic multivector field on the fibre $\pi^{-1}(\zeta)$ of Z .

Definition 3.1. Let M be a hypercomplex manifold. A bivector $\Pi \in \Gamma(\Lambda^2 TM)$ is called a *hyper-Poisson bivector* if

- (i) $\Pi \in \Gamma(\Lambda^2 E \otimes S^2 H)$;
- (ii) for each complex structure I_{ζ} , the corresponding $(2, 0)$ -component $\Pi_{\zeta}^{2,0}$ of Π is a holomorphic Poisson bivector on (M, I_{ζ}) .

Remark 3.2. As explained above, the condition that $\Pi_{\zeta}^{2,0}$ is holomorphic for each ζ is equivalent to $D\Pi = 0$. On the other hand, condition (ii) implies that $[\Pi, \Pi] \in \Gamma(B^3)$, where $[\cdot, \cdot]$ denotes the Schouten-Nijenhuis bracket and B^3 is defined in (3.2).

As usual, given a bivector field on M , we can define a bracket of (real- or complex-valued) functions on M by

$$(3.3) \quad \{f, g\} = \Pi(df, dg).$$

The name “hyper-Poisson” is justified by the following observation, which follows directly from the definition.

Proposition 3.3. *A bivector $\Pi \in \Gamma(\Lambda^2 E \otimes S^2 H)$ is hyper-Poisson if and only if, for each complex structure I_{ζ} , the bracket (3.3) is a Poisson bracket on the sheaf $\mathcal{O}(M, I_{\zeta})$ of I_{ζ} -holomorphic functions.* \square

Remark 3.4. Our notion of “hyper-Poisson” is different from [16]. There, it means a triple (π_1, π_2, π_3) of bivectors, such that $\pi_2 - i\pi_3$ is an I_1 -holomorphic Poisson bivector etc. We do not think there is a danger of confusion, since we talk about *hyper-Poisson bivectors*, while [16] deals with *hyper-Poisson triples*.

Definition 3.5. Let M be a hyperkähler manifold. A hyper-Poisson bivector Π on M is said to be compatible with the hyperkähler structure if, for every complex structure, the holomorphic Poisson bivector $\Pi_{\zeta}^{2,0}$ is compatible with Ω_{ζ}^{-1} , where Ω_{ζ} is the corresponding parallel holomorphic symplectic form.

Example 3.6. Recall that the twistor space Z of a hyperkähler manifold is equipped with a fibrewise $\mathcal{O}(2)$ -valued complex symplectic form ω , i.e. a section of $\Lambda^2 T_{\pi}^* Z \otimes \mathcal{O}(2)$. It can be viewed as a (holomorphic) section of $\Lambda^2(T_{\pi} Z(-1))^*$, i.e. it induces a symplectic form on the bundle E . The fibrewise bivector ω^{-1} is a section of $\Lambda^2(T_{\pi} Z(-1))$ and so multiplying it by a real section of $\pi^* \mathcal{O}(2)$ yields a holomorphic section of $\Lambda^2 T_{\pi} Z$ compatible with the real structure, i.e. a hyper-Poisson bivector. This bivector is simply a constant multiple of ω_{α}^{-1} , where ω_{α} is one of the Kähler forms of the hyperkähler metric (α is determined by the chosen section of $\mathcal{O}(2)$). In other words, for any Kähler form ω_{α} of the hyperkähler metric, the dual bivector ω_{α}^{-1} is a hyper-Poisson bivector compatible with the hyperkähler structure. The corresponding Poisson bracket on $\mathcal{O}(M, I_{\zeta})$ is identically 0 when $I_{\zeta} = \pm I_{\alpha}$.

We can classify hyper-Poisson bivectors on 4-dimensional hyperkähler manifolds.

Theorem 3.7. *Let (M, g, I_1, I_2, I_3) be a 4-dimensional hyperkähler manifold with corresponding Kähler forms $\omega_1, \omega_2, \omega_3$. A bivector Π on M is hyper-Poisson if and only if*

$$(3.4) \quad \Pi = f_1 \omega_1^{-1} + f_2 \omega_2^{-1} + f_3 \omega_3^{-1},$$

for smooth functions $f_1, f_2, f_3 : M \rightarrow \mathbb{R}$ satisfying $I_1 df_1 = I_2 df_2 = I_3 df_3$. Such a bivector is compatible with the hyperkähler structure.

Proof. If $\dim M = 4$, then $\Lambda^2 E \otimes S^2 H$ is spanned at each point by $\omega_1^{-1}, \omega_2^{-1}, \omega_3^{-1}$. Therefore Π must be of the form (3.4). Its $(2, 0)$ -component with respect to I_1 is equal to

$$\frac{1}{2}(f_2 + if_3)(\omega_2^{-1} - i\omega_3^{-1}) = 2(f_2 + if_3)(\omega_2 + i\omega_3)^{-1},$$

Hence, if Π is hyper-Poisson, then $f_2 + if_3$ is I_1 -holomorphic (since $\omega_2 + i\omega_3$ is I_1 -holomorphic). Similarly, $f_3 + if_1$ must be I_2 -holomorphic, and $f_1 + if_2$ must be I_3 -holomorphic. This triple of conditions is equivalent to $I_1 df_1 = I_2 df_2 = I_3 df_3$. Conversely, if the latter condition holds, then, for any complex structure, the $(2, 0)$ -part of Π is holomorphic. The $(2, 0)$ -part is also Poisson, since $\dim_{\mathbb{C}} M = 2$. For the same reason Π is compatible with the hyperkähler structure. \square

Remark 3.8. This result can be, of course, also proved by describing holomorphic sections of the line bundle $\Lambda^2 T_{\pi}^* Z$.

Corollary 3.9. *A 4-dimensional hyperkähler manifold M admits a hyper-Poisson bivector, other than a constant linear combination of $\omega_1^{-1}, \omega_2^{-1}, \omega_3^{-1}$, if and only if M admits a non-zero tri-Hamiltonian vector field. In this case the functions f_1, f_2, f_3 are the three moment maps for this vector field.* \square

Let M^{4d} be a hyperkähler manifold equipped with a compatible hyper-Poisson bivector Π . For each complex structure I_{ζ} , $\zeta \in \mathbb{P}^1$, we have the compatible holomorphic Poisson bivectors $\Pi_1 = \Omega_{\zeta}^{-1}$ and $\Pi_2 = \Pi_{\zeta}^{2,0}$. The Pfaffian polynomials of the corresponding recursion operators combine to define a polynomial $p(\zeta, \lambda)$ of the form

$$(3.5) \quad p(\zeta, \eta) = \lambda^d + p_1(\zeta)\lambda^{d-1} + \cdots + p_d(\zeta),$$

where the degree of p_i is $2i$. Each p_i defines a section of $\mathcal{O}(2i)$ on \mathbb{P}^1 , and (3.5) can be viewed as a holomorphic map

$$Z \longrightarrow \bigoplus_{i=1}^d |\mathcal{O}(2i)|,$$

where $|\mathcal{O}(2i)|$ denotes the total space of $\mathcal{O}(2i)$. This map is compatible with the real structures. In particular $p_1(\zeta)$ is a quadratic polynomial, compatible with the real structure of $|\mathcal{O}(2)|$, and hence p_1 is the hyperkähler moment map for a tri-Hamiltonian vector field X_{Π} . We shall call X_{Π} the *canonical Killing vector field*.

3.3. Hyperkähler transverse Hilbert schemes. Let M be a 4-dimensional hyperkähler manifold with a non-trivial tri-Hamiltonian Killing vector field. The moment map induces a holomorphic map μ from the twistor space Z of M to $|\mathcal{O}(2)|$. Following Atiyah and Hitchin [1] we can perform the transverse Hilbert scheme construction on fibres of $Z \rightarrow \mathbb{P}^1$ and obtain a new twistor space $Z_\mu^{[d]}$. Sections of $Z_\mu^{[d]} \rightarrow \mathbb{P}^1$ are in 1–1-correspondence with 1-dimensional compact complex subspaces C of Z such that:

- (i) the projection $\pi : C \rightarrow \mathbb{P}^1$ is flat with fibres of length d ;
- (ii) the projection μ induces a scheme-theoretic isomorphism between C and its image in $|\mathcal{O}(2)|$.

Observe, that given (ii), (i) simply means that $\mu(C)$ is defined by $p(\zeta, \lambda) = 0$, where p is as in (3.5).

Furthermore, as explained in [4], the normal bundle of the section of $Z_\mu^{[d]}$ corresponding to C splits as $\mathcal{O}(1)^{\oplus 2d}$ if and only if the normal sheaf $\mathcal{N}_{C/Z}$ of the curve C in Z satisfies $H^*(C, \mathcal{N}_{C/Z}(-2)) = 0$. On the Kodaira moduli space of such sections, satisfying in addition a reality condition, we obtain again a (pseudo)-hyperkähler metric. We shall denote this hyperkähler manifold by $M_\mu^{[d]}$ and refer to it as a *hyperkähler transverse Hilbert scheme*.

Consider now the $\mathcal{O}(2)$ -valued complex symplectic form ω on the fibres of Z , which can be viewed as a (holomorphic) section of $\Lambda^2(T_\pi Z(-1))^*$. Performing the construction of section 2.4 fibrewise on Z yields a fibrewise Poisson structure Π_2 on $Z_\mu^{[d]}$, which is a section of $\Lambda^2(T_\pi Z_\mu^{[d]}(-1)) \otimes \mathcal{O}(2) \simeq \Lambda^2 T_\pi Z_\mu^{[d]}$, i.e. it induces a bivector $\hat{\Pi}_2 \in \Lambda^2 E \otimes S^2 H$ on $M_\mu^{[d]}$. We shall see shortly that it is a purely imaginary bivector and, hence, $\Pi = -i\hat{\Pi}_2$ is real and, as discussed above, a hyper-Poisson bivector. Moreover, for each $\zeta \in \mathbb{P}^1$, its $(2,0)$ -component $\Pi_\zeta^{2,0}$ is compatible with Ω_ζ^{-1} , since Π_2 restricted to the fibre Z_ζ is compatible with ω_ζ^{-1} . Thus Π is a hyper-Poisson bivector compatible with the hyperkähler structure on $M_\mu^{[d]}$.

Example 3.10. Consider the case $d = 1$. The $(2,0)$ -part of $\hat{\Pi}_2$ is equal to $\mu\Omega^{-1}$, where Ω is the corresponding parallel holomorphic 2-form and μ is the corresponding holomorphic moment map. Comparing with Theorem 3.7, we conclude that

$$2\hat{\Pi}_2 = \mu_1\omega_1^{-1} + \mu_2\omega_2^{-1} + \mu_3\omega_3^{-1},$$

where $\omega_1, \omega_2, \omega_3$ denote the Kähler forms for complex structures I_1, I_2, I_3 and μ_1, μ_2, μ_3 are the corresponding moment maps. In particular, $\hat{\Pi}_2$ is purely imaginary. Since the real structure on $Z_\mu^{[d]}$ is induced from the one on Z , it follows that $\hat{\Pi}_2$ on $M_\mu^{[d]}$ is also purely imaginary.

Remark 3.11. Let Z be a complex 3-fold with a holomorphic map $\mu : Z \rightarrow |\mathcal{O}(2)|$ such that the composite map to \mathbb{P}^1 is surjective. Suppose further that Z has a fibrewise $\mathcal{O}(2)$ -valued complex symplectic form ω and a real structure covering the natural real structure on $|\mathcal{O}(2)|$. In other words, Z fulfills all conditions of the twistor space of a hyperkähler 4-manifold with a tri-Hamiltonian Killing vector field, except the existence of sections with normal bundle $\mathcal{O}(1) \oplus \mathcal{O}(1)$. In principle, it could happen that Z contains curves of degree $d > 1$, but not of degree 1 (although we do not know such an example). In this case $M_\mu^{[d]}$ is still well defined and a (pseudo)-hyperkähler manifold, although M does not exist. We shall still call $M_\mu^{[d]}$

a hyperkähler transverse Hilbert scheme, since the construction requires only the existence of Z , not necessarily of M .

We can characterise hyperkähler transverse Hilbert schemes arising from 4-manifolds with a locally free tri-Hamiltonian \mathbb{R} -action as follows:

Theorem 3.12. *Let M^{4d} be a hyperkähler manifold, equipped with a compatible hyper-Poisson bivector Π , such that for every complex structure I_ζ , the holomorphic Poisson bivectors $\Pi_1 = \Omega_\zeta^{-1}$ and $\Pi_2 = \Pi_\zeta^{2,0}$ satisfy the assumptions of Theorem 2.8. Then there exists a complex 3-fold Z with properties listed in Remark 3.11 such that M^{4d} is locally isomorphic to $M_\mu^{[d]}$ as a hyper-Poisson hyperkähler manifold. In addition, the holomorphic map $\mu : Z \rightarrow |\mathcal{O}(2)|$ is a submersion.*

Proof. We can perform the construction in the proof of Theorem 2.8 fibrewise on the twistor space of M^{4d} and obtain Z . Its properties follow easily. \square

Remark 3.13. This theorem remains true for pseudo-hyperkähler M^{4d} . We do not know whether the induced metric on a hyperkähler transverse Hilbert scheme is always positive definite.

Remark 3.14. The canonical Killing field X_Π is transverse to the foliation defined in the proof of Theorem 2.8 (on each fibre of the twistor space). Therefore the vertical vector field on Z , which gives the projection to $|\mathcal{O}(2)|$ is induced by X_Π . In particular, if X_Π integrates to an action of \mathbb{R} or S^1 on M^{4d} and the resulting 3-fold Z admits sections, then Z is the twistor of a hyperkähler 4-manifold with a tri-Hamiltonian action of \mathbb{R} or S^1 .

Remark 3.15. One can consider, more generally, hyperkähler transverse Hilbert schemes on 4-manifolds M , the twistor space of which maps to $|\mathcal{O}(2r)|$, $r > 1$, rather than to $|\mathcal{O}(2)|$. The corresponding object arising on $M_\mu^{[d]}$ is then a section Π of $\Lambda^2 E \otimes S^{2r} H$, satisfying $D\Pi = 0$. If we view Π as a $\Lambda^2 E$ -valued polynomial $\Pi(\zeta)$ of degree $2r$, then, for each $\zeta \in \mathbb{P}^1$, $\Pi(\zeta)$ defines an I_ζ -holomorphic Poisson bivector on $M_\mu^{[d]}$, compatible with Ω_ζ^{-1} . Theorem 3.12 remains true, as do the results of the next subsection (with obvious modifications).

3.4. Linear geometry of quaternionic bivectors. Let V be a real vector space of dimension $4n$, equipped with the standard flat quaternionic structure (g, I_1, I_2, I_3) . We denote the corresponding (linear) symplectic forms by $\omega_1, \omega_2, \omega_3$. The complexification $V^\mathbb{C}$ decomposes as $E \otimes H$, where E and H have complex dimensions $2n$ and n , respectively, and are equipped with the standard quaternionic-Hermitian structure, i.e. complex symplectic forms ω_E, ω_H and quaternionic structures σ_E, σ_H , so that $\omega_E(x, \sigma_E(x)) > 0$ and similarly for H .

Let Π be a bivector in $\Lambda^2 V$ belonging to $\Lambda^2 E \otimes S^2 H$. We define an endomorphism $A : V \rightarrow V$ by $\#_\Pi \circ \#_g^{-1}$. Since $\Pi \in \Lambda^2 E \otimes S^2 H$, it is an eigenvector of the operator $\sum_{i=1}^3 I_i \otimes I_i$ with eigenvalue -1 . Thus A is an eigenvector of $\sum_{i=1}^3 I_i \otimes I_i$ on $V^* \otimes V$ with eigenvalue 1 . Haydys [14] calls such endomorphisms *aquaternionic* and shows that they are of the form $A = I_1 A_1 + I_2 A_2 + I_3 A_3$, where each A_i is quaternionic, i.e. it commutes with I_1, I_2, I_3 . Since Π is antisymmetric and real, the A_i are quaternionic matrices which are quaternion-Hermitian, i.e. $A_i^\dagger = A_i$, where \dagger denotes the quaternionic adjoint. In terms of the symplectic structures (cf. Example 3.10).

$$\Pi(\alpha, \cdot) = A_1 \omega_1^{-1}(\alpha, \cdot) + A_2 \omega_2^{-1}(\alpha, \cdot) + A_3 \omega_3^{-1}(\alpha, \cdot).$$

It follows that the $(2, 0)$ component of Π for the complex structure I_1 is

$$\Pi^{2,0}(\alpha, \cdot) = (A_2 + iA_3)(\omega_2 + i\omega_3)^{-1}(\alpha, \cdot),$$

and similarly for other complex structures. Viewing the A_i as endomorphisms of E (which corresponds to the canonical homomorphism $\mathfrak{gl}(n, \mathbb{H}) \hookrightarrow \mathfrak{gl}(2n, \mathbb{C})$) we obtain a quadratic endomorphism

$$A(\zeta) = (A_2 + iA_3) + 2iA_1\zeta - (A_2 - iA_3)\zeta^2$$

of E , where each $A_i \in \mathfrak{gl}(2n, \mathbb{C})$ is symmetric with respect to the symplectic form ω_E . We can consider the sheaf morphism $\eta - A(\zeta) : E \otimes \mathcal{O}(-2) \rightarrow E$ on $T\mathbb{P}^1$. Its cokernel, which we denote by \mathcal{F} , is a 1-dimensional sheaf. Since each $A(\zeta)$ is symmetric with respect to the standard symplectic form on \mathbb{C}^{2n} , the characteristic polynomial of $A(\zeta)$ is of the form $p(\zeta, \eta)^2$, where $p(\zeta, \eta)$ is a polynomial of degree n in η . We call the scheme $C = \{(\zeta, \eta); p(\zeta, \eta) = 0\}$ as the *spectral curve of the bivector* Π and view the sheaf \mathcal{F} as being supported on C . If C is smooth, then \mathcal{F} is a rank 2 vector bundle with $\det \mathcal{F} \simeq K_C(2)$ (cf. [3], which contains more results on vector bundles arising this way). If the bivector Π arises via the hyperkähler transverse Hilbert scheme construction from a 3-dimensional twistor space Z , then the spectral curve C is precisely the curve in Z corresponding to a point in $M_\mu^{[n]}$ and the sheaf \mathcal{F} is isomorphic to $\mathcal{N}_{C/Z}(-1)$.

Remark 3.16. One can associate to Π another spectral object. Since the A_i are quaternion-Hermitian matrices, they are diagonalisable over \mathbb{H} with real eigenvalues. Let us denote the product of eigenvalues of a quaternion-Hermitian matrix X by $\det_{\mathbb{H}} X$ (the so-called *Moore determinant*). We can define a surface in $\mathbb{R}P^3$ as

$$S_{\mathbb{R}} = \{[x_0, x_1, x_2, x_3] \in \mathbb{R}P^3; \det_{\mathbb{H}}(x_0 - x_1A_1 - x_2A_2 - x_3A_3) = 0\}.$$

It is a (ramified) n -fold of $\mathbb{R}P^2$ (via $[x_0, x_1, x_2, x_3] \rightarrow [x_1, x_2, x_3]$), and its complexification is a surface S in $\mathbb{C}P^3$ which can be defined as

$$S = \{[z_0, z_1, z_2, z_3] \in \mathbb{C}P^3; \det(z_0 - z_1A_1 - z_2A_2 - z_3A_3) = 0\},$$

where the A_i are now complex $2n \times 2n$ matrices via the homomorphism $\mathfrak{gl}(n, \mathbb{H}) \hookrightarrow \mathfrak{gl}(2n, \mathbb{C})$. The intersection of S with the quadratic cone

$$x_1 = 2i\zeta, \quad x_2 = 1 - \zeta^2, \quad x_3 = i(1 + \zeta^2)$$

is the doubled spectral curve C . The sheaf \mathcal{F} extends to S and is defined as the cokernel of

$$z_0 - z_1A_1 - z_2A_2 - z_3A_3 : E \otimes \mathcal{O}(-2) \rightarrow E$$

on \mathbb{P}^3 . At present we do not understand the significance of $S_{\mathbb{R}}$ (as opposed to C) for the geometry of hyper-Poisson manifolds.

4. THE HYPER-POISSON BIVECTOR OF THE MONOPOLE MODULI SPACE

We consider the moduli space \mathcal{M}_k of $SU(2)$ -monopoles of charge k , described as the moduli space of $\mathfrak{u}(k)$ -valued solutions of Nahm's equations on $(0, 2)$, with simple poles at $t = 0, 2$ and residues defining the standard irreducible representation of $\mathfrak{su}(2)$. The Nahm equations are $\hat{T}_1 = [T_1, T_0] + [T_2, T_3]$ and two further equations, obtained by cyclic permutations of indices 1, 2, 3. The tangent space at

$[T_0, T_1, T_2, T_3]$ is given by quadruples of smooth maps (t_0, t_1, t_2, t_3) from $[0, 2]$ to $\mathfrak{u}(k)$ satisfying the equations

$$\begin{aligned}\dot{t}_0 &= [t_0, T_0] + [t_1, T_1] + [t_2, T_2] + [t_3, T_3] \\ \dot{t}_1 &= [T_1, t_0] + [t_1, T_0] + [T_2, t_3] + [t_2, T_3] \\ \dot{t}_2 &= [T_2, t_0] + [t_2, T_0] + [T_3, t_1] + [t_3, T_1] \\ \dot{t}_3 &= [T_3, t_0] + [t_3, T_0] + [T_1, t_2] + [t_1, T_2].\end{aligned}$$

The hypercomplex structure is given by the right multiplication by quaternions on $t_0 + t_1 i + t_2 j + t_3 k$ and the Riemannian metric g is

$$\|(t_0, t_1, t_2, t_3)\|^2 = - \int_0^2 \text{tr}(t_0^2 + t_1^2 + t_2^2 + t_3^2).$$

As explained by Atiyah and Hitchin in [1, Ch.6], the hyperkähler manifold \mathcal{M}_k is the hyperkähler transverse Hilbert scheme associated to $S^1 \times \mathbb{R}^3$. Thus, according to §3.3, it possesses a natural hyper-Poisson bivector compatible with the hyperkähler structure. We can identify this bivector as follows:

Theorem 4.1. *The natural hyper-Poisson bivector Π on the moduli space \mathcal{M}_k is given by*

$$\#_g^{-1}\Pi = \frac{-i}{4} \int_0^2 \text{tr} \left(\sum_{i=1}^3 T_i (dT_i \wedge dT_0 - dT_0 \wedge dT_i) + \sum_{i,j,k=1}^3 \epsilon_{ijk} T_i dT_j \wedge dT_k \right),$$

where $(\phi \wedge \psi)(a, b) = \phi(a)\psi(b) - \phi(b)\psi(a)$.

Remark 4.2. It will follow from the proof that the integral is finite.

Proof. We first compute the Kähler forms ω_2 and ω_3 corresponding to the complex structures J and K :

$$\omega_2 = - \int_0^2 \text{tr} (dT_0 \wedge dT_2 + dT_1 \wedge dT_3), \quad \omega_3 = - \int_0^2 \text{tr} (dT_0 \wedge dT_3 + dT_2 \wedge dT_1).$$

The I -holomorphic 2-form $\omega_2 + i\omega_3$ is therefore given by

$$\omega_2 + i\omega_3 = - \int_0^2 \text{tr} d(T_0 - iT_1) \wedge d(T_2 + iT_3).$$

We can now rewrite the integrand in the formula for $\#_g^{-1}\Pi$ as follows:

$$\begin{aligned}\frac{1}{2} \text{tr} ((T_2 + iT_3)(\Phi_2 - i\Phi_3) + \\ + 2T_1(dT_1 \wedge dT_0 - dT_0 \wedge dT_1 + dT_2 \wedge dT_3 - dT_3 \wedge dT_2) + \\ + (T_2 - iT_3)(\Phi_2 + i\Phi_3)),\end{aligned}$$

where

$$\Phi_2 = dT_2 \wedge dT_0 - dT_0 \wedge dT_2 + dT_3 \wedge dT_1 - dT_1 \wedge dT_3,$$

$$\Phi_3 = dT_3 \wedge dT_0 - dT_0 \wedge dT_3 + dT_1 \wedge dT_2 - dT_2 \wedge dT_1.$$

Observe that the first summand is of type $(0, 2)$ for the complex structure I , the second one of type $(1, 1)$, and the third one of type $(2, 0)$. Since $\#_g$ exchanges $(2, 0)$

and $(0, 2)$, we conclude that

$$\begin{aligned}\#_g^{-1}\Pi^{2,0} &= \frac{-i}{8} \int_0^2 \text{tr}(T_2 + iT_3)(\Phi_2 - i\Phi_3) = \\ &= \frac{-i}{8} \int_0^2 \text{tr}(T_2 + iT_3)(d(T_2 - iT_3) \wedge d(T_0 + iT_1) - d(T_0 + iT_1) \wedge d(T_2 - iT_3)).\end{aligned}$$

We now observe that for any bivector π and $\Omega = \omega_2 + i\omega_3$:

$$(4.1) \quad \left(\#_{\Omega}^{-1}\pi\right)(u, v) = \left(\#_g^{-1}\pi\right)(Ju + iKu, Jv + iKv).$$

Computing this for $\pi = \Pi^{2,0}$ we obtain:

$$\#_{\Omega}^{-1}\Pi^{2,0} = \frac{-i}{2} \int_0^2 \text{tr}(T_2 + iT_3)(d(T_2 + iT_3) \wedge d(T_0 - iT_1) - d(T_0 - iT_1) \wedge d(T_2 + iT_3)).$$

Thus, if we set $\beta = T_2 + iT_3$ and $\alpha = T_0 - iT_1$, we obtain

$$(4.2) \quad i\#_{\Omega}^{-1}\Pi^{2,0} = \frac{1}{2} \int_0^2 \text{tr} d(\beta^2) \wedge d\alpha.$$

The complex Nahm equation is the Lax equation $\dot{\beta} = [\beta, \alpha]$. It follows that (β^2, α) also satisfies the Lax equation. Acting by a singular complex gauge transformation which makes α equal to zero (and, consequently, β constant) implies that

$$i\#_{\Omega}^{-1}\Pi^{2,0} = \frac{1}{2} \sum d(\beta_j)^2 \wedge \frac{dp_j}{p_j} = \sum \beta_j d\beta_j \wedge \frac{dp_j}{p_j},$$

where β_j are the poles and p_j the values of the numerator of the rational map corresponding to the given monopole (and the complex structure I). This means that $\Pi^{2,0}$ is precisely the holomorphic bivector obtained from the transverse Hilbert scheme construction applied to $\mathbb{C} \times \mathbb{C}^*$ with the symplectic form $d\beta \wedge \frac{dp}{p}$.

Observe now that, since $\#_g^{-1}\Pi^{2,0} = (\#_g^{-1}\Pi)^{0,2}$ and $Ju + iKu = (Ju) + iI(Ju)$, the formula (4.1) implies that

$$(\#_g^{-1}\Pi)^{0,2}(u, v) = \#_{\Omega}^{-1}\Pi^{2,0}(-Ju, -Jv).$$

Therefore the integral defining $(\#_g^{-1}\Pi)^{0,2}$ is finite. Similarly, the integral defining $(\#_g^{-1}\Pi)^{2,0}$ is finite. Since $(\#_g^{-1}\Pi)^{0,2} + (\#_g^{-1}\Pi)^{2,0}$ is the sum of all terms of the form $T_2 \cdot \phi$ and $T_3 \cdot \phi$ in the formula in the statement, repeating this decomposition for the complex structure J or K shows that the whole integral in the statement is finite. This also shows that

$$\Pi = \frac{1}{2} \left(\Pi_I^{2,0} + \Pi_{-I}^{2,0} + \Pi_J^{2,0} + \Pi_{-J}^{2,0} + \Pi_K^{2,0} + \Pi_{-K}^{2,0} \right),$$

and so, owing to Lemma 4.2 in [21], $\Pi \in \Gamma(\Lambda^2 E \otimes S^2 H)$. Finally, observe that Π is real. \square

Remark 4.3. The canonical Killing vector field X_{Π} (see §3.2) on the moduli space \mathcal{M}_k is $(t_0, t_1, t_2, t_3) = (i, 0, 0, 0)$. We obtain

$$i(X_{\Pi})\#_g^{-1}\Pi = -\frac{1}{2} \int_0^2 \text{tr} \sum_{i=1}^3 T_i dT_i = -\frac{1}{4} dF,$$

where

$$F = \int_0^2 \left(\operatorname{tr} \sum_{i=1}^3 T_i^2 + \frac{k(k^2 - 1)}{4} (s^{-2} + (s - 2)^{-2}) \right).$$

The function F has been shown by Hitchin [15] to essentially give Kähler potentials of the monopole metric: for every complex structure, the sum of F and some linear combination of the coefficients of the spectral curve is a Kähler potential for the corresponding Kähler form. Is this true on a general hyperkähler transverse Hilbert scheme $M_\mu^{[d]}$, i.e. is the 1-form $i(X_\Pi)\#_g^{-1}\Pi$ similarly related to Kähler potentials on $M_\mu^{[d]}$?

REFERENCES

- [1] M.F. Atiyah and N.J. Hitchin, *The geometry and dynamics of magnetic monopoles*, Princeton University Press, Princeton (1988).
- [2] A. Beauville, ‘Variétés Kähleriennes dont la première classe de Chern est nulle’, *J. Differential Geom.* 18 (1983), no. 4, 755–782.
- [3] A. Beauville, ‘Determinantal hypersurfaces’, *Michigan Math. J.* 48 (2000), 39–64.
- [4] R. Bielawski, ‘Hyperkähler manifolds of curves in twistor spaces’, *SIGMA* 10 (2014).
- [5] R. Bielawski, ‘Slices to sums of adjoint orbits, the Atiyah-Hitchin manifold, and Hilbert schemes of points’, *Complex Manifolds* 4 (2017), 16–36.
- [6] R. Bielawski and L. Schwachhöfer, ‘Hypercomplex limits of pluricomplex structures and the Euclidean limit of hyperbolic monopoles’, *Ann. Global Anal. Geom.* 44 (2013), 245–256.
- [7] F. Bottacin, ‘Poisson structures on Hilbert schemes of points of a surface and integrable systems’, *manuscripta math.* 97 (1998), 517–527.
- [8] G.R. Cavalcanti, ‘Examples and counter-examples of log-symplectic manifolds’, *J. Topol.* 10 (2017), 1–21.
- [9] R.L. Fernandes, ‘Completely integrable bi-Hamiltonian systems’, *J. Dyn. Diff. Equat.* 6 (1994), 53–69.
- [10] R. Goto, ‘Rozansky-Witten invariants of log symplectic manifolds’, in *Integrable systems, topology, and physics (Tokyo, 2000)*, Contemp. Math., vol. 309, AMS, Providence, RI, 2002, 69–84.
- [11] M. Gualtieri and S. Li, ‘Symplectic groupoids of log symplectic manifolds’, *IMRN* 2014 (11), 3022–3074, 2014.
- [12] V. Guillemin, E. Miranda, and A. R. Pires, ‘Symplectic and Poisson geometry on b-manifolds’, *Adv. Math.* 264 (2014), 864–896.
- [13] R. Hartshorne, *Deformation Theory*, Springer, New York, 2010.
- [14] A. Haydys, ‘Nonlinear Dirac operator and quaternionic analysis’, *Commun. Math. Phys.* 281 (2008), 251–261.
- [15] N.J. Hitchin, ‘Integrable systems in Riemannian geometry, in: *Surveys in differential geometry: integral systems*, 21–81, Int. Press, Boston, 1998.
- [16] W. Hong and M. Stiénon, ‘From hypercomplex to holomorphic symplectic structures’, *J. Geom. Phys.* 96 (2015), 187–203.
- [17] N. Lora Lamia Donin, ‘Transverse Hilbert schemes and completely integrable systems’, *Complex Manifolds* 4 (2017), 263–272.
- [18] N. Lora Lamia Donin, ‘Hyperkähler manifolds of curves and l-hypercomplex structures’, Ph.D. Thesis, Leibniz Universität Hannover, 2018.
- [19] F. Magri and C. Morosi, ‘A geometrical characterization of integrable Hamiltonian systems through the theory of Poisson-Nijenhuis manifolds’, *Quaderno S.* 19, Univ. of Milan, 1984.
- [20] C. Manolescu, ‘Nilpotent slices, Hilbert schemes, and the Jones polynomial’, *Duke Math. J.* 132 (2006), 311–369.
- [21] S.M. Salamon, ‘Differential geometry of quaternionic manifolds’, *Ann. Sci. Éc. Norm. Supér. Serie 4*, Volume 19 (1986), p. 31–55.
- [22] R.S. Ward, ‘On self-dual gauge fields’, *Phys. Lett. A* 61 (1977), 81–82.