

ON A STOCHASTIC CAMASSA-HOLM TYPE EQUATION WITH HIGHER ORDER NONLINEARITIES

CHRISTIAN ROHDE AND HAO TANG

ABSTRACT. The subject of this paper is a generalized Camassa-Holm equation under random perturbation. We first establish local existence and uniqueness results as well as blow-up criteria for pathwise solutions in the Sobolev spaces H^s with $s > 3/2$. Then we analyze how noise affects the dependence of solutions on initial data. Even though the noise has some already known regularization effects, much less is known concerning the dependence on initial data. As a new concept we introduce the notion of stability of exiting times and construct an example showing that multiplicative noise (in Itô sense) cannot improve the stability of the exiting time, and simultaneously improve the continuity of the dependence on initial data. Finally, we obtain global existence theorems and estimate associated probabilities.

1. INTRODUCTION

We consider a stochastic version of the generalized Camassa-Holm equation. Let t denote the time variable and let x be the one-dimensional space variable. The equation is given for $k \in \mathbb{N}$ by

$$u_t - u_{xxt} + (k+2)u^k u_x - (1 - \partial_{xx}^2)h(t, u)\dot{W} = (k+1)u^{k-1}u_x u_{xx} + u^k u_{xxx}. \quad (1.1)$$

In (1.1), $h : \mathbb{R}^+ \times \mathbb{R} \rightarrow \mathbb{R}$ is some nonlinear function and \dot{W} is a cylindrical Wiener process. We will consider (1.1) on the torus, i.e. $x \in \mathbb{T} = \mathbb{R}/2\pi\mathbb{Z}$.

For $h = 0$ and $k = 1$, equation (1.1) reduces to the deterministic Camassa-Holm (CH) equation given by

$$u_t - u_{xxt} + 3uu_x = 2u_x u_{xx} + uu_{xxx}. \quad (1.2)$$

Fokas & Fuchssteiner [24] introduced (1.2) when studying completely integrable generalizations of the Korteweg-de-Vries (KdV) equation with bi-Hamiltonian structure whereas Camassa & Holm [8] proposed (1.2) to describe the unidirectional propagation of shallow water waves over a flat bottom. Since then, the CH equation (1.2) has been studied intensively, and we cannot even attempt to survey the vast research history here. For the paper at hand it is important to mention the wave-breaking phenomenon which illustrates possible loss of regularity as a fundamental mechanism in the CH equations. In contrast to smooth soliton solutions to the KdV equation [44], solutions to the CH equation remain indeed bounded but their slope can become unbounded in finite time, cf. [14] and related work in [12, 15, 52]. Moreover, for a smooth initial profile, it is possible to predict exactly (by establishing a necessary and sufficient condition) whether wave-breaking occurs for solutions to the Cauchy problem for (1.2) [14, 52]. The other essential feature of the CH equation is the occurrence of traveling waves with a peak at their crest, exactly like that the governing equations for water waves admit the so-called Stokes waves of the greatest height, see [13, 16, 17]. Bressan & Constantin proved the existence of dissipative as well as conservative solutions in [6, 7]. Later, Holden & Raynaud [37, 38] also obtained global conservative and dissipative solutions using Lagrangian transport ideas.

When $h = 0$ and $k = 2$, equation (1.1) forms the cubic equation

$$u_t - u_{xxt} + 4u^2 u_x = 3uu_x u_{xx} + u^2 u_{xxx}, \quad (1.3)$$

which has been derived by Novikov in [55]. It has been proven that (1.3) possesses a bi-Hamiltonian structure with an infinite sequence of conserved quantities and (1.3) admits peaked solutions of explicit

2010 *Mathematics Subject Classification.* Primary: 60H15, 35Q51; Secondary: 35A01, 35B30.

Key words and phrases. Stochastic generalized Camassa-Holm equation; pathwise solution; noise effect; exiting time; dependence on initial data; global existence;

C. R. acknowledges funding by Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC 2075–390740016.

H. T. is supported by the Alexander von Humboldt Foundation.

form $u(t, x) = \pm \sqrt{c} e^{-|x-ct|}$ with $c > 0$ [26], as well as multipeakon solutions with explicit formulas [39]. For the study of other deterministic instances of (1.1) we refer to [9, 32, 66, 67].

Equations like (1.2) are naturally embedded in high-order descriptions with energy-dissipative evolution. A weakly dissipative evolution is given for some parameter $\lambda > 0$ by

$$u_t - u_{xxt} + (k+2)u^k u_x - (1 - \partial_{xx}^2)(\lambda u) = (k+1)u^{k-1}u_x u_{xx} + u^k u_{xxx}, \quad x \in \mathbb{T}, t > 0. \quad (1.4)$$

For instance, the weakly dissipative CH equation, i.e. (1.4) for $k = 1$, has been studied in [49, 65]. For the Novikov equation (1.3) with the same weakly dissipative term $(1 - \partial_{xx}^2)(\lambda u)$, we can also refer to [49].

In this work, we assume that the energy exchange mechanisms are connected with randomness to account for external stochastic influence. We are interested in the case where in the term $(1 - \partial_{xx}^2)(\lambda u)$ the deterministic parameter λ is substituted by a formal cylindrical Wiener process (cf. Section 2 for precise definitions), and where the previously linear dependency on u is replaced by a non-autonomous and nonlinear term $h(t, u)$. Thus, we consider the Cauchy problem for (1.1) on the torus \mathbb{T} , with random initial data $u_0 = u_0(\omega, x)$. Applying the operator $(1 - \partial_{xx}^2)^{-1}$ to (1.1), we reformulate the problem as the stochastic evolution

$$\begin{cases} du + [u^k \partial_x u + F(u)] dt = h(t, u) d\mathcal{W}, & x \in \mathbb{T}, t > 0, k \geq 1, \\ u(\omega, 0, x) = u_0(\omega, x), & x \in \mathbb{T}, \end{cases} \quad (1.5)$$

with $F(u) = F_1(u) + F_2(u) + F_3(u)$ and

$$\begin{aligned} F_1(u) &= (1 - \partial_{xx}^2)^{-1} \partial_x (u^{k+1}), \\ F_2(u) &= \frac{2k-1}{2} (1 - \partial_{xx}^2)^{-1} \partial_x (u^{k-1} u_x^2), \\ F_3(u) &= \frac{k-1}{2} (1 - \partial_{xx}^2)^{-1} (u^{k-2} u_x^3). \end{aligned} \quad (1.6)$$

Here we remark that the operator $(1 - \partial_{xx}^2)^{-1}$ in $F(\cdot)$ is understood as

$$[(1 - \partial_{xx}^2)^{-1} f](x) = [G_{\mathbb{T}} * f](x) \text{ for } f \in L^2(\mathbb{T}) \text{ with } G_{\mathbb{T}} = \frac{\cosh(x - 2\pi [\frac{x}{2\pi}] - \pi)}{2 \sinh(\pi)}, \quad (1.7)$$

where the convolution is on \mathbb{T} only, and the symbol $[x]$ stands for the integer part of x .

The first objective of this paper is to analyze the local existence and uniqueness of pathwise solutions as well as blow-up criteria for problem (1.5) with nonlinear multiplicative noise (see Theorem 2.1). We note that for the CH equation with additive noise, existence and uniqueness has been obtained in [11]. For the stochastic modified CH equation with linear multiplicative noise, we refer to [10]. When the multiplicative noise is given by a one-dimensional Wiener process with H^s diffusion coefficient, the stochastic CH equation was considered in second author's work [58]. In this paper, we consider the local existence and uniqueness of pathwise solutions to (1.5) as well as blow-up criteria for more general noise given by a cylindrical Wiener process with nonlinear coefficient.

The second major objective of the paper is to investigate whether stochastic perturbations can improve the dependence on initial data in the Cauchy problem (1.5). We notice that various instances of transport-type stochastic evolution laws have been studied with respect to the effect of noise on the regularity of their solutions. For example, we refer to [23, 53, 54, 21] for linear stochastic transport equations and to what concerns nonlinear stochastic conservation laws, we refer to [50, 51, 27]. In [46, 29, 57, 58] the dissipation of energy caused by linear multiplicative noise has been analyzed.

In contrast to these works we focus in this paper on the initial-data dependence in the Cauchy problem (1.5). Actually, much less is known on the noise effect with respect to the dependence on initial data. But the question whether (and how) noise can affect initial-data dependence is interesting. Formally speaking, regularization produced by noise may be seen like the deterministic regularization effect induced by adding a Laplacian. However, if one would add a real Laplacian to the governing equations, then by using techniques from the theory for semilinear parabolic equations, the dependence on initial data in some cases turns out to be more regular than being just continuous. For example, for the deterministic incompressible Euler equations, the dependence on initial data cannot be better than continuous [35], but for the deterministic incompressible Navier-Stokes equations with sufficiently large viscosity, it is at least Lipschitz continuous in sufficiently high Sobolev norms, see pp. 79–81 in [31]. This motivates us to study whether (and how) noise can affect initial-data dependence. Actually, to our knowledge, there are very few results in this direction. We can only refer to [59] for the stochastic Euler-Poincaré equation

with respect to this problem. However, in this work, we have higher order nonlinearities, which requires more technical estimates. To analyze initial-data dependence, we introduce the concept of the stability of the exiting time (see Definition 2.2 below), cf. [59]. Then we show under some conditions on $h(t, u)$, that the multiplicative noise (in the Itô sense) cannot improve the stability of the exiting time, *and*, at the same time, improve the continuity of the map $u_0 \mapsto u$ (see Theorem 2.2). It is worth noting that in deterministic cases, the issue of the optimal dependence of solutions (for example, the solution map is continuous but not uniformly continuous) to various nonlinear dispersive and integrable equations has been the subject of many papers. One of the first results of this type dates back at least as far as to Kato [42]. Indeed, Kato [42] proved that the solution map $u_0 \mapsto u$ in $H^s(\mathbb{T})$ ($s > 3/2$), given by the inviscid Burgers equation, is not Hölder continuous regardless of the Hölder exponent. Since then different techniques have been successfully applied to various nonlinear dispersive and integrable equations, see [45, 47, 2] for example. Particularly, for the incompressible Euler equation, we refer to [35, 60], and for CH type equations, we refer to [33, 34, 61, 62, 63] and the references therein.

We conclude the paper with a discussion of time-global well-posedness for (1.5). For general noise terms, it is difficult to determine whether a local pathwise solution to (1.5) is globally defined. As shown in [29, 46, 48, 57, 58], the linear noise $h(t, u)dW = \beta u dW$, with $\beta \in \mathbb{R} \setminus \{0\}$ and W being a standard 1-D Brownian motion, acts dissipatively for many SPDEs. Motivated by these works, we prove some results in Theorems 2.3 and 2.4 for the global dynamics of (1.5) with linear multiplicative noise, that is

$$\begin{cases} du + [u^k \partial_x u + F(u)] dt = b(t)u dW, & x \in \mathbb{T}, t \in \mathbb{R}^+, k \geq 1, \\ u(\omega, 0, x) = u_0(\omega, x), & x \in \mathbb{T}. \end{cases} \quad (1.8)$$

2. PRELIMINARIES AND MAIN RESULTS

2.1. Noise with H^s coefficient. We begin by introducing some notations. Throughout the paper, $(\Omega, \mathcal{F}, \mathbb{P})$ denotes a complete probability space, where \mathbb{P} is the probability measure on Ω and \mathcal{F} is a σ -algebra. Let $t > 0$. $\sigma\left\{(x(\tau), y(\tau))_{\tau \in [0, t]}\right\}$ stands for the completion of the union σ -algebra generated by $(x(\tau), y(\tau))$. All stochastic integrals are defined in the Itô sense and $\mathbb{E} \cdot$ is the mathematical expectation of \cdot with respect to \mathbb{P} . For some separable Banach space X , $\mathcal{B}(X)$ denotes the Borel sets of X , and $\mathcal{P}r(X)$ stands for the collection of Borel probability measures on X . For $E \subseteq X$, $\mathbf{1}_E$ is the indicator function on E .

$L^2(\mathbb{T})$ is the usual square integrable function space on \mathbb{T} . For $s \in \mathbb{R}$, $D^s = (1 - \partial_{xx}^2)^{s/2}$ is defined by $\widehat{D^s f}(k) = (1 + k^2)^{s/2} \widehat{f}(k)$, where \widehat{g} denote the Fourier transform of g on \mathbb{T} . The Sobolev space $H^s(\mathbb{T})$ is defined as

$$H^s(\mathbb{T}) \triangleq \{f \in L^2(\mathbb{T}) : \|f\|_{H^s(\mathbb{T})}^2 = \sum_{k \in \mathbb{Z}} (1 + k^2)^s |\widehat{f}(k)|^2 < +\infty\},$$

and the inner product on $H^s(\mathbb{T})$ is $(f, g)_{H^s} \triangleq \sum_{k \in \mathbb{Z}} (1 + k^2)^s \widehat{f}(k) \cdot \overline{\widehat{g}(k)} = (D^s f, D^s g)_{L^2}$. For function spaces on \mathbb{T} , we will drop \mathbb{T} if there is no ambiguity. We will use \lesssim to denote estimates that hold up to some universal *deterministic* constant which may change from line to line but whose meaning is clear from the context. For linear operators A and B , $[A, B]$ stands for the commutator of A and B , i.e., $[A, B] = AB - BA$.

We briefly recall some aspects of the stochastic analysis theory which we use below. We refer the readers to [19, 25, 40] for an extended treatment of this subject.

We call $\mathcal{S} = (\Omega, \mathcal{F}, \mathbb{P}, \{\mathcal{F}_t\}_{t \geq 0}, \mathcal{W})$ a stochastic basis, where $\{\mathcal{F}_t\}_{t \geq 0}$ is a right-continuous filtration on (Ω, \mathcal{F}) such that $\{\mathcal{F}_0\}$ contains all the \mathbb{P} -negligible subsets and $\mathcal{W}(t) = \mathcal{W}(\omega, t), \omega \in \Omega$ is a cylindrical Brownian motion, defined on an auxiliary Hilbert space U , which is adapted to $\{\mathcal{F}_t\}_{t \geq 0}$. Formally, if $\{e_k\}$ is a complete orthonormal basis of U and $\{W_k\}_{k \geq 1}$ is a sequence of mutually independent standard one-dimensional Brownian motions, then one may define

$$\mathcal{W} = \sum_{k=1}^{\infty} e_k W_k, \quad \mathbb{P} - a.s.$$

To guarantee the convergence of the (formal) summation above, we consider a larger separable Hilbert space U_0 such that the canonical embedding $U \hookrightarrow U_0$ is Hilbert-Schmidt. Then we have that for any

$T > 0$, cf. [19, 25, 41],

$$\mathcal{W} = \sum_{k=1}^{\infty} e_k W_k \in C([0, T]; U_0), \quad \mathbb{P} - a.s.$$

To define the Itô stochastic integral

$$\int_0^\tau G d\mathcal{W} = \sum_{k=1}^{\infty} \int_0^\tau G e_k dW_k \quad (2.1)$$

on H^s , it is required (see e.g. [19, 56]) for the predictable stochastic process G to take values in the space of Hilbert-Schmidt operators from U to H^s , denoted by $L_2(U; H^s)$. For such G , (2.1) is a well-defined continuous H^s -valued square integrable martingale such that for all almost surely bounded stopping times τ and for all $v \in H^s$,

$$\left(\int_0^\tau G d\mathcal{W}, v \right)_{H^s} = \sum_{k=1}^{\infty} \int_0^\tau (G e_k, v)_{H^s} dW_k.$$

Moreover, the desirable Burkholder-Davis-Gundy inequality in our case turns out to be

$$\mathbb{E} \left(\sup_{t \in [0, T]} \left\| \int_0^t G d\mathcal{W} \right\|_{H^s}^p \right) \leq C(p, s) \mathbb{E} \left(\int_0^T \|G\|_{L_2(U; H^s)}^2 dt \right)^{\frac{p}{2}}, \quad p \geq 1.$$

or in terms of the coefficients,

$$\mathbb{E} \left(\sup_{t \in [0, T]} \left\| \sum_{k=1}^{\infty} \int_0^t G e_k dW_k \right\|_{H^s}^p \right) \leq C(p, s) \mathbb{E} \left(\int_0^T \sum_{k=1}^{\infty} \|G e_k\|_{H^s}^2 dt \right)^{\frac{p}{2}}, \quad p \geq 1.$$

Here we remark that the stochastic integral (2.1) does not depend on the choice of the space U_0 , cf. [19, 56]. For example, U_0 can be defined as

$$U_0 = \left\{ v = \sum_{k=1}^{\infty} a_k e_k : \sum_{k=1}^{\infty} \frac{a_k^2}{k^2} < \infty \right\}, \quad \|v\|_{U_0} = \sum_{k=1}^{\infty} \frac{a_k^2}{k^2}.$$

2.2. Definitions of the solutions and stability of the exiting times. We now make the precise notion of a pathwise solution to (1.5).

Definition 2.1 (Pathwise solutions). *Let $\mathcal{S} = (\Omega, \mathcal{F}, \mathbb{P}, \{\mathcal{F}_t\}_{t \geq 0}, \mathcal{W})$ be a fixed stochastic basis. Let $s > 3/2$ and u_0 be an H^s -valued \mathcal{F}_0 -measurable random variable (relative to \mathcal{S}).*

- (1) *A local pathwise solution to (1.5) is a pair (u, τ) , where $\tau \geq 0$ is a stopping time satisfying $\mathbb{P}\{\tau > 0\} = 1$ and $u : \Omega \times [0, \infty] \rightarrow H^s$ is an \mathcal{F}_t -predictable H^s -valued process satisfying*

$$u(\cdot \wedge \tau) \in C([0, \infty); H^s) \quad \mathbb{P} - a.s.,$$

and for all $t > 0$,

$$u(t \wedge \tau) - u(0) + \int_0^{t \wedge \tau} [u^k \partial_x u + F(u)] dt' = \int_0^{t \wedge \tau} h(t', u) d\mathcal{W} \quad \mathbb{P} - a.s.$$

- (2) *The local pathwise solutions are said to be pathwise unique, if given any two pairs of local pathwise solutions (u_1, τ_1) and (u_2, τ_2) with $\mathbb{P}\{u_1(0) = u_2(0)\} = 1$, we have*

$$\mathbb{P}\{u_1(t, x) = u_2(t, x), \quad \forall (t, x) \in [0, \tau_1 \wedge \tau_2] \times \mathbb{T}\} = 1.$$

- (3) *Additionally, (u, τ^*) is called a maximal pathwise solution to (1.5) if $\tau^* > 0$ almost surely and if there is an increasing sequence $\tau_n \rightarrow \tau^*$ such that for any $n \in \mathbb{N}$, (u, τ_n) is a pathwise solution to (1.5) and on the set $\{\tau^* < \infty\}$,*

$$\sup_{t \in [0, \tau_n]} \|u\|_{H^s} \geq n.$$

- (4) *If (u, τ^*) is a maximal pathwise solution and $\tau^* = \infty$ almost surely, then we say that the pathwise solution exists globally.*

A major result of this paper is a (negative) dependence statement for the initial data of the solutions. Precisely, it refers to the stability of a point in time when the solution leaves a certain range. This point in time is called exiting time and we introduce

Definition 2.2 (Stability of exiting time). *Let $s > 3/2$ and $\mathcal{S} = (\Omega, \mathcal{F}, \mathbb{P}, \{\mathcal{F}_t\}_{t \geq 0}, \mathcal{W})$ be a fixed stochastic basis. Let u_0 be an H^s -valued \mathcal{F}_0 -measurable random variable such that $\mathbb{E}\|u_0\|_{H^s}^2 < \infty$. Assume that $\{u_{0,n}\}$ is any sequence of H^s -valued \mathcal{F}_0 -measurable random variables satisfying $\mathbb{E}\|u_{0,n}\|_{H^s}^2 < \infty$. For each n , let u and u_n be the unique pathwise solutions to (1.5) with initial value u_0 and $u_{0,n}$, respectively. For any $R > 0$ and $n \in \mathbb{N}$, we define the R -exiting times*

$$\tau_n^R := \inf \{t \geq 0 : \|u_n(t)\|_{H^s} > R\} \quad \text{and} \quad \tau^R := \inf \{t \geq 0 : \|u(t)\|_{H^s} > R\},$$

where $\inf \emptyset = \infty$. Furthermore,

- (1) If $u_{0,n} \rightarrow u_0$ in H^s \mathbb{P} -a.s. implies that

$$\lim_{n \rightarrow \infty} \tau_n^R = \tau^R \quad \mathbb{P} - \text{a.s.}, \quad (2.2)$$

then the R -exiting time of u is said to be stable.

- (2) If $u_{0,n} \rightarrow u_0$ in $H^{s'}$ for all $s' < s$ almost surely implies that (2.2) holds true, the R -exiting time of u is said to be strongly stable.

2.3. Assumptions. For our results on existence of pathwise solutions, on the stability of exiting times and global well-posedness of (1.1), we rely in the following sections on generic but slightly different assumptions concerning the data in (1.1). These are summarized here, for a comment on possible relaxed versions we refer to Remark 3.1.

Assumption A₁. We assume that $h : [0, \infty) \times H^s \ni (t, u) \mapsto h(t, u) \in L_2(U; H^s)$ for $u \in H^s$ with $s \geq 0$ such that if $u : \Omega \times [0, T] \rightarrow H^s$ is predictable, then $h(t, u)$ is also predictable. Furthermore, we assume the following:

- (1) There is an increasing and locally bounded function $f(\cdot) : [0, +\infty) \rightarrow [0, +\infty)$ with $f(0) = 0$ such that for any $t > 0$ and $u \in H^s$ with $s > \frac{1}{2}$,

$$\|h(t, u)\|_{L_2(U; H^s)} \leq f(\|u\|_{W^{1,\infty}})(1 + \|u\|_{H^s}). \quad (2.3)$$

Particularly, if h does not depend on u , i.e., the additive noise case, then the condition $f(0) = 0$ can be removed.

- (2) There is an increasing locally bounded function $g(\cdot) : [0, +\infty) \rightarrow [0, +\infty)$, such that for any $t > 0$ and $u \in H^s$ with $s > \frac{1}{2}$,

$$\|h(t, u) - h(t, v)\|_{L_2(U; H^s)} \leq g(\|u\|_{H^s} + \|v\|_{H^s})\|u - v\|_{H^s}. \quad (2.4)$$

Assumption A₂. When we consider the initial-data dependence problem for (1.5) in Section 4, we need a similar assumption on $h(t, \cdot)$. For $s \geq 0$ and $u \in H^s$, $h : [0, \infty) \times H^s \ni (t, u) \mapsto h(t, u) \in L_2(U; H^s)$ for $u \in H^s$ with $s \geq 0$ such that if $u : \Omega \times [0, T] \rightarrow H^s$ is predictable, then $h(t, u)$ is also predictable. Moreover, for all $t \geq 0$, we assume

$$\|h(t, u)\|_{L_2(U; H^s)} \leq \|F(u)\|_{H^s}, \quad \|h(t, u) - h(t, v)\|_{L_2(U; H^s)} \leq \|F(u) - F(v)\|_{H^s}, \quad (2.5)$$

where $F(\cdot)$ is defined by (1.6).

Assumption A₃. When considering (1.8) with non-autonomous linear noise $b(t)udW$, we assume that $b(t) \in C([0, \infty))$ such that $0 < b_* \leq b^2(0) \leq \sup_{t \geq 0} b^2(t) \leq b^*$ for some $b_*, b^* \in \mathbb{R}$.

2.4. Main results. In this section we summarize our major contributions providing proofs later in the remainder of the paper.

Theorem 2.1. *Let $s > 3/2$, $k \geq 1$ and let $h(t, u)$ satisfy Assumption A₁. For a given stochastic basis $\mathcal{S} = (\Omega, \mathcal{F}, \mathbb{P}, \{\mathcal{F}_t\}_{t \geq 0}, \mathcal{W})$, if u_0 is an H^s -valued \mathcal{F}_0 -measurable random variable satisfying $\mathbb{E}\|u_0\|_{H^s}^2 < \infty$, then there is a local unique pathwise solution (u, τ) to (1.5) in the sense of Definition 2.1 with*

$$u(\cdot \wedge \tau) \in L^2(\Omega; C([0, \infty); H^s)). \quad (2.6)$$

Moreover, (u, τ) can be extended to a unique maximal pathwise solution (u, τ^*) with

$$\mathbf{1}_{\{\tau^* < \infty\}} = \mathbf{1}_{\{\limsup_{t \rightarrow \tau^*} \|u(t)\|_{W^{1,\infty}} = \infty\}} \quad \mathbb{P} - \text{a.s.} \quad (2.7)$$

Remark 2.1. We remark here that $F_3(u)$ in (1.6) will disappear when $k = 1$. The proof for Theorem 2.1 combines the techniques as employed in the papers [3, 4, 5, 18, 20, 29, 58]. However, the Faedo-Galerkin method used e.g. in [29, 20] cannot be utilized directly since in (1.5), we do not have additional constraints like incompressibility, which guarantees the global existence of an approximate solution (see,

e.g. [22, 29]). Without this, we need to find a positive lower bound for the lifespan of the approximate solutions, which is generally not clear. Particularly, for our case, this difficulty will be overcome by constructing a suitable approximation scheme and establishing an appropriate blow-up criterium, which applies not only for u , but also for the approximate solution u_ε . This idea is transferred from the recent work [18] on deriving blow-up criteria.

The next result addresses the dependence of the solution on initial data giving at least a partial answer.

Theorem 2.2 (Weak instability). *Consider the periodic initial value problem (1.5), where $F(\cdot)$ is given by (1.6) with $k \geq 1$. Let $\mathcal{S} = (\Omega, \mathcal{F}, \mathbb{P}, \{\mathcal{F}_t\}_{t \geq 0}, \mathcal{W})$ be a fixed stochastic basis and let $s > 3/2$. If h satisfies Assumption A₂, then at least one of the following properties holds true.*

- (1) *For any $R \gg 1$, the R -exiting time is **not** strongly stable for the zero solution in the sense of Definition 2.2;*
- (2) *The solution map $u_0 \mapsto u$ defined by solving (1.5) is **not** uniformly continuous as a map from $L^2(\Omega, H^s)$ into $L^2(\Omega; C([0, T]; H^s))$ for any $T > 0$. More precisely, there exist two sequences of solutions $u_{1,n}(t)$ and $u_{2,n}(t)$, and two sequences of stopping times $\tau_{1,n}$ and $\tau_{2,n}$, such that*
 - $\mathbb{P}\{\tau_{i,n} > 0\} = 1$ for each $n > 1$ and $i = 1, 2$. Besides,

$$\lim_{n \rightarrow \infty} \tau_{1,n} = \lim_{n \rightarrow \infty} \tau_{2,n} = \infty \quad \mathbb{P} - a.s. \quad (2.8)$$

- For $i = 1, 2$, $u_{i,n} \in C([0, \tau_{i,n}]; H^s)$ $\mathbb{P} - a.s.$, and

$$\mathbb{E} \left(\sup_{t \in [0, \tau_{1,n}]} \|u_{1,n}(t)\|_{H^s}^2 + \sup_{t \in [0, \tau_{2,n}]} \|u_{2,n}(t)\|_{H^s}^2 \right) \lesssim 1. \quad (2.9)$$

- At $t = 0$,

$$\lim_{n \rightarrow \infty} \mathbb{E} \|u_{1,n}(0) - u_{2,n}(0)\|_{H^s}^2 = 0. \quad (2.10)$$

- For any $T > 0$,

$$\liminf_{n \rightarrow \infty} \mathbb{E} \sup_{t \in [0, T \wedge \tau_{1,n} \wedge \tau_{2,n}]} \|u_{1,n}(t) - u_{2,n}(t)\|_{H^s}^2 \gtrsim \begin{cases} \left(\sup_{t \in [0, T]} |\sin t| \right)^2, & \text{if } k \text{ is odd,} \\ \left(\sup_{t \in [0, T]} \left| \sin \frac{t}{2} \right| \right)^2, & \text{if } k \text{ is even.} \end{cases} \quad (2.11)$$

Remark 2.2. To prove Theorem 2.2, we assume that for some $R_0 \gg 1$, the R_0 -exiting time is strongly stable at the zero solution. Then we will construct an example to show that the solution map $u_0 \mapsto u$ defined by (1.5) is not uniformly continuous. This example involves the construction (for each $s > 3/2$) of two sequences of solutions which are converging at time zero but remain far apart at any later time. Therefore we will first construct two sequences of approximation solutions $u^{i,n}$ ($i \in \{1, 2\}$) such that the actual solutions $u_{i,n}$ starting from $u_{i,n}(0) = u^{i,n}(0)$ satisfy that as $n \rightarrow \infty$,

$$\lim_{n \rightarrow \infty} \mathbb{E} \sup_{[0, \tau_{i,n}]} \|u_{i,n} - u^{i,n}\|_{H^s} = 0, \quad (2.12)$$

where $u_{i,n}$ exists at least on $[0, \tau_{i,n}]$. If we obtain (2.12), then we can estimate the approximation solutions instead of the actual solutions and obtain (2.11). In deterministic case, for other works using the method of approximate solutions for studying dependence on initial data, we refer the reader to [47, 2, 34, 63, 61] and the references therein. However, in contrast to deterministic cases where lifespan estimate can be achieved (see (4.7)–(4.8) in [62] and (3.8)–(3.9) in [63] for example), it is not clear $\inf_n \tau_{i,n} > 0$ almost surely in stochastic setting. Therefore we are motivated to introduce the definition on the stability of the exiting time (see Definition 2.2). Then we find that the property $\inf_n \tau_{i,n} > 0$ can be connected with the stability property of the exiting time of the zero solution. Then we estimate the error in $H^{2s-\sigma}$ and H^σ with suitable σ . Different from [59], the problem (1.5) has nonlinearities of order $k+1$ and (2.12) should depend on k . Therefore more technical estimates are needed (see Lemma 4.1 and (4.17) for example). Finally we use interpolation to derive (2.12). Theorem 2.2 shows that one cannot expect too much for the issue of the dependence on initial data. More precisely, we cannot expect to improve the stability of the exiting time for the zero solution, and simultaneously to improve the continuous dependence of solutions on initial data.

Finally, we focus on (1.8). For the issue of global existence, we have

Theorem 2.3 (Global existence: Case 1). *Let $k \geq 1$ and $s > 3/2$. Let $b(t)$ satisfy Assumption A₃ and $\mathcal{S} = (\Omega, \mathcal{F}, \mathbb{P}, \{\mathcal{F}_t\}_{t \geq 0}, W)$ be a fixed stochastic basis. Assume u_0 to be an H^s -valued \mathcal{F}_0 -measurable random variable satisfying $\mathbb{E}\|u_0\|_{H^s}^2 < \infty$. Let K be a constant such that $\|\cdot\|_{W^{1,\infty}} < K\|\cdot\|_{H^s}$. Then there is a $C = C(s) > 1$ such that for any $R > 1$ and $\lambda_1 > 2$, if $\|u_0\|_{H^s} < \frac{1}{RK} \left(\frac{b_*}{C\lambda_1}\right)^{1/k}$ almost surely, then (1.8) has a maximal pathwise solution (u, τ^*) satisfying for any $\lambda_2 > \frac{2\lambda_1}{\lambda_1 - 2}$ the estimate*

$$\mathbb{P} \left\{ \|u(t)\|_{H^s} < \frac{1}{K} \left(\frac{b_*}{C\lambda_1}\right)^{1/k} e^{-\frac{((\lambda_1-2)\lambda_2-2\lambda_1)}{2\lambda_1\lambda_2} \int_0^t b^2(t') dt'} \text{ for all } t > 0 \right\} \geq 1 - \left(\frac{1}{R}\right)^{2/\lambda_2}.$$

Remark 2.3. Here we notice that if $k = 1$, then $F_3(u)$ in (1.6) will disappear. Motivated by the recent papers [29, 57, 58], where the linear noise $\beta u dW$ with $\beta \in \mathbb{R} \setminus \{0\}$ is considered, we focus on the non-autonomous linear multiplicative noise case, namely (1.8). We transform (1.8) to a non-autonomous random system (5.2). Although the stochastic integral is absent in (5.2), to extend the deterministic results to the stochastic setting, we need to overcome a few technical difficulties since the system is not only random but also non-autonomous. In this work, we manage to gain some estimates and asymptotic limits of the Girsanov type processes (see e.g., (5.6), (5.8), (5.10) and Lemma A.7), which enable us to apply the energy estimate pathwisely (namely for a.e. $\omega \in \Omega$) and obtain Theorem 2.3.

Theorem 2.4 (Global existence: Case 2). *Let $\mathcal{S} = (\Omega, \mathcal{F}, \mathbb{P}, \{\mathcal{F}_t\}_{t \geq 0}, W)$ be a fixed stochastic basis. Let $s > 3$ and $b(t)$ satisfy Assumption A₃. Assume u_0 to be an H^s -valued, \mathcal{F}_0 -measurable random variable satisfying $\mathbb{E}\|u_0\|_{H^s}^2 < \infty$. If u_0 satisfies*

$$\mathbb{P} \{ (1 - \partial_{xx}^2)u_0(x) > 0, \forall x \in \mathbb{T} \} = p, \quad \mathbb{P} \{ (1 - \partial_{xx}^2)u_0(x) < 0, \forall x \in \mathbb{T} \} = q,$$

for some $p, q \in [0, 1]$, then the corresponding maximal pathwise solution (u, τ^) to (1.8) satisfies*

$$\mathbb{P} \{ \tau^* = \infty \} \geq p + q.$$

That is to say, $\mathbb{P} \{ u \text{ exists globally} \} \geq p + q$.

3. PROOF FOR THEOREM 2.1

3.1. Blow-up criteria. Let us postpone the proof for existence and uniqueness of solutions to (1.5) to Section 3.2. We will first prove the blow-up criteria, since some estimates will be used later. Motivated by [18], we first consider the relationship between the blow-up time of $\|u(t)\|_{H^s}$ and the blow-up time of $\|u(t)\|_{W^{1,\infty}}$ for (1.5). Even though one might expect that the $\|u(t)\|_{H^s}$ norm may blow up earlier than $\|u(t)\|_{W^{1,\infty}}$, the following result shows that this is not true.

Lemma 3.1. *Let (u, τ^*) be the unique maximal pathwise solution to (1.5). Then the real-valued stochastic process $\|u\|_{W^{1,\infty}}$ is also \mathcal{F}_t -adapted. Besides, for any $m, n \in \mathbb{Z}^+$, define*

$$\tau_{1,m} = \inf \{ t \geq 0 : \|u(t)\|_{H^s} \geq m \}, \quad \tau_{2,n} = \inf \{ t \geq 0 : \|u(t)\|_{W^{1,\infty}} \geq n \}.$$

Moreover, let $\tau_1 = \lim_{m \rightarrow \infty} \tau_{1,m}$ and $\tau_2 = \lim_{n \rightarrow \infty} \tau_{2,n}$, then we have

$$\tau_1 = \tau_2 \quad \mathbb{P} - a.s.$$

As a corollary, $\mathbf{1}_{\{\lim_{t \rightarrow \tau^} \|u(t)\|_{W^{1,\infty}} = \infty\}} = \mathbf{1}_{\{\tau^* < \infty\}} \quad \mathbb{P} - a.s.$*

Proof. To begin with, we see that $u(\cdot \wedge \tau) \in C([0, \infty); H^s)$ means that for any $t \in [0, \tau]$,

$$[u(t)]^{-1}(Y) = [u(t)]^{-1}(H^s \cap Y), \quad \forall Y \in \mathcal{B}(W^{1,\infty}).$$

Therefore $u(t)$, as a $W^{1,\infty}$ -valued process, is also \mathcal{F}_t -adapted. We then infer from the embedding $H^s \hookrightarrow W^{1,\infty}$ for $s > 3/2$ that for some $M > 0$ and $m \in \mathbb{N}$,

$$\sup_{t \in [0, \tau_{1,m}]} \|u(t)\|_{W^{1,\infty}} \leq M \sup_{t \in [0, \tau_{1,m}]} \|u(t)\|_{H^s} \leq ([M] + 1)m,$$

where $[M]$ means the integer part of M . Therefore we have $\tau_{1,m} \leq \tau_{2,([M]+1)m} \leq \tau_2 \quad \mathbb{P} - a.s.$, which means that $\tau_1 \leq \tau_2 \quad \mathbb{P} - a.s.$ Now we only need to prove $\tau_2 \leq \tau_1 \quad \mathbb{P} - a.s.$ It is easy to see that for all $n_1, n_2 \in \mathbb{Z}^+$,

$$\left\{ \sup_{t \in [0, \tau_{2,n_1} \wedge n_2]} \|u(t)\|_{H^s} < \infty \right\} = \bigcup_{m \in \mathbb{Z}^+} \left\{ \sup_{t \in [0, \tau_{2,n_1} \wedge n_2]} \|u(t)\|_{H^s} < m \right\} \subset \bigcup_{m \in \mathbb{Z}^+} \{ \tau_{2,n_1} \wedge n_2 \leq \tau_{1,m} \}.$$

Since

$$\bigcup_{m \in \mathbb{Z}^+} \{\tau_{2,n_1} \wedge n_2 \leq \tau_{1,m}\} \subset \{\tau_{2,n_1} \wedge n_2 \leq \tau_1\},$$

we see that if

$$\mathbb{P} \left\{ \sup_{t \in [0, \tau_{2,n_1} \wedge n_2]} \|u(t)\|_{H^s} < \infty \right\} = 1, \quad \forall n_1, n_2 \in \mathbb{Z}^+ \quad (3.1)$$

holds true, then for all $n_1, n_2 \in \mathbb{Z}^+$, $\mathbb{P} \{\tau_{2,n_1} \wedge n_2 \leq \tau_1\} = 1$ and

$$\mathbb{P} \{\tau_2 \leq \tau_1\} = \mathbb{P} \left\{ \bigcap_{n_1 \in \mathbb{Z}^+} \{\tau_{2,n_1} \leq \tau_1\} \right\} = \mathbb{P} \left\{ \bigcap_{n_1, n_2 \in \mathbb{Z}^+} \{\tau_{2,n_1} \wedge n_2 \leq \tau_1\} \right\} = 1. \quad (3.2)$$

Since (3.2) requires the assumption (3.1), it suffices to prove (3.1). However, we can not directly apply the Itô formula for $\|u\|_{H^s}^2$ to get control of $\mathbb{E}\|u(t)\|_{H^s}^2$ since we only have $u \in H^s$ and $u^k u_x \in H^{s-1}$. Therefore the well known Itô formula in general Hilbert space can not be used directly, see [19, Theorem 4.32] or [25, Theorem 2.10] for example. We will use the mollifier operator T_ε defined in the appendix to overcome this obstacle. Indeed, applying T_ε to (1.5) and using the Itô formula for $\|T_\varepsilon u\|_{H^s}^2$, we have that for any $t > 0$,

$$\begin{aligned} d\|T_\varepsilon u(t)\|_{H^s}^2 &= (T_\varepsilon h(t, u) dW, T_\varepsilon u)_{H^s} - 2(D^s T_\varepsilon [u^k u], D^s T_\varepsilon u)_{L^2} dt \\ &\quad - 2(D^s T_\varepsilon F(u), D^s T_\varepsilon u)_{L^2} dt + \|T_\varepsilon h(t, u)\|_{L^2(U; H^s)}^2 dt. \end{aligned}$$

Therefore for any $n_1, n_2 \geq 1$ and $t \in [0, \tau_{2,n_1} \wedge n_2]$,

$$\begin{aligned} \|T_\varepsilon u(t)\|_{H^s}^2 - \|T_\varepsilon u(0)\|_{H^s}^2 &= 2 \sum_{j=1}^{\infty} \int_0^t (D^s T_\varepsilon h(t, u) e_j, D^s T_\varepsilon u)_{L^2} dW_j \\ &\quad - 2 \int_0^t (D^s T_\varepsilon [u^k \partial_x u], D^s T_\varepsilon u)_{L^2} dt \\ &\quad - 2 \int_0^t (D^s T_\varepsilon F(u), D^s T_\varepsilon u)_{L^2} dt \\ &\quad + \int_0^t \sum_{k=1}^{\infty} \|D^s T_\varepsilon h(t, u) e_j\|_{L^2}^2 dt \\ &= \int_0^t \sum_{j=1}^{\infty} L_{1,j} dW_j + \sum_{i=2}^4 \int_0^t L_i dt, \end{aligned}$$

where $\{e_k\}$ is the complete orthonormal basis of U . On account of the Burkholder-Davis-Gundy inequality, we arrive at

$$\mathbb{E} \sup_{t \in [0, \tau_{2,n_1} \wedge n_2]} \|T_\varepsilon u(t)\|_{H^s}^2 \leq \mathbb{E} \|T_\varepsilon u_0\|_{H^s}^2 + C \mathbb{E} \left(\int_0^{\tau_{2,n_1} \wedge n_2} \left| \sum_{j=1}^{\infty} L_{1,j} \right|^2 dt \right)^{\frac{1}{2}} + \sum_{i=2}^4 \mathbb{E} \int_0^{\tau_{2,n_1} \wedge n_2} |L_i| dt.$$

Then (2.3), (A.3) and the stochastic Fubini theorem [19] lead to

$$\mathbb{E} \left(\int_0^{\tau_{2,n_1} \wedge n_2} \left| \sum_{j=1}^{\infty} L_{1,j} \right|^2 dt \right)^{\frac{1}{2}} \leq \frac{1}{2} \mathbb{E} \sup_{t \in [0, \tau_{2,n_1} \wedge n_2]} \|T_\varepsilon u\|_{H^s}^2 + C f^2(n_1) \int_0^{n_2} (1 + \mathbb{E} \|u\|_{H^s}^2) dt.$$

For L_2 , we notice that T_ε satisfies (A.1), (A.2) and (A.3). Then it follows from Lemmas A.1, A.2 and A.3, integration by parts and $H^s \hookrightarrow W^{1,\infty}$ that

$$\begin{aligned} &(D^s T_\varepsilon [u^k u_x], D^s T_\varepsilon u)_{L^2} \\ &= ([D^s, u^k] u_x, D^s T_\varepsilon^2 u)_{L^2} + ([T_\varepsilon, u^k] D^s u_x, D^s T_\varepsilon u)_{L^2} + (u^k D^s T_\varepsilon u_x, D^s T_\varepsilon u)_{L^2} \\ &\leq C \|u\|_{W^{1,\infty}}^k \|u\|_{H^s}^2, \end{aligned}$$

which implies

$$\mathbb{E} \int_0^{\tau_2, n_1 \wedge n_2} |L_2| \, dt \leq C n_1^k \int_0^{n_2} (1 + \mathbb{E} \|u\|_{H^s}^2) \, dt.$$

Similarly, it follows from Lemma A.5 and the assumption (2.3) that

$$\mathbb{E} \int_0^{\tau_2, n_1 \wedge n_2} |L_3| + |L_4| \, dt \leq C(n_1^k + f^2(n_1)) \int_0^{n_2} (1 + \mathbb{E} \|u\|_{H^s}^2) \, dt.$$

Therefore we combine the above estimates with using (A.3) to have

$$\mathbb{E} \sup_{t \in [0, \tau_2, n_1 \wedge n_2]} \|T_\varepsilon u(t)\|_{H^s}^2 \leq C \mathbb{E} \|u_0\|_{H^s}^2 + C \int_0^{n_2} \left(1 + \mathbb{E} \sup_{t' \in [0, t \wedge \tau_2, n_1]} \|u(t')\|_{H^s}^2 \right) dt,$$

where $C = C(n_1, k)$ through n_1^k and $n_1^k + f^2(n_1)$. Since the right hand side of the above estimate does not depend on ε , and $T_\varepsilon u$ tends to u in $C([0, T], H^s)$ for any $T > 0$ almost surely as $\varepsilon \rightarrow 0$, one can send $\varepsilon \rightarrow 0$ to obtain

$$\mathbb{E} \sup_{t \in [0, \tau_2, n_1 \wedge n_2]} \|u(t)\|_{H^s}^2 \leq C \mathbb{E} \|u_0\|_{H^s}^2 + C \int_0^{n_2} \left(1 + \mathbb{E} \sup_{t' \in [0, t \wedge \tau_2, n_1]} \|u(t')\|_{H^s}^2 \right) dt. \quad (3.3)$$

Then Gronwall's inequality shows that for each $n_1, n_2 \in \mathbb{Z}^+$, there is a constant $C = C(n_1, n_2, u_0, k) > 0$ such that

$$\mathbb{E} \sup_{t \in [0, \tau_2, n_1 \wedge n_2]} \|u(t)\|_{H^s}^2 < C(n_1, n_2, u_0, k),$$

which gives (3.1). \square

Proof for (2.7). Let us assume the existence and uniqueness first. We notice that for fixed $m, n > 0$, even if $\mathbb{P}\{\tau_{1,m} = 0\}$ or $\mathbb{P}\{\tau_{2,n} = 0\}$ is larger than 0, for a.e. $\omega \in \Omega$, there is $m > 0$ or $n > 0$ such that $\tau_{1,m}, \tau_{2,n} > 0$. By continuity of $\|u(t)\|_{H^s}$ and the uniqueness of u , it is easy to check that $\tau_1 = \tau_2$ is actually the maximal existence time τ^* of u in the sense of Definition 2.1. Consequently, we obtain the desired blow-up criteria. \square

3.2. Sketch of the proof for Theorem 2.1. Since the main idea of proving Theorem 2.1 follows standard ideas and it is very similar to [58, 59], here we only give a sketch of the main steps.

- (1) (Approximate solutions) The first step is to construct a suitable approximation scheme. For any $R > 1$, we let $\chi_R(x) : [0, \infty) \rightarrow [0, 1]$ be a C^∞ function such that $\chi_R(x) = 1$ for $x \in [0, R]$ and $\chi_R(x) = 0$ for $x > 2R$. Then we consider the following cut-off problem on \mathbb{T} ,

$$\begin{cases} du + \chi_R(\|u\|_{W^{1,\infty}}) [u^k \partial_x u + F(u)] \, dt = \chi_R(\|u\|_{W^{1,\infty}}) h(t, u) dW, \\ u(\omega, 0, x) = u_0(\omega, x) \in H^s. \end{cases} \quad (3.4)$$

From Lemma A.5, we see that the nonlinear term $F(u)$ preserves the H^s -regularity of $u \in H^s$ for any $s > 3/2$. However, to apply the theory of SDEs in Hilbert space to (3.4), we will have to mollify the transport term $u^k \partial_x u$ since the product $u^k \partial_x u$ loses one regularity. To this end, we consider the following approximation scheme:

$$\begin{cases} du + G_{1,\varepsilon}(u) dt = G_2(u) dW, & x \in \mathbb{T}, \, t > 0, \\ G_{1,\varepsilon}(u) = \chi_R(\|u\|_{W^{1,\infty}}) [J_\varepsilon((J_\varepsilon u)^k \partial_x J_\varepsilon u) + F(u)], \\ G_2(u) = \chi_R(\|u\|_{W^{1,\infty}}) h(t, u), \\ u(0, x) = u_0(x) \in H^s(\mathbb{T}), \end{cases} \quad (3.5)$$

where J_ε is the Friedrichs mollifier defined in Appendix A. After mollifying the transport term $u^k u_x$, for a fixed stochastic basis $\mathcal{S} = (\Omega, \mathcal{F}, \mathbb{P}, \{\mathcal{F}_t\}_{t \geq 0}, W)$ and for $u_0 \in L^2(\Omega; H^s)$ with $s > 3$, according to the existence theory of SDE in Hilbert space (see for example [56, Theorem 4.2.4 with Example 4.1.3] and [40]), (3.5) admits a unique solution $u_\varepsilon \in C([0, T_\varepsilon], H^s) \, \mathbb{P} - a.s.$ Moreover, the uniform $L^\infty(\Omega; W^{1,\infty})$ condition provided by the cut-off function χ_R enables us to split the expectation $\mathbb{E}(\|u_\varepsilon\|_{H^s}^2 \|u_\varepsilon\|_{W^{1,\infty}})$ to close the *a priori* $L^2(\Omega; H^s)$ estimate for u_ε . Then we can go along the lines as we prove Lemma 3.1 to find that for each fixed ε , if $T_\varepsilon < \infty$, then $\limsup_{t \rightarrow T_\varepsilon} \|u_\varepsilon(t)\|_{W^{1,\infty}} = \infty$. Due to the cut-off in (3.5), for a.e. $\omega \in \Omega$, $\|u_\varepsilon(t)\|_{W^{1,\infty}}$ is always bounded and hence u_ε is actually a global in time solution, that is, $u_\varepsilon \in C([0, \infty), H^s) \, \mathbb{P} - a.s.$

We remark here that the global existence of u_ε is necessary in our framework due to the lack of lifespan estimate in the stochastic setting. Otherwise we will have to prove $\mathbb{P}\{\inf_{\varepsilon>0} T_\varepsilon > 0\} = 1$, which is not clear in general.

- (2) (Pathwise solution to the cut-off problem in H^s with $s > 3$) We pass to the limit $\varepsilon \rightarrow 0$. By applying the stochastic compactness arguments from Prokhorov's and Skorokhod's theorems, we obtain the almost sure convergence for a new approximate solution $(\widetilde{u}_\varepsilon, \widetilde{W}_\varepsilon)$ defined on a *new* probability space. By virtue of a refined martingale representation theorem [36, Theorem A.1], we may send $\varepsilon \rightarrow 0$ in $(\widetilde{u}_\varepsilon, \widetilde{W}_\varepsilon)$ to build a global martingale solution in H^s with $s > 3$ to the cut-off problem. Finally, since F satisfies the estimates as in Lemma A.5 and h satisfies Assumption A₁, one can obtain the pathwise uniqueness easily. Then the Gyöngy–Krylov characterization [30] of the convergence in probability can be applied here to prove the convergence of the original approximate solutions. For more details, we refer to [58, 59].
- (3) (Remove the cut-off and extend the range of s to $s > 3/2$) When $u_0 \in L^\infty(\Omega, H^s)$ with $s > 3/2$, by mollifying the initial data, we obtain a sequence of regular solutions $\{u_n\}_{n \in \mathbb{N}}$ to (1.5). Motivated by [28, 58], one can prove that there is a subsequence (still denoted by u_n) such that for some almost surely positive stopping time τ ,

$$\lim_{n \rightarrow \infty} \sup_{t \in [0, \tau]} \|u_n - u\|_{H^s} = 0 \quad \mathbb{P} - a.s.,$$

and

$$\sup_{t \in [0, \tau]} \|u\|_{H^s} \leq \|u_0\|_{H^s} + 2 \quad \mathbb{P} - a.s. \quad (3.6)$$

Then we can pass the limit $n \rightarrow \infty$ to prove that (u, τ) is a solution to (1.5). Besides, using a cutting argument, as in [29, 28, 3], enables us to remove the $L^\infty(\Omega)$ assumption on u_0 . More precisely, when $\mathbb{E}\|u_0\|_{H^s}^2 < \infty$, we consider the decomposition

$$\Omega_m = \{m-1 \leq \|u_0\|_{H^s} < m\}, \quad m \geq 1.$$

Since $\mathbb{E}\|u_0\|_{H^s}^2 < \infty$, $\bigcup_{m \geq 1} \Omega_m$ is a set of full measure and $1 = \sum_{m \geq 1} \mathbf{1}_{\Omega_m} \quad \mathbb{P} - a.s.$ Therefore we have

$$u_0(\omega, x) = \sum_{m \geq 1} u_{0,m}(\omega, x) = \sum_{m \geq 1} u_0(\omega, x) \mathbf{1}_{\Omega_m} \quad \mathbb{P} - a.s.$$

For each initial value $u_{0,m}$, we let (u_m, τ_m) be the pathwise unique solution to (1.5) satisfying (3.6). Moreover, as $F(0) = 0$ and $h(0) = 0$ (cf. (2.3)), direct computation shows that

$$\left(u = \sum_{m \geq 1} u_m \mathbf{1}_{m-1 \leq \|u_0\|_{H^s} < m}, \quad \tau = \sum_{m \geq 1} \tau_m \mathbf{1}_{m-1 \leq \|u_0\|_{H^s} < m} \right)$$

is the unique pathwise solution to (1.5) corresponding to the initial condition u_0 . Since (u_m, τ_m) satisfies (3.6), we have $u(\cdot \wedge \tau) \in C([0, \infty); H^s) \quad \mathbb{P} - a.s.$ and

$$\begin{aligned} \sup_{t \in [0, \tau]} \|u\|_{H^s}^2 &= \sum_{m=1}^{\infty} \mathbf{1}_{m-1 \leq \|u_0\|_{H^s} < m} \sup_{t \in [0, \tau_m]} \|u_m\|_{H^s}^2 \\ &\leq C \sum_{m=1}^{\infty} \mathbf{1}_{m-1 \leq \|u_0\|_{H^s} < m} \left(4 + \|u_{0,m}\|_{H^s}^2 \right) = C \left(4 + \|u_0\|_{H^s}^2 \right). \end{aligned}$$

Taking expectation in the above inequality, we obtain (2.6). Since the passage from (u, τ) to a unique maximal pathwise solution (u, τ^*) in the sense of Definition 2.1 can be carried out as in [18, 29, 28, 57], we omit the details. The proof for Theorem 2.1 is finished.

Remark 3.1. Changing the growth assumption (2.3) to $\|h(t, u)\|_{L_2(U; H^s)} \leq f(\|u\|_{H^{s'}})(1 + \|u\|_{H^s})$ with $H^s \hookrightarrow H^{s'} \hookrightarrow W^{1, \infty}$ leads to a criticality statement. One can then go along the lines in Lemma 3.1 to find $\mathbf{1}_{\{\tau^* < \infty\}} = \mathbf{1}_{\{\limsup_{t \rightarrow \tau^*} \|u(t)\|_{H^{s'}} = \infty\}} \quad \mathbb{P} - a.s.$ Using another cut-off $\chi_R(\|u\|_{H^{s'}})$ in the approximate scheme (3.4) and in (3.5), the other part of the proof also allows us to establish a local existence and uniqueness result. The difference condition (2.4) is essential to guarantee pathwise uniqueness.

4. PROOF FOR THEOREM 2.2

Now we are going to prove Theorem 2.2. To this end, it suffices to show that if for some $R_0 \gg 1$, the R_0 -exiting time is strongly stable at the zero solution, then the solution map $u_0 \mapsto u$ is not uniformly continuous. We restrict our attention to $k \geq 2$ since the case $k = 1$, i.e., the stochastic CH equation, has been obtained in [59].

4.1. Approximate solutions and associated estimates. Define the approximate solutions as

$$u^{l,n} = ln^{-\frac{1}{k}} + n^{-s} \cos \theta \text{ and } \theta = nx - lt, \quad n \in \mathbb{Z}^+,$$

where

$$l \in \{-1, 1\} \text{ if } k \text{ is odd; } l \in \{0, 1\} \text{ if } k \text{ is even.} \quad (4.1)$$

Substituting $u^{l,n}$ into (1.5), we see that the error $E^{l,n}(t)$ can be defined as

$$E^{l,n}(t) = u^{l,n}(t) - u^{l,n}(0) + \int_0^t \left[(u^{l,n})^k \partial_x u^{l,n} + F(u^{l,n}) \right] dt' - \int_0^t h(t', u^{l,n}) dW. \quad (4.2)$$

Now we analyze the error as follows.

Lemma 4.1. *Let $s > 3/2$. For $n \gg 1$, $\delta \in (1/2, \min\{s-1, 3/2\})$ and any $T > 0$, there is a $C = C(T) > 0$ such that*

$$\mathbb{E} \sup_{t \in [0, T]} \|E^{l,n}(t)\|_{H^\delta}^2 \leq Cn^{-2r_s}.$$

Here r_s is a parameter with

$$0 < r_s = \begin{cases} 2s - \delta - \frac{k+1}{k}, & \text{if } \frac{3}{2} < s \leq \frac{2k+1}{k}, \\ s - \delta + 1, & \text{if } s > \frac{2k+1}{k}. \end{cases} \quad (4.3)$$

Proof. Since (4.1) implies $l^k = l$. This means

$$\begin{aligned} & u^{l,n}(t) - u^{l,n}(0) + \int_0^t (u^{l,n})^k \partial_x u^{l,n} dt' \\ &= u^{l,n}(t) - u^{l,n}(0) + \int_0^t \left(\sum_{j=0}^k \binom{k}{j} \left(ln^{-\frac{1}{k}} \right)^{k-j} n^{-sj} \cos^j \theta \right) (-n^{-s+1} \sin \theta) dt' \\ &= \int_0^t \left(\sum_{j=1}^k \binom{k}{j} \left(ln^{-\frac{1}{k}} \right)^{k-j} n^{-sj} \cos^j \theta \right) (-n^{-s+1} \sin \theta) dt' \triangleq \int_0^t T_{n,k} dt'. \end{aligned}$$

Then we have

$$E^{l,n}(t) - \int_0^t [T_{n,k} + F(u^{l,n})] dt' + \int_0^t h(t', u^{l,n}) dW = 0. \quad (4.4)$$

We first notice that

$$\begin{aligned} \|T_{n,k}\|_{H^\delta} &\lesssim \sum_{j=1}^k n^{\frac{j}{k}-sj-s} \|\cos^j \theta \sin \theta\|_{H^\delta} \\ &\lesssim \sum_{j=1}^k n^{\frac{j}{k}-sj-s-1} \|\cos^{j+1} \theta\|_{H^{\delta+1}} \\ &\lesssim \max_{1 \leq j \leq k} \{n^{\frac{j}{k}-sj-s+\delta}\} \lesssim n^{\frac{1}{k}-2s+\delta} \lesssim n^{1-2s+\delta}. \end{aligned} \quad (4.5)$$

Recall that $F(\cdot)$ is given by (1.6). Since $(1 - \partial_{xx}^2)^{-1}$ is bounded from H^δ to $H^{\delta+2}$, we can use Lemmas A.6 and A.3 to estimate $\|F_i(u^{l,n})\|_{H^\delta}$ ($i = 1, 2, 3$) as follows.

$$\begin{aligned}
\|F_1(u^{l,n})\|_{H^\delta} &\lesssim \sum_{j=0}^k n^{\frac{j}{k}-sj-s} \|\cos^j \theta \sin \theta\|_{H^{\delta-2}} \\
&\lesssim \sum_{j=0}^k n^{\frac{j}{k}-sj-s-1} \|\cos^{j+1} \theta\|_{H^{\delta-1}} \\
&\lesssim \sum_{j=0}^k n^{\frac{j}{k}-sj-s-1} \|\cos^{j+1} \theta\|_{H^\delta} \\
&\lesssim \max_{0 \leq j \leq k} \{n^{\frac{j}{k}-sj-s-1+\delta}\} \lesssim n^{-s+\delta-1}.
\end{aligned} \tag{4.6}$$

$$\begin{aligned}
\|F_2(u^{l,n})\|_{H^\delta} &\lesssim \sum_{j=0}^{k-1} n^{\frac{j+1}{k}-sj-2s+1} \|\cos^j \theta \sin^2 \theta\|_{H^{\delta-1}} \\
&\lesssim \sum_{j=0}^{k-1} n^{\frac{j+1}{k}-sj-2s+1} \|\cos^j \theta \sin^2 \theta\|_{H^\delta} \\
&\lesssim \max_{0 \leq j \leq k-1} \{n^{\frac{j+1}{k}-sj-2s+\delta+1}\} \lesssim n^{-2s+\delta+1+\frac{1}{k}}.
\end{aligned} \tag{4.7}$$

$$\begin{aligned}
\|F_3(u^{l,n})\|_{H^\delta} &\lesssim \sum_{j=0}^{k-2} n^{\frac{j+2}{k}-sj-3s+2} \|\cos^j \theta \sin^3 \theta\|_{H^{\delta-2}} \\
&\lesssim \sum_{j=0}^{k-2} n^{\frac{j+2}{k}-sj-3s+2} \|\cos^j \theta \sin^3 \theta\|_{H^\delta} \\
&\lesssim \max_{0 \leq j \leq k-2} \{n^{\frac{j+2}{k}-sj-3s+\delta+2}\} \lesssim n^{-3s+\delta+\frac{2}{k}+2}.
\end{aligned} \tag{4.8}$$

In the above estimates, we used the fact that $F_3(\cdot)$ appears only for $k \geq 2$. Combining this fact, (4.6), (4.7) and (4.8), we have

$$\|F(u^{l,n})\|_{H^\delta} \lesssim \max \left\{ n^{1-2s+\delta}, n^{-3s+\delta+\frac{2}{k}+2}, n^{-2s+\delta+\frac{1}{k}+1}, n^{-s+\delta-1} \right\} \lesssim n^{-r_s}. \tag{4.9}$$

Then, for any $T > 0$ and $t \in [0, T]$, by virtue of the Itô formula, we arrive at

$$\|E^{l,n}(t)\|_{H^\delta}^2 \leq \left| \left(-2 \int_0^t h(t', u^{l,n}) d\mathcal{W}, E^{l,n} \right)_{H^\delta} \right| + \sum_{i=2}^4 \int_0^t |J_i| dt',$$

where

$$\begin{aligned}
J_2 &= 2 (D^\delta T_{n,k}, D^\delta E^{l,n})_{L^2}, \\
J_3 &= 2 (D^\delta F(u^{l,n}), D^\delta E^{l,n})_{L^2}, \\
J_4 &= \|h(t', u^{l,n})\|_{L_2(U, H^\delta)}^2.
\end{aligned}$$

Taking the supremum with respect to $t \in [0, T]$, using the Burkholder-Davis-Gundy inequality and using (2.5) and (4.9) yield

$$\begin{aligned}
\mathbb{E} \sup_{t \in [0, T]} \left| \left(-2 \int_0^t h(t', u^{l,n}) d\mathcal{W}, E^{l,n} \right)_{H^\delta} \right| &\leq 2 \mathbb{E} \left(\int_0^T \|E^{l,n}(t)\|_{H^\delta}^2 \|F(u^{l,n})\|_{H^\delta}^2 dt \right)^{\frac{1}{2}} \\
&\leq 2 \mathbb{E} \left(\sup_{t \in [0, T]} \|E^{l,n}(t)\|_{H^\delta}^2 \int_0^T \|F(u^{l,n})\|_{H^\delta}^2 dt \right)^{\frac{1}{2}} \\
&\leq \frac{1}{2} \mathbb{E} \sup_{t \in [0, T]} \|E^{l,n}(t)\|_{H^\delta}^2 + CT n^{-2r_s}.
\end{aligned}$$

By virtue of (4.5) and (4.9), we obtain

$$\begin{aligned}
\int_0^T \mathbb{E}|J_2|dt &\leq C \int_0^T \mathbb{E} (\|T_{n,k}\|_{H^\delta} \|E^{l,n}(t)\|_{H^\delta}) dt \\
&\leq C \int_0^T \mathbb{E} \|T_{n,k}\|_{H^\delta}^2 dt + C \int_0^T \mathbb{E} \|E^{l,n}(t)\|_{H^\delta}^2 dt \\
&\leq CTn^{-2r_s} + C \int_0^T \mathbb{E} \|E^{l,n}(t)\|_{H^\delta}^2 dt, \\
\int_0^T \mathbb{E}|J_3|dt &\leq C \int_0^T \mathbb{E} (\|F(u^{l,n})\|_{H^\delta} \|E^{l,n}(t)\|_{H^\delta}) dt \\
&\leq C \int_0^T \mathbb{E} \|F(u^{l,n})\|_{H^\delta}^2 dt + C \int_0^T \mathbb{E} \|E^{l,n}(t)\|_{H^\delta}^2 dt \\
&\leq CTn^{-2r_s} + C \int_0^T \mathbb{E} \|E^{l,n}(t)\|_{H^\delta}^2 dt,
\end{aligned}$$

and by (2.5),

$$\int_0^T \mathbb{E}|J_4|dt \leq C \int_0^T \mathbb{E} \|F(u^{l,n})\|_{H^\delta}^2 dt \leq CTn^{-2r_s}.$$

Collecting the above estimates, we arrive at

$$\mathbb{E} \sup_{t \in [0, T]} \|E^{l,n}(t)\|_{H^\delta}^2 \leq CTn^{-2r_s} + C \int_0^T \mathbb{E} \sup_{t' \in [0, t]} \|E^{l,n}(t')\|_{H^\delta}^2 dt.$$

Obviously, for each $n \geq 1$ and $l \in \{-1, 1\}$, $\mathbb{E} \sup_{t' \in [0, t]} \|E^{l,n}(t')\|_{H^\delta}^2$ is finite. Then we infer from the Grönwall inequality that

$$\mathbb{E} \sup_{t \in [0, T]} \|E^{l,n}(t)\|_{H^\delta}^2 \leq Cn^{-2r_s}, \quad C = C(T).$$

This is the desired result. \square

4.2. Construction of actual solutions. Now we consider the following periodic boundary value problem with deterministic initial data $u^{l,n}(0, x)$, i.e.,

$$\begin{cases} du + [u^k u_x + F(u)] dt = h(t, u) dW, & t > 0, x \in \mathbb{T}, \\ u(0, x) = u^{l,n}(0, x), & x \in \mathbb{T}. \end{cases} \quad (4.10)$$

Since h satisfies (2.5), we see that (2.3) and (2.4) are also verified. Then Theorem 2.1 yields that for each $n \in \mathbb{N}$, (4.10) has a uniqueness maximal pathwise solution $(u_{l,n}, \tau_{l,n}^*)$.

4.3. Estimates on the errors.

Lemma 4.2. *Let $s > \frac{3}{2}$, $\frac{1}{2} < \delta < \min\{s-1, \frac{3}{2}\}$ and $r_s > 0$ be given as in Lemma 4.1. For any $R > 1$ and l satisfying (4.1), we define*

$$\tau_{l,n}^R := \inf \{t \geq 0 : \|u_{l,n}\|_{H^s} > R\}. \quad (4.11)$$

Then for any $T > 0$, when $n \gg 1$, we have that for l satisfying (4.1),

$$\mathbb{E} \sup_{t \in [0, T \wedge \tau_{l,n}^R]} \|u^{l,n} - u_{l,n}\|_{H^\delta}^2 \leq Cn^{-2r_s}, \quad C = C(R, T), \quad (4.12)$$

and

$$\mathbb{E} \sup_{t \in [0, T \wedge \tau_{l,n}^R]} \|u^{l,n} - u_{l,n}\|_{H^{2s-\delta}}^2 \leq Cn^{2s-2\sigma}, \quad C = C(R, T). \quad (4.13)$$

Proof. We first notice that by Lemma A.6, for l satisfying (4.1),

$$\|u^{l,n}(t)\|_{H^s} \lesssim 1, \quad \forall t > 0. \quad (4.14)$$

Let $q = q^{l,n} = \sum_{i=0}^k (u^{l,n})^{k-i} (u_{l,n})^i$ and $v = v^{l,n} = u^{l,n} - u_{l,n}$. In view of (4.2), (4.4) and (4.10), we see that

$$v(t) + \int_0^t \left[\frac{1}{k+1} \partial_x(qv) - F(u_{l,n}) \right] dt' = - \int_0^t h(t', u_{l,n}) d\mathcal{W} + \int_0^t T_{n,k} dt'.$$

For any $T > 0$, we use the Itô formula on $[0, T \wedge \tau_{l,n}^R]$, take the supremum over $t \in [0, T \wedge \tau_{l,n}^R]$ and use the Burkholder-Davis-Gundy inequality with noticing (2.5) to find

$$\mathbb{E} \sup_{t \in [0, T \wedge \tau_{l,n}^R]} \|v(t)\|_{H^\delta}^2 \leq C \mathbb{E} \left(\int_0^{T \wedge \tau_{l,n}^R} |K_1|^2 dt \right)^{\frac{1}{2}} + \sum_{i=2}^5 \mathbb{E} \int_0^{T \wedge \tau_{l,n}^R} |K_i| dt,$$

where

$$\begin{aligned} K_1 &= \|v\|_{H^\delta} \|F(u_{l,n})\|_{H^\delta}, \\ K_2 &= 2 (D^\delta T_{n,k}, D^\delta v)_{L^2}, \\ K_3 &= -\frac{2}{k+1} (D^\delta \partial_x[qv], D^\delta v)_{L^2}, \\ K_4 &= 2 (D^\delta F(u_{l,n}), D^\delta v)_{L^2}, \\ K_5 &= \|h(t, u_{l,n})\|_{L^2(U, H^\delta)}^2. \end{aligned}$$

We can first infer from Lemma A.5 that

$$\begin{aligned} \|F(u_{l,n})\|_{H^\delta}^2 &\lesssim (\|F(u^{l,n}) - F(u_{l,n})\|_{H^\delta} + \|F(u^{l,n})\|_{H^\delta})^2 \\ &\lesssim (\|u^{l,n}\|_{H^s} + \|u_{l,n}\|_{H^s})^2 \|v\|_{H^\delta}^2 + \|F(u^{l,n})\|_{H^\delta}^2. \end{aligned}$$

Therefore, applying Lemmas A.4 and A.5, $H^\delta \hookrightarrow L^\infty$, integrating by part and (4.5), we have

$$|K_1|^2 \lesssim (\|u^{l,n}\|_{H^s} + \|u_{l,n}\|_{H^s})^2 \|v\|_{H^\delta}^4 + \|F(u^{l,n})\|_{H^\delta}^2 \|v\|_{H^\delta}^2,$$

$$|K_2| \lesssim \|T_{n,k}\|_{H^\delta}^2 + \|v\|_{H^\delta}^2 \lesssim n^{-2r_s} + \|v\|_{H^\delta}^2,$$

$$|K_3| \lesssim \|q\|_{H^s} \|v\|_{H^\delta}^2 + \|q_x\|_{L^\infty} \|v\|_{H^\delta}^2 \lesssim \|q\|_{H^s} \|v\|_{H^\delta}^2,$$

$$|K_4| \lesssim (\|u^{l,n}\|_{H^s} + \|u_{l,n}\|_{H^s}) \|v\|_{H^\delta}^2 + \|F(u^{l,n})\|_{H^\delta}^2 + \|v\|_{H^\delta}^2,$$

and

$$|K_5| \lesssim (\|u^{l,n}\|_{H^s} + \|u_{l,n}\|_{H^s})^2 \|v\|_{H^\delta}^2 + \|F(u^{l,n})\|_{H^\delta}^2.$$

By virtue of Lemma A.5, (4.9), (4.11) and (4.14), we have

$$\begin{aligned} & C \mathbb{E} \left(\int_0^{T \wedge \tau_{l,n}^R} |K_1|^2 dt \right)^{\frac{1}{2}} \\ & \leq C \mathbb{E} \left(\sup_{t \in [0, T \wedge \tau_{l,n}^R]} \|v\|_{H^\delta}^2 \int_0^{T \wedge \tau_{l,n}^R} (\|u^{l,n}\|_{H^s} + \|u_{l,n}\|_{H^s})^2 \|v\|_{H^\delta}^2 dt \right)^{\frac{1}{2}} \\ & \quad + C \mathbb{E} \left(\sup_{t \in [0, T \wedge \tau_{l,n}^R]} \|v\|_{H^\delta}^2 \int_0^{T \wedge \tau_{l,n}^R} \|F(u^{l,n})\|_{H^\delta}^2 dt \right)^{\frac{1}{2}} \\ & \leq \frac{1}{2} \mathbb{E} \sup_{t \in [0, T \wedge \tau_{l,n}^R]} \|v\|_{H^\delta}^2 + C_R \mathbb{E} \int_0^{T \wedge \tau_{l,n}^R} \|v(t)\|_{H^\delta}^2 dt + C \mathbb{E} \int_0^{T \wedge \tau_{l,n}^R} \|F(u^{l,n})\|_{H^\delta}^2 dt \\ & \leq \frac{1}{2} \mathbb{E} \sup_{t \in [0, T \wedge \tau_{l,n}^R]} \|v\|_{H^\delta}^2 + C_R \mathbb{E} \int_0^T \sup_{t' \in [0, t \wedge \tau_{l,n}^R]} \|v(t')\|_{H^\delta}^2 dt + CTn^{-2r_s}, \\ & \mathbb{E} \int_0^{T \wedge \tau_{l,n}^R} |K_2| + |K_4| + |K_5| dt \leq CTn^{-2r_s} + C_R \int_0^T \mathbb{E} \sup_{t' \in [0, t \wedge \tau_{l,n}^R]} \|v(t')\|_{H^\delta}^2 dt, \end{aligned}$$

and

$$\mathbb{E} \int_0^{T \wedge \tau_{l,n}^R} |K_3| \, dt \leq C_R \int_0^T \mathbb{E} \sup_{t' \in [0, t \wedge \tau_{l,n}^R]} \|v(t')\|_{H^\delta}^2 \, dt.$$

Consequently,

$$\mathbb{E} \sup_{t \in [0, T \wedge \tau_{l,n}^R]} \|v(t)\|_{H^\delta}^2 \leq CTn^{-2r_s} + C_R \int_0^T \mathbb{E} \sup_{t' \in [0, t \wedge \tau_{l,n}^R]} \|v(t')\|_{H^\delta}^2 \, dt.$$

Using the Grönwall inequality, we have

$$\mathbb{E} \sup_{t \in [0, T \wedge \tau_{l,n}^R]} \|v(t)\|_{H^\delta}^2 \leq Cn^{-2r_s}, \quad C = C(R, T),$$

which is (4.12). For (4.13), we first notice that $u_{l,n}$ is the unique solution to (4.10) with $2s - \delta > 3/2$. For each fixed $n \in \mathbb{Z}^+$, we can go along the lines as we derive (3.3) with using (4.11) to find

$$\mathbb{E} \sup_{t \in [0, T \wedge \tau_{l,n}^R]} \|u_{l,n}(t)\|_{H^{2s-\delta}}^2 \leq C\mathbb{E} \|u^{l,n}(0)\|_{H^{2s-\delta}}^2 + C_R \int_0^T \left(\mathbb{E} \sup_{t' \in [0, t \wedge \tau_{l,n}^R]} \|u_{l,n}(t')\|_{H^{2s-\delta}}^2 \right) dt. \quad (4.15)$$

From (4.15), we can use the Grönwall inequality and Lemma A.6 to infer

$$\mathbb{E} \sup_{t \in [0, T \wedge \tau_{l,n}^R]} \|u_{l,n}(t)\|_{H^{2s-\delta}}^2 \leq C\mathbb{E} \|u^{l,n}(0)\|_{H^{2s-\delta}}^2 \leq Cn^{2s-2\delta}, \quad C = C(R, T).$$

By Lemma A.6 again, we have that for some $C = C(R, T)$ and l satisfying (4.1),

$$\mathbb{E} \sup_{t \in [0, T \wedge \tau_{l,n}^R]} \|v\|_{H^{2s-\delta}}^2 \leq C\mathbb{E} \sup_{t \in [0, T \wedge \tau_{l,n}^R]} \|u_{l,n}\|_{H^{2s-\delta}}^2 + C\mathbb{E} \sup_{t \in [0, T \wedge \tau_{l,n}^R]} \|u^{l,n}\|_{H^{2s-\delta}}^2 \leq Cn^{2s-2\delta},$$

which is (4.13). \square

4.4. Final proof for Theorem 2.2. To begin with, we will show that if the exiting time of the zero solution is strongly stable, then $\{u_{-1,n}\}$ and $\{u_{1,n}\}$ are two sequences of pathwise solutions such that (2.8)–(2.11) are satisfied.

Lemma 4.3. *If for some $R_0 \gg 1$, the R_0 -exiting time of the zero solution to (1.5) is strongly stable, then for l satisfying (4.1) and $\tau_{l,n}^{R_0}$ given in (4.11), we have*

$$\lim_{n \rightarrow \infty} \tau_{l,n}^{R_0} = \infty \quad \mathbb{P} - a.s. \quad (4.16)$$

Proof. We notice that for all $s' < s$, $\lim_{n \rightarrow \infty} \|u^{l,n}(0)\|_{H^{s'}} = \lim_{n \rightarrow \infty} \|u_{l,n}(0) - 0\|_{H^{s'}} = 0$. Since the unique solution with zero initial data to (1.5) is zero and the R_0 -exiting time of the zero solution is ∞ , we see that (4.16) holds true provided the R_0 -exiting time of the zero solution to (1.5) is strongly stable. \square

Proof for Theorem 2.2. For each $n > 1$, for l satisfying (4.1) and for the fixed $R_0 \gg 1$, Lemma A.6 and (4.11) give us $\mathbb{P}\{\tau_{l,n}^{R_0} > 0\} = 1$ and Lemma 4.3 implies (2.8). Besides, Theorem 2.1 and (4.11) show that $u_{l,n} \in C([0, \tau_{l,n}^{R_0}]; H^s) \quad \mathbb{P} - a.s.$ and (2.9) holds true. By virtue of the interpolation inequality, we have

$$\begin{aligned} & \mathbb{E} \sup_{t \in [0, T \wedge \tau_{l,n}^{R_0}]} \|u^{l,n} - u_{l,n}\|_{H^s} \\ & \leq C \left(\mathbb{E} \sup_{t \in [0, T \wedge \tau_{l,n}^{R_0}]} \|u^{l,n} - u_{l,n}\|_{H^\delta} \right)^{\frac{1}{2}} \left(\mathbb{E} \sup_{t \in [0, T \wedge \tau_{l,n}^{R_0}]} \|u^{l,n} - u_{l,n}\|_{H^{2s-\delta}} \right)^{\frac{1}{2}} \\ & \leq C \left(\mathbb{E} \sup_{t \in [0, T \wedge \tau_{l,n}^{R_0}]} \|u^{l,n} - u_{l,n}\|_{H^\delta}^2 \right)^{\frac{1}{4}} \left(\mathbb{E} \sup_{t \in [0, T \wedge \tau_{l,n}^{R_0}]} \|u^{l,n} - u_{l,n}\|_{H^{2s-\delta}}^2 \right)^{\frac{1}{4}}. \end{aligned}$$

For any $T > 0$, combining Lemma 4.2 and the above estimate yields

$$\mathbb{E} \sup_{t \in [0, T \wedge \tau_{l,n}^{R_0}]} \|u^{l,n} - u_{l,n}\|_{H^s} \lesssim n^{-\frac{1}{4} \cdot 2r_s} \cdot n^{\frac{1}{4} \cdot (2s-2\delta)} = n^{r'_s}, \quad (4.17)$$

for some $C = C(R_0, T)$ and l satisfying (4.1). Recalling $k \geq 2$ and (4.3), we have

$$0 > r'_s = -r_s \cdot \frac{1}{2} + (s - \delta) \cdot \frac{1}{2} = \begin{cases} -\frac{s}{2} + \frac{k+1}{2k}, & \text{if } \frac{3}{2} < s \leq \frac{2k+1}{k}, \\ -\frac{1}{2}, & \text{if } s > \frac{2k+1}{k}. \end{cases}$$

Therefore for l satisfying (4.1),

$$\lim_{n \rightarrow \infty} \mathbb{E} \sup_{t \in [0, T \wedge \tau_{l,n}^{R_0}]} \|u^{l,n} - u_{l,n}\|_{H^s} = 0. \quad (4.18)$$

For odd k , (2.10) is given by

$$\|u_{-1,n}(0) - u_{1,n}(0)\|_{H^s} = \|u^{-1,n}(0) - u^{1,n}(0)\|_{H^s} \lesssim n^{-1/k} \rightarrow 0, \text{ as } n \rightarrow \infty.$$

For even k , (2.10) is given by

$$\|u_{0,n}(0) - u_{1,n}(0)\|_{H^s} = \|u^{0,n}(0) - u^{1,n}(0)\|_{H^s} \lesssim n^{-1/k} \rightarrow 0, \text{ as } n \rightarrow \infty.$$

When k is odd, we use (4.18) to find that for any $T > 0$,

$$\begin{aligned} & \liminf_{n \rightarrow \infty} \mathbb{E} \sup_{t \in [0, T \wedge \tau_{-1,n}^{R_0} \wedge \tau_{1,n}^{R_0}]} \|u_{-1,n}(t) - u_{1,n}(t)\|_{H^s} \\ & \geq \liminf_{n \rightarrow \infty} \mathbb{E} \sup_{t \in [0, T \wedge \tau_{-1,n}^{R_0} \wedge \tau_{1,n}^{R_0}]} \|u^{-1,n}(t) - u^{1,n}(t)\|_{H^s} \\ & \quad - \lim_{n \rightarrow \infty} \mathbb{E} \sup_{t \in [0, T \wedge \tau_{-1,n}^{R_0} \wedge \tau_{1,n}^{R_0}]} \|u^{-1,n}(t) - u_{-1,n}(t)\|_{H^s} \\ & \quad - \lim_{n \rightarrow \infty} \mathbb{E} \sup_{t \in [0, T \wedge \tau_{-1,n}^{R_0} \wedge \tau_{1,n}^{R_0}]} \|u^{1,n}(t) - u_{1,n}(t)\|_{H^s} \\ & \gtrsim \liminf_{n \rightarrow \infty} \mathbb{E} \sup_{t \in [0, T \wedge \tau_{-1,n}^{R_0} \wedge \tau_{1,n}^{R_0}]} \|u^{-1,n}(t) - u^{1,n}(t)\|_{H^s} \\ & \gtrsim \liminf_{n \rightarrow \infty} \mathbb{E} \sup_{t \in [0, T \wedge \tau_{-1,n}^{R_0} \wedge \tau_{1,n}^{R_0}]} \|-2n^{-1} + n^{-s} \cos(nx+t) - n^{-s} \cos(nx-t)\|_{H^s} \\ & \gtrsim \liminf_{n \rightarrow \infty} \mathbb{E} \sup_{t \in [0, T \wedge \tau_{-1,n}^{R_0} \wedge \tau_{1,n}^{R_0}]} (n^{-s} \|\sin(nx)\|_{H^s} |\sin t| - \|2n^{-1}\|_{H^s}) \gtrsim \sup_{t \in [0, T]} |\sin t|, \end{aligned} \quad (4.19)$$

where we have used Fatou's lemma. By

$$\left(\mathbb{E} \sup_{t \in [0, T \wedge \tau_{-1,n}^{R_0} \wedge \tau_{1,n}^{R_0}]} \|u_{-1,n}(t) - u_{1,n}(t)\|_{H^s}^2 \right)^{\frac{1}{2}} \geq \mathbb{E} \sup_{t \in [0, T \wedge \tau_{-1,n}^{R_0} \wedge \tau_{1,n}^{R_0}]} \|u_{-1,n}(t) - u_{1,n}(t)\|_{H^s}$$

and (4.19), we obtain (2.11). Similarly, we can also prove (2.11) when k is even. The proof is therefore completed. \square

5. GLOBAL EXISTENCE

In this section, we study the global existence and the blow-up of solutions to (1.8), and estimate the associated probabilities. Motivated by [29, 57, 58], we introduce the following Girsanov type transform

$$v = \frac{1}{\beta(\omega, t)} u, \quad \beta(\omega, t) = e^{\int_0^t b(t') dW_{t'} - \int_0^t \frac{b^2(t')}{2} dt'}. \quad (5.1)$$

Proposition 5.1. *Let $s > 3/2$ and $h(t, u) = b(t)u$ such that $b(t)$ satisfies Assumption A₃. Let $\mathcal{S} = (\Omega, \mathcal{F}, \mathbb{P}, \{\mathcal{F}_t\}_{t \geq 0}, W)$ be fixed in advance. If $u_0(\omega, x)$ is an H^s -valued \mathcal{F}_0 -measurable random variable with $\mathbb{E}\|u_0\|_{H^s}^2 < \infty$ and (u, τ^*) is the corresponding unique maximal pathwise solution to (1.8), then for $t \in [0, \tau^*)$, the process v given in (5.1) solves*

$$\begin{cases} v_t + \beta^k v^k v_x + \beta^k F(v) = 0, & t > 0, x \in \mathbb{T}, \\ v(\omega, 0, x) = u_0(\omega, x), & x \in \mathbb{T}. \end{cases} \quad (5.2)$$

Moreover, one has $v \in C([0, \tau^*); H^s) \cap C^1([0, \tau^*); H^{s-1})$ \mathbb{P} -a.s. and, if $s > 3$, then

$$\mathbb{P}\{\|v(t)\|_{H^1} = \|u_0\|_{H^1}\} = 1. \quad (5.3)$$

Proof. Since $b(t)$ satisfies Assumption A₃, $h(t, u) = b(t)u$ satisfies Assumption A₁. Consequently, Theorem 2.1 implies that (1.8) has a unique maximal pathwise solution (u, τ^*) . Direct computation shows

$$dv = (-\beta^k v^k v_x - \beta^k F(v)) dt. \quad (5.4)$$

Since $v(0) = u_0(\omega, x)$, we see that v satisfies (5.2). Moreover, Theorem 2.1 implies $u \in C([0, \tau^*); H^s) \mathbb{P} - a.s.$, so $v \in C([0, \tau^*); H^s) \mathbb{P} - a.s.$ Besides, from Lemma A.5 and (5.2)₁, we see that for a.e. $\omega \in \Omega$, $v_t = -\beta^k v^k v_x - \beta^k F(v) \in C([0, \tau^*); H^{s-1})$. Hence $v \in C^1([0, \tau^*); H^{s-1}) \mathbb{P} - a.s.$

Notice that if $s > 3$, (5.2)₁ is equivalent to

$$v_t - v_{xxt} + (k+2)\beta^k v^k v_x = (k+1)\beta^k v^{k-1} v_x v_{xx} + \beta^k v^k v_{xxx}, \quad k \geq 1. \quad (5.5)$$

Multiplying both sides of the above equation by v and then integrating the resulting equation on $x \in \mathbb{T}$ with noticing that $(k+1)v^k v_x v_{xx} + v^{k+1} v_{xxx} = \partial_x (v^{k+1} v_{xx})$, we see that for a.e. $\omega \in \Omega$ and for all $t > 0$

$$\frac{d}{dt} \int_{\mathbb{T}} (v^2 + v_x^2) dx = 0,$$

which implies (5.3). \square

5.1. Global existence: Case 1. Now we prove the first global existence result.

Proof for Theorem 2.3. To begin with, we apply the operator D^s to (5.4), multiply both sides of the resulting equation by $D^s v$ and integrate over \mathbb{T} to obtain that for a.e. $\omega \in \Omega$,

$$\frac{1}{2} \frac{d}{dt} \|v(t)\|_{H^s}^2 = -\beta^k(\omega, t) \int_{\mathbb{T}} D^s v \cdot D^s [v^k v_x] dx - \beta^k(\omega, t) \int_{\mathbb{T}} D^s v \cdot D^s F(v) dx.$$

Using Lemmas A.2 and A.5, we conclude that there is a $C = C(s) > 1$ such that for a.e. $\omega \in \Omega$,

$$\frac{d}{dt} \|v(t)\|_{H^s}^2 \leq C\beta^k(t) \|v\|_{W^{1,\infty}}^k \|v\|_{H^s}^2,$$

where β is given in (5.1). Then $w = e^{-\int_0^t b(t') dW_{t'}} u = e^{-\int_0^t \frac{b^2(t')}{2} dt'} v$ satisfies

$$\frac{d}{dt} \|w(t)\|_{H^s} + \frac{b^2(t)}{2} \|w(t)\|_{H^s} \leq C\alpha^k(\omega, t) \|w(t)\|_{W^{1,\infty}}^k \|w(t)\|_{H^s}, \quad \alpha(\omega, t) = e^{\int_0^t b(t') dW_{t'}}.$$

Let $R > 1$ and $\lambda_1 > 2$. Assume $\|u_0\|_{H^s} < \frac{1}{RK} \left(\frac{b_*}{C\lambda_1} \right)^{1/k}$ almost surely. Define

$$\tau_1(\omega) = \inf \left\{ t > 0 : \alpha^k(\omega, t) \|w\|_{W^{1,\infty}}^k = \|u\|_{W^{1,\infty}}^k > \frac{b^2(t)}{C\lambda_1} \right\}. \quad (5.6)$$

Notice that $\|u(0)\|_{W^{1,\infty}}^k \leq K^k \|u(0)\|_{H^s}^k < \frac{b_*}{C\lambda_1}$. Therefore we have $\mathbb{P}\{\tau_1 > 0\} = 1$, and for $t \in [0, \tau_1)$,

$$\frac{d}{dt} \|w(t)\|_{H^s} + \frac{(\lambda_1 - 2)b^2(t)}{2\lambda_1} \|w(t)\|_{H^s} \leq 0.$$

The above inequality and $w = e^{-\int_0^t b(t') dW_{t'}} u$ imply that for a.e. $\omega \in \Omega$, for any $\lambda_2 > \frac{2\lambda_1}{\lambda_1 - 2}$ and for $t \in [0, \tau_1)$,

$$\begin{aligned} \|u(t)\|_{H^s} &\leq \|w_0\|_{H^s} e^{\int_0^t b(t') dW_{t'}} e^{-\int_0^t \frac{(\lambda_1 - 2)b^2(t')}{2\lambda_1} dt'} \\ &= \|u_0\|_{H^s} e^{\int_0^t b(t') dW_{t'}} e^{-\int_0^t \frac{b^2(t')}{\lambda_2} dt'} e^{-\frac{((\lambda_1 - 2)\lambda_2 - 2\lambda_1)}{2\lambda_1 \lambda_2} \int_0^t b^2(t') dt'} \end{aligned} \quad (5.7)$$

Define the stopping time

$$\tau_2 = \tau_2(\omega) = \inf \left\{ t > 0 : e^{\int_0^t b(t') dW_{t'}} e^{-\int_0^t \frac{b^2(t')}{\lambda_2} dt'} > R \right\}. \quad (5.8)$$

Notice that $\mathbb{P}\{\tau_2 > 0\} = 1$. From (5.7), we have that almost surely

$$\begin{aligned} \|u(t)\|_{H^s} &< \frac{1}{RK} \left(\frac{b_*}{C\lambda_1} \right)^{1/k} \times R \times e^{-\frac{((\lambda_1 - 2)\lambda_2 - 2\lambda_1)}{2\lambda_1 \lambda_2} \int_0^t b^2(t') dt'} \\ &= \frac{1}{K} \left(\frac{b_*}{C\lambda_1} \right)^{1/k} e^{-\frac{((\lambda_1 - 2)\lambda_2 - 2\lambda_1)}{2\lambda_1 \lambda_2} \int_0^t b^2(t') dt'} \leq \frac{1}{K} \left(\frac{b_*}{C\lambda_1} \right)^{1/k}, \quad t \in [0, \tau_1 \wedge \tau_2). \end{aligned} \quad (5.9)$$

Combining (5.9) and (5.6), we find that

$$\mathbb{P}\{\tau_1 \geq \tau_2\} = 1. \quad (5.10)$$

Therefore it follows from (5.9) that

$$\mathbb{P}\left\{\|u(t)\|_{H^s} < \frac{1}{K} \left(\frac{b_*}{C\lambda_1}\right)^{1/k} e^{-\frac{((\lambda_1-2)\lambda_2-2\lambda_1)}{2\lambda_1\lambda_2} \int_0^t b^2(t')dt'} \text{ for all } t > 0\right\} \geq \mathbb{P}\{\tau_2 = +\infty\}.$$

We apply (ii) in Lemma A.7 to find that

$$\mathbb{P}\{\tau_2 = +\infty\} > 1 - \left(\frac{1}{R}\right)^{2/\lambda_2},$$

which completes the proof. \square

5.2. Global existence: Case 2. Now we prove Theorem 2.4. Let $\beta(\omega, t)$ be given in (5.1). From Proposition 5.1, we see that for a.e. $\omega \in \Omega$, $v(\omega, t, x)$ solves (5.2) on $[0, \tau^*)$. Moreover, since $H^s \hookrightarrow C^2$ for $s > 3/2$, we have $v, v_x \in C^1([0, \tau^*) \times \mathbb{T})$. Then for a.e. $\omega \in \Omega$, for any $x \in \mathbb{T}$, the problem

$$\begin{cases} \frac{dq(\omega, t, x)}{dt} = \beta^k(\omega, t) v^k(\omega, t, q(\omega, t, x)), & t \in [0, \tau^*), \\ q(\omega, 0, x) = x, & x \in \mathbb{T}, \end{cases} \quad (5.11)$$

has a unique solution $q(\omega, t, x)$ such that $q(\omega, t, x) \in C^1([0, \tau^*) \times \mathbb{T})$ almost surely. Moreover, differentiating (5.11) with respect to x yields that for a.e. $\omega \in \Omega$,

$$\begin{cases} \frac{dq_x}{dt} = k\beta^k(\omega, t) v^{k-1} v_x q_x, & t \in [0, \tau^*), \\ q_x(\omega, 0, x) = 1, & x \in \mathbb{T}. \end{cases}$$

For a.e. $\omega \in \Omega$, we solve the above equation to obtain

$$q_x(\omega, t, x) = \exp\left(\int_0^t k\beta^k(\omega, t') v^{k-1} v_x(\omega, t', q(\omega, t', x)) dt'\right).$$

Thus for all $(t, x) \in [0, \tau^*) \times \mathbb{T}$, we find $q_x > 0$ almost surely.

Lemma 5.1. *Let $s > 3$, $V_0(\omega, x) = (1 - \partial_{xx}^2)u_0(\omega, x)$ and $V(\omega, t, x) = v(\omega, t, x) - v_{xx}(\omega, t, x)$, where $v(\omega, t, x)$ solves (5.2) on $[0, \tau^*)$ \mathbb{P} -a.s. Then for all $(t, x) \in [0, \tau^*) \times \mathbb{T}$,*

$$\text{sign}(v) = \text{sign}(V) = \text{sign}(V_0) \quad \mathbb{P} - \text{a.s.}$$

Proof. We first notice that (5.2)₁ is equivalent to (5.5). Therefore for a.e. $\omega \in \Omega$, $V = v - v_{xx}$ satisfies

$$V_t + \beta^k v^k V_x + (k+1)\beta^k V v^{k-1} v_x = 0.$$

Thus for a.e. $\omega \in \Omega$, we have

$$\begin{aligned} \frac{d}{dt} \left[V(\omega, t, q(\omega, t, x)) q_x^{\frac{k+1}{k}}(\omega, t, x) \right] &= V_t q_x^{\frac{k+1}{k}} + V_x q_t q_x^{\frac{k+1}{k}} + \frac{k+1}{k} V q_x^{\frac{k+1}{k}-1} q_{xt} \\ &= q_x^{\frac{k+1}{k}} \left[V_t + V_x (\beta^k v^k) + \frac{k+1}{k} V q_x^{-1} (k\beta^k v^{k-1} v_x q_x) \right] \\ &= q_x^{\frac{k+1}{k}} [V_t + \beta^k v^k V_x + (k+1)\beta^k V v^{k-1} v_x] = 0. \end{aligned}$$

Notice that $q_x(\omega, 0, x) = 1$ and $q_x > 0$ almost surely. Then we have that $\text{sign}(V) = \text{sign}(V_0)$ \mathbb{P} -a.s. Since $v = G_{\mathbb{T}} * V$ with $G_{\mathbb{T}} > 0$ given in (1.7), we have $\text{sign}(v) = \text{sign}(V)$ \mathbb{P} -a.s. \square

Lemma 5.2. *Let all the conditions as in the statement of Proposition 5.1 hold true. If*

$$\mathbb{P}\{V_0(\omega, x) > 0, \forall x \in \mathbb{T}\} = p, \quad \mathbb{P}\{V_0(\omega, x) < 0, \forall x \in \mathbb{T}\} = q,$$

for some $p, q \in [0, 1]$, then the maximal pathwise solution u to (1.8) satisfies

$$\mathbb{P}\{\|u_x(\omega, t)\|_{L^\infty} \leq \|u(\omega, t)\|_{L^\infty} \lesssim \beta(\omega, t)\|u_0\|_{H^1}, \forall t \in [0, \tau^*)\} \geq p + q.$$

Proof. It is easy to see that for a.e. $\omega \in \Omega$ and for all $(t, x) \in [0, \tau^*) \times \mathbb{T}$,

$$[v + v_x](\omega, t, x) = \frac{1}{2 \sinh(\pi)} \int_0^{2\pi} e^{(x-y-2\pi[\frac{x-y}{2\pi}]-\pi)} V(\omega, t, y) dy, \quad (5.12)$$

$$[v - v_x](\omega, t, x) = \frac{1}{2 \sinh(\pi)} \int_0^{2\pi} e^{(y-x+2\pi[\frac{x-y}{2\pi}]+\pi)} V(\omega, t, y) dy. \quad (5.13)$$

Then one can employ (5.12), (5.13) and Lemma 5.1 to obtain that for a.e. $\omega \in \Omega$ and for all $(t, x) \in [0, \tau^*) \times \mathbb{T}$,

$$\begin{cases} -v(\omega, t, x) \leq v_x(\omega, t, x) \leq v(\omega, t, x), & \text{if } V_0(\omega, x) = (1 - \partial_{xx}^2)u_0(\omega, x) > 0, \\ v(\omega, t, x) \leq v_x(\omega, t, x) \leq -v(\omega, t, x), & \text{if } V_0(\omega, x) = (1 - \partial_{xx}^2)u_0(\omega, x) < 0. \end{cases} \quad (5.14)$$

Notice that

$$\{V_0(\omega, x) > 0\} \cap \{V_0(\omega, x) < 0\} = \emptyset. \quad (5.15)$$

Combining (5.14) and (5.15) yields

$$\mathbb{P}\{|v_x(\omega, t, x)| \leq |v(\omega, t, x)|, \forall (t, x) \in [0, \tau^*) \times \mathbb{T}\} \geq p + q. \quad (5.16)$$

In view of $H^1 \hookrightarrow L^\infty$, (5.3) and (5.16), we arrive at

$$\mathbb{P}\{\|v_x(\omega, t)\|_{L^\infty} \leq \|v(\omega, t)\|_{L^\infty} \lesssim \|v(\omega, t)\|_{H^1} = \|u_0\|_{H^1}, \forall t \in [0, \tau^*)\} \geq p + q.$$

Via (5.1), we obtain the desired estimate. \square

Proof for Theorem 2.4. Let (u, τ^*) be the maximal pathwise solution to (1.8). Then Lemma 5.2 implies that

$$\mathbb{P}\{\|u\|_{W^{1,\infty}} \lesssim 2\beta(\omega, t)\|u_0\|_{H^1}, \forall t \in [0, \tau^*)\} \geq p + q.$$

It follows from (i) in Lemma A.7 that $\sup_{t>0} \beta(\omega, t) < \infty$ \mathbb{P} -a.s. Then we can infer from (2.7) that $\mathbb{P}\{\tau^* = \infty\} \geq p + q$. That is to say, $\mathbb{P}\{u \text{ exists globally}\} \geq p + q$. \square

ACKNOWLEDGEMENT

The authors would like to express their great gratitude to the anonymous referees for their valuable suggestions, which have led to a meaningful improvement of this paper.

APPENDIX A. AUXILIARY RESULTS

In this section we formulate and prove some estimates employed in above proofs. We first recall the Friedrichs mollifier J_ε defined as

$$J_\varepsilon f(x) = j_\varepsilon * f(x).$$

Here $*$ stands for the convolution, $j_\varepsilon(x) = \sum_{k \in \mathbb{Z}} \hat{j}(\varepsilon k) e^{ixk}$ and $j(x)$ is a Schwartz function satisfying $0 \leq \hat{j}(\xi) \leq 1$ for all the $\xi \in \mathbb{R}$ and $\hat{j}(\xi) = 1$ for any $\xi \in [-1, 1]$. We also need another mollifier T_ε on \mathbb{T}^d with $d \geq 1$ as

$$T_\varepsilon f(x) := (1 - \varepsilon^2 \Delta)^{-1} f(x) = \sum_{k \in \mathbb{Z}^d} (1 + \varepsilon^2 |k|^2)^{-1} \hat{f}(k) e^{ix \cdot k}, \quad \varepsilon \in (0, 1).$$

From the construction, we see that J_ε and T_ε enjoy the following estimates, see [58, 59] for example,

$$[D^s, J_\varepsilon] = [D^s, T_\varepsilon] = 0, \quad (A.1)$$

$$(J_\varepsilon f, g)_{L^2} = (f, J_\varepsilon g)_{L^2}, \quad (T_\varepsilon f, g)_{L^2} = (f, T_\varepsilon g)_{L^2}, \quad (A.2)$$

$$\|J_\varepsilon u\|_{H^s}, \|T_\varepsilon u\|_{H^s} \leq \|u\|_{H^s}. \quad (A.3)$$

Lemma A.1 ([59]). *Let $d \geq 1$ and $f, g : \mathbb{T}^d \rightarrow \mathbb{R}^d$ such that $g \in W^{1,\infty}$ and $f \in L^2$. Then for some $C > 0$,*

$$\|[T_\varepsilon, (g \cdot \nabla)]f\|_{L^2} \leq C \|\nabla g\|_{L^\infty} \|f\|_{L^2}.$$

Lemma A.2 ([43]). *If $f \in H^s \cap W^{1,\infty}$, $g \in H^{s-1} \cap L^\infty$ for $s > 0$, then*

$$\|[D^s, f]g\|_{L^2} \leq C_s (\|D^s f\|_{L^2} \|g\|_{L^\infty} + \|\partial_x f\|_{L^\infty} \|D^{s-1} g\|_{L^2}).$$

Lemma A.3 ([43]). *Let $s > 0$, for all $f, g \in H^s \cap L^\infty$,*

$$\|fg\|_{H^s} \leq C_s(\|f\|_{H^s}\|g\|_{L^\infty} + \|f\|_{L^\infty}\|g\|_{H^s}).$$

Lemma A.4 (Proposition 4.2, [64]). *If $\rho > 3/2$ and $0 \leq \eta + 1 \leq \rho$, then for some $c > 0$,*

$$\|[D^\eta \partial_x, f]v\|_{L^2} \leq c\|f\|_{H^\rho}\|v\|_{H^\eta}, \quad \forall f \in H^\rho, v \in H^\eta.$$

Lemma A.5. *For the $F(\cdot)$ defined in (1.6) with $F_3(u)$ disappearing for $k = 1$, we have that for all $k \geq 1$,*

$$\|F(v)\|_{H^s} \lesssim \|v\|_{W^{1,\infty}}^k \|v\|_{H^s}, \quad s > 3/2,$$

$$\|F(u) - F(v)\|_{H^s} \lesssim (\|u\|_{H^s} + \|v\|_{H^s})^k \|u - v\|_{H^s}, \quad s > 3/2,$$

$$\|F(u) - F(v)\|_{H^s} \lesssim (\|u\|_{H^{s+1}} + \|v\|_{H^{s+1}})^k \|u - v\|_{H^s}, \quad 1/2 < s \leq 3/2$$

Proof. Since $s > 3/2$, the desired result follows from Lemma A.3 immediately. We omit the details and we refer to [63] for the CH case. \square

Lemma A.6 ([34, 63]). *Let $\sigma, \alpha \in \mathbb{R}$. If $\lambda \in \mathbb{Z}^+$ and $\lambda \gg 1$, then*

$$\|\sin(\lambda x - \alpha)\|_{H^\sigma} = \|\cos(\lambda x - \alpha)\|_{H^\sigma} \approx \lambda^\sigma.$$

Lemma A.7. *Let Assumption A₃ hold true and let $a(t) \in C([0, \infty))$ be a bounded function. For*

$$X = e^{\int_0^t b(t') dW_{t'} + \int_0^t a(t') - \frac{b^2(t')}{2} dt'},$$

we have the following properties:

(i) *Let $\phi(t) := \int_0^t b^2(t') dt'$ and $\phi^{-1}(t)$ be the inverse function of ϕ . If*

$$\limsup_{t \rightarrow \infty} \frac{1}{\sqrt{2t \log \log t}} \left(\int_0^{\phi^{-1}(t)} a(t') dt' - \frac{t}{2} \right) < -1, \quad (\text{A.4})$$

then

$$\lim_{t \rightarrow \infty} X(t) = 0 \quad \mathbb{P} - a.s. \quad (\text{A.5})$$

If

$$\liminf_{t \rightarrow \infty} \frac{1}{\sqrt{2t \log \log t}} \left(\int_0^{\phi^{-1}(t)} a(t') dt' - \frac{t}{2} \right) > 1, \quad (\text{A.6})$$

then

$$\lim_{t \rightarrow \infty} X(t) = +\infty \quad \mathbb{P} - a.s. \quad (\text{A.7})$$

(ii) *Let $a(t) = \lambda b^2(t)$ with $\lambda < \frac{1}{2}$ and $\tau_R = \inf\{t \geq 0 : X(t) > R\}$ with $R > 1$, then*

$$\mathbb{P}(\tau_R = \infty) \geq 1 - \left(\frac{1}{R} \right)^{1-2\lambda}. \quad (\text{A.8})$$

Proof. Let $Q(t) = \int_0^t b(t') dW_{t'} + \int_0^t a(t') - \frac{b^2(t')}{2} dt'$. Since $\phi : [0, \infty) \ni t \mapsto \phi(t) \in [0, \infty)$ is one-to-one (since ϕ is strictly increasing) and onto (by Assumption A₃), $\phi^{-1}(t)$ is well defined and $\lim_{t \rightarrow \infty} Q(t) = \lim_{t \rightarrow \infty} Q(\phi^{-1}(t))$ $\mathbb{P} - a.s.$ Direct computation (see [1, Exercise 7.7 and Theorem 8.2] for example) shows that

$$B_t := \int_0^{\phi^{-1}(t)} b(t') dW_{t'}$$

is itself a Brownian motion. Then we use the law of the iterated logarithm to get

$$\limsup_{t \rightarrow \infty} \frac{B_t}{\sqrt{2t \log \log t}} = 1, \quad \liminf_{t \rightarrow \infty} \frac{B_t}{\sqrt{2t \log \log t}} = -1 \quad \mathbb{P} - a.s.$$

Therefore (A.4) implies

$$\lim_{t \rightarrow \infty} Q(\phi^{-1}(t)) \leq \limsup_{t \rightarrow \infty} \sqrt{2t \log \log t} \left(\frac{B_t}{\sqrt{2t \log \log t}} + \frac{\int_0^{\phi^{-1}(t)} a(t') dt' - \frac{t}{2}}{\sqrt{2t \log \log t}} \right) = -\infty \quad \mathbb{P} - a.s.,$$

which leads to

$$\lim_{t \rightarrow \infty} X(t) = e^{\lim_{t \rightarrow \infty} Q(\phi^{-1}(t))} = 0 \quad \mathbb{P} - a.s.$$

Hence we obtain (A.5). If (A.6) holds true, then

$$\lim_{t \rightarrow \infty} Q(\phi^{-1}(t)) \geq \liminf_{t \rightarrow \infty} \sqrt{2t \log \log t} \left(\frac{B_t}{\sqrt{2t \log \log t}} + \frac{\int_0^{\phi^{-1}(t)} a(t') dt' - \frac{t}{2}}{\sqrt{2t \log \log t}} \right) = +\infty \quad \mathbb{P} - a.s.,$$

which gives (A.7). Now we consider (A.8) and exploit [29]. It is easy to see that when $a(t) = \lambda b^2(t)$, X is the unique global solution to the problem

$$dX = \lambda b^2(t) X dt + b(t) X dW, \quad X(0) = 1.$$

We apply the Itô formula to X^p with $p > 0$ to obtain that

$$dX^p = \left(p\lambda b^2(t) + \frac{b^2(t)p(p-1)}{2} \right) X^p dt + pb(t) X^p dW.$$

Particularly, if $p = 1 - 2\lambda$, then

$$dX^{1-2\lambda} = (1 - 2\lambda) b(t) X^{1-2\lambda} dW,$$

which means $\mathbb{E}X^{1-2\lambda}(t \wedge \tau_R) = 1$. Hence by the definition of τ_R , continuity of measures and the Chebyshev inequality, we have

$$\begin{aligned} \mathbb{P}\{\tau_R = \infty\} &= \lim_{N \rightarrow \infty} \mathbb{P}(\tau_R > N) \\ &= \lim_{N \rightarrow \infty} \mathbb{P}(X^{1-2\lambda}(N \wedge \tau_R) < R^{1-2\lambda}) \\ &\geq \lim_{N \rightarrow \infty} \left(1 - \frac{\mathbb{E}X^{1-2\lambda}(N \wedge \tau_R)}{R^{1-2\lambda}} \right) = 1 - \frac{1}{R^{1-2\lambda}}, \end{aligned}$$

which is (A.8). □

REFERENCES

- [1] P. Baldi. *Stochastic calculus*. Universitext. Springer, Cham, 2017.
- [2] J. L. Bona and N. Tzvetkov. Sharp well-posedness results for the BBM equation. *Discrete Contin. Dyn. Syst.*, 23(4):1241–1252, 2009.
- [3] D. Breit, E. Feireisl, and M. Hofmanová. Local strong solutions to the stochastic compressible Navier-Stokes system. *Comm. Partial Differential Equations*, 43(2):313–345, 2018.
- [4] D. Breit, E. Feireisl, and M. Hofmanová. *Stochastically forced compressible fluid flows*, volume 3 of *De Gruyter Series in Applied and Numerical Mathematics*. De Gruyter, Berlin, 2018.
- [5] D. Breit and M. Hofmanová. Stochastic Navier-Stokes equations for compressible fluids. *Indiana Univ. Math. J.*, 65(4):1183–1250, 2016.
- [6] A. Bressan and A. Constantin. Global conservative solutions of the Camassa-Holm equation. *Arch. Ration. Mech. Anal.*, 183(2):215–239, 2007.
- [7] A. Bressan and A. Constantin. Global dissipative solutions of the Camassa-Holm equation. *Anal. Appl. (Singap.)*, 5(1):1–27, 2007.
- [8] R. Camassa and D. D. Holm. An integrable shallow water equation with peaked solitons. *Phys. Rev. Lett.*, 71(11):1661–1664, 1993.
- [9] D. Chen, Y. Li, and W. Yan. On the Cauchy problem for a generalized Camassa-Holm equation. *Discrete Contin. Dyn. Syst.*, 35(3):871–889, 2015.
- [10] Y. Chen and H. Gao. Well-posedness and large deviations of the stochastic modified Camassa-Holm equation. *Potential Anal.*, 45(2):331–354, 2016.
- [11] Y. Chen, H. Gao, and B. Guo. Well-posedness for stochastic Camassa-Holm equation. *J. Differential Equations*, 253(8):2353–2379, 2012.
- [12] A. Constantin. On the blow-up of solutions of a periodic shallow water equation. *J. Nonlinear Sci.*, 10(3):391–399, 2000.
- [13] A. Constantin. The trajectories of particles in Stokes waves. *Invent. Math.*, 166(3):523–535, 2006.
- [14] A. Constantin and J. Escher. Wave breaking for nonlinear nonlocal shallow water equations. *Acta Math.*, 181(2):229–243, 1998.
- [15] A. Constantin and J. Escher. Well-posedness, global existence, and blowup phenomena for a periodic quasi-linear hyperbolic equation. *Comm. Pure Appl. Math.*, 51(5):475–504, 1998.
- [16] A. Constantin and J. Escher. Particle trajectories in solitary water waves. *Bull. Amer. Math. Soc. (N.S.)*, 44(3):423–431, 2007.
- [17] A. Constantin and J. Escher. Analyticity of periodic traveling free surface water waves with vorticity. *Ann. of Math. (2)*, 173(1):559–568, 2011.
- [18] D. Crisan, F. Flandoli, and D. D. Holm. Solution Properties of a 3D Stochastic Euler Fluid Equation. *J. Nonlinear Sci.*, 29(3):813–870, 2019.
- [19] G. Da Prato and J. Zabczyk. *Stochastic equations in infinite dimensions*, volume 152 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, second edition, 2014.

- [20] A. Debussche, N. E. Glatt-Holtz, and R. Temam. Local martingale and pathwise solutions for an abstract fluids model. *Phys. D*, 240(14-15):1123–1144, 2011.
- [21] E. Fedrizzi, W. Neves, and C. Olivera. On a class of stochastic transport equations for L^2_{loc} vector fields. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 18(2):397–419, 2018.
- [22] F. Flandoli. An introduction to 3D stochastic fluid dynamics. 1942:51–150, 2008.
- [23] F. Flandoli, M. Gubinelli, and E. Priola. Well-posedness of the transport equation by stochastic perturbation. *Invent. Math.*, 180(1):1–53, 2010.
- [24] B. Fuchssteiner and A. S. Fokas. Symplectic structures, their Bäcklund transformations and hereditary symmetries. *Phys. D*, 4(1):47–66, 1981/82.
- [25] L. Gawarecki and V. Mandrekar. *Stochastic differential equations in infinite dimensions with applications to stochastic partial differential equations*. Probability and its Applications (New York). Springer, Heidelberg, 2011.
- [26] X. Geng and B. Xue. An extension of integrable peakon equations with cubic nonlinearity. *Nonlinearity*, 22(8):1847–1856, 2009.
- [27] B. Gess and P. E. Souganidis. Long-time behavior, invariant measures, and regularizing effects for stochastic scalar conservation laws. *Comm. Pure Appl. Math.*, 70(8):1562–1597, 2017.
- [28] N. Glatt-Holtz and M. Ziane. Strong pathwise solutions of the stochastic Navier-Stokes system. *Adv. Differential Equations*, 14(5-6):567–600, 2009.
- [29] N. E. Glatt-Holtz and V. C. Vicol. Local and global existence of smooth solutions for the stochastic Euler equations with multiplicative noise. *Ann. Probab.*, 42(1):80–145, 2014.
- [30] I. Gyöngy and N. Krylov. Existence of strong solutions for Itô’s stochastic equations via approximations. *Probab. Theory Related Fields*, 105(2):143–158, 1996.
- [31] D. Henry. Geometric theory of semilinear parabolic equations. 840:iv+348, 1981.
- [32] A. A. Himonas and C. Holliman. The Cauchy problem for a generalized Camassa-Holm equation. *Adv. Differential Equations*, 19(1-2):161–200, 2014.
- [33] A. A. Himonas and C. Kenig. Non-uniform dependence on initial data for the CH equation on the line. *Differential Integral Equations*, 22(3-4):201–224, 2009.
- [34] A. A. Himonas, C. Kenig, and G. Misiołek. Non-uniform dependence for the periodic CH equation. *Comm. Partial Differential Equations*, 35(6):1145–1162, 2010.
- [35] A. A. Himonas and G. Misiołek. Non-uniform dependence on initial data of solutions to the Euler equations of hydrodynamics. *Comm. Math. Phys.*, 296(1):285–301, 2010.
- [36] M. Hofmanová. Degenerate parabolic stochastic partial differential equations. *Stochastic Process. Appl.*, 123(12):4294–4336, 2013.
- [37] H. Holden and X. Raynaud. Global conservative solutions of the Camassa-Holm equation—a Lagrangian point of view. *Comm. Partial Differential Equations*, 32(10-12):1511–1549, 2007.
- [38] H. Holden and X. Raynaud. Dissipative solutions for the Camassa-Holm equation. *Discrete Contin. Dyn. Syst.*, 24(4):1047–1112, 2009.
- [39] A. N. W. Hone, H. Lundmark, and J. Szmigielski. Explicit multipeakon solutions of Novikov’s cubically nonlinear integrable Camassa-Holm type equation. *Dyn. Partial Differ. Equ.*, 6(3):253–289, 2009.
- [40] G. Kallianpur and J. Xiong. Stochastic differential equations in infinite-dimensional spaces. 26:vi+342, 1995. Expanded version of the lectures delivered as part of the 1993 Barrett Lectures at the University of Tennessee, Knoxville, TN, March 25–27, 1993, With a foreword by Balram S. Rajput and Jan Rosinski.
- [41] A. Karczewska. Stochastic integral with respect to cylindrical Wiener process. *Ann. Univ. Mariae Curie-Skłodowska Sect. A*, 52(2):79–93, 1998.
- [42] T. Kato. The Cauchy problem for quasi-linear symmetric hyperbolic systems. *Arch. Rational Mech. Anal.*, 58(3):181–205, 1975.
- [43] T. Kato and G. Ponce. Commutator estimates and the Euler and Navier-Stokes equations. *Comm. Pure Appl. Math.*, 41(7):891–907, 1988.
- [44] C. E. Kenig, G. Ponce, and L. Vega. Well-posedness and scattering results for the generalized Korteweg-de Vries equation via the contraction principle. *Comm. Pure Appl. Math.*, 46(4):527–620, 1993.
- [45] C. E. Kenig, G. Ponce, and L. Vega. On the ill-posedness of some canonical dispersive equations. *Duke Math. J.*, 106(3):617–633, 2001.
- [46] J. U. Kim. On the Cauchy problem for the transport equation with random noise. *J. Funct. Anal.*, 259(12):3328–3359, 2010.
- [47] H. Koch and N. Tzvetkov. Nonlinear wave interactions for the Benjamin-Ono equation. *Int. Math. Res. Not.*, (30):1833–1847, 2005.
- [48] I. Kröker and C. Rohde. Finite volume schemes for hyperbolic balance laws with multiplicative noise. *Appl. Numer. Math.*, 62(4):441–456, 2012.
- [49] J. Lenells and M. Wunsch. On the weakly dissipative Camassa-Holm, Degasperis-Procesi, and Novikov equations. *J. Differential Equations*, 255(3):441–448, 2013.
- [50] P.-L. Lions, B. Perthame, and P. E. Souganidis. Scalar conservation laws with rough (stochastic) fluxes. *Stoch. Partial Differ. Equ. Anal. Comput.*, 1(4):664–686, 2013.
- [51] P.-L. Lions, B. Perthame, and P. E. Souganidis. Scalar conservation laws with rough (stochastic) fluxes: the spatially dependent case. *Stoch. Partial Differ. Equ. Anal. Comput.*, 2(4):517–538, 2014.
- [52] H. P. McKean. Breakdown of a shallow water equation. *Asian J. Math.*, 2(4):867–874, 1998. Mikio Sato: a great Japanese mathematician of the twentieth century.
- [53] D. A. C. Mollinedo and C. Olivera. Well-posedness of the stochastic transport equation with unbounded drift. *Bull. Braz. Math. Soc. (N.S.)*, 48(4):663–677, 2017.

- [54] W. Neves and C. Olivera. Wellposedness for stochastic continuity equations with Ladyzhenskaya-Prodi-Serrin condition. *NoDEA Nonlinear Differential Equations Appl.*, 22(5):1247–1258, 2015.
- [55] V. Novikov. Generalizations of the Camassa-Holm equation. *J. Phys. A*, 42(34):342002, 14, 2009.
- [56] C. Prévôt and M. Röckner. *A concise course on stochastic partial differential equations*, volume 1905 of *Lecture Notes in Mathematics*. Springer, Berlin, 2007.
- [57] M. Röckner, R. Zhu, and X. Zhu. Local existence and non-explosion of solutions for stochastic fractional partial differential equations driven by multiplicative noise. *Stochastic Process. Appl.*, 124(5):1974–2002, 2014.
- [58] H. Tang. On the pathwise solutions to the Camassa-Holm equation with multiplicative noise. *SIAM J. Math. Anal.*, 50(1):1322–1366, 2018.
- [59] H. Tang. Noise effects on dependence on initial data and blow-up for stochastic Euler–Poincaré equations. *arXiv:2002.08719*, 2020.
- [60] H. Tang and Z. Liu. Continuous properties of the solution map for the Euler equations. *J. Math. Phys.*, 55(3):031504, 10, 2014.
- [61] H. Tang and Z. Liu. Well-posedness of the modified Camassa-Holm equation in Besov spaces. *Z. Angew. Math. Phys.*, 66(4):1559–1580, 2015.
- [62] H. Tang, S. Shi, and Z. Liu. The dependences on initial data for the b-family equation in critical Besov space. *Monatsh. Math.*, 177(3):471–492, 2015.
- [63] H. Tang, Y. Zhao, and Z. Liu. A note on the solution map for the periodic Camassa-Holm equation. *Appl. Anal.*, 93(8):1745–1760, 2014.
- [64] M. Taylor. Commutator estimates. *Proc. Amer. Math. Soc.*, 131(5):1501–1507, 2003.
- [65] S. Wu and Z. Yin. Global existence and blow-up phenomena for the weakly dissipative Camassa-Holm equation. *J. Differential Equations*, 246(11):4309–4321, 2009.
- [66] Y. Zhao, Y. Li, and W. Yan. Local well-posedness and persistence property for the generalized Novikov equation. *Discrete Contin. Dyn. Syst.*, 34(2):803–820, 2014.
- [67] S. Zhou and C. Mu. The properties of solutions for a generalized b-family equation with peakons. *J. Nonlinear Sci.*, 23(5):863–889, 2013.

INSTITUT FÜR ANGEWANDTE ANALYSIS UND NUMERISCHE SIMULATION, UNIVERSITÄT STUTTGART, PFAFFENWALDRING 57,
70569 STUTTGART, GERMANY

E-mail address: christian.rohde@mathematik.uni-stuttgart.de

INSTITUT FÜR ANGEWANDTE ANALYSIS UND NUMERISCHE SIMULATION, UNIVERSITÄT STUTTGART, PFAFFENWALDRING 57,
70569 STUTTGART, GERMANY

E-mail address, Corresponding author: Hao.Tang@mathematik.uni-stuttgart.de