

TROPICAL LINEAR REPRESENTATIONS OF THE CHINESE MONOID

ZUR IZHAKIAN AND GLENN MERLET

ABSTRACT. We introduce a faithful tropical linear representation of the Chinese monoid, and thus prove that this monoid admits all the semigroup identities satisfied by tropical triangular matrices.

INTRODUCTION

The **Chinese monoid** of rank n is the finitely generated presented monoid

$$\text{Ch}_n := \langle a_1, \dots, a_n \rangle,$$

subject to the relations

$$a_j a_k a_i = a_k a_j a_i = a_k a_i a_j \quad \text{for all } i \leq j \leq k. \quad (\text{CH})$$

Each element $x \in \text{Ch}_n$ has a unique presentation, called the **canonical form**, written as:

$$\mathfrak{C}(x) = b_1 b_2 \cdots b_n \quad (\text{CF.a})$$

with

$$\begin{aligned} b_1 &= a_1^{k_{11}}, \\ b_2 &= (a_2 a_1)^{k_{21}} a_2^{k_{22}}, \\ b_3 &= (a_3 a_1)^{k_{31}} (a_3 a_2)^{k_{32}} a_3^{k_{33}}, \\ &\vdots \\ b_n &= (a_n a_1)^{k_{n1}} (a_n a_2)^{k_{n2}} \cdots (a_n a_{n-1})^{k_{n(n-1)}} a_n^{k_{nn}}, \end{aligned} \quad (\text{CF.b})$$

where all powers k_{ji} are non-negative. This monoid is strongly related to the plactic monoid Plc_n [2], and thereby also to Young tableaux [15, 16], which play a major role in representation theory and algebraic combinatorics [6, 18, 19]. A characterisation of the equivalence classes of Ch_n and a cross-section theorem were provided by an algorithm similar to Schensted's algorithm for Plc_n [20].

The tropical (max-plus) semiring is the set $\mathbb{T} := \mathbb{R} \cup \{-\infty\}$ equipped with the operations of maximum and summation

$$a \vee b := \max\{a, b\}, \quad a + b := \text{sum}\{a, b\},$$

addition and multiplication, respectively. Square matrices over \mathbb{T} form the monoid $\mathcal{M}_n(\mathbb{T})$, whose multiplication is induced from the operations of \mathbb{T} in the familiar way. This matrix monoid has a polynomial growth. While matrices over an infinite field do not admit a semigroup identity [5], by relying on identities for tropical the monoid $\mathcal{U}_n(\mathbb{T})$ of triangular tropical matrices, [12, Theorem 3.7] proves that $\mathcal{M}_n(\mathbb{T})$ satisfies nontrivial semigroup identities. Partial results appear in [7, 8, 10, 11, 21, 17, 4].

Ch_n has polynomial growth of degree $n(n+1)/2$, and was used by Duchamp & Krob (1994) in the their classification of monoids having polynomial growth. Plc_n can be faithfully represented by matrices over the tropical semiring [9, 14]. In this paper, we prove that the same holds for Ch_n , which has the same growth as Plc_n .

Theorem I. *The Chinese monoid Ch_n has a faithful linear representation by tropical triangular matrices in $\mathcal{U}_{n(n+1)}(\mathbb{T})$, more precisely by block-diagonal triangular matrices with blocks of size 2.*

This theorem establishes a useful machinery to determine the canonical form (CF.a) of a given word in Ch_n (Remark 1.6), and also infers that

Corollary I. *Ch_n admits all the semigroup identities satisfied by the monoid $\mathcal{U}_2(\mathbb{T})$ of 2×2 tropical triangular matrices.*

2010 *Mathematics Subject Classification.* Primary: 20M05, 20M07, 20M30, 47D03; Secondary: 16R10, 68Q70, 14T05.

Key words and phrases. Tropical (max-plus) matrices, supertropical matrices, idempotent semirings, semigroup identities, semigroup representations, weighted diagrams.

Acknowledgment: The results within this paper were obtained under the auspices of the Resarch in Pairs program of the Mathematisches Forschungsinstitut Oberwolfach, Germany. The authors thank MFO for the excellent working environment.

The monoid algebra of Ch_n was studied in [13], showing that Ch_n embeds in a product of bicyclic monoids, and thus admits the Adjan identity [1]

$$ab^2a ab ab^2a = ab^2a ba ab^2a. \quad (\text{AD})$$

On the other hand, 2×2 tropical triangular matrices satisfy the same identities as the bicyclic monoid satisfy [3]. Therefore, Corollary I provides an alternative simpler proof of these identities, dismissing the embedding into bicyclic monoids. It also gives the following.

Corollary II. *For any n , the Chinese monoid Ch_n determines the same variety as the bicyclic monoid Bcy .*

Proof. The semigroup identities of Bcy (or equivalently of $\mathcal{U}_2(\mathbb{T})$) are satisfied by Ch_n by previous Corollary. On the other hand, since there is a surjective morphism that sends Ch_2 (which embeds in Ch_n) onto Bcy , the semigroup identities admitted by Ch_n are also satisfied by Ch_2 and Bcy . \square

1. REPRESENTATIONS OF Ch_n

A **tropical linear representation** of a monoid \mathcal{S} is a monoid homomorphism

$$\rho : \mathcal{S} \longrightarrow \mathcal{M}_N(\mathbb{T}), \quad N \in \mathbb{N}, \quad (1.1)$$

i.e., a map satisfying $\rho(xy) = \rho(x)\rho(y)$ for every $x, y \in \mathcal{S}$. A representation ρ is **faithful**, if ρ is injective. When ρ is faithful, \mathcal{S} inherits all the semigroup identities satisfied by $\mathcal{M}_N(\mathbb{T})$. A **semigroup identity** is a pair $\langle u, v \rangle$ of two words $u, v \in \mathcal{A}^*$ on an alphabet \mathcal{A} , and \mathcal{S} **satisfies** $\langle u, v \rangle$, if

$$\phi(u) = \phi(v) \text{ for every semigroup homomorphism } \phi : \mathcal{A}^+ \longrightarrow \mathcal{S}.$$

In other words, the equality $u = v$ holds for any substitution of elements of \mathcal{S} into the letters of u and v .

Given a representation $\rho : \text{Ch}_n \rightarrow \mathcal{M}_N(\mathbb{T})$ of $\text{Ch}_n = \langle a_1, \dots, a_n \rangle$, clearly

$$\begin{aligned} \rho(a_j a_k a_i) &= \rho(a_k a_j a_i) &= \rho(a_k a_i a_j) \\ &= \rho(a_j) \rho(a_k) \rho(a_i) &= \rho(a_k) \rho(a_j) \rho(a_i) &= \rho(a_k) \rho(a_i) \rho(a_j) \end{aligned}$$

for all $i \leq j \leq k$, since ρ is a monoid homomorphism.

We inductively build a faithful linear representation for Ch_{n+1} out of a faithful representation for Ch_n , i.e., a construction by induction on the number of generators. Appendix A presents an explicit faithful representation for the base case Ch_3 .

Lemma 1.1. *Let $\rho : \text{Ch}_n \rightarrow \mathcal{M}_N(\mathbb{T})$ be a representation of Ch_n . For every $\ell = 1, \dots, n$, the map*

$$\rho_\ell : \text{Ch}_{n+1} \longrightarrow \mathcal{M}_N(\mathbb{T}), \quad \rho_\ell : a_j \longmapsto \begin{cases} \rho(a_j) & \text{if } j \leq \ell, \\ \rho(a_{j-1}) & \text{otherwise,} \end{cases} \quad (1.2)$$

defines a representation of Ch_n .

Proof. If $1 \leq i' \leq j' \leq k' \leq n+1$, then $(\rho_\ell(a_{i'}), \rho_\ell(a_{j'}), \rho_\ell(a_{k'})) = (\rho(a_i), \rho(a_j), \rho(a_k))$ for some $1 \leq i \leq j \leq k \leq n$. Therefore, the relations of Ch_{n+1} are satisfied by the images of the generators $\rho_\ell(a_{j'})$ because the relations of Ch_n are satisfied by the images of the generators $\rho(a_j)$. \square

Given a representation $\rho : \text{Ch}_n \rightarrow \mathcal{M}_N(\mathbb{T})$ of Ch_n , $n \geq 3$, using (1.2) we construct the combined representation for Ch_{n+1} :

$$\tilde{\rho} := (\rho_r, \rho_s, \rho_t) : \text{Ch}_{n+1} \longrightarrow (\mathcal{M}_N(\mathbb{T}))^3, \quad r < s < t. \quad (1.3)$$

To prove faithfulness of $\tilde{\rho}$, we need some basic auxiliary lemmas, which are used later to manipulate the canonical form (CF.a) and its matrix image under representations.

Lemma 1.2. *The word $a_k a_i$ commutes with a_i and a_k for any $i \leq k$.*

Proof. Follows from (CH) by taking $j = i$ and $j = k$, respectively. \square

Lemma 1.3. *The words $a_k a_j$ and $a_k a_i$ commute for any $i \leq j \leq k$.*

Proof. Apply (CH) twice to get

$$(a_k a_j)(a_k a_i) = a_k(a_j a_k a_i) = a_k(a_k a_i a_j) = (a_k a_k a_i) a_j = (a_k a_i a_k) a_j = (a_k a_i)(a_k a_j). \quad \square$$

For any element $x \in \text{Ch}_n$, written in the canonical form (CF.a) as $\mathfrak{C}(x)$, there is a one-to-one correspondence between $\mathfrak{C}(x)$ and the $n(n+1)$ -tuple

$$\mathbf{k}(n) := (k_{11}, k_{21}, k_{22}, \dots, k_{i1}, k_{i2}, \dots, k_{(i-1)i}, k_{ii}, \dots, k_{n1}, k_{n2}, \dots, k_{n(n-1)}, k_{nn})$$

with $k_{ij} \in \mathbb{Z}_+$. Accordingly, $\mathfrak{C}(x)$ is written in terms of the n generators a_1, \dots, a_n as

$$\mathfrak{C}(x) = \mathfrak{C}_{\mathbf{k}(n)}(a_1, \dots, a_n), \quad x \in \text{Ch}_n.$$

For short, we write \mathbf{k} for $\mathbf{k}(n)$ and $\tilde{\mathbf{k}}$ for $\mathbf{k}(n+1)$.

Lemma 1.4. $\mathfrak{C}_{\tilde{\mathbf{k}}}(a_1, \dots, a_\ell, a_\ell, a_{\ell+1}, \dots, a_n) = \mathfrak{C}_{\mathbf{k}}(a_1, \dots, a_n)$ where

$$k_{ji} = \begin{cases} \tilde{k}_{ji} & \text{if } j < \ell, \\ \tilde{k}_{\ell i} + \tilde{k}_{(\ell+1)i} & \text{if } j = \ell > i, \\ \tilde{k}_{\ell\ell} + 2\tilde{k}_{(\ell+1)\ell} + \tilde{k}_{(\ell+1)(\ell+1)} & \text{if } j = i = \ell, \\ \tilde{k}_{(j+1)i} & \text{if } j > \ell > i, \\ \tilde{k}_{(j+1)\ell} + \tilde{k}_{(j+1)(\ell+1)} & \text{if } j > i = \ell, \\ \tilde{k}_{(j+1)(i+1)} & \text{if } j, i > \ell. \end{cases} \quad (1.4)$$

Proof. Use (CF.a) to write

$$\begin{aligned} \mathfrak{C}_{\tilde{\mathbf{k}}}(a_1, \dots, a_\ell, a_\ell, a_{\ell+1}, \dots, a_n) &= b_1 \cdots b_\ell b'_{\ell+1} \cdots b'_{n+1} \\ &= b_1 \cdots b_\ell \tilde{b}_\ell \cdots \tilde{b}_n \end{aligned}$$

in terms of b_j 's and b'_j 's, where the b_j 's are defined by (CF.b) and the b'_j 's, for $j \geq \ell$, are given as

$$b'_j = (a_{j-1}a_1)^{k_{j1}} \cdots (a_{j-1}a_\ell)^{k_{j\ell}} (a_{j-1}a_\ell)^{k_{j(\ell+1)}} \cdots (a_{j-1}a_{j-2})^{k_{j(j-1)}} a_{j-1}^{k_{jj}}.$$

By Lemmas 1.2 and 1.3, we see that

$$\begin{aligned} b_\ell \tilde{b}_\ell &= (a_\ell a_1)^{k_{\ell 1}} (a_\ell a_2)^{k_{\ell 2}} \cdots (a_\ell a_{\ell-1})^{k_{\ell(\ell-1)}} a_\ell^{k_{\ell\ell}} (a_\ell a_1)^{k_{(\ell+1)1}} \cdots \\ &\quad \cdots (a_\ell a_{\ell-1})^{k_{(\ell+1)(\ell-1)}} (a_\ell a_\ell)^{k_{(\ell+1)\ell}} a_\ell^{k_{(\ell+1)(\ell+1)}} \\ &= (a_\ell a_1)^{k_{\ell 1} + k_{(\ell+1)1}} \cdots (a_\ell a_{\ell-1})^{k_{\ell(\ell-1)} + k_{(\ell+1)(\ell-1)}} a_\ell^{k_{\ell\ell} + 2k_{(\ell+1)\ell} + k_{(\ell+1)(\ell+1)}}. \end{aligned}$$

For $j > \ell$ we have

$$\begin{aligned} \tilde{b}_j &= (a_j a_1)^{k_{(j+1)1}} \cdots (a_j a_{\ell-1})^{k_{(j+1)(\ell-1)}} (a_j a_\ell)^{k_{(j+1)\ell} + k_{(j+1)(\ell+1)}} (a_j a_{\ell+1})^{k_{(j+1)(\ell+1)}} \cdots \\ &\quad \cdots (a_j a_{j-1})^{k_{(j+1)j}} a_j^{k_{(j+1)(j+1)}}. \end{aligned}$$

□

We can now prove the following lemma.

Lemma 1.5. Let $\rho : \text{Ch}_n \rightarrow \mathcal{M}_N(\mathbb{T})$, $n \geq 3$, be a faithful representation of Ch_n . Then the representation

$$\tilde{\rho} := (\rho_1, \rho_2, \rho_n) : \text{Ch}_{n+1} \longrightarrow (\mathcal{M}_N(\mathbb{T}))^3$$

in (1.3) is faithful. If $\mathbf{k} \mapsto \rho(\mathfrak{C}_{\mathbf{k}}(a_1, \dots, a_n))$ is a restriction of an affine injective map, then $\tilde{\mathbf{k}} \mapsto \tilde{\rho}(\mathfrak{C}_{\tilde{\mathbf{k}}}(a_1, \dots, a_{n+1}))$ is also a restriction of an affine injective map.

Proof. By (1.2) we have

$$\begin{aligned} \rho_\ell(\mathfrak{C}_{\tilde{\mathbf{k}}}(a_1, \dots, a_{n+1})) &= \rho(\mathfrak{C}_{\tilde{\mathbf{k}}}(a_1, \dots, a_\ell, a_\ell, a_{\ell+1}, \dots, a_n)) \\ &= \rho(\mathfrak{C}_{\mathbf{k}}(a_1, \dots, a_\ell, a_{\ell+1}, \dots, a_n)). \end{aligned}$$

Thus, by Lemma 1.4 we see that, if $\mathbf{k} \mapsto \rho(\mathfrak{C}_{\mathbf{k}}(a_1, \dots, a_n))$ is a restriction of an affine map, then $\tilde{\mathbf{k}} \mapsto \tilde{\rho}(\mathfrak{C}_{\tilde{\mathbf{k}}}(a_1, \dots, a_{n+1}))$ is also a restriction of an affine map.

We show that, $\tilde{\mathbf{k}}$ can be determined, assuming that $\rho_\ell(\mathfrak{C}_{\tilde{\mathbf{k}}}(a_1, \dots, a_{n+1}))$ (or the image of some vector $\tilde{\mathbf{k}}$ by the affine map it defines) is known for $\ell = 1, 2, n$. By Lemma 1.4 and the faithfulness of ρ (or the injectivity of the affine map), we know the terms k_{ji} of the $n(n+1)$ -tuple \mathbf{k} , cf. (1.4), and we need to determine the \tilde{k}_{ji} 's explicitly. From (1.4) and the indices $\ell = 1, 2, n$ we deduce: $\ell = n$ gives \tilde{k}_{ji} for $i \leq j < n$, and $\ell = 1$ gives \tilde{k}_{ji} for $i, j > 2$. It remains to determine $\tilde{k}_{n1}, \tilde{k}_{n2}, \tilde{k}_{(n+1)1}, \tilde{k}_{(n+1)2}$. But, the sums $\tilde{k}_{n1} + \tilde{k}_{(n+1)1}$ and $\tilde{k}_{n2} + \tilde{k}_{(n+1)2}$ are known from the case of $\ell = n$, while $\tilde{k}_{n1} + \tilde{k}_{n2}$ and $\tilde{k}_{(n+1)1} + \tilde{k}_{(n+1)2}$ are known from the case of $\ell = 1$. The case of $\ell = 2$ gives the sum $\tilde{k}_{(n+1)2} + \tilde{k}_{(n+1)3}$, from which we deduce $\tilde{k}_{(n+1)2}$, as $\tilde{k}_{(n+1)3}$ is known. Then, by substitution, we obtain the values of the other three terms. □

Proof of Theorem I. Proof by induction of the number of generators of Ch_n , $n \geq 3$. The case of $n = 3$ is given by Lemma A.2 in the appendix. The induction step follows from Lemma 1.5. Thus, $(\mathcal{M}_N(\mathbb{T}))^3$ embeds in $\mathcal{M}_{3N}(\mathbb{T})$ as a 3-block diagonal matrix, each of these blocks by itself is a block diagonal matrix whose blocks are 2×2 triangular matrices. Moreover, $\mathbf{k} \mapsto \rho(\mathfrak{C}_{\mathbf{k}}(a_1, \dots, a_n))$ is a restriction of an affine map

$$\alpha : \mathbb{R}^{\frac{n(n+1)}{2}} \longrightarrow \mathbb{R}^{3^n}.$$

Elementary linear algebra implies that there is a projection ϕ of \mathbb{R}^{3^n} on $\frac{n(n+1)}{2}$ of its coordinates such that $\phi \circ \alpha$ is kept injective. Thus, the representation by the blocks appearing in the projection is faithful and Theorem I is proved. \square

Remark 1.6. *The use of the faithful representation (1.3) allows to compute the canonical form (CF.a) of a given word $x \in \text{Ch}_n$. Indeed, Lemma 1.5 determines inductively all the powers k_{ij} in (CF.a).*

REFERENCES

- [1] S. I. Adjan. *Defining relations and algorithmic problems for groups and semigroups*. Number 85. Proceeding of the Steklov Institute of Mathematics, American Mathematical Society, 1967.
- [2] J. Cassaigne, M. Espie, D. Krob, J.-C. Novelli, F. Hivert. The Chinese monoid, *International J. of Alg. Comp.*, 11(3): 301–334, 2001.
- [3] L. Daviaud, M. Johnson, M. Kambites. Identities in upper triangular tropical matrix semigroups and the bicyclic monoid, *J. Alg.*, 501:503–525, 2018.
- [4] L. Daviaud, M. Johnson. The shortest identities for max-plus automata with two states, In: K. G. Larsen, H. L. Bodlaender, J.-F. Raskin (eds.) *MFCS 2017*, Art. 48:1–13, 2017.
- [5] I. Golubchik, A. Mikhalev. A note on varieties of semiprime rings with semigroup identities, *J. Alg.*, 54:42–45, 1978.
- [6] W. Fulton. *Young Tableaux*. London Mathematical Society Student Texts, vol. 35. Cambridge University Press, 1997.
- [7] Z. Izhakian. Semigroup identities in the monoid of triangular tropical matrices, *Semigroup Forum*, 88(1):145–161, 2014.
- [8] Z. Izhakian. Erratum to: Semigroup identities in the monoid of triangular tropical matrices, *Semigroup Forum*, 92(3):733, 2016.
- [9] Z. Izhakian. Tropical plactic algebra, the cloaktic monoid, and semigroup representations, *J. Alg.*, 524:290–366, 2019.
- [10] Z. Izhakian. Semigroup identities of tropical matrix semigroups of maximal rank, *Semigroup Forum*, 92(3):712–732, 2016.
- [11] Z. Izhakian, S. Margolis. Semigroup identities in the monoid of 2-by-2 tropical matrices. *Semigroup Forum*, 80(2):191–218, 2010.
- [12] Z. Izhakian, G. Merlet. Semigroup identities of tropical matrices through matrix ranks, arXiv:1806.11028, 2018.
- [13] J. Jaszurńska, J. Okninski. Structure of Chinese algebras, *J. Alg.*, 346(1):31–81, 2011.
- [14] M. Johnson, M. Kambites. Tropical representations and identities of plactic monoids, arXiv:1906.03991, 2019.
- [15] A. Lascoux, B. Leclerc, J.-Y. Thibon. The plactic monoid. *Algebraic Combinatorics on Words*, Cambridge University Press, 164–196, 2002.
- [16] A. Lascoux, M.P. Schützenberger. Schubert polynomials and the Littlewood-Richardson rule. *Lett. Math. Phys.*, 10(2–3):111–125, 1985.
- [17] J. Okninski. Identities of the semigroup of upper triangular tropical matrices, *Commun. Alg.*, 43(10):4422–4426, 2015.
- [18] B. E. Sagan. *The Ubiquitous Young Tableau*. 1988.
- [19] B. E. Sagan. *The Symmetric Group: Representations, Combinatorial Algorithms, and Symmetric Functions*. Springer-Verlag, 2000.
- [20] C. Schensted. Longest increasing and decreasing subsequences. *Canad. J. Math.*, 13:179191, 1961.
- [21] Y. Shitov. A semigroup identity for tropical 3x3 matrices, arXiv:1406.2601, 2014.

APPENDIX A. LINEAR REPRESENTATIONS OF Ch_3

Considering the Chinese monoid $\text{Ch}_3 = \langle a, b, c \rangle$ of rank 3, we begin by constructing three linear representations

$$\rho_\ell : \text{Ch}_3 \longrightarrow \mathcal{M}_2(\mathbb{T}), \quad \ell = 1, 2, 3,$$

in terms of generator maps. These representations are later combined together to a faithful representation of Ch_3 . To simplify notations, in the below matrices we write “ $-$ ” for “ $-\infty$ ” and k_i for k_{ii} , $i = 1, 2, 3$. Let x be an element of Ch_3 , written in the canonical form

$$x = (a)^{k_1} (ba)^{k_{21}} b^{k_2} (ca)^{k_{31}} (cb)^{k_{32}} (c)^{k_3}. \quad (\text{A.1})$$

The representations were found by extensive search among triangular matrices of size two with small entries, performed with ScicosLab, a fork from SciLab which includes a max-plus toolbox. The computations below, which finally show that ρ is a faithful representation were performed with Mathematica.

A.1. Representations of Ch_3 .

Representation I: Let ρ_1 be defined by

$$a \mapsto A = \begin{pmatrix} 1 & 0 \\ - & 0 \end{pmatrix}, \quad b \mapsto B = \begin{pmatrix} 0 & 0 \\ - & 1 \end{pmatrix}, \quad c \mapsto C = \begin{pmatrix} 0 & 0 \\ - & 1 \end{pmatrix}, \quad (\text{I.1})$$

for which we have the products

$$\begin{aligned} BCA &= CBA = CAB = \begin{pmatrix} 1 & 1 \\ - & 2 \end{pmatrix}, \\ BBA &= BAB = \begin{pmatrix} 1 & 1 \\ - & 2 \end{pmatrix}, \quad ABA = BAA = \begin{pmatrix} 2 & 1 \\ - & 1 \end{pmatrix}, \\ BCB &= CBB = \begin{pmatrix} 0 & 2 \\ - & 3 \end{pmatrix}, \quad CCB = CBC = \begin{pmatrix} 0 & 2 \\ - & 3 \end{pmatrix}, \\ ACA &= CAA = \begin{pmatrix} 2 & 1 \\ - & 1 \end{pmatrix}, \quad CCA = CAC = \begin{pmatrix} 1 & 1 \\ - & 2 \end{pmatrix}, \\ BAC &= \begin{pmatrix} 1 & 1 \\ - & 2 \end{pmatrix}, \quad ABC = ACB = \begin{pmatrix} 1 & 2 \\ - & 2 \end{pmatrix}, \end{aligned} \quad (\text{I.2})$$

so that relation (CH) is satisfied. Taking powers of A, B, C , we get

$$\begin{aligned} A^{k_1} &= \begin{pmatrix} k_1 & \max\{0, k_1\} - 1 \\ - & 0 \end{pmatrix} = \begin{pmatrix} k_1 & k_1 - 1 \\ - & 0 \end{pmatrix}, \\ B^{k_2} &= \begin{pmatrix} 0 & \max\{0, k_2\} - 1 \\ - & k_2 \end{pmatrix} = \begin{pmatrix} 0 & k_2 - 1 \\ - & k_2 \end{pmatrix}, \\ C^{k_3} &= \begin{pmatrix} 0 & \max\{0, k_3\} - 1 \\ - & k_3 \end{pmatrix} = \begin{pmatrix} 0 & k_3 - 1 \\ - & k_3 \end{pmatrix}, \end{aligned} \quad (\text{I.3})$$

where for $k_1 = k_2 = k_3 = 0$ we have

$$A^0 = B^0 = C^0 = I_1 = \begin{pmatrix} 0 & -1 \\ - & 0 \end{pmatrix},$$

which is an identity matrix for the semigroup generated by A, B , and C .

For products of A, B , and C we have

$$BA = \begin{pmatrix} 1 & 0 \\ - & 1 \end{pmatrix}, \quad CA = \begin{pmatrix} 1 & 0 \\ - & 1 \end{pmatrix}, \quad CB = \begin{pmatrix} 0 & 1 \\ - & 2 \end{pmatrix}, \quad (\text{I.4})$$

have the powers

$$\begin{aligned} (BA)^{k_{21}} &= \begin{pmatrix} k_{21} & k_{21} - 1 \\ - & k_{21} \end{pmatrix}, \\ (CA)^{k_{31}} &= \begin{pmatrix} k_{31} & k_{31} - 1 \\ - & k_{31} \end{pmatrix}, \\ (CB)^{k_{32}} &= \begin{pmatrix} 0 & \max\{0, 2k_{32}\} - 1 \\ - & 2k_{32} \end{pmatrix} = \begin{pmatrix} k_{32} & 2k_{32} - 1 \\ - & 2k_{32} \end{pmatrix}. \end{aligned} \quad (\text{I.5})$$

Computing the matrix product corresponding to the canonical form (CF.a), we obtain

$$\begin{aligned} \rho_1(x) &= \begin{pmatrix} k_1 + k_{21} + k_{31} & k_{21} + k_{31} - 2 + \max \left\{ \begin{array}{l} k_1 + 1, \\ k_1 + k_3 + 2k_{32}, \\ k_1 + k_3 + 2k_{32} + 1, \\ k_2 + k_3 + 2k_{32} + 1, \\ k_1 + k_2 + k_3 + 2k_{32}, \\ k_1 + k_2 + k_3 + 2k_{32} + 1 \end{array} \right\} \\ -\infty & k_2 + k_3 + k_{21} + k_{31} + 2k_{32} \end{pmatrix} \\ &= \begin{pmatrix} k_1 + k_{21} + k_{31} & k_{21} + k_{31} + k_1 + k_2 + k_3 + 2k_{32} - 1 \\ -\infty & k_2 + k_3 + k_{21} + k_{31} + 2k_{32} \end{pmatrix}. \end{aligned} \quad (\text{I.6})$$

Representation II: Let ρ_2 be given by

$$a \mapsto A = \begin{pmatrix} 1 & 0 \\ - & 0 \end{pmatrix}, \quad b \mapsto B = \begin{pmatrix} 1 & 0 \\ - & 0 \end{pmatrix}, \quad c \mapsto C = \begin{pmatrix} 0 & 0 \\ - & 1 \end{pmatrix}, \quad (\text{II.1})$$

for which we have the products

$$\begin{aligned}
BCA &= CBA = CAB = \begin{pmatrix} 2 & 1 \\ - & 1 \end{pmatrix}, \\
BBA &= BAB = \begin{pmatrix} 3 & 2 \\ - & 0 \end{pmatrix}, & ABA &= BAA = \begin{pmatrix} 3 & 2 \\ - & 0 \end{pmatrix}, \\
BCB &= CBB = \begin{pmatrix} 2 & 1 \\ - & 1 \end{pmatrix}, & CCB &= CBC = \begin{pmatrix} 1 & 1 \\ - & 2 \end{pmatrix}, \\
ACA &= CAA = \begin{pmatrix} 2 & 1 \\ - & 1 \end{pmatrix}, & CCA &= CAC = \begin{pmatrix} 1 & 1 \\ - & 2 \end{pmatrix}, \\
BAC &= ABC = \begin{pmatrix} 2 & 2 \\ - & 1 \end{pmatrix}, & ACB &= \begin{pmatrix} 2 & 1 \\ - & 1 \end{pmatrix},
\end{aligned} \tag{II.2}$$

so that relation (CH) is satisfied, where $BAC \neq ABC = ACB$. Taking powers of A, B, C , we get

$$\begin{aligned}
A^{k_1} &= \begin{pmatrix} k_1 & \max\{0, k_1\} - 1 \\ - & 0 \end{pmatrix} = \begin{pmatrix} k_1 & k_1 - 1 \\ - & 0 \end{pmatrix}, \\
B^{k_2} &= \begin{pmatrix} k_2 & \max\{0, k_2\} - 1 \\ - & 0 \end{pmatrix} = \begin{pmatrix} k_2 & k_2 - 1 \\ - & 0 \end{pmatrix}, \\
C^{k_3} &= \begin{pmatrix} 0 & \max\{0, k_3\} - 1 \\ - & k_3 \end{pmatrix} = \begin{pmatrix} 0 & k_3 - 1 \\ - & k_3 \end{pmatrix},
\end{aligned} \tag{II.3}$$

where for $k_1 = k_2 = k_3 = 0$ we have

$$A^0 = B^0 = C^0 = I_2 = \begin{pmatrix} 0 & -1 \\ - & 0 \end{pmatrix},$$

which is an identity matrix for the semigroup generated by A, B , and C .

For products of A, B , and C we have

$$BA = \begin{pmatrix} 2 & 2 \\ - & 1 \end{pmatrix}, \quad CA = \begin{pmatrix} 2 & 2 \\ - & 1 \end{pmatrix}, \quad CB = \begin{pmatrix} 2 & 1 \\ - & 1 \end{pmatrix}, \tag{II.4}$$

with powers

$$\begin{aligned}
(BA)^{k_{21}} &= \begin{pmatrix} 2k_{21} & -1 + \max\{0, 2k_{21}\} \\ - & 0 \end{pmatrix} = \begin{pmatrix} 2k_{21} & 2k_{21} - 1 \\ - & 0 \end{pmatrix}, \\
(CA)^{k_{31}} &= \begin{pmatrix} k_{31} & k_{31} - 1 \\ - & k_{31} \end{pmatrix}, \\
(CB)^{k_{32}} &= \begin{pmatrix} k_{32} & k_{32} - 1 \\ - & k_{32} \end{pmatrix}.
\end{aligned} \tag{II.5}$$

Computing the matrix product corresponding to the canonical form (CF.a), we obtain

$$\begin{aligned}
\rho_2(x) &= \begin{pmatrix} k_1 + k_2 + 2k_{21} + k_{31} + k_{32} & k_{31} + k_{32} - 2 + \max \left\{ \begin{array}{l} k_3 + 1, \\ k_1 + k_2 + 2k_{21} + 1, \\ k_1 + k_2 + k_3 + 2k_{21}, \\ k_1 + k_2 + k_3 + 2k_{21} + 1 \end{array} \right\} \\ -\infty & k_3 + k_{31} + k_{32} \end{pmatrix} \\
&= \begin{pmatrix} k_1 + k_2 + 2k_{21} + k_{31} + k_{32} & k_{31} + k_{32} + k_1 + k_2 + k_3 + 2k_{21} - 1 \\ -\infty & k_3 + k_{31} + k_{32} \end{pmatrix}.
\end{aligned} \tag{II.6}$$

Representation III: Let ρ_3 be given by

$$a \mapsto A = \begin{pmatrix} 1 & 1 \\ - & 0 \end{pmatrix}, \quad b \mapsto B = \begin{pmatrix} 0 & 0 \\ - & 0 \end{pmatrix}, \quad c \mapsto C = \begin{pmatrix} 0 & 1 \\ - & 1 \end{pmatrix}, \tag{III.1}$$

for which we have the products

$$\begin{aligned}
BCA &= CBA = CAB = \begin{pmatrix} 1 & 1 \\ - & 1 \end{pmatrix}, \\
BBA &= BAB = \begin{pmatrix} 1 & 1 \\ - & 0 \end{pmatrix}, & ABA &= BAA = \begin{pmatrix} 2 & 2 \\ - & 0 \end{pmatrix}, \\
BCB &= CBB = \begin{pmatrix} 0 & 1 \\ - & 1 \end{pmatrix}, & CCB &= CBC = \begin{pmatrix} 0 & 2 \\ - & 2 \end{pmatrix}, \\
ACA &= CAA = \begin{pmatrix} 2 & 2 \\ - & 1 \end{pmatrix}, & CCA &= CAC = \begin{pmatrix} 1 & 2 \\ - & 2 \end{pmatrix}, \\
BAC &= ABC = ACB = \begin{pmatrix} 1 & 2 \\ - & 1 \end{pmatrix},
\end{aligned} \tag{III.2}$$

so that relation (CH) is satisfied, where $BAC = ABC \neq BAC$. Taking powers of A, B, C , we get

$$\begin{aligned}
A^{k_1} &= \begin{pmatrix} k_1 & \max\{0, k_1\} \\ - & 0 \end{pmatrix} = \begin{pmatrix} k_1 & k_1 \\ - & 0 \end{pmatrix}, \\
B^{k_2} &= \begin{pmatrix} 0 & 0 \\ - & 0 \end{pmatrix}, \\
C^{k_3} &= \begin{pmatrix} 0 & \max\{0, k_3\} \\ - & k_3 \end{pmatrix} = \begin{pmatrix} 0 & k_3 \\ - & k_3 \end{pmatrix},
\end{aligned} \tag{III.3}$$

where for $k_1 = k_2 = k_3 = 0$ we have

$$A^0 = B^0 = C^0 = I_3 = \begin{pmatrix} 0 & 0 \\ - & 0 \end{pmatrix},$$

which is an identity matrix for the semigroup generated by A, B , and C .

For products of A, B , and C , we have

$$BA = \begin{pmatrix} 1 & 1 \\ - & 0 \end{pmatrix}, \quad CA = \begin{pmatrix} 1 & 1 \\ - & 1 \end{pmatrix}, \quad CB = \begin{pmatrix} 0 & 1 \\ - & 1 \end{pmatrix}, \tag{III.4}$$

have the powers

$$\begin{aligned}
(BA)^{k_{21}} &= \begin{pmatrix} k_{21} & \max\{0, k_{21}\} \\ - & 0 \end{pmatrix} = \begin{pmatrix} k_{21} & k_{21} \\ - & 0 \end{pmatrix}, \\
(CA)^{k_{31}} &= \begin{pmatrix} k_{31} & k_{31} \\ - & k_{31} \end{pmatrix}, \\
(CB)^{k_{32}} &= \begin{pmatrix} 0 & \max\{0, k_{32}\} \\ - & k_{32} \end{pmatrix} = \begin{pmatrix} 0 & k_{32} \\ - & k_{32} \end{pmatrix}.
\end{aligned} \tag{III.5}$$

Computing the matrix product corresponding to the canonical form (CF.a), we obtain

$$\begin{aligned}
\rho_3(x) &= \begin{pmatrix} k_1 + k_{21} + k_{31} & k_{31} - 1 + \max \left\{ \begin{array}{l} k_1 + k_{21} + 1, \\ k_1 + k_3 + k_{21}, \\ k_3 + k_{32} + 1, \\ k_1 + k_3 + k_{32}, \\ k_1 + k_3 + k_{21} + k_{32}, \\ k_1 + k_3 + k_{21} + k_{32} + 1 \end{array} \right\} \\ -\infty & k_3 + k_{31} + k_{32} \end{pmatrix} \\
&= \begin{pmatrix} k_1 + k_{21} + k_{31} & k_{31} + k_1 + k_3 + k_{21} + k_{32} \\ -\infty & k_3 + k_{31} + k_{32} \end{pmatrix}.
\end{aligned} \tag{III.6}$$

Remark A.1. Note that, since $k_1, k_2, k_{21}, k_3, k_{31}, k_{32}$ are non-negative, for all the above representations, the image matrix (I.6), (II.6), and (III.6), of $x \in \text{Ch}_3$ satisfying Equation (A.1) is affine, and its entries are given explicitly in terms of the powers $k_1, k_2, k_{21}, k_3, k_{31}, k_{32}$.

A.2. A faithful representation of Ch_3 . Together, the three representations ρ_1, ρ_2, ρ_3 from above provide the representation

$$\rho := (\rho_1, \rho_2, \rho_3) : \text{Ch}_3 \longrightarrow (\mathcal{M}_2(\mathbb{T}))^3 \tag{A.1}$$

of Ch_3 in terms of generators. It is easily seen from (I.2), (II.2), and (III.2) that

$$\begin{aligned}
\rho(bca) &= \rho(cba) = \rho(cab), & \rho(aba) &= \rho(baa), & \rho(bba) &= \rho(bab), \\
\rho(bcc) &= \rho(cbc), & \rho(bcb) &= \rho(cbb), & \rho(acc) &= \rho(cac), & \rho(aca) &= \rho(caa),
\end{aligned}$$

and all are different from $\rho(bac) \neq \rho(abc) \neq \rho(acb)$, which are also different from each other. $(\mathcal{M}_2(\mathbb{T}))^3$ embeds as diagonal blocks in $\mathcal{M}_6(\mathbb{T})$, so that ρ can be realized as a map $\text{Ch}_3 \rightarrow \mathcal{M}_6(\mathbb{T})$.

Lemma A.2. *The map (A.1) is an injective semigroup homomorphism.*

Proof. Let x be an element of Ch_3 , written in the canonical form

$$x = (a)^{k_1} (ba)^{k_{21}} b^{k_2} (ca)^{k_{31}} (cb)^{k_{32}} (c)^{k_3}.$$

The entries of the matrix images $X = \rho_1(x)$, $Y = \rho_2(x)$, $Z = \rho_3(x)$ give the following system of equations

$$\begin{aligned} X_{1,1} &= k_1 + k_{21} + k_{31}, \\ X_{1,2} &= k_1 + k_2 + k_{21} + k_{31} + k_3 + 2k_{32} - 1, \\ X_{2,2} &= k_2 + k_{21} + k_3 + k_{31} + 2k_{32}, \\ Y_{1,1} &= k_1 + k_2 + 2k_{21} + k_{31} + k_{32}, \\ Y_{1,2} &= k_1 + k_2 + k_3 + 2k_{21} + k_{31} + k_{32} - 1, \\ Y_{2,2} &= k_3 + k_{31} + k_{32}, \\ Z_{1,1} &= k_1 + k_{21} + k_{31}, \\ Z_{1,2} &= k_{31} + k_1 + k_3 + k_{21} + k_{32}, \\ Z_{2,2} &= k_3 + k_{31} + k_{32}, \end{aligned} \tag{A.2}$$

cf. (I.6), (II.6), (III.6) respectively. This affine system has a unique solution, so that the powers $k_1, k_2, k_{21}, k_3, k_{31}, k_{32}$ are uniquely determined. Hence ρ is injective. \square

INSTITUTE OF MATHEMATICS, UNIVERSITY OF ABERDEEN, AB24 3UE, ABERDEEN, UK.
E-mail address: `zzur@abdn.ac.uk`

AIX MARSEILLE UNIV, CNRS, CENTRALE MARSEILLE, I2M, MARSEILLE, FRANCE
E-mail address: `glenn.merlet@univ-amu.fr`