

BOUNDARY REGULARITY FOR MANIFOLD CONSTRAINED $p(x)$ -HARMONIC MAPS

IWONA CHLEBICKA, CRISTIANA DE FILIPPIS, AND LUKAS KOCH

ABSTRACT. We prove partial and full boundary regularity for manifold constrained $p(x)$ -harmonic maps.

CONTENTS

1. Introduction	1
2. Preliminaries	3
2.1. Notation	3
2.2. Main assumptions	3
2.3. Well-known results	6
2.4. Extensions	6
3. Partial boundary regularity	7
3.1. Basic regularity results	8
3.2. Proof of Theorem 1	11
4. Full boundary regularity	23
4.1. Proof of Theorem 2	35
References	36

1. INTRODUCTION

In this paper we complete the partial regularity theory for $p(x)$ -harmonic maps studied in [9] providing partial and full boundary regularity for manifold constrained minima of the variable exponent energy:

$$(1.1) \quad g + \left(W^{1,p(\cdot)}(\Omega, \mathcal{M}) \cap W_0^{1,p(\cdot)}(\Omega, \mathbb{R}^N) \right) \ni w \mapsto \mathcal{E}(w, \Omega) := \int_{\Omega} k(x) |Dw|^{p(x)} \, dx$$

for a suitable boundary datum $g: \bar{\Omega} \rightarrow \mathcal{M}$. Our main accomplishment is that there exists a relatively (to $\bar{\Omega}$) open subset $\Omega_0 \subset \bar{\Omega}$ of full n -dimensional Lebesgue measure on which u is locally Hölder continuous and the singular set $\Sigma_0 := \bar{\Omega} \setminus \Omega_0$ has Hausdorff dimension at the most equal to $n - \gamma_1$, see (2.2)₁ below for more informations on this quantity. This is the content of the following theorem.

Theorem 1. *Under assumptions (2.1), (2.2), (2.3) and (2.6), let $u \in W^{1,p(\cdot)}(\Omega, \mathcal{M})$ be a solution to the Dirichlet problem (1.1) with boundary datum $g \in W^{1,q}(\bar{\Omega}, \mathcal{M})$ satisfying (2.7). Then there exists a relatively (to $\bar{\Omega}$) open subset $\Omega_0 \subseteq \bar{\Omega}$ so that $u \in C_{loc}^{0,1-\frac{n}{q}}(\Omega_0, \mathcal{M})$ with q as in (2.7) and $\mathcal{H}^{n-\gamma_1}(\Sigma_0) = 0$.*

2010 *Mathematics Subject Classification.* 35J25, 35J60, 35J70.

Key words and phrases. Regularity, manifold constrained problems, $p(x)$ -Laplacian.

Acknowledgements. I. Chlebicka is supported by NCN grant no. 2016/23/D/ST1/01072. C. De Filippis and L. Koch are supported by the Engineering and Physical Sciences Research Council (EPSRC): CDT Grant Ref. EP/L015811/1.

Moreover, after strengthening the hypotheses on the variable exponent $p(\cdot)$ and on the boundary datum $g(\cdot)$, we can prove that the singular set of solutions to problem

$$(1.2) \quad g + \left(W^{1,p(\cdot)}(\Omega, \mathcal{M}) \cap W_0^{1,p(\cdot)}(\Omega, \mathbb{R}^N) \right) \ni w \mapsto \mathcal{G}(w, \Omega) := \int_{\Omega} |Dw|^{p(x)} \, dx$$

does not intersect the boundary $\partial\Omega$. In this respect we have

Theorem 2. *Under assumptions (2.1), (2.4) and (2.6), let $u \in W^{1,p(\cdot)}(\Omega, \mathcal{M})$ be a solution to the Dirichlet problem (1.2) with boundary datum $g: \bar{\Omega} \rightarrow \mathcal{M}$ satisfying (2.7). Then there exists a constant $\Upsilon \equiv \Upsilon(\text{data}) \in (0, 1]$ such that if*

$$(1.3) \quad [g]_{0,1-\frac{n}{q};\bar{\Omega}} < \Upsilon,$$

then $\Sigma_0 \Subset \Omega$ and so u is $\left(1 - \frac{n}{q}\right)$ -Hölder continuous in a neighborhood of $\partial\Omega$.

We immediately refer to Section 2.2 for the complete list of assumptions in force concerning the regularity of $\partial\Omega$, the coefficients appearing in the energies displayed in (1.1)-(1.2) and the topology of the manifold \mathcal{M} . The results exposed in Theorems 1-2 are new already in the case $p(\cdot) \equiv \text{const}$. In fact, we recover for the $p(x)$ -Laplacian the boundary regularity theory already available for p -harmonic maps, under weaker assumptions on the boundary datum than those considered in [22, 30, 50]. Let us put our results into the context of the available literature. The regularity theory for vector-valued minimizers of functionals modelled upon the p -Laplacian integral, i.e. variational problems like

$$(1.4) \quad W_{loc}^{1,p}(\Omega, \mathbb{R}^N) \ni w \mapsto \int_{\Omega} F(x, Dw) \, dx$$

$$|z|^p \lesssim F(x, z) \lesssim (1 + |z|^2)^{\frac{p}{2}}, \quad 1 < p < \infty,$$

started with the seminal paper [53] and received several contributions later on, see [23–25, 28, 39, 42] and references therein for an overview of the state of the art concerning p -laplacean type problems. On the other hand, the regularity theory in the case when both minimizers and competitors take values into a manifold $\mathcal{M} \subset \mathbb{R}^N$ faces additional difficulties. The cornerstones of the theory were laid down by the fundamental papers [16, 18, 49, 50] analyzing harmonic maps, i.e., constrained minimizers of the functional in (1.4) for $p = 2$, see also [51]. We mention also the recent works [45, 46] for a fine analysis of the singular set of harmonic maps. The extension of such basic results to the case $p \neq 2$ has been done in the by now classical papers: [20–22, 30, 41]. Moreover, several results have been extended to more general functionals with p -growth, for instance the quasiconvex case has been treated in [36] while a purely PDE approach has been proposed in [15]. The matter of boundary regularity for vectorial problems is rather delicate and received lots of attention in the literature, starting from [37], which covers the case of quadratic functionals. This theory has been extended later on to variational integrals of p -laplacean type, see [13] for the first results in this direction and [3, 14, 26, 28, 38] for general systems with standard p -growth. On the other hand, we notice that energies of the type in (1.1) do not satisfy conditions as in (1.4), but rather, the more general and flexible one

$$(1.5) \quad W_{loc}^{1,p}(\Omega, \mathbb{R}^N) \ni w \mapsto \int_{\Omega} F(x, Dw) \, dx$$

$$|z|^p \lesssim F(x, z) \lesssim (1 + |z|^2)^{\frac{q}{2}} \quad 1 < p \leq q < \infty.$$

The systematic study of functionals as in (1.5) started in [43, 44] and, subsequently, has undergone an intensive development over the last years, see for instance [2, 4–6, 11, 17, 19, 32, 33, 35]. In particular, the energy in (1.1) have been introduced in the setting of Calculus of Variations and Homogenization in the seminal works [54–56]. Energies as in (1.1) also occur in the modelling of electro-rheological fluids, a class of non-newtonian fluids whose viscosity properties are influenced by the presence of external electromagnetic fields [1], see also [12] for the basic properties of the $p(x)$ -Laplacian. As for regularity, the first result in the vectorial case has been obtained in [8], where it is shown that local minimizers of energy (1.2) are locally $C^{1,\beta}$ -regular in the

unconstrained case. Subsequently, the regularity theory of functionals with variable growth has been developed in a series of interesting papers, [47, 48, 52], where the authors established partial regularity results for unconstrained minimizers that are on the other hand obviously related to the constrained case. Especially, in [52] is given an interesting partial regularity result and some singular set estimates for a class of functionals related to the constrained minimization problem in which minimizers are assumed to take values in a single chart. Finally, [9] is devoted to the study of partial inner regularity of manifold constrained $p(x)$ -harmonic maps and to the analysis and dimension-reduction of their singular set.

Organization of the paper. This paper is organized as follows: Section 2 contains our notation, the list of the assumptions which will rule problems (1.1)-(1.2), several by now classical tools in the framework of regularity theory and some results of geometric and topological nature on Lipschitz retractions. Finally, Sections 3-4 are devoted to the proof of Theorem 1 and Theorem 2 respectively.

2. PRELIMINARIES

In this section we display our notation, list the main assumptions in force throughout the paper and collect some useful tools for regularity theory and several well-known results in the framework of manifold-valued maps.

2.1. Notation. Following a usual custom, we denote by c a general constant larger than one. Different occurrences from line to line will be still denoted by c , while special occurrences will be denoted by c_1, c_2, \tilde{c} or the like. Relevant dependencies on parameters will be emphasized using parentheses, i.e., $c \equiv c(p, \nu, L)$ means that c depends on p, ν, L . Given any measurable subset $U \subset \mathbb{R}^n$, we denote by $|U|$ its n -dimensional Lebesgue measure and with $\mathcal{H}^k(U)$ its k -dimensional Hausdorff measure, for some $k \geq 0$. For a point $x_0 \in \mathbb{R}^n$ and a number $\varrho > 0$ we indicate with $B_\varrho(x_0) := \{x \in \mathbb{R}^n : |x - x_0| < \varrho\}$ the open ball centered at x_0 and with radius ϱ and further, $B_\varrho \equiv B_\varrho(0)$. Similarly, for $x_0 \in \mathbb{R}^{n-1} \times \{0\}$ we define the half ball centered at x_0 as: $B_\varrho^+(x_0) := \{x \in B_\varrho(x_0) : x^n > 0\}$. We moreover set $B_\varrho^+ \equiv B_\varrho^+(0)$. We also name $\Gamma_\varrho(x_0)$ the set $\{x \in \mathbb{R}^n : x^n = 0 \text{ and } |x_0 - x| < \varrho\}$ and $\partial^+ B_\varrho^+(x_0) := \partial B_\varrho^+(x_0) \setminus \Gamma_\varrho(x_0)$. As before, $\Gamma_\varrho \equiv \Gamma_\varrho(0)$. With $U \subset \mathbb{R}^n$ being a measurable subset having finite and positive n -dimensional Lebesgue measure, and with $h : U \rightarrow \mathbb{R}^k$, being a measurable map, we shall denote by

$$(h)_U \equiv \int_U h(x) \, dx := \frac{1}{|U|} \int_U h(x) \, dx$$

its integral average. Similarly, with $\gamma \in (0, 1)$ we denote the Hölder seminorm of h as

$$[h]_{0,\gamma;U} := \sup_{x,y \in U, x \neq y} \frac{|h(x) - h(y)|}{|x - y|^\gamma}.$$

It is well known that the quantity defined above is a seminorm and when $[h]_{0,\gamma;U} < \infty$, we will say that h belongs to the Hölder space $C^{0,\gamma}(U, \mathbb{R}^k)$. When clear from the context, we will omit the reference to U , i.e.: $[h]_{0,\gamma;U} \equiv [h]_{0,\gamma}$. Finally, given any set Γ allowing for a trace operator, we denote by $\text{tr}_\Gamma(h)$ the trace of h on Γ .

2.2. Main assumptions. Let us turn to the main assumptions that will characterize our problem. The set $\Omega \subset \mathbb{R}^n$, $n \geq 2$ is open, bounded, connected and

$$(2.1) \quad \partial\Omega \text{ is } C^2\text{-regular.}$$

When considering the functional in (1.1), the exponent $p(\cdot)$ will always satisfy

$$(2.2) \quad \begin{cases} p \in C^{0,\alpha}(\bar{\Omega}) \text{ for some } \alpha \in (0, 1] \\ 1 < \gamma_1 := \inf_{x \in \bar{\Omega}} p(x) \leq p(x) \leq \gamma_2 := \sup_{x \in \bar{\Omega}} p(x) < \infty, \end{cases}$$

while the coefficient $k(\cdot)$ is so that

$$(2.3) \quad \begin{cases} k \in C^{0,\nu}(\bar{\Omega}) & \text{for some } \nu \in (0, 1] \\ 0 < \lambda \leq k(x) \leq \Lambda < \infty & \text{for all } x \in \bar{\Omega} \end{cases}$$

holds true. We anticipate that in the estimates contained in Section 3.2, only $\min\{\alpha, \nu\}$ will be relevant, so, for simplicity, for the proof of Theorem 1 we will assume that $\alpha = \nu$, i.e.: $p(\cdot), k(\cdot) \in C^{0,\alpha}(\bar{\Omega})$. When dealing with the question of full boundary regularity, we need higher regularity for $p(\cdot)$. Precisely, we shall suppose that

$$(2.4) \quad \begin{cases} p \in C^{0,1}(\bar{\Omega}) \\ 2 \leq \gamma_1 \leq p(x) \leq \gamma_2 < \infty, \end{cases}$$

with γ_1 and γ_2 as in (2.2)₂. Given an half ball B_R^+ and a ball $B_\varrho(x_0)$ with $x_0 \in B_R^+$ and $\varrho \in (0, R - |x_0|)$, we denote

$$(2.5) \quad p_1(x_0, \varrho) := \inf_{x \in B_\varrho(x_0) \cap B_R^+} p(x) \quad \text{and} \quad p_2(x_0, \varrho) := \sup_{x \in B_\varrho(x_0) \cap B_R^+} p(x).$$

Since in (2.5) we will always consider the intersection with the same ball B_R^+ , the reference to R in the symbols p_1, p_2 is omitted. When clear from the context, in (2.5) we shall not mention x_0 i.e.: $p_i(x_0, \varrho) \equiv p_i(\varrho)$ for $i \in \{1, 2\}$. With a little abuse, we will adopt the notation in (2.5) also to denote the infimum (resp. the supremum) of $p(\cdot)$ on B_R^+ : the context will remove any ambiguity. Notice that there is no loss of generality in assuming $\gamma_1 < \gamma_2$, otherwise $p(\cdot) \equiv \text{const}$ on $\bar{\Omega}$, and in this case the problem is very well understood, [22, 30, 50]. Furthermore, we need to impose some topological restriction on the manifold \mathcal{M} . Precisely, we ask that

$$(2.6) \quad \begin{cases} \mathcal{M} \text{ is a compact, } m\text{-dimensional, } C^3 \text{ Riemannian submanifold of } \mathbb{R}^N \\ \mathcal{M} \text{ is } [\gamma_2] - 1 \text{ connected} \\ \partial\mathcal{M} = \emptyset. \end{cases}$$

Here $[x]$ denotes the integer part of x and the definition of j -connectedness is given in Section 2.4, Definition 4. Moreover, we assume that the boundary datum satisfies:

$$(2.7) \quad g \in W^{1,q}(\bar{\Omega}, \mathcal{M}) \quad \text{for some } q > \max\{n, \gamma_2\}.$$

Combining (2.7) with Morrey embedding theorem we automatically get that

$$(2.8) \quad g \in C^{0,1-\frac{n}{q}}(\bar{\Omega}, \mathcal{M}).$$

Finally, to shorten the notation we shall collect the main parameters of the problem in the quantities

$$\begin{aligned} \mathbf{data}_{p(\cdot)} &:= (n, N, \mathcal{M}, \lambda, \Lambda, \gamma_1, \gamma_2, q, [p]_{0,\alpha}, \alpha); \\ \mathbf{data} &:= (n, N, \mathcal{M}, \lambda, \Lambda, \gamma_1, \gamma_2, q, [k]_{0,\nu}, [p]_{0,\alpha}, \nu, \alpha). \end{aligned}$$

Any dependencies of the constants appearing in the forthcoming estimates from quantities depending on the characteristics of \mathcal{M} , such as, for instance, the L^∞ -norm of maps with range in \mathcal{M} (which is clearly finite being \mathcal{M} compact) will be simply denoted as a dependency from \mathcal{M} in the form: $c \equiv c(\mathcal{M})$.

Remark 2.1. Assumption (2.1) assures that there exists a positive constant $\hat{r} \equiv \hat{r}(n, \Omega)$ such that $B_\varrho(x_0) \cap \Omega$ is simply connected for all $\varrho \in (0, \hat{r}]$ and any $x_0 \in \partial\Omega$. This renders the existence of a positive constant $c \equiv c(n, \Omega)$ such that

$$\frac{\mathcal{H}^{n-1}(B_\varrho(x_0) \cap \partial\Omega)}{\mathcal{H}^{n-1}(\partial B_\varrho(x_0) \cap \Omega)} > c \quad \text{for all } \varrho \in (0, \hat{r}], x_0 \in \partial\Omega.$$

Moreover, the Ahlfors condition holds

$$|B_\varrho(x_0) \cap \Omega| \sim \varrho^n \quad \text{for all } x_0 \in \bar{\Omega}, \varrho \in (0, \hat{r}],$$

with constants implicit in " \sim " depending on n, Ω . We shall refer to such constants with the term "Ahlfors constants", see [13, Section 2].

As to fully clarify the framework we are going to adopt, we need to introduce some basic terminology on the so-called Musielak-Orlicz-Sobolev spaces. Essentially, these are Sobolev spaces defined by the fact that the distributional derivatives lie in a suitable Musielak-Orlicz space, rather than in a Lebesgue space as usual. Classical Sobolev spaces are then a particular case. Such spaces and related variational problems are discussed for instance in [7, 12, 31, 57], to which we refer for more details. Here, we will consider spaces related to the variable exponent case in both unconstrained and manifold-constrained settings.

Definition 1. *Given an open set $\Omega \subset \mathbb{R}^n$, the Musielak-Orlicz space $L^{p(\cdot)}(\Omega, \mathbb{R}^k)$, $k \geq 1$, with $p(\cdot)$ satisfying (2.2), is defined as*

$$L^{p(\cdot)}(\Omega, \mathbb{R}^k) := \left\{ w: \Omega \rightarrow \mathbb{R}^k \text{ measurable and } \int_{\Omega} |w|^{p(x)} dx < \infty \right\}$$

endowed with the Luxemburg norm $\|w\|_{L^{p(\cdot)}(\Omega, \mathbb{R}^k)} = \inf\{\lambda > 0 : \int_{\Omega} |w/\lambda|^{p(x)} dx < 1\}$. Consequently,

$$W^{1,p(\cdot)}(\Omega, \mathbb{R}^k) := \left\{ w \in W^{1,1}(\Omega, \mathbb{R}^k) \cap L^{p(\cdot)}(\Omega, \mathbb{R}^k) \text{ such that } |Dw| \in L^{p(\cdot)}(\Omega, \mathbb{R}^{k \times n}) \right\}$$

with the norm $\|w\|_{W^{1,p(\cdot)}(\Omega, \mathbb{R}^k)} = \|w\|_{L^{p(\cdot)}(\Omega, \mathbb{R}^k)} + \| |Dw| \|_{L^{p(\cdot)}(\Omega, \mathbb{R}^k)}$. The variant $W_{loc}^{1,p(\cdot)}(\Omega, \mathbb{R}^k)$ is defined as in the classical case, whereas $W_0^{1,p(\cdot)}(\Omega, \mathbb{R}^k)$ is a closure of smooth and compactly supported functions in the norm $\|\cdot\|_{W^{1,p(\cdot)}(\Omega, \mathbb{R}^k)}$.

It is well known that, under assumptions (2.2), the set of smooth maps is dense in $W^{1,p(\cdot)}(\Omega, \mathbb{R}^k)$, see e.g. [17, 57]. Following [9] we also recall the analogous definition of such spaces when mappings take values into \mathcal{M} .

Definition 2. *Let \mathcal{M} be a compact submanifold of \mathbb{R}^k , $k \geq 2$, without boundary and $\Omega \subset \mathbb{R}^n$ an open set. For $p(\cdot)$ satisfying (2.2), the Musielak-Orlicz-Sobolev space $W^{1,p(\cdot)}(\Omega, \mathcal{M})$ of functions into \mathcal{M} can be defined as*

$$W^{1,p(\cdot)}(\Omega, \mathcal{M}) := \left\{ w \in W^{1,p(\cdot)}(\Omega, \mathbb{R}^k) : w(x) \in \mathcal{M} \text{ for a.e. } x \in \Omega \right\}.$$

The local space $W_{loc}^{1,p(\cdot)}(\Omega, \mathcal{M})$ consists of maps belonging to $W^{1,p(\cdot)}(B, \mathcal{M})$ for all open sets $B \Subset \Omega$.

Of course, when $p(\cdot) \equiv \text{const}$, Definitions 1 and 2 reduce to the classical Sobolev spaces $W^{1,p}(\Omega, \mathbb{R}^k)$ and $W^{1,p}(\Omega, \mathcal{M})$ respectively. Since the regularity question in Ω is local in nature, we can choose coordinates $\{x^i\}_{i=1}^n$ centered at $x_0 \in \partial\Omega$ such that locally Ω is the upper half space $\mathbb{R}^n \cap \{x^n > 0\}$, therefore, to avoid unnecessary complications, from now on we will assume that $\Omega \equiv B_1^+$, see [13, 14, 30, 37, 38, 50] for a more detailed discussion on this matter. Let us display the definition of constrained $W^{1,p(\cdot)}$ -minimizer of (1.1) in B_1^+ .

Definition 3. *Let $g \in W^{1,q}(\bar{B}_1^+, \mathcal{M})$ be as in (2.7). A map $u \in W^{1,p(\cdot)}(B_1^+, \mathcal{M})$ with $\text{tr}_{\Gamma_1}(u) = g$, is a constrained minimizer of the functional in (1.1) in the Dirichlet class $C_g^{p(\cdot)}(B_1^+, \mathcal{M})$ provided that:*

$$x \mapsto k(x)|Du(x)|^{p(x)} \in L^1(B_1^+), \quad \text{tr}_{\Gamma_1}(u) = \text{tr}_{\Gamma_1}(g)$$

and

$$\mathcal{E}(u, B_1^+) \leq \mathcal{E}(w, B_1^+)$$

for all maps $w \in W^{1,p(\cdot)}(B_1^+, \mathcal{M})$ so that $(u - w) \in W_0^{1,p(\cdot)}(B_1^+, \mathbb{R}^N)$. The conditions displayed above define class $C_g^{p(\cdot)}(B_1^+, \mathcal{M})$.

To shorten the notation, for $\varrho \in (0, 1]$, $x_0 \in \mathbb{R}^n \cap \{x^n \geq 0\}$, $f \in W^{1,p(\cdot)}(\bar{B}_\varrho^+(x_0), \mathcal{M})$ and a subset $\mathcal{X} \subseteq \mathbb{R}^N$, we also introduce the general Dirichlet class

$$\hat{\mathcal{C}}_f^{p(\cdot)}(B_\varrho^+(x_0), \mathcal{X}) := f + \left(W^{1,p(\cdot)}(B_\varrho^+(x_0), \mathcal{X}) \cap W_0^{1,p(\cdot)}(B_\varrho^+(x_0), \mathbb{R}^N) \right).$$

Clearly, the previous position makes sense also when $p(\cdot) \equiv \text{const}$.

2.3. Well-known results. When dealing with p -Laplacean type problems, we shall often use the auxiliary vector fields $V_{s,t}: \mathbb{R}^{N \times n} \rightarrow \mathbb{R}^{N \times n}$, defined by

$$(2.9) \quad V_{s,t}(z) := (s^2 + |z|^2)^{(t-2)/4} z, \quad t \in (1, \infty) \text{ and } s \in [0, 1]$$

whenever $z \in \mathbb{R}^{N \times n}$. If $s = 0$ we shall simply write $V_{s,t} \equiv V_t$. A useful related inequality is contained in the following

$$(2.10) \quad |V_{s,t}(z_1) - V_{s,t}(z_2)| \approx (s^2 + |z_1|^2 + |z_2|^2)^{(t-2)/4} |z_1 - z_2|,$$

where the equivalence holds up to constants depending only on n, k, t . An important property which is usually related to such field is recorded in the following lemma.

Lemma 2.1. *Let $t > -1$, $s \in [0, 1]$ and $z_1, z_2 \in \mathbb{R}^{N \times n}$ be so that $s + |z_1| + |z_2| > 0$. Then*

$$\int_0^1 \left[s^2 + |z_1 + \lambda(z_2 - z_1)|^2 \right]^{\frac{t}{2}} d\lambda \sim (s^2 + |z_1|^2 + |z_2|^2)^{\frac{t}{2}},$$

with constants implicit in " \sim " depending only on n, N, t .

The next are a couple of simple inequalities which will be used several times throughout the paper. They are elementary, see e.g.: [8, 9, 47, 52].

Lemma 2.2. *The following inequalities hold true.*

- i. *For any $\varepsilon_0 > 0$, there exists a constant $c \equiv c(\varepsilon_0)$ such that for all $t \geq 0$, $l \geq m \geq 1$ there holds $|t^l - t^m| \leq c(l - m) \left(1 + t^{(1+\varepsilon_0)l} \right)$.*
- ii. *For $t \in (0, 1]$, consider the function $g_1(t) := t^{\tilde{c}t^\gamma}$, where \tilde{c} is an absolute real constant and $\gamma \in (0, 1]$. Then $\lim_{t \rightarrow 0} g_1(t) = 1$ and $\sup_{t \in (0, 1]} g_1(t) \leq c(\tilde{c}, \gamma)$. Via the substitution $t \mapsto t^{-1}$, we have an analogous property for the function $[1, \infty) \ni t \mapsto g_2(t) := t^{\tilde{c}t^{-\gamma}}$, for \tilde{c} and γ as before. Precisely there holds that $\lim_{t \rightarrow \infty} g_2(t) = 1$ and $\sup_{t \in [1, \infty)} g_2(t) \leq c(\tilde{c}, \gamma)$.*

We conclude this section by recalling the celebrated iteration lemma, [25].

Lemma 2.3. *Let $h: [\varrho, R_0] \rightarrow \mathbb{R}$ be a non-negative, bounded function and $0 < \theta < 1$, $0 \leq A$, $0 < \beta$. Assume that $h(r) \leq A(d-r)^{-\beta} + \theta h(d)$, for $\varrho \leq r < d \leq R_0$. Then $h(\varrho) \leq cA/(R_0 - \varrho)^{-\beta}$ holds, where $c \equiv c(\theta, \beta) > 0$.*

2.4. Extensions. In this section we shall borrow from [9] some useful lemmas concerning locally Lipschitz retractions. Such results were first introduced in [30] and intensively used in the literature for dealing with possibly non-homogeneous variational problems whose structure is a priori non-compatible with any kind of monotonicity formulae, [10, 36]. We refer to Remark 2.2 below for a quick discussion on this matter. We start with clarifying a key assumption in our paper, which is the concept of j -connectedness.

Definition 4. *Given an integer $j \geq 0$, a manifold \mathcal{M} is said to be j -connected if its first j homotopy groups vanish identically, that is $\pi_0(\mathcal{M}) = \pi_1(\mathcal{M}) = \dots = \pi_{j-1}(\mathcal{M}) = \pi_j(\mathcal{M}) = 0$.*

It is well-known that a compact manifold $\mathcal{M} \subset \mathbb{R}^N$ without boundary admits a tubular neighborhood $\mathcal{M} \subset \omega \subset \mathbb{R}^N$. Identifying \mathcal{M} with its image in \mathbb{R}^N , we say that a neighborhood ω of \mathcal{M} has the nearest point property if for every $x \in \omega$ there is a unique point $\Pi_{\mathcal{M}}(x) \in \mathcal{M}$ such that $\text{dist}(x, \mathcal{M}) = |x - \Pi_{\mathcal{M}}(x)|$. The map $\Pi_{\mathcal{M}}: \omega \rightarrow \mathcal{M}$ is called the retraction onto \mathcal{M} , we shall

refer to it also as "projector". Moreover, the regularity of \mathcal{M} influences the regularity of $\Pi_{\mathcal{M}}$ in the following way:

$$(2.11) \quad \mathcal{M} \text{ is } C^k\text{-regular for } k \geq 2 \Rightarrow \Pi_{\mathcal{M}} \in C^{k-1}(\omega, \mathcal{M}),$$

see [36] for a deeper discussion on this matter. It is important to stress that manifolds endowed with the relatively simple topology described by Definition 4 enjoy good properties in terms of retractions.

Lemma 2.4. *Let $\mathcal{M} \subset \mathbb{R}^N$ be a compact, j -connected submanifold for some integer $j \in \{0, \dots, N-2\}$ contained in an N -dimensional cube Q . Then there exists a closed $(N-j-2)$ -dimensional Lipschitz polyhedron $X \subset Q \setminus \mathcal{M}$ and a locally Lipschitz retraction $\psi: Q \setminus X \rightarrow \mathcal{M}$ such that for any $x \in Q \setminus X$, $|D\psi(x)| \leq c/\text{dist}(x, X)$ holds, for some positive $c \equiv c(N, j, \mathcal{M})$.*

Proof. We refer to [30, Lemma 6.1] for the original proof, or [36, Lemma 4.5] for a simplified version relying on some Lipschitz extensions of maps between Riemannian manifolds. \square

The next lemma allows modifying the image of a map while keeping under control boundary values and $p(\cdot)$ -energy, see also [9, Lemma 5].

Lemma 2.5. *Let \mathcal{M} be as in (2.6) and $U \subseteq B_1^+$ a subset with positive measure and piecewise C^1 -regular boundary. If $w \in W^{1,p(\cdot)}(U, \mathbb{R}^N) \cap L^\infty(U, \mathbb{R}^N)$ is so that its image lies in \mathcal{M} in a neighborhood of ∂U , then there exists $\tilde{w} \in w + \hat{C}_u^{p(\cdot)}(U, \mathcal{M})$ satisfying*

$$\int_U |D\tilde{w}|^{p(x)} dx \leq c \int_U |Dw|^{p(x)} dx,$$

where $c \equiv c(N, \mathcal{M}, \gamma_2)$.

Remark 2.2. When dealing with manifold constrained minima of the p -Laplacean energy it is customary to recover the fundamental Caccioppoli inequality by exploiting the so-called monotonicity formula, see [20–22, 41, 49–51]. This way cannot be used in our case. Even though it is possible to show a monotonicity formula for the $p(x)$ -energy, Lemma 4.2 below, see also [52, Lemma 4.1] or [9, Lemma 12], its proof crucially requires some corollaries of Gehring Lemma, which, in turn, is implied by Caccioppoli inequality, whose proof requires the monotonicity formula. Lemma 2.5 breaks this vicious circle giving the chance of deriving Caccioppoli inequality directly by minimality, as we will see in Section 3.1.

3. PARTIAL BOUNDARY REGULARITY

As mentioned in Section 2.2, to avoid unnecessary complications, we shall take $\Omega \equiv B_1^+$. In fact, since $\partial\Omega$ is C^2 -regular, given any $x_0 \in \partial\Omega$, there exists an open neighborhood B_{x_0} of x_0 and a change of variable $\Psi_0 \in C^2(\bar{B}_{x_0}, \mathbb{R}^n)$ so that in the new coordinates $y^i := \Psi_0^i(x)$ there holds that

$$\Psi_0(x_0) = 0, \quad \Psi_0(\bar{B}_{x_0} \cap \bar{\Omega}) = \bar{B}_1^+, \quad \Psi_0(\bar{B}_{x_0} \cap \partial\Omega) = \Gamma_1.$$

Moreover, there exists a positive constant $c_0 \equiv c_0(n, \partial\Omega)$ such that

$$0 < c_0^{-1} \leq \|D\Psi_0\|_{L^\infty(\bar{B}_{x_0} \cap \bar{\Omega})} \leq c_0 < \infty.$$

We stress that, being $\partial\Omega$ compact, the constant c_0 does not depend from x_0 . A straightforward computation shows that, if $u \in W^{1,p(\cdot)}(\Omega, \mathcal{M})$ solves (1.1), then the map $\tilde{u} := u \circ \Psi_0^{-1}$ solves an analogous problem still satisfying (2.2) and (2.3). Assumption (2.7) on the boundary condition is preserved as well: if $g \in W^{1,q}(\bar{\Omega}, \mathcal{M})$ then $\tilde{g} := g \circ \Psi_0^{-1} \in W^{1,q}(\bar{B}_1^+, \mathcal{M})$. We refer to [13, 30, 37] for more details on this matter. Therefore, keeping Definition 3 in mind, we shall study problem

$$(3.1) \quad C_g^{p(\cdot)}(B_1^+, \mathcal{M}) \ni w \mapsto \min \int_{B_1^+} k(x) |Dw|^{p(x)} dx,$$

with $k(\cdot)$ and $p(\cdot)$ as in (2.3)-(2.2) respectively and g as in (2.7).

3.1. Basic regularity results. We first fix a threshold radius $R_* \in (0, 1]$ so that

$$(3.2) \quad 0 < R_* \leq \min \left\{ 1, \left(\frac{\gamma_1^2}{4n[p]_{0,\alpha}} \right)^{\frac{1}{\alpha}}, \left(\frac{\gamma_1 q \left(1 - \frac{n}{q} \right)}{4n[p]_{0,\alpha}} \right)^{\frac{1}{\alpha}} \right\}$$

and choose a $R \in (0, R_*)$. Further restrictions on the size of R_* will be imposed in Section 3.2. An immediate consequence of (3.2) is that, given any half-ball B_R^+ and all balls $B_\varrho(x_0)$ with $x_0 \in B_R^+$ and $\varrho \in (0, R - |x_0|)$, there holds

$$(3.3) \quad \begin{cases} p_1^*(x_0, \varrho) > p_2(x_0, \varrho) \\ \frac{np_2(x_0, \varrho)}{q} \leq p_1(x_0, \varrho) \end{cases} \quad \text{for all } R \in (0, R_*], \quad \varrho \in (0, R - |x_0|),$$

which is, on the other hand, automatic when $p_1(x_0, \varrho) \geq n$. Obviously, in (3.3) we adopted the usual terminology

$$p^* := \begin{cases} \frac{np}{n-p} & \text{if } 1 < p < n \\ \text{any finite number larger than } p & \text{if } p \geq n. \end{cases}$$

Recall now that, if $B_\varrho(x_0) \Subset B_R^+$ and $w \in W^{1,p}(B_\varrho(x_0), \mathbb{R}^N)$ is such that $w \equiv 0$ on $U \subset B_\varrho(x_0)$ with $|U| > \hat{c}|B_\varrho(x_0)|$ for some positive, absolute \hat{c} , then Sobolev-Poincaré's inequality gives

$$(3.4) \quad \int_{B_\varrho(x_0)} |w/r|^p \, dx \leq c\varrho^{-n(p/p_*-1)} \left(\int_{B_\varrho(x_0)} |Dw|^{p_*} \, dx \right)^{\frac{p}{p_*}},$$

for $c \equiv c(n, N, p, \hat{c})$. Here $p_* := \max \left\{ 1, \frac{np}{n+p} \right\}$. We consider now an intrinsic version of [13, Theorem 2.4].

Proposition 3.1. *Let $U \subset \mathbb{R}^n$ be an open, bounded domain with piecewise C^1 -regular boundary and finite Ahlfors constants depending only from n . Let also $A \subset \bar{U}$ be a closed subset. Consider two non-negative functions $f_1 \in L^1(U)$ and $f_2 \in L^{1+\hat{\sigma}}(U)$ for some $\hat{\sigma} > 0$. With $\theta \in (0, 1)$, assume that there holds*

$$(3.5) \quad \int_{B_{\varrho/2}(x_0) \cap U} f_1 \, dx \leq b \left\{ \left(\int_{B_\varrho(x_0) \cap U} f_1^\theta \, dx \right)^{\frac{1}{\theta}} + \int_{B_\varrho(x_0) \cap U} f_2 \, dx \right\}$$

for almost all $x_0 \in U \setminus A$ with $B_\varrho(x_0) \cap A = \emptyset$ and a positive constant b . Set

$$d(x) := \frac{|B_{\text{dist}(x,A)}(x) \cap U|}{|U|} \quad \text{and} \quad \tilde{f}_1(x) := d(x)f_1(x).$$

Then there exists a positive threshold $\sigma_g \equiv \sigma_g(b, \theta, \hat{\sigma}) \in (0, \hat{\sigma})$ such that

$$\left(\int_U \tilde{f}_1^{1+\sigma} \, dx \right)^{\frac{1}{1+\sigma}} \leq c(n, \theta, b, \hat{\sigma}) \left\{ \left(\int_U f_1 \, dx \right) + \left(\int_U f_2^{1+\sigma} \, dx \right)^{\frac{1}{1+\sigma}} \right\}$$

for all $\sigma \in [0, \sigma_g]$.

Proof. The proof is essentially the same as the one in [13] with minor changes due to the fact that, in our case, (3.5) involves the whole integrand; see also [25, Lemma 6.2]. \square

As a consequence of Proposition 3.1, we derive some higher integrability results for solutions to problem (1.1).

Lemma 3.1. *Under assumptions (2.2), (2.3), (2.6) and (2.7), let $u \in W^{1,p(\cdot)}(B_R^+, \mathcal{M})$ be a solution of problem (3.1). Then, for $x_0 \in \bar{B}_R^+$, with $R \in (0, R_*]$, R_* as in (3.2) and $0 < \varrho < R - |x_0|$, there exists a positive threshold $\sigma_g \equiv \sigma_g(\mathbf{data}_{p(\cdot)}, q) \in (0, \frac{q}{\gamma_2} - 1)$ such that for all $\sigma \in (0, \sigma_g)$ there holds that:*

$$(3.6) \quad \left(\int_{B_{\varrho/2}(x_0) \cap B_R^+} (1 + |Du|^2)^{\frac{p(x)(1+\sigma)}{2}} dx \right)^{\frac{1}{1+\sigma}} \leq c \left[\int_{B_{\varrho}(x_0) \cap B_R^+} (1 + |Du|^2)^{\frac{p(x)}{2}} dx + \left(\int_{B_{\varrho}(x_0) \cap B_R^+} |Dg|^{p(x)(1+\sigma)} dx \right)^{\frac{1}{1+\sigma}} \right].$$

for $c \equiv c(\mathbf{data}_{p(\cdot)}, q)$. If $B_{\varrho}(x_0) \Subset B_R^+$ then, there exists a positive threshold $\sigma'_g \equiv \sigma'_g(\mathbf{data}_{p(\cdot)}) > 0$ so that

$$(3.7) \quad \left(\int_{B_{\varrho/2}(x_0)} (1 + |Du|^2)^{\frac{p(x)(1+\sigma)}{2}} dx \right)^{\frac{1}{1+\sigma}} \leq c \int_{B_{\varrho}(x_0)} (1 + |Du|^2)^{\frac{p(x)}{2}} dx,$$

for all $\sigma \in (0, \sigma'_g)$ with $c \equiv c(\mathbf{data}_{p(\cdot)})$. In particular,

$$(3.8) \quad |Du|^{p(\cdot)(1+\sigma)} \in L^1(B_R^+) \quad \text{for all } \sigma \in \left[0, \min \{ \sigma_g, \sigma'_g \} \right).$$

Proof. We take $x_0 \in \bar{B}_R^+$, $0 < \varrho < R - |x_0|$ and distinguish two cases: $x_0^n \leq \frac{3\varrho}{4}$ and $x_0^n > \frac{3\varrho}{4}$.

Case 1: $x_0^n \leq \frac{3\varrho}{4}$. We fix parameters $\frac{\varrho}{2} < \tau_1 < \tau_2 \leq \varrho$ and a cut-off function $\eta \in C_c^1(B_{\tau_2}(x_0))$ with the following specifics

$$(3.9) \quad \mathbb{1}_{B_{\tau_1}(x_0)} \leq \eta \leq \mathbb{1}_{B_{\tau_2}(x_0)} \quad \text{and} \quad |D\eta| \leq \frac{4}{\tau_2 - \tau_1}.$$

Notice that in this case the intersection $B_{\tau_2}(x_0) \cap \Gamma_R$ can be non-empty and the map $w := u - \eta(u - g)$ agrees with u in the sense of traces on $\partial(B_{\tau_2}(x_0) \cap B_R^+)$. This means that we can use Lemma 2.5 to recover a map $\tilde{w} \in \hat{C}_u^{p(\cdot)}(B_{\tau_2}(x_0) \cap B_R^+)$ satisfying the energy inequality (3.9) and so that

$$\begin{aligned} \int_{B_{\tau_2}(x_0) \cap B_R^+} |Du|^{p(x)} dx &\leq \lambda^{-1} \int_{B_{\tau_2}(x_0) \cap B_R^+} k(x) |Du|^{p(x)} dx \\ &\leq \lambda^{-1} \int_{B_{\tau_2}(x_0) \cap B_R^+} k(x) |D\tilde{w}|^{p(x)} dx \\ &\leq \frac{\Lambda}{\lambda} \int_{B_{\tau_2}(x_0) \cap B_R^+} |D\tilde{w}|^{p(x)} dx \leq c \int_{(B_{\tau_2}(x_0) \setminus B_{\tau_1}(x_0)) \cap B_R^+} |Du|^{p(x)} dx \\ &\quad + c \int_{B_{\tau_2}(x_0) \cap B_R^+} \left[|Dg|^{p(x)} + \left| \frac{u-g}{\tau_2 - \tau_1} \right|^{p(x)} \right] dx, \end{aligned}$$

with $c \equiv c(N, \lambda, \Lambda, \gamma_2, \mathcal{M})$. Once the inequality on the previous display is available, we can use Widmann's hole filling technique, Lemma 2.3 and Lemma 2.2 (ii) to end up with

$$(3.10) \quad \begin{aligned} \int_{B_{\varrho/2}(x_0) \cap B_R^+} |Du|^{p(x)} dx &\leq c \int_{B_{\varrho}(x_0) \cap B_R^+} |Dg|^{p(x)} dx + c\varrho^{-p_2(\varrho)} \int_{B_{\varrho}(x_0) \cap B_R^+} |u-g|^{p(x)} dx \\ &\leq c \int_{B_{\varrho}(x_0) \cap B_R^+} |Dg|^{p(x)} dx + c \int_{B_{\varrho}(x_0) \cap B_R^+} \left| \frac{u-g}{\varrho} \right|^{p(x)} dx \\ &\leq c \int_{B_{\varrho}(x_0) \cap B_R^+} |Dg|^{p(x)} dx + c \int_{B_{\varrho}(x_0) \cap B_R^+} \left| \frac{u-g}{\varrho} \right|^{p_1(\varrho)} dx + c|B_{\varrho}(x_0) \cap B_R^+|, \end{aligned}$$

where $c \equiv c(N, \lambda, \Lambda, \gamma_2, m)$. Now we extend $u = g$ in $B_\varrho(x_0) \setminus B_R^+$, notice that condition $x_0^n \leq 3\varrho/4$ implies that $|B_\varrho(x_0) \setminus B_R^+| \geq c(n)|B_\varrho(x_0)|$ and use (3.4) to bound

$$\begin{aligned} \int_{B_\varrho(x_0) \cap B_R^+} \left| \frac{u-g}{\varrho} \right|^{p_1(\varrho)} dx &\leq c \left(\int_{B_\varrho(x_0) \cap B_R^+} |Du - Dg|^{(p_1(\varrho))^*} dx \right)^{\frac{p_1(\varrho)}{(p_1(\varrho))^*}} \\ &\leq c \left(\int_{B_\varrho(x_0) \cap B_R^+} (1 + |Du|^2)^{\frac{p(x)(p_1(\varrho))^*}{2p_1(\varrho)}} dx \right)^{\frac{p_1(\varrho)}{(p_1(\varrho))^*}} \\ &\quad + c \left(\int_{B_\varrho(x_0) \cap B_R^+} |Dg|^{\frac{p(x)(p_1(\varrho))^*}{p_1(\varrho)}} dx \right)^{\frac{p_1(\varrho)}{(p_1(\varrho))^*}}, \end{aligned}$$

for $c \equiv c(n, \gamma_1, \gamma_2)$. Merging the content of the two previous displays we obtain

$$\begin{aligned} \int_{B_{\varrho/2}(x_0) \cap B_R^+} (1 + |Du|^2)^{\frac{p(x)}{2}} dx &\leq c \int_{B_\varrho(x_0) \cap B_R^+} |Dg|^{p(x)} dx \\ (3.11) \quad &+ c \left(\int_{B_\varrho(x_0) \cap B_R^+} (1 + |Du|^2)^{\frac{p(x)}{2} \cdot \frac{(p_1(\varrho))^*}{p_1(\varrho)}} dx \right)^{\frac{p_1(\varrho)}{(p_1(\varrho))^*}}, \end{aligned}$$

where $c \equiv c(n, N, \lambda, \Lambda, \gamma_1, \gamma_2, m)$.

Case 2: $x_0^n > \frac{3\varrho}{4}$. In this case, we see that $B_{\frac{3\varrho}{4}} \subseteq B_R^+$, so, as in [9, Lemma 9] we recover

$$\begin{aligned} \int_{B_{\varrho/2}(x_0)} (1 + |Du|^2)^{\frac{p(x)}{2}} dx &\leq c \left(\int_{B_{3\varrho/4}(x_0)} (1 + |Du|^2)^{\frac{p(x)}{2} \cdot \frac{(p_1(\varrho))^*}{p_1(\varrho)}} dx \right)^{\frac{p_1(\varrho)}{(p_1(\varrho))^*}} \\ (3.12) \quad &\leq c \left(\int_{B_\varrho(x_0)} (1 + |Du|^2)^{\frac{p(x)}{2} \cdot \frac{(p_1(\varrho))^*}{p_1(\varrho)}} dx \right)^{\frac{p_1(\varrho)}{(p_1(\varrho))^*}}, \end{aligned}$$

for $c \equiv c(n, N, \lambda, \Lambda, \gamma_1, \gamma_2, m)$. Once (3.11)-(3.12) are available, we can apply Proposition 3.1 with $U \equiv B_\varrho(x_0) \cap B_R^+$ and $A \equiv \partial B_\varrho(x_0) \cap B_R^+$ to conclude with (3.6)-(3.7).

Combining (3.6), (3.7) and a standard covering argument, we obtain (3.8) and the proof is complete. \square

Remark 3.1. Since $Dg \in L^q(B_1^+, \mathbb{R}^{N \times n})$, with Hölder inequality we can rearrange (3.6) as follows:

$$\begin{aligned} &\left(\int_{B_{\varrho/2}(x_0) \cap B_R^+} (1 + |Du|^2)^{\frac{p(x)(1+\sigma)}{2}} dx \right)^{\frac{1}{1+\sigma}} \\ (3.13) \quad &\leq c \left[\int_{B_\varrho(x_0) \cap B_R^+} (1 + |Du|^2)^{\frac{p(x)}{2}} dx + \left(\int_{B_\varrho(x_0) \cap B_R^+} |Dg|^q dx \right)^{\frac{p_2(R)}{q}} \right], \end{aligned}$$

for $c \equiv c(\text{data}_{p(\cdot), q})$.

Let us point out a particularly helpful inequality contained in the proof of Lemma 3.1.

Corollary 3.1. *Under assumptions (2.2), (2.3), (2.6) and (2.7), let $u \in W^{1,p(\cdot)}(B_1^+, m)$ be a solution of problem (3.1). Then for any half-ball $B_R \subset \bar{B}_1^+$ and all balls $B_\varrho(x_0)$ with $x_0 \in B_R^+$, $\varrho \in (0, R - |x_0|)$, $R \in (0, R_*]$ and R_* as in (3.2), there holds that*

$$(3.14) \quad \int_{B_{\varrho/2}(x_0) \cap B_R^+} |Du|^{p(x)} dx \leq c \int_{B_\varrho(x_0) \cap B_R^+} \left[\left| \frac{u-g}{r} \right|^{p(x)} + |Dg|^{p(x)} \right] dx$$

with $c \equiv c(\mathbf{data}_{p(\cdot)})$. In case $B_\varrho(x_0) \Subset B_1^+$, the inequality

$$(3.15) \quad \int_{B_{\varrho/2}(x_0) \cap B_R^+} |Du|^{p(x)} dx \leq c \int_{B_\varrho(x_0) \cap B_R^+} \left[\left| \frac{u - (u)_\varrho}{r} \right|^{p(x)} \right] dx,$$

for $c \equiv c(\mathbf{data}_{p(\cdot)})$. Moreover, the following inequalities are satisfied:

$$(3.16) \quad \int_{B_{\varrho/4}(x_0) \cap B_R^+} |Du|^{p(x)} dx \leq c \varrho^{-p_2(\varrho)} \quad \text{and} \quad \int_{B_{\varrho/4}(x_0) \cap B_R^+} |Du|^{p(x)(1+\sigma)} dx \leq c \varrho^{-p_2(\varrho)(1+\sigma)}$$

with $c \equiv c(n, N, \mathcal{M}, \gamma_1, \gamma_2, q, \|Dg\|_{L^q(B_1^+)})$ and for all $\sigma \in [0, \min\{\sigma_g, \frac{q}{n} - 1\}]$, where σ_g is the same higher integrability threshold appearing in Lemma 3.1.

Proof. Inequality (3.14) is the same as (3.10) in the proof of Lemma 3.1, while the proof of (3.15) is contained in [9, Lemma 8]. To prove (3.16) we only need to notice that by (2.7)₂ it immediately follows that

$$(3.17) \quad \begin{aligned} \varrho^{p_2(\varrho)} \int_{B_\varrho(x_0) \cap B_R^+} |Dg|^{p(x)(1+\sigma)} dx &\leq c \left[\varrho^{p_2(\varrho)} + \varrho^{p_2(\varrho)} \left(\int_{B_\varrho(x_0) \cap B_R^+} |Dg|^{p_2(\varrho)(1+\sigma)} dx \right) \right] \\ &\leq c \left[\varrho^{p_2(\varrho)} + \varrho^{p_2(\varrho)(1-\frac{n(1+\sigma)}{q})} \left(1 + \|Dg\|_{L^q(B_1^+)}^{2\gamma_2} \right) \right] \leq c(n, \gamma_2, \|Dg\|_{L^q(B_1^+)}). \end{aligned}$$

Using this information together with (3.14) and (2.6)₁, we obtain (3.16)₁. Combining (3.6)-(3.7) with (3.16)₁ and (3.17) we get (3.16)₂ and the proof is complete. \square

By Proposition 3.1 with $A \equiv \emptyset$, we can prove a globally higher integrability result for p -harmonic functions, see e.g. [9, Lemma 10] or [13, Lemma 3.3].

Lemma 3.2. *Let $R \in (0, 1]$, $x_0 \in \Gamma_R$ and $\varrho \in (0, R - |x_0|)$. Assume (2.2)₂, (2.3)₂ and (2.6), take $p \in [\gamma_1, \gamma_2]$ and $f \in W^{1,p}(\bar{B}_\varrho^+(x_0) \cap \bar{B}_R^+, \mathcal{M})$ so that $|Df|^p \in L^{1+\delta}(\bar{B}_\varrho^+(x_0) \cap \bar{B}_R^+)$. If $v \in W^{1,p}(B_\varrho^+(x_0) \cap B_R^+, \mathcal{M})$ is a solution of the Dirichlet problem*

$$(3.18) \quad \hat{\mathcal{C}}_f^p(B_\varrho^+(x_0) \cap B_R^+, \mathcal{M}) \ni w \mapsto \min \int_{B_\varrho^+(x_0) \cap B_R^+} k(x) |Dw|^p dx,$$

then there exists a positive threshold $\delta_g \equiv \delta_g(n, N, \mathcal{M}, \gamma_1, \gamma_2, \lambda, \Lambda) \in (0, \hat{\delta})$ so that

$$(3.19) \quad \left(\int_{B_\varrho^+(x_0) \cap B_R^+} |Dv|^{p(1+\delta)} dx \right)^{\frac{1}{1+\delta}} \leq c \left\{ \int_{B_\varrho(x_0) \cap B_R^+} |Dv|^p dx + \left(\int_{B_\varrho(x_0) \cap B_R^+} |Df|^{p(1+\delta)} dx \right)^{\frac{1}{1+\delta}} \right\}$$

for all $\delta \in [0, \delta_g)$. In (3.19), $c \equiv c(n, N, \mathcal{M}, \gamma_1, \gamma_2, \lambda, \Lambda)$.

3.2. Proof of Theorem 1. The proof of Theorem 1 relies on the following result.

Proposition 3.2. *Under assumptions (2.1), (2.2), (2.3) and (2.6), let $u \in W^{1,p(\cdot)}(B_1^+, \mathcal{M})$ be a solution of problem (3.1) with boundary datum $g: \bar{B}_1^+ \rightarrow \mathcal{M}$ satisfying (2.7). Then, there exist a threshold radius $R_* \equiv R_*(\mathbf{data}) \in (0, 1]$ and a smallness parameter $\varepsilon \equiv \varepsilon(\mathbf{data}) \in (0, 1]$ such that if*

$$(3.20) \quad \left(\varrho^{p_2(x_0, \varrho) - n} \int_{B_\varrho(x_0) \cap B_R^+} |Du|^{p_2(x_0, \varrho)} dx \right)^{\frac{1}{p_2(x_0, \varrho)}} + \left(\varrho^{q-n} \int_{B_\varrho(x_0) \cap B_R^+} |Dg|^q dx \right)^{\frac{1}{q}} < \varepsilon,$$

for some $R \in (0, R_*)$, $x_0 \in B_R^+$ and $\varrho \in (0, R - |x_0|)$, then

$$u \in C_{loc}^{0, 1 - \frac{n}{q}} \left((B_\varrho(x_0) \cap \bar{B}_R^+) \setminus \Sigma_0(u, B_\varrho(x_0) \cap \bar{B}_R^+), \mathcal{M} \right),$$

where $\Sigma_0(u, B_\varrho(x_0) \cap \bar{B}_R^+) \subset \bar{B}_R^+$ is a closed subset with $\dim_{\mathcal{H}}(\Sigma_0(u, B_\varrho(x_0) \cap \bar{B}_R^+)) < n - \gamma_1$.

Proof. For the sake of simplicity, we split the proof into six steps.

Step 1: Setting a threshold radius. As mentioned in Section 3.1, there is no loss of generality in reducing the size of the half ball we are working on. Precisely, in addition to (3.2), we choose a radius $R \in (0, R_*]$, where now it is

$$(3.21) \quad 0 < R_* < \min \left\{ 1, \left[\frac{\gamma_1^2}{4n[p]_{0,\alpha}} \right]^{\frac{1}{\alpha}}, \left(\frac{\gamma_1 q \left(1 - \frac{n}{q}\right)}{4n[p]_{0,\alpha}} \right)^{\frac{1}{\alpha}}, \left(\frac{\sigma_0 \gamma_1}{2[p]_{0,\alpha}(2 + \sigma_0)} \right)^{\frac{1}{\alpha}} \right\},$$

for $\sigma_0 \in (0, 1)$ defined as

$$(3.22) \quad \sigma_0 := \min \left\{ \frac{1}{4}, \frac{\sigma'_g}{2}, \frac{\sigma_g}{2}, \frac{2}{\gamma_2 - 1}, \frac{\alpha}{\gamma_2}, \frac{q - \gamma_2}{\gamma_2} \right\}.$$

In (3.22), σ_g and σ'_g are the higher integrability thresholds appearing Lemma 3.1, therefore, given an half-ball $B_R^+ \subset B_1^+$, by (3.8) there holds that

$$(3.23) \quad |Du|^{p(\cdot)(1+\sigma)} \in L^1(B_R^+) \quad \text{for all } \sigma \in [0, \sigma_0].$$

Moreover, in addition to (3.3), another straightforward consequence of the restriction imposed in (3.21) yields that

$$(3.24) \quad p_2(x_0, \varrho) < p_2(x_0, \varrho) \left(1 + \frac{\sigma_0}{2} \right) \leq (1 + \sigma_0)p_1(x_0, \varrho),$$

whenever $x_0 \in B_R^+$ and $\varrho \in (0, R - |x_0|)$. Hence, combining (3.23) and (3.24) we can conclude that

$$(3.25) \quad |Du|^{p_2(x_0, \varrho)} \in L^1(B_\varrho(x_0) \cap B_R^+).$$

Let us stress that by continuity, for any point $\bar{x} \in \bar{B}_R^+$ for which $p(\bar{x}) \geq n$, we can find a small ball $B_{\varrho_{\bar{x}}}(\bar{x}) \subset \bar{B}_R^+$ so that $p(x) > n - \frac{\sigma_0}{2}$ for all $x \in B_{\varrho_{\bar{x}}}(\bar{x})$. Combining this information with (3.7) and observing that, by (3.22)

$$\left(n - \frac{\sigma_0}{2} \right) (1 + \sigma_0) > n + \frac{\sigma_0}{4},$$

with Sobolev-Morrey embedding theorem we obtain that $u \in C^{0, \frac{\sigma_0}{4n+\sigma_0}}(B_{\varrho_{\bar{x}}/2}(\bar{x}) \cap B_1^+, \mathcal{M})$. Therefore, for the rest of the paper, we shall assume that $\gamma_2 < n$. Moreover, since from now on we work on sets of the type $B_\varrho(x_0) \cap B_R^+$ with $x_0 \in B_R^+$ and $\varrho \in (0, R - |x_0|)$, we shall simplify the notation in (2.5) as follows: $p_1(x_0, \varrho) \equiv p_1(\varrho)$ and $p_2(x_0, \varrho) \equiv p_2(\varrho)$.

Step 2: Comparison, first time. Let $u \in W^{1,p(\cdot)}(B_1^+, \mathcal{M})$ be a solution to the minimization problem (3.1) with (2.7) in force. We introduce the extensions

$$(3.26) \quad \tilde{u}(x) := \begin{cases} u(x', x^n) - g(x', x^n) & \text{if } x^n \geq 0 \\ -(u(x', -x^n) - g(x', -x^n)) & \text{if } x^n < 0. \end{cases}$$

Since $\text{tr}_{\Gamma_1}(u) = \text{tr}_{\Gamma_1}(g)$, it easily follows that $\tilde{u} \in W^{1,p(\cdot)}(B_1, \mathbb{R}^N)$ and, by (3.23), for all $B_\varrho(x_0) \subseteq B_R \subset B_{R_*}$ with R_* as in (3.21) there holds that

$$(3.27) \quad \int_{B_\varrho(x_0)} |D\tilde{u}|^{p_2(R)} \, dx \leq c \int_{B_\varrho(x_0) \cap B_R^+} [|Du|^{p_2(R)} + |Dg|^{p_2(R)}] \, dx$$

with $c \equiv c(\gamma_1, \gamma_2)$. Before going on we define the following quantities:

$$\phi(x_0, \varrho, p) := \left(\varrho^p \int_{B_\varrho(x_0)} (1 + |D\tilde{u}|^2)^{p/2} \, dx \right)^{\frac{1}{p}};$$

$$\begin{aligned}\phi^+(x_0, \varrho, p) &:= \left(\varrho^p \int_{B_\varrho(x_0) \cap B_R^+} (1 + |Du|^2)^{p/2} \, dx \right)^{\frac{1}{p}}; \\ \psi(x_0, \varrho) &:= \phi(x_0, \varrho, p_2(\varrho)), \quad \psi^+(x_0, \varrho) := \phi^+(x_0, \varrho, p_2(\varrho)); \\ \chi^+(x_0, \varrho) &:= \psi^+(x_0, \varrho) + \left(\varrho^{q-n} \int_{B_\varrho(x_0) \cap B_R^+} |Dg|^q \, dx \right)^{\frac{1}{q}},\end{aligned}$$

where x_0 , ϱ and R satisfy the usual relation $R \in (0, R_*]$, $x_0 \in B_R^+$ and $\varrho \in (0, R - |x_0|)$. In the definition of $\psi(x_0, \varrho)$, $p_2(\varrho)$ is defined as in (2.5). We shall start our analysis by considering a point $x_0 \in \Gamma_R$ and imposing (3.20) on $B_\varrho(x_0) \cap B_R^+ \equiv B_\varrho^+(x_0)$, which, with the terminology introduced above reads as

$$(3.28) \quad \chi^+(x_0, \varrho) < \varepsilon,$$

where $\varepsilon \in (0, 1)$ is a small parameter whose size will be suitably reduced along the proof. Notice that, as done in the case of (3.27), for all balls $B_\varrho(x_0) \subset B_R$, by Hölder inequality we have

$$\begin{aligned}(3.29) \quad \chi^+(x_0, \varrho) &\leq c' \left[\psi(x_0, \varrho) + \left(\varrho^{p_2(\varrho)-n} \int_{B_\varrho(x_0) \cap B_R^+} |Dg|^{p_2(\varrho)} \, dx \right)^{\frac{1}{p_2(\varrho)}} \right] \\ &\quad + c' \left(\varrho^{q-n} \int_{B_\varrho(x_0) \cap B_R^+} |Dg|^q \, dx \right)^{\frac{1}{q}} \\ &\leq c' \left[\psi(x_0, \varrho) + \left(\varrho^{q-n} \int_{B_\varrho(x_0) \cap B_R^+} |Dg|^q \, dx \right)^{\frac{1}{q}} \right],\end{aligned}$$

for $c' \equiv c'(n, \gamma_1, \gamma_2, q)$. Now we compare u to a solution $v \in W^{1, p_2(\varrho)}(B_{\varrho/2}^+(x_0), \mathcal{M})$ of the Dirichlet problem

$$(3.30) \quad \hat{C}_u^{p_2(\varrho)}(B_{\varrho/2}^+(x_0), \mathcal{M}) \ni w \mapsto \min \int_{B_{\varrho/2}^+(x_0)} k(x) |Dw|^{p_2(\varrho)} \, dx.$$

Such a solution exists, given that by (3.25), class $\hat{C}_u^{p_2(\varrho)}(B_{\varrho/2}^+(x_0), \mathcal{M})$ is non-empty. The minimality of v in class $\hat{C}_u^{p_2(\varrho)}(B_{\varrho/2}^+(x_0), \mathcal{M})$ yields that it satisfies the Euler-Lagrange equation

$$(3.31) \quad 0 = \int_{B_{\varrho/2}^+(x_0)} k(x) p_2(\varrho) |Dv|^{p_2(\varrho)-2} [Dv \cdot D\varphi - A_v(Dv, Dv)\varphi] \, dx,$$

for any $\varphi \in W_0^{1, p_2(\varrho)}(B_{\varrho/2}^+(x_0), \mathbb{R}^N) \cap L^\infty(B_{\varrho/2}^+(x_0), \mathbb{R}^N)$, where, for $y \in \mathcal{M}$, $A_y : T_y \mathcal{M} \times T_y \mathcal{M} \rightarrow (T_y \mathcal{M})^\perp$ denotes the second fundamental form of \mathcal{M} . In particular, by tangentiality,

$$(3.32) \quad \nabla^2 \Pi(v)(Dv, Dv) = -A_v(Dv, Dv) \quad \text{and} \quad |A_v(Dv, Dv)| \leq c_{\mathcal{M}} |Dv|^2,$$

where $c_{\mathcal{M}}$ depends only on the geometry of \mathcal{M} , see [51, Appendix to Chapter 2]. Let us quantify the $L^{p_2(\varrho)}$ -distance between Du and Dv . We first notice that, by (3.25), the map $\varphi := u - v$ is admissible as a test in (3.31), thus exploiting the monotonicity properties of the integrand in (3.30), (2.10) and Lemma 2.1 we obtain

$$\begin{aligned}(3.33) \quad &\int_{B_{\varrho/2}^+(x_0)} |V_{p_2(\varrho)}(Du) - V_{p_2(\varrho)}(Dv)|^2 \, dx \\ &\leq c \int_{B_{\varrho/2}^+(x_0)} k(x) [|Du|^{p_2(\varrho)} - |Dv|^{p_2(\varrho)}] \, dx + c \int_{B_{\varrho/2}^+(x_0)} |Dv|^{p_2(\varrho)} |u - v| \, dx\end{aligned}$$

where $c \equiv c(n, N, \gamma_1, \gamma_2, \lambda, \Lambda, \mathcal{M})$. Let us estimate the two quantities appearing on the right-hand side of (3.33). Notice that, being v a solution of (3.30), it satisfies the assumptions of Lemma

3.2 with $p = p_2(\varrho)$, $f = u$ and $\hat{\delta} = \frac{\sigma_0}{2}$, therefore, choosing any $\sigma' \in (0, \min\{\delta_g, \hat{\delta}\})$, by Hölder inequality we control:

$$\begin{aligned} \int_{B_{\varrho/2}^+(x_0)} |Dv|^{p_2(\varrho)} |u - v| \, dx &\leq c \varrho^n \left(\int_{B_{\varrho/2}^+(x_0)} |Dv|^{p_2(\varrho)(1+\sigma')} \, dx \right)^{\frac{1}{1+\sigma'}} \\ &\cdot \left(\int_{B_{\varrho/2}^+(x_0)} |u - v|^{\frac{1+\sigma'}{\sigma'}} \, dx \right)^{\frac{\sigma'}{1+\sigma'}} =: c(n) \varrho^n [(I) \cdot (II)]. \end{aligned}$$

By (3.6), (3.19), (3.24), the minimality of v in class $\hat{C}_u^{p_2(\varrho)}(B_{\varrho/2}^+(x_0), m)$, Hölder inequality, (3.28) and Lemma 2.2 (ii.) we bound

$$\begin{aligned} (I) &\leq c \left(\int_{B_{\varrho/2}^+(x_0)} |Du|^{p_2(\varrho)(1+\sigma')} \, dx \right)^{\frac{1}{1+\sigma'}} \\ &\leq c \left(\int_{B_{\varrho/2}^+(x_0)} |Du|^{p_1(\varrho)(1+\sigma_0)} \, dx \right)^{\frac{p_2(\varrho)}{p_1(\varrho)(1+\sigma_0)}} \\ &\leq c \left[\int_{B_{\varrho}^+(x_0)} (1 + |Du|^2)^{\frac{p(x)}{2}} \, dx + \left(\int_{B_{\varrho}^+(x_0)} |Dg|^{p(x)(1+\sigma_0)} \, dx \right)^{\frac{1}{1+\sigma_0}} \right]^{\frac{p_2(\varrho)}{p_1(\varrho)}} \\ &\leq c \left[\int_{B_{\varrho}^+(x_0)} (1 + |Du|^2)^{\frac{p(x)}{2}} \, dx + \left(\int_{B_{\varrho}^+(x_0)} |Dg|^q \, dx \right)^{\frac{p_2(\varrho)}{q}} \right]^{\frac{p_2(\varrho)}{p_1(\varrho)}} \\ &\leq c \varepsilon^{\frac{p_2(\varrho)-p_1(\varrho)}{p_1(\varrho)}} \left[\int_{B_{\varrho}^+(x_0)} (1 + |Du|^2)^{\frac{p_2(\varrho)}{2}} \, dx + \left(\int_{B_{\varrho}^+(x_0)} |Dg|^q \, dx \right)^{\frac{p_2(\varrho)}{q}} \right] \end{aligned}$$

with $c \equiv c(\text{data}_{p(\cdot)})$. With Poincaré inequality, (3.22), the minimality of v in class $\hat{C}_u^{p_2(\varrho)}(B_{\varrho/2}^+(x_0), m)$ and (3.28) we get

$$\begin{aligned} (II) &\leq c \left(\varrho^{p_2(\varrho)} \int_{B_{\varrho/2}^+(x_0)} |Du - Dv|^{p_2(\varrho)} \, dx \right)^{\frac{\sigma'}{1+\sigma'}} \\ &\leq c \left(\varrho^{p_2(\varrho)-n} \int_{B_{\varrho}^+(x_0)} (1 + |Du|^2)^{\frac{p_2(\varrho)}{2}} \, dx \right)^{\frac{\sigma'}{1+\sigma'}} \leq c \varepsilon^{\frac{\gamma_1 \sigma'}{1+\sigma'}}, \end{aligned}$$

where $c \equiv c(n, m, \gamma_1, \gamma_2, \lambda, \Lambda)$. Finally, by (3.13), (3.19), Lemma 2.2 (i.) with $\varepsilon_0 = \sigma'$, the minimality of v in class $\hat{C}_u^{p_2(\varrho)}(B_{\varrho/2}^+(x_0), m)$, (3.22) and (3.28) we have

$$\begin{aligned} \int_{B_{\varrho/2}^+(x_0)} k(x) \left[|Du|^{p_2(\varrho)} - |Dv|^{p_2(\varrho)} \right] \, dx &\leq \int_{B_{\varrho/2}^+(x_0)} k(x) \left| |Du|^{p_2(\varrho)} - |Du|^{p(x)} \right| \, dx \\ &+ \int_{B_{\varrho/2}^+(x_0)} k(x) \left| |Dv|^{p(x)} - |Dv|^{p_2(\varrho)} \right| \, dx \\ &\leq c \varrho^{n+\alpha} \left[\int_{B_{\varrho/2}^+(x_0)} (1 + |Du|^2)^{\frac{p_2(\varrho)}{2}(1+\sigma')} \, dx + \int_{B_{\varrho/2}^+(x_0)} (1 + |Dv|^2)^{\frac{p_2(\varrho)}{2}(1+\sigma')} \, dx \right] \end{aligned}$$

$$\begin{aligned}
&\leq c\varrho^{n+\alpha} \left(\int_{B_{\varrho/2}^+(x_0)} (1 + |Du|^2)^{\frac{p_1(\varrho)(1+\sigma_0)}{2}} dx \right) \\
&\leq c\varrho^{n+\alpha} \left[\left(\int_{B_{\varrho}^+(x_0)} (1 + |Du|^2)^{\frac{p_2(\varrho)}{2}} dx \right) + \left(\int_{B_{\varrho}^+(x_0)} |Dg|^q dx \right)^{\frac{p_2(\varrho)}{q}} \right]^{1+\sigma_0} \\
&\leq c\varrho^{n+\alpha-\sigma_0 p_2(\varrho)} \left[\varrho^{p_2(\varrho)-n} \int_{B_{\varrho}^+(x_0)} (1 + |Du|^2)^{\frac{p_2(\varrho)}{2}} dx + \left(\varrho^{q-n} \int_{B_{\varrho}^+(x_0)} |Dg|^q dx \right)^{\frac{p_2(\varrho)}{q}} \right]^{\sigma_0} \\
&\quad \cdot \left[\int_{B_{\varrho}^+(x_0)} (1 + |Du|^2)^{\frac{p_2(\varrho)}{2}} dx + \left(\int_{B_{\varrho}^+(x_0)} |Dg|^q dx \right)^{\frac{p_2(\varrho)}{q}} \right] \\
&\leq c\varepsilon^{\sigma_0 \gamma_1} \left[\int_{B_{\varrho}^+(x_0)} (1 + |Du|^2)^{\frac{p_2(\varrho)}{2}} dx + \varrho^{n(1-\frac{p_2(\varrho)}{q})} \left(\int_{B_{\varrho}^+(x_0)} |Dg|^q dx \right)^{\frac{p_2(\varrho)}{q}} \right],
\end{aligned}$$

where $c \equiv c(\mathbf{data}_{p(\cdot)})$. Merging the content of all the previous displays we end up with

$$\begin{aligned}
&\int_{B_{\varrho/2}^+(x_0)} |V_{p_2(\varrho)}(Du) - V_{p_2(\varrho)}(Dv)|^2 dx \\
(3.34) \quad &\leq c\varepsilon^{\frac{\sigma' \gamma_1}{1+\sigma'}} \left[\int_{B_{\varrho}^+(x_0)} (1 + |Du|^2)^{\frac{p_2(\varrho)}{2}} dx + \varrho^{n(1-\frac{p_2(\varrho)}{q})} \left(\int_{B_{\varrho/2}^+(x_0)} |Dg|^q dx \right)^{\frac{p_2(\varrho)}{q}} \right],
\end{aligned}$$

where $c \equiv c(\mathbf{data}_{p(\cdot)})$. If $p_2(\varrho) \geq 2$, by (2.10) and (3.34) we directly obtain that

$$\begin{aligned}
&\int_{B_{\varrho/2}^+(x_0)} |Du - Dv|^{p_2(\varrho)} dx \leq \int_{B_{\varrho/2}^+(x_0)} |V_{p_2(\varrho)}(Du) - V_{p_2(\varrho)}(Dv)|^2 dx \\
&\leq c\varepsilon^{\frac{\sigma' \gamma_1}{1+\sigma'}} \left[\int_{B_{\varrho}^+(x_0)} (1 + |Du|^2)^{\frac{p_2(\varrho)}{2}} dx + \varrho^{n(1-\frac{p_2(\varrho)}{q})} \left(\int_{B_{\varrho}^+(x_0)} |Dg|^q dx \right)^{\frac{p_2(\varrho)}{q}} \right],
\end{aligned}$$

while, when $1 < p_2(\varrho) < 2$, via Hölder inequality, (3.34), the minimality of v in class $\hat{\mathcal{C}}_u^{p_2(\varrho)}(B_{\varrho/2}^+(x_0), \mathcal{M})$ and (2.10) we can conclude that

$$\begin{aligned}
&\int_{B_{\varrho/2}^+(x_0)} |Du - Dv|^{p_2(\varrho)} dx \\
&\leq \left(\int_{B_{\varrho/2}^+(x_0)} |Du - Dv|^2 (|Du|^2 + |Dv|^2)^{\frac{p_2(\varrho)-2}{2}} dx \right)^{\frac{p_2(\varrho)}{2}} \\
&\quad \cdot \left(\int_{B_{\varrho/2}^+(x_0)} (|Du|^2 + |Dv|^2)^{\frac{p_2(\varrho)}{2}} dx \right)^{\frac{2-p_2(\varrho)}{2}} \\
&\leq c\varepsilon^{\frac{\sigma' \gamma_1^2}{2(1+\sigma')}} \left[\int_{B_{\varrho}^+(x_0)} (1 + |Du|^2)^{\frac{p_2(\varrho)}{2}} dx + \varrho^{n(1-\frac{p_2(\varrho)}{q})} \left(\int_{B_{\varrho}^+(x_0)} |Dg|^q dx \right)^{\frac{p_2(\varrho)}{q}} \right].
\end{aligned}$$

All in all, setting $\kappa := \frac{\gamma_1 \sigma'}{1 + \sigma'} \min \{1, \frac{\gamma_1}{2}\}$, we get

$$(3.35) \quad \begin{aligned} \int_{B_{\varrho/2}^+(x_0)} |Du - Dv|^{p_2(\varrho)} \, dx &\leq c\varepsilon^\kappa \int_{B_\varrho^+(x_0)} (1 + |Du|^2)^{\frac{p_2(\varrho)}{2}} \, dx \\ &+ c\varepsilon^\kappa \varrho^{n(1 - \frac{p_2(\varrho)}{q})} \left(\int_{B_\varrho^+(x_0)} |Dg|^q \, dx \right)^{\frac{p_2(\varrho)}{q}}, \end{aligned}$$

for $c \equiv c(\mathbf{data}_{p(\cdot)})$.

Step 3: Comparison, second time. Set $k_0 := k(x_0)$. We confront v with the solution $h \in W^{1,p_2(\varrho)}(B_{\varrho/4}^+(x_0), \mathbb{R}^N)$ of the Dirichlet problem

$$(3.36) \quad \hat{C}_v^{p_2(\varrho)}(B_{\varrho/4}^+(x_0), \mathbb{R}^N) \ni w \mapsto \int_{B_{\varrho/4}^+(x_0)} k_0 |Dw|^{p_2(\varrho)} \, dx.$$

Furthermore, h solves the Euler-Lagrange equation

$$(3.37) \quad 0 = \int_{B_{\varrho/4}^+(x_0)} k_0 p_2(\varrho) |Dh|^{p_2(\varrho)-2} Dh \cdot D\varphi \, dx$$

for all $\varphi \in W_0^{1,p_2(\varrho)}(B_{\varrho/4}^+(x_0), \mathbb{R}^N)$. Notice that, by the results in [40] there holds that

$$(3.38) \quad \|h\|_{L^\infty(B_{\varrho/4}^+(x_0))} \leq c(N) \|v\|_{L^\infty(B_{\varrho/4}^+(x_0))} \leq c(N, m).$$

Recalling [13, Lemma 3.4], see also [37, Proof of Lemma 2] there holds that

$$(3.39) \quad \begin{aligned} \int_{B_\varsigma^+(x_0)} |Dh|^{p_2(\varrho)} \, dx &\leq c \left(\frac{\varsigma}{\varrho} \right)^\vartheta \int_{B_{\varrho/2}^+(x_0)} |Du|^{p_2(\varrho)} \, dx \\ &+ c\varsigma^{n(1 - \frac{p_2(\varrho)}{q})} \left(\int_{B_{\varrho/2}^+(x_0)} |Dg|^q \, dx \right)^{\frac{p_2(\varrho)}{q}}, \end{aligned}$$

for all $\vartheta \in \left(n \left(1 - \frac{p_2(\varrho)}{q} \right), n \right)$ with $c \equiv c(n, N, \gamma_1, \gamma_2, \lambda, \Lambda, q)$. For (3.39) we also used that, by (3.36) and (3.30),

$$\mathbf{tr}_{\Gamma_1}(h) = \mathbf{tr}_{\Gamma_{R/4}}(v) = \mathbf{tr}_{\Gamma_{R/4}}(u) = \mathbf{tr}_{\Gamma_{R/4}}(g),$$

the minimality of h in class $\hat{C}_v^{p_2(\varrho)}(B_{\varrho/4}^+(x_0), \mathbb{R}^N)$ and the one of v in class $\hat{C}_u^{p_2(\varrho)}(B_{\varrho/2}^+(x_0), m)$. Exploiting now the monotonicity properties of the integrand in (3.36), Lemma 2.1, (3.31), (3.37), Hölder inequality and the minimality of h in class $\hat{C}_v^{p_2(\varrho)}(B_{\varrho/4}^+(x_0), \mathbb{R}^N)$, we estimate

$$\begin{aligned} &c \int_{B_{\varrho/4}^+(x_0)} |V_{p_2(\varrho)}(Dv) - V_{p_2(\varrho)}(Dh)|^2 \, dx \\ &\leq \int_{B_{\varrho/4}^+(x_0)} k_0 p_2(\varrho) \left(|Dv|^{p_2(\varrho)-2} Dv - |Dh|^{p_2(\varrho)-2} Dh \right) \cdot (Dv - Dh) \, dx \\ &= \int_{B_{\varrho/4}^+(x_0)} (k_0 - k(x)) p_2(\varrho) |Dv|^{p_2(\varrho)-2} Dv \cdot (Dv - Dh) \, dx \\ &\quad + \int_{B_{\varrho/4}^+(x_0)} k(x) p_2(\varrho) |Dv|^{p_2(\varrho)-2} Dv \cdot (Dv - Dh) \, dx \\ &\leq c\varrho^\alpha \int_{B_{\varrho/4}^+(x_0)} |Dv|^{p_2(\varrho)-1} |Dv - Dh| \, dx + c \int_{B_{\varrho/4}^+(x_0)} |Dv|^{p_2(\varrho)} |v - h| \, dx \\ &\leq c\varrho^\alpha \int_{B_{\varrho/4}^+(x_0)} |Dv|^{p_2(\varrho)} \, dx + c \int_{B_{\varrho/4}^+(x_0)} |Dv|^{p_2(\varrho)} |v - h| \, dx =: c [\varrho^\alpha (\text{I}) + (\text{II})], \end{aligned}$$

with $c \equiv c(n, N, \lambda, \Lambda, \gamma_1, \gamma_2, [k]_{0,\alpha}, \alpha)$. The minimality of v in class $\hat{C}_u^{p_2(\varrho)}(B_{\varrho/2}^+(x_0), m)$ yields that

$$(I) \leq \int_{B_{\varrho}^+(x_0)} |Du|^{p_2(\varrho)} \, dx$$

and, recalling also (3.28), we see that

$$\begin{aligned} \left(\frac{\varrho}{2}\right)^{p_2(\varrho)-n} \int_{B_{\varrho/2}^+(x_0)} |Dv|^{p_2(\varrho)} \, dx &\leq 2^{n-\gamma_1} \varrho^{p_2(\varrho)-n} \int_{B_{\varrho/2}^+(x_0)} |Du|^{p_2(\varrho)} \, dx \\ &\leq 2^{n-\gamma_1} \left[\varrho^{p_2(\varrho)-n} \int_{B_{\varrho}^+(x_0)} (1 + |Du|^2)^{\frac{p_2(\varrho)}{2}} \, dx + \left(\varrho^{q-n} \int_{B_{\varrho}^+(x_0)} |Dg|^q \, dx \right)^{\frac{p_2(\varrho)}{q}} \right] \\ (3.40) \quad &< 2^{n-\gamma_1} \varepsilon^{p_2(\varrho)}. \end{aligned}$$

By Hölder inequality, the minimality of h in class $\hat{C}_v^{p_2(\varrho)}(B_{\varrho/4}^+(x_0), \mathbb{R}^N)$ and the one of v in class $\hat{C}_u^{p_2(\varrho)}(B_{\varrho/2}^+(x_0), m)$, Lemma 3.2, (3.7), (3.38) and (3.40) we bound

$$\begin{aligned} (II) &\leq c \varrho^n \left(\int_{B_{\varrho/4}^+(x_0)} |Dv|^{p_2(\varrho)(1+\sigma')} \, dx \right)^{\frac{1}{1+\sigma'}} \left(\int_{B_{\varrho/4}^+(x_0)} |v-h|^{p_2(\varrho)} \, dx \right)^{\frac{\sigma'}{1+\sigma'}} \\ &\leq c \varrho^n \left[\int_{B_{\varrho/2}^+(x_0)} |Du|^{p_2(\varrho)} \, dx + \left(\int_{B_{\varrho/2}^+(x_0)} |Dg|^q \, dx \right)^{\frac{p_2(\varrho)}{q}} \right] \\ &\quad \cdot \left(\varrho^{p_2(R)-n} \int_{B_{\varrho/4}^+(x_0)} |Dv - Dh|^{p_2(\varrho)} \, dx \right)^{\frac{\sigma'}{1+\sigma'}} \\ &\leq c \varepsilon^{\frac{\gamma_1 \sigma'}{1+\sigma'}} \left[\int_{B_{\varrho}^+(x_0)} (1 + |Du|^2)^{\frac{p_2(\varrho)}{2}} \, dx + \varrho^{1-\frac{p_2(\varrho)}{q}} \left(\int_{B_{\varrho}^+(x_0)} |Dg|^q \, dx \right)^{\frac{p_2(\varrho)}{q}} \right], \end{aligned}$$

for $c \equiv c(\mathbf{data}_{p(\cdot)})$. Merging the content of the two previous displays and proceeding as in the last part of *Step 2* we end up with

$$\begin{aligned} &\int_{B_{\varrho/4}^+(x_0)} |Dv - Dh|^{p_2(\varrho)} \, dx \\ (3.41) \quad &\leq c [\varepsilon^\kappa + \varrho^\alpha] \left[\int_{B_{\varrho}^+(x_0)} (1 + |Du|^2)^{\frac{p_2(\varrho)}{2}} \, dx + \varrho^{n(1-\frac{p_2(\varrho)}{q})} \left(\int_{B_{\varrho}^+(x_0)} |Dg|^q \, dx \right)^{\frac{p_2(\varrho)}{q}} \right], \end{aligned}$$

with $c \equiv c(\mathbf{data})$. Collecting inequalities (3.35) and (3.41) we obtain

$$\begin{aligned} &\int_{B_{\varrho/4}^+(x_0)} |Du - Dh|^{p_2(\varrho)} \, dx \\ (3.42) \quad &\leq c [\varepsilon^\kappa + \varrho^\alpha] \left[\int_{B_{\varrho}^+(x_0)} (1 + |Du|^2)^{\frac{p_2(\varrho)}{2}} \, dx + \varrho^{n(1-\frac{p_2(\varrho)}{q})} \left(\int_{B_{\varrho}^+(x_0)} |Dg|^q \, dx \right)^{\frac{p_2(\varrho)}{q}} \right], \end{aligned}$$

where $c \equiv c(\mathbf{data})$.

Step 4: Morrey decay estimates at the boundary. Let $\varsigma \in (0, \frac{\varrho}{4})$ and estimate, via (3.27), (3.39) and (3.42),

$$\begin{aligned}
& \int_{B_\varsigma(x_0)} (1 + |D\tilde{u}|^2)^{\frac{p_2(\varrho)}{2}} dx \leq c \left[\int_{B_\varsigma^+(x_0)} |Du|^{p_2(\varrho)} dx + \int_{B_\varsigma^+(x_0)} |Dg|^{p_2(\varrho)} dx \right] + c\varsigma^n \\
& \leq c \left[\int_{B_\varsigma^+(x_0)} |Du - Dh|^{p_2(\varrho)} dx + \int_{B_\varsigma^+(x_0)} |Dh|^{p_2(\varrho)} dx + \int_{B_\varsigma^+(x_0)} |Dg|^{p_2(\varrho)} dx \right] + c\varsigma^n \\
& \leq c \left[\left(\frac{\varsigma}{\varrho} \right)^n + \varepsilon^\kappa + \varrho^\alpha + \left(\frac{\varsigma}{\varrho} \right)^\vartheta \right] \\
& \quad \cdot \left[\int_{B_\varrho^+(x_0)} (1 + |Du|^2)^{\frac{p_2(\varrho)}{2}} dx + \varrho^{n(1-\frac{p_2(\varrho)}{q})} \left(\int_{B_\varrho^+(x_0)} |Dg|^q dx \right)^{\frac{p_2(\varrho)}{q}} \right] \\
& \quad + c \left(\frac{\varsigma}{\varrho} \right)^{n(1-\frac{p_2(\varrho)}{q})} \left(\varrho^{q-n} \int_{B_\varrho^+(x_0)} |Dg|^q dx \right)^{\frac{p_2(\varrho)}{q}},
\end{aligned}$$

where $c \equiv c(\mathbf{data})$. Now recall that $\vartheta > n \left(1 - \frac{p_2(\varrho)}{q} \right)$, so we can always find $\hat{\nu} \in \left(n \left(1 - \frac{p_2(\varrho)}{q} \right), \vartheta \right)$. Moreover, set

$$\hat{p}_2(\varrho) := n \left(1 - \frac{p_2(\varrho)}{q} \right) \quad \text{and} \quad \tilde{p}_2(\varrho) := p_2(\varrho) - n + \hat{\nu}$$

and choose $\varsigma = \tau\varrho$ for some $\tau \in (0, \frac{1}{4})$. Multiplying both sides of the previous inequality by $(\tau\varrho)^{p_2(\varrho)-n}$ we obtain

$$\begin{aligned}
& (\tau\varrho)^{p_2(\varrho)-n} \int_{B_{\tau\varrho}(x_0)} (1 + |D\tilde{u}|^2)^{\frac{p_2(\varrho)}{2}} dx \\
& \leq \tau^{\tilde{p}_2(\varrho)} \left[c\tau^{n-\nu} + c\tau^{-\nu} (\varepsilon^\kappa + R_*^\alpha) + c\tau^{\vartheta-\nu} \right] \\
& \quad \cdot \left[\varrho^{p_2(\varrho)-n} \int_{B_\varrho^+(x_0)} (1 + |Du|^2)^{\frac{p_2(\varrho)}{2}} dx + \left(\varrho^{q-n} \int_{B_\varrho^+(x_0)} |Dg|^q dx \right)^{\frac{p_2(\varrho)}{q}} \right] \\
& \quad + c\tau^{\hat{p}_2(\varrho)} \left(\varrho^{q-n} \int_{B_\varrho^+(x_0)} |Dg|^q dx \right)^{\frac{1}{q}},
\end{aligned} \tag{3.43}$$

for $c \equiv c(\mathbf{data})$. With the notation introduced in *Step 2*, the inequality in (3.43) reads as

$$\begin{aligned}
& \phi(x_0, \tau\varrho, p_2(\varrho)) \leq \tau^{\frac{\tilde{p}_2(\varrho)}{p_2(\varrho)}} \left[c\tau^{\frac{(n-\nu)}{p_2(\varrho)}} + c\tau^{-\frac{\nu}{p_2(\varrho)}} \left(\varepsilon^{\frac{\kappa}{p_2(\varrho)}} + R_*^{\frac{\alpha}{p_2(\varrho)}} \right) + c\tau^{\frac{\vartheta-\nu}{p_2(\varrho)}} \right] \\
& \quad \cdot \left[\psi^+(x_0, \varrho) + \left(\varrho^{q-n} \int_{B_\varrho^+(x_0)} |Dg|^q dx \right)^{\frac{1}{q}} \right] + c\tau^{1-\frac{n}{q}} \left(\varrho^{q-n} \int_{B_\varrho^+(x_0)} |Dg|^q dx \right)^{\frac{1}{q}},
\end{aligned} \tag{3.44}$$

therefore since

$$\phi(x_0, r, p) \leq \phi(x_0, r, q) \quad \text{for } p \leq q, \tag{3.45}$$

we obtain from (3.44):

$$\psi(x_0, \tau\varrho) \leq \tau^{\frac{\tilde{p}_2(\varrho)}{p_2(\varrho)}} \left[c\tau^{\frac{(n-\nu)}{p_2(\varrho)}} + c\tau^{-\frac{\nu}{p_2(\varrho)}} \left(\varepsilon^{\frac{\kappa}{p_2(\varrho)}} + R_*^{\frac{\alpha}{p_2(\varrho)}} \right) + c\tau^{\frac{\vartheta-\nu}{p_2(\varrho)}} \right]$$

$$\begin{aligned} & \cdot \left[\psi^+(x_0, \varrho) + \left(\varrho^{q-n} \int_{B_\varrho^+(x_0)} |Dg|^q \, dx \right)^{\frac{p_2(\varrho)}{q}} \right] \\ & + c\tau^{1-\frac{n}{q}} \left(\varrho^{q-n} \int_{B_\varrho^+(x_0)} |Dg|^q \, dx \right)^{\frac{1}{q}} \end{aligned}$$

with $c \equiv c(\mathbf{data})$. Recalling that $\tau \in (0, \frac{1}{4})$, it is easy to see that

$$\begin{aligned} & \tau^{\frac{\bar{p}_2(\varrho)}{p_2(\varrho)}} \left[c\tau^{\frac{(n-\nu)}{p_2(\varrho)}} + c\tau^{-\frac{\nu}{p_2(\varrho)}} \left(\varepsilon^{\frac{\kappa}{p_2(\varrho)}} + R_*^{\frac{\alpha}{p_2(\varrho)}} \right) + c\tau^{\frac{\vartheta-\nu}{p_2(\varrho)}} \right] \\ & \leq \tau^{\frac{\bar{p}_2(\varrho)}{p_2(\varrho)}} \left[c\tau^{\frac{n-\vartheta}{\gamma_2}} + c\tau^{-\frac{\vartheta}{\gamma_1}} \left(\varepsilon^{\frac{\kappa}{\gamma_2}} + R_*^{\frac{\alpha}{\gamma_2}} \right) + c\tau^{\frac{\vartheta-\nu}{\gamma_2}} \right], \end{aligned}$$

therefore, merging the content of the two above displays we obtain

$$\begin{aligned} \psi(x_0, \tau\varrho) & \leq \tau^{\frac{\bar{p}_2(\varrho)}{p_2(\varrho)}} \left[c\tau^{\frac{n-\vartheta}{\gamma_2}} + c\tau^{-\frac{\vartheta}{\gamma_1}} \left(\varepsilon^{\frac{\kappa}{\gamma_2}} + R_*^{\frac{\alpha}{\gamma_2}} \right) + c\tau^{\frac{\vartheta-\nu}{\gamma_2}} \right] \\ & \cdot \left[\psi^+(x_0, \varrho) + \left(\varrho^{q-n} \int_{B_\varrho^+(x_0)} |Dg|^q \, dx \right)^{\frac{p_2(\varrho)}{q}} \right] \\ & + c\tau^{1-\frac{n}{q}} \left(\varrho^{q-n} \int_{B_\varrho^+(x_0)} |Dg|^q \, dx \right)^{\frac{1}{q}} \\ & \leq \tau^{\frac{\bar{p}_2(\varrho)}{p_2(\varrho)}} \left[c\tau^{\frac{n-\vartheta}{\gamma_2}} + c\tau^{-\frac{\vartheta}{\gamma_1}} \left(\varepsilon^{\frac{\kappa}{\gamma_2}} + R_*^{\frac{\alpha}{\gamma_2}} \right) + c\tau^{\frac{\vartheta-\nu}{\gamma_2}} \right] \\ & \cdot \left[\psi(x_0, \varrho) + \left(\varrho^{q-n} \int_{B_\varrho^+(x_0)} |Dg|^q \, dx \right)^{\frac{p_2(\varrho)}{q}} \right] \\ & + c\tau^{1-\frac{n}{q}} \left(\varrho^{q-n} \int_{B_\varrho^+(x_0)} |Dg|^q \, dx \right)^{\frac{1}{q}} \end{aligned} \tag{3.46}$$

with $c \equiv c(\mathbf{data})$. Select τ , ε and R_* so small that

$$\begin{aligned} \tau^{\frac{\bar{p}_2(\varrho)}{p_2(\varrho)}} & \leq \frac{1}{8}, & c'\tau^{\frac{n-\vartheta}{\gamma_2}} & \leq \frac{1}{3}, & c'\tau^{-\frac{\vartheta}{\gamma_1}} \left(\varepsilon^{\frac{\kappa}{\gamma_2}} + R_*^{\frac{\alpha}{\gamma_2}} \right) & \leq \frac{1}{3} \\ c'\tau^{\frac{\vartheta-\nu}{\gamma_2}} & \leq \frac{1}{3}, & (c' + c)\tau^{1-\frac{n}{q}} & \leq \frac{1}{8}, \end{aligned} \tag{3.47}$$

where c' is the same constant appearing in (3.29). By (3.29) and (3.28), with the choice made above we can conclude that

$$\begin{aligned} \chi^+(x_0, \tau\varrho) & \leq \frac{1}{2} \left[\psi^+(x_0, \varrho) + \left(\varrho^{q-n} \int_{B_\varrho^+(x_0)} |Dg|^q \, dx \right)^{\frac{p_2(\varrho)}{q}} \right] \\ & + \frac{1}{2} \left(\varrho^{q-n} \int_{B_\varrho^+(x_0)} |Dg|^q \, dx \right)^{\frac{1}{q}} < \varepsilon, \end{aligned}$$

so iterations are legal. Moreover, combining (3.46) and (3.47) we have

$$(3.48) \quad \psi(x_0, \tau \varrho) \leq \tau^{\frac{\bar{p}_2(\varrho)}{p_2(\varrho)}} \psi(x_0, \varrho) + c \varrho^{1-\frac{n}{q}} \left(\int_{B_\varrho^+(x_0)} |Dg|^q \, dx \right)^{\frac{1}{q}}$$

for $c \equiv (\mathbf{data}, q)$. Iterating (3.48) we end up with

$$(3.49) \quad \begin{aligned} \psi(x_0, \tau^k \varrho) &\leq \tau^{k \frac{\bar{p}_2(\varrho)}{p_2(\varrho)}} \psi(x_0, \varrho) \\ &+ c \left(\int_{B_\varrho^+(x_0)} |Dg|^q \, dx \right)^{\frac{1}{q}} \varrho^{1-\frac{n}{q}} \tau^{(k-1)(1-\frac{n}{q})} \sum_{j=0}^{k-1} \tau^{j(\frac{\bar{p}_2(\varrho)}{p_2(\varrho)} - 1 + \frac{n}{q})}. \end{aligned}$$

Since $\frac{\bar{p}_2(\varrho)}{p_2(\varrho)} - 1 + \frac{n}{q} > 0$, the series on the right-hand side of (3.48) converges, so we have

$$(3.50) \quad \psi(x_0, \tau^k \varrho) \leq \tau^{k \frac{\bar{p}_2(\varrho)}{p_2(\varrho)}} \psi(x_0, \varrho) + c \left(\int_{B_\varrho^+(x_0)} |Dg|^q \, dx \right)^{\frac{1}{q}} \varrho^{1-\frac{n}{q}} \tau^{(k-1)(1-\frac{n}{q})}.$$

Whenever $0 < \varsigma < \varrho$ we can find $k \in \mathbb{N}$ so that $\tau^{k+1} \varrho \leq \varsigma < \tau^k \varrho$, so using (3.50) we obtain

$$(3.51) \quad \begin{aligned} \psi(x_0, \varsigma) &\leq \tau^{1-\frac{n}{p_2(\varrho)}} \psi(x_0, \tau^k \varrho) \\ &\leq \tau^{1-\frac{n}{p_2(\varrho)}} \left[\tau^{k \frac{\bar{p}_2(\varrho)}{p_2(\varrho)}} \psi(x_0, \varrho) + c \varrho^{1-\frac{n}{q}} \tau^{(k-1)(1-\frac{n}{q})} \left(\int_{B_\varrho^+(x_0)} |Dg|^q \, dx \right)^{\frac{1}{q}} \right] \\ &\leq c \tau^{-1-\frac{n}{\gamma_1}} \left[\left(\frac{\varsigma}{\varrho} \right)^{\frac{\bar{p}_2(\varrho)}{p_2(\varrho)}} \psi(x_0, \varrho) + \varrho^{1-\frac{n}{q}} \left(\int_{B_\varrho^+(x_0)} |Dg|^q \, dx \right)^{\frac{1}{q}} \right] \\ &\leq c \left[\left(\frac{\varsigma}{\varrho} \right)^{1-\frac{n}{q}} \psi(x_0, \varrho) + \varsigma^{1-\frac{n}{q}} \left(\int_{B_\varrho^+(x_0)} |Dg|^q \, dx \right)^{\frac{1}{q}} \right], \end{aligned}$$

for $c \equiv c(\mathbf{data})$. To summarize, we just got that, if $x_0 \in \Gamma_R$ is any point satisfying (3.28) on $B_\varrho^+(x_0)$ for some $\varrho \in (0, R - |x_0|)$ then

$$(3.52) \quad \begin{aligned} \psi(x_0, \varsigma) &\leq c \left[\left(\frac{\varsigma}{\varrho} \right)^{1-\frac{n}{q}} \psi(x_0, \varrho) + \varsigma^{1-\frac{n}{q}} \left(\int_{B_\varrho^+(x_0)} |Dg|^q \, dx \right)^{\frac{1}{q}} \right] \\ &\leq c \left[\left(\frac{\varsigma}{\varrho} \right)^{1-\frac{n}{q}} \left(\varrho^{p_2(\varrho)} \int_{B_\varrho^+(x_0)} (1 + |Du|^2)^{\frac{p_2(\varrho)}{2}} \, dx \right)^{\frac{1}{p_2(\varrho)}} + \varsigma^{1-\frac{n}{q}} \left(\int_{B_\varrho^+(x_0)} |Dg|^q \, dx \right)^{\frac{1}{q}} \right] \\ &\leq c \left[\left(\frac{\varsigma}{\varrho} \right)^{1-\frac{n}{q}} \chi^+(x_0, \varrho) + \varsigma^{1-\frac{n}{q}} \|Dg\|_{L^q(B_1^+)} \right] \leq c \left[\left(\frac{\varsigma}{\varrho} \right)^{1-\frac{n}{q}} + \varsigma^{1-\frac{n}{q}} \right] \leq c \left(\frac{\varsigma}{\varrho} \right)^{1-\frac{n}{q}} \end{aligned}$$

for $c \equiv c(\mathbf{data}, \|Dg\|_{L^q(B_1^+)})$. In (3.52) we also used (3.28) to control $\chi^+(x_0, \varrho)$ with $\varepsilon \in (0, 1]$.

Step 5: Partial Hölder continuity. Now we aim to prove an estimate analogous to (3.52) valid also for points $x_0 \in \bar{B}_R^+$ not necessarily belonging to Γ_R . As in [37, Proof of Lemma 2], we fix $\iota = \frac{1}{1000}$ and $x_0 \in B_R^+$ satisfying (3.28) for some $\varrho \in (0, R - |x_0|)$. For $0 < \sigma < \varrho$ we distinguish two main cases: $\varsigma < \iota \varrho$ or $\varsigma \geq \iota \varrho$.

Case 1: $\varsigma < \iota \varrho$. We take $\hat{x} \in \Gamma_R$ so that $d := \text{dist}(x_0, \Gamma_R) = |x_0 - \hat{x}|$. Now, if $\iota \varrho \geq d$ we notice that

$$B_d(x_0) \subset B_{2d}(\hat{x}) \subset B_{\varrho/2}(x_0) \subset B_\varrho(\hat{x}),$$

therefore

$$\chi^+ \left(\hat{x}, \frac{\varrho}{2} \right) \leq c\chi^+(x_0, \varrho) \leq c(n, \gamma_1, \gamma_2, q)\varepsilon,$$

so reducing the size of $\varepsilon \equiv \varepsilon(\mathbf{data})$ determined in (3.47) to $\varepsilon' := \frac{\varepsilon}{2c}$ we end up with

$$(3.53) \quad \chi^+ \left(\hat{x}, \frac{\varrho}{2} \right) < \varepsilon'.$$

If $\iota\varrho > \varsigma \geq d$ we immediately notice that

$$B_\varsigma(x_0) \subset B_{4\varsigma}(\hat{x}) \subset B_{\varrho/4}(\hat{x}) \subset B_\varrho(x_0),$$

and, since (3.53) legalizes (3.52) with x_0 replaced by \hat{x} , we obtain

$$\psi(x_0, \varsigma) \leq c\psi(\hat{x}, 4\varsigma) \leq c8^{1-\frac{n}{q}} \left(\frac{\varsigma}{\varrho} \right)^{1-\frac{n}{q}},$$

for $c \equiv c(\mathbf{data}, \|Dg\|_{L^q(B_1^+)})$. On the other hand, if $\iota\varrho \geq d > \varsigma$ we have that

$$B_\varsigma(x_0) \subset B_{2d}(\hat{x}) \subset B_{\varrho/4}(\hat{x}) \subset B_\varrho(x_0),$$

so, by (3.53) and (3.52) (with x_0 replaced by \hat{x} of course) we get

$$\begin{aligned} \psi(x_0, \varsigma) &\leq c\frac{\varsigma}{d}\psi(\hat{x}, 2d) \leq c\frac{\varsigma}{d} \left[\left(\frac{4d}{\varrho} \right)^{1-\frac{n}{q}} + (2d)^{1-\frac{n}{q}} \right] \\ &\leq c \left(\frac{\varsigma}{d} \right)^{\frac{n}{q}} \left[\left(\frac{\varsigma}{\varrho} \right)^{1-\frac{n}{q}} + \varsigma^{1-\frac{n}{q}} \right] \leq c \left[\left(\frac{\varsigma}{\varrho} \right)^{1-\frac{n}{q}} + \varsigma^{1-\frac{n}{q}} \right] \leq c \left(\frac{\varsigma}{\varrho} \right)^{1-\frac{n}{q}} \end{aligned}$$

also in this case, with $c \equiv c(\mathbf{data}, \|Dg\|_{L^q(B_1^+)})$. Now we consider the occurrence $\varsigma < \iota\varrho < d$. It follows that $B_{\iota\varrho}(x_0) \subseteq B_R^+$ and

$$2\iota\varrho \left(\int_{B_{2\iota\varrho}(x_0)} (1 + |Du|^2)^{\frac{p_2(2\iota\varrho)}{2}} dx \right)^{\frac{1}{p_2(2\iota\varrho)}} \leq c\iota^{1-\frac{n}{\gamma_1}} \chi^+(x_0, \varrho) < c(n, \gamma_1, \gamma_2, q)\varepsilon.$$

Reducing the size of ε in such a way that $c\varepsilon \leq \varepsilon_0$, where ε_0 is the smallness threshold appearing in [9, (3.16)] we obtain

$$2\iota\varrho \left(\int_{B_{2\iota\varrho}(x_0)} (1 + |Du|^2)^{\frac{p_2(2\iota\varrho)}{2}} dx \right)^{\frac{1}{p_2(2\iota\varrho)}} < \varepsilon_0.$$

Then, [9, estimates (3.40)-(3.43)] apply, thus getting

$$\left(\varsigma^{p_2(\varsigma)} \int_{B_\varsigma(x_0)} (1 + |Du|^2)^{p_2(\varsigma)} dx \right)^{\frac{1}{p_2(\varsigma)}} \leq \iota^{-\beta_0} c(\mathbf{data}, \beta_0) \left(\frac{\varsigma}{\varrho} \right)^{\beta_0},$$

for all $\beta_0 \in (0, 1)$. Recalling the explicit expression of $\psi(x_0, \cdot)$ we then bound

$$\begin{aligned} \psi(x_0, \sigma) &\leq c \left(\varsigma^{p_2(\varsigma)} \int_{B_\varsigma(x_0)} (1 + |Du|^2)^{\frac{p_2(\varsigma)}{2}} dx \right)^{\frac{1}{p_2(\varsigma)}} \\ &\quad + c \left(\varsigma^{q-n} \int_{B_\varsigma(x_0) \cap B_R^+} |Dg|^q dx \right)^{\frac{1}{q}} \leq c \left[\left(\frac{\varsigma}{\varrho} \right)^{\beta_0} + \varsigma^{1-\frac{n}{q}} \right], \end{aligned}$$

with $c \equiv c(\mathbf{data}, \|Dg\|_{L^q(B_1^+)}, \beta_0)$. The desired estimate follows by choosing $\beta_0 = 1 - \frac{n}{q}$ in the previous display.

Case 2: $\varsigma \geq \iota\varrho$. Estimate (3.52) trivially holds with a constant $c \equiv c(\mathbf{data}, \|Dg\|_{L^q(B_1^+)})$.

All in all, we have just proved that if $x_0 \in \bar{B}_R^+$ satisfies (3.28) on $B_\varrho(x_0) \cap B_R^+$ for some $\varrho \in (0, R - |x_0|)$, then

$$(3.54) \quad \psi(x_0, \varsigma) \leq c(\mathbf{data}, \|Dg\|_{L^q(B_1^+)}) \left(\frac{\varsigma}{\varrho}\right)^{1-\frac{n}{q}}.$$

Now, by the continuity of Lebesgue's integral and of the mapping $x_0 \mapsto p_2(x_0, \varrho)$, we can conclude that if (3.28) holds for x_0 on $B_\varrho(x_0) \cap B_R^+$ then it holds also on $B_\varrho(y) \cap B_R^+$ for all $y \in \bar{B}_1^+$ belonging to a sufficiently small, relatively open neighborhood of x_0 , say, $B_{x_0} \subset \bar{B}_R^+$. Then the set

$$D_0 := \left\{ y \in B_{x_0} : \chi^+(y, \varrho) < \varepsilon \text{ on } B_\varrho(y) \cap B_R^+, R \in (0, R_*], \varrho \in (0, R - |y|) \right\}$$

is relatively open, so via (3.54) we can conclude that

$$(3.55) \quad \left(\varsigma^{-n(1-\frac{n}{q})} \int_{B_\varsigma(x_0)} |D\tilde{u}|^{\gamma_1} dx \right)^{\frac{1}{\gamma_1}} \leq \psi(x_0, \varsigma) \leq c \left(\frac{\varsigma}{\varrho} \right)^{1-\frac{n}{q}},$$

where $c \equiv c(\mathbf{data}, \|Dg\|_{L^q(B_1^+)})$. By (3.55) and the well-known characterization of Hölder continuity due to Campanato and Meyers we can conclude that \tilde{u} is $(1 - \frac{n}{q})$ -Hölder continuous in a neighborhood of D_0 , which in turn implies that $u \in C_{loc}^{0, 1-\frac{n}{q}}(D_0, \mathcal{M})$.

Step 6: Hausdorff dimension of the singular set. Given the characterization of D_0 , we easily see that the singular set $\Sigma_0(u, B_\varrho(x_0) \cap B_R^+)$ can be defined as

$$\Sigma_0(u, B_\varrho(x_0) \cap B_R^+) := \bar{B}_R^+ \cap \bar{B}_\varrho(x_0) \setminus D_0.$$

Moreover, for $y \in B_\varrho(x_0) \cap B_R^+$, being $g \in C^{0, 1-\frac{n}{q}}(\bar{B}_1^+, \mathcal{M})$ we see that

$$\begin{aligned} \limsup_{\varsigma \rightarrow 0} \chi^+(y, \varsigma) &\leq \limsup_{\varsigma \rightarrow 0} \left(\varsigma^{p_2(y, \varsigma)-n} \int_{B_\varsigma(y) \cap B_R^+} (1 + |Du|^2)^{\frac{p_2(y, \varsigma)}{2}} dx \right)^{\frac{1}{p_2(y, \varsigma)}} \\ &\quad + \limsup_{\varsigma \rightarrow 0} \left(\varsigma^{q-n} \int_{B_\varsigma(y) \cap B_R^+} |Dg|^q dx \right)^{\frac{1}{q}} \\ &\leq \limsup_{\varsigma \rightarrow 0} \left(\varsigma^{p_2(y, \varsigma)-n} \int_{B_\varsigma(y) \cap B_R^+} (1 + |Du|^2)^{\frac{p_2(y, \varsigma)}{2}} dx \right)^{\frac{1}{p_2(y, \varsigma)}}, \end{aligned}$$

therefore

$$\Sigma_0(u, B_\varrho(x_0) \cap B_R^+) \subset \left\{ y \in \bar{B}_\varrho(x_0) \cap \bar{B}_R^+ : \limsup_{\varsigma \rightarrow 0} \psi^+(y, \varsigma) > 0 \right\}.$$

Now, notice that, as in (3.24),

$$(3.56) \quad p_2(y, \varsigma) \leq (1 + \sigma_0)p_1(x_0, R_*) \text{ for all } 0 < \varsigma \leq R_*, \quad B_\varsigma(y) \cap B_R^+ \subset B_\varrho(x_0) \cap B_R^+,$$

so we obtain,

$$\begin{aligned} &\left(\varsigma^{p_2(y, \varsigma)} \int_{B_\varsigma(y) \cap B_R^+} (1 + |Du|^2)^{\frac{p_2(y, \varsigma)}{2}} dx \right)^{\frac{1}{p_2(y, \varsigma)}} \\ &\leq \left(\varsigma^{p_1(x_0, R_*)(1+\sigma_0)} \int_{B_\varsigma(y) \cap B_R^+} (1 + |Du|^2)^{\frac{p_1(x_0, R_*)(1+\sigma_0)}{2}} dx \right)^{\frac{1}{p_1(x_0, R_*)(1+\sigma_0)}}, \end{aligned}$$

which by (3.6) is finite. This allows concluding that $\Sigma_0(u, B_\varrho(x_0) \cap B_R^+)$ is contained into the set

$$D_1 := \left\{ y \in \bar{B}_\varrho(x_0) \cap \bar{B}_R^+ : \limsup_{\varsigma \rightarrow 0} \phi^+(y, \varsigma, p_1(x_0, R_*)(1 + \sigma_0))^{p_1(x_0, R_*)(1+\sigma_0)} > 0 \right\}.$$

By [25, Proposition 2.7] it follows that $\dim_{\mathcal{H}}(D_1) \leq n - p_1(x_0, R_*)(1 + \sigma_0)$, so, by (2.2)₂ we easily have that $\dim_{\mathcal{H}}(D_1) < n - \gamma_1$ and so $\dim_{\mathcal{H}}(\Sigma_0(u, B_\varrho(x_0) \cap B_R^+)) < n - \gamma_1$. The proof is complete. \square

Once Proposition 3.2 is available, we can cover B_1^+ with balls having the same features of $B_\varrho(x_0) \cap B_R^+$ and remembering that, by (2.2)₂, $p_1(x_0, R_*) \geq \gamma_1$, we obtain that $\dim_{\mathcal{H}}(\Sigma_0(u)) \leq n - \gamma_1(1 + \delta_0) < n - \gamma_1$, and so $\dim_{\mathcal{H}}(\Sigma_0(u)) < n - \gamma_1$. Via a standard covering argument, we can conclude that $u \in C_{loc}^{0,1-\frac{n}{q}}(\bar{B}_1^+ \setminus \Sigma_0(u), \mathcal{M})$ and the proof of Theorem 1 is complete.

4. FULL BOUNDARY REGULARITY

In this section we recover a regularity criterion based on the result in Theorem 1. The main preliminary step consists in proving compactness of sequences of minimizers of (3.1) under uniform assumptions, see [9, 13, 47].

Remark 4.1. We will always assume that $\gamma_2 < n$, otherwise, as stressed in *Step 1* of the proof of Theorem 1, we would have u Hölder continuous in a small neighborhood of any point $\bar{x} \in \bar{B}_1^+$ so that $p(\bar{x}) \geq n$ for free by Morrey's embedding theorem.

Lemma 4.1. *Let $\{k_j\}, \{p_j\}$ be two sequences of Hölder continuous functions satisfying*

$$(4.1) \quad \begin{cases} \sup_{j \in \mathbb{N}} [k_j]_{0,\nu} < c_k \quad \text{for some } \nu \in (0, 1] \\ \lambda \leq k_j(x) \leq \Lambda \quad \text{for all } x \in \bar{B}_1^+ \\ \|k_j - k_0\|_{L^\infty(\bar{B}_1^+)} \rightarrow 0, \quad k_0(\cdot) \in C^{0,\nu}(\bar{B}_1^+) \end{cases}$$

and

$$(4.2) \quad \begin{cases} \sup_{j \in \mathbb{N}} [p_j]_{0,\alpha} < c_p \quad \text{for some } \alpha \in (0, 1] \\ p_j(x) \geq \gamma_1 > 1 \quad \text{for all } x \in \bar{B}_1^+, j \in \mathbb{N} \\ \|p_j - p_0\|_{L^\infty(\bar{B}_1^+)} \rightarrow 0, \quad p_0 \geq \gamma_1 > 1 \text{ constant,} \end{cases}$$

respectively. For each $j \in \mathbb{N}$, let $u_j \in W^{1,p_j(\cdot)}(B_1^+, \mathcal{M})$ be a constrained minimizer of

$$\mathcal{E}_j(w, B_1^+) := \int_{B_1^+} k_j(x) |Dw|^{p_j(x)} \, dx,$$

in class $C_{g_j}^{p_j(\cdot)}(B_1^+, \mathcal{M})$, where the manifold \mathcal{M} is as in (2.6) and the sequence $\{g_j\} \subset W^{1,q}(\bar{B}_1^+, \mathcal{M})$, uniformly satisfying (2.7), is weakly convergent to some $g_0 \in W^{1,q}(\bar{B}_1^+, \mathcal{M})$. Then, there exists a subsequence, still denoted by $\{u_j\}$, such that

$$(4.3) \quad u_j \rightharpoonup u_0 \quad \text{weakly in } W^{1,(1+\tilde{\sigma})p_0}(B_R^+, \mathcal{M})$$

for some $\tilde{\sigma} > 0$ and any $R \in (0, 1)$. In particular, u_0 is a constrained minimizer of the functional

$$\mathcal{E}_0(w, B_R^+) := \int_{B_R^+} k_0(x) |Dw|^{p_0} \, dx$$

in class $C_{g_0}^{p_0}(B_R^+, \mathcal{M})$. Moreover,

$$\mathcal{E}_j(u_j, B_R^+) \rightarrow \mathcal{E}_0(u_0, B_R^+) \quad \text{for all } R \in (0, 1).$$

Finally, if x_j is a singular point of u_j and $x_j \rightarrow x_0$, then x_0 is a singular point for u_0 .

Proof. For the reader's convenience, we split the proof into three steps.

Step 1: Weak convergence. By assumption, the sequence $\{g_j\}$ is weakly convergent in $W^{1,q}(\bar{B}_1, \mathcal{M})$, so, we can find a positive, finite constant $M \equiv M(n, \mathcal{M}, q)$ so that

$$(4.4) \quad \sup_{j \in \mathbb{N}} \|g_j\|_{W^{1,q}(B_1^+)} \leq M.$$

Since the whole sequence $\{u_j\}$ has image contained into \mathcal{M} , which, by (2.6)₁ is compact, we immediately have that $\sup_{j \in \mathbb{N}} \|u_j\|_{L^\infty(B_1^+)} \leq c(\mathcal{M}) < \infty$, thus, up to extracting a non-relabelled subsequence,

$$(4.5) \quad u_j \rightharpoonup u_0 \quad \text{weakly in } L^t(B_1^+, \mathcal{M}) \quad \text{for all } t \in (1, \infty).$$

Moreover, being the assumptions in (4.1)-(4.2) uniform in $j \in \mathbb{N}$, we deduce that Lemma 3.1 and Lemma 3.2 for the associated frozen problem hold with constants independent from j . In particular, recalling the uniform features of the g_j 's and combining (3.8) with a standard covering argument we can conclude that $\{u_j\} \subset W_{loc}^{1,p(\cdot)(1+\sigma)}(B_1^+, \mathcal{M})$ for all $\sigma \in \left[0, \min\{\sigma_g, \sigma'_g\}\right)$. Now we take any ball $B_\varrho(x_0) \subset B_1$ with $\varrho \in (0, \frac{1}{4} \min\{1 - |x_0|, R_*\}]$ and R_* as in (3.21), so we can apply (3.16)₂ with any $\sigma \in \left(0, \min\left\{\sigma_g, \frac{n-\gamma_2}{\gamma_2}, \frac{q}{n} - 1\right\}\right)$ to deduce that

$$(4.6) \quad \int_{B_\varrho(x_0) \cap B_1^+} |Du_j|^{p_j(x)(1+\sigma)} dx \leq c\varrho^{n-p_2(x_0, \varrho)(1+\sigma)} \leq c(n, N, \mathcal{M}, \gamma_1, \gamma_2, q).$$

In (4.6) we also used (4.4) to incorporate the dependency of the constant from $\|Dg_j\|_{L^q(B_1^+)}$ into the one from (n, \mathcal{M}, q) . Now set

$$\hat{\sigma}_g := \frac{1}{4} \min \left\{ \sigma_g, \sigma'_g, \delta_g, \frac{n-\gamma_2}{\gamma_2}, \frac{q}{n} - 1 \right\},$$

where σ_g, σ'_g and δ_g are the same higher integrability threshold determined in Lemmas 3.1, 3.2 respectively and choose any $\sigma \in (0, \hat{\sigma}_g)$. Because of the uniform convergence of the p_j 's to the constant p_0 , taking $j \in \mathbb{N}$ sufficiently large we can find positive constants $\gamma_1 \leq q_1 \leq q_2 \leq \gamma_2$ such that

$$(4.7) \quad 1 < q_1 \leq p_j(\cdot) \leq q_2 < \infty \text{ on } \bar{B}_1^+, \quad q_2 \left(1 + \frac{\sigma}{2}\right) \leq q_1(1 + \sigma), \quad q_2 \leq p_0 \left(1 + \frac{\sigma}{2}\right)$$

and

$$(4.8) \quad 0 \leq q_2 - q_1 < \frac{\delta_g \gamma_1}{16} \quad \text{and} \quad 1 \leq \frac{q_2}{q_1} < 2.$$

Combining (4.6), (4.7) and the choice of $\sigma > 0$ we made, we can conclude that

$$(4.9) \quad \int_{B_\varrho(x_0) \cap B_1^+} |Du_j|^{q_2(1+\frac{\sigma}{2})} dx \leq c(n, \mathcal{M}, \gamma_1, \gamma_2, q).$$

By (4.5) and (4.9) we derive the uniform boundedness of the u_j 's in $W^{1,(1+\sigma/2)q_2}(B_\varrho(x_0) \cap B_1^+, \mathcal{M})$, so, up to extract a (non relabelled) subsequence, we obtain that $u_j \rightharpoonup \bar{u}_0$ weakly in $W^{1,(1+\sigma/2)q_2}(B_\varrho(x_0) \cap B_1^+, \mathcal{M})$, for some $\bar{u}_0 \in W^{1,(1+\sigma/2)q_2}(B_\varrho(x_0) \cap B_1^+, \mathcal{M})$. Anyway, by (4.5), $\bar{u}_0(x) = u_0(x)$, $u_0(x) \in \mathcal{M}$ for a.e. $x \in B_\varrho(x_0) \cap B_1^+$ and, by Rellich-Kondrachov theorem,

$$(4.10) \quad u_j \rightarrow u_0 \quad \text{strongly in } L^{(1+\sigma/2)q_2}(B_\varrho(x_0) \cap B_1^+, \mathcal{M}),$$

$$(4.11) \quad Du_j \rightharpoonup Du_0 \quad \text{weakly in } L^{(1+\sigma/2)q_2}(B_\varrho(x_0) \cap B_1^+, \mathbb{R}^{N \times n}).$$

From (4.7)₁ and (4.2)₃, we see that $q_2 \geq p_0$, therefore (4.3) is proved for instance with

$$(4.12) \quad \tilde{\sigma} = \frac{\hat{\sigma}_g}{4}.$$

Using the lower semicontinuity of the norm, we also have that

$$(4.13) \quad \int_{B_\varrho(x_0) \cap B_1^+} |Du_0|^{q_2(1+\frac{\tilde{\sigma}}{2})} dx \leq c(\mathbf{data}_{p(\cdot)}).$$

Inequality (4.13) and the convergence in (4.10)-(4.11) hold on $B_\varrho(x_0) \cap B_1^+$, but can show that they actually hold on half balls having any radius $R \in (0, 1)$. In fact, being B_1^+ compact, we can find $m \equiv m(n)$ and a finite family of balls $\{B_{\varrho_k}(x_k)\}_{k=1}^m$ so that $\{\varrho_k\} \subset (0, \frac{R_*}{4})$ and $B_1^+ \subseteq \bigcup_{k=1}^m B_{\varrho_k}(x_k)$. Then, given any measurable subset $U \subseteq B_R^+$ with $R \in (0, 1)$, we trivially have that $U \subseteq \bigcup_{k=1}^m (B_{\varrho_k}(x_k) \cap B_1^+)$ and, recalling (4.10), (4.11) and (4.13),

$$(4.14) \quad \int_U |Du_0|^{q_2(1+\frac{\sigma}{2})} dx \leq \sum_{k=1}^m \int_{B_{\varrho_k}(x_k) \cap B_1^+} |Du_0|^{q_2(1+\frac{\sigma}{2})} dx \leq mc \leq c(\mathbf{data}_{p(\cdot)})$$

$$(4.15) \quad \|Du_j\|_{L^{q_2(1+\frac{\sigma}{2})}(U)} \leq \sum_{k=1}^m \|Du_j\|_{L^{q_2(1+\frac{\sigma}{2})}(B_{\varrho_k}(x_k) \cap B_1^+)} \leq c(\mathbf{data}_{p(\cdot)})$$

$$(4.16) \quad \|u_j - u_0\|_{L^{q_2(1+\frac{\sigma}{2})}(U)} \leq \sum_{k=1}^m \|u_j - u_0\|_{L^{q_2(1+\frac{\sigma}{2})}(B_{\varrho_k}(x_k) \cap B_1^+)} \rightarrow 0,$$

so (4.3) is completely proved. Notice that, (4.3) and the weak continuity of the trace operator yield in particular that

$$(4.17) \quad \mathbf{tr}_{\Gamma_R}(u_0) = \mathbf{tr}_{\Gamma_R}(g_0) \quad \text{for all } R \in (0, 1).$$

Step 2: Compactness. We fix $R \in (0, 1)$ and, as a first step towards the proof of the minimality of $\mathcal{E}_0(u_0, B_R^+)$ in class $C_{g_0}^{p_0}(B_R^+, \mathcal{M})$ we show that

$$(4.18) \quad \mathcal{E}_0(u_0, B_R^+) \leq \liminf_{j \rightarrow \infty} \mathcal{E}_j(u_j, B_R^+).$$

Since

$$\mathcal{E}_j(u_j, B_R^+) = \left(\mathcal{E}_j(u_j, B_R^+) - \mathcal{E}_0(u_j, B_R^+) \right) + \mathcal{E}_0(u_j, B_R^+)$$

and, by weak lower semicontinuity and (4.5) there holds that

$$(4.19) \quad \mathcal{E}_0(u_0, B_R^+) \leq \liminf_{j \rightarrow \infty} \mathcal{E}_0(u_j, B_R^+),$$

we only need to show that

$$(4.20) \quad |\mathcal{E}_j(u_j, B_R^+) - \mathcal{E}_0(u_j, B_R^+)| \rightarrow 0,$$

which is a consequence of (4.1)₃, (4.2)₃, Lemma 2.2 (i.) with $\varepsilon_0 = \frac{\sigma}{2}$ and (4.9). In fact,

$$\begin{aligned} |\mathcal{E}_j(u_j, B_R^+) - \mathcal{E}_0(u_j, B_R^+)| &\leq \left| \int_{B_R^+} (k_j(x) - k_0(x)) |Du_j|^{p_j(x)} dx \right| \\ &+ \left| \int_{B_R^+} k_0(x) \left[|Du|^{p_j(x)} - |Du_j|^{p_0} \right] dx \right| \leq \|k_j - k_0\|_{L^\infty(B_R^+)} \int_{B_R^+} |Du|^{p_j(x)} dx \\ &+ c \|p_j - p_0\|_{L^\infty(B_R^+)} \int_{B_R^+} (1 + |Du_j|^2)^{\frac{q_2}{2}(1+\frac{\sigma}{2})} dx \\ &\leq c \left[\|k_j - k_0\|_{L^\infty(B_R^+)} + \|p_j - p_0\|_{L^\infty(B_R^+)} \right] \rightarrow 0. \end{aligned}$$

The constant c appearing in the previous display depends only from $\mathbf{data}_{p(\cdot)}$. Combining (4.20) and (4.19) we end up with (4.18). Now, let $\tilde{u}_0 \in W^{1,p_0}(B_R^+, \mathcal{M})$ be a solution of the Dirichlet problem

$$(4.21) \quad \hat{C}_{u_0}^{p_0}(B_R^+, \mathcal{M}) \ni w \mapsto \min \mathcal{E}_0(w, B_R^+).$$

As in [9, 13, 29] we fix any $\theta \in (0, 1)$, a cut-off function $\eta \in C_c^1(B_R)$ satisfying

$$(4.22) \quad \mathbf{1}_{B_{(1-\theta)R}} \leq \eta \leq \mathbf{1}_{B_R} \quad \text{and} \quad |D\eta| \lesssim \frac{1}{R\theta},$$

and consider a bi-Lipschitz transformation $\Phi: \bar{B}_R^+ \rightarrow \bar{B}_R$ so that

$$(4.23) \quad \Phi|_{\partial^+ B_R^+} = \mathbb{I}_{\partial^+ B_R^+} \quad \text{and} \quad \Phi(\Gamma_R) = \{x \in \partial B_R: x^n < 0\}.$$

Being Φ bi-Lipschitz, if \mathcal{G}_Φ is its jacobian, we have that

$$(4.24) \quad 0 < c(n)^{-1} \leq |\mathcal{G}_\Phi(x)| \leq c(n) < \infty.$$

Let us look at the function

$$\tilde{u}_j(x) := \tilde{u}_0(x) + (1 - \eta(\Phi(x))) (u_j(x) - u_0(x)) \quad \text{for } x \in B_R^+.$$

By (4.22)₁, (4.23) and (4.17) we see that

$$(4.25) \quad B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\} \subseteq B_R^+ \cap \Phi^{-1}(\bar{B}_R \setminus \bar{B}_{(1-\theta)R}),$$

and since,

$$\partial \left(B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\} \right) = \partial B_R^+ \cup \partial \{ \eta(\Phi(x)) = 1 \},$$

also that

$$(4.26) \quad \text{in a neighborhood of } \partial \left(B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\} \right), \tilde{u}_j \text{ takes values in } \mathcal{M}.$$

In particular, according to (4.17) and to the definition given in (4.21),

$$(4.27) \quad \text{tr}_{\Gamma_R}(\tilde{u}_j) = \text{tr}_{\Gamma_R}(g_j), \quad \text{tr}_{\partial^+ B_R^+} \tilde{u}_j = \text{tr}_{\partial^+ B_R^+}(u_j), \quad \text{tr}_{\partial \{ \eta(\Phi(x)) = 1 \}} = \text{tr}_{\partial \{ \eta(\Phi(x)) = 1 \}}(\tilde{u}_0).$$

Conditions (4.26)-(4.27) legalize the application of Lemma 2.5 on the set $B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\}$ to end up with a function $\bar{w}_j \in W_{loc}^{1,p_j(\cdot)}(B_1^+, \mathcal{M})$ satisfying

$$(4.28) \quad \begin{cases} \bar{w}_j \left(\partial \left(B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\} \right) \right) \subset \mathcal{M} \\ \text{tr}_{\Gamma_R}(\bar{w}_j) = \text{tr}_{\Gamma_R}(g_j) \\ \text{tr}_{\partial^+ B_R^+}(\bar{w}_j) = \text{tr}_{\partial^+ B_R^+}(u_j) \\ \text{tr}_{\partial \{ \eta(\Phi(x)) = 1 \}}(\bar{w}_j) = \text{tr}_{\partial \{ \eta(\Phi(x)) = 1 \}}(g_0), \\ \int_{B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\}} |D\bar{w}_j|^{p_j(x)} dx \lesssim \int_{B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\}} |D\tilde{u}_j|^{p_j(x)} dx \end{cases}$$

with constants implicit in " \lesssim " depending by $(N, \mathcal{M}, \gamma_2)$. Finally, define

$$\tilde{w}_j(x) := \begin{cases} \tilde{u}_0(x) & \text{if } x \in B_R^+ \cap \{ \eta(\Phi(x)) = 1 \} \\ \bar{w}_j(x) & \text{if } x \in B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\}. \end{cases}$$

Now, notice that the choices we made in (4.8) and (4.12) imply that

$$(4.29) \quad \frac{q_2}{p_0} \left(1 + \frac{\tilde{\sigma}}{2} \right) = 1 + \left[\frac{q_2 - p_0}{p_0} + \frac{q_2 \tilde{\sigma}}{2p_0} \right] < 1 + \frac{\delta_g}{8},$$

so by Lemma 3.2, (4.29), (4.14) and the minimality of \tilde{u}_0 in class $\hat{\mathcal{C}}_{u_0}^{p_0}(B_R^+, \mathcal{M})$, we get

$$(4.30) \quad \begin{aligned} & \int_{B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\}} |D\tilde{u}_0|^{p_j(x)} dx \leq |B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\}| \\ & \quad + \int_{B_R^+} |D\tilde{u}_0|^{q_2(1+\frac{\tilde{\sigma}}{2})} dx \\ & \leq |B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\}| + c \int_{B_R^+} |Du_0|^{q_2(1+\frac{\tilde{\sigma}}{2})} dx < \infty, \end{aligned}$$

for $c \equiv c(\text{data}_{p(\cdot)})$. In (4.30) we used, in particular, that

$$(4.31) \quad \int_{B_R^+} |D\tilde{u}_0|^{q_2(1+\frac{\tilde{\sigma}}{2})} dx \leq c \int_{B_R^+} |Du_0|^{q_2(1+\frac{\tilde{\sigma}}{2})} dx.$$

Via (4.24), (4.25) and a straightforward change of variables we have

$$\begin{aligned}
 (4.32) \quad & \left| B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\} \right| = \int_{B_R^+} \mathbb{1}_{\{0 \leq \eta(\Phi(x)) < 1\}} \, dx \\
 & \leq \int_{B_R^+ \cap \{\Phi^{-1}(\bar{B}_R \setminus B_{(1-\theta)R})\}} \, dx \leq \int_{B_R \setminus \bar{B}_{(1-\theta)R}} |\mathcal{G}_\Phi(x)|^{-1} \, dx \\
 & \leq c(n) |\bar{B}_R \setminus \bar{B}_{(1-\theta)R}| \rightarrow 0 \quad \text{as } \theta \rightarrow 0.
 \end{aligned}$$

We then estimate

$$\begin{aligned}
 (4.33) \quad & \mathcal{E}_j(u_j, B_R^+) \leq \mathcal{E}_j(\tilde{u}_j, B_R^+) \\
 & \leq \mathcal{E}_j(\tilde{u}_j, B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\}) + \mathcal{E}_j(\tilde{u}_0, B_R^+ \cap \{\eta(\Phi(x)) = 1\}) \\
 & =: (\text{I})_j + (\text{II})_j.
 \end{aligned}$$

In the previous display, we used that, in view of (4.28)_{2,3}, \tilde{u}_j is a legitimate comparison map to u_j . The bounds in (4.30), (4.28)₄ and (4.32) then legalize the following estimate:

$$\begin{aligned}
 (\text{I})_j & \leq c \int_{B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\}} \left[|D\tilde{u}_0|^{p_j(x)} + |Du_j - Du_0|^{p_j(x)} + \left| \frac{u_j - u_0}{R\theta} \right|^{p_j(x)} \right] \, dx \\
 & \leq c \int_{B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\}} |D\tilde{u}_0|^{p_j(x)} \, dx + c \int_{B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\}} \left[|Du_j|^{p_j(x)} + |Du_0|^{p_j(x)} \right] \, dx \\
 & \quad + c \int_{B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\}} \left| \frac{u_j - u_0}{R\theta} \right|^{p_j(x)} \, dx =: c \left[(\text{I})_j^1 + (\text{I})_j^2 + (\text{I})_j^3 \right]
 \end{aligned}$$

where $c \equiv c(N, \mathcal{M}, \gamma_1, \gamma_2)$. Let us bound the three terms appearing on the right-hand side of the above inequality. By Lemma 2.2 (i.) with $\varepsilon_0 = \frac{\tilde{\sigma}}{2}$, (4.31), (4.7), (3.19), (4.32), (4.2)₃ and the absolute continuity of Lebesgue's integral we have

$$\begin{aligned}
 (\text{I})_j^1 & \leq c \int_{B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\}} \left[|D\tilde{u}_0|^{p_j(x)} - |D\tilde{u}_0|^{p_0(x)} \right] \, dx + c \int_{B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\}} |D\tilde{u}_0|^{p_0} \, dx \\
 & \leq c \|p_j - p_0\|_{L^\infty(B_1^+)} \int_{B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\}} |D\tilde{u}_0|^{q_2(1+\frac{\tilde{\sigma}}{2})} \, dx + o(\theta) \\
 & \leq c \|p_j - p_0\|_{L^\infty(B_1^+)} \int_{B_R^+} |Du_0|^{q_2(1+\frac{\tilde{\sigma}}{2})} \, dx + o(\theta) = o(j) + o(\theta),
 \end{aligned}$$

with $c \equiv c(\text{data}_{p(\cdot)})$. By (4.7), (4.14), (4.15), (4.32) we get that

$$(\text{I})_j^2 = o(\theta).$$

Moreover, using (4.16), Hölder inequality and (4.32) we have

$$\begin{aligned}
 (\text{I})_j^3 & \leq |B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\}| + \int_{B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\}} \left| \frac{u_j - u_0}{r\theta} \right|^{q_2} \, dx \\
 & \leq o(\theta) + (R\theta)^{-q_2} |B_R^+ \cap \{0 \leq \eta(\Phi(x)) < 1\}|^{1 - \frac{q_2}{q_2(1+\frac{\tilde{\sigma}}{2})}} \|u_j - u_0\|_{L^{q_2(1+\frac{\tilde{\sigma}}{2})}(B_R^+)}^{q_2} \\
 & \leq o(\theta) + (R\theta)^{-q_2} o(j),
 \end{aligned}$$

and, trivially,

$$(\text{II})_j \leq \mathcal{E}_j(\tilde{u}_0, B_R^+).$$

Finally, by (4.1)₃, (4.2)₃, (4.30) and (4.31) we get

$$\begin{aligned}
 |\mathcal{E}_j(\tilde{u}_0, B_R^+) - \mathcal{E}_0(\tilde{u}_0, B_R^+)| & \leq \left[\|k_j - k_0\|_{L^\infty(B_1^+)} + \|p_j - p_0\|_{L^\infty(B_1^+)} \right] \left(1 + \int_{B_R^+} |Du_0|^{q_2(1+\frac{\tilde{\sigma}}{2})} \, dx \right) \\
 & \leq c(\text{data}_{p(\cdot)}) \left[\|k_j - k_0\|_{L^\infty(B_1^+)} + \|p_j - p_0\|_{L^\infty(B_1^+)} \right] = o(j).
 \end{aligned}$$

Plugging the content of all the previous estimates in (4.33) we end up with

$$\varepsilon_j(u_j, B_R^+) \leq \varepsilon_0(\tilde{u}_0, B_R^+) + o(j) + o(\theta) + (R\theta)^{-q_2} o(j).$$

By (4.18) we can take the liminf as $j \rightarrow \infty$ in the above display to obtain

$$\begin{aligned} \varepsilon_0(u_0, B_R^+) &\leq \liminf_{j \rightarrow \infty} \varepsilon_j(u_j, B_R^+) \\ &\leq \limsup_{j \rightarrow \infty} \left[\varepsilon_0(\tilde{u}_0, B_R^+) + o(j) + o(\theta) + (R\theta)^{-q_2} o(j) \right] \\ (4.34) \quad &\leq \varepsilon_0(\tilde{u}_0, B_R^+) + o(\theta). \end{aligned}$$

Sending $\theta \rightarrow 0$ in (4.34) and using the minimality of \tilde{u}_0 in class $\hat{\mathcal{C}}_{u_0}^{p_0}(B_R^+, \mathcal{M})$, we end up with

$$\varepsilon_0(u_0, B_R^+) \leq \varepsilon_0(\tilde{u}_0, B_R^+) \leq \varepsilon_0(w, B_R^+)$$

for all $w \in \hat{\mathcal{C}}_{u_0}^{p_0}(B_R^+, \mathcal{M})$, so, by Definition 3 and (4.17), the minimality of u_0 in class $\mathcal{C}_{g_0}^{p_0}(B_R^+, \mathcal{M})$ is proved. Finally, combining (4.34) with the minimality of \tilde{u}_0 in class $\hat{\mathcal{C}}_{u_0}^{p_0}(B_R^+, \mathcal{M})$, we can conclude that $\varepsilon_j(u_j, B_R^+) \rightarrow \varepsilon_0(u_0, B_R^+)$.

Step 3. Singular points. Let $\{x_j\} \subset \bar{B}_1^+$ be the sequence of singular points in the statement. The interior case $x_0 \in B_1^+$ has already been analyzed in [9, Section 4.1], so we can assume that $x_0 \in \Gamma_1$. Up to choose $j \in \mathbb{N}$ sufficiently large and then relabel, we can also suppose that $\{x_j\} \subset B_R^+$ for some $R \in (0, \frac{R_*}{2})$, $x_0 \in \Gamma_R$ and (4.7)-(4.8) are in force. By Theorem 1 and (4.4), we can find a radius $\tilde{R} > 0$ and a positive constant $\tilde{\varepsilon}$, both independent on $j \in \mathbb{N}$ so that if x_j is a singular point of u_j , then

$$(4.35) \quad \left(\varrho^{p_{2,j}(\varrho)-n} \int_{B_\varrho^+(x_j)} (1 + |Du_j|^2)^{\frac{p_{2,j}(\varrho)}{2}} dx \right)^{\frac{1}{p_{2,j}(\varrho)}} > \tilde{\varepsilon} > 0$$

for all $\varrho \in (0, \frac{1}{2} \min \{ \tilde{R}, R_* - R \})$, with R_* as in (3.21). In the above display, we denoted $p_{2,j}(\varrho) := \sup_{x \in B_\varrho(x_j) \cap B_R^+} p_j(x)$. Set $\sigma' := \min \{ \tilde{\sigma}, \frac{\alpha}{\gamma_2} \}$. By Lemma 2.2 (i.) with $\varepsilon_0 = \frac{\sigma'}{2}$ and (3.16)₂, we estimate

$$\begin{aligned} &\left| \varrho^{p_{2,j}(\varrho)-n} \int_{B_\varrho^+(x_j)} \left[(1 + |Du_j|^2)^{\frac{p_{2,j}(\varrho)}{2}} - (1 + |Du_j|^2)^{\frac{p_j(x)}{2}} \right] dx \right|^{\frac{1}{p_{2,j}(\varrho)}} \\ (4.36) \quad &\leq c\varrho^{1+\frac{\alpha}{\gamma_1}} \left(\int_{B_\varrho^+(x_j)} (1 + |Du_j|^2)^{\frac{p_{2,j}(\varrho)}{2} \left(1 + \frac{\sigma'}{2} \right)} dx \right)^{\frac{1}{p_{2,j}(\varrho)}} \leq c\varrho^{-\frac{\sigma'}{2} + \frac{\alpha}{\gamma_2}} \rightarrow 0, \end{aligned}$$

for $c \equiv c(n, N, \mathcal{M}, \gamma_1, \gamma_2, q)$. By (4.35), (4.36), (4.4) and (3.14) we then get

$$\begin{aligned} \tilde{\varepsilon} &< c\varrho^{-\frac{\sigma'}{2} + \frac{\alpha}{\gamma_2}} + c \left(\varrho^{p_{2,j}(\varrho)-n} \int_{B_\varrho^+(x_j)} (1 + |Du_j|^2)^{\frac{p_j(x)}{2}} dx \right)^{\frac{1}{p_{2,j}(\varrho)}} \\ &\leq c\varrho^{-\frac{\sigma'}{2} + \frac{\alpha}{\gamma_2}} + c\varrho + c\varrho^{1-\frac{n}{p_{2,j}(\varrho)}} \left[\int_{B_{2\varrho}^+(x_j)} \left| \frac{u_j - g_j}{\varrho} \right|^{p_j(x)} dx + \int_{B_{2\varrho}^+(x_j)} |Dg_j|^{p_j(x)} dx \right]^{\frac{1}{p_{2,j}(\varrho)}} \\ &\leq c\varrho^{-\frac{\sigma'}{2} + \frac{\alpha}{\gamma_2}} + c\varrho + c\varrho^{1-\frac{n}{q}} \left(\int_{B_\varrho^+(x_j)} |Dg_j|^q dx \right)^{\frac{1}{q}} \\ &\quad + c\varrho^{1-\frac{n}{p_{2,j}(\varrho)}} \left[\int_{B_{2\varrho}^+(x_j)} \left| \frac{u_j - u_0}{\varrho} \right|^{q_2 \left(1 + \frac{\tilde{\sigma}}{2} \right)} dx + \int_{B_{2\varrho}^+(x_j)} \left| \frac{g_j - u_0}{\varrho} \right|^{p_j(x)} dx \right]^{\frac{1}{p_{2,j}(\varrho)}} \end{aligned}$$

(4.37)

$$\leq c\varrho^{\sigma''} + c \left[\int_{B_{2\varrho}^+(x_j)} |u_j - u_0|^{q_2(1+\frac{\tilde{\sigma}}{2})} dx + \int_{B_{2\varrho}^+(x_j)} |g_j - u_0|^{p_j(x)} dx \right]^{\frac{1}{p_{2,j}(\varrho)}},$$

where we set $\sigma'' := \min \left\{ 1 - \frac{n}{q}, \frac{\alpha}{\gamma_2} - \frac{\sigma'}{2} \right\}$ and $c \equiv c(n, N, M, \gamma_1, \gamma_2, q)$. By (4.16) we get

$$(4.38) \quad \int_{B_{2\varrho}^+(x_j)} |u_j - u_0|^{q_2(1+\frac{\tilde{\sigma}}{2})} dx \rightarrow 0 \quad \text{as } j \rightarrow \infty.$$

Since $g_j \rightharpoonup g_0$ weakly in $W^{1,q}(\bar{B}_1^+, M)$, then by Rellich-Kondrachov theorem there holds that, up to subsequences, $g_j \rightarrow g_0$ strongly in $L^q(\bar{B}_1^+, M)$ and pointwise a.e., therefore, keeping also (4.2)₃ in mind, we can apply the dominated convergence theorem to end up with

$$(4.39) \quad \int_{B_{2\varrho}^+(x_j)} |g_j - u_0|^{p_j(x)} dx \rightarrow \int_{B_{2\varrho}^+(x_j)} |g_0 - u_0|^{p_0} dx \quad \text{as } j \rightarrow \infty.$$

By (4.2)₃, (4.38) and (4.39) we can take the limit superior on both sides of the inequality in (4.37) to obtain

$$(4.40) \quad \tilde{\varepsilon} \leq c\varrho^{\sigma''} + c \left(\int_{B_{2\varrho}^+(x_j)} |g_0 - u_0|^{p_0} dx \right)^{\frac{1}{p_0}}.$$

We finally pass to the limit superior for $\varrho \rightarrow \infty$ in (4.40) and have

$$0 < \bar{\varepsilon}^{p_0} \leq \limsup_{\varrho \rightarrow 0} \int_{B_{2\varrho}^+(x_j)} |u_0 - g_0|^{p_0} dx,$$

meaning that x_0 is a singular point for u_0 . In the previous display, we set $\bar{\varepsilon} := \tilde{\varepsilon}/c$. \square

The next lemma is a monotonicity formula in the spirit of [9, 22, 50, 52].

Lemma 4.2. *Under assumptions (2.3), (2.4), (2.6) and (2.7), let $u \in W^{1,p(\cdot)}(B_1^+, M)$ be a solution of problem (3.1). Suppose also that*

$$(4.41) \quad k(0) = 1.$$

Then, there exist $\Upsilon \equiv \Upsilon(n, N, M, \gamma_1, \gamma_2, q) \in (0, 1]$ and a threshold $T \equiv T(\mathbf{data}, \kappa) \in (0, 1]$ such that if

$$(4.42) \quad [g]_{0, 1-\frac{n}{q}; \bar{B}_1^+} < \Upsilon,$$

then for all $\kappa \in (0, 1 - \frac{n}{q}]$, the map $\Phi: (0, \frac{T}{4}) \rightarrow [0, \infty)$ defined as

$$(4.43) \quad \Phi(\tau) := \exp \left(\frac{\tilde{c}}{\beta''} \tau^{\beta''} \right) \left[\tau^{p_2(\tau)-n} \int_{B_1^+} k(x) |D\tilde{u}|^{p_2(\tau)} dx + c \frac{\tau^\kappa}{\kappa} \right],$$

with \tilde{u} as in (3.26), $\beta'' \equiv \beta''(n, q, \nu)$ and $c, \tilde{c} \equiv c, \tilde{c}(\mathbf{data}, \|Dg\|_{L^q(B_1^+)}, \kappa)$, is monotone non-decreasing. Moreover, the following inequality holds true

$$(4.44) \quad \int_{\partial B_1^+} |u(Rx) - u(\varrho x)|^{p_2(\varrho)} d\mathcal{H}^{n-1}(x) \leq c \log(R/\varrho) \left[\varrho^{p_2(\varrho)-p_2(R)} (\Phi(R) - \Phi(\varrho)) \right] + c(R - \varrho)^{\gamma_1(1-\frac{n}{q})},$$

for $c \equiv c(\mathbf{data}, \|Dg\|_{L^q(B_1^+)}, \kappa)$.

Proof. Let $u \in W^{1,p(\cdot)}(B_1^+, \mathcal{M})$ be a solution of problem (3.1), $\kappa \in (0, 1)$ be a fixed constant and select $T \in (0, 1]$ so that

$$0 < T \leq \min \left\{ R_*, \frac{1-\kappa}{16[p]_{0,1}}, \left(\frac{\lambda}{4[k]_{0,\nu}} \right)^{\frac{2}{\nu}} \right\},$$

where R_* is as in (3.21). Such a position assures that, whenever $\tau \in \left(0, \frac{T}{4}\right]$, (3.23)-(3.25) hold with R replaced by τ , moreover,

$$(4.45) \quad p_2(4\tau) - p_1(\tau) \leq \frac{1-\kappa}{2} \quad \text{and} \quad 4[k]_{0,\nu} \tau^{\frac{\kappa}{2}} \leq \lambda,$$

with ν as in (2.3)₁. For $\tau \in \left(0, \frac{T}{4}\right]$, we introduce the functional

$$W^{1,p_2(\tau)}(B_\tau^+, \mathcal{M}) \ni w \mapsto \mathcal{E}_\tau(w, B_\tau^+) := \int_{B_\tau^+} k(x) |Dw|^{p_2(\tau)} \, dx$$

and let $v \in W^{1,p_2(\tau)}(B_\tau^+, \mathcal{M})$ be a solution of problem

$$(4.46) \quad \hat{\mathcal{C}}_u^{p_2(\tau)}(B_\tau^+, \mathcal{M}) \ni w \mapsto \min \mathcal{E}_\tau(w, B_\tau^+).$$

By the minimality of v in class $\hat{\mathcal{C}}_u^{p_2(\tau)}(B_\tau^+, \mathcal{M})$ and that of u in class $\mathcal{C}_g^{p(\cdot)}(B_1^+, \mathcal{M})$ we bound

$$\begin{aligned} |\mathcal{E}_\tau(u, B_\tau^+) - \mathcal{E}_\tau(v, B_\tau^+)| &= \mathcal{E}_\tau(u, B_\tau^+) - \mathcal{E}_\tau(v, B_\tau^+) \\ &\leq |\mathcal{E}_\tau(u, B_\tau^+) - \mathcal{E}(u, B_\tau^+)| + |\mathcal{E}_\tau(v, B_\tau^+) - \mathcal{E}(v, B_\tau^+)| =: \text{(I)} + \text{(II)}. \end{aligned}$$

Let

$$(4.47) \quad \sigma'' := \frac{1}{4} \min \left\{ \sigma_g, \delta_g, \frac{n-\gamma_2}{\gamma_2}, \frac{1-\kappa}{2\gamma_2} \right\},$$

where σ_g and δ_g are the higher integrability threshold from Lemmas 3.1-3.2 respectively. Combining (2.2)₁, (2.3)₁, Lemma 3.1, Lemma 2.2 (i.) with $\varepsilon_0 = \sigma''$ and (3.16)₂ we end up with

$$\text{(I)} \leq c\tau \int_{B_\tau^+} (1 + |Du|^2)^{\frac{p_2(\tau)(1+\sigma'')}{2}} \, dx \leq c\tau^{1+n-p_2(4\tau)(1+\sigma'')},$$

with $c \equiv c(\text{data}_{p(\cdot)}, \|Dg\|_{L^q(B_1^+)})$. In a totally similar way, using this time Lemma 3.2, (3.16)₂ and Lemma 2.2 (ii.) with $\varepsilon_0 = \sigma''$ we get

$$\begin{aligned} \text{(II)} &\leq c\tau \int_{B_\tau^+} (1 + |Dv|^2)^{\frac{p_2(\tau)(1+\sigma'')}{2}} \, dx \\ &\leq c\tau \int_{B_\tau^+} (1 + |Dv|^2)^{\frac{p_2(\tau)(1+\sigma'')}{2}} \, dx \leq c\tau^{1+n-p_2(4\tau)(1+\sigma'')}, \end{aligned}$$

for $c \equiv c(\text{data}_{p(\cdot)}, \|Dg\|_{L^q(B_1^+)})$. Merging the content of the previous displays we obtain

$$(4.48) \quad \mathcal{E}_\tau(u, B_\tau^+) \leq \mathcal{E}_\tau(v, B_\tau^+) + c\tau^{1+n-p_2(4\tau)(1+\sigma'')}.$$

Now, for τ as above, define $x_\tau := \tau \frac{x}{|x|}$. As in [50, Lemma 1.3] we consider the following comparison map

$$w_\tau(x) := \begin{cases} u(x) & \text{if } x \in B_1^+ \setminus B_\tau^+ \\ \tilde{u}(x_\tau) + g(x) & \text{if } x \in B_\tau^+, \end{cases}$$

where \tilde{u} is defined in (3.26). Notice that, by (3.24)-(3.25) there holds that

$$(4.49) \quad w_\tau \in u + W_0^{1,p(\cdot)}(B_\tau^+, \mathbb{R}^N) \quad \text{and} \quad w_\tau \in W^{1,p_2(\tau)}(B_\tau^+, \mathbb{R}^N).$$

Moreover, since (2.11) and (4.42) are in force, we see that

$$\text{dist}(w_\tau, \mathcal{M}) \leq c(n, q, \beta_0) \Upsilon \tau^{\beta_0} \quad \text{with} \quad \beta_0 := 1 - \frac{n}{q},$$

therefore, choosing Υ small enough, and thus determining the dependency $\Upsilon \equiv \Upsilon(n, N, \mathcal{M}, \gamma_1, \gamma_2, q)$, we can project w_τ onto \mathcal{M} thus obtaining a map $\bar{w}_\tau := \Pi_{\mathcal{M}}(w_\tau)$ satisfying

$$(4.50) \quad \bar{w}_\tau \in \hat{\mathcal{C}}_u^{p_2(\tau)}(B_\tau^+, \mathcal{M}) \quad \text{and} \quad \int_{B_\tau^+} |D\bar{w}_\tau|^{p_2(\tau)} dx \leq (1 + c\Upsilon\tau^{\beta_0}) \int_{B_\tau^+} |Dw_\tau|^{p_2(\tau)} dx,$$

for $c \equiv c(n, N, \mathcal{M}, \gamma_1, \gamma_2, q)$. Notice that, by the mean value theorem applied to the function $[0, \infty) \ni s \mapsto (t+s)^{p_2(\tau)}$ there holds that

$$(4.51) \quad (|D\tilde{u}| + |Dg|)^{p_2(\tau)} \leq |D\tilde{u}|^{p_2(\tau)} + p_2(\tau)(|D\tilde{u}| + |Dg|)^{p_2(\tau)-1}|Dg|,$$

so by Hölder inequality with conjugate exponents $\left(\frac{p_2(\tau)}{p_2(\tau)-1}, p_2(\tau)\right)$, (4.42) and (4.50) we get

$$\begin{aligned} \int_{B_\tau^+} |Dw_\tau|^{p_2(\tau)} dx &\leq \int_{B_\tau^+} (|D\tilde{u}(x_\tau)| + |Dg|)^{p_2(\tau)} dx \\ &\leq (1 + c\tau^{\beta_0}) \int_{B_\tau^+} |D\tilde{u}(x_\tau)|^{p_2(\tau)} dx \\ &\quad + c \left[\tau^{-\beta_0(p_2(\tau)-1)} \int_{B_\tau^+} |Dg|^{p_2(\tau)} dx + \int_{B_\tau^+} |Dg|^{p_2(\tau)} dx \right] \\ &\leq (1 + c\tau^{\beta_0}) \int_{B_\tau^+} |D\tilde{u}(x_\tau)|^{p_2(\tau)} dx \\ &\quad + c \left[\tau^{-\beta_0(p_2(\tau)-1)+n\left(1-\frac{p_2(\tau)}{q}\right)} + \tau^{n\left(1-\frac{p_2(\tau)}{q}\right)} \right] \|Dg\|_{L^q(B_1^+)} \\ (4.52) \quad &\leq (1 + c\tau^{\beta_0}) \int_{B_\tau^+} |D\tilde{u}(x_\tau)|^{p_2(\tau)} dx + c\tau^{n\left(1-\frac{1}{q}\right)+1-p_2(\tau)} \end{aligned}$$

for $c \equiv c(n, \gamma_1, \gamma_2, q, \|Dg\|_{L^q(B_1^+)})$. In the previous expression, we also used the original value of β_0 . By (2.3), (4.41) and (4.50) we can refine (4.52) as

$$\begin{aligned} \int_{B_\tau^+} k(x)|D\bar{w}_\tau|^{p_2(\tau)} dx &\leq (1 + 4[k]_{0,\nu}\tau^\nu) \int_{B_\tau^+} |D\bar{w}_\tau|^{p_2(\tau)} dx \\ (4.53) \quad &\leq (1 + c\tau^{\beta'}) \int_{B_\tau^+} |D\tilde{u}(x_\tau)|^{p_2(\tau)} dx + c\tau^{n\left(1-\frac{1}{q}\right)+1-p_2(\tau)}, \end{aligned}$$

where $\beta' := \min\{\beta_0, \nu\}$ and $c \equiv c(n, N, \mathcal{M}, \gamma_1, \gamma_2, q, \|Dg\|_{L^q(B_1^+)})$. Let us evaluate the $p_2(\tau)$ -energy of \tilde{u} . First, recall that if $\frac{\partial \tilde{u}}{\partial r} := D\tilde{u} \cdot \frac{x}{|x|}$ denotes the radial derivative of \tilde{u} , then

$$(4.54) \quad \left| \frac{\partial \tilde{u}}{\partial r} \right| \leq |D\tilde{u}|.$$

Moreover, if $p_2(\tau) \geq 2$ and $t \geq s \geq 0$ there holds that

$$(4.55) \quad (t-s)^{p_2(\tau)} \leq t^{p_2(\tau)} - s^{p_2(\tau)}.$$

A straightforward computation renders, for $x \in B_\tau^+$ that

$$|D\tilde{u}(x_\tau)|^2 = \frac{\tau^2}{|x|^2} \left[|D\tilde{u}(x_\tau)|^2 - \left| D\tilde{u}(x_\tau) \cdot \frac{x_\tau}{|x_\tau|} \right|^2 \right],$$

so by (4.54), (4.55), area formula, (2.3), (4.41) and (4.45)₂ we obtain

$$\begin{aligned} \int_{B_\tau^+} |D\tilde{u}(x_\tau)|^{p_2(\tau)} dx &= \frac{\tau}{n-p_2(\tau)} \int_{\partial B_\tau^+} \left[|D\tilde{u}(x)|^2 - \left| \frac{\partial \tilde{u}}{\partial r} \right|^2 \right]^{\frac{p_2(\tau)}{2}} d\mathcal{H}^{n-1}(x) \\ &\leq \frac{\tau}{n-p_2(\tau)} \left[\int_{\partial B_\tau^+} |D\tilde{u}(x)|^{p_2(\tau)} - \left| \frac{\partial \tilde{u}}{\partial r} \right|^{p_2(\tau)} \right] d\mathcal{H}^{n-1}(x) \end{aligned}$$

$$(4.56) \quad \leq \frac{\tau}{n-p_2(\tau)} \left[(1 + \tau^{\frac{\nu}{2}}) \int_{\partial B_\tau^+} k(x) |D\tilde{u}(x)|^{p_2(\tau)} d\mathcal{H}^{n-1}(x) - \int_{\partial B_\tau^+} \left| \frac{\partial \tilde{u}}{\partial r} \right|^{p_2(\tau)} d\mathcal{H}^{n-1}(x) \right].$$

Recalling the position made in (3.26), by (4.51), Hölder inequality with conjugate exponents $\left(\frac{p_2(\tau)}{p_2(\tau)-1}, p_2(\tau)\right)$, (4.42), (2.3), (4.48) and the minimality of v in class $\hat{C}_u^{p_2(\tau)}(B_\tau^+, m)$ with (4.50)₁ we have

$$(4.57) \quad \begin{aligned} \mathcal{E}_\tau(\tilde{u}, B_\tau^+) &\leq (1 + c\tau^{\beta_0})\mathcal{E}_\tau(u, B_\tau^+) + c\tau^{-\beta_0(p_2(\tau)-1)} \int_{B_\tau^+} |Dg|^{p_2(\tau)} dx \\ &\leq (1 + c\tau^{\beta_0})\mathcal{E}_\tau(v, B_\tau^+) + c \left[\tau^{n(1-\frac{1}{q})+1-p_2(\tau)} + \tau^{n+1-p_2(4\tau)(1+\sigma'')} \right] \\ &\leq (1 + c\tau^{\beta_0})\mathcal{E}_\tau(\bar{w}_\tau, B_\tau^+) + c \left[\tau^{n(1-\frac{1}{q})+1-p_2(\tau)} + \tau^{n+1-p_2(4\tau)(1+\sigma'')} \right] \\ &\leq (1 + c\tau^{\beta'}) \int_{B_\tau^+} |D\tilde{u}(x_\tau)|^{p_2(\tau)} dx + c \left[\tau^{n(1-\frac{1}{q})+1-p_2(\tau)} + \tau^{n+1-p_2(4\tau)(1+\sigma'')} \right]. \end{aligned}$$

with $c(\mathbf{data}_{p(\cdot)}, \|Dg\|_{L^q(B_1^+)})$. Merging (4.57) with (4.56) and using (4.54), (2.3) and (4.45)₂ we obtain

$$(4.58) \quad \begin{aligned} \mathcal{E}_\tau(\tilde{u}, B_\tau^+) &\leq \frac{\tau}{n-p_2(\tau)} \left[(1 + c\tau^{\beta''}) \int_{\partial B_\tau^+} k(x) |D\tilde{u}|^{p_2(\tau)} d\mathcal{H}^{n-1}(x) \right. \\ &\quad \left. - \int_{\partial B_\tau^+} \left| \frac{\partial \tilde{u}}{\partial r} \right|^{p_2(\tau)} d\mathcal{H}^{n-1}(x) + c\tau^{\beta'} (\tau^{\frac{\nu}{2}} + 1) \int_{\partial B_\tau^+} k(x) |D\tilde{u}|^{p_2(\tau)} d\mathcal{H}^{n-1}(x) \right] \\ &\quad + c \left[\tau^{n(1-\frac{1}{q})+1-p_2(\tau)} + \tau^{n+1-p_2(4\tau)(1+\sigma'')} \right] \\ &\leq \frac{\tau(1 + c\tau^{\beta''})}{n-p_2(\tau)} \int_{\partial B_\tau^+} k(x) |D\tilde{u}|^{p_2(\tau)} d\mathcal{H}^{n-1}(x) \\ &\quad - \frac{\tau}{n-p_2(\tau)} \int_{\partial B_\tau^+} \left| \frac{\partial \tilde{u}}{\partial r} \right|^{p_2(\tau)} d\mathcal{H}^{n-1}(x) + c \left[\tau^{n(1-\frac{1}{q})+1-p_2(\tau)} + \tau^{n+1-p_2(4\tau)(1+\sigma'')} \right], \end{aligned}$$

with $\beta'' := \min\{\frac{\nu}{2}, \beta'\}$ and $c \equiv c(\mathbf{data}, \|Dg\|_{L^q(B_1^+)})$. To summarize, we got

$$(4.59) \quad \begin{aligned} \tau \int_{\partial B_\tau^+} k(x) |D\tilde{u}|^{p_2(\tau)} d\mathcal{H}^{n-1}(x) &\geq \frac{n-p_2(\tau)}{1 + c\tau^{\beta''}} \int_{B_\tau^+} k(x) |D\tilde{u}|^{p_2(\tau)} dx \\ &\quad + \frac{\tau}{1 + c\tau^{\beta''}} \int_{\partial B_\tau^+} \left| \frac{\partial \tilde{u}}{\partial r} \right|^{p_2(\tau)} d\mathcal{H}^{n-1}(x) \\ &\quad - \frac{c(n-p_2(\tau))}{1 + c\tau^{\beta''}} \left[\tau^{n(1-\frac{1}{q})+1-p_2(\tau)} + \tau^{n+1-p_2(4\tau)(1+\sigma'')} \right] \end{aligned}$$

for $c \equiv c(\mathbf{data}, \|Dg\|_{L^q(B_1^+)}, \beta_0)$. Now, set

$$(4.60) \quad \left(0, \frac{T}{4}\right) \ni \tau \mapsto f(\tau) := \tau^{p_2(\tau)-n} \int_{B_\tau^+} k(x) |D\tilde{u}|^{p_2(\tau)} dx.$$

Multiplying both sides of (4.59) by $\tau^{p_2(\tau)-n-1}$ and using (4.45)₁ and (4.47) we obtain

$$(4.61) \quad \begin{aligned} \tau^{p_2(\tau)-n} \int_{\partial B_\tau^+} k(x) |D\tilde{u}|^{p_2(\tau)} d\mathcal{H}^{n-1}(x) &\geq \frac{n-p_2(\tau)}{1 + c\tau^{\beta''}} \left[\tau^{-1} f(\tau) - c \left(\tau^{\kappa-1} + \tau^{-\frac{n}{q}} \right) \right] \\ &\quad + \frac{\tau^{p_2(\tau)-n}}{1 + c\tau^{\beta''}} \int_{\partial B_\tau^+} \left| \frac{\partial \tilde{u}}{\partial r} \right|^{p_2(\tau)} d\mathcal{H}^{n-1}(x) \end{aligned}$$

for $c \equiv c(\mathbf{data}, \|Dg\|_{L^q(B_1^+)}, \kappa)$. From (2.4) follows that $(0, \frac{T}{4}) \ni \tau \mapsto p_2(\tau)$ is differentiable with bounded, non-negative first derivative $0 \leq p'(\tau) \leq c(n, [p]_{0,1})$. We compute:

$$\begin{aligned} f'(\tau) &= (p_2(\tau) - n)\tau^{p_2(\tau)-n-1} \int_{B_\tau^+} k(x)|D\tilde{u}|^{p_2(\tau)} dx \\ &\quad + \tau^{p_2(\tau)-n} \int_{\partial B_\tau^+} k(x)|D\tilde{u}|^{p_2(\tau)} d\mathcal{H}^{n-1}(x) \\ &\quad + p'_2(\tau) \log(\tau) \tau^{p_2(\tau)-n} \int_{B_\tau^+} k(x)|D\tilde{u}|^{p_2(\tau)} dx \\ &\quad + p'_2(\tau) \tau^{p_2(\tau)-n} \int_{B_\tau^+} k(x) \log(|D\tilde{u}|) |D\tilde{u}|^{p_2(\tau)} dx. \end{aligned}$$

We record that, for all $\varepsilon_0 \in (0, 1)$ there holds that

$$(4.62) \quad |\log(t)| \leq c(\varepsilon_0)(1+t)t^{-\varepsilon_0} \quad \text{for any } t > 0.$$

Let us estimate the last two terms appearing in the expansion of $f'(\tau)$. Using (4.62) with $\varepsilon_0 = 1 - \beta''$ we bound

$$p'_2(\tau) \log(\tau) \tau^{p_2(\tau)-n} \int_{B_\tau^+} k(x)|D\tilde{u}|^{p_2(\tau)} dx \leq cp'_2(\tau) \tau^{\beta''-1+p_2(\tau)-n} \int_{B_\tau^+} k(x)|D\tilde{u}|^{p_2(\tau)} dx.$$

By (4.62) with $\varepsilon_0 = 1 - p_2(\tau) \min\left\{\frac{\sigma_0}{2}, \frac{1-\kappa}{2\gamma_2}\right\}$, (keep (3.22)-(3.24) in mind) and (3.16)₂ we obtain

$$\begin{aligned} p'_2(\tau) \tau^{p_2(\tau)-n} \int_{B_\tau^+} k(x) \log(|D\tilde{u}|) |D\tilde{u}|^{p_2(\tau)} dx &\leq c\tau^{p_2(\tau)} \int_{B_\tau^+} (1 + |D\tilde{u}|)^{p_2(\tau)(1+\varepsilon_0)} dx \\ &\leq c\tau^{p_2(\tau)-p_2(4\tau)(1+\varepsilon_0)} \leq c\tau^{\kappa-1}, \end{aligned}$$

for $c \equiv c(\mathbf{data}_{p(\cdot)}, \|Dg\|_{L^1(B_1^+)}, \kappa)$. All in all, we got the following lower bound for $f'(\tau)$:

$$\begin{aligned} f'(\tau) &\geq (p_2(\tau) - n)\tau^{p_2(\tau)-n-1} \int_{B_\tau^+} k(x)|D\tilde{u}|^{p_2(\tau)} dx \\ &\quad + \tau^{p_2(\tau)-n} \int_{\partial B_\tau^+} k(x)|D\tilde{u}|^{p_2(\tau)} d\mathcal{H}^{n-1}(x) \\ &\quad - cp'_2(\tau) \tau^{\beta''-1+p_2(\tau)-n} \int_{B_\tau^+} k(x)|D\tilde{u}|^{p_2(\tau)} dx - c\tau^{\kappa-1} \\ &= \tau^{p_2(\tau)-n} \int_{\partial B_\tau^+} k(x)|D\tilde{u}|^{p_2(\tau)} d\mathcal{H}^{n-1}(x) \\ (4.63) \quad &\quad + \left(p_2(\tau) - n - cp'_2(\tau) \tau^{\beta''}\right) \frac{f(\tau)}{\tau} - c\tau^{\kappa-1}, \end{aligned}$$

with $c \equiv c(\mathbf{data}, \|Dg\|_{L^q(B_1^+)}, \kappa)$. Set

$$\varphi(\tau) := n - p_2(\tau) + cp'_2(\tau) \tau^{\beta''}.$$

Merging (4.61) and (4.63) we obtain

$$\begin{aligned} f'(\tau) + \left(\varphi(\tau) - \frac{n - p_2(\tau)}{1 + c\tau^{\beta''}}\right) \frac{f(\tau)}{\tau} &\geq \frac{\tau^{p_2(\tau)-n}}{1 + c\tau^{\beta''}} \int_{\partial B_\tau^+} \left| \frac{\partial \tilde{u}}{\partial r} \right|^{p_2(\tau)} d\mathcal{H}^{n-1}(x) \\ &\quad - c\tau^{\kappa-1} \left[\frac{2(n - p_2(\tau))}{1 + c\tau^{\beta''}} + 1 \right], \end{aligned}$$

where $c \equiv c(\mathbf{data}, \|Dg\|_{L^q(B_1^+)}, \kappa)$. In the previous display we also used that $\kappa \leq \beta_0$. It is easy to see that

$$\left| \varphi(\tau) - \frac{n - p_2(\tau)}{1 + c\tau^{\beta''}} \right| \leq \tilde{c}(\mathbf{data}, \|Dg\|_{L^q(B_1^+)}, \kappa) \tau^{\beta''},$$

therefore we get

$$(4.64) \quad f'(\tau) + \tilde{c}\tau^{\beta''-1}f(\tau) + c\tau^{\kappa-1} \geq \frac{\tau^{p_2(\tau)-n}}{1+c\tau^{\beta''}} \int_{\partial B_r^+} \left| \frac{\partial \tilde{u}}{\partial r} \right|^{p_2(\tau)} d\mathcal{H}^{n-1}(x).$$

Let $\Phi(\cdot)$ be the function defined in (4.43). Combining (4.64) and the fact that $\tau \in (0, 1]$, we immediately see that

$$\begin{aligned} \Phi'(\tau) &\geq \exp \left\{ \frac{\tilde{c}\tau^{\beta''}}{\beta''} \right\} \left[\tilde{c}\tau^{\beta''-1}f(\tau) + f'(\tau) + c\tau^{\kappa-1} \right] \\ &\geq \frac{\tau^{p_2(\tau)-n}}{1+c} \int_{\partial B_r^+} \left| \frac{\partial \tilde{u}}{\partial r} \right|^{p_2(\tau)} d\mathcal{H}^{n-1}(x), \end{aligned}$$

with $c \equiv c(\mathbf{data}, \|Dg\|_{L^q(B_1^+)}, \kappa)$. At this stage, we integrate the inequality in the previous display over $\tau \in (\varrho, R)$ with $0 < \varrho < R \leq T \leq 1$ to get

$$\begin{aligned} \Phi(R) - \Phi(\varrho) &\geq \frac{1}{1+c} \int_{\varrho}^R \tau^{p_2(\tau)-n} \left(\int_{\partial B_r^+} \left| \frac{\partial \tilde{u}}{\partial r} \right|^{p_2(\tau)} d\mathcal{H}^{n-1}(x) \right) d\tau \\ (4.65) \quad &\geq \frac{\varrho^{p_2(R)-p_2(\varrho)}}{1+c} \int_{\varrho}^R \tau^{p_2(\varrho)-n} \left(\int_{\partial B_r^+} \left| \frac{\partial \tilde{u}}{\partial r} \right|^{p_2(\tau)} d\mathcal{H}^{n-1}(x) \right) d\tau. \end{aligned}$$

Once (4.65) is available, we can proceed exactly as in [52, Lemma 4.1] to end up with

$$\begin{aligned} &\int_{\partial B_1^+} |\tilde{u}(Rx) - \tilde{u}(\varrho x)|^{p_2(\varrho)} d\mathcal{H}^{n-1}(x) \\ &\leq \log(R/\varrho) \int_{\varrho}^R \tau^{p_2(\varrho)-n} \left(\int_{\partial B_r^+} \left| \frac{\partial \tilde{u}}{\partial r} \right|^{p_2(\varrho)} d\mathcal{H}^{n-1}(x) \right) d\tau \\ (4.66) \quad &\leq \frac{1+c}{\varrho^{p_2(R)-p_2(\varrho)}} \log(R/\varrho) [\Phi(R) - \Phi(\varrho)], \end{aligned}$$

for $c \equiv c(\mathbf{data}, \|Dg\|_{L^q(B_1^+)}, \kappa)$. Finally, keeping in mind (2.7) and position (3.26) we bound via (4.66):

$$\begin{aligned} &\int_{\partial B_1^+} |u(Rx) - u(\varrho x)|^{p_2(\varrho)} d\mathcal{H}^{n-1}(x) \leq c \int_{\partial B_1^+} |\tilde{u}(Rx) - \tilde{u}(\varrho x)|^{p_2(\varrho)} d\mathcal{H}^{n-1}(x) \\ &\quad + c \int_{\partial B_1^+} |g(Rx) - g(\varrho x)|^{p_2(\varrho)} d\mathcal{H}^{n-1}(x) \\ &\leq c \log(R/\varrho) \left[\varrho^{p_2(\varrho)-p_2(R)} (\Phi(R) - \Phi(\varrho)) \right] + c(R - \varrho)^{\gamma_1 \left(1 - \frac{n}{q}\right)}, \end{aligned}$$

with $c \equiv c(\mathbf{data}, \|Dg\|_{L^q(B_1^+)}, \kappa)$ and the proof is complete. \square

Before going on, let us stress that, as in Section 3, we can reduce problem (1.2) to an equivalent one defined on the half-ball B_1^+ . In fact, in the proof of Theorem 2 we shall consider $u \in W^{1,p(\cdot)}(B_1^+, m)$ solution to

$$(4.67) \quad \mathcal{C}_g^{p(\cdot)}(B_1^+, m) \ni w \mapsto \int_{B_1^+} |Dw|^{p(x)} dx,$$

with boundary datum $g(\cdot)$ as in (2.7) (of course $\bar{\Omega}$ is replaced by \bar{B}_1^+). Now we are ready to prove Theorem 2.

4.1. Proof of Theorem 2. As a consequence of Theorem 1, we know that $u \in C_{loc}^{0,\beta_0}(\bar{B}_1^+ \setminus \Sigma_0(u), \mathcal{M})$, for a closed, negligible set $\Sigma_0 \subset \bar{B}_1^+$. Let us prove that $\Sigma_0 \cap \partial B_1^+ = \emptyset$. By contradiction, assume that $x_0 \in \Gamma_1$ is a singular point for $u \in W^{1,p(\cdot)}(B_1^+, \mathcal{M})$, solution to (4.67). Up to translations, there is no loss of generality in assuming $x_0 = 0$. Now, for $j \in \mathbb{N}$, define the rescaled maps

$$u_j(x) := u(x/j), \quad p_j(x) := p(x/j), \quad k_j(x) := j^{p_j(x)-p(0)}, \quad g_j(x) := g(x/j).$$

Since $u \in W^{1,p(\cdot)}(B_1^+, \mathcal{M})$ solves (4.67), we deduce that each $u_j \in W^{1,p_j(\cdot)}(B_j^+, \mathcal{M})$ solves problem

$$(4.68) \quad \mathcal{C}_{g_j}^{p_j(\cdot)}(B_j^+, \mathcal{M}) \ni w \mapsto \min \int_{B_j^+} k_j(x) |Dw|^{p_j(x)} \, dx,$$

therefore it is easy to see that it also solves

$$(4.69) \quad \mathcal{C}_{g_j}^{p_j(\cdot)}(B_1^+, \mathcal{M}) \ni w \mapsto \min \int_{B_1^+} k_j(x) |Dw|^{p_j(x)} \, dx.$$

Notice that, whenever $x \in \bar{B}_1^+$, a straightforward computation shows that

$$(4.70) \quad \{p_j\} \text{ and } \{k_j\} \text{ are Lipschitz continuous uniformly in } j \in \mathbb{N} \text{ in } \bar{B}_1^+.$$

Again for $x \in \bar{B}_1^+$, recalling Morrey's embedding theorem we see that

$$(4.71) \quad \sup_{x \in \bar{B}_1^+} |p_j(x) - p(0)| \leq 4[p]_{0,1} |x/j| \leq 4[p]_{0,1} (1/j) \rightarrow 0$$

$$(4.72) \quad \sup_{x \in \bar{B}_1^+} |k_j(x) - 1| \leq \max \left\{ \exp \left(\frac{4[p]_{0,1} \log(j)}{j} \right) - 1, 1 - \exp \left(\frac{-4[p]_{0,1} \log(j)}{j} \right) \right\} \rightarrow 0$$

$$(4.73) \quad \sup_{x \in \bar{B}_1^+} |g_j(x) - g(0)| \leq 4[g]_{0,1-\frac{n}{q}} |x/j|^{1-\frac{n}{q}} \leq 4[g]_{0,1-\frac{n}{q}} (1/j)^{1-\frac{n}{q}} \rightarrow 0.$$

Furthermore, recalling (2.7) and (4.73) we see that

$$\int_{B_1^+} |Dg_j|^q \, dx \leq j^{q-n} \|Dg\|_{L^q(B_1^+)}^q \, dx \rightarrow 0,$$

so

$$(4.74) \quad g_j \rightarrow g(0) \text{ in } W^{1,q}(\bar{B}_1^+, \mathcal{M}).$$

Collecting (4.68) and (4.70)-(4.74) we see that the assumptions of Lemma 4.1 are satisfied in B_1^+ , so, in particular $u_j \rightharpoonup u_0$ weakly in $W_{loc}^{1,(1+\sigma)p(0)}(B_1^+, \mathcal{M})$, u_0 is a solution of problem

$$(4.75) \quad \mathcal{C}_{g(0)}^{p(0)}(B_R^+, \mathcal{M}) \ni w \mapsto \min \int_{B_R^+} |Dw|^{p(0)} \, dx$$

for any $R \in (0, 1)$ and, since $x_0 = 0$ is a singular point of all the u_j 's, then it is also a singular point for u_0 . We fix $0 < \mu_1 < \mu_2 < 1$ and let $j \in \mathbb{N}$ be so large that $j^{-1} < \frac{T}{4}$ with T as in Lemma 4.2. Recalling also (1.3) (on \bar{B}_1^+ of course), we see that the assumptions of Lemma 4.2 are satisfied, we can apply (4.44) with $\varrho = \mu_1/j$ and $R = \mu_2/j$ to get

$$(4.76) \quad \begin{aligned} & \int_{\partial B_1^+} |u_j(\mu_1 x) - u_j(\mu_2 x)|^{p_2(\mu_1/j)} \, d\mathcal{H}^{n-1}(x) \\ &= \int_{\partial B_1^+} |u(j^{-1}\mu_1 x) - u(j^{-1}\mu_2 x)|^{p_2(\mu_1/j)} \, d\mathcal{H}^{n-1}(x) \\ &\leq c \log(\mu_2/\mu_1) (\Phi(\mu_2/j) - \Phi(\mu_1/j)) + c j^{-\gamma_1(1-\frac{n}{q})} (\mu_2 - \mu_1)^{\gamma_1(1-\frac{n}{q})}, \end{aligned}$$

with $\Phi(\cdot)$ defined as in (4.43) with $k(\cdot) \equiv 1$. By Lemma 4.2, we deduce that

$$\lim_{j \rightarrow \infty} \Phi(\mu_1/j) = \lim_{j \rightarrow \infty} \Phi(\mu_2/j) = L \text{ for some finite } L \geq 0,$$

thus

$$(4.77) \quad c \log(\mu_2/\mu_1) (\Phi(\mu_2/j) - \Phi(\mu_1/j)) + c j^{-\gamma_1(1-\frac{n}{q})} (\mu_2 - \mu_1)^{\gamma_1(1-\frac{n}{q})} \rightarrow 0.$$

Furthermore, in light of (4.3) we have that $u_j \rightarrow u_0$ almost everywhere in B_1^+ , so, recalling also (4.71) we get

$$(4.78) \quad |u_j(\mu_2 x) - u_j(\mu_1 x)|^{p_2(\mu_1/j)} \rightarrow |u_0(\mu_2 x) - u_0(\mu_1 x)|^{p(0)} \quad \text{for a.e. } x \in B_1^+.$$

Combining (4.78), (2.6)₁ and the dominated convergence theorem, we obtain

$$(4.79) \quad \begin{aligned} & \lim_{j \rightarrow \infty} \int_{\partial B_1^+} |u_j(\mu_2 x) - u_j(\mu_1 x)|^{p_2(\mu_1/j)} d\mathcal{H}^{n-1}(x) \\ &= \int_{\partial B_1^+} |u_0(\mu_2 x) - u_0(\mu_1 x)|^{p(0)} d\mathcal{H}^{n-1}(x). \end{aligned}$$

Inserting (4.79) and (4.77) in (4.76), we end up with

$$\int_{\partial B_1^+} |u_0(\mu_2 x) - u_0(\mu_1 x)|^{p(0)} d\mathcal{H}^{n-1}(x) = 0,$$

which in turn implies that u_0 is homogeneous of degree zero. Recalling that u_0 is a solution of (4.75), by [30, Theorem 5.7] we can conclude that u_0 is constant, so $x_0 = 0$ cannot be a singular point. This means that $\Sigma_0 \subseteq B_1^+$ and the proof is complete.

REFERENCES

- [1] E. Acerbi, G. Mingione, Regularity results for stationary electro-rheological fluids. *Arch. Ration. Mech. Anal.* 164, no. 3, 213-259, (2002).
- [2] P. Baroni, Riesz potential estimates for a general class of quasilinear equations. *Calc. Var. & PDE* 53(3-4), 12, pp. 803-846, (2015).
- [3] L. Beck, Partial regularity for weak solutions of nonlinear elliptic systems: The subquadratic case. *Manuscripta Math.* 123, (4), 453-491, (2007).
- [4] L. Beck, G. Mingione, Lipschitz bounds and non-uniform ellipticity. *Comm. Pure Appl. Math.*, (2020). <https://doi.org/10.1002/cpa.21880>
- [5] P. Bella, M. Schaffner, Local boundedness and Harnack inequality for solutions of linear non-uniformly elliptic equations. *Comm. Pure Appl. Math.*, to appear.
- [6] P. Bella, M. Schaffner, On the regularity of scalar integral functionals with (p, q) -growth. *Anal. PDE*, to appear.
- [7] I. Chlebicka, A pocket guide to nonlinear differential equations in Musielak–Orlicz spaces, *Nonl. Analysis*, 175, 1-27, (2018).
- [8] A. Coscia, G. Mingione, Hölder continuity of the gradient of $p(x)$ -harmonic mappings. *C. R. Acad. Sci. Paris Sér. I Math* 328 (4), 363-368, (1999).
- [9] C. De Filippis, Partial regularity for manifold constrained $p(x)$ -harmonic maps. *Calc. Var. & PDE* 58:47, (2019).
- [10] C. De Filippis, G. Mingione, Manifold constrained non-uniformly elliptic problems. *J. Geom. Analysis*, 1-63, (2019). <https://doi.org/10.1007/s12220-019-00275-3>
- [11] C. De Filippis, G. Mingione, On the Regularity of Minima of Non-autonomous Functionals. *J. Geom. Analysis*, 1-43, (2019). <https://doi.org/10.1007/s12220-019-00225-z>
- [12] L. Diening, P. Hästö, M. Ružička, Lebesgue and Sobolev spaces with variable exponents. *Lecture Notes in Mathematics* 2017, Springer-Verlag, Berlin, (2011).
- [13] F. Duzaar, J. F. Grotowski, M. Kronz, Partial and Full Boundary Regularity for Minimizers of Functionals with Nonquadratic Growth. *J. Convex Analysis* 11, 2, 437-476, (2004).
- [14] F. Duzaar, J. Kristensen, G. Mingione, The existence of regular boundary points for non-linear elliptic systems. *J. Reine Angew. Math.* 602, 17-58, (2007).
- [15] F. Duzaar, G. Mingione, The p -harmonic approximation and the regularity of p -harmonic maps. *Calc. Var. Partial Differential Equations* 20, no. 3, 235-256, (2004).
- [16] J. Eells, J. H. Sampson, Harmonic mappings of Riemannian manifolds. *Amer. J. Math.* 86, 109-160, (1964).
- [17] L. Esposito, F. Leonetti, G. Mingione, Sharp regularity for functionals with (p, q) growth. *J. Differential Equations* 204, 5-55, (2004).
- [18] L. C. Evans, R. F. Gariepy, Partial regularity for constrained minimizers of convex or quasiconvex functionals. *Rend. Sem. Mat. Univ. Pol. Torino*, vol. 47, (1989).
- [19] I. Fonseca, J. Malý, G. Mingione, Scalar minimizers with fractal singular sets. *Arch. Ration. Mech. Anal.* 172, 295-307, (2004).

- [20] M. Fuchs, p -Harmonic Obstacle Problems, I. *Annali di Matematica* 156 (4), 127-158, (1990).
- [21] M. Fuchs, p -Harmonic Obstacle Problems, II. *Manuscripta Math.* 63, 381-419, (1989).
- [22] M. Fuchs, p -Harmonic Obstacle Problems, III. *Annali di Matematica* 156 (4), 159-180, (1990).
- [23] M. Giaquinta, L. Martinazzi, An Introduction to the Regularity Theory for Elliptic Systems, Harmonic Maps and Minimal Graphs. *Edizioni della Normale*, (2012).
- [24] M. Giaquinta, G. Modica, J. Souček, Cartesian Currents in the Calculus of Variations II. [*Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics*] vol. 38, Springer-Verlag, Berlin, 1998, Variational integrals. MR 1645082 (2000b:49001b)
- [25] E. Giusti, Direct Methods in the calculus of variations. *World Scientific Publishing Co., Inc., River Edge*, (2003).
- [26] J. F. Grotowski, Boundary regularity for nonlinear elliptic systems. *Calc. Var. & PDE* 15, 353-388, (2002).
- [27] C. Hamburger, Partial boundary regularity of solutions of nonlinear superelliptic systems. *Bollettino UMI* 1, Ser. 8, 63-81, (2007).
- [28] C. Hamburger, Regularity of differential forms minimizing degenerate elliptic functionals. *J. Reine Angew. Math., (Crelles J.)*, 431, 7-64, (1992).
- [29] R. Hardt, D. Kinderlehrer, F.-H. Lin, Stable defects of minimizers of constrained variational principles. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 5, 297-322, (1988).
- [30] R. Hardt, F.-H. Lin, Mappings Minimizing the L^p Norm of the Gradient. *Comm. Pure Appl. Math.* 40, 555-588, (1987).
- [31] P. Harjulehto, P. Hästö, U. V. Le, M. Nuortio, Overview of differential equations with non-standard growth, *Nonlinear Anal.* 72 (12), 4551-4574, (2010).
- [32] P. Harjulehto, P. Hästö, Double Phase image restoration. *Preprint*, (2019). <https://arxiv.org/pdf/1906.09837.pdf>
- [33] P. Hästö, J. Ok, Maximal regularity for local minimizers of non-autonomous functionals. *Preprint*, (2019). <https://arxiv.org/pdf/1902.00261.pdf>
- [34] A. Hatcher, Algebraic Topology. *Cambridge University Press*, Cambridge, (2002).
- [35] J. Hirsch, M. Schäffner, Growth conditions and regularity, an optimal local boundedness result. *Preprint*, (2019). <https://arxiv.org/pdf/1911.12822.pdf>
- [36] C. Hopper, Partial Regularity for Holonomic Minimizers of Quasiconvex Functionals. *Arch. Rational Mech. Anal.* 222, 91-141, (2016).
- [37] J. Jost, M. Meier, Boundary regularity for Minima of Certain Quadratic Functionals. *Math. Ann.* 262, 549-561, (1983).
- [38] J. Kristensen, G. Mingione, Boundary Regularity in Variational Problems. *Arch. Rational Mech. Anal.* 198, 369-455, (2010).
- [39] T. Kuusi, G. Mingione, Vectorial nonlinear potential theory. *J. Europ. Math. Soc. (JEMS)* 20, 929-1004, (2018).
- [40] F. Leonetti, F. Siepe, Maximum Principle for Vector Valued Minimizers. *J. Convex Anal.* 12, 267-278, (2005).
- [41] S. Luckhaus, Partial Hölder continuity for minima of certain energies among maps into a Riemannian manifold. *Indiana Univ. Math. J.* 37, 349-367, (1988).
- [42] J. J. Manfredi, Regularity of the gradient for a class of nonlinear possibly degenerate elliptic equations. *Ph.D. Thesis, University of Washington, St. Louis*, (1986).
- [43] P. Marcellini, Regularity of minimizers of integrals of the calculus of variations with non standard growth conditions. *Arch. Rat. Mech. Anal.* 105, 267-284, (1989).
- [44] P. Marcellini, On the definition and the lower semicontinuity of certain quasiconvex integrals. *Annales de l'I.H.P. Analyse non linéaire* 3, nr. 5, 391-409, (1986).
- [45] M. Mazowiecka, M. Miśkiewicz, A. Schikorra, On the size of the singular set of minimizing harmonic maps into the 2-sphere in dimension four and higher. *Preprint*, (2019), <https://arxiv.org/pdf/1902.03161.pdf>
- [46] A. Naber, D. Valtorta, Rectifiable-Reifenberg and the regularity of stationary and minimizing harmonic maps. *Ann. of Math.* (2) 185, no. 1, 131-227, (2017).
- [47] M. A. Ragusa, A. Tachikawa, Boundary regularity of minimizers of $p(x)$ -energy functionals. *Ann. I. H. Poincaré - AN* 33, 451-476, (2016).
- [48] M. A. Ragusa, A. Tachikawa, H. Takabayashi, Partial regularity of $p(x)$ -harmonic maps. *Transaction of the AMS* 365, nr. 6, 3329-3353, (2013).
- [49] R. Schoen, K. Uhlenbeck, A Regularity Theory for Harmonic Maps. *J. Differential Geometry* 17, 307-335, (1982).
- [50] R. Schoen, K. Uhlenbeck, Boundary regularity and the Dirichlet problem for harmonic maps. *J. Differential Geometry* 18, 253-268, (1983).
- [51] L. Simon, Theorems on Regularity and Singularity of Energy Minimizing Maps. *Lectures in Mathematics*, ETH Zürich, Birkhäuser, Basel, (1996).
- [52] A. Tachikawa, On the singular set of $p(x)$ -energy. *Calc. Var. & PDE* 50, 145-169, (2014).
- [53] K. Uhlenbeck, Regularity for a class of non-linear elliptic systems. *Acta Math.* 138, 219-240, (1977).
- [54] V. V. Zhikov, On some variational problems. *Russian J. Math. Phys.* 5, 105-116, (1997).

- [55] V. V. Zhikov, Averaging of functionals of the calculus of variations and elasticity theory, *Izv. Akad. Nauk SSSR Ser. Mat.* 50, 675-710, (1986).
- [56] V. V. Zhikov, On Lavrentiev's Phenomenon, *Russian J. Math. Phys.* 3, 249-269, (1995).
- [57] V. V. Zhikov, S. M. Kozlov, O. A. Oleinik: Homogenization of differential operators and integral functionals. *Springer-Verlag, Berlin*, (1994).

IWONA CHLEBICKA, FACULTY OF MATHEMATICS, INFORMATICS AND MECHANICS, UNIVERSITY OF WARSAW,
UL. BANACHA 2, 02-097 WARSAW, POLAND
E-mail address: `iskrzypczak@mimuw.edu.pl`

CRISTIANA DE FILIPPIS, MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, ANDREW WILES BUILDING,
RADCLIFFE OBSERVATORY QUARTER, WOODSTOCK ROAD, OXFORD, OX26GG, OXFORD, UNITED KINGDOM
E-mail address: `Cristiana.DeFilippis@maths.ox.ac.uk`

LUKAS KOCH, MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, ANDREW WILES BUILDING, RADCLIFFE
OBSERVATORY QUARTER, WOODSTOCK ROAD, OXFORD, OX26GG, OXFORD, UNITED KINGDOM
E-mail address: `Lukas.Koch@maths.ox.ac.uk`