

# Rationality of inner twisted flags of type $A_n$

Saša Novaković

January 2020

**Abstract.** We prove that  $\mathrm{rcodim}(X) \geq 2$  if  $X$  is a rational inner twisted flag of type  $A_n$ .

## 1. INTRODUCTION

A potential measure for rationality was introduced by Bernardara and Bolognesi [6] with the notion of categorical representability. We use the definition given in [1]. A  $k$ -linear triangulated category  $\mathcal{T}$  is said to be *representable in dimension  $m$*  if there is a semiorthogonal decomposition (see Section 3 for the definition)  $\mathcal{T} = \langle \mathcal{A}_1, \dots, \mathcal{A}_n \rangle$  and for each  $i = 1, \dots, n$  there exists a smooth projective connected variety  $Y_i$  with  $\dim(Y_i) \leq m$ , such that  $\mathcal{A}_i$  is equivalent to an admissible subcategory of  $D^b(Y_i)$ . We set

$$\mathrm{rdim}(\mathcal{T}) := \min\{m \mid \mathcal{T} \text{ is representable in dimension } m\},$$

whenever such a finite  $m$  exists. Let  $X$  be a smooth projective  $k$ -variety. One says  $X$  is *representable in dimension  $m$*  if  $D^b(X)$  is representable in dimension  $m$ . We will use the following notations:

$$\mathrm{rdim}(X) := \mathrm{rdim}(D^b(X)), \quad \mathrm{rcodim}(X) := \dim(X) - \mathrm{rdim}(X).$$

Note that when the base field  $k$  of a variety  $X$  is not algebraically closed, the existence of  $k$ -rational points on  $X$  and whether  $X$  is  $k$ -rational is a major open question in arithmetic geometry. We recall the following question which was formulated in [7]:

**Question.** *Let  $X$  be a smooth projective variety over  $k$  of dimension at least 2. Suppose  $X$  is  $k$ -rational. Do we have  $\mathrm{rcodim}(X) \geq 2$  ?*

There are several results suggesting that this question has a positive answer (see [1], [2], [7], [15], [16], [17] and references therein). In this context, we prove:

**Theorem 1.1.** *Let  $A$  be a central simple  $k$ -algebra of degree  $n$  over a field of characteristic zero. We fix a sequence of integers  $0 < n_1 < n_2 < \dots < n_r < n$  and let  $X = \mathrm{BS}(n_1, \dots, n_r, A)$  be a twisted flag of dimension at least two. If  $X$  is rational over  $k$ , then  $\mathrm{rcodim}(X) \geq 2$ .*

**Acknowledgement.** I would like to thank Nikita Semenov for answering questions concerning twisted flag varieties. This research was conducted in the framework of the research training group GRK 2240: Algebro-geometric Methods in Algebra, Arithmetic and Topology, which is funded by the DFG.

## 2. INNER TWISTED FLAGS OF TYPE $A_n$

Recall that a finite-dimensional  $k$ -algebra  $A$  is called *central simple* if it is an associative  $k$ -algebra that has no two-sided ideals other than 0 and  $A$  and if its center equals  $k$ . If the algebra  $A$  is a division algebra it is called *central division algebra*. Note that  $A$  is a central simple  $k$ -algebra if and only if there is a finite field extension  $k \subset L$ , such that  $A \otimes_k L \simeq M_n(L)$ . This is also equivalent to  $A \otimes_k \bar{k} \simeq M_n(\bar{k})$ . An extension  $k \subset L$  such that  $A \otimes_k L \simeq M_n(L)$  is called *splitting field* for  $A$ . The *degree* of a central simple algebra  $A$  is defined to be  $\deg(A) := \sqrt{\dim_k A}$ . According to the *Wedderburn Theorem*, for any central simple  $k$ -algebra  $A$  there is a unique integer  $n > 0$  and a division  $k$ -algebra  $D$  such

that  $A \simeq M_n(D)$ . The division algebra  $D$  is also central and unique up to isomorphism. The degree of the unique central division algebra  $D$  is called the *index* of  $A$  and is denoted by  $\text{ind}(A)$ . Two central simple algebras  $A$  and  $B$  are said to be *Brauer-equivalent* if there are positive integers  $r, s$  such that  $M_r(A) \simeq M_s(B)$ .

A Brauer–Severi variety of dimension  $n$  can also be defined as a scheme  $X$  of finite type over  $k$  such that  $X \otimes_k L \simeq \mathbb{P}^n$  for a finite field extension  $k \subset L$ . A field extension  $k \subset L$  for which  $X \otimes_k L \simeq \mathbb{P}^n$  is called *splitting field* of  $X$ . Clearly,  $k^s$  and  $\bar{k}$  are splitting fields for any Brauer–Severi variety. In fact, every Brauer–Severi variety always splits over a finite Galois extension. It follows from descent theory that  $X$  is projective, integral and smooth over  $k$ . For details we refer to [3] and [10].

To a central simple  $k$ -algebra  $A$  one can also associate twisted forms of Grassmannians. Let  $A$  be of degree  $n$  and  $1 \leq d \leq n$ . Consider the subset of  $\text{Grass}_k(d \cdot n, A)$  consisting of those subspaces of  $A$  that are left ideals  $I$  of dimension  $d \cdot n$ . This subset can be given the structure of a projective variety which turns out to be a generalized Brauer–Severi variety. It is denoted by  $\text{BS}(d, A)$ . After base change to some splitting field  $L$  of  $A$  the variety  $\text{BS}(d, A)$  becomes isomorphic to  $\text{Grass}_L(d, n)$ . If  $d = 1$  the generalized Brauer–Severi variety is the Brauer–Severi variety associated to  $A$ . For details see [9].

We also recall the basics of generalized Brauer–Severi schemes (see [14]). Let  $X$  be a noetherian  $k$ -scheme and  $\mathcal{A}$  a sheaf of Azumaya algebras of rank  $n^2$  over  $X$  (see [11], [12] for details on Azumaya algebras). For an integer  $1 \leq n_1 < n$  the generalized Brauer–Severi scheme  $p : \text{BS}(n_1, \mathcal{A}) \rightarrow X$  is defined as the scheme representing the functor  $F : \text{Sch}/X \rightarrow \text{Sets}$ , where  $(\psi : Y \rightarrow X)$  is mapped to the set of left ideals  $\mathcal{J}$  of  $\psi^* \mathcal{A}$  such that  $\psi^* \mathcal{A}/\mathcal{J}$  is locally free of rank  $n(n - n_1)$ . By definition, there is an étale covering  $U \rightarrow X$  and a locally free sheaf  $\mathcal{E}$  of rank  $n$  with the following trivializing diagram:

$$\begin{array}{ccc} \text{Grass}(n_1, \mathcal{E}) & \xrightarrow{\pi} & \text{BS}(n_1, \mathcal{A}) \\ q \downarrow & & \downarrow p \\ U & \xrightarrow{g} & X \end{array}$$

In the same way one defines the twisted relative flag  $\text{BS}(n_1, \dots, n_r, \mathcal{A})$  as the scheme representing the functor  $F : \text{Sch}/X \rightarrow \text{Sets}$ , where  $(\psi : Y \rightarrow X)$  is mapped to the set of left ideals  $\mathcal{J}_1 \subset \dots \subset \mathcal{J}_r$  of  $\psi^* \mathcal{A}$  such that  $\psi^* \mathcal{A}/\mathcal{J}_i$  is locally free of rank  $n(n - n_i)$ . As for the generalized Brauer–Severi schemes, there is an étale covering  $U \rightarrow X$  and a locally free sheaf  $\mathcal{E}$  of rank  $n$  with diagram

$$\begin{array}{ccc} \text{Flag}_U(n_1, \dots, n_r, \mathcal{E}) & \xrightarrow{\pi} & \text{BS}(n_1, \dots, n_r, \mathcal{A}) \\ q \downarrow & & \downarrow p \\ U & \xrightarrow{g} & X \end{array}$$

Note that the usual Brauer–Severi schemes are obtained from the generalized one by setting  $n_1 = 1$ . In this case one has a well known one-to-one correspondence between sheaves of Azumaya algebras of rank  $n^2$  on  $X$  and Brauer–Severi schemes of relative dimension  $n - 1$  via  $\tilde{H}^1(X_{\text{ét}}, \text{PGL}_n)$  (see [11]). Note that if the base scheme  $X$  is a point a sheaf of Azumaya algebras on  $X$  is a central simple  $k$ -algebra and the generalized Brauer–Severi schemes are the generalized Brauer–Severi varieties from above. Consider a twisted flag  $X = \text{SB}(n_1, \dots, n_r, A) \rightarrow \text{Spec}(k)$ . Such an  $X$  is an *inner form* of a partial flag variety. That is, there is a cartesian square of the form

$$\begin{array}{ccc} \text{Grass}_L(n_1, \dots, n_r, V) & \xrightarrow{\pi} & \text{BS}(n_1, \dots, n_r, A) \\ q \downarrow & & \downarrow p \\ \text{Spec}(L) & \xrightarrow{\pi} & \text{Spec}(k) \end{array}$$

where  $L/k$  is a Galois extension and the 1-cocycle  $\text{Gal}(L/k) \rightarrow \text{Aut}(\text{Grass}_L(n_1, \dots, n_r, V))$  factors through  $\text{PGL}(V)$ .

### 3. SEMIORTHOGONAL DECOMPOSITIONS

Let  $\mathcal{D}$  be a triangulated category and  $\mathcal{C}$  a triangulated subcategory. The subcategory  $\mathcal{C}$  is called *thick* if it is closed under isomorphisms and direct summands. For a subset  $A$  of objects of  $\mathcal{D}$  we denote by  $\langle A \rangle$  the smallest full thick subcategory of  $\mathcal{D}$  containing the elements of  $A$ . For a smooth projective variety  $X$  over  $k$ , we denote by  $D^b(X)$  the bounded derived category of coherent sheaves on  $X$ . Moreover, if  $B$  is an associated  $k$ -algebra, we write  $D^b(B)$  for the bounded derived category of finitely generated left  $B$ -modules.

**Definition 3.1.** Let  $A$  be a division algebra over  $k$ , not necessarily central. An object  $\mathcal{E}^\bullet \in D^b(X)$  is called *A-exceptional* if  $\text{End}(\mathcal{E}^\bullet) = A$  and  $\text{Hom}(\mathcal{E}^\bullet, \mathcal{E}^\bullet[r]) = 0$  for  $r \neq 0$ . By *generalized exceptional object*, we mean  $A$ -exceptional for some division algebra  $A$  over  $k$ . If  $A = k$ , the object  $\mathcal{E}^\bullet$  is called *exceptional*.

**Definition 3.2.** A totally ordered set  $\{\mathcal{E}_1^\bullet, \dots, \mathcal{E}_n^\bullet\}$  of generalized exceptional objects on  $X$  is called a *generalized exceptional collection* if  $\text{Hom}(\mathcal{E}_i^\bullet, \mathcal{E}_j^\bullet[r]) = 0$  for all integers  $r$  whenever  $i > j$ . A generalized exceptional collection is *full* if  $\langle \{\mathcal{E}_1^\bullet, \dots, \mathcal{E}_n^\bullet\} \rangle = D^b(X)$  and *strong* if  $\text{Hom}(\mathcal{E}_i^\bullet, \mathcal{E}_j^\bullet[r]) = 0$  whenever  $r \neq 0$ . If the set  $\{\mathcal{E}_1^\bullet, \dots, \mathcal{E}_n^\bullet\}$  consists of exceptional objects it is called *exceptional collection*.

Recall that a full thick triangulated subcategory  $\mathcal{D}$  of  $D^b(X)$  is called *admissible* if the inclusion  $\mathcal{D} \hookrightarrow D^b(X)$  has a left and right adjoint functor.

**Definition 3.3.** Let  $X$  be a smooth projective variety over  $k$ . A sequence  $\mathcal{D}_1, \dots, \mathcal{D}_n$  of full admissible triangulated subcategories of  $D^b(X)$  is called *semiorthogonal* if  $\mathcal{D}_j \subset \mathcal{D}_i^\perp = \{\mathcal{F}^\bullet \in D^b(X) \mid \text{Hom}(\mathcal{G}^\bullet, \mathcal{F}^\bullet) = 0, \forall \mathcal{G}^\bullet \in \mathcal{D}_i\}$  for  $i > j$ . Such a sequence defines a *semiorthogonal decomposition* of  $D^b(X)$  if the smallest thick full subcategory containing all  $\mathcal{D}_i$  equals  $D^b(X)$ .

For a semiorthogonal decomposition we write  $D^b(X) = \langle \mathcal{D}_1, \dots, \mathcal{D}_n \rangle$ .

**Remark 3.4.** Let  $\mathcal{E}_1^\bullet, \dots, \mathcal{E}_n^\bullet$  be a full generalized exceptional collection on  $X$ . It is easy to verify that by setting  $\mathcal{D}_i = \langle \mathcal{E}_i^\bullet \rangle$  one gets a semiorthogonal decomposition  $D^b(X) = \langle \mathcal{D}_1, \dots, \mathcal{D}_n \rangle$ .

### 4. PROOF OF THEOREM 1.1

**Lemma 4.1.** *Let  $X$  be the inner twisted flag from Theorem 1.1. Then  $D^b(X)$  admits a semiorthogonal decomposition with components being equivalent to  $D^b(A^{\otimes j})$  for suitable positive integers  $j \geq 0$ .*

*Proof.* On a relative flag  $\text{Grass}(n_1, \dots, n_r, \mathcal{E})$  one has tautological subbundles sitting in the following sequence

$$0 \longrightarrow \mathcal{R}_1 \longrightarrow \dots \longrightarrow \mathcal{R}_r \longrightarrow q^* \mathcal{E} \longrightarrow \mathcal{T}_1 \longrightarrow \dots \longrightarrow \mathcal{T}_r \longrightarrow 0.$$

In our case  $\mathcal{E}$  is a  $k$ -vector space of dimension  $n$  denoted by  $V$ . Recall that  $X = \text{BS}(n_1, \dots, n_r, A)$  is the inner twisted flag obtained from  $\text{Grass}(n_1, \dots, n_r, V)$  by descent. Denote by  $\alpha(1), \dots, \alpha(r)$  partitions of the forms

$$(\alpha_1, \dots, \alpha_{l_1}), (\alpha_1, \dots, \alpha_{l_2}), \dots, (\alpha_1, \dots, \alpha_{l_r}),$$

with  $0 \leq \alpha_i \leq l_2 - l_1, \dots, 0 \leq \alpha_i \leq l_m - l_{m-1}, 0 \leq \alpha_i \leq n - l_m$ . Let  $|\alpha| = \sum_{i=1}^r |\alpha(i)|$ , where  $|\alpha(j)| = \sum_{s=1}^{l_j} \alpha_s$  for  $1 \leq j \leq r$ . Then the sheaf  $(S^{\alpha(1)} \mathcal{R}_1 \otimes \dots \otimes S^{\alpha(r)} \mathcal{R}_r)^{\oplus n \cdot |\alpha|}$  descends to a sheaf  $\mathcal{J}_\alpha$  on  $X$  (see [14], Section 4). As in the case of generalized Brauer–Severi varieties, one can show that  $\text{End}(\mathcal{J}_\alpha)$  is Brauer-equivalent to  $A^{\otimes |\alpha|}$ . Note that  $D^b(\text{End}(\mathcal{J}_\alpha))$  is equivalent to  $D^b(A^{\otimes |\alpha|})$  by Morita equivalence. Now it is proved in [4] that the subcategories  $\langle \mathcal{J}_\alpha \rangle \simeq D^b(\text{End}(\mathcal{J}_\alpha))$  form a semiorthogonal decomposition.  $\square$

**Proposition 4.2.** *Let  $X$  be the inner twisted flag from above. Then  $\mathrm{rdim}(X) \leq \mathrm{ind}(A) - 1$ .*

*Proof.* According to Lemma 4.1,  $D^b(X)$  admits a semiorthogonal decomposition with components being equivalent to  $D^b(A^{\otimes j})$  for suitable positive integers  $j \geq 0$ . Now let  $A = M_s(D)$  according to the Wedderburn theorem. Since any of the derived categories  $D^b(A^{\otimes j})$  can be embedded into  $D^b(Y)$ , where  $Y$  is the Brauer–Severi variety corresponding to  $D$  (see [5]), the assertion follows from the fact that  $\deg(D) = \mathrm{ind}(A)$  and  $\dim(Y) = \mathrm{ind}(A) - 1$ .  $\square$

**Lemma 4.3.** *Let  $\alpha \geq 1$  be an integer. Then  $\alpha x^2 - x - 1 \geq 0$  for all  $x \geq 2$ .*

*Proof.* The function  $\alpha x^2 - x - 1$  is strictly monotonically increasing for  $x > 1/2\alpha$  and has its positive root at  $(1/2\alpha) \cdot (1 + \sqrt{1 + 4\alpha})$  which is  $< 2$  for all  $\alpha \geq 1$ .  $\square$

*Proof.* (of Theorem 1.1)

Recall the well known fact that  $X(k) \neq \emptyset$  iff  $\mathrm{ind}(A) \mid d$ , where  $d = \gcd(n, n_1, \dots, n_r)$  denotes the greatest common divisor. So if  $X$  is  $k$ -rational, it has a  $k$ -rational point and therefore  $\mathrm{ind}(A) \mid d$ . Now let  $n = b \cdot \mathrm{ind}(A)$  and  $n_i = b_i \cdot \mathrm{ind}(A)$  for  $i = 1, \dots, r$ . Then  $b, b_1, \dots, b_r$  are positive integers satisfying  $0 < b_1 < b_2 < \dots < b_r < b$ . This implies that  $a \geq 1$  for

$$a = b_1(b - b_1) + \sum_{i=2}^r (b_i - b_{i-1})(b - b_i).$$

Assume  $\mathrm{ind}(A) > 1$  and let  $x = \mathrm{ind}(A)$ . With  $\alpha = a$ , Lemma 4.3 implies

$$a \cdot \mathrm{ind}(A)^2 - \mathrm{ind}(A) - 1 \geq 0.$$

It is easy to see that

$$a \cdot \mathrm{ind}(A)^2 = n_1(n - n_1) + \sum_{i=2}^r (n_i - n_{i-1})(n - n_i).$$

This implies

$$n_1(n - n_1) + \sum_{i=2}^r (n_i - n_{i-1})(n - n_i) \geq (\mathrm{ind}(A) - 1) + 2.$$

Now, since

$$\dim(X) = n_1(n - n_1) + \sum_{i=2}^r (n_i - n_{i-1})(n - n_i)$$

and since  $\mathrm{rdim}(X) \leq \mathrm{ind}(A) - 1$  according to Proposition 4.2, we conclude

$$\mathrm{rcodim}(X) = \dim(X) - \mathrm{rdim}(X) \geq 2.$$

It remains to consider the case  $\mathrm{ind}(A) = 1$ . In this case the inner twisted flag  $X$  is split, i.e. is isomorphic to  $\mathrm{Flag}_k(n_1, \dots, n_r, V)$ , where  $V$  is a  $n$ -dimensional  $k$ -vector space. From Kapranov [13], we know that  $X$  admits a full exceptional collection. But then  $\mathrm{rdim}(X) = 0$  according to [1], Proposition 6.1.6 and hence  $\mathrm{rcodim}(X) \geq 2$ .  $\square$

**Remark 4.4.** If the inner twisted flag  $X$  from Theorem 1.1 is of dimension one, it is actually a Brauer–Severi curve. In this case [15] shows that rationality of  $X$  is equivalent to  $\mathrm{rdim}(X) = 0$ . Hence  $\mathrm{rcodim}(X) = 1$ . This is one reason why the question in the introduction assumes  $\dim(X) \geq 2$ .

## REFERENCES

- [1] A. Auel and M. Bernardara: Cycles, derived categories, and rationality, in *Surveys on Recent Developments in Algebraic Geometry*, Proceedings of Symposia in Pure Mathematics 95 (2017), 199-266.
- [2] A. Auel and M. Bernardara: Semiorthogonal decompositions and birational geometry of del Pezzo surfaces over arbitrary fields. *Proc. London Math. Soc.* 117 (2018) 1-64.
- [3] M. Artin: Brauer-Severi varieties. Brauer groups in ring theory and algebraic geometry, *Lecture Notes in Math.* 917, Notes by A. Verschoren, Berlin, New York: Springer-Verlag (1982), 194-210
- [4] S. Baek: Semiorthogonal decompositions for twisted Grassmannians. *arXiv:1205.1175v1 [math.AG]* (2012).
- [5] M. Bernardara: A semiorthogonal decomposition for Brauer–Severi schemes. *Math. Nachr.* 282 (2009), 1406-1413.
- [6] M. Bernardara and M. Bolognesi: Categorical representability and intermediate Jacobians of Fano threefolds. *EMS Ser. Congr. Rep.*, Eur. Math. Soc. (2013), 1-10.
- [7] M. Bernardara: Semiorthogonal decompositions and noncommutative motives in algebraic geometry. Habilitation thesis, available at <https://www.math.univ-toulouse.fr/~mbernard/> (2016).
- [8] M. Bernardara: Categorical dimension of birational automorphisms and filtrations of Cremona maps, *arXiv:1701.06803v3* (2017).
- [9] A. Blanchet: Function fields of generalized Brauer–Severi varieties. *Comm. Algebra*. Vol. 19 (1991), 97-118.
- [10] P. Gille and T. Szamuely: *Central simple algebras and Galois cohomology*. Cambridge Studies in advanced Mathematics. 101. Cambridge University Press. (2006)
- [11] A. Grothendieck: Le group de Brauer I: Algebras d Azumaya et interpretations diverses, *Seminaire Bourbaki*. No. 290 (1964).
- [12] A. Grothendieck: Le group de Brauer II: Theorie cohomologique, *Seminaire Bourbaki*. No. 297 (1965).
- [13] M.M. Kapranov: On the derived categories of coherent sheaves on some homogeneous spaces. *Invent. Math.* Vol. 92 (1988), 479-508.
- [14] M. Levine, V. Srinivas and J. Weyman: K-Theory of Twisted Grassmannians. *K-Theory* Vol. 3 (1989), 99-121.
- [15] S. Novaković: Non-existence of exceptional collections on twisted flags and categorical representability via noncommutative motives. *arXiv:1607.01043v1 [math.AG]* (2016).
- [16] S. Novaković: On non-existence of full exceptional collections on some relative flags. *arXiv:1607.04834v1 [math.AG]* (2016).
- [17] S. Novaković: Rational points on symmetric powers and categorical representability. *arXiv:1704.02474v1 [math.AG]* (2017).

MATHEMATISCHES INSTITUT, HEINRICH-HEINE-UNIVERSITÄT 40225 DÜSSELDORF,  
GERMANY

E-mail adress: novakovic@math.uni-duesseldorf.de