

# ANALYTIC CLASSIFICATION OF GERMS OF PARABOLIC ANTIHOLOMORPHIC DIFFEOMORPHISMS OF CODIMENSION $k$

JONATHAN GODIN AND CHRISTIANE ROUSSEAU

**ABSTRACT.** We investigate the local dynamics of antiholomorphic diffeomorphisms around a parabolic fixed point. We first give a normal form. Then we give a complete classification including a modulus space for antiholomorphic germs with a parabolic fixed point under analytic conjugacy. We then study some geometric applications: existence of real analytic invariant curve, existence of holomorphic and antiholomorphic roots of holomorphic and antiholomorphic parabolic germs, commuting holomorphic and antiholomorphic parabolic germs.

## CONTENTS

|  |    |
|--|----|
| 1. Introduction  | 1  |
| 2. Antiholomorphic Parabolic Fixed Points                      | 5  |
| 3. Properties of the Formal Normal Form                        | 10 |
| 4. Fatou Coordinates   | 11 |
| 5. Modulus of Analytic Classification                          | 17 |
| 6. Space of Orbits and Classification Under Analytic Conjugacy | 24 |
| 7. Applications of the Classification Theorem                  | 29 |
| References   | 41 |

## 1. INTRODUCTION

In this paper, we are interested in the local dynamics of antiholomorphic diffeomorphisms with a parabolic fixed point, i.e. a fixed point of multiplicity  $k + 1$  (i.e. of codimension  $k$ ). We study the classification

---

*Date:* February 18, 2022.

*2010 Mathematics Subject Classification.* 37F45, 32H50.

*Key words and phrases.* Antiholomorphic Dynamics, Antiholomorphic Parabolic Fixed Point, Local Classification under Analytic Conjugation, Space of Orbits near a Fixed Point.

The first author was supported by a FRQ-NT PhD scholarship. The second author is supported by NSERC in Canada.

under conjugacy by analytic changes of coordinate of a germ of an antiholomorphic diffeomorphism  $f$  with a parabolic fixed point. In local coordinate, it may be chosen in the form

$$(1) \quad f(z) = \bar{z} + \frac{1}{2}\bar{z}^{k+1} + o(\bar{z}^{k+1})$$

for some integer  $k \geq 1$ .

The classification of parabolic fixed points in the holomorphic case for a germ

$$(2) \quad g(z) = z + z^{k+1} + \left(\frac{k+1}{2} - b\right) z^{2k+1} + o(z^{2k+1})$$

is well known. (See e.g. [5] or [6].) The dynamics of  $g$  (see Figure 1) is determined by a topological invariant, the integer  $k$ , a formal invariant, the complex number  $b$ , and an analytic invariant given by an equivalence class of  $2k$  germs of diffeomorphisms which are the transition functions on the space of orbits of  $g$  (the Écalle horn maps). Two germs  $g_1$  and  $g_2$  are formally equivalent if and only if they have the same topological invariant and formal invariant. Furthermore, they are analytically equivalent if and only if they also have the same analytic invariant.




FIGURE 1. Dynamics of a holomorphic parabolic germ with topological invariant  $k = 3$

The goal of this paper is to establish a local classification of antiholomorphic parabolic germs under the analytic conjugation and to describe the space of orbits of such a germ and, more generally, to explore the geometric properties of antiholomorphic parabolic germs which are invariant under analytic conjugation. This is done for fixed points of any multiplicity. It allows us to provide a solution to the following problems.

*Questions.* 1. (*Antiholomorphic Root Extraction*) The second iterate of an antiholomorphic parabolic germ  $f$  as in equation (1)

is a holomorphic germ which is parabolic. When is the converse true: Given a parabolic germ of holomorphic diffeomorphism  $g$ , when is it possible to write it as  $g = f \circ f$ , for some antiholomorphic parabolic germ  $f$ ? We call  $f$  an *antiholomorphic square root* of  $g$ . More generally, when does  $g$  have an antiholomorphic root of some order?

2. Analogously, when does an antiholomorphic parabolic germ have an antiholomorphic root? When are the roots unique?
3. (*Embedding*) Let  $\{v^t\}_t$ , where  $v^t: z \mapsto v^t(z)$ , be the flow of the differential equation  $\dot{z} = v(z) = \frac{z^{k+1}}{1+bz^k}$ . Each  $v^t$  ( $t \neq 0$ ) is a holomorphic germ with a parabolic fixed point at the origin. Then  $\overline{v^{\frac{1}{2}}(\cdot)}$  is an antiholomorphic germ, and any antiholomorphic parabolic germ is formally conjugate to such a germ. Given an antiholomorphic parabolic germ  $f$ , when is it analytically conjugate to some  $\overline{v^{\frac{1}{2}}(\cdot)}$ ? In that case, it allows to embed  $f$  in the family  $\overline{v^t}$ .
4. When does an antiholomorphic germ preserve a germ of real analytic curve? This is equivalent to say that the germ is analytically conjugate to a germ with real coefficients.
5. (*Centralizer*) Can we describe all the antiholomorphic parabolic germs  $f$  that commute with a holomorphic parabolic germ  $g$ ? If  $f$  and  $g$  commute, then  $f$  sends the orbits of  $g$  on the orbits of  $g$ . This greatly restricts the possible  $f$ . In an analogous way, can we describe all the holomorphic and the antiholomorphic germs that commute with an antiholomorphic parabolic germ?

The above problems are questions about the equivalence classes of germs under analytic conjugacy. Therefore, the answer should be read in the modulus of classification, which will be introduced in Section 5.

The local dynamics of an antiholomorphic parabolic germ has similarities with the holomorphic case: indeed the  $n$ -th iterate  $f^{\circ n}$  is holomorphic for  $n$  even. We find that the dynamics is determined by the same topological and formal invariants, but the analytic invariant is composed of  $k$  germs of diffeomorphisms, instead of  $2k$ . This is explained by the fact that an orbit of  $f$  will usually jump between two Fatou petals of its associated holomorphic parabolic germ  $f^{\circ 2}$  (see Figure 2), so that the dynamics in those petals are not independent.

We observe other differences from the holomorphic case. A holomorphic germ has  $2k$  formal separatrices. The antiholomorphic germ has instead a privileged unique direction; a *formal symmetry axis*. There




FIGURE 2. An orbit of  $f$  jumping between two petals. An orbit of the second iterate  $f^{\circ 2}$  will remain either in the upper petal or the lower petal.

is also a topological difference between the cases where the codimension is odd or even. When  $k$  is even, the rotation  $z \mapsto -z$  is a formal symmetry of  $f$ , whilst it is not for  $k$  odd.

Antiholomorphic dynamics has been considered before in the context of anti-polynomials, that is a polynomial function of  $\bar{z}$ ,  $p(z) = \bar{z}^n + \dots + a_0$ . Iteration of anti-polynomials was studied by Nakane and Schleicher in [8], Hubbard and Schleicher in [4], and Mukherjee, Nakane and Schleicher in [7]. Their focus is mostly on the family of anti-polynomials  $p_c(z) = \bar{z}^d + c$  and the description of the connectedness locus  $\mathcal{M}_d^*$  called the *multicorn*. The context is global in nature, but the local analysis contributes significantly.

An important role is played by periodic orbits of  $p_c$  of odd period  $k$ , because when  $k$  is odd,  $p_c^{\circ k}$  is antiholomorphic. In this case, all indifferent periodic orbits are parabolic and they occur along real analytic arcs in the parameter space, as proved in [4] and [7]. However, only points of codimension 1 and 2 are observed. This is due to a choice of a special subfamily of anti-polynomials of degree  $d$ . Indeed, higher codimension is already observed in the 2-parameter family  $\bar{z}^d + c_1 \bar{z} + c_0$ , e.g. when  $c_1 = 1$  and  $c_0 = 0$ .

One of the tools used for anti-polynomials is called the *Écalle height*, introduced by Hubbard and Schleicher in [4]. In codimension 1, on the Écalle cylinder of the attractive petal, the imaginary part of an orbit is intrinsic, and this is used to prove that landing points of curves in the parameter space where  $p_c$  has a parabolic periodic orbit of odd period are points of codimension 2. When studying the space of orbits of an antiholomorphic germ of parabolic diffeomorphism of any codimension, we see that the Écalle height has a meaning only on the Écalle cylinder of the petals containing the formal symmetry axis of  $f$ . This is seen by describing the space of orbits on a neighbourhood of a parabolic fixed point, which we do in Section 6.2.

The paper is organized as follows. In Section 2, we define the topological and formal invariants of  $f$ . We also establish a formal normal form for  $f$ .

In Section 3, we study the formal normal form.

In Section 4, we introduce the rectifying coordinate and the Fatou coordinates in order to define the transition functions (Def. 5.0.4) in Section 5, which is the analytic invariant. This leads to the modulus of classification of  $f$ .

In Section 6.1, we recall a description of the space of orbits in the holomorphic case using  $2k$  spheres (or Écalle cylinders) glued with the horn maps (these are the expressions of the transition functions in the coordinates of the spheres). We use this space of orbits in Section 6.2 to identify the space of orbits of  $f$  to a manifold of real dimension 2 by quotienting the space of orbits of  $f \circ f$  by the action of  $f$ .

After describing the space of orbits, we state, in Section 6.3, the main result of the paper: the Classification Theorem 6.3.1. The idea in spirit is that two germs are equivalent if and only if their space of orbits are equivalent; the Classification Theorem is a way to rigorously express this statement.

Finally, with the Classification Theorem in hands, in Section 7 we answer Questions 1 to 5 of the introduction.

## 2. ANTIHOLOMORPHIC PARABOLIC FIXED POINTS

**Notation.** For the whole paper, we will use the following notation :

- $\sigma(z) = \bar{z}$  is the complex conjugation;
- $\tau(w) = \frac{1}{\bar{w}}$  is the antiholomorphic inversion;
- $T_C(Z) = Z + C$  is the translation by  $C \in \mathbb{C}$ ;
- $L_c(w) = cw$  is the linear transformation with multiplier  $c \in \mathbb{C}$ ;
- $v^t$  is the time- $t$  of the vector field

$$(3) \quad \dot{z} = v(z) = \frac{z^{k+1}}{1 + bz^k}.$$

A function  $f: U \rightarrow \mathbb{C}$  defined on a domain  $U \subseteq \mathbb{C}$  is antiholomorphic if  $\frac{\partial f}{\partial z} \equiv 0$  on  $U$ . From this definition, together with the chain rule, it follows that antiholomorphy is an intrinsic property of  $f$  under holomorphic changes of variable. Equivalently,  $f: z \mapsto f(z)$  is antiholomorphic if  $f \circ \sigma: z \mapsto f(\bar{z})$  is holomorphic, therefore  $f(z)$  expands in a power series in terms of  $\bar{z}$ .

Note that the multiplier at a fixed point of an antiholomorphic function is not intrinsic, only its modulus is. Indeed, a scaling of  $\lambda$  will add a factor of  $\frac{\lambda}{\bar{\lambda}}$  to the multiplier.

**Definition 2.0.1** (Parabolic fixed point). A germ of antiholomorphic diffeomorphism fixing the origin  $f: (\mathbb{C}, 0) \rightarrow (\mathbb{C}, 0)$  has a *parabolic fixed point* at 0 if 0 is an isolated fixed point and

$$\left| \frac{\partial f}{\partial \bar{z}}(0) \right| = 1.$$

We will also say that  $f$  is an *antiholomorphic parabolic germ*.

**Proposition 2.0.2.** *Let  $f(z) = a_1 \bar{z} + a_2 \bar{z}^2 + a_3 \bar{z}^3 + \dots$ . If  $|a_1| = 1$ , then  $f$  is formally conjugate to a formal power series*

$$(4) \quad f^\dagger(w) = \bar{w} + \sum_{n=2}^{\infty} A_n \bar{w}^n$$

with real coefficients  $A_n$ . If there exists  $n \geq 2$  such that  $A_n \neq 0$ , then 0 is a parabolic fixed point of  $f$ . Let  $n_0 = k + 1$  be the minimum such  $n$ . Then a scaling brings  $A_{k+1}$  to  $\frac{1}{2}$  if  $k$  is odd (resp.  $\pm \frac{1}{2}$  if  $k$  is even).

*Proof.* The proof is a mere computation. Let  $w = \hat{h}(z) = \sum_{n \geq 1} b_n z^n$  be a formal change of coordinate and suppose  $f^\dagger(w) = \bar{w} + \sum_{n \geq 2} A_n \bar{w}^n$ . If we compare  $h \circ f(z) = f^\dagger \circ h(z)$  degree by degree, we find an expression for the coefficients of the form

$$\begin{cases} b_1 a_1 = \bar{b}_1 \\ A_n = b_n - \bar{b}_n + a_n + P_n(A_1, \dots, A_{n-1}, a_1, \dots, a_{n-1}, b_1, \dots, b_{n-1}), \end{cases}$$

where  $P_n$  is some polynomial. Hence, we have  $\arg b_1 = -\frac{1}{2} \arg a_1 + \ell\pi$ , with  $\ell \in \mathbb{Z}$ . With a recursive argument, if  $A_1, \dots, A_{n-1}$  are real, for  $A_n$  to be real, we may choose  $\Im b_n = \frac{1}{2} \Im(a_n + P_n)$ , since  $P_n$  depends only on terms that were fixed in the previous steps.  $\square$

*Remark 2.0.3.* The formal change of coordinate  $\hat{h}$  is not unique. Indeed, only the imaginary part of the coefficients are determined, leaving their real part free. However, the order of the first non linear term is well defined. This leads to the following definition.

**Definition 2.0.4.** We say that  $f$  is parabolic of *codimension  $k$*  if the first non linear term of  $f^\dagger$  is of order  $k + 1$ .

*Remarks 2.0.5.* 1. The formal series with real coefficients preserves the real axis. This indicates that  $f$  has a privileged unique direction which we will call a *formal symmetry axis*. Hence a conjugacy between two antiholomorphic parabolic germs must preserve the formal symmetry axis. We can of course suppose that this formal axis is the real axis. Note however that in the

case of even codimension, there is no canonical orientation of the formal symmetry axis.

2. The dynamics nearby the formal symmetry axis is a topological invariant. When  $k$  is odd, a rotation of angle  $\pi$  will flip the attractive semi-axis with the repulsive one. When  $k$  is even, both semi-axes are either attractive (when  $A_{k+1} < 0$ ) or repulsive (when  $A_{k+1} > 0$ ) (see Figures 3 and 4). In this paper, **we will only consider the case**  $A_{k+1} > 0$ . Indeed, when  $A_{k+1} < 0$ , i.e.  $f$  is of *negative type*, then  $f^{-1}$  will be of positive type and classifying  $f^{-1}$  is equivalent to classifying  $f$ .

**Definition 2.0.6.** When the codimension  $k$  is even, we say  $f$  is of *positive type* (resp. *negative type*) if  $A_{k+1} > 0$  (resp.  $A_{k+1} < 0$ ), where  $A_{k+1}$  is the first non zero coefficient in (4).




FIGURE 3. Dynamics near the formal symmetry axis of  $f(z) = \bar{z} + o(\bar{z})$  in *odd* codimension. One sector is attractive and the other repulsive; this yields the two possibilities above.




FIGURE 4. Dynamics near the formal symmetry axis of  $f(z) = \bar{z} + o(\bar{z})$  in *even* codimension. The possibilities are: on the left, both sectors are repulsive (positive type) and, on the right, both are attractive (negative type).

The composition of two antiholomorphic germs is holomorphic. Therefore, we will look at  $g := f \circ f$ , which is a holomorphic parabolic germ. Recall that in the holomorphic case, the codimension of  $g$  is the order of the first non zero term of  $g(z) - z$ . It is linked to the multiplicity of the fixed point:  $g$  is of codimension  $k$  if and only if the fixed point has multiplicity  $k + 1$ .

**Corollary 2.0.7.**  $f$  is of codimension  $k$  if and only if  $g = f \circ f$  is of codimension  $k$ .

The case when  $f \circ f = id$  is seen as a degenerate case where  $f$  is of “codimension infinity”. Indeed, it only happens if  $f$  is analytically conjugate to the complex conjugation, as is shown below. This case was excluded from our definition of parabolic point, since the fixed point of  $\sigma$  at the origin is not isolated.

**Proposition 2.0.8.** *Let  $f(z) = a_1\bar{z} + a_2\bar{z}^2 + a_3\bar{z}^3 + \dots$  be an antiholomorphic germ at the origin. The following statements are equivalent:*

1.  $f$  is formally conjugate to  $\sigma$ ;
2.  $f$  is analytically conjugate to  $\sigma$ ;
3.  $f \circ f = id$ .

*Proof.* 1.  $\Rightarrow$  3. Since there is a formal change of coordinate  $m$  such that  $m \circ f \circ m^{-1} = \sigma$ , we have  $m \circ f \circ f \circ m^{-1} = id$  formally, which yields  $f \circ f = id$ .

3.  $\Rightarrow$  2. Let us suppose that  $f \circ f = id$ . In particular,  $|a_1| = 1$ . We can of course suppose  $f$  is already in a coordinate such that  $a_1 = 1$ .

Let  $F_1(x, y) = \Re f(z)$  and  $F_2(x, y) = \Im f(z)$ , then we have

$$F(x, y) = \begin{pmatrix} F_1(x, y) \\ F_2(x, y) \end{pmatrix} = \begin{pmatrix} x + O(|(x, y)|^2) \\ -y + O(|(x, y)|^2) \end{pmatrix}.$$

We are interested in the fixed points of  $F$ , that is the zeros of  $F - id$ . Since  $\frac{\partial}{\partial y}(F_2(x, y) - y)|_{(0,0)} = -2$ , by the Implicit Function Theorem, there exists an analytic curve  $\gamma: t \mapsto t + i\eta(t)$  such that  $F_2(x, y) - y = 0$  if and only if  $y = \eta(x)$ .

We complexify  $t$  to obtain a change of coordinate  $t = \gamma^{-1}(z) = u + iv$  that rectifies the curve  $\gamma$  on the real line. Let  $\tilde{f} = \gamma^{-1} \circ f \circ \gamma$ . In the new coordinate,  $\tilde{F} = \gamma^{-1} \circ F \circ \gamma$  has now the form

$$\tilde{F}(u, v) = \begin{pmatrix} u + r(u, v) \\ -v(1 + O(|(u, v)|^2)) \end{pmatrix},$$

where  $r(u, v) = O(|(u, v)|^2)$ . The equation for fixed points  $\tilde{F} = id$  is equivalent to  $v = 0$  and  $r(u, 0) = 0$ . If  $r(u, 0) = au^s + o(u^s)$ ,  $a \neq 0$ , then this would contradict the fact that we must have  $\tilde{F} \circ \tilde{F}(u, 0) = \begin{pmatrix} u \\ 0 \end{pmatrix}$ . Therefore  $r(u, 0) \equiv 0$ , in other words  $r(u, v) = vp(u, v)$ .

We see that the real axis is a line of fixed points for  $\tilde{f}$  near the origin. By the Identity Theorem, because  $\tilde{f} \circ \sigma - id = 0$  on the real axis near the origin, we have  $\tilde{f} \equiv \sigma$ .

2.  $\Rightarrow$  1. This is immediate.  $\square$

The formal power series with real coefficients is used to determine a formal normal form for  $f$ . Recall that a formal normal form for



$g := f \circ f$  may be taken as the time-1 map of the flow of (see [6])

$$(5) \quad \dot{z} = \frac{z^{k+1}}{1 + bz^k}$$

for some constant  $b \in \mathbb{C}$ . We will call this constant  $b$  the *formal invariant*. It is also sometimes called the “résidu itératif” and, as mentioned in [4], it is determined by the holomorphic fixed point index, that is, the residue of  $\frac{1}{z-g(z)}$  at the origin.

When  $g = f \circ f$  is of codimension  $k$ , it is possible to get rid of the terms of degree  $k+1 < j < 2k+1$  by an analytic change of coordinate. In this coordinate,  $g$  is written

$$(6) \quad g(z) = z + z^{k+1} + \left( \frac{k+1}{2} - b \right) z^{2k+1} + o(z^{2k+1}),$$

where  $b \in \mathbb{C}$  is the formal invariant of  $g$ . When  $g$  is in this form, we will say that it is *prenormalized*.

**Definition 2.0.9.** The *formal invariant* of  $f$  is the formal invariant of  $f \circ f$ , which is the constant  $b$  in (6).

As the name suggests,  $b$  is invariant under formal changes of coordinate. Since  $g^\dagger := f^\dagger \circ f^\dagger$  and  $g$  have the same formal invariant, where  $f^\dagger$  is as in (4), it follows that  $b$  is real because all the coefficients of  $g^\dagger$  are real.

An important consequence of this, is that the time- $t$  map  $v^t$  of (5) for  $t \in \mathbb{R}$  has a power series at 0 with real coefficients, that is the complex conjugation  $\sigma$  and  $v^t$  commute.

**Proposition 2.0.10.** *Let  $v^{\frac{1}{2}}$  be the time- $\frac{1}{2}$  of the vector field (5) for some  $b$ . If  $f$  is of codimension  $k$ , of positive type if  $k$  is even and if  $f$  has formal invariant  $b$ , then  $f$  and  $\sigma \circ v^{\frac{1}{2}}$  are formally conjugate.*

*Proof.* Let  $f^\dagger$  be the formal power series with real coefficients formally conjugate to  $f$  in Proposition 2.0.2. Then  $f^\dagger \circ \sigma$  is a parabolic formal power series of  $z$ . A formal normal form may be chosen as  $v^{\frac{1}{2}}$ , the time- $\frac{1}{2}$  of the vector field (5). Since both the coefficients of  $v^{\frac{1}{2}}$  and  $f^\dagger$  are real, the formal change of coordinate  $h$  commutes with  $\sigma$ , provided that  $h'(0) = 1$ , so that  $v^{\frac{1}{2}} \circ \sigma$  is formally conjugate to  $f$ .  $\square$

The formal change of coordinate conjugating  $f$  to its formal normal form can always be truncated at the  $(2k+2)$ -th term, which yields a holomorphic change of coordinate taking  $f$  to the form

$$(7) \quad f(z) = \bar{z} + \frac{1}{2} \bar{z}^{k+1} + \left( \frac{k+1}{8} - \frac{b}{2} \right) \bar{z}^{2k+1} + o(\bar{z}^{2k+1}),$$

that is  $f$  and  $\sigma \circ v^{\frac{1}{2}}$  have the same first three terms.

**Definition 2.0.11.** When  $f$  is in the form (7), we will say that it is *prenormalized*.

*Remark 2.0.12.* In even codimension,  $f$  may only be prenormalized as in (7) when  $A_{k+1} > 0$ . In odd codimension,  $f$  may always be prenormalized as in (7).

The formal normal form is a model to which the germs can be compared. Now that this form has been established, we describe its properties.

### 3. PROPERTIES OF THE FORMAL NORMAL FORM

Let us start with the following observations.

**Proposition 3.0.1.** *Let  $v$  be the vector field (5) of codimension  $k$  and formal invariant  $b$ .*

1.  $v$  is invariant under the rotations of order  $k$ .
2.  $v$  is invariant under the complex conjugation  $\sigma$  when  $b$  is real.

The holomorphic and antiholomorphic formal normal forms are respectively

$$(8) \quad v^1(z) = z + z^{k+1} + \left( \frac{k+1}{2} - b \right) z^{2k+1} + o(z^{2k+1})$$

$$(9) \quad \sigma \circ v^{\frac{1}{2}}(z) = \bar{z} + \frac{1}{2} \bar{z}^{k+1} + \left( \frac{k+1}{8} - \frac{b}{2} \right) \bar{z}^{2k+1} + o(\bar{z}^{2k+1}),$$

where  $v^t$  is the time- $t$  of  $v$ .

We see that the real axis is a symmetry axis. We introduce a notation for the other symmetry axes.

**Definition 3.0.2.** Let  $\sigma_\ell$  denote the reflection

$$(10) \quad \sigma_\ell(z) := e^{\frac{2i\pi\ell}{k}} \bar{z}, \quad \text{for } \ell = 0, \dots, k-1.$$

**Corollary 3.0.3.**

1.  $v$  is invariant under  $\sigma_\ell$  for  $\ell = 0, \dots, k-1$  when  $b$  is real;
2.  $v^1$  commutes with any rotation of order  $k$ , and when  $b$  is real, it commutes with  $\sigma_\ell$  for  $\ell = 0, \dots, k-1$ ;
3. When  $k$  is even,  $\sigma \circ v^{\frac{1}{2}}$  commutes with  $z \mapsto -z$ .

We will only be interested in real values of  $b$ .

**Proposition 3.0.4** (Roots of the normal forms).

1. For  $n$  even,  $v^1$  has  $k$  one-parameter families of antiholomorphic  $n$ -th roots given by  $\sigma_\ell \circ v^{\frac{1}{n}+iy}$  for  $y \in \mathbb{R}$ ,  $\ell = 0, \dots, k-1$ .
2. For  $n$  odd,  $\sigma \circ v^{\frac{1}{2}}$  has exactly one antiholomorphic  $n$ -th root given by  $\sigma \circ v^{\frac{1}{2n}}$ .

We ask the following questions, which will be answered in Section 7.2.

*Question 3.0.5.* For a holomorphic parabolic germ  $g$ , how many distinct antiholomorphic  $n$ -th roots ( $n$  even) does it have?

*Question 3.0.6.* For an antiholomorphic parabolic germ  $f$  and  $n$  odd, when is the formal  $n$ -th root convergent?

#### 4. FATOU COORDINATES

For the whole section, when the codimension of  $f$  is even, we will suppose  $f$  is of positive type (see Definition 2.0.6). The formal normal form  $\sigma \circ v^{\frac{1}{2}}$  is a model to which it is natural to compare the antiholomorphic germ  $f$ . In the holomorphic study of parabolic germs, we use holomorphic diffeomorphisms called Fatou coordinates defined on sectors covering the origin on which the germ is conjugated to its normal form, i.e. changes of coordinates to the normal form. We then compare Fatou coordinates on the intersection of the sectors, thus yielding a conformal invariant describing the space of orbits of the germ. See [5] or [6] for the details.

The same approach can be adapted to the antiholomorphic case. It will be necessary to find a sectorial normalization (Fatou coordinates) of the antiholomorphic germ  $f$ . However, instead of adapting the construction of the holomorphic case, we will prove that it is possible to choose Fatou coordinates of  $f \circ f$ , which is holomorphic, that are also Fatou coordinates of  $f$ .

**4.1. Rectifying Coordinates and sectors.** Suppose that an antiholomorphic parabolic germ  $f$  is of codimension  $k$  for  $k \geq 1$  with a formal invariant  $b$  (see Def. 2.0.9). The Fatou coordinates  $\varphi_j$  are often constructed in the rectifying coordinate given by the time of the vector field (5). Since  $v^{\frac{1}{2}}$  and  $v^1$  are the time maps of the vector field (5), we define the time coordinate by

$$(11) \quad Z(z) = \int_{z_0}^z \frac{1 + b\zeta^k}{\zeta^{k+1}} d\zeta = -\frac{1}{kz^k} + b \log z + \frac{1}{kz_0^k} - b \log z_0$$

which is multi-valued. See Figure 5 for its Riemann surface. It is the inverse of the flow of (5) with starting point  $z_0$ . We will single out the

following  $2k + 1$  charts of  $Z$ :

$$(12) \quad Z_j(z) = -\frac{1}{kz^k} + b \log z - \frac{ji\pi b}{k}$$

for  $j = -k, \dots, -1, 0, 1, \dots, k$ ,

where  $\log z$  is determined by  $\arg z \in (-\pi, \pi)$  for  $-k < j < k$ , and for  $Z_k$  (resp.  $Z_{-k}$ ),  $\arg(\cdot)$  will be the continuation in  $(0, 2\pi)$  (resp. in  $(-2\pi, 0)$ ). In particular, we see that  $Z_k = Z_{-k}$ , and that *both  $Z_0$  and  $Z_k$  commute with the complex conjugation*.




FIGURE 5. The Riemann surface of the time coordinate  $Z$ . The hole in the middle correspond to the image of  $\mathbb{C} \setminus D(0, r)$  in the  $z$ -coordinate, while a neighbourhood of  $z = 0$  is sent to a neighbourhood of infinity. A curve going  $k$  times around the hole in the  $Z$ -coordinate will turn one time around  $\infty$  in the  $z$ -coordinate.

Now we define the sectors in the  $z$ -space (see Figure 7). On the Riemann surface of  $Z_j$ , we write  $G_j$  for the expression of  $g := f \circ f$  in the  $Z_j$ -coordinate. Let  $z_j^* = \delta e^{\frac{ij\pi}{k}}$  for  $-k \leq j \leq k$  and some small enough  $\delta > 0$ . Let  $Z_j^*$  be the image of  $Z_j(z_j^*)$ . We consider a vertical line  $\ell_j$  passing through  $Z_j^*$  and its image  $G_j(\ell_j)$ . Let  $B_j$  be the domain bounded by  $\ell_j$  and  $G_j(\ell_j)$  and containing  $\ell_j$  and  $G_j(\ell_j)$ . The sector in the  $Z_j$ -coordinate is then obtained by

$$U_j = \{Z_j \mid \exists n \in \mathbb{Z}, G_j^{\circ n}(Z_j) \in B_j\}$$

for  $-k \leq j \leq k$  (see Figure 6). We see that  $U_{-k} = U_k$ , since  $Z_k = Z_{-k}$ . The sector  $S_j$  in the  $z$ -coordinate is  $Z_j^{-1}(U_j)$  (see Figure 7). These sectors are sometimes called *Fatou petals*. They are described in great details in [2], although the authors only consider *attractive* petals. Note that there are  $2k$  petals, with half of them being *repulsive* (see Figure 8). Also,  $S_k$  and  $S_{-k}$  are the same petal.




FIGURE 6. A chart  $U_j$  on the Riemann surface with the vertical strip  $B_j$

*Remark 4.1.1.* Note that

$$2i\pi b = \int_{\partial D(0,\delta)} \frac{1+bz^k}{z^{k+1}} dz = \sum_j \int_{Z_j(\gamma_j)} dZ_j = \sum_{j=-k+1}^k \left( Z_j(z_{j+1}) - Z_j(z_j) \right)$$

where  $\gamma_j$  is an arc of the circle  $\partial D(0,\delta)$  in  $S_j$ , with endpoints  $z_{j+1}$  and  $z_j$ , where  $z_j = \delta e^{\frac{i(2j-1)\pi}{2k}}$ . The  $Z_j$  defined as in (12) satisfy this condition.




FIGURE 7. The particular case of  $z = z^4$ . On the right, the sector  $U_0$  in the  $Z_0$ -coordinate, obtained from a strip (in dark gray). On the left,  $S_0 = Z^{-1}(U_0)$  the sector in  $z$ , with the preimage of the strip (in dark gray).

The sectors are ordered as in Figure 9. Note in particular that  $S_0$  intersects the positive real axis, and  $S_k = S_{-k}$ , the negative real axis.

**Definition 4.1.2.** The *time coordinate* is the Riemann surface obtained from the disjoint union of the  $U_j$ , glued together by the transition functions: the charts are the  $U_j \hookrightarrow \mathbb{C}$ , with the diffeomorphism  $Z_j: S_j \rightarrow U_j$  given by

$$Z_j(z) = -\frac{1}{kz^k} + b \log z - \frac{ji\pi b}{k},$$

and the transition functions are  $Z_j \circ Z_{j-1}^{-1} = T_{\frac{-i\pi b}{k}}$  for  $-k < j \leq k$  where the composition is defined.




FIGURE 8. Petals for the holomorphic map  $f \circ f$ . Dynamics inside a repulsive petal on the left. Dynamics inside an attracting petal on the right.




FIGURE 9. Ordering of the sectors for  $k = 3$ .

The time coordinate is conformally equivalent to a punctured disk of the origin.

Now we define the complex conjugation on the time coordinate. Note that on a subdomain  $S'_0 \subseteq S_0$  such that  $\sigma(S'_0) = S'_0$ , we have  $Z_0(\bar{z}) = \overline{Z_0(z)}$ . The complex conjugation on the time coordinate is then obtained by analytic continuation on the other charts  $U_j$ .

**Proposition 4.1.3** (Complex conjugation). *For  $z \in S_j$ , let  $\ell$  be such that  $\sigma(z) = \bar{z} \in S_\ell$ . We define the complex conjugation  $\Sigma$  on the time coordinate in the charts by*

$$\Sigma_{\ell,j} \circ Z_j(z) = Z_\ell \circ \sigma(z).$$

*Then  $\Sigma$  is well-defined and  $\Sigma \circ \Sigma = id$ .*

*Proof.* The proof consists of showing that  $\Sigma$  is compatible on both charts when  $Z_j \in U_j \cap U_{j+1}$  or when  $\Sigma(Z_j) \in U_\ell \cap U_{\ell \pm 1}$ . It is a simple computation. Note that for a subdomain  $S'_j \subset S_j$  such that  $\sigma(S'_j) \subset S_{-j}$ , then in the charts, we have  $\Sigma_{-j,j}(Z_j) = \overline{Z_j}$ .  $\square$

This allows us to talk about the normal form  $\sigma \circ v^{\frac{1}{2}}$  in the time coordinate. It is the antiholomorphic map  $\Sigma \circ T_{\frac{1}{2}}$ .

**4.2. Fatou Coordinates.** Let us call the petals of the normal form  $S_j^v$ . The orbits of the normal form  $\sigma \circ v^{\frac{1}{2}}$  jump from  $S_j^v$  to  $S_{-j}^v$ . This means that the dynamics of those two petals is no longer independent, unlike the holomorphic case. See Figure 10.

In its prenormalized form  $f$  is close to  $\sigma \circ v^{\frac{1}{2}}$  in the sense that  $|f - \sigma \circ v^{\frac{1}{2}}| = o(|z|^{2k+1})$ . In the following lemma, we prove that it is also true that  $F$  and  $\Sigma \circ T_{\frac{1}{2}}$  are close in the time coordinate.




FIGURE 10. On the left, an orbit of  $\sigma \circ v^{\frac{1}{2}}$  in the  $z$  coordinate. The orbit jumps between the sectors  $S_j^v$  and  $S_{-j}^v$  of  $\sigma \circ v^{\frac{1}{2}}$ . On the right, the same orbit is represented in the time coordinate; it is the orbit of  $\Sigma \circ T_{\frac{1}{2}}$ .

**Lemma 4.2.1.** *Let  $f$  be in its prenormalized form (7) and let  $F$  (resp.  $\Sigma \circ T_{\frac{1}{2}}$ ) be the expressions of  $f$  (resp.  $\sigma \circ v^{\frac{1}{2}}$ ) in the time coordinate. On each chart  $U_j$ , we have  $|F - \Sigma \circ T_{\frac{1}{2}}| = O(|Z|^{-1})$ .*

*Proof.* The proof is similar to that found in [6]. Let  $m(z) = f \circ (\sigma \circ v^{\frac{1}{2}})^{-1}(z) = z + o(z^{2k+1})$ . We see that

$$\begin{aligned} Z_j \circ m(z) &= -\frac{1}{kz^k}(1 + o(z^{2k})) + b \log z + o(z^{2k}) - \frac{j\pi b}{k} \\ &= Z_j(z) + o(z^k). \end{aligned}$$

Since  $z^k Z_j(z) \rightarrow -\frac{1}{k}$  when  $z \rightarrow 0$ , and because  $Z_j$  is invertible and  $|Z_j(z)| \rightarrow \infty$  when  $z \rightarrow 0$ , it follows that  $o(z^k)$  is  $O(|Z_j|^{-1})$  when  $|Z_j| \rightarrow \infty$ . Therefore,  $F \circ (\Sigma \circ T_{\frac{1}{2}})^{-1}(Z_j) = Z_j + O(|Z_j|^{-1})$ .  $\square$

We now present the existence of the Fatou coordinates. Note that Hubbard and Schleicher proved their existence in [4] (Lemma 2.3) in the codimension 1 case for a map with a parabolic periodic orbit of odd period  $n$ . We recover their case by considering  $f^{on}$ . This corresponds

for us to a germ of antiholomorphic parabolic diffeomorphism of codimension 1. The proof in higher codimension is in the same spirit with an adaptation, since we need to work with pairs of sectors  $(U_j, U_{-j})$ .

**Proposition 4.2.2.** *Let  $F$  and  $\Sigma$  be the expression of  $f$  and  $\sigma$  in the time coordinates respectively. Recall that  $U_j = Z_j(S_j)$ . On each  $U_j$ , there exists a holomorphic diffeomorphism  $\Phi_j: U_j \rightarrow \mathbb{C}$  such that*

$$(13) \quad \Phi_j \circ F \circ \Phi_{-j}^{-1} = \Sigma \circ T_{\frac{1}{2}},$$

whenever the composition is defined.

Moreover, if  $\tilde{\Phi}_j$  are other Fatou coordinates, then there exists  $C_j \in \mathbb{C}$  for  $j = 1, \dots, k-1$  and  $C_0, C_k \in \mathbb{R}$  such that  $\Phi_j \circ \tilde{\Phi}_j^{-1} = T_{C_j}$  and  $\Phi_{-j} \circ \tilde{\Phi}_{-j}^{-1} = T_{\bar{C}_j}$  for  $j \geq 0$ .

*Proof.* The proof makes use of the rigidity of the conformal structure of the doubly punctured sphere  $S^2 \setminus \{0, \infty\}$ , as in the proof of the uniqueness in the holomorphic case.

Let  $\Phi_j$  denote the Fatou coordinate of  $f \circ f$  on  $U_j$ , that is

$$\Phi_j \circ (F \circ F) \circ (\Phi_j)^{-1} = T_1.$$

(We know that it exists since  $f \circ f$  is holomorphic and that it is unique up to left-composition with a translation.) In the space of the Fatou coordinate  $W_j = \Phi_j(Z_j)$ , an orbit  $\{(f \circ f)^{on}(z)\}$  corresponds to  $\{W_j + n\}$ . Note also that  $\Phi_j(Z_j) = Z_j + D_j + O(|Z_j|^{-1})$ , for some constant  $D_j \in \mathbb{C}$  (see [6]).

We first note that each  $\Phi_j(U_j)$  contains a vertical strip  $B_j$  of width 1, by construction of the time coordinate  $U_j$  and of the Fatou coordinate. We define

$$Q_j = \Phi_{-j} \circ F \circ (\Phi_j)^{-1} \quad \text{for } -k \leq j \leq k.$$

Then we see that  $Q_j \circ Q_{-j} = T_1$  since  $\Phi_j$  are Fatou coordinates of  $F \circ F$ . It follows that  $Q_j$  commutes with  $T_1$ , since

$$\begin{aligned} T_1 \circ Q_j &= (Q_j \circ Q_{-j}) \circ Q_j \\ &= Q_j \circ (Q_{-j} \circ Q_j) \\ &= Q_j \circ T_1. \end{aligned}$$

Indeed,  $Q_j$  represents  $F$  in the Fatou coordinates. It is therefore natural that  $Q_j$  commutes with  $T_1$ , which represents  $F \circ F$  in the Fatou coordinates.

Because  $Q_j$  is the composition of an antiholomorphic germ by a holomorphic diffeomorphism,  $Q_j$  is antiholomorphic. In particular,  $\Sigma \circ$



$Q_j$  is holomorphic, and  $\Sigma \circ Q_j - id$  is 1-periodic and holomorphic, so it has a Fourier expansion

$$\Sigma \circ Q_j(W_j) - W_j = \sum_{n=-\infty}^{\infty} c_{n,j} e^{2i\pi n W_j}.$$

Moreover, by lemma 4.2.1, we have  $\Sigma \circ Q_j(W) = W + M_j + O(|W|^{-1})$ , where  $M_j \in \mathbb{C}$  is a constant. Therefore,  $|\Sigma \circ Q_j - id|$  is bounded when  $|W| \rightarrow \infty$ , so we must have  $c_{n,j} = 0$  for  $n \in \mathbb{Z}^*$ .

We conclude that  $Q_j(W_j) = \overline{W_j} + \overline{c_{j,0}}$ . Since  $Q_j \circ Q_j = T_1$ , it follows that  $c_{j,0} = \frac{1}{2} + iy$ . We then adjust all the  $\Im c_{j,0}$  to 0 by choosing the appropriate Fatou coordinates (i.e. composing them with a translation).

The uniqueness comes from a combination of the uniqueness of the Fatou coordinates for the holomorphic  $f \circ f$  and having to preserve the constants  $c_{j,0} = \frac{1}{2}$ .  $\square$

## 5. MODULUS OF ANALYTIC CLASSIFICATION

If two antiholomorphic parabolic germs are analytically conjugate, then they have the same space of orbits. The space of orbits of an antiholomorphic parabolic germ  $f$  is a quotient of the set of orbits of the associated holomorphic parabolic germ  $g = f \circ f$ . Hence we start by describing the space of orbits of  $g$ ; on a Fatou coordinate, it is the quotient by  $T_1$ , which is a bi-infinite cylinder. We also need to identify some orbits represented in two different Fatou coordinates. This is done by means of the transition maps (the horn maps of Écalle).

We will describe the space of orbits of  $f$  in Section 6.1 and classify the antiholomorphic germs in Section 6.3. To do both of these, we will need the *transition functions*, which will form an analytic invariant.

The transition functions we describe here are the same as for the holomorphic case. We will introduce what we need here; all the details are found in [6] or [5].

In the time coordinate, if  $U_j$  is a repelling (resp. attractive) petal, then  $U_j$  and  $U_{j+1}$  intersect on a domain containing an upper half-plane (resp. a lower half-plane), see Figure 11. We can compare the Fatou coordinates  $\Phi_j$  and  $\Phi_{j+1}$  by looking at

$$\Psi_j: V_j \xrightarrow{\Phi_j^{-1}} U_j \cap U_{j+1} \xrightarrow{\Phi_{j+1}} V_{j+1},$$

resp.

$$\Psi_j: V_{j+1} \xrightarrow{\Phi_{j+1}^{-1}} U_{j+1} \cap U_j \xrightarrow{\Phi_j} V_j,$$

where  $V_j = \Phi_j(U_j)$  for all  $j$ . This yields a diffeomorphism defined on a domain of  $V_j$  (resp.  $V_{j+1}$ ) containing an upper-half plane (resp.

lower half-plane) with its image in  $V_{j+1}$  (resp.  $V_j$ ) also containing some upper-half plane (resp. lower-half plane).




FIGURE 11. Charts  $U_{j-1}$  and  $U_j$  on the time coordinate. They intersect on a region containing (in this case) an upper half-plane.

Notice the order of the composition for  $\Psi_j$ : we choose the convention that these functions will go from a repulsive petal to an attractive petal. Figure 12 is an illustration of the direction of the arrows in the  $z$ -coordinate for  $k = 3$ , where  $\xi_j$  is the expression of  $\Psi_j$  in the  $z$ -coordinate.

**Definition 5.0.1.** Let  $\Phi_j$  be a Fatou coordinate of  $f$  on  $U_j$ . The *transition functions* (equivalent to the Écalle horn maps) of  $f$  are the  $2k$  functions  $\Psi_j$  for  $j = 1, \dots, k$  and  $j = -1, \dots, -k$  obtained by

$$(14) \quad \Psi_j = \begin{cases} \Phi_j \circ \Phi_{j-\text{sgn}(j)}^{-1}, & \text{for } j \text{ odd;} \\ \Phi_{j-\text{sgn}(j)} \circ \Phi_j^{-1}, & \text{for } j \text{ even;} \end{cases}$$

where the composition is defined. Here,  $\text{sgn}(j)$  is the sign of  $j$ .




FIGURE 12. Direction of the transition functions  $\{\xi_j\}_j$  represented in the  $z$ -coordinates.

By the uniqueness of Proposition 4.2.2, we may change  $\Phi_{\pm j}$  by  $T_{C_j} \circ \Phi_j$  and  $T_{\overline{C_j}} \circ \Phi_{-j}$  for some  $C_j \in \mathbb{C}$  for  $j = 1, \dots, k-1$ , or  $\Phi_j$  by  $T_{R_j} \circ \Phi_j$  for some  $R_j \in \mathbb{R}$  for  $j = 0, k$ . This will yield another set of  $2k$  transition functions. We will identify together these different possible choices of transition functions at the end of this section.

The following proposition is the first step towards the geometric invariant. The transition functions allow to describe the space of orbits of  $F$  and  $F \circ F$ .

**Proposition 5.0.2.** *Let  $(\Psi_{-k}, \dots, \Psi_{-1}, \Psi_1, \dots, \Psi_k)$  be transition functions of  $f$ . They satisfy the equation*

$$(15) \quad \Sigma \circ T_{\frac{1}{2}} \circ \Psi_j = \Psi_{-j} \circ \Sigma \circ T_{\frac{1}{2}}.$$

*In particular, they are transition functions of  $f \circ f$  and satisfy*

$$(16) \quad T_1 \circ \Psi_j = \Psi_j \circ T_1.$$

*Proof.* The proof is identical to the holomorphic case; it follows from the definition of  $\Psi_j$  and (13).  $\square$

Equation (15) says that the orbits of  $\Sigma \circ T_{\frac{1}{2}}$  in one Fatou coordinate are sent by the  $\Psi_j$  on the orbits of  $\Sigma \circ T_{\frac{1}{2}}$  in another Fatou coordinate. In those coordinates, the orbits of  $\Sigma \circ T_{\frac{1}{2}}$  correspond to those of  $f$ . Therefore, the transition functions allow to identify the same orbits of  $f$  in different coordinates.

We can rewrite Equation (15) as

$$(17) \quad \Psi_{-j} = \Sigma \circ T_{\frac{1}{2}} \circ \Psi_j \circ \Sigma \circ T_{-\frac{1}{2}},$$

so that  $\Psi_{-j}$  is determined by  $\Psi_j$ . Thus we only need half of the transition functions of  $f$  to determine all of them. For the rest of the paper, we will work with  $\Psi_1, \dots, \Psi_k$ , knowing that  $\Psi_{-1}, \dots, \Psi_{-k}$  are obtained from Equation (17).

The transition functions in the holomorphic case are well known and their properties are described in [5] by Ilyashenko. Because the transition functions of  $f$  are also those of  $f \circ f$ , they share the properties which we describe now.

Each  $\Psi_j$  satisfies Equation (16); it follows that  $\Psi_j - id$  is 1-periodic and has a Fourier expansion

$$(18) \quad \begin{aligned} \Psi_j(W) - W &= c_j + \sum_{n=1}^{\infty} c_{n,j} e^{2i\pi n W} & \text{for } j > 0 \text{ odd;} \\ \Psi_j(W) - W &= c_j + \sum_{n=-1}^{-\infty} c_{n,j} e^{2i\pi n W} & \text{for } j > 0 \text{ even.} \end{aligned}$$

In particular, we see that  $|\Psi_j - id - c_j|$  is exponentially decreasing when  $\Im W \rightarrow \infty$  and  $j$  is odd (resp.  $\Im W \rightarrow -\infty$  and  $j$  is even).

Since the Fatou coordinates are not unique, we may change them and obtain new transition functions. This will change the constants  $c_j$  and  $c_{n,j}$ , but the following alternating sum will always be preserved

$$(19) \quad \begin{aligned} & (-1)^{k-1}c_{-k} + (-1)^{k-2}c_{-k+1} \pm \cdots - c_{-2} + c_{-1} \\ & - c_1 + c_2 - \cdots + (-1)^{k-1}c_{k-1} + (-1)^k c_k = 2i\pi b. \end{aligned}$$

We successively change  $\Phi_1$  by  $T_{-c_1 - \frac{i\pi b}{k}} \circ \Phi_1$ ,  $\Phi_2$  by  $T_{-c_1 + c_2 - \frac{2i\pi b}{k}}$ ,  $\Phi_3$  by  $T_{-c_1 + c_2 - c_3 - \frac{3i\pi b}{k}} \circ \Phi_3$ , and so on. The constant terms of the new transition functions will satisfy,

$$(20) \quad c_j = \begin{cases} -\frac{i\pi b}{k}, & \text{for } j > 0 \text{ odd;} \\ \frac{i\pi b}{k}, & \text{for } j > 0 \text{ even.} \end{cases}$$

for  $j = 1, 2, \dots, k$ . Note that, together with Equation (17), the constant terms  $c_j$  for  $j = -k, \dots, -1$  are given by  $c_j = \overline{c_{-j}}$ .

**Definition 5.0.3** (Normalized transition functions). The transition functions  $(\Psi_1, \dots, \Psi_k)$  are said to be *normalized* if the constant terms  $c_j$  of Equation (18) satisfy (20) for  $j = 1, 2, \dots, k$ .

Even when normalized, the transition functions are not uniquely determined. There is still a remaining degree of freedom: we may change the source and target space of each  $\Psi_j$  by the same translation  $T_C$ , with  $C \in \mathbb{R}$ .

In the case of even codimension, if  $f$  is prenormalized, then  $f_1(z) = -f(-z)$  is also prenormalized, and  $f$  and  $f_1$  are conjugate. Hence, we will need to identify their moduli.

**Definition 5.0.4** (Modulus of classification). We define the equivalence relation  $\sim$  on normalized transition functions of the form (18) by

$$(21) \quad (\Psi_1, \dots, \Psi_k) \sim (\Psi'_1, \dots, \Psi'_k) \Leftrightarrow \exists C \in \mathbb{R}, \Psi_j = T_C \circ \Psi'_j \circ T_{-C}.$$

The *modulus of classification* of  $f$  of codimension  $k$  (and of positive type when  $k$  is even) (see Def. 2.0.6) is defined by the triple  $(k, b, [\Psi_1, \dots, \Psi_k])$ , where  $k$  is the codimension (Def. 2.0.4),  $b$  is the formal invariant (Def. 2.0.9), and  $[\Psi_1, \dots, \Psi_k]$  is the equivalence class of normalized transition functions of  $f$  (Defs. 5.0.1 and 5.0.3), where the equivalence class is defined according to the following two cases:

1. If  $k$  is odd, then  $[\Psi_1, \dots, \Psi_k]$  is the equivalence class under the Relation (21);
2. If  $k$  is even, then  $[\Psi_1, \dots, \Psi_k]$  is the equivalence under the Relation (21) and the additional relation

$$(22) \quad (\Psi_1, \dots, \Psi_k) \cong (\Psi'_1, \dots, \Psi'_k) \iff \Psi'_j = \Sigma \circ T_{\frac{1}{2}} \circ \Psi_{k-j+1} \circ \Sigma \circ T_{-\frac{1}{2}}.$$

This equivalence class is called the *analytic invariant*.

**5.1. Remarks on the Écalle-Voronin Modulus.** In the holomorphic case, the modulus of classification is known as the *Écalle-Voronin modulus* (see [5] and [6]). For any holomorphic parabolic germ  $g$  (not necessarily of the form  $g = f \circ f$ ), we can obtain its Écalle-Voronin modulus the same way as described above, but without equation (15), so that the  $2k$  transition functions are needed. There are  $2k$  degrees of freedom, so we normalize the transition functions by choosing the constant terms as in (20) and with  $c_{-j} = -c_j$ ; the remaining degree of freedom is a translation in every Fatou coordinate by a constant  $C \in \mathbb{C}$ . Therefore, we quotient by the equivalence relation

$$(23) \quad (\Psi_{-k}, \dots, \Psi_{-1}, \Psi_1, \dots, \Psi_k) \sim_{\mathbb{C}} (\Psi'_{-k}, \dots, \Psi'_{-1}, \Psi'_1, \dots, \Psi'_k) \\ \iff \exists C \in \mathbb{C}, \Psi_j = T_C \circ \Psi'_j \circ T_{-C} \\ \text{for } j \in \{-k, \dots, -1\} \cup \{1, \dots, k\}.$$

We also quotient by the action of the rotations of order  $k$ . If we note the indices  $\Psi_{-j} = \Psi_{2k-j+1}$  for  $j = 1, \dots, k$ , then we have the identification

$$(24) \quad (\Psi_{k+1}, \dots, \Psi_{2k}, \Psi_1, \dots, \Psi_k) \\ \sim (\Psi'_{k+1+2m}, \dots, \Psi'_{2k+2m}, \Psi'_{1+2m}, \dots, \Psi'_{k+2m}) \\ \text{for } m = 0, \dots, k-1,$$

where indices are mod  $2k$ . We will note the equivalence class of both of these identifications by  $[\Psi_{-k}, \dots, \Psi_{-1}, \Psi_1, \dots, \Psi_k]$ . The modulus of  $g$  is then  $(k, b, [\Psi_{-k}, \dots, \Psi_{-1}, \Psi_1, \dots, \Psi_k])$ .

We describe the link between the Écalle-Voronin modulus of  $g$  and the modulus of classification of an antiholomorphic parabolic germ  $f$  of positive type such that  $g = f \circ f$ , when such a  $f$  exists. For  $k$  odd (resp.  $k$  even), the symmetry axis of  $f$  appears along one of the  $k$  (resp.  $\frac{k}{2}$ ) “symmetry axes” of  $g$  of the form  $e^{\frac{2i\ell\pi}{k}}\mathbb{R}$ ,  $\ell = 0, \dots, k-1$ . Therefore, we associate to the analytic invariant of  $g$  the regular  $k$ -gon with the  $k$  symmetry axes  $e^{\frac{i\ell\pi}{k}}\mathbb{R}$  (see Figure 13) in the following way. Divide the  $k$ -gon by its  $k$  symmetry axes to produce  $2k$  sectors in the  $k$ -gon, then starting with the sector above the horizontal line on the right side, we identify this sector with  $\Psi_1$  and going anti-clockwise, we associate

$\Psi_2, \dots, \Psi_k, \Psi_{-k}, \dots, \Psi_{-1}$  to the subsequent sectors, as in Figure 13. The dihedral group  $D_{2k}$  acts on the sectors of the  $k$ -gon. The action is defined for an element  $u \in D_{2k}$  by mapping a sector to its image by the linear application represented by  $u$  (a rotation or a reflection).

**Definition 5.1.1.** Let  $u \in D_{2k}$  and let  $\Delta_j$  be the sector of the  $k$ -gon associated with  $\Psi_j$ . The action of  $u$  on the sectors of the regular  $k$ -gon defines a permutation on  $\{-k, \dots, -1, 1, \dots, k\}$  also noted  $u$  by abuse of notation, defined so that  $u^{-1}(\Delta_j) = \Delta_{u(j)}$  (see Figure 13).




FIGURE 13. Pentagon with 10 sectors and the transition functions, for codimension 5, with the symmetry axis of  $f$  in bold in (a) and (b).

Let  $(\Psi_{-k}, \dots, \Psi_{-1}, \Psi_1, \dots, \Psi_k)$  be a representative of the analytic invariant of  $g$ . Let us suppose that the symmetry axis of  $f$  is along the reflection axis of  $s \in D_{2k}$ . We rotate the coordinate by, say, an angle of  $\frac{2\ell i\pi}{k}$ , so that the symmetry axis corresponds to the (repulsive) real axis (this is possible because  $f$  is of positive type). The rotation corresponds to an element  $r \in D_{2k}$  and a permutation of indices, thus we obtain  $r^{-1}sr = s_0$ , where  $s_0(j) = -j$  is the permutation for  $s$  in the new coordinate. The representative of the modulus of  $g$  is permuted:  $(\Psi_{r(-k)}, \dots, \Psi_{r(-1)}, \Psi_{r(1)}, \dots, \Psi_{r(k)})$  as in (24); it now has an equivalent representative  $(\Psi'_{r(-k)}, \dots, \Psi'_{r(-1)}, \Psi'_{r(1)}, \dots, \Psi'_{r(k)})$  under Relation (23) obtained from the Fatou coordinates of  $f$ , so that it satisfies

$$\Psi'_{r(j)} \circ \Sigma \circ T_{\frac{1}{2}} = \Sigma \circ T_{\frac{1}{2}} \circ \Psi'_{r(-j)}.$$

or equivalently

$$(25) \quad \Psi'_j \circ \Sigma \circ T_{\frac{1}{2}} = \Sigma \circ T_{\frac{1}{2}} \circ \Psi'_{s(j)}.$$

The analytic invariant of  $f$  is then  $[\Psi'_{r(1)}, \dots, \Psi'_{r(k)}]$ .

Lastly, we will talk about the modulus of the inverse of a holomorphic parabolic germ  $g$ . The following proposition is probably well-known, but we could not find it in the literature, so the proof is included.

**Proposition 5.1.2.** *Let  $g$  be a holomorphic parabolic germ. Let the modulus of  $g$  be  $(k, b, [\Psi_{-k}, \dots, \Psi_{-1}, \Psi_1, \dots, \Psi_k])$ . Then the modulus of  $g^{-1}$  is given by*

$$(26) \quad (k, -b, [\tilde{\Psi}_{-k}, \dots, \tilde{\Psi}_{-1}, \tilde{\Psi}_1, \dots, \tilde{\Psi}_k]),$$

where  $\tilde{\Psi}_j$  stands for  $L_{-1} \circ \Psi_{r_1^{-1}(j)}^{-1} \circ L_{-1}$ , with  $L_{-1}: Y \mapsto -Y$  and with  $r_1$ , the rotation of the indices induced by  $y \mapsto e^{\frac{i\pi}{k}} y$  (see Definition 5.1.1).

*Proof.* Suppose  $g$  is prenormalized. The dynamics of  $g^{-1}$  is reversed, therefore the dynamics of  $g^{-1}$  in the first sector  $S_0$  is attractive, but the dynamics in  $S_{-1}$  is repulsive. We apply the change of coordinate  $y = L_\lambda(z) = \lambda z$ , where  $\lambda = e^{\frac{i\pi}{k}}$ . Let  $\tilde{g}^{-1} = L_\lambda \circ g^{-1} \circ L_\lambda^{-1}$ .

At the formal level, we apply  $(z, t) \mapsto (y, -t)$  to the vector field (5) to obtain

$$(27) \quad \dot{y} = w(y) = \frac{y^{k+1}}{1 - by^k}.$$

The formal normal form of  $\tilde{g}^{-1}$  is the time-1 map  $w^1$  of (27). In particular,  $\tilde{g}^{-1}$  has formal invariant  $-b$ . We will denote the sectors of  $\tilde{g}^{-1}$  in the  $y$ -coordinate by  $\tilde{S}_j$ . Note that  $\tilde{S}_j = L_\lambda^{-1}(S_j) = S_{j-1}$ .

The time coordinate on the sector  $\tilde{S}_j$  is defined as

$$(28) \quad Y_j(y) = \frac{-1}{ky^k} - b \log y + \frac{j\pi b}{k}.$$

We have the relation  $L_{-1} \circ Y_j = Z_{j-1} \circ L_\lambda^{-1}$ , where the  $Z_j$ 's are the time coordinates of  $g$ . Indeed, we see that

$$Z_{j-1}(e^{\frac{-i\pi}{k}} y) = \frac{1}{ky^k} + b \log y - \frac{i\pi b}{k} - \frac{(j-1)\pi b}{k} = -Y_j(y).$$

Let  $\Phi_j$  be a Fatou coordinate of  $g$  in  $S_j$ . We know that  $\Phi_j \circ Z_j \circ g^{-1} \circ Z_j^{-1} \circ \Phi_j^{-1} = T_{-1}$ . We will show that  $L_{-1} \circ \Phi_j \circ L_{-1}$  is a Fatou coordinate of  $\tilde{g}^{-1}$  on  $\tilde{S}_{j+1}$ . Indeed, we have

$$\begin{aligned} (L_{-1} \circ \Phi_j \circ L_{-1}) \circ Y_{j+1} \circ \tilde{g}^{-1} \circ Y_{j+1}^{-1} \circ (L_{-1} \circ \Phi_j \circ L_{-1})^{-1} \\ = L_{-1} \circ \Phi_j \circ Z_j \circ g^{-1} \circ Z_j^{-1} \circ \Phi_j^{-1} \circ L_{-1} \\ = L_{-1} \circ T_{-1} \circ L_{-1} = T_1. \end{aligned}$$

It follows that the transition functions  $\tilde{\Psi}_j$  of  $\tilde{g}^{-1}$  are given by, for  $j \neq 1, k$  odd,

$$\begin{aligned}\tilde{\Psi}_j &= \tilde{\Phi}_j \circ (\tilde{\Phi}_{j-\text{sgn}(j)})^{-1} \\ &= L_{-1} \circ \Phi_{j-1} \circ \Phi_{j-1-\text{sgn}(j-1)}^{-1} \circ L_{-1} \\ &= L_{-1} \circ \Psi_{j-1}^{-1} \circ L_{-1}.\end{aligned}$$

The other values of  $j$  are done similarly. Note however that for  $j = 1$ , the equation becomes  $\tilde{\Psi}_1 = L_{-1} \circ \Psi_{-1}^{-1} \circ L_{-1}$ , and that for  $j = k$ , we have  $\tilde{\Psi}_{-k} = L_{-1} \circ \Psi_k^{-1} \circ L_{-1}$ .  $\square$

## 6. SPACE OF ORBITS AND CLASSIFICATION UNDER ANALYTIC CONJUGACY

When two antiholomorphic parabolic germs are analytically conjugate, it is clear that their space of orbits are essentially the same. Our guiding principle is that *the space of orbits completely describes the dynamics of the germs*; when two germs have the same space of orbits, they should be analytically conjugate. A formal statement will be given in the form of the Classification Theorem 6.3.1 in Section 6.3.

**6.1. Description of the Space of Orbits.** The space of orbits in the holomorphic case is well known. It is briefly described in [5]. We will introduce the objects from the holomorphic case needed for the antiholomorphic description.

In each Fatou coordinate of  $f$  (and  $f \circ f$ ), we may choose a fundamental domain of  $g = f \circ f$  by taking any vertical strip  $B_j$  of width 1. We quotient by the action of  $T_1$ , obtaining the bi-infinite Écalle cylinders. Some orbits of  $g$  will appear in two consecutive cylinders. Since  $\Psi_j \circ T_1 = T_1 \circ \Psi_j$ , we identify together those orbits by identifying  $W_j$  with  $\Psi_j(W_j)$  (or  $\Psi_j^{-1}(W_j)$  depending on  $j$ ).

The universal covering  $E: \mathbb{C} \rightarrow \mathbb{C}^*$  given by  $w = E(W) = \exp(-2i\pi W)$  is a biholomorphism of each cylinder onto  $\mathbb{C}^*$ . This allows us to see the Écalle cylinders as Riemann spheres punctured at 0 and infinity  $\mathbb{S}_j^2 \setminus \{0, \infty\}$  (see Figure 14). We will define the horns maps using this universal covering.

**Definition 6.1.1.** The *horns maps*  $\psi_j$ , for  $j = 1, \dots, k$ , are defined by  $\psi_j = E \circ \Psi_j \circ E^{-1}$ , where  $\Psi_j$  is a transition function and

$$(29) \quad E(W) = \exp(-2i\pi W).$$

For positive  $j$  odd (resp.  $j$  even), the horn maps are defined on a punctured neighbourhood of the origin (resp. of infinity) with their






FIGURE 14. A fundamental domain obtained by  $Z_0^{-1}(B_0)$  in the  $z$ -coordinate (in gray) and the sphere it represents

image in a neighbourhood punctured of the origin (resp. of infinity). By the Riemann Removable Singularity Theorem, they extend to

$$\psi_j: \begin{cases} (\mathbb{C}, 0) \rightarrow (\mathbb{C}, 0), & \text{for } j > 0 \text{ odd;} \\ (\mathbb{C}, \infty) \rightarrow (\mathbb{C}, \infty), & \text{for } j > 0 \text{ even.} \end{cases}$$

To retrieve the  $\psi_j$  for  $j < 0$ , we use Equation (17) in the coordinate  $w = E(W)$

$$(30) \quad \psi_{-j} = L_{-1} \circ \tau \circ \psi_j \circ L_{-1} \circ \tau,$$

where  $L_{-1}(w) = -w$  and  $\tau(w) = \frac{1}{w}$ .

The space of orbits of  $g = f \circ f$  is described by the  $2k$  spheres with identifications at the origin or at infinity, as seen in Figure 15.




FIGURE 15. Space of orbits for  $f \circ f$  of codimension 4

*Remark 6.1.2.* By differentiating  $E \circ \Psi_j \circ E^{-1}$ , we obtain

$$(31) \quad \begin{aligned} (\psi_j)'(\infty) &= e^{\frac{2\pi^2 b}{k}} && \text{for } j \text{ odd;} \\ (\psi_j)'(0) &= e^{\frac{2\pi^2 b}{k}} && \text{for } j \text{ even;} \end{aligned}$$

so the product of those derivatives for  $j \in \{-k, \dots, -1\} \cup \{1, \dots, k\}$  is  $e^{4\pi^2 b}$ .

**6.2. The space of orbits of  $f$ .** The complex conjugation  $\Sigma$  becomes  $\tau(w) = \frac{1}{\bar{w}}$  on the spheres, where  $w \in \mathbb{S}_j^2$  and  $\tau(w) \in \mathbb{S}_{-j}^2$ . The translation  $T_{\frac{1}{2}}$  becomes  $L_{-1}(w) = -w$ . In the Fatou coordinates,  $f$  is  $\Sigma \circ T_{\frac{1}{2}}$ , so that on the spheres,  $f$  is  $L_{-1} \circ \tau(w) = -\frac{1}{\bar{w}}$ . To obtain the space of orbits of  $f$ , we identify  $w$  and  $L_{-1} \circ \tau(w)$  in the space of orbits of  $f \circ f$ , that is on the  $2k$  spheres above.

Let us first consider the case of codimension 1, so that we have two sectors  $S_0$  and  $S_1 = S_{-1}$  in the  $z$ -coordinate and two spheres  $\mathbb{S}_0^2$  and  $\mathbb{S}_1^2$ . Recall that for  $z \in S_0$ ,  $f^{-1}(z)$  is still in  $S_0$ . This means that  $L_{-1} \circ \tau$  acts on  $\mathbb{S}_0^2$ . It sends 0 to  $\infty$ , the northern hemisphere to the southern hemisphere, and the equator on itself. On the equator  $|w| = 1$ , we identify  $w$  to  $-w$ . The resulting surface is the real projective space  $\mathbb{RP}^2$ . This is also true for  $S_1$  and  $\mathbb{S}_1^2$ . The equators of both spheres play a special role; they each represent orbits along an invariant half-curve that each forms a “semi-axis” of reflection for  $f$ .


Therefore, in codimension 1, the space of orbits is two real projective spaces with one germ of holomorphic diffeomorphism

$$[\psi_1]: (\mathbb{RP}^2, [0]) \rightarrow (\mathbb{RP}^2, [0]),$$

where  $[\psi_1]$  is the equivalence class of  $\psi_1$  under the quotient of  $\mathbb{S}^2$  to  $\mathbb{RP}^2$  and  $[0]$  is equivalence class of the points  $\{0, \infty\}$  identified together. The class  $[\psi_1]$  is well-defined since  $\psi_1 \circ L_{-1} \circ \tau = \tau \circ L_{-1} \circ \psi_{-1}$ .

In codimension  $k > 1$ , the spheres  $\mathbb{S}_0$  and  $\mathbb{S}_k$  both quotient to a real projective space, but the other spheres are identified in pairs  $(\mathbb{S}_j, \mathbb{S}_{-j})$ , so that the quotient of the union of the two spheres is diffeomorphic to a sphere. The space of orbits is then described by two real projective spaces together with  $k - 1$  spheres and  $k$  equivalence classes of horn maps  $[\psi_j] = \{\psi_j, \psi_{-j}\}$ . The class  $[\psi_j]$  defines a germ at  $[0] = [\infty]$  on the quotiented spheres  $(\mathbb{S}_j, \mathbb{S}_{-j})$ , since the representatives satisfy  $\psi_j \circ L_{-1} \circ \tau = L_{-1} \circ \tau \circ \psi_{-j}$ .

On the two extreme projective spaces we have a distinguished curve given by the equator. The only changes of coordinates on  $\mathbb{D}$  preserving the equator and sending opposite points to opposite points are the linear maps  $L_c$  with  $|c| = 1$ . Hence the lines  $\Im W = y$  are invariant. This

FIGURE 16. The space of orbits of  $f$  in codimension 4

$y$  is the generalization of the *Écalle height* introduced in [4]. However, on the other spheres, there does not appear to be a quantity preserved by changes of coordinates.

The two projective spaces correspond to the orbits of two Fatou petals containing the formal symmetry axis of  $f$ . The equator of each projective space corresponds to a semi-axis of symmetry in each respective petal. These axes meet at the origin, but generally they cannot be extended into a real analytic curve. We will see in Theorem 7.3.1 exactly when they extend into a real analytic curve. This formal symmetry axis explains the existence of the *Écalle height* in the two projective spaces and why the *Écalle height* does not exist in the other spheres.

**6.3. Classification Under Analytic Conjugacy.** We can now state and prove the main theorem of this paper.

**Classification Theorem 6.3.1.** *For antiholomorphic parabolic germs of codimension  $k$  (and of positive type when  $k$  is even), we have:*

1. *The modulus of classification is a complete invariant of analytic classification under holomorphic conjugacy;*
2. *The moduli space is the set of  $(k, b, [\Psi_1, \dots, \Psi_k])$ , where the constant terms of  $\Psi_j$  satisfy (20), and  $[\Psi_1, \dots, \Psi_k]$  is the equivalence class under the Relation (21) (and (22) for  $k$  even).*

*Proof.* The proof is analogous to that in the holomorphic case, which can be found in [5] and [6].

1. Let  $f_\ell: (\mathbb{C}, 0) \rightarrow (\mathbb{C}, 0)$  be a germ of antiholomorphic diffeomorphism with a parabolic fixed point at the origin and let  $(k_\ell, b_\ell, [\Psi_{1,\ell}, \dots, \Psi_{k,\ell}])$  be its modulus of classification, for  $\ell = 1, 2$ . We can of course suppose that  $f_\ell$  is prenormalized.

Suppose first that  $f_2(u) = h \circ f_1 \circ h^{-1}(u)$  for some germ of analytic diffeomorphism  $u = h(z)$ . The germs  $f_1$  and  $f_2$  must have the same codimension and formal invariant, since they are topological and formal

invariants. For the analytic invariant, first let  $F_1$ ,  $F_2$  and  $H$  denote the expressions of  $f_1$ ,  $f_2$  and  $h$  in the time coordinate. If  $\Phi_j$  is a Fatou coordinate of  $F_2$  on  $U_j$ , then  $\Phi_j \circ H$  is a Fatou coordinate of  $F_1$ . It follows that they have the same transition functions.

Conversely, suppose  $f_1$  and  $f_2$  have the same modulus. We can choose a common normalized representative  $(\Psi_1, \dots, \Psi_k)$  for both classes and Fatou coordinates  $\Phi_{j,\ell}$  for  $f_\ell$  ( $\ell = 1, 2$ ), such that the  $\Psi_j$  are the transition maps for these Fatou coordinates. Let  $\varphi_{j,\ell} = Z_j^{-1} \circ \Phi_{j,\ell} \circ Z_j$ , for  $\ell = 1, 2$ , be the expression of Fatou coordinates in the  $z$ -coordinate. Then we define

$$h_j(z) = \varphi_{j,1}^{-1} \circ \varphi_{j,2}(z)$$

on  $S_j$ . It sends the orbits of  $f_2$  to those of  $f_1$  on  $S_j$ . On the intersection of two consecutive sectors  $S_j \cap S_{j+1}$ , we see that

$$h_{j+1} \circ h_j^{-1}(z) = id,$$

so the  $h_j$ 's agree on the intersection. We may define  $h$  on  $\bigcup_j S_j$  by  $h(z) = h_j(z)$  if  $z \in S_j$ . It sends a punctured neighbourhood of the origin into another punctured neighbourhood of the origin, so by the Riemann Bounded Extension Theorem,  $h$  extends to a holomorphic diffeomorphism of a neighbourhood of the origin.

Finally, we see that  $h \circ f_2 = f_1 \circ h$  since it sends the orbits of  $f_2$  to the orbits of  $f_1$  on a whole neighbourhood of the origin.

2. The proof is in two steps. First we construct an abstract Riemann surface  $S$  on which  $\Sigma \circ T_{\frac{1}{2}}$  is well defined and we prove that this surface has the conformal type of a punctured disk. Then we prove that  $\Sigma \circ T_{\frac{1}{2}}$  is the germ we are looking for on the disk.

Consider a triple  $(k, b, [\Psi_1, \dots, \Psi_k])$ . We must find a parabolic germ of antiholomorphic diffeomorphism with this modulus of classification.

Let  $(\Psi_1, \dots, \Psi_k)$  be a representative of the analytic invariant and let  $\Psi_{-1}, \dots, \Psi_{-k}$  be the other transition functions obtained from (17). Let  $\sigma \circ v^{\frac{1}{2}}$  be the normal form of codimension  $k$  with formal invariant  $b$ . Let  $U_j$  be the charts in the time coordinate of  $v$ .

We consider the transition functions defined on those charts. More precisely,  $\Psi_1$  is defined on a domain of  $U_0$  containing an upper-half plane with its image in  $U_1$ ;  $\Psi_2$  is defined on a domain of  $U_2$  containing a lower half-plane with its image in  $U_1$ , and so on. We define the Riemann surface  $S$  by

$$S = \bigsqcup_{j=-k}^k U_j / \sim$$

where  $\sim$  identifies  $W_j \in U_j$  with its image by  $\Psi_j$  or  $\Psi_{j-1}$  (depending on  $j$ ). As it is done in [6], we can build a smooth quasi-conformal mapping  $P: S \rightarrow \mathbb{C}^*$ , and from the Ahlfors-Bers Theorem (see [1]), find a diffeomorphism  $Q: D \rightarrow (\mathbb{C}, 0)$ , where  $D = P(S) \cup \{0\}$ , so that the composition  $H = Q \circ P: S \rightarrow (\mathbb{C}, 0)$  is holomorphic. In fact,  $H$  is a biholomorphism of  $S$  onto some punctured disk of the origin.

By (15), we know that  $\Sigma \circ T_{\frac{1}{2}}$  is well defined on  $S$ . The map  $f = H \circ \Sigma \circ T_{\frac{1}{2}} \circ H^{-1}$  extends to the origin by  $f(0) = 0$ , since it is bounded around 0 (we can apply the Riemann Removable Singularity Theorem to  $f \circ \sigma$  to see this). Lastly,  $T_1$  is also well-defined on  $S$ , so we set  $g = H \circ T_1 \circ H^{-1}$ . Since  $T_1 = (\Sigma \circ T_{\frac{1}{2}}) \circ (\Sigma \circ T_{\frac{1}{2}})$ , it follows that  $g = f \circ f$ , and by the chain rule,  $|\frac{\partial f}{\partial \bar{z}}|^2 = g'(0)$ . By the holomorphic case, we know  $g$  is a holomorphic parabolic germ of codimension  $k$ , so it follows that  $f$  is parabolic and of codimension  $k$ . The formal invariant of  $f$  is  $b$ , since it is determined by (19).

It remains only to prove that  $f$  is of positive type. The formal symmetry axis of  $f$  is the real line, since the  $\Psi_1$  and  $\Psi_{-1}$  are defined on  $U_0$  and  $\Psi_j \circ \Sigma \circ T_{\frac{1}{2}} = \Sigma \circ T_{\frac{1}{2}} \circ \Psi_{-j}$ . Moreover, the petal  $S_0$  of  $f$  is repulsive, since  $H|_{U_0} \circ Z_0: U_0 \rightarrow U_0$  is a Fatou coordinate of  $f$  and  $U_0$  contains a half-plane  $\{\Re Z < R\}$ , i.e.  $S_0$  contains all the backward iterates of  $f$ . Therefore,  $f$  is of positive type.

We conclude  $f$  is a germ with modulus  $(k, b, [\Psi_1, \dots, \Psi_k])$ .  $\square$

## 7. APPLICATIONS OF THE CLASSIFICATION THEOREM

Here we will solve Questions 1 to 5 of the introduction and other related questions.

**7.1. Embedding in a Flow or the Complex Conjugate of a Flow.** If a holomorphic germ  $g$  is conjugate to the normal form, that is  $g = h \circ v^1 \circ h^{-1}$ , then it is embedded in the family  $g_t = h \circ v^t \circ h^{-1}$ . Similarly, we will say that  $f$  is *embeddable* if it is embedded in the family  $f_t := h \circ \sigma \circ v^t \circ h^{-1}$ . When is an antiholomorphic parabolic germ embeddable? This corresponds to Question 3. The answer is read in the modulus of classification.

**Theorem 7.1.1** (Embedding in a flow). *An antiholomorphic parabolic germ  $f$  is analytically conjugate to its normal form (i.e. embeddable) if and only if the transition functions of  $f$  are translations.*

*Proof.* The transition functions of the normal form are translations since the time charts  $Z_j$  are Fatou coordinates. Therefore, it follows from the Classification Theorem 6.3.1.  $\square$

In Proposition 2.0.2, we proved that  $f$  is always formally conjugate to the sum of a formal germ with real coefficients. We ask the related question.

*Question 7.1.2.* When is a parabolic antiholomorphic germ analytically conjugate to a series with real coefficients?

Of course, this is the case for any embeddable germ. But we will show in Section 7.3 that the embeddable germs form a subset of infinite codimension in the set of antiholomorphic parabolic germs conjugate to a germ with real coefficients. We will answer this question in Section 7.3. We first tackle Question 1.

**7.2. Antiholomorphic  $n$ -th Root Problem.** Question 1 and its restatement Question 3.0.5, as well as Question 3.0.6 are concerned with the existence of antiholomorphic roots and their uniqueness.

**Theorem 7.2.1** (Antiholomorphic Root Extraction Problem). *Let  $g$  be a holomorphic parabolic germ and let  $(k, b, [\Psi_{-k}, \dots, \Psi_{-1}, \Psi_1, \dots, \Psi_k])$  be its Écalle-Voronin modulus (see Section 5.1), then*

1.  $g$  has  $k$  one-parameter families of formal antiholomorphic  $n$ -th roots, one for each of the  $k$  formal axes of reflection;
2.  $g$  has an antiholomorphic  $n$ -th root ( $n$  even) tangent to the formal symmetry axis  $e^{\frac{i\pi\ell}{k}}\mathbb{R}$  if and only if there exists a representative  $(\Psi_{-k}, \dots, \Psi_{-1}, \Psi_1, \dots, \Psi_k)$  of the equivalence relation (23) which has a symmetry property with respect to the formal symmetry axis  $e^{\frac{i\pi\ell}{k}}\mathbb{R}$ , namely

$$(32) \quad \Psi_j \circ \Sigma \circ T_{\frac{1}{n}} = \Sigma \circ T_{\frac{1}{n}} \circ \Psi_{s_\ell(j)},$$

where  $s_\ell$  is the reflection of indices with respect to  $e^{\frac{i\pi\ell}{k}}\mathbb{R}$  (see Definition 5.1.1);

3. If  $g$  is not analytically conjugate to its normal form, then
  - i) Each family has at most one convergent root, so  $g$  has at most  $k$  distinct antiholomorphic  $n$ -th roots  $f_j$ ;
  - ii) If  $g$  has  $m$  distinct antiholomorphic roots  $f_{\ell_1}, \dots, f_{\ell_m}$  with distinct linear parts  $e^{\frac{2i\pi\ell_j}{k}}\bar{z}$ ,  $j = 1, \dots, m$ , then the modulus of  $g$  has  $\gcd(\ell_m - \ell_1, \dots, \ell_m - \ell_{m-1}, k)$  independent transition functions.

*Proof.* 1. That  $g$  has a 1-parameter family of formal antiholomorphic  $n$ -th root on each symmetry axis is a consequence of the fact that  $g$  is formally conjugate to the normal form  $v^1$ , and Proposition 3.0.4.

2. We first prove this for the case  $k$  odd. Let

$$\Psi = (\Psi_{-k}, \dots, \Psi_{-1}, \Psi_1, \dots, \Psi_k)$$

be a representative of the analytic invariant of  $g$ .

First, we suppose that the representative satisfies (32). To realize an antiholomorphic root tangent to the symmetry axis  $e^{\frac{2i\pi\ell}{k}}\mathbb{R}$ , we rotate the coordinate by  $R_\ell(z) = e^{-\frac{2i\pi\ell}{k}}z$  so that this axis is on the real line and the dynamics of  $R_\ell \circ g \circ R_\ell^{-1}$  on the side of the positive real axis is repulsive. Let  $g_\ell = R_\ell \circ g \circ R_\ell^{-1}$  and let  $r$  correspond to the permutation of indices defined by  $R_\ell$  (see Definition 5.1.1). The representative is permuted into  $(\Psi_{r(-k)}, \dots, \Psi_{r(-1)}, \Psi_{r(1)}, \dots, \Psi_{r(k)})$ . Also, we have that  $r^{-1}s_\ell r = s_0$ , where  $s_0(j) = -j$  and  $s_\ell$  is the reflection of indices induced by  $\sigma_\ell$ . Equation (32) becomes

$$\Psi_{r(j)} \circ \Sigma \circ T_{\frac{1}{2}} = \Sigma \circ T_{\frac{1}{2}} \circ \Psi_{r(-j)}.$$

Now we may repeat the proof of part two of Theorem 6.3.1 to obtain an antiholomorphic germ  $f_\ell: (\mathbb{C}, 0) \rightarrow (\mathbb{C}, 0)$  with analytic invariant  $[\Psi_{r(1)}, \dots, \Psi_{r(k)}]$  such that  $f_\ell$  is  $\Sigma \circ T_{\frac{1}{n}}$  in each Fatou coordinate of  $g_\ell$ . Since  $(\Sigma \circ T_{\frac{1}{n}})^n = T_1$  in the Fatou coordinates, it follows that  $f_\ell^{\circ n} = g_\ell$ .

Conversely, suppose  $g$  has an  $n$ -th root  $f_\ell$  tangent to the reflection axis  $e^{\frac{2i\pi\ell}{k}}\mathbb{R}$ . We rotate the coordinate by  $R_\ell$ . Let  $f = R_\ell \circ f_\ell \circ R_\ell^{-1}$ . Now in every Fatou coordinate of  $g_\ell$ ,  $f$  has the form  $\Sigma \circ T_{\frac{1}{n}+iy}$ , by Proposition 3.0.4. By changing Fatou coordinates, we may obtain  $\Sigma \circ T_{\frac{1}{n}}$ . Those Fatou coordinates give us a representative  $(\Psi'_{-k}, \dots, \Psi'_{-1}, \Psi'_1, \dots, \Psi'_k)$  that is equivalent to  $\Psi$  under the Relations (23) and (24) and that satisfies

$$\Psi'_j \circ \Sigma \circ T_{\frac{1}{2}} = \Sigma \circ T_{\frac{1}{2}} \circ \Psi'_{-j}.$$

It follows that  $(\Psi'_{r^{-1}(-k)}, \dots, \Psi'_{r^{-1}(-1)}, \Psi'_{r^{-1}(1)}, \dots, \Psi'_{r^{-1}(k)})$  is a representative equivalent to  $\Psi$  under Relation (23) which satisfies (32), using the fact that  $(r^{-1}s_\ell r)(j) = -j$ .

In the case  $k$  even, we may have antiholomorphic roots of positive and negative type. Those of positive type are tangent to an axis  $e^{\frac{i\pi\ell}{k}}\mathbb{R}$  with  $\ell$  even and are done as before. Those of negative type are tangent to an axis  $e^{\frac{i\pi\ell}{k}}\mathbb{R}$  with  $\ell$  odd. For such a root  $f_\ell$ ,  $f_\ell^{-1}$  will be a root of positive type of  $g^{-1}$ . By Proposition 5.1.2, the modulus of  $g^{-1}$  is

$$(k, -b, [\tilde{\Psi}_{-k}, \dots, \tilde{\Psi}_{-1}, \tilde{\Psi}_1, \dots, \tilde{\Psi}_k]),$$

where  $\tilde{\Psi}_j = L_{-1} \circ \Psi_{r_1^{-1}(j)}^{-1} \circ L_{-1}$  and  $r_1$  is the rotation of indices induced by  $z \mapsto e^{\frac{i\pi}{k}}z$ . By the previous two paragraphs,  $g^{-1}$  has an  $n$ -th antiholomorphic root of positive type tangent to  $e^{\frac{i\pi\ell}{k}}\mathbb{R}$  if and only if there

exists a representative  $(\tilde{\Psi}_{-k}, \dots, \tilde{\Psi}_{-1}, \tilde{\Psi}_1, \dots, \tilde{\Psi}_k)$  such that

$$\tilde{\Psi}_j \circ \Sigma \circ T_{\frac{1}{n}} = \Sigma \circ T_{\frac{1}{n}} \circ \tilde{\Psi}_{s_\ell(j)},$$

By simplifying the  $L_{-1}$  and taking the inverse on both sides, we obtain (32).

3. *i)* Suppose  $g$  has two roots  $f_1$  and  $f_2$  from the same family. In particular, they have the same linear term, so we may suppose they are tangent to  $\sigma$ , modulo conjugating  $g$  by a rotation of order  $k$ . In the Fatou coordinates, they take the form  $\Sigma \circ T_{\frac{1}{2}+iy_j}$ , for  $j = 1, 2$ , by Proposition 3.0.4. Since  $\Sigma \circ T_{\frac{1}{2}+iy_j}$  satisfies (15) for  $j = 1, 2$ , and by combining with (18), we see that either  $y_1 = y_2$  or the  $\Psi_j$ 's are translations.

*ii)* Lastly, consider the dihedral group  $D_{2k}$  with its action on the regular  $k$ -gon. Recall that we can divide a regular  $k$ -gon by its  $k$  symmetry axes to form  $2k$  sectors, which we associate to the transition functions (see Section 5.1 and Figure 13).

Let  $H = \langle s_1, \dots, s_m \rangle$  be the subgroup of  $D_{2k}$  generated by the reflections along the symmetry axes of  $f_{\ell_1}, \dots, f_{\ell_m}$ . Each element of  $H$  acts on the modulus by introducing relations of the type

$$\Sigma \circ T_{\frac{1}{2}} \circ \Psi_j = \Psi_{s_\ell(j)} \circ \Sigma \circ T_{\frac{1}{2}},$$

thus reducing the number of independent  $\Psi_j$  (see Figure 17). The orbit by  $H$  of a sector represents the transition functions tied together, therefore the number of independent transition functions corresponds to  $|D_{2k} : H|$ . To compute this, we first observe that  $H$  must itself be a dihedral group, so that  $H = D_{2j}$  for some  $j$  (see [3]). In fact,  $j$  is given by

$$j = \frac{k}{\gcd(\ell_m - \ell_1, \dots, \ell_m - \ell_{m-1}, k)},$$

since any rotation  $s_m s_p$  has order  $k / \gcd(\ell_m - \ell_p, k)$ . It follows that  $|D_{2k} : H| = \gcd(\ell_m - \ell_1, \dots, \ell_m - \ell_{m-1}, k)$ .  $\square$

**Theorem 7.2.2.** *Let  $f: (\mathbb{C}, 0) \rightarrow (\mathbb{C}, 0)$  be an antiholomorphic parabolic germ of codimension  $k$ , and of positive type when  $k$  is even. Let  $(k, b, [\Psi_1, \dots, \Psi_k])$  be the modulus of  $f$  and let  $n \geq 3$  be an odd number. Then  $f$  has a single antiholomorphic formal  $n$ -th root ( $n$  odd). Moreover,  $f$  has an antiholomorphic  $n$ -root ( $n$  odd) if and only if*

$$(33) \quad \Psi_j \circ T_{\frac{1}{n}} = T_{\frac{1}{n}} \circ \Psi_j.$$

*Proof.* That  $f$  has a unique formal  $n$ -th root follows from Proposition 3.0.4.






FIGURE 17. The orbit of one sector by the subgroup  $H$  generated by two reflections (bold axes) in codimension 12.

For the second part,  $f$  has an antiholomorphic  $n$ -th root if and only if the transition functions satisfy  $\Psi_j \circ \Sigma \circ T_{\frac{1}{2n}} = \Sigma \circ T_{\frac{1}{2n}} \circ \Psi_j$ , since we may realize  $\Sigma \circ T_{\frac{1}{2n}}$  on the Riemann surface of  $f \circ \bar{f}$ . We combine this last equation with (15) to obtain  $\Psi_j \circ T_{\frac{1}{2n} - \frac{1}{2}} = T_{\frac{1}{2n} - \frac{1}{2}} \circ \Psi_j$ . To conclude, we note that  $\gcd(1-n, 2n) = 2$  because  $n$  is odd, so there exists  $p, q \in \mathbb{Z}$  such that  $(1-n)p - 2nq = 2$ . In other words, we have  $\frac{1}{n} = \frac{(1-n)}{2n}p - q$ . Since  $\Psi_j$  commutes with  $T_{\frac{(1-n)}{2n}p}$  and with  $T_q$  (because  $q$  is an integer), it follows that  $\Psi_j$  commutes with  $T_{\frac{1}{n}}$ .  $\square$

**Corollary 7.2.3.**  *$f$  has an antiholomorphic  $n$ -th root for  $n$  odd if and only if  $f$  is the square root of  $g$  and  $g$  has a holomorphic  $n$ -th root, with  $n$  odd.*

*Proof.* Equation (33) is independent of the representative and it is equivalent to a holomorphic parabolic germ having a holomorphic  $n$ -th root (see [5]).

Suppose  $g = f \circ f$  and  $g$  has a holomorphic  $n$ -th root. Then the modulus of  $g$  and  $f$  satisfies (33), so that  $f$  has an antiholomorphic  $n$ -th root.

The converse is direct.  $\square$

**7.3. Germs with an Invariant Real Analytic Curve.** An antiholomorphic germ with real coefficients preserves the real axis. Any

germ  $f$  analytically conjugate to the latter will preserve a real analytic curve; it is a property of the equivalence class of  $f$ . Therefore, Question 7.1.2 is equivalent to asking when does an antiholomorphic parabolic germ  $f$  preserve a real analytic curve.

Like the embedding problem, to preserve a germ of real axis is a condition of codimension infinity, but it is a “smaller” infinity, i.e. not every transition functions needs to be a translation, but we will see that there are infinitely many conditions of the form  $c_{n,j} = \overline{c_{-n,-j}}$  for the Fourier coefficients in (18).

**Theorem 7.3.1.** *Let  $f: (\mathbb{C}, 0) \rightarrow (\mathbb{C}, 0)$  be an antiholomorphic parabolic germ and  $(k, b, [\Psi_1, \dots, \Psi_k])$  be its modulus of classification. The following statements are equivalent*

1.  $f$  preserves a real analytic curve at the origin;
2.  $f$  is analytically conjugate to a germ with real coefficients;
3. each representative  $(\Psi_1, \dots, \Psi_k)$  satisfies  $\Psi_j \circ T_{\frac{1}{2}} = T_{\frac{1}{2}} \circ \Psi_j$ ;
4. each representative  $(\Psi_1, \dots, \Psi_k)$  satisfies  $\Psi_j \circ \Sigma = \Sigma \circ \Psi_{-j}$ , where  $\Psi_{-j}$  is defined by (17).

*Proof.* 1.  $\Rightarrow$  2. Let  $\gamma$  be the germ of real analytic parametrization of the invariant curve of  $f$ . We can extend  $\gamma$  on a disk around the origin. Then  $\gamma^{-1} \circ f \circ \gamma$  fixes a germ of the real axis, so its power series has real coefficients.

2.  $\Rightarrow$  3. We can of course suppose that the power series of  $f$  has real coefficients, so that  $\sigma \circ f = f \circ \sigma$ . It follows that  $\sigma \circ f$  is a holomorphic square root of  $g = f \circ f$ . Therefore, we have  $\Psi_j \circ T_{\frac{1}{2}} = T_{\frac{1}{2}} \circ \Psi_j$ .

3.  $\Leftrightarrow$  4. It follows from equation (15).

4.  $\Rightarrow$  1. This is the harder part of the proof. The hypothesis implies that  $\Sigma$  is well defined on the Riemann surface  $S$  constructed in the proof of part 2 of the Classification Theorem 6.3.1. Let  $z = H(W)$  be the coordinate given by  $H: S \rightarrow (\mathbb{C}, 0) \setminus \{0\}$ , the biholomorphism of  $S$  to a punctured neighbourhood of the origin. Let  $\sigma'$  and  $f'$  be the expression of  $\Sigma$  and  $\Sigma \circ T_{\frac{1}{2}}$  in this coordinate (note that  $f$  and  $f'$  are analytically conjugate). Since on  $S$ ,  $\Sigma$  and  $\Sigma \circ T_{\frac{1}{2}}$  commute, it follows that  $\sigma'$  and  $f'$  commute also. So it remains only to show that  $\sigma'$  preserves a germ of real analytic curve at the origin.

We can extend  $\sigma'$  by  $\sigma'(0) = 0$  by the Riemann Removable Singularity Theorem (this is true for antiholomorphic functions, since we simply apply it to  $\sigma' \circ \sigma$ , where  $\sigma(z) = \bar{z}$ ). In the charts  $U_0$  and  $U_k$ ,  $\Sigma$  fixes the real axis. These two curves carry in the  $z$ -coordinate and meet at the origin to form a continuous curve  $\gamma$  fixed by  $\sigma'$ . In fact,  $\gamma$  is a  $C^1$

curve, as  $H$  can be extended to a  $C^1$  diffeomorphism  $\tilde{H}: \tilde{S} \rightarrow (\mathbb{C}, 0)$ , where  $\tilde{S}$  is obtained from  $S$  with the point  $\infty = H^{-1}(0)$  added.

This curve divides a small disk  $D(0, \delta)$  in two connected components  $A$  and  $B$ . By the Riemann Mapping Theorem, there exists a biholomorphism  $\varphi$  of  $A$  to the upper half-plane that sends continuously the boundary of  $A$  on the real line, see Figure 18. The image of  $\gamma$  corresponds to an interval  $[a, b]$ . We can extend  $\varphi$  to  $A \cup \gamma \cup B$  by

$$\tilde{\varphi}(z) = \begin{cases} \varphi(z) & \text{if } z \in A \cup \gamma; \\ \sigma \circ \varphi \circ \sigma'(z) & \text{if } z \in B. \end{cases}$$

This is holomorphic on  $B$ , since it is the composition of a holomorphic map with two antiholomorphic maps, and it is continuous on  $A \cup \gamma \cup B$ .




FIGURE 18. Mapping  $\varphi$  extended on  $A \cup \gamma \cup B$

The argument to prove that  $\tilde{\varphi}$  is holomorphic is similar to that of the Schwarz Reflection Principle. The idea is to show that the integral of  $\tilde{\varphi}$  along any triangle in  $A \cup \gamma \cup B$  vanishes, and it will follow from Morera's theorem. If a simple closed curve is in  $A \cup \gamma$  or  $B \cup \gamma$ , then it follows from Cauchy's theorem (we may take a limit of closed curves in  $A$  or  $B$  converging to the initial one). Then, we can divide any triangle in  $A \cup \gamma \cup B$  along  $\gamma$  to obtain a finite number of closed curves in  $A \cup \gamma$  and in  $B \cup \gamma$ , as in Figure 19. Thus,  $\tilde{\varphi}|_{(a,b)}^{-1}$  is a real analytic parametrization of  $\gamma$ .  $\square$




FIGURE 19. Triangle divided in two closed curves

When  $f$  is conjugate to a germ with real coefficients, then so is the holomorphic germ  $g = f \circ f$ . However, it is not true that every holomorphic germ analytically conjugate to a germ with real coefficients must have a germ  $f$  such that  $g = f \circ f$ , as the next theorem will show (see also Proposition 7.3.3). Even though this is not a property directly linked to the antiholomorphic parabolic germs, we will still prove the following necessary and sufficient condition for a holomorphic parabolic germ to preserve a real analytic curve.

**Theorem 7.3.2.** *Let  $g: (\mathbb{C}, 0) \rightarrow (\mathbb{C}, 0)$  be a parabolic holomorphic germ and let  $(k, b, [\Psi_{-k}, \dots, \Psi_{-1}, \Psi_1, \dots, \Psi_k])$  be its Écalle-Voronin modulus (see Section 5.1). The following statements are equivalent*

1.  $g$  preserves a germ of real analytic curve at the origin;
2.  $g$  is analytically conjugate to a germ with real coefficients;
3. For each representative  $(\Psi_{-k}, \dots, \Psi_{-1}, \Psi_1, \dots, \Psi_k)$ , there exists  $y \in \mathbb{R}$  such that the transition functions satisfy  $\Sigma \circ T_{iy} \circ \Psi_j = \Psi_{-j} \circ \Sigma \circ T_{iy}$ ;
4. There exists a representative  $(\Psi_{-k}, \dots, \Psi_{-1}, \Psi_1, \dots, \Psi_k)$  such that  $\Psi_j \circ \Sigma = \Sigma \circ \Psi_{-j}$ .

*Proof.* 1.  $\Rightarrow$  2. and 4.  $\Rightarrow$  1. are the same as in the previous theorem.

2.  $\Rightarrow$  3. Suppose  $g$  is in a coordinate such that  $g = \sigma \circ g \circ \sigma$ . Let  $\Phi_j$  be a Fatou coordinate on  $U_j$  for  $-k \leq j \leq k$ , where  $\Phi_{-k} = \Phi_k$ . Then  $\Sigma \circ \Phi_{-j} \circ \Sigma$  is also a Fatou coordinate of  $g$  on  $U_j$ . By the uniqueness of the Fatou coordinate (for the holomorphic case), there exists a constant  $C \in \mathbb{C}$  such that

$$\Sigma \circ \Phi_{-j} \circ \Sigma \circ \Phi_j^{-1} = T_C$$

for all  $j$ . In particular, for  $j = 0$ , by taking the inverse and conjugating both sides by  $\Sigma$  of the previous equation, we obtain  $T_C = T_{-\bar{C}}$ , so  $C$  must be pure imaginary, say  $iy$  with  $y \in \mathbb{R}$ . For  $j > 0$  and odd, we conclude with

$$\Psi_j = \Phi_j \circ \Phi_{j-1}^{-1} = T_{-iy} \circ \Sigma \circ \Phi_{-j} \circ \Phi_{-j+1}^{-1} \circ \Sigma \circ T_{iy} = T_{-iy} \circ \Sigma \circ \Psi_{-j} \circ \Sigma \circ T_{iy}.$$

The other values of  $j$  are done similarly.

3.  $\Leftrightarrow$  4. It follows from the fact that for any two representatives  $(\Psi_{-k}, \dots, \Psi_{-1}, \Psi_1, \dots, \Psi_k)$  and  $(\Psi'_{-k}, \dots, \Psi'_{-1}, \Psi'_1, \dots, \Psi'_k)$ , there exists  $C \in \mathbb{C}$  such that  $\Psi_j \circ T_C = T_C \circ \Psi'_j$ . We choose  $C = -\frac{iy}{2}$  to get a representative satisfying  $\Psi'_j \circ \Sigma = \Sigma \circ \Psi'_{-j}$ .  $\square$

**Proposition 7.3.3.** *There exists  $g$  with real coefficients that has no antiholomorphic square root (see section 7.2).*

*Proof.* The holomorphic germ realized by the Écalle-Voronin modulus  $(1, 0, [W + e^{2i\pi W}, W + e^{-2i\pi W}])$  has no antiholomorphic square root, but preserves a germ of real analytic curve.  $\square$

**7.4. Centralizer in the Group of Holomorphic and Antiholomorphic Germs.** Let  $\text{Diff}_1(0, \mathbb{C})$  (resp.  $\overline{\text{Diff}_1(0, \mathbb{C})}$ ) be the set of germs of holomorphic (resp. antiholomorphic) diffeomorphisms with a fixed point at the origin with multiplier 1. Note in particular that an antiholomorphic germ must be tangent to  $\sigma$ . We set

$$\text{Diff}_1(0, \mathbb{C}, \overline{\mathbb{C}}) = \text{Diff}_1(0, \mathbb{C}) \cup \overline{\text{Diff}_1(0, \mathbb{C})}.$$

It forms a group with  $\text{Diff}_1(0, \mathbb{C})$  as a subgroup. Next, let  $\mathcal{A}_{k,b} \subset \text{Diff}_1(0, \mathbb{C})$  (resp.  $\overline{\mathcal{A}_{k,b}} \subset \overline{\text{Diff}_1(0, \mathbb{C})}$  when  $b$  is real) be the set of germs of holomorphic (resp. antiholomorphic) diffeomorphisms with a parabolic fixed point of codimension  $k$  at the origin and with formal invariant  $b$ .

We will study the centralizer of  $g \in \mathcal{A}_{k,b}$  and of  $f \in \overline{\mathcal{A}_{k,b}}$  in  $\text{Diff}_1(0, \mathbb{C}, \overline{\mathbb{C}})$ . Let us note the following: if  $g \in \mathcal{A}_{k,b}$  commutes with  $f \in \overline{\text{Diff}_1(0, \mathbb{C})}$ , then the formal invariant  $b$  of  $g$  is automatically real. In fact, we have that  $g$  and  $\sigma \circ g \circ \sigma$  are analytically conjugate by means of  $(\sigma \circ f)$ . So we are interested only in  $b$  real, since if  $b$  is not real, the centralizer of  $g$  in  $\text{Diff}_1(0, \mathbb{C}, \overline{\mathbb{C}})$  is the same as the centralizer of  $g$  in  $\text{Diff}(0, \mathbb{C})$ , which is already known (see [5]).

**Lemma 7.4.1.** *Let  $g \in \mathcal{A}_{k,b}$  and let  $m \in \text{Diff}_1(0, \mathbb{C}, \overline{\mathbb{C}})$  be a germ that commutes with  $g \in \mathcal{A}_{k,b}$ . Then either  $m$  is the identity, or analytically conjugate to  $\sigma$ , or  $m \in \mathcal{A}_{k,b} \cup \overline{\mathcal{A}_{k,b}}$ . Moreover, in the Fatou coordinates of  $g$ ,  $m$  will be of the form*

$$(34) \quad T_C \quad \text{or} \quad \Sigma \circ T_C \quad \text{for } C \in \mathbb{C}.$$

*Proof.* If  $m$  is not the identity or analytically conjugate to  $\sigma$ , then  $m$  is parabolic. It must have codimension  $k$ , since it maps the orbits of  $g$  on the orbits of  $g$  and the Fatou petals of  $g$  on the Fatou petals of  $g$ . To see that  $m$  has the same formal invariant, we compare degree by degree the power series on both sides of the equation  $m \circ g(z) = g \circ m(z)$ .

To see  $m$  has one of the forms of (34), the first part of the proof of Proposition 4.2.2 applies almost identically to  $m$ .  $\square$

The obvious germs in the centralizer of  $f$  or  $g$  are the iterates, the roots, and the iterates of the roots, which we define below as the *fractional iterates*. We prove in Theorem 7.4.3 and 7.4.4 that these are all the possible elements of the centralizer, provided that  $f$  or  $g$  are not conjugate to their respective normal form.

**Definition 7.4.2.** Let  $f \in \overline{\mathcal{A}_{k,b}}$  (resp.  $g \in \mathcal{A}_{k,b}$ ). We say that  $m \in \text{Diff}_1(0, \mathbb{C}, \overline{\mathbb{C}})$  is a *fractional iterate of order  $p$*  of  $f$  (resp.  $g$ ) if there exists  $q \in \mathbb{Z}$  such that  $\gcd(p, q) = 1$  and  $m^{op} = f^{oq}$  (resp.  $m^{op} = g^{oq}$ ).

Note that a fractional iterate of order one is just an iterate of  $f$  or of  $g$ .

**Theorem 7.4.3.** *Let  $g \in \mathcal{A}_{k,b}$  with  $b \in \mathbb{R}$ . Then we have one of the following cases:*

1.  *$g$  is embeddable, i.e.  $g = h \circ v^1 \circ h^{-1}$ . Let  $g_t = h \circ v^t \circ h^{-1}$ . Then the centralizer of  $g$  is*

$$Z_g = \{g_t \mid t \in \mathbb{C}\} \cup \{h \circ \sigma \circ v^t \circ h^{-1} \mid t \in \mathbb{C}\}.$$

2.  *$g$  is not embeddable, then  $Z_g$  contains only holomorphic and antiholomorphic fractional iterates of  $g$  and Schwarz reflections. More precisely, there exists  $p \in \mathbb{N} \setminus \{0\}$  such that the centralizer of  $g$  is one of the following:*

- $Z_g = \bigcup_{d|p} \{\text{fractional iterates of order } d \text{ of } g\},$
- $Z_g = \bigcup_{d|p} \{\text{hol. fractional iterates of order } d \text{ of } g\} \cup \bigcup_{\substack{d|2p \\ d \text{ even}}} \{\text{antihol. fractional iterates of order } d \text{ of } g\},$
- $Z_g = \bigcup_{d|p} \{\text{hol. fractional iterates of order } d \text{ of } g\} \cup \bigcup_{\substack{d|p \\ d \text{ even}}} \{\text{antihol. fractional iterates of order } d \text{ of } g\} \cup \{\text{Schwarz reflection tangent to } \sigma\}.$

*The centralizer includes a Schwarz reflection tangent to  $\sigma$  if and only if  $g$  is analytically conjugate to a germ with real coefficients (see Theorem 7.3.2).*

*Proof.* 1. Since  $g$  is embeddable, each of its transition functions is a translation, so any  $T_t$  will commute with them. Each  $T_t$  represents a germ analytically conjugate to  $g^t$  for some  $t \in \mathbb{C}$ . In the case  $m$  is antiholomorphic,  $\Sigma \circ T_t$  is compatible with the transition functions with no restriction on  $t$ . It corresponds to  $h \circ \sigma \circ v^t \circ h^{-1}$  for  $t \in \mathbb{C}$ .

2. Since  $g$  is not embeddable, one of the transition functions is not a translation. Suppose  $g_1 \in \text{Diff}_1(0, \mathbb{C}, \overline{\mathbb{C}})$  is holomorphic and commutes with  $g$ . By Lemma 7.4.1,  $g_1$  becomes  $T_t$  in the Fatou coordinates for some  $t \in \mathbb{C}$ . Then  $T_t$  must commute with the transition functions.

This will only happen if the transition functions expand in the form

$$(35) \quad \Psi_j(W) = W + (-1)^j \frac{i\pi b}{k} + \sum_{\ell \in \mathbb{Z}^*} c_{\ell p, j} e^{2i\pi \ell p W}$$

and  $t = \frac{a}{p}$  for some  $a, p \in \mathbb{N}$ . This corresponds to  $g_1$  being a holomorphic fractional iterate of order  $p$  of  $g$ . Let  $p$  denote the maximal order of holomorphic roots of  $g$ .

Suppose  $f_1 \in \text{Diff}_1(0, \mathbb{C}, \overline{\mathbb{C}})$  is antiholomorphic and commutes with  $g$ . We can suppose  $f_1$  is not a Schwarz reflection, since this is covered in Theorem 7.3.2. Then by Lemma 7.4.1,  $f_1$  is  $\Sigma \circ T_t$  in the Fatou coordinates and it must be compatible with the transition functions. Therefore,  $T_{2\Re t}$  commutes with the  $\Psi_j$ 's, and by the previous case, we have  $2\Re t = \frac{a}{r}$ , with  $\gcd(a, r) = 1$  and  $r \mid p$ . We have  $\Sigma \circ T_t \circ \Psi_j = \Psi_{-j} \circ \Sigma \circ T_t$  for some representative of the modulus if and only if

$$(36) \quad \overline{c_{n,j}} = e^{-i\pi a n/r} e^{2n\pi y} c_{-n, -j}$$

and  $t = \frac{a}{2r} + iy$  ( $y \in \mathbb{R}$ ), where  $c_{n,j}$  are the Fourier coefficients of  $\Psi_j$  from (18). Of course, generically there exists no  $y$  that satisfy (36), so there are no antiholomorphic germs that commute with  $g$  in the generic case. When such a  $y$  exists, by changing the  $\Psi_j$ 's by  $T_{-\frac{iy}{2}} \circ \Psi_j \circ T_{\frac{iy}{2}}$ , we can suppose  $y = 0$ .

Since  $T_{\frac{1}{p}}$  commutes with the  $\Psi_j$ 's, we have  $c_{n,j} = 0$  if  $p \nmid n$ , so that (36) becomes  $\overline{c_{sp,j}} = e^{-i\pi s a p/r} c_{-sp, -j}$ .

If  $p/r$  and  $a$  are odd, then (36) becomes

$$\overline{c_{sp,j}} = \begin{cases} -c_{sp,j}, & s \text{ odd}, \\ c_{sp,j}, & s \text{ even}. \end{cases}$$

This is precisely the condition for  $g$  to have an antiholomorphic root of order  $2p$ .

If  $p/r$  or  $a$  is even, then  $e^{-i\pi s a p/r} = 1$  for all  $s$ , so we obtain the condition necessary for  $\Sigma$  to be compatible with the transition functions, and then there is a Schwarz reflection in the centralizer of  $g$ . In that case, then the highest orders of the holomorphic and antiholomorphic iterates coincide.  $\square$

Lastly, we will study the centralizer of  $f \in \overline{\mathcal{A}_{k,b}}$  in  $\text{Diff}_1(0, \mathbb{C}, \overline{\mathbb{C}})$ . Of course, we have  $Z_f \subseteq Z_{f \circ f}$ , so the details are similar to the previous theorem.

**Theorem 7.4.4.** *Let  $f \in \overline{\mathcal{A}_{k,b}}$ . Then we have one of the following cases:*

1.  *$f$  is embeddable, i.e.  $f = h \circ \sigma \circ v^{\frac{1}{2}} \circ h^{-1}$ , then its centralizer is*

$$Z_f = \{h \circ v^t \circ h^{-1} \mid t \in \mathbb{R}\} \cup \{h \circ \sigma \circ v^t \circ h^{-1} \mid t \in \mathbb{R}\}.$$

2.  *$f$  is not embeddable, then  $Z_f$  contains only holomorphic and antiholomorphic fractional iterates of  $f$  and Schwarz reflections. More precisely, there exists  $p \in \mathbb{N} \setminus \{0\}$  such that the centralizer of  $f$  is one of the following:*

- *if  $p$  is odd, then*

$$Z_f = \bigcup_{d|p} \{\text{hol. fractional iterates of order } d \text{ of } f \circ f\} \cup \bigcup_{\substack{d|p \\ d \text{ odd}}} \{\text{antihol. fractional iterates of order } d \text{ of } f\};$$

- *if  $p$  is even, then*

$$Z_f = \bigcup_{d|p} \{\text{hol. fractional iterates of order } d \text{ of } f \circ f\} \cup \bigcup_{\substack{d|p \\ d \text{ odd}}} \{\text{antihol. fractional iterates of order } d \text{ of } f\};$$

$$\cup \{\text{Schwarz reflection tangent to } \sigma\}.$$

*The centralizer includes a Schwarz reflection tangent to  $\sigma$  if and only if  $f$  is analytically conjugate to a germ with real coefficients (see Theorem 7.3.1).*

*Proof.* 1. In the Fatou coordinates,  $T_t$  must commute with  $\Sigma \circ T_{\frac{1}{2}}$ , so  $t$  must be real. The rest of the details are exactly as in the proof of Theorem 7.4.3.

2. Let  $p \in \mathbb{N}$  be the highest order of the holomorphic roots of  $f \circ f$ . This number might be 1, in which case  $Z_f$  contains only iterates of  $f$ . Indeed, if  $p = 1$ ,  $Z_f$  does not contain a Schwarz reflection, otherwise  $f \circ f$  would have the holomorphic root  $\sigma \circ f$ , which would contradict that the highest order of the holomorphic root is 1. Let us suppose that  $p > 1$ .

Let  $(\Psi_1, \dots, \Psi_k)$  be a representative of the analytic invariant of  $f$ . Let  $\Psi_{-j}$  be determined by (17) for  $j = 1, \dots, k$ . Suppose that the



transition functions have a Fourier expansion

$$\Psi_j(W) = W + C_j + \sum_n c_{n,j} e^{2i\pi n W},$$

for all  $j$ , where  $\overline{C_{-j}} = C_j = (-1)^j \frac{i\pi b}{k}$  for  $j > 0$ . Because the  $\Psi_j$ 's satisfy  $\Psi_j \circ \Sigma \circ T_{\frac{1}{2}} = \Sigma \circ T_{\frac{1}{2}} \circ \Psi_{-j}$ , we obtain

$$(37) \quad \overline{c_{n,j}} = e^{i\pi n} c_{-n,-j}.$$

Since  $f \circ f$  has a root of order  $p$ , we also have  $T_{\frac{1}{p}} \circ \Psi_j = \Psi_j \circ T_{\frac{1}{p}}$ , which means that  $c_{n,j} = 0$  for all  $j$  when  $p \nmid n$ . If we write  $n = \ell p$  in (37), it becomes

$$\overline{c_{\ell p,j}} = e^{i\pi \ell p} c_{-\ell p,-j}, \quad (\ell \in \mathbb{Z}).$$

We have two cases.

If  $p$  is odd, then we have

$$\overline{c_{\ell p,j}} = \begin{cases} c_{-\ell p,-j} & \text{if } \ell \text{ is odd,} \\ -c_{-\ell p,-j} & \text{if } \ell \text{ is even.} \end{cases}$$

This corresponds to the necessary condition for  $f$  to have an antiholomorphic  $p$ -th root. Therefore, the maximal order of antiholomorphic root of  $f$  is at least  $p$ . To see that it is exactly  $p$ , we simply note that if  $f_1$  is an antiholomorphic root of order  $q \geq p$ , then  $f_1 \circ f_1$  is a holomorphic root of order  $q$  of  $f \circ f$ , and because  $p$  is maximal, we must have  $p = q$ . We must also prove that antiholomorphic fractional iterates of  $f \circ f$  are fractional iterates of  $f$ . Suppose  $f \circ f$  has an antiholomorphic root of order  $q$ , where  $q$  must be even. Then we have  $\frac{q}{2} \mid p$ . Since  $p$  is odd, we know that  $\frac{q}{2}$  must be odd. It follows that  $(\Sigma \circ T_{\frac{1}{q}})^{\circ \frac{q}{2}} = \Sigma \circ T_{\frac{1}{2}}$ .

If  $p$  is even, then we have  $\overline{c_{n,j}} = c_{-n,-j}$ , which corresponds to the condition necessary for  $\Sigma$  to be compatible with the  $\Psi_j$ 's. In this case, the centralizer of  $f$  contains a Schwarz reflection, and it follows that the highest order of antiholomorphic fractional iterates of  $f$  and the highest order of holomorphic fractional iterates of  $f \circ f$  coincide.  $\square$

## REFERENCES

- [1] L.V. Ahlfors. *Lectures on Quasiconformal Mappings*, by Lars V. Ahlfors. Manuscript Prepared with the Assistance of Clifford J. Earle, Jr. Van Nostrand mathematical studies, 10. Van Nostrand, 1966.
- [2] A.F. Beardon. *Iteration of rational functions: complex analytic dynamical systems*. Graduate texts in mathematics. Springer-Verlag, 1991.
- [3] L.C. Grove and C.T. Benson. *Finite Reflection Groups*. Graduate Texts in Mathematics. Springer New York, 2013.

- [4] J.H. Hubbard and D. Schleicher. Multicorns are not path connected. In *Frontiers in complex dynamics*, volume 51 of *Princeton Math. Ser.*, pages 73–102. Princeton Univ. Press, Princeton, NJ, 2014.
- [5] Y. Ilyashenko. Nonlinear Stokes phenomena. In *Nonlinear Stokes phenomena*, volume 14 of *Adv. Soviet Math.*, pages 1–55. Amer. Math. Soc., Providence, RI, 1993.
- [6] Y. Ilyashenko and S. Yakovenko. *Lectures on analytic differential equations*. Graduate studies in mathematics, v. 86. American Mathematical Society, Providence, R.I., 2008.
- [7] S. Mukherjee, S. Nakane, and D. Schleicher. On multicorns and unicorns II: bifurcations in spaces of antiholomorphic polynomials. *Ergodic Theory Dynam. Systems*, 37(3):859–899, 2017.
- [8] S. Nakane and D. Schleicher. On multicorns and unicorns. I. Antiholomorphic dynamics, hyperbolic components and real cubic polynomials. *Internat. J. Bifur. Chaos Appl. Sci. Engrg.*, 13(10):2825–2844, 2003.

*E-mail address:* godinj@dms.umontreal.ca

*E-mail address:* rousseac@dms.umontreal.ca

UNIVERSITÉ DE MONTRÉAL, C.P. 6128, SUCC. CENTRE-VILLE, MONTRÉAL,  
QC, CANADA, H3C 3J7