

Log-concavity and log-convexity of moments of averages of i.i.d. random variables

Philip Lamkin* and Tomasz Tkocz†

21st April 2020

Abstract

We show that the sequence of moments of order less than 1 of averages of i.i.d. positive random variables is log-concave. For moments of order at least 1, we conjecture that the sequence is log-convex and show that this holds eventually for integer moments (after neglecting the first p^2 terms of the sequence).

2010 Mathematics Subject Classification. Primary 05A20; Secondary 60E15, 26D15.

Key words. log-convexity, log-concavity, moment comparison, sums of independent random variables

1 Introduction and results

Suppose X_1, X_2, \dots are i.i.d. copies of a positive random variable and f is a nonnegative function. This article is concerned with certain combinatorial properties of the sequence

$$a_n = \mathbb{E}f\left(\frac{X_1 + \dots + X_n}{n}\right), \quad n = 1, 2, \dots \quad (1)$$

For instance, $f(x) = x^p$ is a fairly natural choice leading to the sequence of moments of averages of the X_i . Since we have the identity

$$\sum_{i=1}^{n+1} x_i = \sum_{i=1}^{n+1} \frac{\sum_{j:j \neq i} x_j}{n},$$

we conclude that the sequence $(a_n)_{n=1}^\infty$ is nonincreasing when f is convex. What about inequalities involving more than two terms?

Such inequalities have been studied to some extent. One fairly general result is due to Boland, Proschan and Tong from [1] (with applications in reliability theory). It asserts in particular that for $n = 2, 3, \dots$,

$$\mathbb{E}\phi(X_1 + \dots + X_n, X_{n+1} + \dots + X_{2n}) \leq \mathbb{E}\phi(X_1 + \dots + X_{n-1}, X_n + \dots + X_{2n}) \quad (2)$$

for a symmetric (invariant under permuting coordinates) continuous random vector $X = (X_1, \dots, X_{2n})$ with nonnegative components and a symmetric convex function $\phi : [0, +\infty)^2 \rightarrow \mathbb{R}$.

*Carnegie Mellon University; Pittsburgh, PA 15213, USA. Email: plamkin@andrew.cmu.edu

† Carnegie Mellon University; Pittsburgh, PA 15213, USA. Email: ttkocz@math.cmu.edu. Research supported in part by the Collaboration Grants from the Simons Foundation.

We obtain a satisfactory answer to a natural question of log-convexity/concavity of sequences (a_n) for completely monotone functions, also providing insights into the case of power functions.

Recall that a nonnegative sequence $(x_n)_{n=1}^\infty$ supported on a set of contiguous integers is called log-convex (resp. log-concave) if $x_n^2 \leq x_{n-1}x_{n+1}$ (resp. $x_n^2 \geq x_{n-1}x_{n+1}$) for all $n \geq 2$ (for background on log-convex/concave sequences, see for instance [8, 12]). One of the crucial properties of log-convex sequences is that log-convexity is preserved by taking sums (which follows from the Cauchy-Schwarz inequality, see for instance [8]). Recall that an infinitely differentiable function $f : (0, \infty) \rightarrow (0, \infty)$ is called completely monotone if we have $(-1)^n f^{(n)}(x) \geq 0$ for all positive x and $n = 1, 2, \dots$; equivalently, by Bernstein's theorem (see for instance [7]), the function f is the Laplace transform of a nonnegative Borel measure μ on $[0, +\infty)$, that is

$$f(x) = \int_0^\infty e^{-tx} d\mu(t). \quad (3)$$

For example, when $p < 0$, the function $f(x) = x^p$ is completely monotone. Such integral representations are at the heart of our first two results.

Theorem 1. *Let $f : (0, \infty) \rightarrow (0, \infty)$ be a completely monotone function. Let X_1, X_2, \dots be i.i.d. positive random variables. Then the sequence $(a_n)_{n=1}^\infty$ defined by (1) is log-convex.*

Theorem 2. *Let $f : [0, \infty) \rightarrow [0, \infty)$ be such that $f(0) = 0$ and its derivative f' is completely monotone. Let X_1, X_2, \dots be i.i.d. nonnegative random variables. Then the sequence $(a_n)_{n=1}^\infty$ defined by (1) is log-concave.*

In particular, applying these to the functions $f(x) = x^p$ with $p < 0$ and $0 < p < 1$ respectively, we obtain the following corollary.

Corollary 3. *Let X_1, X_2, \dots be i.i.d. positive random variables. The sequence*

$$b_n = \mathbb{E} \left(\frac{X_1 + \dots + X_n}{n} \right)^p, \quad n = 1, 2, \dots \quad (4)$$

is log-convex when $p < 0$ and log-concave when $0 < p < 1$.

For $p > 1$, we pose the following conjecture.

Conjecture 1. Let $p > 1$. Let X_1, X_2, \dots be i.i.d. nonnegative random variables. Then the sequence (b_n) defined in (4) is log-convex.

We offer a partial result supporting this conjecture.

Theorem 4. *Let X_1, X_2, \dots be i.i.d. nonnegative random variables, let p be a positive integer and let b_n be defined by (4). Then for every $n \geq p^2$, we have $b_n^2 \leq b_{n-1}b_{n+1}$.*

Remark 5. When $p = 2$, we have $b_n = \frac{n\mathbb{E}X_1^2 + n(n-1)(\mathbb{E}X_1)^2}{n^2} = (\mathbb{E}X_1)^2 + n^{-1}\text{Var}(X_1)$, which is clearly a log-convex sequence (as a sum of two log-convex sequences). The

following argument for $p = 3$ was kindly communicated to us by Krzysztof Oleszkiewicz: when $p = 3$, we can write

$$b_n = (\mathbb{E}X_1)^3 + (\mathbb{E}X_1^3 + (\mathbb{E}X_1)^3 - 2(\mathbb{E}X_1^2)(\mathbb{E}X_1))n^{-2} + (\mathbb{E}X_1) \text{Var}(X_1)(3n^{-1} - n^{-2}).$$

The sequences (n^{-2}) and $(3n^{-1} - n^{-2})$ are log-convex. By the Cauchy-Schwarz inequality the factor at n^{-2} is nonnegative,

$$\mathbb{E}X_1^2 \leq \sqrt{\mathbb{E}X_1^3} \sqrt{\mathbb{E}X_1} \leq \frac{\mathbb{E}X_1^3}{2\mathbb{E}X_1} + \frac{(\mathbb{E}X_1)^2}{2},$$

so again (b_n) is log-convex as a sum of three log-convex sequences. It remains elusive how to group terms and proceed along these lines in general. Our proof of Theorem 4 relies on this idea, but uses a straightforward way of rearranging terms.

Remark 6. It would be tempting to use the aforementioned result of Boland et al. with $\phi(x, y) = (xy)^p$ to resolve Conjecture 1. However, this function is neither convex nor concave on $(0, +\infty)^2$ for $p > \frac{1}{2}$. For $0 < p < \frac{1}{2}$, the function is concave and (2) yields $(b_n n^p)^2 \geq b_{n-1}(n-1)^p b_{n+1}(n+1)^p$, $n \geq 2$, equivalently, $b_n^2 \geq \left(\frac{n^2-1}{n^2}\right)^p b_{n-1} b_{n+1}$. Corollary 3 improves on this by removing the factor $\left(\frac{n^2-1}{n^2}\right)^p < 1$. For $p < 0$, ϕ is convex, so (2) gives $b_n^2 \leq \left(\frac{n^2-1}{n^2}\right)^p b_{n-1} b_{n+1}$ and Corollary 3 removes the factor $\left(\frac{n^2-1}{n^2}\right)^p > 1$.

Concluding this introduction, it is of significant interest to study the log-behaviour of various sequences, particularly those emerging from algebraic, combinatorial, or geometric structures, which has involved and prompted the development of many deep and interesting methods, often useful beyond the original problems (see, e.g., [3, 4, 5, 6, 10, 11, 12, 13]). We propose to consider sequences of moments of averages of i.i.d. random variables arising naturally in probabilistic limit theorems. For moments of order less than 1, we employ an analytical approach exploiting integral representations for power functions. For moments of order higher than 1, our Conjecture 1, besides refining the monotonicity property of the sequence (b_n) (resulting from convexity), would furnish new examples of log-convex sequences. For instance, neither does it seem trivial, nor handled by known techniques, to determine whether the sequence obtained by taking the Bernoulli distribution with parameter $\theta \in (0, 1)$, $b_n = \sum_{k=0}^n \binom{n}{k} \left(\frac{k}{n}\right)^p \theta^k (1-\theta)^{n-k}$ is log-convex. In the case of integral p , we have $b_n = \sum_{k=0}^p S(p, k) \frac{n!}{(n-k)! n^p} \theta^k$, where $S(p, k)$ is the Stirling number of the second kind.

The rest of this paper is occupied with the proofs of Theorems 1, 2, 4 (in their order of statement) and then we conclude with additional remarks and conjectures.

2 Proofs

2.1 Proof of Theorem 1

Suppose that f is completely monotone. Using (3) and independence, we have

$$a_n = \mathbb{E}f\left(\frac{X_1 + \dots + X_n}{n}\right) = \int_0^\infty \left[\mathbb{E}e^{-tX_1/n}\right]^n d\mu(t).$$

Let $u_n(t) = [\mathbb{E}e^{-tX_1/n}]^n$. It suffices to show that for every positive t , the sequence $(u_n(t))$ is log-convex (because sums/integrals of log-convex sequences are log-convex: the Cauchy-Schwarz inequality applied to the measure μ yields

$$\left(\int \sqrt{u_{n-1}(t)u_{n+1}(t)} d\mu(t) \right)^2 \leq \left(\int u_{n-1}(t) d\mu(t) \right) \left(\int u_{n+1}(t) d\mu(t) \right),$$

which combined with $u_n(t) \leq \sqrt{u_{n-1}(t)u_{n+1}(t)}$, gives $a_n^2 \leq a_{n-1}a_{n+1}$. The log-convexity of $(u_n(t))$ follows from Hölder's inequality,

$$\mathbb{E}e^{-tX_1/n} = \mathbb{E}e^{-\frac{n-1}{2n} \frac{tX_1}{n-1}} e^{-\frac{n+1}{2n} \frac{tX_1}{n+1}} \leq \left(\mathbb{E}e^{-\frac{tX_1}{n-1}} \right)^{\frac{n-1}{2n}} \left(\mathbb{E}e^{-\frac{tX_1}{n+1}} \right)^{\frac{n+1}{2n}},$$

which finishes the proof.

2.2 Proof of Theorem 2

Suppose now that $f(0) = 0$ and f' is completely monotone, say $f'(x) = \int_0^\infty e^{-tx} d\mu(t)$ for some nonnegative Borel measure μ on $(0, \infty)$ (by (3)). Introducing a new measure $d\nu(t) = \frac{1}{t} d\mu(t)$ we can write

$$f(y) = f(y) - f(0) = \int_0^y f'(x) dx = \int_0^\infty \int_0^\infty t e^{-tx} \mathbf{1}_{\{0 < x < y\}} dx d\nu(t).$$

Integrating against dx gives

$$f(y) = \int_0^\infty [1 - e^{-ty}] d\nu(t).$$

Let F be the Laplace transform of X_1 , that is

$$F(t) = \mathbb{E}e^{-tX_1}, \quad t > 0.$$

Then

$$\mathbb{E}f\left(\frac{X_1 + \dots + X_n}{n}\right) = \int_0^\infty [1 - F(t/n)^n] d\nu(t) = \int_0^\infty G(n, t) d\nu(t),$$

where, to shorten the notation, we introduce the following nonnegative function

$$G(\alpha, t) = 1 - F(t/\alpha)^\alpha, \quad \alpha, t > 0.$$

To show the inequality

$$\left[\mathbb{E}f\left(\frac{X_1 + \dots + X_n}{n}\right) \right]^2 \geq \mathbb{E}f\left(\frac{X_1 + \dots + X_{n-1}}{n-1}\right) \cdot \mathbb{E}f\left(\frac{X_1 + \dots + X_{n+1}}{n+1}\right)$$

it suffices to show that pointwise

$$G(n, s)G(n, t) \geq \frac{1}{2}G(n-1, s)G(n+1, t) + \frac{1}{2}G(n+1, s)G(n-1, t),$$

for all $s, t > 0$. This follows from two properties of the function G :

- 1) for every fixed $t > 0$ the function $\alpha \mapsto G(\alpha, t)$ is nondecreasing,
- 2) the function $G(\alpha, t)$ is concave on $(0, \infty) \times (0, \infty)$.

Indeed, by 2) we have

$$G(n, s)G(n, t) \geq \frac{G(n-1, s) + G(n+1, s)}{2} \cdot \frac{G(n-1, t) + G(n+1, t)}{2}$$

(in fact we only use concavity in the first argument). It thus suffices to prove that

$$\begin{aligned} & G(n-1, s)G(n-1, t) + G(n+1, s)G(n+1, t) \\ & - G(n-1, s)G(n+1, t) - G(n+1, s)G(n-1, t) \\ & = [G(n-1, s) - G(n+1, s)] \cdot [G(n-1, t) - G(n+1, t)] \end{aligned}$$

is nonnegative, which follows by 1).

It remains to prove 1) and 2). To prove the former, first notice that $F(t/\alpha)^\alpha = (\mathbb{E}e^{-tX/\alpha})^\alpha = \|e^{-tX}\|_{1/\alpha}$ is the $L_{1/\alpha}$ -norm of e^{-tX} . By convexity, for an arbitrary random variable Z , $p \mapsto (\mathbb{E}|Z|^p)^{1/p} = \|Z\|_p$ is nondecreasing, so $F(t/\alpha)^\alpha = \|e^{-tX}\|_{1/\alpha}$ is nonincreasing and thus $G(\alpha, t) = 1 - F(t/\alpha)^\alpha$ is nondecreasing. To prove the latter, notice that by Hölder's inequality the function $t \mapsto \log F(t)$ is convex,

$$F(\lambda s + (1-\lambda)t) = \mathbb{E}[(e^{-sX})^\lambda (e^{-tX})^{1-\lambda}] \leq (\mathbb{E}e^{-sX})^\lambda (\mathbb{E}e^{-tX})^{1-\lambda} = F(s)^\lambda F(t)^{1-\lambda}.$$

Therefore its perspective function $H(\alpha, t) = \alpha \log F(t/\alpha)$ is convex (see, e.g. Ch. 3.2.6 in [2]), which implies that $F(t/\alpha)^\alpha = e^{H(\alpha, t)}$ is also convex.

2.3 Proof of Theorem 4

We recall a standard combinatorial formula: first by the multinomial theorem and independence, we have

$$\mathbb{E}(X_1 + \dots + X_n)^p = \sum \frac{p!}{p_1! \dots p_n!} \mathbb{E}(X_1^{p_1} \dots X_n^{p_n}) = \sum \frac{p!}{p_1! \dots p_n!} \mu_{p_1} \dots \mu_{p_n},$$

where the sum is over all sequences (p_1, \dots, p_n) of nonnegative integers such that $p_1 + \dots + p_n = p$ and we denote $\mu_k = \mathbb{E}X_1^k$, $k \geq 0$. Now we partition the summation according to the number m of positive terms in the sequence (p_1, \dots, p_n) . Let Q_m be the set of integer partitions of p into exactly m (nonempty) parts, that is the set of m -element multisets $q = \{q_1, \dots, q_m\}$ with positive integers q_j summing to p , $q_1 + \dots + q_m = p$. Then

$$\mathbb{E}(X_1 + \dots + X_n)^p = \sum_{m=1}^p \sum_{q \in Q_m} \frac{p!}{q_1! \dots q_m!} \frac{n!}{\alpha(q) \cdot (n-m)!} \mu_{q_1} \dots \mu_{q_m},$$

where $\alpha(q) = l_1! \dots l_h!$ for $q = \{q_1, \dots, q_m\}$ with exactly h distinct terms such that there are l_1 terms of type 1, l_2 terms of type 2, etc., so $l_1 + \dots + l_h = m$ (e.g. for $q = \{2, 2, 2, 3, 4, 4\} \in Q_6$, we have $h = 3$, $l_1 = 3$, $l_2 = 1$, $l_3 = 2$). The factor $\frac{n!}{\alpha(q) \cdot (n-m)!}$

arises because given a multiset $q \in Q_m$, there are exactly

$$\begin{aligned} & \binom{n}{l_1} \binom{n-l_1}{l_2} \binom{n-l_1-l_2}{l_3} \cdots \binom{n-l_1-\cdots-l_{h-1}}{l_h} \\ &= \frac{n!}{l_1! \cdots l_h! \cdot (n-l_1-\cdots-l_h)!} = \frac{n!}{\alpha(q) \cdot (n-m)!} \end{aligned}$$

many nonnegative integer-valued sequences (p_1, \dots, p_n) with $\mu_{p_1} \cdots \mu_{p_n} = \mu_{q_1} \cdots \mu_{q_m}$ (equivalently, $\{p_1, \dots, p_n\} = \{q_1, \dots, q_m, 0\}$, as sets).

We have obtained

$$b_n = \mathbb{E} \left(\frac{X_1 + \cdots + X_n}{n} \right)^p = \sum_{m=1}^p \frac{n!}{n^p (n-m)!} \sum_{q \in Q_m} \beta(q) \mu(q), \quad (5)$$

where $\beta(q) = \frac{p!}{\alpha(q) \cdot q_1! \cdots q_m!}$ and $\mu(q) = \mu_{q_1} \cdots \mu_{q_m}$. By homogeneity, we can assume that $\mu_1 = \mathbb{E}X_1 = 1$. Note that when X_1 is constant, we get from (5) that

$$1 = \sum_{m=1}^p \frac{n!}{n^p (n-m)!} \sum_{q \in Q_m} \beta(q).$$

Since Q_p has only one element, namely $\{1, \dots, 1\}$ and $\mu(\{1, \dots, 1\}) = 1$, when we subtract the two equations, the terms corresponding to $m = p$ cancel and we get

$$b_n - 1 = \sum_{m=1}^{p-1} \frac{n!}{n^p (n-m)!} \sum_{q \in Q_m} \beta(q) (\mu(q) - 1).$$

By the monotonicity of moments, $\mu(q) \geq 1$ for every q , so (b_n) is a sum of the constant sequence $(1, 1, \dots)$ and the sequences $(u_n^{(m)}) = (\frac{n!}{n^p (n-m)!})$, $m = 1, \dots, p-1$, multiplied respectively by the nonnegative factors $\sum_{q \in Q_m} \beta(q) (\mu(q) - 1)$. Since sums of log-convex sequences are log-convex, it remains to verify that for each $1 \leq m \leq p-1$, we have $(u_n^{(m)})^2 \leq u_{n-1}^{(m)} u_{n+1}^{(m)}$, $n \geq p^2$. The following lemma finishes the proof.

Lemma 7. *Let $p \geq 2$, $1 \leq m \leq p-1$ be integers. Then the function*

$$f(x) = \log \frac{x(x-1) \cdots (x-m+1)}{x^p}$$

is convex on $[p^2 - 1, \infty)$.

Proof. The statement is clear for $m = 1$. Let $2 \leq m \leq p-1$ and $p \geq 3$. We have

$$x^2 f''(x) = p-1 - x^2 \sum_{k=1}^{m-1} \frac{1}{(x-k)^2}.$$

To see that this is positive for every $x \geq p^2 - 1$ and $2 \leq m \leq p-1$, it suffices to consider $m = p-1$ and $x = p^2 - 1$ (writing $\frac{x}{x-k} = 1 + \frac{k}{x-k}$, we see that the right hand side is

increasing in x). Since

$$\sum_{k=1}^{p-2} \frac{1}{(p^2-1-k)^2} = \sum_{k=p^2-p+1}^{p^2-2} \frac{1}{k^2} \leq \sum_{k=p^2-p+1}^{p^2-2} \left(\frac{1}{k-1} - \frac{1}{k} \right) = \frac{1}{p^2-p} - \frac{1}{p^2-2},$$

we have

$$x^2 f''(x) \geq p-1 - (p^2-1)^2 \left(\frac{1}{p^2-p} - \frac{1}{p^2-2} \right) = \frac{(p-1)(p+2)}{p(p^2-2)},$$

which is clearly positive. \square

3 Final remarks

Remark 8. Using majorization type arguments (see, e.g. [9]), Conjecture 1 can be verified in a rather standard but lengthy way for every $p > 1$ and $n = 2$. The idea is to establish a pointwise inequality: we conjecture that for nonnegative numbers x_1, \dots, x_{2n} and a convex function $\phi : [0, \infty) \rightarrow [0, \infty)$ we have

$$\frac{1}{\binom{2n}{n}} \sum_{|I|=n} \phi \left(\frac{x_I x_{I^c}}{n^2} \right) \leq \frac{1}{\binom{2n}{n+1}} \sum_{|I|=n+1} \phi \left(\frac{x_I x_{I^c}}{n^2-1} \right),$$

where for a subset I of the set $\{1, \dots, 2n\}$ we denote $x_I = \sum_{i \in I} x_i$. We checked that this holds for $n = 2$. Taking the expectation on both sides for $\phi(x) = x^p$ gives the desired result that $b_n^2 \leq b_{n-1} b_{n+1}$.

Remark 9. It is tempting to ask for generalisations of Conjecture 1 beyond the power functions, say to ask whether the sequence (a_n) defined in (1) is log-convex for every convex function f . This is false, as can be seen by taking the function f of the form $f(x) = \max\{x-a, 0\}$ and the X_i to be i.i.d Bernoulli random variables.

Acknowledgments. We thank Krzysztof Oleszkiewicz for his great help, valuable correspondence, feedback and for allowing us to use his slick proof for $p = 3$ from Remark 5. We also thank Marta Strzelecka and Michał Strzelecki for the helpful discussions. We are indebted to the referees for their careful reading and useful comments.

We are grateful for the excellent working conditions provided by the Summer Undergraduate Research Fellowship 2019 at Carnegie Mellon University during which this work was initiated.

References

- [1] Boland, P., Proschan, F., Tong, Y., Some majorization inequalities for functions of exchangeable random variables. *Topics in statistical dependence (Somerset, PA, 1987)*, 85–91, IMS Lecture Notes Monogr. Ser., 16, *Inst. Math. Statist., Hayward, CA*, 1990.

- [2] Boyd, S., Vandenberghe, L., Convex optimization. *Cambridge University Press, Cambridge*, 2004.
- [3] Brenti, F., Unimodal, log-concave and Pólya frequency sequences in combinatorics. *Mem. Amer. Math. Soc.* 81 (1989), no. 413, viii+106 pp.
- [4] Došlić, T., Seven (lattice) paths to log-convexity. *Acta Appl. Math.* 110 (2010), no. 3, 1373–1392.
- [5] Došlić, T., Veljan, D., Logarithmic behavior of some combinatorial sequences. *Discrete Math.* 308 (2008), no. 11, 2182–2212.
- [6] Fadnavis, S., On Brenti’s conjecture about the log-concavity of the chromatic polynomial. *European J. Combin.* 33 (2012), no. 8, 1842–1846.
- [7] Feller, W., An introduction to probability theory and its applications. Vol. II. Second edition, *John Wiley & Sons, Inc., New York-London-Sydney* 1971.
- [8] Liu, L., Wang, Y., On the log-convexity of combinatorial sequences. *Adv. in Appl. Math.* 39 (2007), no. 4, 453–476.
- [9] Marshall, A., Olkin, I., Arnold, B., Inequalities: theory of majorization and its applications. Second edition. Springer Series in Statistics. *Springer, New York*, 2011.
- [10] Nayar, P., Oleszkiewicz, K., Khinchine type inequalities with optimal constants via ultra log-concavity. *Positivity* 16 (2012), no. 2, 359–371.
- [11] Sagan, B., Inductive and injective proofs of log concavity results. *Discrete Math.* 68 (1988), no. 2–3, 281–292.
- [12] Stanley, R., Log-concave and unimodal sequences in algebra, combinatorics, and geometry. *Graph theory and its applications: East and West (Jinan, 1986)*, 500–535, Ann. New York Acad. Sci., 576, *New York Acad. Sci., New York*, 1989.
- [13] Stanley, R., Positivity problems and conjectures in algebraic combinatorics. *Mathematics: frontiers and perspectives*, 295–319, *Amer. Math. Soc., Providence, RI*, 2000.