

# K-NN active learning under local smoothness assumption

Boris Ndjia Njike · Xavier Siebert

Received: date / Accepted: date

**Abstract** There is a large body of work on convergence rates either in passive or in active learning. Here we first outline some of the main results that have been obtained, more specifically in a nonparametric setting under assumptions about the smoothness of the regression function (or the boundary between classes) and the margin noise. We discuss the relative merits of these underlying assumptions by putting active learning in perspective with recent work on passive learning. In particular, the regression function is assumed to be smooth with respect to the marginal probability of the data. This is more general than what was previously used in the literature. Our contribution is an active nearest neighbor learning algorithm, that is able to deal with this setting, and that outperforms its passive counterpart. Our algorithm works for a larger class of probability distributions than those previously used, especially for distributions of probability for which the density function is not necessarily bounded below, and also for discrete distributions.

**Keywords** Nonparametric learning · active learning · nearest-neighbors · smoothness condition.

---

F. Author

Affiliation: University of Mons, Department of Mathematics and Operational Research

Tel.: +32 65-374677

Fax: +32 65-374677

E-mail: borisedgar.ndjianjike@umons.ac.be

ORCID ID: <https://orcid.org/0000-0002-2122-994>

S. Author

Affiliation: University of Mons, Department of Mathematics and Operational Research

Tel.: +32 65-374690

Fax: +32 65-374690

E-mail: xavier.siebert@umons.ac.be

## 1 Introduction

Active learning is a machine learning approach for reducing the data labeling effort. Given an instance space  $\mathcal{X}$  or a pool of unlabeled data  $\{X_1, \dots, X_w\}$  provided by a distribution  $P_X$ , the learner focuses its labeling effort only on the most “informative” points so that a model built from them can achieve the best possible guarantees (Dasgupta, 2011). Such guarantees are particularly interesting when they are significantly better than those obtained in passive learning (Hanneke and Yang, 2015). In the context of this work, we consider binary classification (where the label  $Y$  of  $X$  takes its value in  $\{0, 1\}$ ) in a nonparametric setting. Extensions to multiclass classification and adaptive algorithms are discussed at the end of this paper (Section 6).

The nonparametric setting has the advantage of providing guarantees with many informations such as the dependence on the dimensional and distributional parameters by using some hypotheses on the regularity of the decision boundary (Castro and Nowak, 2008), on the regression function (Minsker, 2012; Locatelli et al., 2017), and on the geometry of instance space (called strong density assumption) (Audibert and Tsybakov, 2007; Locatelli et al., 2017; Minsker, 2012). One of the initial works on nonparametric active learning (Castro and Nowak, 2008) assumed that the decision boundary is the graph of a smooth function, that a margin assumption very similar to Tsybakov’s noise assumption (Mammen and Tsybakov, 1999) holds, and that distribution  $P_X$  is uniform. This led to a better guarantee than in passive learning. Instead of the assumption on the decision boundary, other works (Minsker, 2012; Locatelli et al., 2017) supposed rather that the regression function is smooth (in some sense). This assumption, along with Tsybakov’s noise assumption and the strong density assumption also gave a better guarantee than in passive learning. Moreover, unlike in (Castro and Nowak, 2008), they provided algorithms that are adaptive with respect to the margin’s noise and to the smoothness parameters.

However, recent work (Chaudhuri and Dasgupta, 2014) pointed out some disadvantages of the preceding smoothness assumption, and extended it in the context of passive learning with  $k$ -nearest neighbors ( $k$ -NN) by using a more general smoothness assumption that is able to sharply characterize the rate of convergence for all probability distributions that satisfy it.

In this paper, we thus extend the work of (Chaudhuri and Dasgupta, 2014) to the active learning setting, and provide a novel algorithm that outputs a classifier with the same rate of convergence as other recent algorithms with more restrictive hypotheses, as for example (Minsker, 2012; Locatelli et al., 2017). Section 2 introduces general definitions, Section 3 presents previous work on convergence rates in active and passive non-parametric learning, with a special emphasis on the assumptions related to our work. Section 4 provides an outline of our algorithm while Section 5 describes its theoretical motivations and Section 6 contains the conclusion and some perspectives for future work.

## 2 Preliminaries

We begin with some general definitions and notations about active learning in binary classification, then recall the concept of  $k$ -NN classifiers. Finally, the main assumptions that are used in nonparametric active learning are explained.

### 2.1 Active learning setting

Let  $(\mathcal{X}, \rho)$  be a metric space. In this paper we set  $\mathcal{X} \subset \mathbb{R}^d$  and refer to it as the instance space, and take  $\rho$  as the Euclidean metric. Let  $\mathcal{Y} = \{0, 1\}$  the label space. We assume that the pairs  $(X, Y)$  are random variables distributed according to an unknown probability  $P$  over  $\mathcal{X} \times \mathcal{Y}$ . Let us denote  $P_X$  the marginal distribution of  $P$  over  $\mathcal{X}$ .

Given  $w \in \mathbb{N}$  and an i.i.d. sample  $(X_1, Y_1), \dots, (X_w, Y_w)$  drawn according to probability  $P$ , the learning problem consists in minimizing the risk  $\mathcal{R}(f) = P(Y \neq f(X))$  over all measurable functions, called classifiers,  $f : \mathcal{X} \rightarrow \mathcal{Y}$ .

In active learning, the labels are not available from the beginning but we can request iteratively at a certain cost (to a so-called oracle) a given number  $n$  of samples, called the budget ( $n \leq w$ ). In passive learning, all labels are available and  $n = w$ . At any time, we choose to request the label of a point  $X$  according to the previous observations. The point  $X$  is chosen to be most “informative”, which amounts to belonging to a region where classification is difficult and requires more labeled data to be collected. Therefore, the goal of active learning is to design a sampling strategy that outputs a classifier  $\hat{f}_{n,w}$  whose excess risk (see below) is as small as possible with high probability over the requested samples, as reviewed in (Dasgupta, 2011; Hanneke and Yang, 2015; Dasgupta, 2017).

Given  $x$  in  $\mathcal{X}$ , let us introduce the regression function  $\eta(x) = \mathbb{E}(Y|X = x) = P(Y = 1|X = x)$ . It is easy to show (Lugosi, 2002) that the function  $f^*(x) = \mathbb{1}_{\eta(x) \geq 1/2}$  achieves the minimum risk and that  $\mathcal{R}(f^*) = \mathbb{E}_X(\min(\eta(X), 1 - \eta(X)))$ . Because  $P$  is unknown, the function  $f^*$  is unreachable and thus the aim of a learning algorithm is to return a classifier  $\hat{f}_{n,w}$  with minimum excess risk  $\mathcal{R}(\hat{f}_{n,w}) - \mathcal{R}(f^*)$  with high probability over the sample  $(X_1, Y_1), \dots, (X_w, Y_w)$ .

### 2.2 $k$ -Nearest Neighbors ( $k$ -NN) classifier

Given two integers  $k, n$  such that  $k < n$ , and a test point  $X \in \mathcal{X}$ , the  $k$ -NN classifier predicts the label of  $X$  by giving the majority vote of its  $k$  nearest neighbors amongst the sample  $X_1, \dots, X_n$ . For  $k = 1$ , the  $k$ -NN classifier returns the label of the nearest neighbor of  $X$  amongst the sample  $X_1, \dots, X_n$ . If  $k$  is allowed to grow with  $n$ , the method is called  $k_n$ -NN. For a complete discussion of nearest neighbors classification, see for example (Biau and Devroye, 2015; Shalev-Shwartz and Ben-David, 2014; Chaudhuri and Dasgupta, 2014).

### 2.3 Regularity, noise and strong density assumptions

Let  $B(x, r) = \{x' \in \mathcal{X}, \rho(x, x') < r\}$  and  $\bar{B}(x, r) = \{x' \in \mathcal{X}, \rho(x, x') \leq r\}$  the open and closed balls (with respect to the Euclidean metric  $\rho$ ), respectively, centered at  $x \in \mathcal{X}$  with radius  $r > 0$ . Let  $\text{supp}(P_X) = \{x \in \mathcal{X}, \forall r > 0, P_X(B(x, r)) > 0\}$  the support of the marginal distribution  $P_X$ .

**Definition 1a (Hölder continuity)**

Let  $\eta : \mathcal{X} \rightarrow [0, 1]$  be the regression function defined as  $\eta(x) = P(Y = 1|X = x)$ . We say that  $\eta$  is  $(\alpha, L)$ -**Hölder continuous** ( $0 < \alpha \leq 1$ , and  $L \geq 1$ ) if  $\forall x, x' \in \mathcal{X}$ ,

$$|\eta(x) - \eta(x')| \leq L\rho(x, x')^\alpha. \quad (\text{H1a})$$

The notion of Hölder continuity ensures that the proximity between two closest (according to the metric  $\rho$ ) points is reflected in a similar value of the conditional probability  $\eta$ .

This definition remains true for a general metric space, but when  $\rho$  is the Euclidean metric, we should always have  $0 < \alpha \leq 1$ , otherwise  $\eta$  becomes constant (Pugh, 2002).

In most of the previous works (for example (Audibert and Tsybakov, 2007; Minsker, 2012; Gaïffas, 2007)), the definition H1a is used along with the following notion (H1b) for technical reasons.

**Definition 1b (Strong density)**

Let  $P$  be the probability distribution defined over  $\mathcal{X} \times \mathcal{Y}$  and  $P_X$  the marginal distribution of  $P$  over  $\mathcal{X}$ . We say that  $P$  satisfies the **strong density** assumption if there exists some constants  $r_0 > 0$ ,  $c_0 > 0$ ,  $p_{\min} > 0$  such that for all  $x \in \text{supp}(P_X)$ :

$$\begin{aligned} \lambda(B(x, r) \cap \text{supp}(P_X)) &\geq c_0 \lambda(B(x, r)), \quad \forall r \leq r_0 \\ \text{and } p_X(x) &> p_{\min}, \end{aligned} \quad (\text{H1b})$$

where  $p_X$  is the density function of the marginal distribution  $P_X$  and  $\lambda$  is the Lebesgue measure.

The strong density assumption ensures that, given a realisation  $X = x$  according to  $P_X$ , there exists an infinite number of realisations  $X_1 = x_1, \dots, X_m = x_m, \dots$  in a neighborhood of  $x$ .

Sometimes, the notion of strong density is used to geometrically characterize the set where the classification is difficult (Locatelli et al., 2017), and then combined with the following definition of *Margin noise*, allows to nicely control the error of classification beyond a given number of label requests.

**Definition 2 (Margin noise)**

The probability distribution  $P$  satisfies the **margin noise** assumption (sometimes called **Tsybakov's noise** assumption (Audibert and Tsybakov, 2007)) with parameter  $\beta \geq 0$  if for all  $0 < \epsilon \leq 1$ , there is  $C = C(\beta) \in [1, +\infty[$  such that

$$P_X(x \in \mathcal{X}, |\eta(x) - 1/2| \leq \epsilon) < C\epsilon^\beta. \quad (\text{H2})$$

The margin noise assumption gives a bound on the probability that the label of the points in the neighborhood of a point  $x$  differs from the label of  $x$  given by the conditional probability  $\eta(x)$ . It also describes the behavior of the regression function in the vicinity of the decision boundary  $\eta(x) = \frac{1}{2}$ . When  $\beta$  goes to infinity, we observe a “jump” of  $\eta$  around the decision boundary, and then we obtain Massart’s noise condition (Massart and Nédélec, 2006). Small values of  $\beta$  allow for  $\eta$  to “cuddle”  $\frac{1}{2}$  when we approach the decision boundary.

**Definition 3** ( $(\alpha, L)$ -smooth)

Let  $0 < \alpha \leq 1$  and  $L > 1$ . The regression function is  $(\alpha, L)$ -**smooth** if for all  $x, z \in \text{supp}(P_X)$  we have:

$$|\eta(x) - \eta(z)| \leq L \cdot P_X(B(x, \rho(x, z)))^{\alpha/d}, \quad (\text{H3})$$

where  $d$  is the dimension of the instance space.

Equivalently, (H3) can be rewritten as:

$$|\eta(x) - \eta(z)| \leq L \cdot \min \left( P_X(B(x, \rho(x, z)))^{\alpha/d}, P_X(B(z, \rho(x, z)))^{\alpha/d} \right).$$

It is important to note that the  $(\alpha, L)$ -smooth assumption (H3) is more general than the Hölder continuity assumption (H1a), as stated in Theorem 1 below.

**Theorem 1** (Chaudhuri and Dasgupta, 2014)

Suppose that  $\mathcal{X} \subset \mathbb{R}^d$ , that the regression function  $\eta$  is  $(\alpha_h, L_h)$ -Hölder continuous, and that  $P_X$  satisfies H1b. Then there is a constant  $L > 1$  such that for any  $x, z \in \text{supp}(P_X)$ , we have:

$$|\eta(x) - \eta(z)| \leq L \cdot P_X(B(x, \rho(x, z)))^{\alpha_h/d}.$$

**Definition 4** (Doubling-probability)

The marginal distribution  $P_X$  is a **doubling-probability** if there exists a constant  $C_{db} > 0$  such that for any  $x \in \mathcal{X}$ , and  $r > 0$ , we have:

$$P_X(B(x, r)) \leq C_{db} P_X(B(x, r/2)). \quad (\text{H4})$$

This notion was initially introduced for geometric purposes in the setting of measure theory (Heinonen, 2012; Federer, 2014). It helps for constructing a subcover of a metric space by also minimizing the overlap between the elements of the subcover. Doubling-probability has been used in a machine learning context, particularly  $k$ -NN classification (or regression), where the constant  $C_{db}$  is interpreted as the intrinsic dimension of the region where the data belong (Kpotufe, 2011). This allows to reduce considerably the complexity of the classification problem and to bypass the so-called curse of dimension. Also, it is also proved (Kpotufe, 2011) that the notion of doubling-probability generalizes the strong density assumption H1b. It is thus more universal, and does not require a probability density.

In this paper, doubling-probability is used only for geometrical purposes. It is later relaxed, so that it becomes sufficient to consider only balls  $B(x, r)$  with  $P_X(B(x, r))$  sufficiently large to satisfy the doubling-probability condition (H4).

### 3 Convergence rates in nonparametric active learning

#### 3.1 Previous work

Active learning theory has been mostly studied during the last decades in a parametric setting, see for example (Balcan et al., 2010; Hanneke, 2011; Dasgupta, 2017) and references therein. One of the pioneering works studying the achievable limits in active learning in a nonparametric setting (Castro and Nowak, 2008) required that the decision boundary is the graph of a Hölder continuous function with parameter  $\alpha$  (H1a). Using a notion of margin noise (with parameter  $\beta$ ) very similar to (H2), the following minimax rate was obtained:

$$O\left(n^{-\frac{\beta}{2\beta+\gamma-2}}\right), \quad (1)$$

where  $\gamma = \frac{d-1}{\alpha}$  and  $d$  is the dimension of instance space ( $\mathcal{X} = \mathbb{R}^d$ ).

Note that this result assumes the knowledge of the smoothness and margin noise parameters, whereas an algorithm that achieves the same rate, but that adapts to these parameters was proposed recently in (Locatelli et al., 2018). In passive learning, by assuming that the regression function is Hölder continuous (H1a), along with (H1b) and (H2), the following minimax rate was established (Audibert and Tsybakov, 2007):

$$O\left(n^{-\frac{\alpha(\beta+1)}{2\alpha+d}}\right). \quad (2)$$

In active learning, using the same assumptions (H1a), (H1b) and (H2), with the additional condition  $\alpha\beta < d$ , the following minimax rate was obtained (Locatelli et al., 2017)

$$\tilde{O}\left(n^{-\frac{\alpha(\beta+1)}{2\alpha+d-\alpha\beta}}\right), \quad (3)$$

where  $\tilde{O}$  indicates that there may be additional logarithmic factors. This active learning rate given by (3) thus represents an improvement over the passive learning rate (2) that uses the same hypotheses.

With another assumption on the regression function relating the  $L_2$  and  $L_\infty$  approximation losses of certain piecewise constant or polynomial approximations of  $\eta$  in the vicinity of the decision boundary, the same rate (3) was also obtained (Minsker, 2012).

#### 3.2 Link with $k$ -NN classifiers

For practical applications, an interesting question is whether  $k$ -NN classifiers attain the rate given by (2) in passive learning and by (3) in active learning. In passive learning, under assumptions (H1a), (H1b) and (H2), and for suitable  $k_n$ , it was shown in (Chaudhuri and Dasgupta, 2014) that  $k_n$ -NN indeed achieves the rate (2).

In active learning a pool-based algorithm that outputs a  $k$ -NN classifier has been proposed in (Kontorovich et al., 2016), but its assumptions differ from ours in terms of smoothness and noise. Similarly, the algorithm proposed in (Hanneke) outputs a 1-NN classifier based on a subsample of a given pool of data, such that the label of each instance of this subsample is determined with high probability by the labels of its neighbors within the pool. The number of neighbors is adaptively chosen for each instance in the subsample, leading to the minimax rate (3) under the same assumptions as in (Locatelli et al., 2017).

To obtain more general results on the rate of convergence for  $k$ -NN classifiers in metric spaces under minimal assumptions, the more general smoothness assumption given by (H3) was used in (Chaudhuri and Dasgupta, 2014). By using a  $k$ -NN algorithm, and under assumptions (H2) and (H3), the rate of convergence obtained in (Chaudhuri and Dasgupta, 2014) is also of the order of (2). Additionally, using assumption (H3) instead of (H1a) removes the need for the strong density assumption (H1b), which therefore allows for more probability classes.

### 3.3 Contributions of the current work

In this work, we will use the assumptions that were used in the context of passive learning in (Chaudhuri and Dasgupta, 2014), and show that it is possible to use them in active learning as well.

For the sake of clarity, let us restate here these assumptions that will be used throughout this paper. We assume that the assumptions (H3), (H2), (H4) simultaneously hold respectively with parameters  $(\alpha, L)$ ,  $(\beta, C)$ ,  $C_{db}$ .

In this paper, we provide an active learning algorithm under assumptions (H3), (H2) that were used in passive learning in (Chaudhuri and Dasgupta, 2014). We additionally assume that the underlying marginal probability  $P_X$  satisfies (H4) mostly for geometrical convenience. Our algorithm has several advantages:

- The assumption H3 involves a dependence on the marginal distribution  $P_X$ , and holds for any pair of distributions  $P_X$  and  $\eta$ , which allows the use of discrete probabilities. However in active learning the Hölder continuity notion (H1a) is typically used, along with the strong density notion (H1b) (Castro and Nowak, 2008; Locatelli et al., 2017; Minsker, 2012), which implies assuming the existence of the density  $p_X$  of the marginal probability  $P_X$ . By using assumption (H3) instead of (H1a) and thereby avoiding (H1b), our algorithm removes unnecessary restrictions on the distribution that would exclude important densities (e.g., Gaussian) as noticed in (Döring et al., 2017).
- The rate of convergence of our algorithm is better than those obtained in passive learning under (H3) and (H2).

- According to the assumption (H2), as we will see, our algorithm also (as in (Minsker, 2012)) covers the most interesting case where the regression function is allowed to cross the boundary decision  $\{x, \eta(x) = \frac{1}{2}\}$ .

In the following, we will show that the rate of convergence of our algorithm remains the same as (3), despite the use of more general hypotheses.

## 4 KALLS algorithm

### 4.1 Setting

As explained in Section 2.1, we consider an active learning setting with a pool of i.i.d. unlabeled examples  $\mathcal{K} = \{X_1, X_2, \dots, X_w\}$ . Let  $n \leq w$  the budget, that is the maximum number of points whose label we are allowed to query to the oracle. The objective of the algorithm is to build a 1-NN classifier, based on a labelled set  $\mathcal{S}_{ac}$  of carefully chosen points. This set contains a subset of most *informative* points in  $\mathcal{K}$  and is called the *active set*. More precisely, a point  $X_t$  is considered *informative* if its label cannot be inferred (see below) from the previous observations  $X_{t'}$  (with  $t' < t$ ). The set  $\mathcal{S}_{ac}$  starts with  $X_{t_1} = X_1$  chosen arbitrarily in  $\mathcal{K}$  and stops when the budget  $n$  is reached or when  $X_w$  is attained.

When a point  $X_t$  is *informative*, instead of requesting directly its label to the (noisy) oracle, we infer it by requesting the labels of its nearest neighbors in  $\mathcal{K}$ , as was done in (Hanneke). This is reasonable for practical situations where the uncertainty about the label of  $X_t$  has to be overcome, and it is related to the assumption (H3). Note that it differs from the setting of (Locatelli et al., 2018), where the label of  $X_t$  is requested several times. The number of neighbors  $k_t$  used for inferring that label of  $X_t$  is determined such that, while respecting the budget, we can predict with high confidence the true label as  $f^*(X_t)$  of  $X_t$  by the empirical mean of the labels of its  $k_t$  nearest neighbors.

The labelled active set  $\mathcal{S}_{ac}$  output by the algorithm will comprise only the informative points on which we have sufficient guarantees when considering the inferred label as the right label. Finally, we show that the labelled active set  $\mathcal{S}_{ac}$  is sufficient to predict the label of any new point with a 1-NN classification rule  $\hat{f}_{n,w}$ .

### 4.2 Algorithm

The KALLS algorithm (Algorithm 1) aims at determining the *active set* defined in Section 4.1 and the related 1-NN classifier  $\hat{f}_{n,w}$  under the assumption (H3) and (H2).

Before beginning the description of KALLS, let us introduce some variables and notations, whose precise form will be justified in Section 5. The latter contains the proof sketch of the convergence of KALLS, while the complete proofs are in Appendix A.



For  $\epsilon, \delta \in (0, 1)$ ,  $k \geq 1$ , set:

$$b_{\delta,k} = \sqrt{\frac{2}{k} \left( \log \left( \frac{1}{\delta} \right) + \log \log \left( \frac{1}{\delta} \right) + \log \log(e.k) \right)}. \quad (4)$$

$$k(\epsilon, \delta) = \frac{c}{\Delta^2} \left\lceil \log \left( \frac{1}{\delta} \right) + \log \log \left( \frac{1}{\delta} \right) + \log \log \left( \frac{512\sqrt{e}}{\Delta} \right) \right\rceil. \quad (5)$$

where

$$\Delta = \max \left( \frac{\epsilon}{2}, \left( \frac{\epsilon}{2C} \right)^{\frac{1}{\beta+1}} \right), \quad c \geq 7.10^6. \quad (6)$$

Let

$$\phi_n = \sqrt{\frac{1}{n} \left( \log \left( \frac{1}{\delta} \right) + \log \log \left( \frac{1}{\delta} \right) \right)}. \quad (7)$$

For  $X_s \in \mathcal{K} = \{X_1, \dots, X_w\}$ , we denote henceforth by  $X_s^{(k)}$  its  $k$ -th nearest neighbor in  $\mathcal{K}$ , and  $Y_s^{(k)}$  the corresponding label.

For an integer  $k \geq 1$ , let

$$\hat{\eta}_k(X_s) = \frac{1}{k} \sum_{i=1}^k Y_s^{(i)}, \quad \bar{\eta}_k(X_s) = \frac{1}{k} \sum_{i=1}^k \eta(X_s^{(i)}). \quad (8)$$

The inputs of **KALLS** are a pool  $\mathcal{K}$  of unlabelled data of size  $w$ , the budget  $n$ , the smoothness parameters  $(\alpha, L)$  from (H3), the margin noise parameters  $(\beta, C)$  from (H2), a confidence parameter  $\delta \in (0, 1)$  and an accuracy parameter  $\epsilon \in (0, 1)$ . For the moment, these parameters are fixed from the beginning but adaptive algorithms such as (Locatelli et al., 2017) could be exploited, in particular for the  $\alpha$  and  $\beta$  parameters.

At any given stage, the current version of the labelled active set  $\mathcal{S}_{ac}$  is denoted by  $\hat{\mathcal{S}}$ . Based on  $\mathcal{S}_{ac}$ , with high confidence, the 1-NN classifier  $\hat{f}_{n,w}$  agrees with the Bayes classifier at points that lie beyond some margin  $\Delta_o > 0$  of the decision boundary. Formally, given  $x \in \mathcal{X}$  such that  $|\eta(x) - 1/2| > \Delta_o$ , we have  $\hat{f}_{n,w}(x) = \mathbb{1}_{\eta(x) \geq 1/2}$  with high confidence. We will show in Section 5 that, with a suitable choice of  $\Delta_o$ , the assumption (H2) leads to the desired rate of convergence (3).

**KALLS** uses two main subroutines : **Reliable** and **ConfidentLabel**, which are detailed below in Sections 4.3 and 4.4, respectively.

#### 4.3 **Reliable** subroutine

The **Reliable** subroutine is a binary test that checks if the label of a current point  $X$  can be inferred with high confidence from some previously informative points before reaching  $X$ . These points are obtained via a set  $\hat{\mathcal{S}}$  called *current active set*. Each element of  $\hat{\mathcal{S}}$  can be seen as a triplet  $(X', \hat{Y}', c)$  where  $X'$  is an informative point,  $\hat{Y}'$  its inferred label, and  $c > 0$  can be thought as

**Algorithm 1:**  $k$ -NN Active Learning under Local Smoothness (KALLS)

---

**Input:** a pool  $\mathcal{K} = \{X_1, \dots, X_w\}$ , label budget  $n$ , smoothness parameters  $(\alpha, L)$ , margin noise parameters  $(\beta, C)$ , confidence parameter  $\delta$ , accuracy parameter  $\epsilon$ .

**Output:** 1-NN classifier  $\hat{f}_{n,w}$

```

1   $s = 1$  ▷ index of point currently examined
2   $\hat{\mathcal{S}} = \emptyset$  ▷ current active set
3   $t = n$  ▷ current label budget
4   $I = \emptyset$  ▷ Set of informative point indexes (used for theoretical proofs)
5  while  $t > 0$  and  $s < w$  do
6    Let  $\delta_s = \frac{\delta}{32s^2}$ 
7     $T = \text{Reliable}(X_s, \delta_s, \alpha, L, \hat{\mathcal{S}})$ 
8    if  $T = \text{True}$  then
9       $s = s + 1$ 
10   else
11      $[\hat{Y}_s, Q_s] = \text{confidentLabel}(X_s, k(\epsilon, \delta_s), t, \delta)$ 
12      $\widehat{LB}_s = \left| \frac{1}{|Q_s|} \sum_{(X,Y) \in Q_s} Y - \frac{1}{2} \right| - b_{\delta_s, |Q_s|}$  ▷ Lower bound guarantee on
13      $|\eta(X_s) - \frac{1}{2}|$ 
14      $t = t - |Q_s|$ 
15      $I = I \cup \{s\}$ 
16     if  $\widehat{LB}_s \geq 0.1b_{\delta_s, |Q_s|}$  then
17        $\hat{\mathcal{S}} = \hat{\mathcal{S}} \cup \{(X_s, \hat{Y}_s, \widehat{LB}_s)\}$ 
18    $s = s + 1$ 
19   $\mathcal{S}_{ac} = \{(X_s, \hat{Y}_s), (X_s, \hat{Y}_s, \widehat{LB}_s) \in \hat{\mathcal{S}}\}$ 
20   $\hat{f}_{n,w} \leftarrow \text{1-NN}(\mathcal{S}_{ac})$ 

```

---

a guarantee for predicting the right label  $Y$  of  $X'$  as  $\hat{Y}'$ . Formally, we have  $O(c) \leq |\eta(X') - \frac{1}{2}|$  when  $(X', \hat{Y}', c) \in \hat{\mathcal{S}}$  and  $X'$  is relatively far from the decision boundary. If  $\text{Reliable}(X, \delta, \alpha, L, \hat{\mathcal{S}})$  outputs **True**, the point  $X$  is not considered to be informative, and  $\hat{\mathcal{S}}$  will not be updated. By convention,  $\text{Reliable}(X, \delta, \alpha, L, \emptyset)$  always returns *False*.

The inputs are the current point  $X$ , a confidence parameter  $\delta$ , the smoothness parameters  $(\alpha, L)$  from (H3), and the set  $\hat{\mathcal{S}}$  before examining the point  $X$ .

If  $|\eta(X) - \frac{1}{2}|$  entails the same confidence lower bound  $O(c)$  as that of some previous informative point  $X'$  (with  $(X', \hat{Y}', c) \in \hat{\mathcal{S}}$ ), there is a low degree of uncertainty on the label of  $X$ , and  $X$  is considered to be uninformative.

Using the assumption (H3), it suffices to have

$$\min(P_X(B(X, \rho(X', X))), P_X(B(X', \rho(X', X)))) \leq O(c^{d/\alpha}). \quad (9)$$

Because the  $P_X$  appearing in (9) are unknown, it has to be replaced by an estimate. We will show that it can be estimated with arbitrary precision and confidence using only unlabelled data from  $\mathcal{K}$ .

The **Reliable** subroutine uses **EstProb** $(X, r, \epsilon_o, 50, \delta)$  (inspired from (Kontorovich et al., 2016)) as follows:

**Algorithm 2: Reliable subroutine**


---

**Input:** an instance  $X$ , a confidence parameter  $\delta$ , smoothness parameters  $\alpha, L$ , a set  $\widehat{\mathcal{S}} \subset \mathcal{X} \times \mathcal{Y} \times \mathbb{R}^+$

**Output:**  $T$

```

1 for  $(X', Y', c) \in \widehat{\mathcal{S}}$  do
2    $\widehat{p}_{X'} = \text{EstProb}(X', \rho(X, X'), (\frac{c}{64L})^{d/\alpha}, 50, \delta)$ 
3    $\widehat{p}_X = \text{EstProb}(X, \rho(X, X'), (\frac{c}{64L})^{d/\alpha}, 50, \delta)$ 
4 if  $\exists (X', Y, c) \in \widehat{\mathcal{S}}$  such that  $(\widehat{p}_{X'} \leq \frac{75}{94} (\frac{c}{64L})^{d/\alpha} \text{ OR } \widehat{p}_X \leq \frac{75}{94} (\frac{c}{64L})^{d/\alpha})$  then
5    $T = \text{True}$ 
6 else
7    $T = \text{False}$ 

```

---

**Algorithm 3: EstProb subroutine**


---

**Input:** an instance  $x \in \mathcal{X}$ , a positive number  $r > 0$ , an accuracy parameter  $\epsilon_o$ , an integer parameter  $u$ , a confidence parameter  $\delta$

**Output:**  $\widehat{p}_X$   $\triangleright$  An estimate of  $P_X(B(x, r))$

```

1 Set  $p \sim \mathbb{1}_{B(x, r)}$  a Bernoulli variable
2  $\widehat{p}_X = \text{BerEst}(\epsilon_o, \delta, u)$   $\triangleright$  in BerEst subroutine, to draw a single  $p_i$ , randomly sample  $X_i \in \mathcal{K}$ , and set  $p_i = \mathbb{1}_{X_i \in B(x, r)}$ .

```

---

**Algorithm 4: BerEst subroutine (Bernoulli Estimation)**


---

**Input:** accuracy parameter  $\epsilon_o$ , confidence parameter  $\delta'$ , budget parameter  $u$ .  $\triangleright u$  does not depend on the label budget  $n$

**Output:**  $\widehat{p}$

```

1 Sample  $p_1, \dots, p_4$   $\triangleright$  with respect to  $\sim p$ 
2  $S = \{p_1, \dots, p_4\}$ 
3  $K = \frac{4u}{\epsilon_o} \log(\frac{8u}{\delta'\epsilon_o})$ 
4 for  $i = 3 : \log_2(u \log(2K/\delta')/\epsilon_o)$  do
5    $m = 2^i$ 
6    $S = S \cup \{p_{m/2+1}, \dots, p_m\}$ 
7    $\widehat{p} = \frac{1}{m} \sum_{j=1}^m p_j$ 
8   if  $\widehat{p} > u \log(2m/\delta')/m$  then
9     Break
10 Output  $\widehat{p}$ 

```

---

1. Call the subroutine  $\text{BerEst}(\epsilon_o, \delta, 50)$ .
2. To draw a single  $p_i$  in  $\text{BerEst}(\epsilon_o, \delta, 50)$ , sample randomly an example  $X_i$  from  $\mathcal{K}$ , and set  $p_i = \mathbb{1}_{X_i \in B(x, r)}$ .

The subroutine **BerEst** consists in estimating adaptively with high probability the expectation of a Bernoulli variable  $Z \sim p$ . In our setting, we estimate a probability-ball, so that a realisation of  $Z$  can be set as  $p_i = \mathbb{1}_{X_i \in B(x, r)}$ . The variables  $p_1, \dots, p_4$  are sampled at the beginning for theoretical analysis where we want a concentration inequality to hold for a number of samples

greater than 4 (see (Kontorovich et al., 2016; Maurer and Pontil, 2009) for more details).

However, it is not dramatic if  $P_X$  is supposed to be known by the learner. This is not a limitation, since it can be assumed that the pool  $\mathcal{K}$  of data is large enough such that  $P_X$  can be estimated to any desired accuracy.

#### 4.4 **ConfidentLabel** subroutine

If a point  $X$  is considered informative, it is introduced in the **ConfidentLabel** (Algorithm(5)), along with an integer  $k'$ , a budget parameter  $t$  and a confidence parameter  $\delta$ . This subroutine infers with high confidence (at least  $1 - \delta$ ) the label of  $X$ , by using the labels of its  $k'$  nearest neighbors, knowing that we can request at most  $t$  labels. The parameter  $k'$  is chosen such that, with high probability, the empirical majority of the  $k'$ -NN labels differs from the majority in expectation by less than some margin, and all the  $k'$ -NN are at most at some distance from  $X$ . The **ConfidentLabel** subroutine outputs  $\hat{Y}$ ,  $Q$  where  $Q$  represents the set of labeled nearest neighbors in the subroutine, and  $\hat{Y}$  represents the majority label in  $Q$ .

---

##### Algorithm 5: **confidentLabel** subroutine

---

**Input:** an instance  $X$ , integer  $k'$ , budget parameter  $t \geq 1$ , confidence parameter  $\delta$ .  
**Output:**  $\hat{Y}$ ,  $Q$

```

1  $Q = \emptyset$ 
2  $k = 1$ 
3 while  $k \leq \min(k', t)$  do
4   Request the label  $Y^{(k)}$  of  $X^{(k)}$ 
5    $Q = Q \cup \{(X^{(k)}, Y^{(k)})\}$ 
6   if  $\left| \frac{1}{k} \sum_{i=1}^k Y^{(i)} - \frac{1}{2} \right| > 2b_{\delta,k}$  then
7     exit ▷ cut-off condition
8   ▷  $b_{\delta,k}$  is defined in (4)
9    $k = k + 1$ 
10  $\hat{\eta} \leftarrow \frac{1}{|Q|} \sum_{(X,Y) \in Q} Y$ 
11  $\hat{Y} = \mathbb{1}_{\hat{\eta} \geq 1/2}$ 
```

---

## 5 Theoretical motivations

This Section provides the main results and theoretical motivations behind the KALLS algorithm. Let us recall  $\mathcal{K} = \{X_1, \dots, X_w\}$  is the pool of unlabeled data and  $n$  is the budget.

Let us denote by  $\mathcal{A}_{a,w}$  the set of active learning algorithms on  $\mathcal{K}$ , and  $\mathcal{P}(\alpha, \beta)$  the set of probabilities that satisfy assumption (H3) and (H2).

Additionally, let us introduce the set of probabilities  $\mathcal{P}'(\alpha, \beta)$  on  $\mathcal{X} \times \mathcal{Y}$ . A probability  $P \in \mathcal{P}'(\alpha, \beta)$  if  $P \in \mathcal{P}(\alpha, \beta)$  and its marginal probability  $P_X$  is a doubling-probability. For  $A \in \mathcal{A}_{a,w}$ , we denote by  $\hat{f}_{A,n,w} := \hat{f}_{n,w}$  the classifier that is provided by  $A$ .

Theorem 2 and its equivalent form in Theorem 3 are the main results of this paper. They provide bounds on the excess risk for the KALLS algorithm in terms of the set  $\mathcal{P}'(\alpha, \beta)$ . The main idea of the proof is sketched in Section 5.2, while a detailed proof can be found in Appendix A.

## 5.1 Main results

### Theorem 2 (Excess risk for the KALLS algorithm.)

Let the set  $\mathcal{P}'(\alpha, \beta)$  such that  $\alpha\beta < d$  where  $d$  is the dimension of the input space  $\mathcal{X} \subset \mathbb{R}^d$ . For  $P \in \mathcal{P}'(\alpha, \beta)$ , if  $\hat{f}_{n,w}$  is the  $1 - \text{NN}$  classifier provided by KALLS, then we have:

$$\sup_{P \in \mathcal{P}'(\alpha, \beta)} \mathbb{E}_n \left[ R(\hat{f}_{n,w}) - R(f^*) \right] \leq \tilde{O} \left( n^{-\frac{\alpha(\beta+1)}{2\alpha+d-\alpha\beta}} \right), \quad (10)$$

where  $\mathbb{E}_n$  is with respect to the randomness of the KALLS algorithm.

The result (10) is also stated below (Theorem 3) in a more practical form using label complexity. This latter form will be used in the proof.

### Theorem 3 (Label complexity for the KALLS algorithm.)

Let the set  $\mathcal{P}'(\alpha, \beta)$  such that  $\alpha\beta < d$ . Let  $\epsilon, \delta \in (0, 1)$ . For all  $n, w \in \mathbb{N}$  such that:

if

$$n \geq \tilde{O} \left( \left( \frac{1}{\epsilon} \right)^{\frac{2\alpha+d-\alpha\beta}{\alpha(\beta+1)}} \right), \quad (11)$$

$$w \geq \tilde{O} \left( \left( \frac{1}{\epsilon} \right)^{\frac{2\alpha+d}{\alpha(\beta+1)}} \right) \quad (12)$$

and

$$w \geq \frac{400 \log \left( \frac{12800w^2}{\delta \left( \frac{1}{64L} \bar{c} \phi_n \right)^{d/\alpha}} \right)}{\left( \frac{1}{64L} \bar{c} \phi_n \right)^{d/\alpha}}, \quad (13)$$

where  $L$  appears in (H3),  $\bar{c} = 0.1$  and  $\phi_n$  is defined by (7),

then with probability at least  $1 - \delta$  we have:

$$\sup_{P \in \mathcal{P}'(\alpha, \beta)} \left[ R(\hat{f}_{n,w}) - R(f^*) \right] \leq \epsilon. \quad (14)$$

Before proving this theorem, a couple of important remarks should be made:

1. The rate of convergence (10) obtained in Theorem 2 is an improvement over the passive learning counterpart. For  $P \in \mathcal{P}(\alpha, \beta)$ , if  $\hat{f}_n$  is the classifier provided by a passive learning algorithm, we have (Chaudhuri and Dasgupta, 2014):

$$\sup_{P \in \mathcal{P}(\alpha, \beta)} \mathbb{E}_n \left[ R(\hat{f}_n) - R(f^*) \right] \leq \tilde{O} \left( n^{-\frac{\alpha(\beta+1)}{2\alpha+d}} \right). \quad (15)$$

Because  $\mathcal{P}'(\alpha, \beta) \subset \mathcal{P}(\alpha, \beta)$ , we also have:

$$\sup_{P \in \mathcal{P}'(\alpha, \beta)} \mathbb{E}_n \left[ R(\hat{f}_n) - R(f^*) \right] \leq \tilde{O} \left( n^{-\frac{\alpha(\beta+1)}{2\alpha+d}} \right). \quad (16)$$

2. The rate (10) is also minimax. Indeed, let us introduce the set of probabilities  $\bar{\mathcal{P}}(\alpha, \beta)$  that satisfy the Hölder continuous assumption (H1a) (with parameter  $\alpha$ ), strong density assumption (H1b), and margin noise assumption (H2).

Let us assume that  $\alpha\beta < d$ . It was proven in (Minsker, 2012) that if  $\text{supp}(P_X) \subset [0, 1]^d$ , there exists a constant  $\gamma > 0$  such that for all  $n$  large enough and for any active classifier  $\hat{f}_n$ , we have:

$$\sup_{P \in \bar{\mathcal{P}}(\alpha, \beta)} \left[ R(\hat{f}_n) - R(f^*) \right] \geq \gamma n^{-\frac{\alpha(\beta+1)}{2\alpha+d-\alpha\beta}}. \quad (17)$$

Moreover, the strong density assumption implies the doubling-probability assumption (Kpotufe, 2011), and according to Theorem 1, the lower bound obtained in (17) is also valid for the family of probabilities  $\mathcal{P}'(\alpha, \beta)$ .

## 5.2 Proof sketch of Theorem (3)

For a classifier  $\hat{f}_{n,w}$ , it is well known (Lugosi, 2002) that the excess of risk is:

$$R(\hat{f}_{n,w}) - R(f^*) = \int_{\{x, \hat{f}_{n,w}(x) \neq f^*(x)\}} |2\eta(x) - 1| dP_X(x). \quad (18)$$

We thus aim to prove that (11) is a sufficient condition to guarantee (with probability  $\geq 1 - \delta$ ), that  $\hat{f}_{n,w}$  agrees with  $f^*$  on the set  $\{x, |\eta(x) - 1/2| > \Delta_o\}$ , for a suitable choice of  $\Delta_o > 0$ .

Introducing  $\Delta_o$  in (18) leads to:

$$R(\hat{f}_{n,w}) - R(f^*) \leq 2\Delta_o P_X(|\eta(x) - 1/2| < \Delta_o). \quad (19)$$

Therefore, if  $\Delta_o \leq \frac{\epsilon}{2}$  then we have immediately,  $R(\hat{f}_{n,w}) - R(f^*) \leq \epsilon$ . On the other hand, if  $\Delta_o > \frac{\epsilon}{2}$ , by hypothesis (H2), we have  $R(\hat{f}_{n,w}) - R(f^*) \leq 2C\Delta_o^{\beta+1}$ . In the latter case, setting  $\Delta_o = \left(\frac{\epsilon}{2C}\right)^{\frac{1}{\beta+1}}$  guarantees  $R(\hat{f}_{n,w}) - R(f^*) \leq \epsilon$ . Altogether, using for  $\Delta_o$  the value  $\Delta = \max\left(\frac{\epsilon}{2}, \left(\frac{\epsilon}{2C}\right)^{\frac{1}{\beta+1}}\right)$  guarantees  $R(\hat{f}_{n,w}) - R(f^*) \leq \epsilon$ . This explains the expression (6).

We present the proof sketch of Theorem 3 in three main steps, and refer to the corresponding Lemmas and Theorems in the Appendix A for more details.

### 1. Adaptive label requests on informative points:

We design two events  $A_1, A_2$  with  $P(A_1 \cap A_2) \geq 1 - \frac{3\delta}{16}$ , such that:

- Given an informative point  $X_s$ , if  $|\eta(X_s) - \frac{1}{2}| \geq \frac{1}{2}\Delta$  and if the budget allows ( $n = w = +\infty$ ), on  $A_1 \cap A_2$ , the cut-off condition used in Algorithm 5

$$\left| \frac{1}{k} \sum_{i=1}^k Y_s^{(i)} - \frac{1}{2} \right| \leq 2b_{\delta_s, k}$$

will be violated after at most  $\tilde{k}(\epsilon, \delta_s)$  requests, with  $\tilde{k}(\epsilon, \delta_s) \leq k(\epsilon, \delta_s)$ . Also, the label inferred after  $\tilde{k}(\epsilon, \delta_s)$  label requests corresponds to the true label  $f^*(X_s)$ . The intuition behind is to adapt the number of labels requested with respect to the noise; i.e., fewer label requests on a less noisy point (i.e.,  $|\eta(X_s) - \frac{1}{2}| \geq \frac{1}{2}\Delta$ ), and more label requests on a noisy point. This provides significant savings in the number of requests needed to predict with high probability the correct label.

- In the event  $A_1 \cap A_2$ , any informative point  $X_s$  falls in a high density region such that all the  $k(\epsilon, \delta_s)$  nearest neighbors of  $X_s$  are within at most some distance to  $X_s$ , and the condition (12) is sufficient to have  $k(\epsilon, \delta_s) \leq w$ .

### 2. Condition to be an informative point

We design an event  $A_3$  with  $P(A_3) \geq 1 - \delta/16$  and such that, on this event  $A_3$ , for  $X_s$  ( $s \leq w$ ) a point whose informativeness we want to check, and  $\hat{S}$  the version of the current active set just before reaching the point  $X_s$  in KALLS (Algorithm (1)), the following holds.

If there exists  $s' < s$  such that  $X_{s'}$  is an informative point and  $(X_{s'}, \hat{Y}_{s'}, \widehat{LB}_{s'}) \in \hat{S}$ , and that satisfies

$$\left( \hat{p}_{X_{s'}} \leq \frac{75}{94} \left( \frac{1}{64L} \widehat{LB}_{s'} \right)^{d/\alpha} \text{ or } \hat{p}_{X_s} \leq \frac{75}{94} \left( \frac{1}{64L} \widehat{LB}_{s'} \right)^{d/\alpha} \right) \quad (20)$$

where  $\hat{p}_{X_{s'}}$  and  $\hat{p}_{X_s}$  are defined in Algorithm2, then,

$$\min(P_X(B(X_s, \rho(X_s, X_{s'}))), P_X(B(X_{s'}, \rho(X_s, X_{s'})))) \leq \left( \frac{1}{64L} \widehat{LB}_{s'} \right)^{d/\alpha}. \quad (21)$$

In this case, let  $X_{s'}$  be such a point that satisfies (20) and (21), we can easily prove that when  $X_{s'}$  is relatively far from the boundary, i.e.,  $|\eta(X_{s'}) - \frac{1}{2}| \geq \frac{\Delta}{32}$ , on  $A_1 \cap A_2 \cap A_3$ , we have the lower bound guarantee

$$|\eta(X_{s'}) - \frac{1}{2}| \geq \frac{32}{63} \widehat{LB}_{s'}, \quad (22)$$

and easily deduce by using the smoothness assumption, (21) and (22), that the points  $X_s$  and  $X_{s'}$  have the same label, then we do not need to use  $X_s$  in the subroutine **ConfidentLabel** (Algorithm 5) and  $X_s$  is an uninformative point. In addition, (13) is a sufficient condition such that the number of points used in  $\text{Estprob}(X_s, \rho(X_s, X_{s'}), \left(\frac{1}{64L}\widehat{LB}_{s'}\right)^{d/\alpha}, 50, \delta_s)$  (respectively in  $\text{Estprob}(X_{s'}, \rho(X_s, X_{s'}), \left(\frac{1}{64L}\widehat{LB}_{s'}\right)^{d/\alpha}, 50, \delta_s)$ ) is lower than  $w$ .

### 3. Label the instance space and label complexity

The set  $I$  is introduced in **KALLS** (Algorithm(1)) as the set of informative points indexes. Let  $s_I = \max I$ , the index of the last informative point. For  $\epsilon, \delta \in (0, \frac{1}{2})$ ,  $\Delta$  defined in (6), and  $(\alpha, L)$  the smoothness parameters, let us introduce

$$T_{\epsilon, \delta} = \frac{1}{\tilde{p}_\epsilon} \ln\left(\frac{8}{\delta}\right), \text{ and } \tilde{p}_\epsilon = \left(\frac{\Delta}{128L}\right)^{d/\alpha}.$$

We design two events  $A_4$  and  $A_5$ , with  $P(A_4 \cap A_5) \geq 1 - \delta/4$ , such that on  $A_1 \cap A_2 \cap A_3 \cap A_4$ , if

$$s_I \geq T_{\epsilon, \delta} \quad (23)$$

and equations (13) and (12) hold, then, for all  $x \in \text{supp}(P_X)$  with  $|\eta(x) - \frac{1}{2}| > \Delta$ , we have:

$$|\eta(X_x^{(1)}) - \frac{1}{2}| \geq \frac{1}{2}\Delta \quad \text{and} \quad \hat{f}_{n,w}(x) = f^*(x) = f^*(X_x^{(1)}), \quad (24)$$

where  $X_x^{(1)}$  is the nearest neighbor of  $x$  in  $\hat{S}_{ac}$ , and  $\hat{f}_{n,w}$  the 1-NN classifier on  $\hat{S}_{ac}$ . Additionally, on  $A_1 \cap A_2 \cap A_3 \cap A_4 \cap A_5$  we prove that (11) is sufficient to obtain (23).

Finally, if (11), (12) and (13) hold simultaneously, then, on  $A_1 \cap A_2 \cap A_3 \cap A_4 \cap A_5$ , the final classifier  $\hat{f}_{n,w}$  agrees with the Bayes classifier  $f^*$  on  $\{x, |\eta(x) - 1/2| > \Delta\}$ . Thus, (14) holds with probability at least  $1 - \left(\frac{\delta}{16} + \frac{\delta}{8} + \frac{\delta}{16} + \frac{\delta}{8} + \frac{\delta}{8}\right) = 1 - \delta/2 > 1 - \delta$ .

## 6 Conclusion and future work

In this paper we first reviewed the main results for convergence rates in a non-parametric setting for active learning, with a special emphasis on the relative merits of the assumptions about the smoothness and the margin noise. Then, by putting active learning in perspective with recent work on passive learning, we provided a novel active learning algorithm with a particular smoothness assumption customized for  $k$ -NN.

We showed that our algorithm has a convergence rate comparable to state-of-the-art active learning algorithms, but using less restrictive assumptions.



This removes unnecessary restrictions on the distribution that would exclude important densities (e.g., Gaussian).

Additionally, our algorithm can readily be extended to multi-class classification, and then compared to recent results obtained in passive learning by (Reeve and Brown, 2017) which extended the work of (Chaudhuri and Dasgupta, 2014) to multi-class classification.

Finally, an important direction for further work is to extend our results to the case where the key parameters of the problem (defining smoothness and noise) are unknown. Ongoing work in this direction builds upon previous results in an adaptive setting (Locatelli et al., 2017), (Minsker, 2012), (Balcan and Hanneke, 2012), (Hanneke, 2011).

## References

- Jean-Yves Audibert and Alexandre B Tsybakov. Fast learning rates for plug-in classifiers. *The Annals of statistics*, 35(2):608–633, 2007.
- Maria Florina Balcan and Steve Hanneke. Robust interactive learning. In *Conference on Learning Theory*, pages 20–1, 2012.
- Maria-Florina Balcan, Steve Hanneke, and Jennifer Wortman Vaughan. The true sample complexity of active learning. *Machine learning*, 80(2-3):111–139, 2010.
- G rard Biau and Luc Devroye. *Lectures on the nearest neighbor method*. Springer, 2015.
- Rui M Castro and Robert D Nowak. Minimax bounds for active learning. *IEEE Transactions on Information Theory*, 54(5):2339–2353, 2008.
- Kamalika Chaudhuri and Sanjoy Dasgupta. Rates of convergence for nearest neighbor classification. In *Advances in Neural Information Processing Systems*, pages 3437–3445, 2014.
- Sanjoy Dasgupta. Two faces of active learning. *Theoretical computer science*, 412(19):1767–1781, 2011.
- Sanjoy Dasgupta. Active learning theory. *Encyclopedia of Machine Learning and Data Mining*, pages 14–19, 2017.
- Maik D r ng, L szl  Gy rfi, and Harro Walk. Rate of convergence of k-nearest-neighbor classification rule. *The Journal of Machine Learning Research*, 18(1):8485–8500, 2017.
- Gerald A Edgar. Packing measure in general metric space. *Real Analysis Exchange*, 26(2):831–852, 2000.
- Herbert Federer. *Geometric measure theory*. Springer, 2014.
- St phane Ga ffas. Sharp estimation in sup norm with random design. *Statistics & probability letters*, 77(8):782–794, 2007.
- Steve Hanneke. Nonparametric active learning, part 1: Smooth regression functions.
- Steve Hanneke. Rates of convergence in active learning. *The Annals of Statistics*, 39(1):333–361, 2011.
- Steve Hanneke and Liu Yang. Minimax analysis of active learning. *The Journal of Machine Learning Research*, 16(1):3487–3602, 2015.
- Juha Heinonen. *Lectures on analysis on metric spaces*. Springer Science & Business Media, 2012.
- Wassily Hoeffding. Probability inequalities for sums of bounded random variables. *Journal of the American Statistical Association*, 58(301):13–30, 1963.
- Emilie Kaufmann, Olivier Capp , and Aur lien Garivier. On the complexity of best-arm identification in multi-armed bandit models. *The Journal of Machine Learning Research*, 17(1):1–42, 2016.
- Aryeh Kontorovich, Sivan Sabato, and Ruth Urner. Active nearest-neighbor learning in metric spaces. In *Advances in Neural Information Processing Systems*, pages 856–864, 2016.
- Samory Kpotufe. k-nn regression adapts to local intrinsic dimension. In *Advances in Neural Information Processing Systems*, pages 729–737, 2011.

- Andrea Locatelli, Alexandra Carpentier, and Samory Kpotufe. Adaptivity to noise parameters in nonparametric active learning. *Proceedings of Machine Learning Research vol.*, 65:1–34, 2017.
- Andrea Locatelli, Alexandra Carpentier, and Samory Kpotufe. An adaptive strategy for active learning with smooth decision boundary. In *Algorithmic Learning Theory*, pages 547–571, 2018.
- Gábor Lugosi. Pattern classification and learning theory. In *Principles of nonparametric learning*, pages 1–56. Springer, 2002.
- Enno Mammen and Alexandre B Tsybakov. Smooth discrimination analysis. *The Annals of Statistics*, 27(6):1808–1829, 1999.
- Pascal Massart and Élodie Nédélec. Risk bounds for statistical learning. *The Annals of Statistics*, 34(5):2326–2366, 2006.
- Andreas Maurer and Massimiliano Pontil. Empirical bernstein bounds and sample variance penalization. *arXiv preprint arXiv:0907.3740*, 2009.
- Stanislav Minsker. Plug-in approach to active learning. *Journal of Machine Learning Research*, 13(Jan):67–90, 2012.
- Wolfgang Mulzer. Five proofs of Chernoff’s bound with applications. *arXiv preprint arXiv:1801.03365*, 2018.
- Charles Chapman Pugh. *Real mathematical analysis*. Springer, 2002.
- Henry WJ Reeve and Gavin Brown. Minimax rates for cost-sensitive learning on manifolds with approximate nearest neighbours. In *International Conference on Algorithmic Learning Theory*, pages 11–56, 2017.
- Shai Shalev-Shwartz and Shai Ben-David. *Understanding machine learning: From theory to algorithms*. Cambridge university press, 2014.
- Mathukumalli Vidyasagar. *Learning and generalisation: with applications to neural networks*. Springer Science & Business Media, 2013.

## A Detailed proof of Theorem 3

This Appendix is organized as follows: in Section A.1, we introduce some additional notations. In Section A.2 we adaptively determine the number of label requests needed to accurately predict the label of an informative point that is relatively far from the boundary decision. In Section A.4, we provide some lemmas that give a sufficient condition for a point to be informative. In Section A.5, we give theorems that allow us to classify each instance relatively far from the decision boundary. Finally in Section A.6, we provide the label complexity and establish Theorem 3.

### A.1 Notations

Some notations that will be used throughout the proofs are listed here for convenience.

As defined in Section 2.3, let  $B(x, r) = \{x' \in \mathcal{X}, \rho(x, x') < r\}$  and  $\bar{B}(x, r) = \{x' \in \mathcal{X}, \rho(x, x') \leq r\}$  the open and closed balls with respect to the Euclidean metric  $\rho$ , respectively, centered at  $x \in \mathcal{X}$  with radius  $r > 0$ . Let  $\text{supp}(P_X) = \{x \in \mathcal{X}, \forall r > 0, P_X(B(x, r)) > 0\}$  the support of the marginal distribution  $P_X$ .

For  $p \in (0, 1]$ , and  $x \in \text{supp}(P_X)$ , let us define

$$r_p(x) = \inf\{r > 0, P_X(B(x, r)) \geq p\}. \quad (25)$$

Let us recall for  $X_s \in \mathcal{K} = \{X_1, \dots, X_w\}$ , we denote by  $X_s^{(k)}$  its  $k$ -th nearest neighbor in  $\mathcal{K}$ , and  $Y_s^{(k)}$  the corresponding label.

For an integer  $k \geq 1$ , let

$$\hat{\eta}_k(X_s) = \frac{1}{k} \sum_{i=1}^k Y_s^{(i)}, \quad \bar{\eta}_k(X_s) = \frac{1}{k} \sum_{i=1}^k \eta(X_s^{(i)}). \quad (26)$$

### A.2 Adaptive label requests on informative points

#### Lemma 1 (Chernoff bounds, (Mulzer, 2018))

Suppose  $X_1, \dots, X_m$  are independent random variables taking value in  $\{0, 1\}$ . Let  $X$  denote their sum and  $\mu = E(X)$  its expected value. Then,

- For any  $\delta \in (0, 1)$ ,

$$P_m(X \leq (1 - \delta)\mu) \leq \exp(-\delta^2 \mu / 2), \quad (27)$$

where  $P_m$  is the probability with respect to the sample  $X_1, \dots, X_m$ .

- Additionally, for any  $\delta' \geq 1$ , we have:

$$P_m(X \geq (1 + \delta')\mu) \leq \exp(-\delta' \mu / 4). \quad (28)$$

#### Lemma 2 (Logarithmic relationship, (Vidyasagar, 2013))

Suppose  $a, b, c > 0$ ,  $abe^{c/a} > 4 \log_2(e)$ , and  $u \geq 1$ . Then:

$$u \geq 2c + 2a \log(ab) \Rightarrow u > c + a \log(bu).$$

#### Lemma 3 (Chaudhuri and Dasgupta, 2014)

For  $p \in (0, 1]$ , and  $x \in \text{supp}(P_X)$ , let us define  $r_p(x) = \inf\{r > 0, P_X(B(x, r)) \geq p\}$ . For all  $p \in (0, 1]$ , and  $x \in \text{supp}(P_X)$ , we have:

$$P_X(B(x, r_p(x)) \geq p.$$

**Theorem 4**

Let  $\epsilon, \delta \in (0, 1)$ . Set  $\Delta = \max(\epsilon, (\frac{\epsilon}{2C})^{\frac{1}{\beta+1}})$ , and  $p_\epsilon = \left(\frac{31\Delta}{1024L}\right)^{d/\alpha}$ , where  $\alpha, L, \beta, C$  are parameters used in (H2) and (H3).

For  $p \in (0, 1]$ , and  $x \in \text{supp}(P_X)$ , let us introduce  $r_p(x) = \inf\{r > 0, P_X(B(x, r)) \geq p\}$  and  $k_s := k(\epsilon, \delta_s)$  defined in (5) (where  $\delta_s = \frac{\delta}{32s^2}$ ).

For  $k, s \geq 1$ , set  $\tau_{k,s} = \sqrt{\frac{2}{k} \log(\frac{32s^2}{\delta})}$ . There exists an event  $A_1$  with probability at least  $1 - \frac{\delta}{16}$ , such that on  $A_1$ , for all  $1 \leq s \leq w$ , if

$$k_s \leq (1 - \tau_{k,s})p_\epsilon(w - 1) \quad (29)$$

then the  $k_s$  nearest neighbors of  $X_s$  (in the pool  $\mathcal{K}$ ) belong to the ball  $B(X_s, r_{p_\epsilon}(X_s))$ . Additionally, the condition

$$w \geq \bar{O}\left(\left(\frac{1}{\epsilon}\right)^{\frac{2\alpha+d}{\alpha(\beta+1)}}\right) \quad (30)$$

is sufficient to have (29).

*Proof*

Fix  $x \in \text{supp}(P_X)$ . For  $k \in \mathbb{N}$ , let us denote  $X_x^{(k)}$ , the  $k^{\text{th}}$  nearest neighbor of  $x$  in the pool. we have,

$$P(\rho(x, X_x^{(k_s+1)}) > r_{p_\epsilon}(x)) \leq P\left(\sum_{i=1}^w \mathbb{1}_{X_i \in B(x, r_{p_\epsilon}(x))} \leq k_s\right).$$

Then, by using Lemma 1 and Lemma 3, and if  $k_s$  satisfies (29), we have:

$$\begin{aligned} P(\rho(x, X_x^{(k_s+1)}) > r_{p_\epsilon}(x)) &\leq P\left(\sum_{i=1}^w \mathbb{1}_{X_i \in B(x, r_{p_\epsilon}(x))} \leq (1 - \tau_{k,s})p_\epsilon(w - 1)\right) \\ &\leq P\left(\sum_{i=1}^w \mathbb{1}_{X_i \in B(x, r_{p_\epsilon}(x))} \leq (1 - \tau_{k,s})P_X(B(x, r_{p_\epsilon}(x)))(w - 1)\right) \\ &\leq \exp(-\tau_{k,s}^2(w - 1)P_X(B(x, r_{p_\epsilon}(x)))/2) \\ &\leq \exp(-\tau_{k,s}^2(w - 1)p_\epsilon/2) \\ &\leq \exp(-\tau_{k,s}^2 k_s/2) \\ &\leq \exp(-\log(32s^2/\delta)) \\ &= \frac{\delta}{32s^2}. \end{aligned}$$

Fix  $x = X_s$ . Given  $X_s$ , there exists an event  $A_{1,s}$ , such that  $P(A_{1,s}) \geq 1 - \delta/(32s^2)$ , and on  $A_{1,s}$ , if

$$k_s \leq (1 - \tau_{k,s})p_\epsilon(w - 1),$$

we have  $B(X_s, r_{p_\epsilon}(X_s)) \cap \{X_1, \dots, X_w\} \geq k_s$ . By setting  $A_1 = \cap_{s \geq 1} A_{1,s}$ , we have  $P(A_1) \geq 1 - \delta/16$ , and on  $A_1$ , for all  $1 \leq s \leq w$ , if  $k_s \leq (1 - \tau_{k,s})p_\epsilon(w - 1)$ , then  $B(X_s, r_{p_\epsilon}(X_s)) \cap \{X_1, \dots, X_w\} \geq k_s$ .

Now, let us proof that the condition (30) is sufficient to guarantee (29).

The relation (29) implies

$$w \geq \frac{k_s}{(1 - \tau_{k,s})p_\epsilon} + 1. \quad (31)$$

We can see by a bit of calculus, that  $\tau_{k_s, s} \leq \frac{1}{2}$ , and then

$$\begin{aligned} \frac{k_s}{(1 - \tau_{k_s, s})p_\epsilon} + 1 &\leq \frac{2k_s}{p_\epsilon} + 1 \\ &\leq 4 \frac{k_s}{p_\epsilon} \quad \left( \text{because } \frac{k_s}{p_\epsilon} \geq 1 \right) \\ &= \frac{4c}{p_\epsilon \Delta^2} \left[ \log\left(\frac{32s^2}{\delta}\right) + \log \log\left(\frac{32s^2}{\delta}\right) + \log \log\left(\frac{512\sqrt{e}}{\Delta}\right) \right] \\ &= \frac{b}{\Delta^{2+\frac{d}{\alpha}}} \left[ \log\left(\frac{32s^2}{\delta}\right) + \log \log\left(\frac{32s^2}{\delta}\right) + \log \log\left(\frac{512\sqrt{e}}{\Delta}\right) \right], \end{aligned}$$

where  $b = 4c \left( \frac{1024L}{31} \right)^{d/\alpha}$ .

$$\begin{aligned} \frac{k_s}{(1 - \tau_{k_s, s})p_\epsilon} + 1 &\leq \bar{C} \left( \frac{1}{\epsilon} \right)^{\frac{2\alpha+d}{\alpha(\beta+1)}} \left[ \log\left(\frac{32s^2}{\delta}\right) + \log \log\left(\frac{32s^2}{\delta}\right) + \log \log\left(\frac{512\sqrt{e}}{\Delta}\right) \right] \\ &\quad \text{as } \Delta = \max(\epsilon, \left( \frac{\epsilon}{2C} \right)^{\frac{1}{\beta+1}}), \text{ where } \bar{C} = b(2C)^{\frac{2\alpha+d}{\alpha(\beta+1)}} \\ &\leq \bar{C} \left( \frac{1}{\epsilon} \right)^{\frac{2\alpha+d}{\alpha(\beta+1)}} \left[ 2 \log\left(\frac{32s^2}{\delta}\right) + \log\left(\frac{512\sqrt{e}}{\epsilon}\right) \right] \\ &\quad \text{as } \log(x) \leq x, \text{ and } \Delta \geq \epsilon \\ &\leq 2\bar{C} \left( \frac{1}{\epsilon} \right)^{\frac{2\alpha+d}{\alpha(\beta+1)}} \left[ \log(s^2) + \log\left(\frac{16384\sqrt{e}}{\delta\epsilon}\right) \right] \\ &\leq 4\bar{C} \left( \frac{1}{\epsilon} \right)^{\frac{2\alpha+d}{\alpha(\beta+1)}} \left[ \log(s) + \log\left(\frac{16384\sqrt{e}}{\delta\epsilon}\right) \right] \\ &\leq 4\bar{C} \left( \frac{1}{\epsilon} \right)^{\frac{2\alpha+d}{\alpha(\beta+1)}} \left[ \log(w) + \log\left(\frac{16384\sqrt{e}}{\delta\epsilon}\right) \right]. \end{aligned} \tag{32}$$

Now, we are going to apply the Lemma 2. If we set in Lemma 2

$$a = 4\bar{C} \left( \frac{1}{\epsilon} \right)^{\frac{2\alpha+d}{\alpha(\beta+1)}}, \quad c = 4\bar{C} \left( \frac{1}{\epsilon} \right)^{\frac{2\alpha+d}{\alpha(\beta+1)}} \log\left(\frac{16384\sqrt{e}}{\delta\epsilon}\right), \quad b = 1$$

we can easily see that  $c \geq a$ ,  $a \geq 4$  and then

$$abe^{c/a} \geq 4e > \log_2(e).$$

Then, the relation

$$w \geq 4\bar{C} \left( \frac{1}{\epsilon} \right)^{\frac{2\alpha+d}{\alpha(\beta+1)}} \left( \log\left(\frac{16384\sqrt{e}}{\delta\epsilon}\right) + \log\left(4\bar{C} \left( \frac{1}{\epsilon} \right)^{\frac{2\alpha+d}{\alpha(\beta+1)}}\right) \right)$$

is sufficient to guarantee (32).

Let us note that the guarantee obtained in the preceding theorem corresponds to that obtained in passive setting ( $w = n$ ).

A.3 Motivation for choosing  $k_s$  for  $X_s$ **Lemma 4 (Hoeffding's inequality, (Hoeffding, 1963))**

– **First version:**

Let  $X$  be a random variable with  $E(X) = 0$ ,  $a \leq X \leq b$ , then for  $v > 0$ ,

$$E(e^{vX}) \leq e^{v^2(b-a)^2/8}.$$

– **Second version:**

Let  $X_1, \dots, X_m$  be independent random variables such that  $-1 \leq X_i \leq 1$ , ( $i = 0, \dots, m$ ). We define the empirical mean of these variables by

$$\bar{X} = \frac{1}{m} \sum_{i=1}^m X_i.$$

Then we have:

$$P(|\bar{X} - E(\bar{X})| \geq t) \leq \exp(-mt^2/2)$$

**Lemma 5 (Kaufmann et al., 2016)**

Let  $\zeta(u) = \sum_{k \geq 1} k^{-u}$ . Let  $X_1, X_2, \dots$  be independent random variables, identically distributed, such that, for all  $v > 0$ ,  $E(e^{vX_1}) \leq e^{v^2\sigma^2/2}$ . For every positive integer  $t$ , let  $S_t = X_1 + \dots + X_t$ . Then, for all  $\gamma > 1$  and  $r \geq \frac{8}{(e-1)^2}$ :

$$P\left(\bigcup_{t \in \mathbb{N}^*} \left\{|S_t| > \sqrt{2\sigma^2 t(r + \gamma \log \log(et))}\right\}\right) \leq \sqrt{e}\zeta\left(\gamma\left(1 - \frac{1}{2r}\right)\right)\left(\frac{\sqrt{r}}{2\sqrt{2}} + 1\right)^\gamma \exp(-r).$$

**Lemma 6**

Let  $m \geq 1$  and  $u \geq 20$ . Then we have:

$$m \geq 2u \log(\log(u)) \implies m \geq u \log(\log(m)).$$

*Proof*

Define  $\phi(m) = m - u \log(\log(m))$ , and let  $m_0 = 2u \log(\log(u))$ . We have:

$$\begin{aligned} \phi(m_0) &= 2u \log(\log(u)) - u(\log(\log(2u \log(\log(u)))) \\ &= 2u \log(\log(u)) - u \log(\log(2u) + \log(\log(\log(u)))) \end{aligned}$$

It can be shown numerically that  $\phi(m_0) \geq 0$  for  $u \geq 20$ .

Also, we have:  $\phi'(m) = \frac{m \log(m) - u}{m \log(m)} \geq 0$  for all  $m \geq m_0$  (notice that  $m_0 \geq u$  for  $u \geq 20$ ).

Then it is easy to see that  $\phi(m) \geq \phi(m_0)$  for all  $m \geq m_0$ . This establishes the lemma.

**Theorem 5**

Let  $\delta \in (0, 1)$ , and  $\epsilon \in (0, 1)$ . Let us assume that  $w$  satisfies (12). For  $X_s$ , set  $\tilde{k}(\epsilon, \delta_s)$  (with  $\delta_s = \frac{\delta}{32s^2}$ ) as

$$\tilde{k}(\epsilon, \delta_s) = \frac{c}{4|\eta(X_s) - \frac{1}{2}|^2} \left[ \log\left(\frac{32s^2}{\delta}\right) + \log \log\left(\frac{32s^2}{\delta}\right) + \log \log\left(\frac{256\sqrt{e}}{|\eta(X_s) - \frac{1}{2}|}\right) \right],$$

where  $c \geq 7.10^6$ . For  $k \geq 1$ ,  $s \leq w$ , let  $\Delta = \max(\frac{\epsilon}{2}, (\frac{\epsilon}{2C})^{\frac{1}{\beta+1}})$  and  $b_{\delta_s, k}$  defined in (4). Then, there exists an event  $A_2$ , such that  $P(A_2) \geq 1 - \delta/8$ , and on  $A_1 \cap A_2$ , we have:

1. For  $k \geq 1$ ,  $\hat{\eta}_k(X_s)$  and  $\bar{\eta}_k(X_s)$  defined in (26), for all  $s \in \{1, \dots, w\}$ ,

$$|\hat{\eta}_k(X_s) - \bar{\eta}_k(X_s)| \leq b_{\delta_s, k}. \quad (33)$$

2. For all  $s \leq w$ , if  $|\eta(X_s) - \frac{1}{2}| \geq \frac{1}{2}\Delta$ , then,  $\tilde{k}(\epsilon, \delta_s) \leq k(\epsilon, \delta_s)$ , and the subroutine  $\text{ConfidentLabel}(X_s) := \text{ConfidentLabel}(X_s, k(\epsilon, \delta_s), t = \infty, \delta_s)$  uses at most  $\tilde{k}(\epsilon, \delta_s)$  label requests. We also have

$$\left| \frac{1}{\bar{k}_s} \sum_{i=1}^{\bar{k}_s} Y_s^{(i)} - \frac{1}{2} \right| \geq 2b_{\delta_s, \bar{k}_s} \quad (34)$$

and

$$f^*(X_s) = \mathbb{1}_{\hat{\eta}_{\bar{k}_s}(X_s) \geq \frac{1}{2}}, \quad (35)$$

Where  $\bar{k}_s$  is the number of requests made in  $\text{ConfidentLabel}(X_s)$ .

*Proof*

1. Let us begin with the proof of the first part of Theorem 5.

Here, we follow the proof of Theorem 8 in (Kaufmann et al., 2016), with few additional modifications.

Let  $s \in \{1, \dots, w\}$ . Set  $S_k = \sum_{i=1}^k (Y_s^{(i)} - \eta(X_s^{(i)}))$ . Given  $\{X_1, \dots, X_w\}$ ,  $E(Y_s^{(k)} - \eta(X_s^{(k)})) = 0$ , and the random variables  $\{Y_s^{(i)} - \eta(X_s^{(i)}), i = 1, \dots, k\}$  are independent. Then by Lemma 4, given  $\{X_1, \dots, X_w\}$ , as  $Y_s^{(1)} - \eta(X_s^{(1)})$  takes values in  $[-1, 1]$ , we have  $E(e^{v(Y_s^{(1)} - \eta(X_s^{(1)}))}) \leq e^{v^2/2}$  for all  $v > 0$ . Furthermore, set  $z = \log(\frac{32s^2}{\delta})$ , and  $r = z + 3\log(z)$ . We have  $r \geq \frac{8}{(e-1)^2}$ , and by Lemma 5, with  $\gamma = 3/2$ , we have:

$$\begin{aligned} P\left(\bigcup_{k \in \mathbb{N}^*} \left\{|S_k| > \sqrt{2k(r + \gamma \log \log(ek))}\right\}\right) &\leq \sqrt{e}\zeta(3/2)(1 - \frac{1}{2r})\left(\frac{\sqrt{r}}{2\sqrt{2}} + 1\right)^{3/2} \exp(-r) \\ &= \frac{\sqrt{e}}{8}\zeta\left(\frac{3}{2} - \frac{3}{4(z + 3\log(z))}\right) \frac{(\sqrt{z + 3\log(z)} + \sqrt{8})^{3/2}}{z^3} \frac{\delta}{32s^2} \end{aligned}$$

It can be shown numerically that for  $z \geq 2.03$ , which holds for all  $\delta \in (0, 1)$ ,  $s \geq 1$ ,

$$\frac{\sqrt{e}}{8}\zeta\left(\frac{3}{2} - \frac{3}{4(z + 3\log(z))}\right) \frac{(\sqrt{z + 3\log(z)} + \sqrt{8})^{3/2}}{z^3} \leq 1.$$

Then, we have, given  $s \in \{1, \dots, w\}$ , there exists an event  $A'_{2,s}$  such that  $P(A'_{2,s}) \geq 1 - \delta/32s^2$ , and simultaneously for all  $k \geq 1$ , we have:

$$|S_k| \leq \sqrt{2k \left( \log\left(\frac{32s^2}{\delta}\right) + \log \log\left(\frac{32s^2}{\delta}\right) + \log \log(ek) \right)}.$$

By setting  $A'_2 = \cap_{s \geq 1} A'_{2,s}$ , we have  $P(A'_2) \geq 1 - \delta/16$ , and on  $A'_2$ , we have for all  $s \in \{1, \dots, w\}$ , for all  $k \geq 1$ ,

$$|\hat{\eta}_k(X_s) - \bar{\eta}_k(X_s)| \leq b_{\delta_s, k}.$$

2. For the proof of the second part of Theorem 5, we are going to show that there exists an event  $A'_2$  such that (34) and (35) hold on  $A'_2 \cap A''_2 \cap A_1$ . Given  $\{X_1, \dots, X_w\}$ , and  $X_s \in \{X_1, \dots, X_w\}$ , by Lemma 4, there exists an event  $A''_{2,s}$ , with  $P(A''_{2,s}) \geq 1 - \delta/32s^2$ , and on  $A''_{2,s}$ , we have:

$$|\hat{\eta}_k(X_s) - \bar{\eta}_k(X_s)| \leq \sqrt{\frac{2 \log(\frac{32s^2}{\delta})}{k}}.$$



This implies that:

$$|\widehat{\eta}_k(X_s) - \frac{1}{2}| \geq |\bar{\eta}_k(X_s) - \frac{1}{2}| - \sqrt{\frac{2 \log(\frac{32s^2}{\delta})}{k}}. \quad (36)$$

On the event  $A_1$ , we have, for all  $k \leq k_s$ , by the  $\alpha$ -smoothness assumption (H3),

$$|\eta(X_s) - \eta(X_s^{(k)})| \leq \frac{31}{1024} \Delta. \quad (37)$$

And then, if  $|\eta(X_s) - \frac{1}{2}| \geq \frac{1}{2} \Delta$ , then  $|\eta(X_s) - \frac{1}{2}| \geq \frac{1}{32} \Delta$ . The relation (37) becomes

$$|\eta(X_s^{(k)}) - \frac{1}{2}| \geq \frac{1}{1024} |\eta(X_s) - \frac{1}{2}|.$$

Then (36) becomes:

$$|\widehat{\eta}_k(X_s) - \frac{1}{2}| \geq \frac{1}{1024} |\eta(X_s) - \frac{1}{2}| - \sqrt{\frac{2 \log(\frac{32s^2}{\delta})}{k}}. \quad (38)$$

A sufficient condition for  $k$  to satisfy (34), is

$$\frac{1}{1024} |\eta(X_s) - \frac{1}{2}| - \sqrt{\frac{2 \log(\frac{32s^2}{\delta})}{k}} \geq 2b_{\delta_s, k}$$

and then:

$$\frac{1}{1024} |\eta(X_s) - \frac{1}{2}| - \sqrt{\frac{2 \log(\frac{32s^2}{\delta})}{k}} \geq 2\sqrt{\frac{2}{k} \left( \log\left(\frac{32s^2}{\delta}\right) + \log \log\left(\frac{32s^2}{\delta}\right) + \log \log(ek) \right)}$$

this implies:

$$k \geq \frac{1024}{|\eta(X_s) - \frac{1}{2}|^2} \left( \sqrt{2 \log\left(\frac{32s^2}{\delta}\right)} + 2\sqrt{2 \left( \log\left(\frac{32s^2}{\delta}\right) + \log \log\left(\frac{32s^2}{\delta}\right) + \log \log(ek) \right)} \right)^2. \quad (39)$$

On the other hand, the right-hand side is smaller than:

$$\frac{1024}{|\eta(X_s) - \frac{1}{2}|^2} \left( \sqrt{2 \log\left(\frac{32s^2}{\delta}\right)} + 2\sqrt{2 \log\left(\frac{32s^2}{\delta}\right)} + 2\sqrt{2 \log \log\left(\frac{32s^2}{\delta}\right)} + 2\sqrt{2 \log \log(ek)} \right)^2.$$

To deduce (39), it suffices to have the expression into brackets lower than:

$$\frac{\sqrt{k}}{32} |\eta(X_s) - \frac{1}{2}|.$$

Then, it suffices to have simultaneously:

$$\begin{aligned} \sqrt{2 \log\left(\frac{32s^2}{\delta}\right)} &\leq \frac{1}{9} \frac{\sqrt{k}}{32} |\eta(X_s) - \frac{1}{2}| \\ \sqrt{2 \log \log\left(\frac{32s^2}{\delta}\right)} &\leq \frac{1}{6} \frac{\sqrt{k}}{32} |\eta(X_s) - \frac{1}{2}| \\ \sqrt{2 \log \log(ek)} &\leq \frac{1}{6} \frac{\sqrt{k}}{32} |\eta(X_s) - \frac{1}{2}| \end{aligned}$$

Equivalently, we have:

$$k \geq \frac{1024}{|\eta(X_s) - \frac{1}{2}|^2} 162 \log\left(\frac{32s^2}{\delta}\right) \quad (40)$$

$$k \geq \frac{1024}{|\eta(X_s) - \frac{1}{2}|^2} 72 \log \log\left(\frac{32s^2}{\delta}\right) \quad (41)$$

$$k \geq \frac{1024}{|\eta(X_s) - \frac{1}{2}|^2} 72 \log \log(ek) \quad (42)$$

We can apply the Lemma 6 in (42) by taking:  $m = ek$  and  $u = \frac{73728e}{|\eta(X_s) - \frac{1}{2}|^2}$ . We have  $m \geq 1$  and  $u \geq 20$  and then, a sufficient condition to have (42) is:

$$k \geq 2 \frac{73728e}{|\eta(X_s) - \frac{1}{2}|^2} \log \log\left(\frac{73728e}{|\eta(X_s) - \frac{1}{2}|^2}\right)$$

or

$$k \geq 4 \frac{73728e}{|\eta(X_s) - \frac{1}{2}|^2} \log \log\left(\frac{\sqrt{73728e}}{|\eta(X_s) - \frac{1}{2}|}\right) \quad (43)$$

We can easily see that  $\tilde{k}_s := \tilde{k}(\epsilon, \delta_s)$  satisfies (40), (41), (43). Then

$$\left| \frac{1}{\tilde{k}_s} \sum_{i=1}^{\tilde{k}_s} Y_s^{(i)} - \frac{1}{2} \right| \geq 2b_{\delta_s, \tilde{k}_s}. \quad (44)$$

As  $|\eta(X_s) - \frac{1}{2}| \geq \frac{1}{2}\Delta$ , we can easily see that  $\tilde{k}(\epsilon, \delta_s) \leq k(\epsilon, \delta_s)$ . By taking the minimum value  $\bar{k}_s = \tilde{k}(\epsilon, \delta_s)$  that satisfies (44), we can see that when the budget allows us, the subroutine **ConfidentLabel** requests  $\bar{k}_s$  labels, and we have:

$$\left| \frac{1}{\bar{k}_s} \sum_{i=1}^{\bar{k}_s} Y_s^i - \frac{1}{2} \right| \geq 2b_{\delta_s, \bar{k}_s}. \quad (45)$$

By setting  $A'_2 = \cap_{s \geq 1} A'_{2,s}$ , we have  $P(A'_2) \geq 1 - \delta/16$ , and we can deduce (34).

We have on  $A'_2$ , for all  $s \leq w$ ,  $k \leq k(\epsilon, \delta_s)$ ,

$$|\hat{\eta}(X_s) - \bar{\eta}_k(X_s)| \leq b_{\delta_s, k}.$$

And then, on  $A_1 \cap A'_2$ , we have for all  $s \leq w$ ,  $k \leq k(\epsilon, \delta_s)$  :

$$\begin{aligned} |\eta(X_s) - \hat{\eta}_k(X_s)| &\leq |\eta(X_s) - \bar{\eta}_k(X_s)| + |\bar{\eta}_k(X_s) - \hat{\eta}_k(X_s)| \\ &\leq \frac{31}{1024} \Delta + b_{\delta_s, k}. \end{aligned} \quad (46)$$

Assume without loss of generality that  $\eta(X_s) \geq \frac{1}{2}$ , which leads to:

$$\begin{aligned} \hat{\eta}_{\bar{k}_s}(X_s) - \frac{1}{2} &= \hat{\eta}_{\bar{k}_s}(X_s) - \eta(X_s) + \eta(X_s) - \frac{1}{2} \\ &\geq -|\hat{\eta}_{\bar{k}_s}(X_s) - \eta(X_s)| + \eta(X_s) - \frac{1}{2}. \end{aligned} \quad (47)$$

If  $\eta(X_s) - \frac{1}{2} \geq \frac{1}{2}\Delta$ , with (46), the expression (47) becomes:

$$\begin{aligned}
\widehat{\eta}_{\bar{k}_s}(X_s) - \frac{1}{2} &\geq -\frac{31}{1024}\Delta - b_{\delta_s, \bar{k}_s} + \frac{1}{2}\Delta \\
&= \frac{481}{1024}\Delta - b_{\delta_s, \bar{k}_s} \\
&\geq -b_{\delta_s, \bar{k}_s}
\end{aligned} \tag{48}$$

On the other hand, we have by (34),

$$|\widehat{\eta}_{\bar{k}_s}(X_s) - \frac{1}{2}| \geq 2b_{\delta_s, \bar{k}_s},$$

that is to say:

$$\widehat{\eta}_{\bar{k}_s}(X_s) - \frac{1}{2} \geq 2b_{\delta_s, \bar{k}_s} \quad \text{or} \quad \widehat{\eta}_{\bar{k}_s}(X_s) - \frac{1}{2} \leq -2b_{\delta_s, \bar{k}_s}.$$

By (48), we have necessarily  $\widehat{\eta}_{\bar{k}_s}(X_s) - \frac{1}{2} \geq 2b_{\delta_s, \bar{k}_s}$ , and then:

$$\widehat{\eta}_{\bar{k}_s} - \frac{1}{2} \geq \max(-b_{\delta_s, \bar{k}_s}, 2b_{\delta_s, \bar{k}_s}) = 2b_{\delta_s, \bar{k}_s} \geq 0,$$

Thus we can easily deduce (35).

By setting  $A_2 = A'_2 \cap A''_2$ , we have  $P(A_2) \geq 1 - \delta/8$  and on  $A_1 \cap A_2$ , the item 1 and item 2 hold simultaneously.

#### A.4 Sufficient condition to be an informative point

As noticed in Section 4.3, a sufficient condition for a point  $X_t$  (with  $t \leq w$ ) to be considered as *not* informative is:

$$\min(P_X(B(X_t, \rho(X_t, X_s))), P_X(B(X_s, \rho(X_t, X_s)))) \leq O((\widehat{LB}_s)^{d/\alpha}). \tag{49}$$

for some previous informative point  $X_s$  (with  $(X_s, \widehat{Y}_s, \widehat{LB}_s) \in \widehat{S}$  the *current active set* just before attaining  $X_t$  in  $\text{KALLS}(\text{Algorithm}(1))$ ). Because  $P_X$  is unknown, we provide a computational scheme sufficient to obtain (49).

Firstly we follow the general procedure used in (Kontorovich et al., 2016) to estimate adaptively the expectation of a Bernoulli random variable. And secondly, we apply it to the Bernoulli variable  $\mathbb{1}_A$  where  $A = \{x, x \in B(X, r)\}$  for  $r > 0$  and  $X \in \mathcal{X}$ .

**Lemma 7** (Kontorovich et al., 2016)

Let  $\delta' \in (0, 1)$ ,  $\epsilon_o > 0$ ,  $t \geq 7$  and set  $g(t) = 1 + \frac{8}{3t} + \sqrt{\frac{2}{t}}$ . Let  $p_1, p_2, \dots \in \{0, 1\}$  be i.i.d Bernoulli random variables with expectation  $p$ . Let  $\widehat{p}$  be the output of  $\text{BerEst}(\epsilon_o, \delta', t)$ . There exists an event  $A'$ , such that  $P(A') \geq 1 - \delta'$ , and on  $A'$ , we have:

1. If  $\widehat{p} \leq \frac{\epsilon_o}{g(t)}$  then  $p \leq \epsilon_o$ , otherwise, we have  $p \geq \frac{2-g(t)}{g(t)}\epsilon_o$ .
2. The number of random draws in the  $\text{BerEst}$  subroutine (Algorithm 4) is at most  $\frac{8t \log(\frac{8t}{\delta' \psi})}{\psi}$ , where  $\psi := \max(\epsilon_o, \frac{p}{g(t)})$ .

**Lemma 8**

Let  $\epsilon, \delta \in (0, 1)$ ,  $r > 0$ . Let us assume that  $w$  satisfies (13).

There exists an event  $A_3$ , such that  $P(A_3) \geq 1 - \delta/16$ , we have, on  $A_3$ , for all  $s \leq w$ :

If there exists  $1 \leq s' < s$ , such that  $X_{s'}$  is an informative point, and  $(X_{s'}, \widehat{Y}_{s'}, \widehat{LB}_{s'}) \in \widehat{S}$  (the current active set just before attaining  $X_s$  defined in  $\text{KALLS}(\text{Algorithm}(1))$ ), and that satisfies:

$$\left( \widehat{p}_{X_{s'}} \leq \frac{75}{94} \left( \frac{1}{64L} \widehat{LB}_{s'} \right)^{d/\alpha} \quad \text{or} \quad \widehat{p}_{X_s} \leq \frac{75}{94} \left( \frac{1}{64L} \widehat{LB}_{s'} \right)^{d/\alpha} \right) \tag{50}$$

where

$$\widehat{p}_{X_{s'}} := \text{Estprob}(X_{s'}, \rho(X_s, X_{s'}), \left(\frac{1}{64L} \widehat{LB}_{s'}\right)^{d/\alpha}, 50, \delta_s)$$

and

$$\widehat{p}_{X_s} := \text{Estprob}(X_s, \rho(X_s, X_{s'}), \left(\frac{1}{64L} \widehat{LB}_{s'}\right)^{d/\alpha}, 50, \delta_s)$$

then

$$\min(P_X(B(X_s, \rho(X_{s'}, X_s))), P_X(B(X_{s'}, \rho(X_{s'}, X_s)))) \leq \left(\frac{1}{64L} \widehat{LB}_{s'}\right)^{d/\alpha}. \quad (51)$$

Otherwise, if (50) does not hold, i.e:

$$\min(\widehat{p}_{X_{s'}}, \widehat{p}_{X_s}) > \frac{75}{94} \left(\frac{1}{64L} \widehat{LB}_{s'}\right)^{d/\alpha},$$

then

$$\min(P_X(B(X_s, \rho(X_{s'}, X_s))), P_X(B(X_{s'}, \rho(X_{s'}, X_s)))) \geq \frac{28}{47} \left(\frac{1}{64L} \widehat{LB}_{s'}\right)^{d/\alpha}. \quad (52)$$

*Proof*

By following the scheme of subroutine **Estprob**, this Lemma is a direct application of Lemma 7 by taking for all  $s \leq w$ ,  $t = 50$ ,  $\epsilon_o = \left(\frac{1}{64L} \widehat{LB}_{s'}\right)^{d/\alpha}$ ,  $\delta' = \delta_s$ ,  $r = \rho(X_s, X_{s'})$ ,  $A_{3,s} := A'$ . And then, if we set  $A_3 = \bigcap_{s \geq 1} A_{3,s}$ , we have  $P(A_3) \geq 1 - \delta/16$ , and on the event  $A_3$ , we can easily deduce (51) and (52) in each cases.

On the other hand, for all  $s \leq w$ , the number of draws in  $\text{Estprob}(X_s, \rho(X_s, X_{s'}), \left(\frac{1}{64L} \widehat{LB}_{s'}\right)^{d/\alpha}, 50, \delta_s)$  (respectively  $\text{Estprob}(X_{s'}, \rho(X_s, X_{s'}), \left(\frac{1}{64L} \widehat{LB}_{s'}\right)^{d/\alpha}, 50, \delta_s)$ ) is always lower than  $w$ . Indeed, by Lemma 7, the number of draws is at most:

$$N := \frac{400 \log\left(\frac{12800s^2}{\delta\psi}\right)}{\psi} \quad \text{where} \quad \psi = \max\left(\left(\frac{1}{64L} \widehat{LB}_{s'}\right)^{d/\alpha}, \frac{75}{94} P_X(B(X_s, \rho(X_s, X_{s'})))\right).$$

Then we have:

$$\begin{aligned} N &\leq \frac{400 \log\left(\frac{12800s^2}{\delta\left(\frac{1}{64L} \widehat{LB}_{s'}\right)^{d/\alpha}}\right)}{\left(\frac{1}{64L} \widehat{LB}_{s'}\right)^{\frac{d}{\alpha}}} \\ &\leq \frac{400 \log\left(\frac{12800s^2}{\delta\left(\frac{1}{64L} \bar{c} b_{\delta_{s'}, |Q_{s'}|}\right)^{d/\alpha}}\right)}{\left(\frac{1}{64L} \bar{c} b_{\delta_{s'}, |Q_{s'}|}\right)^{d/\alpha}} \quad (\text{as } \widehat{LB}_{s'} \geq \bar{c} b_{\delta_{s'}, |Q_{s'}|}, \text{ with } \bar{c} = 0.1) \quad (53) \\ &\leq \frac{400 \log\left(\frac{12800w^2}{\delta\left(\frac{1}{64L} \bar{c} \phi_n\right)^{d/\alpha}}\right)}{\left(\frac{1}{64L} \bar{c} \phi_n\right)^{d/\alpha}} \quad (\text{we can easily see that } b_{\delta_{s'}, |Q_{s'}|} \geq \phi_n) \\ &\leq w \quad (\text{by (13)}). \end{aligned}$$

In equation (53),  $b_{\delta_{s'}, |Q_{s'}|}$  is defined by (4), and  $|Q_{s'}|$  represents the number of label requests used in the subroutine **ConfidentLabel**(Algorithm (5)) at the stage  $s'$ .

## A.5 Label the instance space

**Theorem 6**

Let  $\epsilon, \delta \in (0, 1)$ . Let

$$T_{\epsilon, \delta} = \frac{1}{\tilde{p}_\epsilon} \log\left(\frac{8}{\delta}\right), \text{ and } \tilde{p}_\epsilon = \left(\frac{\Delta}{128L}\right)^{d/\alpha}, \text{ with } \Delta = \max\left(\frac{\epsilon}{2}, \left(\frac{\epsilon}{2C}\right)^{\frac{1}{\beta+1}}\right) \quad (54)$$

Let  $I$  the set of indexes of informative points used in KALLS (Algorithm 1). Let us consider its last update in KALLS (Algorithm 1) and also denoted it by  $I$ .

Then, set  $s_I = \max I$  the index of the last informative point. Let  $\hat{S}_{ac}$  be the active set obtained in KALLS (Algorithm 1) and denote by  $\hat{f}_{n,w}$  the output  $\text{INN}(\hat{S}_{ac})$ . There exists an event  $A_4$  such that  $P(A_4) \geq 1 - \delta/8$ , and on  $A_1 \cap A_2 \cap A_3 \cap A_4$ , we have

$$1. \quad \sup_{x \in \text{supp}(P_X)} \min_{\bar{X} \in \{X_1, \dots, X_{T_{\epsilon, \delta}}\}} P_X(B(x, \rho(\bar{X}, x))) \leq \tilde{p}_\epsilon. \quad (55)$$

2. If  $w$  satisfies (12) and (13) and the following condition holds

$$s_I \geq T_{\epsilon, \delta}, \quad (56)$$

then, for all  $x \in \text{supp}(P_X)$  such that  $|\eta(x) - \frac{1}{2}| > \Delta$ , there exists  $s := s(x) \in I$  such that:

$$|\eta(X_s) - \frac{1}{2}| \geq \frac{1}{2}\Delta \quad (57)$$

and

$$f^*(x) = f^*(X_s). \quad (58)$$

In addition, we have

$$\hat{f}_{n,w}(x) = f^*(x). \quad (59)$$

*Proof*

This proof is based on results from (Hanneke) with few additional modifications.

1. Let us begin by proving the first part of Theorem 6.

For  $x \in \text{supp}(P_X)$ , let us introduce

$$r_{\tilde{p}_\epsilon}(x) = \inf\{r > 0, P_X(B(x, r)) \geq \tilde{p}_\epsilon\}.$$

By Lemma 3, we have  $P_X(B(x, r_{\tilde{p}_\epsilon}(x))) \geq \tilde{p}_\epsilon$ . Then each  $\bar{X} \in \{X_1, \dots, X_{T_{\epsilon, \delta}}\}$  belongs to  $B(x, r_{\tilde{p}_\epsilon}(x))$  with probability at least  $\tilde{p}_\epsilon$ . If we denote  $\hat{P}$  the probability over the data, we have:

$$\begin{aligned} & \hat{P}(\exists \bar{X} \in \{X_1, \dots, X_{T_{\epsilon, \delta}}\}, P_X(B(x, \rho(x, \bar{X}))) \leq \tilde{p}_\epsilon) \\ &= 1 - \hat{P}(\forall \bar{X} \in \{X_1, \dots, X_{T_{\epsilon, \delta}}\}, P_X(B(x, \rho(x, \bar{X}))) > \tilde{p}_\epsilon) \\ &= 1 - \prod_{i=1}^{T_{\epsilon, \delta}} \hat{P}(P_X(B(x, \rho(x, X_i))) > \tilde{p}_\epsilon) \\ &\geq 1 - \prod_{i=1}^{T_{\epsilon, \delta}} \hat{P}(\rho(x, X_i) > r_{\tilde{p}_\epsilon}(x)) \\ &= 1 - \prod_{i=1}^{T_{\epsilon, \delta}} (1 - \hat{P}(\rho(x, X_i) \leq r_{\tilde{p}_\epsilon}(x))) \\ &\geq 1 - (1 - \tilde{p}_\epsilon)^{T_{\epsilon, \delta}} \\ &\geq 1 - \exp(-T_{\epsilon, \delta} \tilde{p}_\epsilon) \\ &= 1 - \delta/8. \end{aligned}$$

Then, there exists an event  $A_4$ , such that  $P(A_4) \geq 1 - \delta/8$  and (55) holds on  $A_4$ . And then, we can easily conclude the first part.

2. For the second part of Theorem 6, let  $x \in \text{supp}(P_X)$ . By (55), on  $A_4$  there exists  $X_x \in \{X_1, \dots, X_{T_{\epsilon, \delta}}\}$  such that:

$$P_X(B(x, \rho(X_x, x))) \leq \tilde{p}_\epsilon. \quad (60)$$

By assumption (H3), we have:

$$|\eta(x) - \eta(X_x)| \leq \frac{1}{128} \Delta < \frac{1}{32} \Delta. \quad (61)$$

Then if  $|\eta(x) - \frac{1}{2}| > \Delta$ , we have:

$$(1 - \frac{1}{32})\Delta < |\eta(X_x) - \frac{1}{2}| < (1 + \frac{1}{32})\Delta. \quad (62)$$

As  $s_I \geq T_{\epsilon, \delta}$ , then there exists  $s'$  such that  $X_x := X_{s'}$  and  $X_{s'}$  passes through the subroutine **Reliable**.

We have two cases:

- a)  $X_{s'}$  is **uninformative**. Then there exists  $s < s'$ , such that  $X_s$  is an informative point, and

$$\widehat{LB}_s \geq 0.1b_{\delta_s, |Q_s|} \text{ and } \min(\widehat{p}_{X_s}, \widehat{p}_{X_{s'}}) \leq \frac{75}{94} \left( \frac{1}{64L} \widehat{LB}_s \right)^{d/\alpha}$$

where  $\widehat{p}_{X_s} := \text{Estprob}(X_s, \rho(X_s, X_{s'}), \left( \frac{1}{64L} \widehat{LB}_s \right)^{d/\alpha}, 50, \delta_s)$ , and

$\widehat{p}_{X_{s'}} := \text{Estprob}(X_{s'}, \rho(X_s, X_{s'}), \left( \frac{1}{64L} \widehat{LB}_s \right)^{d/\alpha}, 50, \delta_s)$  then by Lemma 8,

$$\min(P_X(B(X_s, \rho(X_s, X_{s'}))), P_X(B(X_{s'}, \rho(X_s, X_{s'})))) \leq \left( \frac{1}{64L} \widehat{LB}_s \right)^{d/\alpha}. \quad (63)$$

Necessary, we have  $|\eta(X_s) - \frac{1}{2}| \geq \frac{1}{32} \Delta$ . Indeed, if  $|\eta(X_s) - \frac{1}{2}| < \frac{1}{32} \Delta$ , then on  $A_1 \cap A_2$ , by denoting  $\bar{k}_s$  the number of request labels in  $\text{ConfidentLabel}(X_s) := \text{ConfidentLabel}(X_s, k(\epsilon, \delta_s), t, \delta_s)$ , (where  $t = n - \sum_{s_i \in I, s_i < s} |Q_{s_i}|$  and  $|Q_{s_i}|$  the number of label requests used in  $\text{ConfidentLabel}(X_{s_i})$ )

We have:

$$\begin{aligned} \widehat{LB}_s &= |\widehat{\eta}_{\bar{k}_s}(X_s) - \frac{1}{2}| - b_{\delta_s, \bar{k}_s} \\ &\leq |\widehat{\eta}_{\bar{k}_s}(X_s) - \bar{\eta}_{\bar{k}_s}(X_s)| + |\bar{\eta}_{\bar{k}_s}(X_s) - \frac{1}{2}| - b_{\delta_s, \bar{k}_s} \\ &\leq |\bar{\eta}_{\bar{k}_s}(X_s) - \frac{1}{2}| \quad (\text{by (33)}) \\ &\leq |\eta(X_s) - \frac{1}{2}| \\ &\quad + \frac{1}{32} (1 - \frac{1}{32}) \Delta \quad (\text{by assumption (H3) and Theorem 4}) \end{aligned} \quad (64)$$

$$\begin{aligned} &< \frac{1}{32} \Delta + \frac{1}{32} (1 - \frac{1}{32}) \Delta \\ &= \frac{63}{1024} \Delta \end{aligned} \quad (65)$$

By assumption (H3) and (63), we have:

$$\begin{aligned}
|\eta(X_{s'}) - \frac{1}{2}| &\leq |\eta(X_s) - \frac{1}{2}| + \frac{1}{64} \widehat{LB}_s \\
&< \frac{1}{32} \Delta + \frac{1}{64} \cdot \frac{63}{1024} \Delta \quad (\text{by (65)}) \\
&= \left( \frac{1}{32} + \frac{1}{64} \cdot \frac{63}{1024} \right) \Delta \\
&\leq \left( 1 - \frac{1}{32} \right) \Delta
\end{aligned}$$

that contradicts (62), then we have  $|\eta(X_s) - \frac{1}{2}| \geq \frac{1}{32} \Delta$ . Therefore, by (63), (64), we have:

$$\begin{aligned}
P_X(B(X_{s'}, \rho(X_s, X_{s'}))) &\leq \left( \frac{1}{64L} \widehat{LB}_s \right)^{d/\alpha} \\
&\leq \left( \frac{1}{64L} \left( |\eta(X_s) - \frac{1}{2}| + \frac{1}{32} \left( 1 - \frac{1}{32} \right) \Delta \right) \right)^{d/\alpha} \\
&\leq \left( \frac{1}{64L} \left( |\eta(X_s) - \frac{1}{2}| + \left( 1 - \frac{1}{32} \right) |\eta(X_s) - \frac{1}{2}| \right) \right)^{d/\alpha} \\
&= \left( \frac{1}{64L} \left( 2 - \frac{1}{32} \right) |\eta(X_s) - \frac{1}{2}| \right)^{d/\alpha} \\
&= \left( \frac{63}{2048L} |\eta(X_s) - \frac{1}{2}| \right)^{d/\alpha}. \tag{66}
\end{aligned}$$

On the other hand, by (60), we have:

$$\begin{aligned}
P_X(B(x, \rho(X_{s'}, x))) &\leq \tilde{p}_\epsilon \\
&= \left( \frac{1}{128L} \Delta \right)^{d/\alpha} \\
&\leq \left( \frac{1}{128L} |\eta(x) - \frac{1}{2}| \right)^{d/\alpha}. \tag{67}
\end{aligned}$$

We have:

$$\begin{aligned}
|\eta(x) - \eta(X_s)| &\leq |\eta(x) - \eta(X_{s'})| + |\eta(X_{s'}) - \eta(X_s)| \\
&\leq L \cdot P_X(B(x, \rho(X_{s'}, x)))^{\alpha/d} + L \cdot P_X(B(X_{s'}, \rho(X_{s'}, X_s)))^{\alpha/d} \quad (\text{by assumption (H3)}) \\
&\leq \frac{1}{128} |\eta(x) - \frac{1}{2}| + \frac{63}{2048} |\eta(X_s) - \frac{1}{2}| \quad (\text{by (66) and (67)}) \tag{68} \\
&\leq \frac{1}{128} |\eta(x) - \frac{1}{2}| + \frac{\frac{63}{2048}}{1 - \frac{63}{2048}} |\eta(X_{s'}) - \frac{1}{2}| \quad (\text{by assumption (H3) and (66)}) \\
&\leq \frac{1}{128} |\eta(x) - \frac{1}{2}| + \frac{63}{1985} \left( 1 + \frac{1}{128} \right) |\eta(x) - \frac{1}{2}| \quad (\text{by (61)}) \\
&= \frac{79}{1985} |\eta(x) - \frac{1}{2}| \tag{69}
\end{aligned}$$

- b)  $X_{s'}$  **is informative**. In this case,  $s = s'$  and then we always obtains the equation (69), which becomes

$$|\eta(X_s) - \frac{1}{2}| \geq \left(1 - \frac{79}{1985}\right) |\eta(x) - \frac{1}{2}| \quad (70)$$

$$\begin{aligned} &\geq \left(1 - \frac{79}{1985}\right) \Delta \\ &\geq \frac{1}{2} \Delta \end{aligned} \quad (71)$$

Then

$$|\eta(X_s) - \frac{1}{2}| \geq \frac{1}{2} \Delta \quad (72)$$

On  $A_1 \cap A_2$ , by Theorem 5, the subroutine **ConfidentLabel**( $X_s$ ) uses at most  $\tilde{k}(\epsilon, \delta_s)$  request labels, and returns the correct label (with respect to the Bayes classifier) of  $X_s$ . Let us proof that  $f^*(x) = f^*(X_s)$ . Let us assume without loss of generality that  $\eta(X_s) - \frac{1}{2} \geq 0$ . We will show that  $\eta(x) - \frac{1}{2} \geq 0$ . We have:

$$\begin{aligned} \eta(x) - \frac{1}{2} &= \eta(x) - \eta(X_s) + \eta(X_s) - \frac{1}{2} \\ &\geq \eta(X_s) - \frac{1}{2} - \frac{79}{1985} |\eta(x) - \frac{1}{2}| \quad (\text{by (69)}) \\ &\geq \left(1 - \frac{79}{1985}\right) |\eta(x) - \frac{1}{2}| \\ &\quad - \frac{79}{1985} |\eta(x) - \frac{1}{2}| \quad (\text{by (69)}) \\ &= \frac{1827}{1985} |\eta(x) - \frac{1}{2}| \\ &\geq 0 \end{aligned}$$

Then  $f^*(x) = f^*(X_s)$ .

As  $|\eta(X_s) - \frac{1}{2}| \geq \frac{1}{2} \Delta$ , by using Theorem 5 (the second part), we can easily see that  $(X_s, \hat{Y}_s) \in \hat{S}_{ac}$  (where  $\hat{Y}_s$  is the inferred label of  $X_s$  provided by the subroutine **ConfidentLabel** in **KALLS**(Algorithm (1))).

Let  $X_x^{(1)}$  the nearest neighbor of  $x$  in  $\hat{S}_{ac}$ . We have:

$$\begin{aligned} |\eta(x) - \eta(X_x^{(1)})| &\leq L.P_X(B(x, \rho(x, X_x^{(1)})))^{\alpha/d} \\ &\leq L.P_X(B(x, \rho(x, X_s)))^{\alpha/d} \\ &\leq L.P_X(B(x, \rho(x, X_{s'})))^{\alpha/d} + L.P_X(B(X_{s'}, \rho(X_{s'}, X_s)))^{\alpha/d} \\ &\leq \frac{79}{1985} |\eta(x) - \frac{1}{2}| \quad \text{by (69)} \end{aligned} \quad (73)$$

Then,  $|\eta(X_x^{(1)}) - \frac{1}{2}| \geq \left(1 - \frac{79}{1985}\right) |\eta(x) - \frac{1}{2}| \geq \frac{1}{2} \Delta$  and by Theorem 5, the subroutine **ConfidentLabel**( $X_x^{(1)}$ ) outputs

$$\hat{Y}_x^{(1)} = f^*(X_x^{(1)}). \quad (74)$$

Furthermore, (73) implies

$$|\eta(x) - \eta(X_x^{(1)})| \leq |\eta(x) - \frac{1}{2}|$$

then  $f^*(x) = f^*(X_x^{(1)})$ . With (74), we easily deduce that:

$$f_{n,w}(x) = \hat{Y}_x^{(1)} = f^*(X_x^{(1)}) = f^*(x).$$



## A.6 Label complexity

### Lemma 9

Let us assume that  $w$  satisfies (12), (13), and  $w \geq T_{\epsilon, \delta}$ . Then, there exists an event  $A_5$  such that  $P(A_5) \geq 1 - \delta/8$ , and on  $A_1 \cap A_2 \cap A_3 \cap A_5$ . The condition (11) is sufficient to guarantee (56).

Before beginning the proof, let us define a notion that will be used through the proof.

### Definition 5

Let a set  $\mathcal{F} \subset \text{supp}(P_X)$ . Let  $\{x_1, \dots, x_m\} \subset \mathcal{F}$  and  $p(0, 1]$ . We say that the set  $\{x_1, \dots, x_m\} \subset \mathcal{F}$  is a  $p$ -probability-packing set of  $\mathcal{F}$  if:

$$\forall s, s' \leq m, s \neq s' \implies \rho(x_s, x_{s'}) > r_p(x_s) \vee r_p(x_{s'}) \quad (75)$$

where  $r_p$  is defined by (25), and  $a \vee b = \max(a, b)$  for  $a, b \in \mathbb{R}$

This notion of  $p$ -probability-packing comes from the Definition 1.4 in (Edgar, 2000). It will be used on a particular set of the form  $\{x \in \text{supp}(P_X), \gamma \leq |\eta(x) - \frac{1}{2}| \leq \gamma'\}$  (where  $0 < \gamma < \gamma'$ ). This allows us to upper bound the number of informative points where we have a very high confidence for inferring their labels.

### Proof

Let us consider the last update of  $I$ , the set of indexes of informative points used in KALLS(Algorithm 1).

Set  $s_I = \max I$ , the index of the last informative point. We consider two cases:

1. **First case:**  $s_I = w$ : we can easily see that (56) is satisfied, and we have trivially that the condition (11) is sufficient to guarantee (56).
2. **Second case:**  $s_I < w$ : then the total number of label requests up to  $s_I$  is:

$$\sum_{s \in I} |Q_s| \quad (76)$$

where  $|Q_s|$  is the number of label requests used in the subroutine **ConfidentLabel**(Algorithm(5)) with input  $X_s$ . Let  $s \in I$ . For brevity, let us denote  $\text{ConfidentLabel}(X_s, t) := \text{ConfidentLabel}(X_s, k(\epsilon, \delta_s), t, \delta_s)$ , (where  $t = n - \sum_{s_i \in I, s_i < s} |Q_{s_i}|$  the budget parameter). If  $s \neq s_I$ , the subroutine

**ConfidentLabel**( $X_s, t$ ) implicitly assumes that the process of label request do not takes into account the constraint related to the budget  $n$  (very large budget with respect to  $k(\epsilon, \delta_s)$ ) such that  $\text{ConfidentLabel}(X_s, t) = \text{ConfidentLabel}(X_s, t = \infty)$ . Then we have:

$$n > \sum_{\substack{s \in I \\ s < s_I}} |Q_s| \quad (77)$$

On the other hand, we want to guarantee the condition (56). For this, necessary for all  $s \in I$ , such that  $s \leq T_{\epsilon, \delta}$ , and  $s < s_I$ , at the end of the subroutine **ConfidentLabel**( $X_s, t$ ), the budget  $n$  is not yet reached and then we can replace the relation (77) by

$$n > \sum_{\substack{s \in I \\ s < s_I \\ s \leq T_{\epsilon, \delta}}} |Q_s| \quad (78)$$

Then, necessarily, (56) holds when (78) holds.

Also, for  $s \in I$ , by theorem5, if we assume that  $|\eta(X_s) - \frac{1}{2}| \geq \frac{1}{2}\Delta$ , we have that  $|Q_s| \leq \tilde{k}(\epsilon, \delta_s)$ , and the subroutine **ConfidentLabel**( $X_s, t$ ), (with  $t = n - \sum_{s_i \in I, s_i < s} |Q_{s_i}|$ ) terminates when the cut-off condition (34) is satisfied. The right hand side of (78) is equal to:

$$\sum_{\substack{s \in I \\ s < s_I \\ s \leq T_{\epsilon, \delta} \\ |\eta(X_s) - \frac{1}{2}| \geq \frac{1}{2}\Delta}} |Q_s| + \sum_{\substack{s \in I \\ s < s_I \\ s \leq T_{\epsilon, \delta} \\ |\eta(X_s) - \frac{1}{2}| \leq \frac{1}{2}\Delta}} |Q_s| \quad (79)$$

Firstly, let us consider the first term in (79) and denote it by  $T_1$ . Let us denote by  $B_s$  the event:

$$B_s = \{|\eta(X_s) - \frac{1}{2}| \geq \frac{1}{2}\Delta\}.$$

We have

$$\mathbb{1}_{B_s} = \sum_{j=1}^{m_\epsilon} \mathbb{1}_{B_{s,j}} \quad (80)$$

where

$$B_{s,j} = \{2^{j-1}\frac{1}{2}\Delta \leq |\eta(X_s) - \frac{1}{2}| \leq 2^j\frac{1}{2}\Delta\} \quad \text{and } m_\epsilon = \left\lceil \log_2 \left( \frac{1}{\frac{1}{2}\Delta} \right) \right\rceil.$$

Then,

$$\begin{aligned} T_1 &\leq \sum_{\substack{s \in I \\ s < s_I \\ s \leq T_{\epsilon, \delta} \\ |\eta(X_s) - \frac{1}{2}| \geq \frac{1}{2}\Delta}} \tilde{k}(\epsilon, \delta_s) \quad \text{by Theorem 5} \\ &= \sum_{\substack{s \in I \\ s < s_I \\ s \leq T_{\epsilon, \delta}}} \sum_{j=1}^{m_\epsilon} \tilde{k}(\epsilon, \delta_s) \mathbb{1}_{B_{s,j}} \end{aligned} \quad (81)$$

On  $B_{s,j}$ ,

$$\begin{aligned} \tilde{k}(\epsilon, \delta_s) &\leq \frac{c}{2^{2j}\Delta^2} \left[ \log\left(\frac{32s^2}{\delta}\right) + \log\log\left(\frac{32s^2}{\delta}\right) + \log\log\left(\frac{512\sqrt{\epsilon}}{2^j\Delta}\right) \right] \\ &\leq \frac{c}{2^{2j}\Delta^2} \left[ 2\log\left(\frac{32s^2}{\delta}\right) + \log\log\left(\frac{512\sqrt{\epsilon}}{\Delta}\right) \right] \end{aligned} \quad (82)$$

Then (81) becomes:

$$T_1 \leq \frac{c}{\Delta^2} \left[ 2\log\left(\frac{32T_{\epsilon, \delta}^2}{\delta}\right) + \log\log\left(\frac{512\sqrt{\epsilon}}{\Delta}\right) \right] \sum_{j=1}^{m_\epsilon} 2^{-2j} \sum_{\substack{s \in I \\ s < s_I \\ s \leq T_{\epsilon, \delta}}} \mathbb{1}_{B_{s,j}} \quad (83)$$

In (83), the term  $N_j = \sum_{\substack{s \in I \\ s < s_I \\ s \leq T_{\epsilon, \delta}}} \mathbb{1}_{B_{s,j}}$  represents the numbers of informative points that belong to the set

$$I_j = \{x, \gamma_{j-1} \leq |\eta(x) - \frac{1}{2}| \leq \gamma_j\} \quad (84)$$

(where  $\gamma_j = 2^j \cdot \frac{\Delta}{2}$ ,  $j = 1 \dots, m_\epsilon$ ). We will prove that

$$N_j \leq O\left((\gamma_j)^{\beta - \frac{d}{\alpha}}\right) \quad (85)$$

We proceed in two steps:

- The set of informative points that belong to  $I_j$  forms a  $p_j$ -probability-packing set (for  $p_j$  well chosen) of  $I_j$ .
  - The cardinal of any  $p_j$ -probability-packing set satisfies (85)
- (a) Let us begin with first step:  
 Let  $X_s, X_{s'}$  any two informative points that belong to  $I_j$ . Let us assume that  $s < s'$ .  
 As  $X_s \in I_j$ , we have  $|\eta(X_s) - \frac{1}{2}| \geq \frac{\Delta}{2}$  and by Theorem 5, the number of label requests  $\bar{k}_s$  used in **ConfidentLabel**( $X_s$ ) satisfies:

$$|\hat{\eta}_{\bar{k}_s} - \frac{1}{2}| \geq 2b_{\delta_s, \bar{k}_s} \quad (86)$$

where  $\hat{\eta}_{\bar{k}_s} := \hat{\eta}_{\bar{k}_s}(X_s)$  and  $b_{\delta_s, \bar{k}_s}$  are respectively defined by (26) and (4).  
 Then

$$LB_s := |\hat{\eta}_{\bar{k}_s} - \frac{1}{2}| - b_{\delta_s, \bar{k}_s} \geq 0.1b_{\delta_s, \bar{k}_s} \quad (87)$$

Additionally, as  $X_s$  and  $X_{s'}$  are both informative points, by Lemma 8, we necessary have on event  $A_3$  (see Lemma 8), that

$$\min(\hat{p}_X, \hat{p}_{X'}) \geq \frac{75}{94} \left( \frac{1}{64L} \widehat{LB_s} \right)^{d/\alpha} \quad (88)$$

On the event  $A_3$ , the equations (87), (88), necessary imply:

$$\min(P_X(B(X_s, \rho(X_{s'}, X_s))), P_X(B(X_{s'}, \rho(X_s, X_{s'})))) \geq \frac{28}{47} \left( \frac{1}{64L} \widehat{LB_{s'}} \right)^{d/\alpha} \quad (89)$$

Let us introduction the quantity  $\bar{\eta}_{\bar{k}_s} := \bar{\eta}_{\bar{k}_s}(X_s)$  defined by (26). We have, by Theorem 5, on the event  $A_2$  (see Theorem 5),

$$|\bar{\eta}_{\bar{k}_s} - \hat{\eta}_{\bar{k}_s}| \leq b_{\delta_s, \bar{k}_s}.$$

Then, on the event  $A_1 \cap A_2$ , we have:

$$\begin{aligned} |\bar{\eta}_{\bar{k}_s} - \hat{\eta}_{\bar{k}_s}| \leq b_{\delta_s, \bar{k}_s} &\Rightarrow |\hat{\eta}_{\bar{k}_s} - \frac{1}{2}| \geq |\bar{\eta}_{\bar{k}_s} - \frac{1}{2}| - b_{\delta_s, \bar{k}_s} \\ &\geq |\eta(X_s) - \frac{1}{2}| - \frac{31}{1024}\Delta - b_{\delta_s, \bar{k}_s} \\ &\quad \text{by smoothness assumption (see for e.g (37))} \\ &\geq |\eta(X_s) - \frac{1}{2}| - \frac{62}{1024}|\eta(X_s) - \frac{1}{2}| - b_{\delta_s, \bar{k}_s} \\ &\quad \text{as } |\eta(X_s) - \frac{1}{2}| \geq \frac{1}{2}\Delta \end{aligned} \quad (90)$$

$$= \frac{481}{512}|\eta(X_s) - \frac{1}{2}| - b_{\delta_s, \bar{k}_s} \quad (91)$$

Therefore, we have on  $A_1 \cap A_2$ :

$$\begin{aligned}
LB_s &= |\widehat{\eta}_{\bar{k}_s} - \frac{1}{2}| - b_{\delta_s, \bar{k}_s} \\
&= |\widehat{\eta}_{\bar{k}_s} - \frac{1}{2}| - \frac{4}{3}b_{\delta_s, \bar{k}_s} + \frac{1}{3}b_{\delta_s, \bar{k}_s} \\
&\geq \frac{1}{3}|\widehat{\eta}_{\bar{k}_s} - \frac{1}{2}| + \frac{1}{3}b_{\delta_s, \bar{k}_s} \\
&\quad \text{by using (86)} \\
&\geq \frac{1}{3} \left( \frac{481}{512} |\eta(X_s) - \frac{1}{2}| - b_{\delta_s, \bar{k}_s} \right) + \frac{1}{3}b_{\delta_s, \bar{k}_s} \\
&\quad \text{by using (91)} \\
&= \frac{481}{1536} |\eta(X_s) - \frac{1}{2}| \\
&\geq \frac{481}{1536} \gamma_{j-1} \text{ as } X_s \in I_j \\
&= \frac{481}{3072} \gamma_j
\end{aligned} \tag{92}$$

Then, the equation (92) becomes:

$$\min(P_X(B(X_s, \rho(X_{s'}, X_s))), P_X(B(X_{s'}, \rho(X_{s'}, X_s)))) \geq \frac{28}{47} \left( \frac{1}{L} \frac{481}{196608} \gamma_j \right)^{d/\alpha} \tag{93}$$

As the same way, we also obtain (93) if  $s' < s$ .

Then, if we set

$$p_j = \frac{28}{47} \left( \frac{1}{L} \frac{481}{196608} \gamma_j \right)^{d/\alpha} \tag{94}$$

we have that, by (25) and (93):

$$\rho(X_{s'}, X_s) \geq r_{p_j}(X_s) \vee r_{p_j}(X_{s'}) \tag{95}$$

Therefore, the set of informative points that belong to  $I_j = \{x, \gamma_{j-1} \leq |\eta(x) - \frac{1}{2}| \leq \gamma_j\}$  forms an  $p_j$ -probability-packing set.

- (b) As second step, let us determine an upper bound of the cardinal of any  $p_j$ -probability-packing set of  $I_j$ . Let  $\Lambda_j = \{x_1, \dots, x_{M_j}\}$  any  $p_j$ -probability-packing set of  $I_j$ .

For all  $s, s' \leq M_j$ , we obviously have:

$$s \neq s' \implies B(x_s, \frac{r_{p_j}(x_s)}{2}) \cap B(x_{s'}, \frac{r_{p_j}(x_{s'})}{2}) = \emptyset \tag{96}$$

Then, we have:

$$\begin{aligned}
P_X\left(\bigcup_{s=1}^{M_j} B(x_s, r_{p_j}(x_s)/2)\right) &= \sum_{i=1}^{M_j} P_X(B(x_s, r_{p_j}(x_s)/2)) \\
&\geq C_{db} \sum_{i=1}^{M_j} P_X(B(x_s, r_{p_j}(x_s))) \\
&\quad \text{by assumption (H4)} \\
&\geq C_{db} M_j p_j \\
&\quad \text{by (25)}
\end{aligned} \tag{97}$$

On the other hand, if  $z \in B(x_s, r_{p_j}(x_s)/2)$  for some  $s \leq M_j$ , by the assumption (H3), equation (25), and the fact that  $|\eta(x_s) - \frac{1}{2}| \leq \gamma_j$ , we have:

$$\begin{aligned}
|\eta(z) - \frac{1}{2}| &\leq \gamma_j + L(p_j)^{\alpha/d} \\
&= \gamma_j + \left(\frac{28}{47}\right)^{\alpha/d} \frac{481}{196608} \gamma_j \\
&= \gamma_j \left(1 + \left(\frac{28}{47}\right)^{\alpha/d} \frac{481}{196608}\right) \\
&= \tilde{c} \gamma_j
\end{aligned} \tag{98}$$

Where

$$\tilde{c} = \left(1 + \left(\frac{28}{47}\right)^{\alpha/d} \frac{481}{196608}\right) \tag{99}$$

Now we can upper bound  $M_j$ : by using (98), (97), and assumption (H2),

$$\begin{aligned}
C_{db} M_j p_j &\leq P_X\left(\bigcup_{s=1}^{M_j} B(x_s, r_{p_j}(x_s)/2)\right) \leq P_X\left(z \mid |\eta(z) - \frac{1}{2}| \leq \tilde{c} \gamma_j\right) \\
&\leq C(\tilde{c} \gamma_j)^\beta
\end{aligned} \tag{100}$$

Then,

$$\begin{aligned}
M_j &\leq \frac{C}{C_{db}} \frac{(\tilde{c} \gamma_j)^\beta}{p_j} \\
&= \tilde{b} (\gamma_j)^{\beta - \frac{d}{\alpha}}
\end{aligned} \tag{101}$$

where  $\tilde{b} = \frac{C}{C_{db}} \tilde{c}^{\beta - \frac{d}{\alpha}} \frac{47}{28} \left(\frac{196608}{481}\right)^{d/\alpha}$ .

Then, the cardinal of any  $p_j$ -probability-packing set of  $I_j$  is upper bound by  $O\left((\gamma_j)^{\beta - \frac{d}{\alpha}}\right)$ , consequently, equation (85) holds.

The equation (83) becomes:

$$\begin{aligned}
T_1 &\leq \frac{\tilde{c} \tilde{b}}{\Delta^2} \left[ 2 \log\left(\frac{32T_{\epsilon, \delta}^2}{\delta}\right) + \log \log\left(\frac{512\sqrt{e}}{\Delta}\right) \right] \sum_{j=1}^{m_\epsilon} 2^{-2j} (\gamma_j)^{\beta - \frac{d}{\alpha}} \\
&= \frac{\tilde{c} \tilde{b}}{2} \Delta^{\beta - \frac{d}{\alpha} - 2} \left[ 2 \log\left(\frac{32T_{\epsilon, \delta}^2}{\delta}\right) + \log \log\left(\frac{512\sqrt{e}}{\Delta}\right) \right] \sum_{j=1}^{m_\epsilon} 2^{(-2 + \beta - \frac{d}{\alpha})j} \\
T_1 &= \frac{\tilde{c} \tilde{b}}{\Delta^2} \left[ 2 \log\left(\frac{32T_{\epsilon, \delta}^2}{\delta}\right) + \log \log\left(\frac{512\sqrt{e}}{\Delta}\right) \right] \sum_{j=1}^{m_\epsilon} 2^{-2j} (\gamma_j)^{\beta - \frac{d}{\alpha}} \\
&= b_0 \left(\frac{1}{\epsilon}\right)^{\frac{2\alpha + d - \alpha\beta}{\alpha(\beta+1)}} \left[ 2 \log\left(\frac{32T_{\epsilon, \delta}^2}{\delta}\right) + \log \log\left(\frac{512\sqrt{e}}{\Delta}\right) \right] m_\epsilon
\end{aligned} \tag{102}$$

where  $b_0 = \tilde{c} \tilde{b} (2C)^{\frac{2\alpha + d - \alpha\beta}{\alpha(\beta+1)}}$ . Equation (102) holds because we have  $\alpha\beta \leq d$ ,  $\Delta = \max\left(\frac{\epsilon}{2}, \left(\frac{\epsilon}{2C}\right)^{\frac{1}{\beta+1}}\right)$ .

Now, it remains to upper bound the second term in (79). Denote it by  $T_2$ . By Lemma 1, (equation(28)), there exists an event  $A_5$  such that  $P(A_5) \geq 1 - \delta/8$ , and on  $A_5$ , we have:

$$T_2 \leq \sum_{s \leq T_{\epsilon, \delta}} |Q_s| \mathbb{1}_{\tilde{B}_s} \leq k(\epsilon, \delta) \left(1 + \frac{4}{P_X(\tilde{B})T_{\epsilon, \delta}} \log\left(\frac{8}{\delta}\right)\right) P_X(\tilde{B})T_{\epsilon, \delta}$$

Because  $|Q_s| \leq k(\epsilon, \delta)$  (according to the subroutine **ConfidentLabel**) for all  $s \leq T_{\epsilon, \delta}$  and where  $\tilde{B} = \{x, |\eta(x) - \frac{1}{2}| \leq \frac{\Delta}{2}\}$  and  $k(\epsilon, \delta)$  is defined in (5). Consequently, we have:

$$\begin{aligned}
T_2 &\leq k(\epsilon, \delta) \left( P_X(\tilde{B})T_{\epsilon, \delta} + 4 \log \left( \frac{8}{\delta} \right) \right) \\
&\leq k(\epsilon, \delta) \left( T_{\epsilon, \delta} \frac{1}{2^\beta} C \Delta^\beta + 4 \log \left( \frac{8}{\delta} \right) \right) \quad \text{by assumption (H2)} \\
&= k(\epsilon, \delta) \left( \left( \frac{128L}{\Delta} \right)^{d/\alpha} \log \left( \frac{8}{\delta} \right) \frac{1}{2^\beta} C \Delta^\beta + 4 \log \left( \frac{8}{\delta} \right) \right) \quad \text{by (54)} \\
&= k(\epsilon, \delta) \log \left( \frac{8}{\delta} \right) \left( (128)^{d/\alpha-\beta} 64^\beta C \left( \frac{1}{\Delta} \right)^{d/\alpha-\beta} + 4 \right) \quad (103)
\end{aligned}$$

As  $\alpha\beta \leq d$ ,  $\Delta \leq 1$ ,  $C \geq 1$ , the term  $(128)^{d/\alpha-\beta} 64^\beta C \left( \frac{1}{\Delta} \right)^{d/\alpha-\beta}$  in (103) is greater than 1. Thus, (103) becomes:

$$\begin{aligned}
T_2 &\leq 5k(\epsilon, \delta) \log \left( \frac{8}{\delta} \right) (128)^{d/\alpha-\beta} 64^\beta C \left( \frac{1}{\Delta} \right)^{d/\alpha-\beta} \\
&= 5c \left[ \log \left( \frac{1}{\delta} \right) + \log \log \left( \frac{1}{\delta} \right) + \log \log \left( \frac{512\sqrt{e}}{\Delta} \right) \right] \log \left( \frac{8}{\delta} \right) (128)^{d/\alpha-\beta} 64^\beta C \left( \frac{1}{\Delta} \right)^{d/\alpha-\beta+2} \\
&\quad \text{see (5), where } k(\epsilon, \delta) \text{ is defined} \\
&\leq \left( \frac{1}{\epsilon} \right)^{\frac{2\alpha+d-\alpha\beta}{\alpha(\beta+1)}} \left[ \log \left( \frac{1}{\delta} \right) + \log \log \left( \frac{1}{\delta} \right) + \log \log \left( \frac{512\sqrt{e}}{\Delta} \right) \right] \log \left( \frac{8}{\delta} \right) \tilde{u} \quad (104)
\end{aligned}$$

Where  $\tilde{u} = 5c(2C)^{\frac{2\alpha+d-\alpha\beta}{\alpha(\beta+1)}} 64^\beta (128)^{d/\alpha-\beta} C$ . The equation (104) holds by using the definition of  $\Delta$  (6).

By combining (104) and (102), the term obtained in (79) is less than:

$$\begin{aligned}
&b_0 \left( \frac{1}{\epsilon} \right)^{\frac{2\alpha+d-\alpha\beta}{\alpha(\beta+1)}} \left[ 2 \log \left( \frac{32T_{\epsilon, \delta}^2}{\delta} \right) + \log \log \left( \frac{512\sqrt{e}}{\Delta} \right) \right] m_\epsilon + \\
&\left( \frac{1}{\epsilon} \right)^{\frac{2\alpha+d-\alpha\beta}{\alpha(\beta+1)}} \left[ \log \left( \frac{1}{\delta} \right) + \log \log \left( \frac{1}{\delta} \right) + \log \log \left( \frac{512\sqrt{e}}{\Delta} \right) \right] \log \left( \frac{8}{\delta} \right) \tilde{u}
\end{aligned}$$

Thus, if the label budget  $n$  satisfies

$$n \geq b_0 \left( \frac{1}{\epsilon} \right)^{\frac{2\alpha+d-\alpha\beta}{\alpha(\beta+1)}} \left[ \left[ 2 \log \left( \frac{32T_{\epsilon, \delta}^2}{\delta} \right) + \log \log \left( \frac{512\sqrt{e}}{\Delta} \right) \right] m_\epsilon + \log \left( \frac{1}{\delta} \right) + \log \log \left( \frac{1}{\delta} \right) + \log \log \left( \frac{512\sqrt{e}}{\Delta} \right) \right] \log \left( \frac{8}{\delta} \right) \tilde{u} \quad (105)$$

We have that  $n$  satisfies (78), and (56) is necessary satisfied.

### Proof of Theorem 3

Finally, we are able to prove Theorem 3.

Previously, we have designed five events  $A_1, A_2, A_3, A_4, A_5$ , such that if the event  $A :=$

$A_1 \cap A_2 \cap A_3 \cap A_4 \cap A_5$ , and if the label budget  $n \geq \tilde{O} \left( \left( \frac{1}{\epsilon} \right)^{\frac{2\alpha+d-\alpha\beta}{\alpha(\beta+1)}} \right)$  (105) (where  $\tilde{O}$

assumes the existence of polylogarithmic factor in  $\frac{1}{\epsilon}$  and  $\frac{1}{\delta}$ ), then (56) is necessary satisfied, therefore, by Theorem 6, if  $w$  satisfies (12) and (13), and the event  $A$  holds, the classifier provided by the algorithm KALLS agrees with the Bayes classifier on  $\{x, |\eta(x) - \frac{1}{2}| > \Delta\}$ . Consequently, we have:

$$\begin{aligned}
R(\hat{f}_{n,w}) - R(f^*) &= \int_{\text{supp}(P_X)} \mathbf{1}_{\hat{f}_{n,w} \neq f^*(x)} |2\eta(x) - 1| dP_X(x) \\
&\leq \int_{\{x, |\eta(x) - \frac{1}{2}| \leq \Delta\} \cap \text{supp}(P_X)} 2\Delta dP_X(x) \\
&\leq 2C\Delta^{\beta+1} \quad \text{by assumption (H2)} \\
&\leq \epsilon \quad \text{by using the definition of } \Delta \text{ (6)}
\end{aligned}$$

Thus with probability at least  $1 - P(A^c) \geq 1 - \delta$ , (14) holds.