

Spectral conditions for graph rigidity in the Euclidean plane

Sebastian M. Cioabă^{*}, Sean Dewar[†] and Xiaofeng Gu[‡]

March 12, 2024

Abstract

Rigidity is the property of a structure that does not flex. It is well studied in discrete geometry and mechanics, and has applications in material science, engineering and biological sciences. A bar-and-joint framework is a pair (G, p) of graph G together with a map p of the vertices of G into the Euclidean plane. We view the edges of (G, p) as bars and the vertices as universal joints. The vertices can move continuously as long as the distances between pairs of adjacent vertices are preserved. The framework is rigid if any such motion preserves the distances between all pairs of vertices. In 1970, Laman obtained a combinatorial characterization of rigid graphs in the Euclidean plane. In 1982, Lovász and Yemini discovered a new characterization and proved that every 6-connected graph is rigid. Combined with a characterization of global rigidity, their proof actually implies that every 6-connected graph is globally rigid. Consequently, if Fiedler's algebraic connectivity is greater than 5, then G is globally rigid. In this paper, we improve this bound and show that for a graph G with minimum degree $\delta \geq 6$, if its algebraic connectivity is greater than $2 + \frac{1}{\delta-1}$, then G is rigid and if its algebraic connectivity is greater than $2 + \frac{2}{\delta-1}$, then G is globally rigid. Our results imply that every connected regular Ramanujan graph with degree at least 8 is globally rigid. We also prove a more general result giving a sufficient spectral condition for the existence of k edge-disjoint spanning rigid subgraphs. The same condition implies that a graph contains k edge-disjoint spanning 2-connected subgraphs. This result extends previous spectral conditions for packing edge-disjoint spanning trees.

MSC: 05C50, 52C25, 05C70, 05C40

Key words: eigenvalue, algebraic connectivity, connectivity, rigidity, redundant rigidity, global rigidity

^{*}Department of Mathematical Sciences, University of Delaware, Newark, DE 19716, USA. E-mail: cioaba@udel.edu

[†]Johann Radon Institute for Computational and Applied Mathematics (RICAM), Austrian Academy of Sciences, 4040 Linz, Austria. E-mail: sean.dewar@ricam.oeaw.ac.at

[‡]Department of Mathematics, University of West Georgia, Carrollton, GA 30118, USA. E-mail: xgu@westga.edu; Corresponding author.

1 Introduction

In this paper, we consider finite undirected simple graphs. Throughout the paper, k denotes a positive integer and G denotes a simple graph with vertex set $V(G)$ and edge set $E(G)$.

Rigidity is the property of a structure that does not flex. Arising from mechanics, rigidity has been studied in discrete geometry and combinatorics (see, among others, [14, 25]) and has applications in material science, engineering and biological sciences (see [9, 11, 37] for example). A **d -dimensional framework** is a pair (G, p) , where G is a graph and p is a map from $V(G)$ to \mathbb{R}^d . Roughly speaking, it is a straight line realization of G in \mathbb{R}^d . Two frameworks (G, p) and (G, q) are **equivalent** if $\|p(u) - p(v)\| = \|q(u) - q(v)\|$ holds for every edge $uv \in E(G)$, where $\|\cdot\|$ denotes the Euclidean norm in \mathbb{R}^d . Two frameworks (G, p) and (G, q) are **congruent** if $\|p(u) - p(v)\| = \|q(u) - q(v)\|$ holds for every $u, v \in V(G)$. A framework (G, p) is **generic** if the coordinates of its points are algebraically independent over the rationals. The framework (G, p) is **rigid** if there exists an $\varepsilon > 0$ such that if (G, p) is equivalent to (G, q) and $\|p(u) - q(u)\| < \varepsilon$ for every $u \in V(G)$, then (G, p) is congruent to (G, q) . As observed in [1], a generic realization of G is rigid in \mathbb{R}^d if and only if every generic realization of G is rigid in \mathbb{R}^d . Hence the generic rigidity can be considered as a property of the underlying graph. A graph is called **rigid** in \mathbb{R}^d if every/some generic realization of G is rigid in \mathbb{R}^d .

A d -dimensional framework (G, p) is **globally rigid** if every framework that is equivalent to (G, p) is congruent to (G, p) . This can be considered as the framework being rigid with ε being chosen to be as large as you wish. In [13] it was proven that if there exists a generic framework (G, p) in \mathbb{R}^d that is globally rigid, then any other generic framework (G, q) in \mathbb{R}^d will also be globally rigid. Following from this, we define a graph G to be **globally rigid** in \mathbb{R}^d if there exists a globally rigid generic framework (G, p) in \mathbb{R}^d . A closely related concept to global rigidity is redundant rigidity. A graph G is **redundantly rigid** in \mathbb{R}^d if $G - e$ is rigid in \mathbb{R}^d for every edge $e \in E(G)$. It was proven by Hendrickson [20] that any globally rigid graph in \mathbb{R}^d is $(d + 1)$ -connected and redundantly rigid in \mathbb{R}^d . Hendrickson [20] also conjectured the converse. It can be shown easily that it is true for $d = 1$, however the conjecture is not true for $d \geq 3$ [8]. The final case $d = 2$ of the conjecture was confirmed to be true by [7] and [22].

In the following, we will focus on rigidity and global rigidity only in \mathbb{R}^2 , unless otherwise stated. For a subset $X \subseteq V(G)$, let $G[X]$ be the subgraph of G induced by X and $E(X)$ denote the edge set of $G[X]$. A graph G is **sparse** if $|E(X)| \leq 2|X| - 3$ for every $X \subseteq V(G)$ with $|X| \geq 2$. By definition, any sparse graph is simple. If in addition $|E(G)| = 2|V(G)| - 3$, then G is called **(2, 3)-tight**. Laman [27] proved that a graph G is rigid in \mathbb{R}^2 if G contains a spanning (2, 3)-tight subgraph. Thus a (2, 3)-tight graph is also called a **minimally rigid** graph. In

history, it was first discovered by Pollaczek-Geiringer [33,34] who made notable progress on the properties of minimally rigid graphs, but her work was forgotten. Laman [27] rediscovered the characterization of minimally rigid graphs in \mathbb{R}^2 . Since then, a minimally rigid graph is known as a **Laman graph**.

Lovász and Yemini [29] gave a new characterization of rigid graphs using matroid theory and showed that 6-connected graphs are rigid. In fact they proved that every 6-connected graph is rigid even with the removal of any three edges. It implies that 6-connected graphs are redundantly rigid and thus globally rigid. They also constructed infinitely many 5-connected graphs that are not rigid.

We will study rigidity and global rigidity from spectral graph theory viewpoint. We describe the matrices and the eigenvalues of our interest below. If G is an undirected simple graph with $V(G) = \{v_1, v_2, \dots, v_n\}$, its **adjacency matrix** is the n by n matrix $A(G)$ with entries $a_{ij} = 1$ if there is an edge between v_i and v_j and $a_{ij} = 0$ otherwise, for $1 \leq i, j \leq n$. Let $D(G) = (d_{ij})_{1 \leq i, j \leq n}$ be the **degree matrix** of G , that is, the n by n diagonal matrix with d_{ii} being the degree of vertex v_i in G for $1 \leq i \leq n$. The matrix $L(G) = D(G) - A(G)$ is called the **Laplacian matrix** of G . For $1 \leq i \leq n$, we use $\mu_i(G)$ to denote the i -th smallest eigenvalue of $L(G)$. It is not difficult to see that $\mu_1(G) = 0$. The second smallest eigenvalue of $L(G)$, $\mu_2(G)$, is known as the **algebraic connectivity** of G .

Fiedler [10] proved that the vertex-connectivity of G is at least $\mu_2(G)$. Thus, the theorem of Lovász and Yemini [29] that every 6-connected graph is (globally) rigid implies that if $\mu_2(G) > 5$, then G is (globally) rigid. In this paper, we will improve this sufficient condition to “ $\mu_2(G) > 2 + \frac{1}{\delta-1}$ ” for rigidity (Corollary 1.4) and to “ $\mu_2(G) > 2 + \frac{2}{\delta-1}$ ” for global rigidity (Corollary 1.6). Actually we obtain more general sufficient spectral conditions for the existence of k edge-disjoint spanning rigid subgraphs in Theorem 1.1 and Corollary 1.2.

Theorem 1.1. *Let G be a graph with minimum degree $\delta(G) \geq 6k$. If*

$$(1) \mu_2(G) > \frac{6k-1}{\delta(G)+1},$$

$$(2) \mu_2(G-u) > \frac{4k-1}{\delta(G-u)+1} \text{ for every } u \in V(G), \text{ and}$$

$$(3) \mu_2(G-v-w) > \frac{2k-1}{\delta(G-v-w)+1} \text{ for every } v, w \in V(G),$$

then G contains at least k edge-disjoint spanning rigid subgraphs.

Theorem 1.1 has the following weaker, but neater, corollary.

Corollary 1.2. *Let G be a graph with minimum degree $\delta \geq 6k$. If*

$$\mu_2(G) > 2 + \frac{2k-1}{\delta-1},$$

then G contains at least k edge-disjoint spanning rigid subgraphs.

When $k = 1$, we obtain the following spectral condition for rigid graphs.

Corollary 1.3. *Let G be a graph with minimum degree $\delta(G) \geq 6$. If*

$$(1) \mu_2(G) > \frac{5}{\delta(G) + 1},$$

$$(2) \mu_2(G - u) > \frac{3}{\delta(G - u) + 1} \text{ for every } u \in V(G), \text{ and}$$

$$(3) \mu_2(G - v - w) > \frac{1}{\delta(G - v - w) + 1} \text{ for every } v, w \in V(G),$$

then G is rigid.

This result is similar in spirit and motivated by the work of Jackson and Jordán [23], in which they proved that a simple graph G is (globally) rigid if G is 6-edge-connected, $G - u$ is 4-edge-connected for every vertex u and $G - \{v, w\}$ is 2-edge-connected for any vertices $v, w \in V(G)$. Corollary 1.3 involves several conditions and we can show that the condition “ $\mu_2(G) > \frac{5}{\delta(G)+1}$ ” is essentially best possible. A family of examples will be constructed in a later section.

As before, we can also obtain the following weaker, but easier to state and verify, condition for a graph to be rigid.

Corollary 1.4. *Let G be a graph with minimum degree $\delta \geq 6$. If*

$$\mu_2(G) > 2 + \frac{1}{\delta - 1},$$

then G is rigid.

Using the same method as used in Theorem 1.1 when $k = 1$, we can prove the following similar results for redundant rigidity and global rigidity.

Theorem 1.5. *Let G be a graph with minimum degree $\delta(G) \geq 6$. If*

$$(1) \mu_2(G) > \frac{6}{\delta(G) + 1},$$

$$(2) \mu_2(G - u) > \frac{4}{\delta(G - u) + 1} \text{ for every } u \in V(G), \text{ and}$$

$$(3) \mu_2(G - v - w) > \frac{2}{\delta(G - v - w) + 1} \text{ for every } v, w \in V(G),$$

then G is redundantly rigid.

Corollary 1.6. *Let G be a graph with minimum degree $\delta \geq 6$. If*

$$\mu_2(G) > 2 + \frac{2}{\delta - 1},$$

then G is globally rigid.

In the next section, we will present some preliminaries that will be used in our proofs. The proofs of the main results will be presented in Section 3. A family of examples will be constructed in Section 4 to show the best possible bound of $\mu_2(G)$ in Corollary 1.3. In the last section, we will make some concluding remarks and give some applications of our results on rigidity and global rigidity of pseudo-random graphs and Ramanujan graphs, as well as a spectral sufficient condition for the existence of edge-disjoint spanning 2-connected graphs. Some questions are posted.

2 Preliminaries

The following theorem obtained by Lovász and Yemini [29] plays a very important role in graph rigidity in \mathbb{R}^2 .

Theorem 2.1 (Lovász and Yemini [29]). *A graph G is rigid if and only if $\sum_{X \in \mathcal{G}} (2|V(X)| - 3) \geq 2|V(G)| - 3$ for every collection \mathcal{G} of induced subgraphs of G whose edges partition $E(G)$.*

A characterization of global rigidity in \mathbb{R}^2 came from the combination of a result of Connelly [7] and a result of Jackson and Jordán [22].

Theorem 2.2 (Connelly [7], Jackson and Jordán [22]). *A graph G is globally rigid if and only if G is 3-connected and redundantly rigid, or G is a complete graph on at most three vertices.*

Sufficient conditions for the existence of edge-disjoint spanning rigid subgraphs have also been well studied. Jordán [24] showed that every $6k$ -connected graph contains k edge-disjoint spanning rigid subgraphs. Cheriyan, Durand de Gevigney and Szigeti [3] proved that a simple graph G contains k edge-disjoint spanning rigid subgraphs if $G - Z$ is $(6k - 2k|Z|)$ -edge-connected for every $Z \subset V(G)$. In fact, they proved a stronger result of packing spanning rigid subgraphs and spanning trees in [3]. The results were extended to a more general case in [16]. Motivated by the spanning tree packing theorem of Nash-Williams [32] and Tutte [36], the third author [17] obtained a partition condition for packing spanning rigid subgraphs, described below.

For any partition π of $V(G)$, $e_G(\pi)$ denotes the number of edges of G whose ends lie in two different parts of π . A part of π is **trivial** if it consists of a single vertex. Let $Z \subset V(G)$ and π be a partition of $V(G - Z)$ with n_0 trivial parts u_1, u_2, \dots, u_{n_0} . We define $n_Z(\pi)$ to be $\sum_{1 \leq i \leq n_0} |Z_i|$ where Z_i is the set of vertices in Z that are adjacent to u_i for $1 \leq i \leq n_0$. If $Z = \emptyset$, then $n_Z(\pi) = 0$.

Theorem 2.3 (Gu [17]). *A graph G contains k edge-disjoint spanning rigid subgraphs if for every $Z \subset V(G)$ and every partition π of $V(G - Z)$ with n_0 trivial parts and n'_0 nontrivial parts,*

$$e_{G-Z}(\pi) \geq k(3 - |Z|)n'_0 + 2kn_0 - 3k - n_Z(\pi).$$

Now we introduce some useful tools for Laplacian eigenvalues. Fiedler [10] applied Cauchy interlacing to the Laplacian matrix and obtained the following result (see also [2, Section 1.7] and [12, Thm. 13.5.1]).

Theorem 2.4 (Fiedler [10]). *If S is a subset of vertices of the graph G , then*

$$\mu_2(G) \leq \mu_2(G - S) + |S|.$$

For any subset $U \subset V(G)$, $\partial_G(U)$ or simply $\partial(U)$ denotes the set of edges in G , each of which has one end in U and the other end in $V(G) \setminus U$. We will also need the following result.

Lemma 2.5 (Liu et al. [28]). *Suppose that $X, Y \subset V(G)$ with $X \cap Y = \emptyset$. Let $e(X, Y)$ denote the number of edges with one end in X and the other in Y . If $\mu_2(G) \geq \max\{\frac{|\partial(X)|}{|X|}, \frac{|\partial(Y)|}{|Y|}\}$, then*

$$[e(X, Y)]^2 \geq |X||Y| \left(\mu_2(G) - \frac{|\partial(X)|}{|X|} \right) \left(\mu_2(G) - \frac{|\partial(Y)|}{|Y|} \right).$$

The following combinatorial lemma is well known. It was used in [19, 26] for example. For the sake of completeness, we include a short proof below.

Lemma 2.6. *Let G be a graph with minimum degree δ and U be a non-empty proper subset of $V(G)$. If $|\partial(U)| \leq \delta - 1$, then $|U| \geq \delta + 1$.*

Proof: We argue by contradiction and assume that $|U| \leq \delta$. Then $|U|(|U| - 1) + |\partial(U)| \geq |U|\delta$ by counting the total degrees of vertices in U . But $|U|(|U| - 1) + |\partial(U)| \leq \delta(|U| - 1) + (\delta - 1) \leq |U|\delta - 1$, contrary to the fact that $|U|(|U| - 1) + |\partial(U)| \geq |U|\delta$. Thus $|U| \geq \delta + 1$. \square

3 The proofs of the main results

In this section, we present the proofs of Theorem 1.1, Corollary 1.2, Theorem 1.5 and Corollary 1.6. We first restate Theorem 1.1 below and present its proof.

Theorem 3.1. *Let G be a graph with minimum degree $\delta(G) \geq 6k$. If*

$$\mu_2(G - Z) > \frac{6k - 2k|Z| - 1}{\delta(G - Z) + 1}$$

for every $Z \subset V(G)$ with $|Z| \leq 2$, then G has at least k edge-disjoint spanning rigid subgraphs.

Proof. Let $H = G - Z$. By Theorem 2.3, it suffices to show that for any partition π of $V(H)$ with n_0 trivial parts and n'_0 nontrivial parts,

$$e_H(\pi) \geq k(3 - |Z|)n'_0 + 2kn_0 - 3k - n_Z(\pi), \quad (1)$$

for every $Z \subset V(G)$.

We first prove that if $|Z| \geq 3$, then (1) is always true. Actually, for every trivial part (a single vertex) u_j , its degree $d_H(u_j)$ in H must satisfy the inequality $d_H(u_j) \geq \delta(G) - |Z_j| \geq 6k - |Z_j|$, where Z_j is the set of neighbors of u_j in Z . Recall that $n_Z(\pi) = \sum_{1 \leq j \leq n_0} |Z_j|$. If $|Z| \geq 3$, then

$$\begin{aligned} e_H(\pi) &\geq \frac{1}{2} \sum_{1 \leq j \leq n_0} d_H(u_j) \\ &\geq \frac{1}{2} \sum_{1 \leq j \leq n_0} \delta(G) - \frac{1}{2} \sum_{1 \leq j \leq n_0} |Z_j| \\ &\geq 3kn_0 - \frac{1}{2} n_Z(\pi) \\ &\geq k(3 - |Z|)n'_0 + 2kn_0 - 3k - n_Z(\pi). \end{aligned}$$

We assume that $|Z| \leq 2$ from now on. If $V_1, V_2, \dots, V_{n'_0}$ are the nontrivial parts in the partition π of H and u_1, u_2, \dots, u_{n_0} are the trivial parts of π , then

$$\sum_{1 \leq j \leq n_0} d_H(u_j) \geq \sum_{1 \leq j \leq n_0} (\delta(G) - |Z_j|) \geq 6kn_0 - n_Z(\pi). \quad (2)$$

Without loss of generality, we may assume that $|\partial_H(V_1)| \leq |\partial_H(V_2)| \leq \dots \leq |\partial_H(V_{n'_0})|$. For convenience, we will use ∂ for ∂_H in the following. If $|\partial(V_2)| \geq 6k - 2k|Z|$, then

$$\begin{aligned} e_H(\pi) &= \frac{1}{2} \left(\sum_{1 \leq i \leq n'_0} |\partial(V_i)| + \sum_{1 \leq j \leq n_0} d_H(u_j) \right) \\ &\geq \frac{1}{2} ((6k - 2k|Z|)(n'_0 - 1) + 6kn_0 - n_Z(\pi)) \\ &\geq k(3 - |Z|)n'_0 + 2kn_0 - 3k - n_Z(\pi), \end{aligned}$$

and we are done. Thus, we assume that $n'_0 \geq 2$ and $|\partial(V_2)| \leq 6k - 2k|Z| - 1$.

Let q be the largest index such that $|\partial(V_q)| \leq 6k - 2k|Z| - 1$. Then $2 \leq q \leq n'_0$. Therefore,

$$|\partial(V_i)| \geq 6k - 2k|Z| \text{ for } q < i \leq n'_0, \quad (3)$$

whenever such an i exists.

For $1 \leq i \leq q$, since $|\partial(V_i)| \leq 6k - 2k|Z| - 1 \leq \delta(G) - 2k|Z| - 1 \leq \delta(G - Z) - 1 = \delta(H) - 1$, Lemma 2.6 implies that $|V_i| \geq \delta(H) + 1$. As $\mu_2(H) > \frac{6k-2k|Z|-1}{\delta(H)+1}$, it follows that $|V_i|\mu_2(H) > 6k - 2k|Z| - 1$ for $1 \leq i \leq q$. By Lemma 2.5, for $2 \leq i \leq q$,

$$\begin{aligned} [e(V_1, V_i)]^2 &\geq |V_1||V_i| \left(\mu_2(H) - \frac{|\partial(V_1)|}{|V_1|} \right) \left(\mu_2(H) - \frac{|\partial(V_i)|}{|V_i|} \right) \\ &= (|V_1|\mu_2(H) - |\partial(V_1)|) (|V_i|\mu_2(H) - |\partial(V_i)|) \\ &> (6k - 2k|Z| - 1 - |\partial(V_1)|) (6k - 2k|Z| - 1 - |\partial(V_i)|) \\ &\geq (6k - 2k|Z| - 1 - |\partial(V_i)|)^2. \end{aligned}$$

Thus $e(V_1, V_i) > 6k - 2k|Z| - 1 - |\partial(V_i)|$, and so $e(V_1, V_i) \geq 6k - 2k|Z| - |\partial(V_i)|$. We get that

$$|\partial(V_1)| \geq \sum_{2 \leq i \leq q} e(V_1, V_i) \geq (6k - 2k|Z|)(q - 1) - \sum_{2 \leq i \leq q} |\partial(V_i)|,$$

and thus

$$\sum_{1 \leq i \leq q} |\partial(V_i)| = |\partial(V_1)| + \sum_{2 \leq i \leq q} |\partial(V_i)| \geq (6k - 2k|Z|)(q - 1). \quad (4)$$

Using (2), (3) and (4), we obtain that

$$\begin{aligned} e_H(\pi) &= \frac{1}{2} \left(\sum_{1 \leq i \leq n'_0} |\partial(V_i)| + \sum_{1 \leq j \leq n_0} d_H(u_j) \right) \\ &= \frac{1}{2} \left(\sum_{1 \leq i \leq q} |\partial(V_i)| + \sum_{q < i \leq n'_0} |\partial(V_i)| + \sum_{1 \leq j \leq n_0} d_H(u_j) \right) \\ &\geq \frac{1}{2} \left((6k - 2k|Z|)(q - 1) + (6k - 2k|Z|)(n'_0 - q) + 6kn_0 - n_Z(\pi) \right) \\ &\geq k(3 - |Z|)n'_0 + 2kn_0 - 3k - n_Z(\pi), \end{aligned}$$

which completes the proof. \square

Corollary 1.2 follows directly from Theorem 1.1 and the following lemma.

Lemma 3.2. *Let G be a graph with minimum degree $\delta \geq 6k$. If $\mu_2(G) > 2 + \frac{2k-1}{\delta-1}$, then $\mu_2(G - Z) > \frac{6k-2k|Z|-1}{\delta(G-Z)+1}$ for every $Z \subset V(G)$ with $|Z| \leq 2$.*

Proof. Notice that $\delta(G - u) \geq \delta - 1$ for every $u \in V(G)$ and $\delta(G - v - w) \geq \delta - 2$ for every $v, w \in V(G)$. It suffices to show that $\mu_2(G) > \frac{6k-1}{\delta+1}$, $\mu_2(G - u) > \frac{4k-1}{\delta}$ and $\mu_2(G - v - w) > \frac{2k-1}{\delta-1}$. Because $\delta \geq 6k$, it is not hard to verify that $2 + \frac{2k-1}{\delta-1} > \frac{6k-1}{\delta+1}$ and $1 + \frac{2k-1}{\delta-1} > \frac{4k-1}{\delta}$. Thus $\mu_2(G) > 2 + \frac{2k-1}{\delta-1} > \frac{6k-1}{\delta+1}$. By Theorem 2.4, $\mu_2(G - u) \geq \mu_2(G) - 1 > 1 + \frac{2k-1}{\delta-1} > \frac{4k-1}{\delta}$ and $\mu_2(G - v - w) \geq \mu_2(G) - 2 > \frac{2k-1}{\delta-1}$. \square

The proofs of Theorem 1.5 and Corollary 1.6 are quite similar to the above. We first restate Theorem 1.5 as below and give a quick proof.

Theorem 3.3. *Let G be a graph with minimum degree $\delta(G) \geq 6$. If*

$$\mu_2(G - Z) > \frac{6 - 2|Z|}{\delta(G - Z) + 1}$$

for every $Z \subset V(G)$ with $|Z| \leq 2$, then G is redundantly rigid.

Proof. We need to show that for any edge $f \in E(G)$, $G - f$ is rigid. By Theorem 2.3, it suffices to show that for any partition π of $V(G - f - Z)$ with n_0 trivial parts and n'_0 nontrivial parts,

$$e_{G-f-Z}(\pi) \geq (3 - |Z|)n'_0 + 2n_0 - 3 - n_Z(\pi),$$

for every $Z \subset V(G)$. If π consists of exactly one part, then $n_0 = 0$, $n'_0 = 1$, $e_{G-f-Z}(\pi) = 0$, and clearly the above inequality holds. Thus we may assume that π contains at least two parts in the following.

Notice that $e_{G-f-Z}(\pi) \geq e_{G-Z}(\pi) - 1$, and it suffices to show that

$$e_{G-Z}(\pi) \geq (3 - |Z|)n'_0 + 2n_0 - 2 - n_Z(\pi). \quad (5)$$

First, (5) will hold if $|Z| \geq 3$. The proof is the same as that of Theorem 3.1 for $k = 1$, and thus will be omitted here. We assume that $|Z| \leq 2$ in the remaining proof.

Case 1: $n'_0 \leq 1$. As $\delta(G) \geq 6$, we have $2e_{G-Z}(\pi) \geq \delta(G - Z)n_0 \geq (6 - |Z|)n_0$. If (5) does not hold then

$$(3 - |Z|)n'_0 + 2n_0 - 2 - n_Z(\pi) > e_{G-Z}(\pi) \geq \frac{1}{2}(6 - |Z|)n_0,$$

which yields

$$(6 - 2|Z|)n'_0 - 4 - 2n_Z(\pi) > (2 - |Z|)n_0.$$

Given that $n'_0 \leq 1$ and $|Z| \leq 2$, the above inequality holds only when $|Z| = 0$, $n_0 = 0$ and $n'_0 = 1$. However this implies that π consists of exactly one part, violating our assumption. Hence (5) must hold.

Case 2: $n'_0 \geq 2$. This case is similar to the proof of Theorem 3.1, and thus will be omitted. \square

Lemma 3.4. *Let G be a graph with minimum degree $\delta \geq 6k$. If $\mu_2(G) > 2 + \frac{2}{\delta-1}$, then $\mu_2(G - Z) > \frac{6-2|Z|}{\delta(G-Z)+1}$ for every $Z \subset V(G)$ with $|Z| \leq 2$.*

Proof. Notice that $\delta(G-u) \geq \delta-1$ for every $u \in V(G)$ and $\delta(G-v-w) \geq \delta-2$ for every $v, w \in V(G)$. It suffices to show that $\mu_2(G) > \frac{6}{\delta+1}$, $\mu_2(G-u) > \frac{4}{\delta}$ and $\mu_2(G-v-w) > \frac{2}{\delta-1}$. Because $\delta \geq 6k$, it is not hard to verify that $2 + \frac{2}{\delta-1} > \frac{6}{\delta+1}$ and $1 + \frac{2}{\delta-1} > \frac{4}{\delta}$. Thus $\mu_2(G) > 2 + \frac{2}{\delta-1} > \frac{6}{\delta+1}$. By Theorem 2.4, $\mu_2(G-u) \geq \mu_2(G) - 1 > 1 + \frac{2}{\delta-1} > \frac{4}{\delta}$ and $\mu_2(G-v-w) \geq \mu_2(G) - 2 > \frac{2}{\delta-1}$. \square

The proof of Corollary 1.6. By Theorem 3.3 and Lemma 3.4, G is redundantly rigid. Since $\mu_2(G) > 2$, the vertex-connectivity of G is at least 3. Hence G is globally rigid by Theorem 2.2. \square

4 Examples

In this section, we construct a family of graphs to show that the condition “ $\mu_2(G) > \frac{5}{\delta(G)+1}$ ” in Corollary 1.3 is essentially best possible.




Figure 1: An example of \mathcal{H}_d when $d = 10$

The family of graphs was initially constructed in [6]. Let $d \geq 6$ be an integer and let

H_1, H_2, H_3, H_4, H_5 be 5 vertex-disjoint copies of a graph obtained from K_{d+1} by deleting two disjoint edges. Suppose that the deleted edges are $a_i b_i$ and $u_i v_i$ in H_i for $1 \leq i \leq 5$. Let \mathcal{H}_d be the d -regular graph whose vertex set is $\bigcup_{i=1}^5 V(H_i)$ and whose edge set is the union of $\bigcup_{i=1}^5 E(H_i)$ with the set $F = \{b_1 a_2, b_2 a_3, b_3 a_4, b_4 a_5, b_5 a_1, u_1 v_3, u_3 v_5, u_5 v_2, u_2 v_4, u_4 v_1\}$. An example is shown in Figure 1 when $d = 10$. By the computation in [6], it follows that $\frac{5}{d+3} < \mu_2(\mathcal{H}_d) \leq \frac{5}{d+1}$ for $d \geq 6$. However, we can show that \mathcal{H}_d is not rigid as below.

Let $X_i = V(H_i)$ for $1 \leq i \leq 5$, and for $6 \leq i \leq 15$, X_i be the vertex set induced by a single edge in F . Let $\mathcal{G} = \{X_i : 1 \leq i \leq 15\}$. Clearly $\{E(X), X \in \mathcal{G}\}$ partitions $E(G)$. Then

$$\sum_{X \in \mathcal{G}} (2|X| - 3) = 5(2(d+1) - 3) + 10(2 \times 2 - 3) = 10d + 5.$$

Notice that $|V(G)| = 5d + 5$ and it follows that $2|V| - 3 = 10d + 7$. By Theorem 2.1, \mathcal{H}_d is not rigid.

5 Concluding remarks

In this paper, we discovered improved spectral conditions for rigid graphs and globally rigid graphs in \mathbb{R}^2 from Laplacian eigenvalues. Corollaries 1.4 and 1.6 give simple conditions for rigidity and global rigidity, respectively. However we do not know whether they are best possible. Corollary 1.6 might be close to best possible, as a necessary condition for globally rigid graphs in \mathbb{R}^2 is 3-connectedness. It would be interesting to see how large can $\mu_2(G)$ be for non-rigid graphs. Another problem of interest would be obtaining a spectral condition for a graph to contain a spanning (a, b) -tight subgraph for other values of a and b .

One immediate application of our results is on packing spanning 2-connected subgraphs. Since every rigid graph with at least 3 vertices is 2-connected, by Corollary 1.2, we have the following result on edge-disjoint spanning 2-connected subgraphs. This result can be seen as a spectral analogue of Jordán's combinatorial sufficient condition [24] for packing spanning rigid subgraphs and hence spanning 2-connected subgraphs. It also extends the spectral conditions for vertex-connectivity of [5, 10, 18, 26], and the spectral conditions for k edge-disjoint spanning trees of [5, 6, 15, 19, 21, 28], to k edge-disjoint spanning 2-connected subgraphs.

Proposition 5.1. *Let G be a graph with minimum degree $\delta \geq 6k$. If $\mu_2(G) > 2 + \frac{2k-1}{\delta-1}$, then G has at least k edge-disjoint spanning 2-connected subgraphs.*

We may also consider eigenvalues of other matrices. The matrix $Q(G) = D(G) + A(G)$ is called the **signless Laplacian matrix** of G . For $1 \leq i \leq n$, we use $\lambda_i := \lambda_i(G)$ and

$q_i := q_i(G)$ to denote the i -th largest eigenvalue of $A(G)$ and $Q(G)$, respectively. By Courant-Weyl inequalities (on page 29 of [2]), it is not hard to see that $\mu_2 + \lambda_2 \geq \delta$ and $\delta + \lambda_2 \leq q_2$. Thus all results involving μ_2 in the paper will imply sufficient conditions using λ_2 and q_2 . For example, by Corollary 1.6, it follows that for a graph G with minimum degree $\delta \geq 6$, if $\lambda_2(G) < \delta - 2 - \frac{2}{\delta-1}$ (or alternatively, if $q_2(G) < 2\delta - 2 - \frac{2}{\delta-1}$), then G is globally rigid. Other results of the paper can be modified in similar ways.

One reason to switch to eigenvalues of other matrices stems from pseudo-random graphs. Define $\lambda(G) = \max_{2 \leq i \leq n} |\lambda_i(G)| = \max\{|\lambda_2(G)|, |\lambda_n(G)|\}$. We call $\lambda(G)$ the second largest absolute eigenvalue of G . It is known that a d -regular graph on n vertices with *small* $\lambda(G)$ (compared to d ; for example, $\lambda(G) = \Theta(\sqrt{d})$) has edge distribution similar to the random graph of same edge density, namely it is a kind of **pseudo-random graph** (see [26] for more details). Clearly, the results in this paper imply sufficient conditions for the rigidity and global rigidity of pseudo-random graphs.

A connected d -regular graph G is called a **Ramanujan graph** if $|\lambda_i(G)| \leq 2\sqrt{d-1}$ for all $\lambda_i(G) \neq \pm d$ with $2 \leq i \leq n$. By definition, if G is a d -regular Ramanujan graph, then $\mu_2(G) \geq d - 2\sqrt{d-1}$. The rigidity of Ramanujan graphs was investigated by Servatius [35]. By Corollary 1.6, we can conclude that any connected d -regular Ramanujan graph with $d \geq 8$ is globally rigid in \mathbb{R}^2 . It is unknown whether this holds for smaller values of d , although as all cubic graphs (with the exception of the complete graph on four vertices) are flexible, we know that the lowest value of d where it could hold is either 4, 5, 6 or 7. We plan to focus on these specific graphs in more detail in future work [4].

Acknowledgments

The authors would like to thank the anonymous referees for reviewing the manuscript and providing insightful comments. The first author is supported by NSF grants DMS-1600768, CIF-1815922 and a JSPS Invitational Fellowship for Research in Japan S19016. The second author is supported by Austrian Science Fund (FWF): P31888. The third author is supported by a grant from the Simons Foundation (522728, XG).

References

- [1] L. Asimov and B. Roth, The rigidity of graphs, *Trans. Amer. Math. Soc.* **245** (1978), 279–289.
- [2] A. E. Brouwer and W. H. Haemers, *Spectra of Graphs*, Springer Universitext, 2012.

- [3] J. Cheriyan, O. Durand de Gevigney and Z. Szigeti, Packing of rigid spanning subgraphs and spanning trees, *J. Combin. Theory Ser. B* **105** (2014), 17–25.
- [4] S. M. Cioabă, S. Dewar and X. Gu, Graph rigidity properties of the Ramanujan graphs, manuscript in preparation.
- [5] S. M. Cioabă and X. Gu, Connectivity, toughness, spanning trees of bounded degrees, and spectrum of regular graphs, Special Issue dedicated to Professor Miroslav Fiedler, *Czechoslovak Math. J.* **66** (2016), 913–924.
- [6] S. M. Cioabă and W. Wong, Edge-disjoint spanning trees and eigenvalues of regular graphs, *Linear Algebra Appl.* **437** (2012), 630–647.
- [7] R. Connelly, Generic global rigidity, *Discrete Comput. Geom.* **33** (2005) 549–563.
- [8] R. Connelly, W. J. Whiteley, Global Rigidity: The Effect of Coning, *Discrete Comput. Geom.* **43** (2010), 717–735.
- [9] M. Cucuringu, A. Singer and D. Cowburn, Eigenvector synchronization, graph rigidity and the molecule problem, *Inf. Inference* **1** (2012), no. 1, 21–67.
- [10] M. Fiedler, Algebraic connectivity of graphs, *Czechoslovak Math. J.* **23(98)** (1973), 298–305.
- [11] M. Gáspár and P. Csermely, Rigidity and flexibility of biological networks, *Brief Funct Genomics.* **11(6)** (2012), 443–456.
- [12] C. Godsil and G. Royle, *Algebraic Graph Theory*, Springer-Verlag, New York, 2001.
- [13] S. J. Gortler, A. Healy, D. Thurston, Characterizing generic global rigidity, *Amer. J. Math.* **132(4)** (2010), 897–939.
- [14] J. Graver, B. Servatius and H. Servatius, *Combinatorial Rigidity*, Graduate Studies in Mathematics, Volume 2, American Mathematical Society (1993).
- [15] X. Gu, *Connectivity and Spanning Trees of Graphs*, PhD Dissertation, West Virginia University, 2013.
- [16] X. Gu, Packing spanning trees and spanning 2-connected k -edge-connected essentially $(2k - 1)$ -edge-connected subgraphs, *J. Comb. Optim.* **33** (2017), 924–933.

- [17] X. Gu, Spanning rigid subgraph packing and sparse subgraph covering, *SIAM J. Discrete Math.* **32** (2018), 1305–1313.
- [18] X. Gu, Toughness in pseudo-random graphs, *European J. Combin.* **92** (2021), 103255.
- [19] X. Gu, H. -J. Lai, P. Li and S. Yao, Edge-disjoint spanning trees, edge connectivity and eigenvalues in graphs, *J. Graph Theory* **81** (2016), 16–29.
- [20] B. Hendrickson, Conditions for unique graph realizations, *SIAM J. Comput.* **12** (1992) 65–84.
- [21] Y. Hong, X. Gu, H. -J. Lai and Q. Liu, Fractional spanning tree packing, forest covering and eigenvalues, *Discrete Appl. Math.* **213** (2016), 219–223.
- [22] B. Jackson and T. Jordán, Connected rigidity matroids and unique realizations of graphs, *J. Combin. Theory Ser. B* **94** (2005) 1–29.
- [23] B. Jackson and T. Jordán, A sufficient connectivity condition for generic rigidity in the plane, *Discrete Appl. Math.* **157** (2009), 1965–1968.
- [24] T. Jordán, On the existence of k edge-disjoint 2-connected spanning subgraphs, *J. Combin. Theory Ser. B* **95** (2005), 257–262.
- [25] T. Jordán, Combinatorial rigidity: graphs and matroids in the theory of rigid frameworks, *Discrete geometric analysis*, 33–112, *MSJ Mem.*, 34, *Math. Soc. Japan, Tokyo*, 2016.
- [26] M. Krivelevich and B. Sudakov, Pseudo-random graphs. More sets, graphs and numbers, 199–262, *Bölyai Soc. Math. Stud.*, 15, Springer, Berlin, 2006.
- [27] G. Laman, On graphs and rigidity of plane skeletal structures, *J. Engrg. Math.* **4** (1970), 331–340.
- [28] Q. Liu, Y. Hong, X. Gu and H. -J. Lai, Note on edge-disjoint spanning trees and eigenvalues, *Linear Algebra Appl.* **458** (2014), 128–133.
- [29] L. Lovász and Y. Yemini, On generic rigidity in the plane, *SIAM J. Algebraic Discrete Methods* **3** (1982), 91–98.
- [30] A. Lubotzky, R. Phillips and P. Sarnak, Ramanujan graphs, *Combinatorica* **8(3)** (1988), 261–277.
- [31] W. Mader. Minimale n -fach zusammenhängende Graphen, *Math. Ann.* **191** (1971), 21–28.

- [32] C. St. J. A. Nash-Williams, Edge-disjoint spanning trees of finite graphs, *J. London Math. Soc.* **36** (1961), 445–450.
- [33] H. Pollaczek-Geiringer, Über die Gliederung ebener Fachwerke, *Z. Angew. Math. Mech.* **7(1)** (1927), 58–72.
- [34] H. Pollaczek-Geiringer, Zur Gliederungstheorie Räumlicher Fachwerke, *Z. Angew. Math. Mech.* **12(6)** (1932), 369–376.
- [35] B. Servatius, On the rigidity of Ramanujan graphs, *Ann. Univ. Sci. Budapest. Eötvös Sect. Math.* **43** (2000), 165–170.
- [36] W. T. Tutte, On the problem of decomposing a graph into n factors, *J. London Math. Soc.* **36** (1961), 221–230.
- [37] D. Zelazo, A. Franchi, H. H. Bühlhoff and P. R. Giordano, Decentralized rigidity maintenance control with range measurements for multi-robot systems, *Int. J. Robot. Res.* **34(1)** (2015), 105–128.