

ABOUT THE SYMMETRY OF GENERAL RELATIVITY

SERHII SAMOKHVALOV

Communicated by XXX

Abstract. Generalized deformed gauge groups are used for investigation of symmetry of general relativity (GR). GR is formulated in generalized reference frames, which are represented by affine frames. The general principle of relativity is extended to the requirement of invariance of the theory with respect to the group GL^g of local linear transformations of affine frames. GR is interpreted as the gauge theory of the gauge group of translations T_M^g . The groups GL^g and T_M^g are united into the group S_M^g , which is their semidirect product and is the complete symmetry group of GR. By GL^g -gauge fixing one can obtain: Einstein gravity, GR in an orthogonal frame or teleparallel equivalent of GR, dilaton gravity, unimodular gravity, etc.

MSC: 70S10, 83C22, 83C40

Keywords: Affine frames, conservation law, general relativity, generalized deformed gauge groups, quasilocality

Contents

1 Introduction	2
2 Quasilocality of Gauge Charges	4
3 General Relativity in an Affine Frame	7
3.1 Field variables and Lagrangian of GRAF	7
3.2 Palatini equation	9
3.3 Field equations	10
4 Consequences of Gauge Invariance of GRAF	12
4.1 Generalized principle of relativity	13
4.2 Gauge translational invariance	15
4.3 Complete symmetry of the general relativity	18

5	Special Gauge Conditions	20
5.1	Einstein gravity	21
5.2	General relativity in an orthonormal frame	24
5.3	Dilaton gravity, unimodular gravity, and so on	25
6	Discussion and Conclusions	26

1. Introduction

There are large number theories of gravity locally equivalent to the general relativity. These theories differ in number of field variables and the more or less broad groups of gauge symmetries [7]. A broader group of gauge symmetry gives more opportunities to choose physically acceptable global solutions, for example devoid of singularities.

General relativity falls out of the general scheme of theories of gauge interactions [2]. Discussions about the gauge group that should form the basis of the gauge theory of gravity, as well as the geometric structure that should be consistent with it, continue today [6], [16]. One might even find an assertion that general relativity is not a gauge theory and does not have a gauge group because any theory can be written in a covariant form, and so diffeomorphisms are empty of dynamical meaning [1].

The main reason for the difficulty of interpreting gravity as a gauge interaction is that the gauge groups of internal symmetry do not act on the space-time manifold, but for gauge gravity this restriction is obviously meaningless because a gravitational field is born by an energy-momentum which associated with space-time translations.

A generalization of gauge groups for the case of their non-trivial action on the space-time continuum was proposed in [18]. Gauge fields, corresponding to an internal or external (space-time) symmetry, both acquire single interpretation as deformation parameters of suitable generalized deformed gauge groups. The most simple and transparent introduction to this subject is given in [19]. The dynamic and geometric meaning of the deformed group of diffeomorphisms with all completeness and mathematical rigor are discussed in [20] and [23].

It should be noted here that group-theoretical description of gauge fields is more fundamental than geometrical description because according to Klein's Erlangen Program geometrical structures are defined by the groups, which acts on manifolds. Thus, the use of a powerful apparatus of the theory of deformed gauge groups has

helped to solve certain long-standing problems that relate to the foundations of geometry [23].

In this work we use generalized deformed gauge groups for investigation of symmetry of general relativity. By the general relativity (GR) we understand the theory of gravity in the (pseudo)Riemannian space-time with the Hilbert's Lagrangian. However, we shall not restrict ourselves to holonomic reference frames associated with certain coordinate systems as in Einstein gravity (we shall call such coordinates Lagrangian coordinates), and we shall consider a theory of gravity in general anholonomic reference frames (affine frames or affine tetrads).

Transitions between affine frames form the gauge group of linear transformations GL^g . The general relativity in an affine frame (GRAF) is invariant under these GL^g -transformations. Transitions between holonomic reference frames (or between corresponding Lagrangian coordinates) form the group H^g that is isomorphic to the group of space-time diffeomorphisms and is a special subgroup of the group GL^g . In this sense *GRAF is based on the more wide principle of relativity than the Einstein gravity (EG)* [21].

On the space-time manifold M one can define Eulerian coordinates for numeration of its points. The group T_M^g of gauge translations on M is the group of diffeomorphisms of Eulerian coordinates in the additive parameterization [20]. GRAF is formulated as a gauge theory of translations and so is T_M^g -invariant.

Note that Eulerian coordinates have a different physical meaning unlike Lagrangian ones. If Eulerian coordinates numerate points on the space-time manifold, then Lagrangian ones define the holonomic reference frames. So T_M^g -transformations and H^g -transformations have a different physical meaning of gauge translations and changes of holonomic reference frames respectively. The identification of these transformations leads to known difficulties in determining the energy-momentum tensor in EG. On the other hand, separation of gauge translations and changes of reference frames (which takes place in GRAF) permit to define the energy-momentum of gravitational field as a T_M^g -tensor [12], [5]. When changing the reference frame (under the GL^g -transformations or H^g -transformations in the holonomic case), the energy-momentum varies according to the non-tensor law, as it should be [11]. It should be noted that the terms "Lagrangian and Eulerian coordinates" in the space-time continuum are similar (but not identical) to the corresponding concepts in the classical continuum mechanics. The analogy is that at translations the values of Eulerian coordinates change, and the values of Lagrangian coordinates does not.

The gauge group of linear transformations GL^g of affine frames and the generalized gauge group of translations T_M^g are united to the generalized gauge group

$S_M^g = GL^g \rtimes T_M^g$ that is their semidirect product. The group S_M^g is the group of symmetry of GRAF that defines a structure of GRAF.

In this article, first of all, in Section 2, we prove the fact that the currents which are conserved as a result of any generalized gauge symmetries G_M^g are quasilocal that means they have superpotentials. Moreover, we show that equations for the gauge fields in the gauge theories of generalized gauge groups G_M^g are equivalent to the expressions of currents (which are conserved as result of G_M^g -symmetries) by the superpotentials.

Then, in Section 3, we describe GRAF: field variables, the Lagrangian, field equations. It is shown, that Palatini equation (equality to zero of variational derivatives of Lagrangian with respect to connection coefficients) is equivalent to metricity and non-torsionity conditions and performed in (pseudo)Riemannian space-time automatically. It is also shown that the equation of gravitational field in GRAF is the Einstein equation in an affine frame.

In Section 4 we study consequences of S_M^g -invariance of GRAF. This is the central part of the work. Here we show that the Palatini equation is a strong Noether identity which follows from the GL^g -invariance of GRAF. Then we find a tensor density of energy-momentum of gravitational field in GRAF and its superpotential. Noether identities which follow from the gauge translational invariance (T_M^g -invariance) of GRAF are written down. Equations of gravity written down in the form that is similar to the electromagnetic ones (as Maxwell equations). It is also shown that the complete symmetry group of GRAF $S_M^g = GL^g \rtimes T_M^g$ has many deformed forms that lead to a lot of expressions for energy-momentum tensors of gravitational field. Among these forms is one highlighted, so called group of parallel transports in the Riemannian space [23], that corresponds to the case of reconciling the geometric structure given by the action of the group with the geometric structure of the Riemannian space. The energy-momentum of gravitational field in this case is localizable.

In section 5 we show that by limiting admissible reference frames (by GL^g -gauge fixing) from GRAF, in addition to EG, one can obtain other local equivalent formulations of GR: general relativity in an orthonormal frame (GROF) [5], [17] or teleparallel equivalent of general relativity (TEGR) [12], [1], [11], dilaton gravity (DG) [4], unimodular gravity (UG) [27], etc.

2. Quasilocality of Gauge Charges

In this section it is proved that *quantities which are conserved as a result of gauge symmetries are quasilocal* [22]. Quasilocality of the quantity Q means, that the

current J^μ , that relate to the quantity $Q = \int J^0 dV$, has the form $J^\mu = \partial_\sigma S^{\sigma\mu}$, where $S^{\sigma\mu} = -S^{\mu\sigma}$ is its superpotential [26]. Here x^μ are the Eulerian coordinates on the space-time manifold M , for which we will use the Greek indices of the middle of the alphabet. Our proof is based on the generalized Noether theorem, which generalizes the former Noether theorem [14] for the case of existence of a surface part of the Lagrangian.

Let the system of fields $q^i(x)$ (with indices of the middle of the Latin alphabet) be given on the space-time manifold M . Let the action be given: $S = \int \Lambda dx$, where dx is an element of coordinate volume in M , with the Lagrangian

$$\Lambda = L + \partial_\sigma V^\sigma \quad (1)$$

which consists of the volume L and the surface $\partial_\sigma V^\sigma$ parts ($\partial_\sigma := \partial/\partial x^\sigma$), and L and V^σ depend on the fields q and their partial derivatives ∂q . Therefore, the complete Lagrangian is linearly dependent also on the second derivatives of the fields $\partial^2 q$, which fold in $\partial_\sigma V^\sigma$. The surface part $\partial_\sigma V^\sigma$ does not affect the equations of motion determined solely by the bulk Lagrangian L . The surface part characterizes the boundary-value problem as well as the symmetry defined by the boundary-value problem.

At the transformations

$$x'^\mu = x^\mu + \delta x^\mu, \quad q'^i(x) = q^i(x) + \delta q^i$$

the action is transformed: $\bar{\delta}S = \int \delta' \Lambda dx$, where $\delta' \Lambda := \Lambda'(x')J - \Lambda(x)$ is the integral variation of the Lagrangian Λ , $J := |\partial_\rho x'^\sigma|$. Here $\Lambda'(x')$ is the Lagrangian, calculated with $q'^i(x')$. According to (1) we obtain

$$\delta' \Lambda = \delta' L + \partial_\sigma \delta' V^\sigma \quad (2)$$

where

$$\delta' V^\sigma := \partial_{\rho'} x^\sigma V'^{\rho'}(x')J - V^\sigma(x). \quad (3)$$

Here we have taken into account that $\partial_\sigma(\partial_{\rho'} x^\sigma J) \equiv 0$, where $\partial_{\rho'} := \partial/\partial x'^{\rho'}$. We obtain

$$\delta' L = \nabla_{q^i} L \delta q^i + \partial_\sigma (P_i^\sigma \delta q^i + L \delta x^\sigma)$$

where

$$\nabla_{q^i} L := Q_i - \partial_\sigma P_i^\sigma \quad (4)$$

is the variational derivatives of the volume Lagrangian, $Q_i := \partial_{q^i} L$, $P_i^\sigma := \partial_{\partial_\sigma q^i} L$, where $\partial_{q^i} := \partial/\partial q^i$, $\partial_{\partial_\nu q^i} := \partial/\partial \partial_\nu q^i$. Thus, taking into account (2), we obtain

$$\delta' \Lambda = \nabla_{q^i} L \delta q^i + \partial_\sigma (P_i^\sigma \delta q^i + L \delta x^\sigma + \delta' V^\sigma). \quad (5)$$

Let be given generalized gauge group G_M^g [18], that is parameterized by the functions $g^a(x)$ (here we will use indexes from the beginning of the Latin alphabet) and its infinitesimal transformations δ_g are defined

$$\delta_g x^\mu = h_a^\mu g^a, \quad \delta_g q^i = a_a^i g^a + b_a^{i\mu} \partial_\mu g^a, \quad \delta'_g V^\sigma = c_a^\sigma g^a + d_a^{\mu\sigma} \partial_\mu g^a \quad (6)$$

where a_a^i , $b_a^{i\mu}$, c_a^σ , $d_a^{\mu\sigma}$ and h_a^μ are the functions, that depend on x explicitly or implicitly via the fields q and their derivatives. Formulas (6) define the group G_M^g as the transformations group. We will omit the display of dependence on x of the fields q^i and the parameters g^a of the group G_M^g , where this does not lead to confusion.

Substitution of expressions (6) in (5) allows us to express the integral variation of the total Lagrangian through the parameters of the group G_M^g and their derivatives

$$\delta'_g \Lambda = \nabla_{q^i} L (a_a^i g^a + b_a^{i\mu} \partial_\mu g^a) - \partial_\sigma (J_a^\sigma g^a + S_a^{\mu\sigma} \partial_\mu g^a) \quad (7)$$

where the definition is entered

$$J_a^\sigma := -P_i^\sigma a_a^i - L h_a^\sigma - c_a^\sigma, \quad S_a^{\mu\sigma} := -P_i^\sigma b_a^{i\mu} - d_a^{\mu\sigma} \quad (8)$$

for the generalized Noether current J_a^σ and its superpotential $S_a^{\mu\sigma}$.

In the case of invariance of the action S with respect to the transformations of the group G_M^g , we have $\bar{\delta}_g S = 0$ and hence $\delta'_g \Lambda = 0$. Since the parameters of the group G_M^g and their derivatives are independent functions, equating of coefficients at them in expression (7) to zero allows us to find the following generalized strong (performed everywhere, not only on shell) Noether identities

$$\nabla_{q^i} L a_a^i = \partial_\sigma J_a^\sigma, \quad \nabla_{q^i} L b_a^{i\mu} = J_a^\mu + \partial_\sigma S_a^{\mu\sigma}, \quad S_a^{\mu\nu} = -S_a^{\nu\mu}. \quad (9)$$

Using expression (4) for the variational derivatives, second identity (9) can be represented as

$$J_a^\mu = Q_i b_a^{i\mu} + P_i^\sigma \partial_\sigma b_a^{i\mu} + \partial_\sigma d_a^{\mu\sigma} \quad (10)$$

which often makes it easier to find the current J_a^μ , especially in the case of $b_a^{i\mu} = const$ and $d_a^{\mu\sigma} = const$, thus $J_a^\mu = Q_i b_a^{i\mu}$.

On shell second identity (9) gives

$$J_a^\mu = -\partial_\sigma S_a^{\mu\sigma}$$

so current J_a^μ is *quasilocal* and $S_a^{\mu\sigma}$ is its *superpotential*.

In gauge theories part of fields q^i are the gauge fields A_μ^a , for them $b_{a\mu}^{\nu b} \sim \delta_a^b \delta_\mu^\nu$ and others coefficients $b_a^{i\mu}$ equal to zero. In this case $J_a^\mu \sim \partial_{A_\mu^a} L$. Moreover, second identity (9) gives

$$\nabla_{A_\mu^a} L \sim J_a^\mu + \partial_\sigma S_a^{\mu\sigma}$$

so equation of motion for gauge fields $\nabla_{A_\mu^a} L = 0$ is reduced to an expression of the current by the superpotential: $J_a^\mu = -\partial_\sigma S_a^{\mu\sigma}$. From this formula, taking into account the antisymmetry of the superpotential by the upper indices (9), immediately follows the weak Noether identity $\partial_\sigma J_a^\sigma = 0$. That is the conservation law of the current that follows also from first identity (9) on shell.

On the infinitesimal level deformations of the group $G_M^g \rightarrow G_M^{gH}$ are reduced to gauge linear transformations of parameters $g^a = H_\alpha^a g^\alpha$ with $|H_\alpha^a| \neq 0$ [18]. Under deformations currents and superpotentials change according to the formulas

$$J_\alpha^\mu = H_\alpha^a J_a^\mu + \partial_\sigma H_\alpha^a S_a^{\sigma\mu}, \quad S_\alpha^{\mu\nu} = H_\alpha^a S_a^{\mu\nu}. \quad (11)$$

Thus deformations lead to redistributions between J_a^μ and $\partial_\sigma S_a^{\mu\sigma}$ in the expression $J_a^\mu + \partial_\sigma S_a^{\mu\sigma}$.

3. General Relativity in an Affine Frame

In this section we *generalize the general principle of relativity to use anholonomic reference frames (affine frames)*. Here we describe general relativity in an affine frame (GRAF): main field variables of the theory, Lagrangian and field equations [21].

3.1. Field variables and Lagrangian of GRAF

Suppose that in the space-time M an affine frame is given $h_m = h_m^\mu \partial_\mu$, which vectors are numbered in Latin letters from the middle of the alphabet. Suppose also that space-time is (pseudo) Riemannian, and therefore a scalar product is given $h_m \cdot h_n = g_{mn}$. Replacement of coordinate indices to frame indices and vice versa will be performed using matrices h_m^μ and inverse ones h_μ^m , in particular $g_{\mu\nu} = h_\mu^m h_\nu^n g_{mn}$. We will also assume that in M is given an affine connection with connection 1-form $\gamma_n^m = \gamma_{\mu n}^m dx^\mu$, where $\gamma_{\mu n}^m$ is the connection coefficients in the affine frame h_m . Anholonomic coefficients F_{mn}^k of the reference fields h_m are defined by the expression

$$[h_m, h_n] = F_{mn}^k h_k$$

where square brackets denote the commutator of the vector fields. The group-theoretic meaning of the affine frame as a set of generators of a deformed group of diffeomorphisms, and of the anholonomic coefficients as its structural functions, similar to the structural constants of the Lie group, is important [18].

The condition of absence of torsion (non-torsionity condition) is

$$F^k_{mn} = \gamma^k_{mn} - \gamma^k_{nm} \quad (12)$$

and the metricity condition is

$$\partial_k g_{mn} = \gamma_{mkn} + \gamma_{nkm} \quad (13)$$

where $\partial_k g_{mn} := h^\mu_k \partial_\mu g_{mn}$. Under the conditions (12), (13), the connection coefficients are decomposed into the sum $\gamma_{mkn} = \omega_{mkn} + \sigma_{mkn}$, where

$$\omega_{mkn} = \frac{1}{2}(F^k_{mnn} + F^k_{mkn} - F^k_{nkm}), \quad \sigma_{mkn} = \frac{1}{2}(\partial_n g_{mk} + \partial_k g_{mn} - \partial_m g_{kn}) \quad (14)$$

with symmetry properties $\omega_{mkn} = -\omega_{nkm}$, $\sigma_{mnk} = \sigma_{mkn}$. The point indicates from where the index was lowered (or upped).

The main field variables in GRAF are the transition coefficients h^m_μ between affine and coordinate frames, the metric g_{mn} and the connection coefficients $\gamma^m_{\mu n}$ in the affine frame (frame-metric-affine gravity theory: generalized Einstein-Palatini model). This choice of field variables dictates the breakup of the Hilbert's Lagrangian

$$\Lambda = -\frac{1}{2\kappa} e R \quad (15)$$

to a bulk part L_γ (the truncated Hilbert's Lagrangian) that does not depend on the derivatives of the connection coefficients $\gamma^m_{\mu n}$, and a surface part having the form of divergence $\partial_\sigma V_\gamma^\sigma$ into which derivatives of $\gamma^m_{\mu n}$ are folded

$$\Lambda = L_\gamma + \partial_\sigma V_\gamma^\sigma \quad (16)$$

where κ is the Einstein gravitational constant (which we will hereafter assume equal to 1), $e := \sqrt{|g_{\mu\nu}|}$, and R is the scalar curvature of the space-time M . Indeed, since

$$\frac{1}{2}eR = \Sigma_m^{n\mu\nu} (\partial_\mu \gamma_{\nu n}^m + \gamma_{\mu s}^m \gamma_{\nu n}^s) \quad (17)$$

where $\Sigma_m^{n\mu\nu} := \frac{1}{2}e\delta_{ms}^{\mu\nu}g^{sn}$, $\delta_{mn}^{\mu\nu} := h_n^\mu h_n^\nu - h_n^\mu h_m^\nu$, throwing in the first term of formula (17) the derivative to the first factor and changing the sign, we obtain

$$\Lambda = \partial_\mu \Sigma_m^{n\mu\nu} \gamma_{\nu n}^m - \Sigma_m^{n\mu\nu} \gamma_{\mu s}^m \gamma_{\nu n}^s - \partial_\mu (\Sigma_m^{n\mu\nu} \gamma_{\nu n}^m).$$

So

$$L_\gamma = \partial_\mu \Sigma_m^{n\mu\nu} \gamma_{\nu n}^m - H_\gamma, \quad H_\gamma := \Sigma_m^{n\mu\nu} \gamma_{\mu s}^m \gamma_{\nu n}^s \quad (18)$$

$$V_\gamma^\sigma = \Sigma_m^{n\nu\sigma} \gamma_{\nu n}^m = e \gamma_n^{[n\sigma]}.$$

Square brackets here mean antisymmetrization by the indexes contained therein.

All addends of the partition (16), like Hilbert's own Lagrangian (15), are scalar densities with respect to changing of coordinates x^μ . A general-covariant method of separating a surface term from the Hilbert's Lagrangian is possible due to the use of frames fields that do not change when changing coordinates.

3.2. Palatini equation

In this subsection we show that the *Palatini equation obtained by variation of Hilbert's Lagrangian with respect to coefficients of affine connection in the affine frame is the identity due to the metricity and non-torsionity conditions.*

The integral variation of Lagrangian L_γ is determined by the expression

$$\begin{aligned} \delta' L_\gamma &= \partial_{\gamma^m_{\nu n}} L_\gamma \delta \gamma^m_{\nu n} + \nabla_{h^p_\nu} L_\gamma \delta h^p_\nu + \nabla_{g_{mn}} L_\gamma \delta g_{mn} \\ &+ \partial_\mu (\delta \Sigma_m^{n\mu\nu} \gamma^m_{\nu n} + L_\gamma \delta x^\mu). \end{aligned} \quad (19)$$

Let us Consider the terms in formula (19). The derivatives of the Lagrangian L_γ with respect to $\gamma^m_{\nu n}$ are

$$\partial_{\gamma^m_{\nu n}} L_\gamma = \partial_\sigma \Sigma_m^{n\sigma\nu} - M_m^{n\nu}$$

where

$$M_m^{n\nu} := \partial_{\gamma^m_{\nu n}} H_\gamma = \Sigma_m^{s\nu\sigma} \gamma^n_{\sigma s} - \Sigma_s^{n\nu\sigma} \gamma^s_{\sigma m}.$$

Equating these derivatives to zero, we obtain equation which we will call the Palatini equation in the affine frame

$$\partial_\sigma \Sigma_m^{n\sigma\nu} - M_m^{n\nu} = 0. \quad (20)$$

With metricity condition (13), Palatini equation (20) goes into the non-torsionity condition (12) and vice versa. Therefore, when these two conditions are fulfilled simultaneously, Palatini equation is fulfilled as identity.

Indeed, a direct calculation gives

$$\begin{aligned} \partial_\sigma \Sigma_m^{n\sigma\nu} &= \frac{1}{2} e [F_{m s}^\nu - \delta_{m s}^{\nu\sigma} (R_\sigma + S_\sigma) + \delta_{m k}^{\nu\sigma} g^{kp} \partial_\sigma g_{ps}] g^{sn} \\ M_m^{n\nu} &= -\frac{1}{2} e (\gamma_{s m}^{\cdot\nu} + \gamma_{s m}^\nu - h_m^\nu \gamma_{s p}^{\cdot p} - h_s^\nu \tilde{R}_m) g^{sn} \end{aligned} \quad (21)$$

where $R_m := F_{s m}^s$, $S_m := \frac{1}{2} g^{ps} \partial_m g_{ps}$, $\tilde{R}_m := \gamma_{s m}^s$, so the Palatini equation (20) is reduced to the equation

$$\begin{aligned} F_{m n}^\nu + \gamma_{n m}^{\cdot\nu} + \gamma_{n m}^\nu - h_s^\nu g^{sp} \partial_m g_{pn} + h_n^\nu (R_m + S_m - \tilde{R}_m) \\ - h_m^\nu (R_n + S_n - g^{sp} \partial_s g_{pn} + \gamma_{np}^{\cdot p}) = 0. \end{aligned} \quad (22)$$

Due to the metricity condition (13), we have $\tilde{R}_n = g^{sp} \partial_s g_{pn} - \gamma_{np}^{\cdot p}$, $\tilde{S}_m = S_m$, where $\tilde{S}_m := \gamma_{m s}^s$, and equation (22) is reduced to the equation

$$F_{m n}^\nu = \gamma_{m n}^\nu - \gamma_{n m}^\nu + \delta_{m n}^{\nu s} (R_s + S_s - \tilde{R}_s).$$

Summing up this equation by ν and m (with help of h_ν^m) we obtain $(d-2)(R_n + S_n - \tilde{R}_n) = 0$, where d is the dimension of space-time M . Therefore, when $d \neq 2$ equation (22) is reduced to the non-torsionity condition (12).

Now let the initial assumption be the non-torsionity condition (12), from which follows $R_s = \tilde{R}_s - \tilde{S}_s$. In this case, equation (22) is reduced to the equation

$$\begin{aligned} \partial_m g_{kn} &= \gamma_{kmn} + \gamma_{nmk} + g_{kn}(S_m - \tilde{S}_m) \\ &\quad - g_{km}(\tilde{R}_n + S_n - \tilde{S}_n - g^{sp}\partial_s g_{pn} + \gamma_{np}^{\cdot p}). \end{aligned}$$

Summing up this equation by k and m (with help of g^{km}) we obtain

$$(1-d)(\tilde{R}_n + S_n - \tilde{S}_n - g^{sp}\partial_s g_{pn} + \gamma_{np}^{\cdot p}) = 0$$

so (for $d \neq 1$) previous equation is reduced to the equation

$$\partial_m g_{kn} = \gamma_{kmn} + \gamma_{nmk} + g_{kn}(S_m - \tilde{S}_m).$$

And at last, summing up this equation by k and n (with help of g^{kn}) we obtain $(d-2)(S_m - \tilde{S}_m) = 0$, so for $d > 2$ this equation is reduced to the metricity condition (13).

Thus, due to the Palatini equation in the affine frame (20), the metricity condition (13) and the non-torsionity condition (12) follow one from the other.

In what follows, we assume fulfilling conditions (12) and (13) (which always take place in the Riemannian space), and hence the Palatini equation (20).

3.3. Field equations

In this subsection we show that the *field equation in GRAF is the Einstein equation in an affine frame*.

Both variational derivatives in formula (19) are expressed by the symmetric Einstein tensor in our case (under the conditions (12) and (13)) $G_{mn} = R_{mn} - \frac{1}{2}g_{mn}R$

$$\nabla_{h_\nu^n} L_\gamma = eG_n^\nu, \quad \nabla_{g_{mn}} L_\gamma = \frac{1}{2}eG^{mn}$$

so $h_\nu^m \nabla_{h_\nu^n} L_\gamma = 2g_{ns} \nabla_{g_{ms}} L_\gamma$. This indicates that the number of fields h_μ^m and g_{mn} is redundant to find the equation of motion of the gravitational field in GRAF. For example, we can choose h_μ^m as basic field variables and consider g_{mn} as parameters. In particular, g_{mn} may be a metric η_{mn} of a flat space-time corresponding to the choice of orthonormal frame fields, or a metric of any other (pseudo)Riemannian space-time, such as AdS space-time. The space-time with

the metric g_{mn} can be considered as a vacuum, in which gravitational phenomena are described by the potentials h_μ^m . In the group-theoretical interpretation of gravitational field as a gauge field [18], the gravitational field acts as a further deformation of the background (vacuum) space-time.

All addends with variational derivatives in formula (19) are reduced to one

$$eG_n^\nu \delta h_\nu^n + \frac{1}{2}eG^{mn} \delta g_{mn} = \frac{1}{2}eG^{\mu\nu} \delta g_{\mu\nu}. \quad (23)$$

Since $\Sigma_m^{n\mu\nu}$ depends only on the fields h_σ^s and g_{ps} and not on their derivatives, we have

$$\delta \Sigma_m^{n\mu\nu} = \partial_{h_\sigma^s} \Sigma_m^{n\mu\nu} \delta h_\sigma^s + \partial_{g_{ps}} \Sigma_m^{n\mu\nu} \delta g_{ps}.$$

Directly from the definition of $\Sigma_m^{n\mu\nu}$ we find

$$\begin{aligned} \partial_{h_\sigma^s} \Sigma_m^{n\mu\nu} &= h_s^\sigma \Sigma_m^{n\mu\nu} + h_s^\mu \Sigma_m^{n\nu\sigma} + h_s^\nu \Sigma_m^{n\sigma\mu} \\ \partial_{g_{ps}} \Sigma_m^{n\mu\nu} &= -g^{n\{p} \Sigma_m^{s\}\mu\nu} + \frac{1}{2}g^{ps} \Sigma_m^{n\mu\nu}. \end{aligned} \quad (24)$$

Braces here mean symmetrization by the indexes contained therein. So

$$\delta \Sigma_m^{n\mu\nu} \gamma_{\nu n}^m = B_s^{\sigma\mu} \delta h_\sigma^s + \frac{1}{2}D^{ps\mu} \delta g_{ps},$$

where

$$\begin{aligned} B_s^{\sigma\mu} &:= \partial_{\partial_\mu h_\sigma^s} L_\gamma = \partial_{h_\sigma^s} \Sigma_m^{n\mu\nu} \gamma_{\nu n}^m \\ \frac{1}{2}D^{ps\mu} &:= \partial_{\partial_\mu g_{ps}} L_\gamma = \partial_{g_{ps}} \Sigma_m^{n\mu\nu} \gamma_{\nu n}^m \end{aligned} \quad (25)$$

- generalized momentums of fields h_σ^s and g_{ps} respectively. Using (24) we obtain

$$\begin{aligned} B_s^{\sigma\mu} &= e\gamma_{s.}^{[\sigma\mu]} - h_s^\sigma V_\gamma^\mu + h_s^\mu V_\gamma^\sigma \\ D^{ps\mu} &= -e\gamma^{\mu\{ps\}} + e\gamma_n^{\{p.g^s\}\mu} - g^{ps} V_\gamma^\mu. \end{aligned} \quad (26)$$

Formula (19), taking into account (20), (23) and (25), takes the form

$$\delta' L_\gamma = \frac{1}{2} eG^{\mu\nu} \delta g_{\mu\nu} + \partial_\mu (B_s^{\sigma\mu} \delta h_\sigma^s + \frac{1}{2}D^{ps\mu} \delta g_{ps} + L_\gamma \delta x^\mu). \quad (27)$$

It is interesting to note that generalized momentums can be found solely from the vector density V_γ^μ , which determines the surface part of Hilbert's Lagrangian

$$B_s^{\sigma\mu} = -\partial_{h_\sigma^s} V_\gamma^\mu, \quad \frac{1}{2}D^{ps\mu} = -\partial_{g_{ps}} V_\gamma^\mu.$$

We will note the formula

$$M_m^{n\sigma} = D_m^{n\sigma} - B_m^{n\sigma}$$

which is proved with the help of formulas (21), (26) by the direct substitution.

From the Palatine equation (20) follows $M_m^{n\nu}\gamma_{\nu n}^m = \partial_\mu \Sigma_m^{n\mu\nu} \gamma_{\nu n}^m = 2H_\gamma$, so, provided that it is satisfied

$$L_\gamma = H_\gamma = \Sigma_m^{n\mu\nu} \gamma_{\mu s}^m \gamma_{\nu n}^s. \quad (28)$$

In a coordinate frame, when $\gamma_{\mu n}^s$ is converted to Christoffel symbols $\Gamma_{\mu\nu}^\sigma$, the Lagrangian L_γ coincides with the truncated Hilbert's Lagrangian L_Γ [10]. In an orthonormal frame, when $\gamma_{\mu n}^m$ goes into the Ricci rotational coefficients $\Omega_{\mu n}^m$, the Lagrangian L_γ is the same as Lagrangian L_Ω that was suggested in [5], [3] (this Lagrangian is used in the teleparallel equivalent of general relativity (TEGR), too [11]). In the general case of an affine frame, when applying formula (28) to the Lagrangian L_γ , conditions (12) and (13) must be taken into account. Therefore, the connection coefficients will no longer be arbitrary in this case, and formulas (14) for their constituents must be applied when they are varied.

4. Consequences of Gauge Invariance of GRAF

Hilbert's Lagrangian Λ exhibits uniquely wide gauge symmetry. First of all, Λ is invariant with respect to GL^g -transformations of the affine frame fields, which corresponds to the transition between general reference frames. In addition, Λ is invariant with respect to the gauge translations of the space-time, that is, T_M^g -transformations. Both these groups combine into a semidirect product $S_M^g = GL^g \times T_M^g$, forming a complete symmetry group of GRAF. This symmetry of GRAF is a consequence of the fact that when transforming frame fields

$$h'_{m'}(x') = L_{m'}^n(x) h_n(x), \quad x'^\mu = x^\mu + t^\mu(x) \quad (29)$$

which are parameterized by the functions $g^a(x) = (L_{m'}^n(x), t^\mu(x)) \in S_M^g$, the scalar curvature remains unchanged $R'(x') = R(x)$, and hence $\delta'_g \Lambda = \Lambda'(x')J - \Lambda(x) = 0$. As a result, the equations, which are obtained from the Lagrangian L_γ , are S_M^g -invariant.

Transformations (29) uniquely determine the way in which the semidirect product of the groups GL^g and T_M^g is formed, defining action $\phi_{t^\mu(x)} L_{m'}^n(x) = L_{m'}^n(x + t(x))$. The transformations from the group S_M^g for the field variables of GRAF are

$$\begin{aligned} h'^{m'}_\mu(x') &= L'^{m'}_k(x) h^k_\nu(x) \partial'_\mu x^\nu, & g'^{m'n'}(x') &= L'^k_{m'}(x) L'^l_{n'}(x) g_{kl}(x) \\ \gamma'^{m'}_{\mu\nu}(x') &= [L'^{m'}_k(x) L'^l_{n'}(x) \gamma^k_{\nu l}(x) + L'^{m'}_k(x) \partial_\nu L'^k_{n'}(x)] \partial'_\mu x^\nu \end{aligned} \quad (30)$$

where $L_k^{m'}$ and $L_{n'}^k$ are mutually inverse matrices $L_k^{m'} L_{n'}^k = \delta_{n'}^{m'}$ which are distinguished from each other by the order of indexes.

Note that transformations from S_M^g are considered here from an active point of view, that is, as real local linear transformations of tangent spaces TM_x (rather than their reparameterization) and real displacements of points in the space-time manifold M in fixed coordinates (rather than changing coordinates).

4.1. Generalized principle of relativity

In this subsection we study the consequences of GL^g -invariance of GRAF. Here we show that *the Palatini equation is the strong Noether identity which follows from GL^g -invariance of GRAF.*

A distinctive feature of GRAF is the interpretation of reference frames as anholonomic in the general case affine frame fields (*generalized reference frames*), which is broader than in the EG. This extension of the concept of reference frames in GRAF leads to the extension of the general principle of relativity, which in the framework of GRAF is interpreted as the invariance of gravitational field equations with respect to the transitions between the generalized reference frames, which form the group GL^g of local linear transformations of affine frame fields. The group GL^g is broader than the group H^g of diffeomorphisms of Lagrangian coordinates describing transitions between holonomic frame fields accepted in the EG as reference frames, or than the group O^g of local Lorentz transformations describing transitions between orthonormal frame fields accepted as reference frames in GROF. Both of these groups are subgroups of the group GL^g , so the symmetry of GRAF is wider than the symmetry of EG or GROF.

The generalized principle of (general) relativity states that the gravitational field equations must have the same appearance in all generalized reference frames (affine frames), that is, be GL^g -invariant.

Consider the GL^g -transformations $L_k^m \approx \delta_k^m + l_k^m$ with infinitesimal parameters l_k^m . In this case formulas (30) take the form

$$\begin{aligned} \delta_l h_\mu^m &= l_k^m h_\mu^k, & \delta_l g_{mn} &= -l_m^k g_{kn} - l_n^k g_{mk} \\ \delta_l \gamma_{\mu n}^m &= l_k^m \gamma_{\mu n}^k - l_n^k \gamma_{\mu k}^m - \partial_\mu l_n^m. \end{aligned} \quad (31)$$

From these formulas, taking into account (3) and (18), it follows

$$\delta_l V_\gamma^\sigma = -\sum_m^{\nu\sigma} \partial_\nu l_n^m. \quad (32)$$

Here δ_l is the variation of the shape of fields at infinitesimal GL^g -transformations, and, due to $\delta_l x^\mu = 0$, $\delta'_l = \delta_l$.

The transformations (6) for $g^a \sim l^m_n$ take the form

$$\delta_l h^m_\mu = a^m_{\mu n} l^n_k, \quad \delta_l g_{mn} = a_{mnp} l^p_k, \quad \delta_l V^\sigma = d_m^{n\nu\sigma} \partial_\nu l^m_n$$

where in our case of transformations (31), (32)

$$a^m_{\mu n} = \delta^m_n h^k_\mu, \quad a_{mnp}{}^k = -g_{pm} \delta^k_n - g_{pn} \delta^k_m, \quad d_m^{n\nu\sigma} = -\Sigma_m^{n\nu\sigma} \quad (33)$$

and all the relevant coefficients $b^{i\mu}_a = 0$ and $c^\sigma_a = 0$.

From the definitions (8), taking into account (25), subsequent expressions for the Noether currents and their superpotentials related to GL^g -transformations follow

$$J_m^{n\mu} = -B_s^{\sigma\mu} a_{\sigma m}^s{}^n - \frac{1}{2} D^{ps\mu} a_{psm}{}^n, \quad S_m^{n\mu\nu} = -d_m^{n\mu\nu}$$

which with the help of (33) are specified as follows

$$J_m^{n\mu} = D_m^{n\mu} - B_m^{n\mu} = M_m^{n\mu}, \quad S_m^{n\mu\nu} = \Sigma_m^{n\mu\nu}. \quad (34)$$

The partition (16) of the Hilbert's Lagrangian Λ into the volume L_γ and the surface $\partial_\sigma V_\gamma^\sigma$ parts is not GL^g -invariant and when changing a general reference frame between them there is redistribution with conservation

$$\delta'_l \Lambda = \delta'_l L_\gamma + \partial_\sigma \delta'_l V_\gamma^\sigma = 0. \quad (35)$$

Since for GL^g -transformations $\delta_l x^\mu = 0$ and $\delta_l g_{\mu\nu} = 0$, from (27), taking into account the expressions (31), it follows: $\delta_l L_\gamma = -\partial_\sigma (M_m^{n\sigma} l^m_n)$, which together with (32) and (35), gives the GL^g -invariance condition of the Hilbert's Lagrangian in (pseudo)Riemannian space

$$\delta_l \Lambda = \partial_\sigma (\Sigma_m^{n\nu\sigma} \delta_l \gamma^m_{\nu n}) = -\partial_\sigma (M_m^{n\sigma} l^m_n + \Sigma_m^{n\nu\sigma} \partial_\nu l^m_n) = 0. \quad (36)$$

It is interesting to note that due to the fact that $\delta_l g_{\mu\nu} = 0$, in expression (36) there are no variational derivatives, so all the identities obtained from (36) are strong

$$\partial_\sigma M_m^{n\sigma} = 0, \quad \partial_\sigma \Sigma_m^{n\nu\sigma} + M_m^{n\nu} = 0, \quad \Sigma_m^{n\nu\sigma} = -\Sigma_m^{n\sigma\nu}. \quad (37)$$

The second identity (37) coincides with the Palatini equation (20), which, as we can see, is a consequence of GL^g -invariance of the theory and is an expression of the corresponding Noether current through the superpotential.

Condition (36) of GL^g -invariance of GRAF for contravariant parameters l^{mn} of group GL^g takes the form

$$\delta_l \Lambda = -\partial_\sigma (N_{mn}{}^\sigma l^{mn} + \Sigma_{mn}{}^{\nu\sigma} \partial_\nu l^{mn}) = 0 \quad (38)$$

where $N_{mn}{}^\sigma := M_{mn}{}^\cdot{}^\sigma + \Sigma_m{}^{s\nu\sigma} \partial_\nu g_{sn}$. So the Palatini equation can be represented as

$$\partial_\sigma \Sigma_{mn}{}^\cdot{}^{\nu\sigma} + N_{mn}{}^\nu = 0. \quad (39)$$

By the metricity condition (13) we obtain

$$N_{mn}{}^\sigma = \Sigma_{ms}{}^\cdot{}^{\nu\sigma} \gamma^s{}_{\nu n} - \Sigma_{ns}{}^\cdot{}^{\nu\sigma} \gamma^s{}_{\nu m}$$

hence $N_{mn}{}^\sigma$, like $\Sigma_{mn}{}^\cdot{}^{\nu\sigma}$, are antisymmetric in the lower indices. In this case, condition (38) of the GL^g -invariance of GRAF includes only the antisymmetric components $\omega^{mn} := l^{[mn]}$ of parameters of the group GL^g . The transformations with parameters ω^{mn} , according to (31), do not change the metric in the affine frame $\delta_\omega g_{mn} = -\omega_{mn} - \omega_{nm} = 0$, and therefore correspond to local (pseudo)rotations. Thus, current $N_{mn}{}^\nu$ has the physical meaning of the tensor density of the internal angular momentum (spin) of the gravitational field, and $\Sigma_{mn}{}^\cdot{}^{\nu\sigma}$, according to (39), has the physical meaning of its superpotential. Given also the non-torsionity condition (12) we obtain

$$N_{mn}{}^\sigma = \frac{\epsilon}{2} (F_{mn}^\sigma - h_m^\sigma \tilde{R}_n + h_n^\sigma \tilde{R}_m).$$

Transformations with symmetric components $\sigma^{mn} := l^{\{mn\}}$ of parameters of the group GL^g lead to deformations of the metric in the affine frame $\delta_\sigma g_{mn} = -2\sigma_{mn}$ with the infinitesimal strain tensor σ_{mn} . The Noether current

$$J_{\{mn\}}{}^\sigma = M_{\{mn\}}{}^\cdot{}^\sigma + \partial_\nu g_{s\{m} \Sigma_n{}^{s\nu\sigma} \quad (40)$$

which is connected with σ^{mn} , in the case of metricity (13) equals zero. This is another argument in favor of the possibility of arbitrary choice of vacuum (with metric g_{mn}) on the basis of which gravitational phenomena are considered. This additional possibility may be useful when looking for new solutions in gravity theory.

4.2. Gauge translational invariance

A gravitational field is born by an energy-momentum which associated with space-time translations. So gravity is a gauge theory of the translations group. This, regardless of the particular type of Lagrangian, determines the structure of theory. In particular, it permits to *represent the gravitational field equations in superpotential form*, that is, in the form that analogous to the form of Maxwell dynamic equations.

In this subsection we study the consequences of T_M^g -invariance of GRAF.

From formulas (30) follows, that T_M^g -transformations with infinitesimal parameters $\delta_t x^\mu = t^\mu$ are

$$\begin{aligned}\delta_t h_\mu^m &= -t^\sigma \partial_\sigma h_\mu^m - h_\sigma^m \partial_\mu t^\sigma, & \delta_t g_{mn} &= -t^\sigma \partial_\sigma g_{mn} \\ \delta_t \gamma_{\mu n}^m &= -t^\sigma \partial_\sigma \gamma_{\mu n}^m - \gamma_{\sigma n}^m \partial_\mu t^\sigma.\end{aligned}\quad (41)$$

The partition (16) of the Hilbert's Lagrangian Λ is T_M^g -invariant, because $\delta'_t L_\gamma = 0$ and $\delta'_t V_\gamma^\sigma = 0$. Therefore, the surface addend of the Hilbert's Lagrangian is not affected on the consequences of T_M^g -invariance of the GRAF equations. The parameters t^μ describe the infinitesimal displacements along the Eulerian coordinates. They correspond to displacements $t^m = h_\mu^m t^\mu$ along the affine basis vectors that parameterize the deformed gauge group of translations T_M^{gh} [20]. Transformations (41) in terms of parameters t^m of the group T_M^{gh} take the form

$$\delta_t h_\mu^m = -F_{\mu n}^m t^n - \partial_\mu t^m, \quad \delta_t g_{mn} = -\partial_s g_{mn} t^s. \quad (42)$$

Transformations (6) for $g^a \sim t^m$ give

$$\delta_t h_\mu^m = a_{\mu n}^m t^n + b_{\mu n}^{m\nu} \partial_\nu t^n, \quad \delta_t g_{mn} = a_{mns} t^s.$$

So for T_M^{gh} -transformations (42) we obtain

$$a_{\mu n}^m = -F_{\mu n}^m, \quad a_{mns} = -\partial_s g_{mn}, \quad b_{\mu n}^{m\nu} = -\delta_n^m \delta_\mu^\nu \quad (43)$$

and all other coefficients (except h_m^μ) equal zero. From the definitions (8), taking into account (13), (25), (43), follow the subsequent expressions for the Noether current t_m^μ and superpotential $S_m^{\mu\nu}$ related to T_M^{gh} -transformations

$$t_m^\mu = B_s^{\sigma\mu} F_{\sigma m}^s + D_s^{\cdot p\mu} \gamma_{mp}^s - L_\gamma h_m^\mu, \quad S_m^{\mu\nu} = B_m^{\mu\nu}. \quad (44)$$

Therefore, t_m^μ determines the *energy-momentum of the gravitational field* in the affine frame h_m . The current t_m^μ (as well as the superpotential $B_m^{\mu\nu}$) is a mixed coordinate-frame tensor density with respect to the gauge translations (at which diffeomorphisms of Eulerian coordinates occur) and global linear transformations of the frame fields. In the case of gauge linear transformations of frame fields, the current t_m^μ (as well as the superpotential $B_m^{\mu\nu}$) is transformed by the non-tensor law [15]

$$\begin{aligned}t_{m'}^\mu &= L_{m'}^m (t_m^\mu - \partial_\sigma \Delta B_m^{\mu\sigma}) - \partial_\sigma L_{m'}^m (B_m^{\mu\sigma} + \Delta B_m^{\mu\sigma}) \\ B_{m'}^{\mu\sigma} &= L_{m'}^m (B_m^{\mu\sigma} + \Delta B_m^{\mu\sigma})\end{aligned}\quad (45)$$

where $\Delta B_m^{\mu\sigma} := e(\Delta\gamma_m^{[\mu\sigma]} - h_m^\mu \Delta\gamma_n^{[n\sigma]} + h_m^\sigma \Delta\gamma_n^{[n\mu]})$, $\Delta\gamma_{\nu n}^m := \partial_\nu L_{s'}^m L_n^{s'}$. This is due to the extra energy-momentum that must be transmitted for the transition between mutually accelerated reference frames [11].

Note that to denote quantities having different tensor dimensions, one letter can be used without danger of confusing these quantities, such as the translation parameters t^m and the tensor density t_m^σ of the energy-momentum of the gravitational field.

Taking into account (27) and (42), the T_M^{gh} -invariance condition of GRAF is written as

$$\delta'_t \Lambda = -eG_n^\nu (\gamma_{\nu m}^n t^m + \partial_\nu t^n) - \partial_\nu (t_m^\nu t^m + B_m^{\mu\nu} \partial_\mu t^m) = 0 \quad (46)$$

so we obtain the following strong Noether identities

$$-eG_n^\nu \gamma_{\nu m}^n = \partial_\nu t_m^\nu, \quad -eG_m^\mu = \partial_\nu B_m^{\mu\nu} + t_m^\mu, \quad B_m^{\mu\nu} = -B_m^{\nu\mu}. \quad (47)$$

The second identity (47) can be represented in the form (10), which in our case gives

$$t_m^\nu = -\partial_{h_\nu^m} L_\gamma.$$

In the presence of matter fields with the tensor density τ_m^μ of the energy-momentum, when Einstein equation $eG_m^\mu = \tau_m^\mu$ are fulfilled, the second identity (47) takes the form

$$\partial_\nu B_m^{\mu\nu} = -T_m^\mu \quad (48)$$

where $T_m^\mu = t_m^\mu + \tau_m^\mu$ is the total tensor density of the energy-momentum of the gravitational and matter fields. Conversely, from the equation (48), given second identity (47), the Einstein equation follows. Therefore, *equation (48) is the Einstein equation in the affine frame*, which is written in the superpotential form, that is, in the form similar to that of Maxwell's dynamic equations (and equations in other gauge theories). The tensor density $B_m^{\mu\nu}$, called the *induction of the gravitational field* in the GRAF, acts as a superpotential of the total energy-momentum T_m^μ of the gravitational and matter fields.

Representation of the Einstein equation in the form (48), similar to the form of field equations of any other gauge theory, is a consequence of T_M^{gh} -symmetry of the theory of gravity. This is convincingly illustrated the fact that gravity, as an interaction generated by energy-momentum, is the gauge theory of the translation group, and the superpotential of the total energy-momentum T_m^μ (induction of the gravitational field $B_m^{\mu\nu}$) acts as the power characteristic of the gravitational field (but not a curvature, as in EG, or a torsion, as in TEGR). Such an interpretation opens new avenues for unification of gravity with other gauge theories by uniting appropriate generalized deformed gauge groups [24], which may be nontrivial [25].

From the equation (48), considering (47), the conservation law follows

$$\partial_\mu T_m^\mu = 0. \quad (49)$$

On the gravitational extremal $eG_m^\mu = \tau_m^\mu$, taking into account (49) $\partial_\sigma t_m^\sigma = -\partial_\sigma \tau_m^\sigma$, first identity (47) goes into the equation of motion of (macro)matter

$$\partial_\mu \tau_m^\mu = \tau_s^\mu \gamma_{\mu s}^s. \quad (50)$$

For the dust matter $\tau_m^\mu = e\mu u^\mu u_m$, where μ is the density of its mass, $u^\mu = \dot{x}^\mu$. Provided that the mass is conserved $\partial_\mu(e\mu u^\mu) = 0$ (that is, in the absence of reactive forces), equation (50) is the equation of geodesic $\dot{u}^m + \gamma_{sn}^m u^s u^n = 0$ on which the particles move.

4.3. Complete symmetry of the general relativity

Under energy-momentum we understand a Noether current which is conserved as a result of the space-time translational invariance. But translations can simultaneously cause GL^g -transformations of the reference frame (as in the case of EG), which lead to the energy-momentum renormalization. This situation corresponds to the deformation of the group $S_M^{gh} \rightarrow S_M^{gH}$ [23]. The multiplicity of deformed specimens of the group S_M^{gH} leads to the multiplicity of variants of Noether currents which are conserved due to translational invariance of GRAF, hence leading to a *plurality of ways of determining the energy-momentum of the gravitational field and the corresponding superpotential*. Among all deformed groups S_M^{gH} there is one highlighted, so-called group of parallel transports DP , transformations of which are consistent with the geometric structure of a (pseudo)Riemannian space [23]. This group corresponds to the energy-momentum tensor density, which is characterized by its dependence of the derivatives of connection coefficients, so the energy-momentum in this case is localizable.

In this subsection, we explore these issues.

Consider the infinitesimal variant of transformations of a complete symmetry group $S_M^{gh} = GL^g \times T_M^{gh}$, considering both transformations (31) of the subgroup GL^g and transformations (42) of the subgroup T_M^{gh} simultaneously $\delta_g = \delta_l + \delta_t$

$$\delta_g h_\mu^m = l^m_k h_\mu^k - F_{\mu n}^m t^n - \partial_\mu t^m, \quad \delta_g g_{mn} = -l^k_m g_{kn} - l^k_n g_{mk} - \partial_s g_{mn} t^s. \quad (51)$$

The S_M^{gh} -invariance condition of GRAF is obtained by combining conditions (36) and (46)

$$\begin{aligned} \delta'_g \Lambda = & -eG_n^\nu (\gamma_{\nu m}^n t^m + \partial_\nu t^n) \\ & - \partial_\sigma (t_m^\sigma t^m + B_m^{\nu\sigma} \partial_\nu t^m + M_m^{n\sigma} l_n^m + \Sigma_m^{n\nu\sigma} \partial_\nu l_n^m) = 0. \end{aligned} \quad (52)$$

We deform the group $S_M^{gh} \rightarrow S_M^{gH}$, assuming that an additional linear transformation $dl_n^m = -\lambda_{sn}^m t^s$ occurs at the translations, that is, introducing instead of l_n^m the new parameters of infinitesimal linear transformations

$$r_n^m = l_n^m + \lambda_{sn}^m t^s \quad (53)$$

with arbitrary functions λ_{sn}^m . In this parameterization (that is, for the group S_M^{gH}), transformations (51) take the form

$$\begin{aligned} \delta_g h_\mu^m &= r_k^m h_\mu^k - (F_{\mu n}^m + \lambda_{n\mu}^m) t^n - \partial_\mu t^m \\ \delta_g g_{mn} &= -r_m^k g_{kn} - r_n^k g_{mk} - (\partial_s g_{mn} - \lambda_{msn} - \lambda_{nsm}) t^s. \end{aligned} \quad (54)$$

The deformation of the group S_M^{gh} changes the way of forming the semidirect product of the groups GL^g and T_M^{gh} , which is determined by the transformations (54) [23].

The S_M^{gH} -invariance condition of the GRAF can be written in the form

$$\begin{aligned} \delta'_g \Lambda &= -eG_n^\nu (\gamma_{\nu m}^n t^m + \partial_\nu t^n) \\ &\quad - \partial_\sigma \left(\left(t_m^\sigma + \partial_\nu \left(e\lambda_{m\cdot}^{[\sigma \nu]} \right) \right) t^m + \left(B_m^{\nu\sigma} - e\lambda_{m\cdot}^{[\nu \sigma]} \right) \partial_\nu t^m \right) \\ &\quad + \partial_\sigma (M_m^{n\sigma} r_n^m + \Sigma_m^{n\nu\sigma} \partial_\nu r_n^m) = 0 \end{aligned} \quad (55)$$

which follows from condition (52), taking into account (53) and the Palatini equation (20). Therefore, during the deformation of the group S_M^{gh} , the energy- momentum of the gravitational field and correspondent superpotential are renormalized

$$\begin{aligned} t_m^\nu &\rightarrow J_m^\nu = t_m^\nu + \partial_\sigma \left(e\lambda_{m\cdot}^{[\nu \sigma]} \right), \quad B_m^{\nu\sigma} \rightarrow S_m^{\nu\sigma} = B_m^{\nu\sigma} - e\lambda_{m\cdot}^{[\nu \sigma]} \\ S_m^{\nu\sigma} &= e\gamma_{m\cdot}^{[\nu \sigma]} - e\lambda_{m\cdot}^{[\nu \sigma]} - h_m^\nu V_\gamma^\sigma + h_m^\sigma V_\gamma^\nu. \end{aligned} \quad (56)$$

Taking into account second identity (47), we see that the deformation of the group S_M^{gh} leads to the redistribution between the divergence of superpotential and the tensor density of the energy-momentum of the gravitational field in the Einstein tensor

$$-eG_m^\mu = \partial_\nu B_m^{\mu\nu} + t_m^\mu = \partial_\nu S_m^{\mu\nu} + J_m^\mu$$

illustrating, in the case of the group S_M^{gH} , the general position that take place for generalized deformed gauge groups and which was indicated by formulas (11) and the text after them.

Special choice of λ_{sn}^m allows to discover a lot of "new internal gauge symmetries" in the theory of gravity [13]. Now let

$$\lambda_{sn}^m = \gamma_{sn}^m. \quad (57)$$

In this case, given the non-torsionity (12) and metricity (13) conditions, transformations (54) are greatly simplified

$$\delta_g h_\mu^m = r^m_k h_\mu^k - D_\mu t^m, \quad \delta_g g_{mn} = -r^k_m g_{kn} - r^k_n g_{mk} \quad (58)$$

where $D_\mu t^m = \partial_\mu t^m + \gamma_{\mu n}^m t^n$ is the covariant derivative of the vector t^m . Note that in pure translations from the group S_M^{gH} (at $r^m_k = 0$) the metric in the affine frame does not change

$$\delta_t g_{mn} = 0.$$

In addition, from (58) in this case we obtain

$$h'_m(x') = h_m(x) + t^s \gamma_{sm}^n h_n(x),$$

which corresponds to the definition of the affine connection by the method of Cartan's moving frame. The group S_M^{gH} , provided (57), is the group DP of parallel transports in the Riemannian space [23]. Transformations from the DP set on the M the structure of the (pseudo)Riemannian space with the metric g_{mn} in the affine frame h_m . Note that translations do not form a subgroup in DP since $[D_\mu D_\mu]_n^m = R_{n\mu\nu}^m \in AlGL^g$.

From (56) it follows that for DP

$$J_m^\nu = t_m^\nu + e D_\sigma \gamma_{m \cdot}^{[\nu \sigma]}, \quad S_m^{\nu\sigma} = -h_m^\nu V_\gamma^\sigma + h_m^\sigma V_\gamma^\nu = -\delta_{m\mu}^{\nu\sigma} V_\gamma^\mu \quad (59)$$

where $D_\sigma \gamma_{m \cdot}^{[\nu \sigma]} = \frac{1}{e} \partial_\sigma (e \gamma_{m \cdot}^{[\nu \sigma]})$ is the covariant divergence of the antisymmetric T_M^g -tensor $\gamma_{m \cdot}^{[\nu \sigma]}$. It is interesting to note that in case of group DP , superpotential $S_m^{\nu\sigma}$ depends solely of vector density V_γ^μ , that determines the surface part of Hilbert's Lagrangian Λ . In addition, in expression (59) for J_m^ν there are derivatives of $\gamma_{m \cdot}^{[\nu \sigma]}$, so the energy-momentum of the gravitational field in this case is localizable.

5. Special Gauge Conditions

The broad symmetry of GRAF is provided by a sufficient number of field variables of the theory. Limiting such symmetry by imposing gauge conditions reduces the number of field variables. *Different variants of the gauge conditions lead to different locally equivalent formulations of the general relativity*, the most characteristic of which we consider in this section.

Keeping the status of gravity as a gauge theory of a translation group, we will not consider the gauge conditions that violate the T_M^g -invariance of the theory. We

restrict ourselves to the gauge conditions that narrow the GL^g -invariance of GRAF, which corresponds to the choice of special classes of generalized reference frames (affine frames).

5.1. Einstein gravity

From the outset, general relativity was formulated by Einstein in holonomic reference frames, which are described by frame fields with zero anholonomic objects, and can therefore be implemented as coordinate frame fields of certain coordinate systems. In this case, the gauge condition that limits the GL^g -invariance of the theory is the holonomicity condition

$$F_{\mu\nu}^m = \partial_\nu h_\mu^m - \partial_\mu h_\nu^m = 0 \quad (60)$$

which ensures that there are functions $y^m(x)$ such that $h_\mu^m = \partial_\mu y^m$. The quantities y^m which give holonomic reference frames play the role of Lagrangian coordinates, in contrast to the coordinates x^μ in space-time, which numerate its points and play the role of Eulerian coordinates. Therefore, the functions $y^m(x)$ are scalars with respect to the transformations of the coordinates $x'^\mu = x'^\mu(x)$, that is, invariant under the gauge translations of the space-time $x'^\mu = x^\mu + \delta x^\mu(x)$: $y'^m(x') = y^m(x)$, which, in fact, allows them to be interpreted as Lagrangian coordinates of the gravitational system. The vectors h_m of holonomic frames are the coordinate vectors ∂_m of the coordinates y^m .

Unlike Eulerian coordinates x^μ , numerating space-time points and which changes should not alter any physical quantities, changes of Lagrangian coordinates y^m alter the physical reference frame and certain physical quantities may vary. Thus, during the transition between mutually accelerated reference frames, the energy-momentum must change. If the transformations of the world coordinates x^μ correspond to the gauge translations in space-time in a fixed general reference frame and belong to the group T_M^g , then with the changes of the Lagrangian coordinates y^m the translations do not occur, but transformations of holonomic reference frames take place. Such transformations belong to another group, namely the group GL^g , forming its subgroup H^g which is isomorphic to the group of diffeomorphisms of space-time T_M^g . Despite the isomorphism of the groups H^g and T_M^g , they have different physical meaning and their identification leads to known difficulties in determining the energy-momentum of the gravitational field.

The symmetry group of EG is $\bar{S}_M^g = H^g \times T_M^g$.

Due to the condition (60), in holonomic reference frames $\omega_{kn}^m = 0$, and connection coefficients γ_{kn}^m are reduced to $\Gamma_{kn}^m = \sigma_{kn}^m$, which are Christoffel symbols in Lagrangian coordinates y^m , and thus behave as scalars when transformation of

Eulerian coordinates x^μ occurs. As for an arbitrary affine frame, this fact provides a general-covariance (gauge translational invariance) of the partition (16) of the Hilbert's Lagrangian into bulk and surface parts $\Lambda = L_\Gamma + \partial_\sigma V_\Gamma^\sigma$ [10], where

$$L_\Gamma = \Sigma_m^{n\mu\nu} \Gamma_{\mu s}^m \Gamma_{\nu n}^s = e \Gamma_{m s}^{[m} \Gamma_n^{s|n]}, \quad V_\Gamma^\sigma = \Sigma_m^{n\nu\sigma} \Gamma_{\nu n}^m = e \Gamma_n^{[n\sigma]} \quad (61)$$

(vertical bars highlight indices that are skipped during antisymmetrization).

The action associated with the volume Lagrangian L_Γ can be represented as an integral with respect to Lagrangian coordinates y^m

$$S_\Gamma = \int L_\Gamma dx = \frac{1}{2} \int \delta_{mn}^{kp} \Gamma_{k s}^m \Gamma_p^{s n} v dy$$

where dx and $dy = \partial y dx$ are the coordinate volumes of the Eulerian and Lagrangian coordinates, $v := \sqrt{|g_{mn}|}$, $\partial y := |\partial_\mu y^m|$, $e = \partial y v$. So, action S_Γ is obviously invariant under the transformations of the world (Eulerian) coordinates x^μ .

The tensor density of the energy-momentum of the gravitational field (44), by condition (60), is given by

$$t_m^\mu = \partial_p x^\mu (D_s^{n \cdot p} \Gamma_{nm}^s - L_\Gamma \delta_m^p).$$

In our case, second formula (26) is reduced to

$$D^{nsp} = \partial y v (-\Gamma^{pns} \cdot + \Gamma_k^{k \{n} g^{s\} p} - g^{ns} \Gamma_k^{[k p]})$$

therefore

$$t_m^\mu = \partial y \partial_p x^\mu \bar{t}_m^p \quad (62)$$

where

$$\bar{t}_m^p = v (-\Gamma_{sn}^p \Gamma_m^{sn} + \Gamma_{kn}^k \Gamma_m^{\{pn\}} - \Gamma_n^{[np]} \Gamma_{km}^k - \Gamma_{ks}^{[k} \Gamma_n^{s|n]} \delta_m^p). \quad (63)$$

The induction of the gravitational field (26) in holonomic frames is specified as follows

$$B_m^{\mu\sigma} = \partial y \partial_p x^\mu \partial_s x^\sigma \bar{B}_m^{ps} \quad (64)$$

where

$$\bar{B}_m^{ps} = v (\Gamma_m^{[ps]} \cdot - \delta_m^p \Gamma_n^{[n s]} + \delta_m^s \Gamma_n^{[np]}). \quad (65)$$

In relation to the transformations of Eulerian coordinates x^μ , the energy-momentum of the gravitational field, according to (62), is described by four vector densities t_m^σ , and the induction of the gravitational field given by formula (64) is described by four antisymmetric tensor densities $B_m^{\sigma\mu}$, which correspond to four Lagrangian coordinates y^m . In the case of nonlinear transformations of Lagrangian coordinates

$y^{m'} = y^{m'}(y)$, both t_m^σ and $B_m^{\sigma\mu}$ are transformed by non-tensor laws specifying formulas (45) for our case with $L_m^{m'} = \partial_m y^{m'}$.

In the case of identification of Lagrangian coordinates with Eulerian ones $y^m = x^m$, t_m^σ goes to the Einstein's complex \bar{t}_m^p of the energy-momentum of the gravitational field, and $B_m^{\mu\sigma}$ to the Freud's superpotential \bar{B}_m^{ps} [22] (this fact was first noted in [15]). In this case, the tensor properties of \bar{t}_m^p and \bar{B}_m^{ps} at the transformations of the coordinates x^m are violated precisely because of the identification $y^m = x^m$, which rigidly links translations in space-time with the changes of reference frames. Einstein's equation in superpotential form $\partial_\nu B_m^{\mu\nu} = -t_m^\mu - \tau_m^\mu$ (48) in holonomic frames can be represented solely in Lagrangian coordinates by means of Freud's superpotential \bar{B}_m^{ps} and the Einstein's complex \bar{t}_m^p as follows

$$\partial_s \bar{B}_m^{ps} = -\bar{t}_m^p - \bar{\tau}_m^p \quad (66)$$

where $\bar{\tau}_m^p = \partial y^{-1} \partial_\mu y^p \tau_m^\mu$ [22].

Taking into account the fact that $\Gamma_{sn}^s = \frac{1}{2} g^{pk} \partial_n g_{pk} = \frac{1}{v} \partial_n v$, as well as condition (60), in holonomic reference frames the tensor density of the internal angular momentum (spin) of the gravitational field and its superpotential can be represented as $N_{mn}{}^\nu = -\partial y \partial_{[m} x^\nu \partial_{n]} v$, $\Sigma_{mn}{}^{\nu\sigma} = \frac{1}{2} \partial y v \delta_{mn}^{\nu\sigma}$.

Since holonomic reference frames are completely determined by the functions $y^m = y^m(x)$ which bind the Lagrangian and Eulerian coordinates, they can be selected as independent field variables. The Lagrangian L_Γ (61), like the complete Lagrangian of the gravitational system $L = L_\Gamma + L_\psi$ (where L_ψ is the Lagrangian of matter fields), does not depend on the functions $y^m(x)$ itself, but only on their derivatives. Therefore, the fields $y^m(x)$ are cyclic coordinates of the system and the equations of motion for them are reduced to the condition of preserving the corresponding generalized momentum $P_m^\sigma = \partial_{\partial_\sigma y^m} L$: $\partial_\sigma P_m^\sigma = 0$, which (up to an overall sign) coincides with the tensor density of energy-momentum of the gravitational system: $P_m^\sigma = -T_m^\sigma$.

At the end of this subsection, we consider the transformation of gauge translations along Lagrangian coordinates, which illustrates the significant physical difference between Lagrangian and Eulerian coordinates.

Gauge translations along the Lagrangian coordinates $\delta_d y^m = d^m$ form ordinary gauge group $D^g = \sum_{x \in M} T \subset GL^g$ and correspond to the transformations of the frame fields with infinitesimal parameters $l_n^m = \partial_n d^m$, therefore they are given by the formulas

$$\delta_d x^\mu = 0, \quad \delta_d h_\mu^m = \partial_\mu d^m, \quad \delta_d g_{mn} = -2g_{k\{m} \partial_{n\}} d^k$$

which follow from formulas (31) with $l_n^m = \partial_n d^m$ (compare with (42)). Invariance with respect to such transformations leads to identity

$$\delta_d \Lambda = -\partial_\sigma (C_m^{\mu\sigma} \partial_\mu d^m + \Sigma_m^{\mu\nu\sigma} \partial_{\mu\nu}^2 d^m) = 0$$

which follows from identity (36) for $l_n^m = \partial_n d^m$, where $C_m^{\mu\sigma} = M_m^{\mu\sigma} + \Sigma_m^{\nu\sigma} \partial_\nu h_n^\mu$. It follows that the current associated with D^g -transitions along the Lagrangian coordinates, like the corresponding superpotential $S_m^{\mu\nu} = C_m^{\mu\nu} + \partial_\sigma \Sigma_m^{\mu\nu\sigma}$, is equal to zero.

5.2. General relativity in an orthonormal frame

The disadvantage of holonomic reference frames is their inevitable deformation in the curve space during the transition between the reference frames with the infinitesimal strain tensor $\sigma^{mn} = g^{s\{m} \partial_s \delta y^{n\}}$. Such a drawback is obviously absent in general reference frames for which $g_{mn} = \text{const}$, in particular in (pseudo)orthonormal reference frames, when $g_{mn} = \eta_{mn}$. In this case $\sigma_{kn}^m = 0$, $e = h := |h_\mu^m|$ and the connection coefficients γ_{kn}^m are reduced to the quantities $\Omega_{kn}^m = \omega_{kn}^m$, which is Ricci rotation coefficients. When we consider only orthonormal reference frames, the group GL^g narrows to the group O^g of gauge Lorentz transformations. The symmetry group of GROF is $\hat{S}_M^g = O^g \times T_M^g$.

The Lagrangian of general relativity in an orthogonal frame and the corresponding vector defining the surface term are written as [12], [5]

$$L_\Omega = \Sigma_m^{\nu\mu} \Omega_{\mu s}^m \Omega_{\nu n}^s = e \Omega_{m s}^{[m} \Omega_n^{s]n}, \quad V_\Omega^\sigma = \Sigma_m^{\nu\sigma} \Omega_{\nu n}^m = e R^\sigma$$

where $R_n = \nabla_\sigma h_n^\sigma = F_{s n}^s$ is a vector that has an important geometric meaning and determines the rate of change of the local coordinate volume along the directions of basis vectors of the selected orthonormal frame. When writing these formulas, equality $\Omega_{s n}^s = R_n$ is taken into account.

The induction of the gravitational field in orthonormal reference frames is given by

$$B_m^{\mu\sigma} = e (\Omega_{m \cdot}^{\mu \sigma} - h_m^\mu R^\sigma + h_m^\sigma R^\mu). \quad (67)$$

A distinctive feature of orthonormal reference frames is the fact that the connection coefficients are uniquely determined by the induction of the gravitational field, that is, formula (67) can be reversed. Indeed, from (67) follows $B_n^{n\mu} = -(d-2)e R^\mu$, hence

$$e \Omega_{m \cdot}^{\mu \sigma} = B_m^{\mu\sigma} - (h_m^\mu B_n^{n\sigma} - h_m^\sigma B_n^{n\mu}) / (d-2).$$

The energy-momentum tensor density of gravitational field (44) in our case due to $D_s^{n\sigma} = 0$ is determined by the expression [12], [5]

$$t_m^\mu = B_s^{n\mu} F_{nm}^s - L_\Omega h_m^\mu. \quad (68)$$

Einstein equation in GROF specifies equation (48)

$$\partial_\sigma B_m^{\mu\sigma} = -t_m^\mu - \tau_m^\mu$$

where $B_m^{\mu\sigma}$ and t_m^μ are given by formulas (67) and (68) respectively. Note that in this form Einstein equation is represented in TEGR [11].

The transformations of the deformed group $\hat{S}_M^{gH} = O^g \times T_M^g$ under condition (57) (which in our case is $\lambda_{sn}^m = \Omega_{sn}^m$) have the form

$$\delta_g h_\mu^m = \omega^m_k h_\mu^k - D_\mu t^m, \quad \delta_g g_{mn} = 0$$

and coincide (in the absence of torsion) with generalized Poincare gauge transformations [8], [9]. In this case, the energy-momentum of the gravitational field and the corresponding superpotential (59) are given by

$$J_m^\nu = t_m^\nu + e D_\sigma \Omega_m^{\nu\sigma}, \quad S_m^{\nu\sigma} = -e \delta_{m\mu}^{\nu\sigma} R^\mu.$$

The symmetry with respect to O^g -transformations determine the gravitational field spin, which current density and the corresponding superpotential are given by the expressions $N_{mn}^\sigma = \frac{e}{2}(F_{mn}^\sigma - h_m^\sigma R_n + h_n^\sigma R_m)$, $\Sigma_{mn}^{\sigma\nu} = \frac{e}{2} \delta_{mn}^{\sigma\nu}$.

5.3. Dilaton gravity, unimodular gravity, and so on

The Weyl transformations

$$\bar{h}_m \rightarrow h_m = \bar{h}_m / \phi \quad (69)$$

form a subgroup W^g of the group GL^g with parameters $L_m^{m'} = \phi \delta_m^{m'}$ [4]. So dilation gravity as a theory with Weyl symmetry W^g is a special case of GRAF. In this case $\Delta \gamma_{\nu n}^m = -\partial_\nu \ln \phi \delta_n^m$, so $\Delta B_m^{\mu\nu} = 0$, and formulas (45) for the energy-momentum and suitable superpotential of gravitational field in DG give

$$t_m^\mu = (\bar{t}_m^\mu + \partial_\sigma \ln \phi \bar{B}_m^{\mu\sigma}) / \phi, \quad B_m^{\mu\sigma} = \bar{B}_m^{\mu\sigma} / \phi \quad (70)$$

where \bar{t}_m^μ and $\bar{B}_m^{\mu\sigma}$ apply to the frame \bar{h}_m . In the case where $\bar{h}_m = \partial_m$ is the coordinate frame of the Lagrangian coordinates y^m , the relation (70) can be represented with the help of the Einstein energy-momentum complex \bar{t}_m^p (63) and Freud's superpotential \bar{B}_m^{ps} (65) by substituting in formulas (70) $\bar{t}_m^\mu \rightarrow \bar{t}_m^p$, $\bar{B}_m^{\mu\sigma} \rightarrow \bar{B}_m^{ps}$ and entering the quantities $t_m^\mu \rightarrow t_m^p = \partial y^{-1} \partial_\mu y^p t_m^\mu$ and $B_m^{\mu\sigma} \rightarrow B_m^{ps} =$

$\partial y^{-1} \partial_\mu y^p \partial_\sigma y^s B_m^{\mu\sigma}$: the energy-momentum complex and the corresponding superpotential of the gravitational field in DG in the Lagrangian coordinates y^m . In their terms, the gravitational field equation in DG takes the form $\partial_s B_m^{ps} = -t_m^p - \tau_m^p$, where $\tau_m^p = \bar{\tau}_m^p / \phi$.

The symmetry group of DG is $\tilde{S}_M^g = C^g \rtimes T_M^g$, where $C^g = H^g \otimes W^g$.

The scalar field ϕ acts in DG as a finite parameter of dilation symmetry, so equation of motion with respect to ϕ is fulfilled automatically by the Palatini equation.

Moreover, Noether current and corresponding superpotential linked to Weyl transformations (69) are zero due to the equality to zero of the current $J_{\{mn\}}^\sigma$ (40).

Unimodular gravity is a theory where the determinant of the metric is fixed [27]. As mentioned above (in the subsection 3.3), in the GRAF the metric g_{mn} in the affine frame h_m can be considered as an arbitrary parameter of the theory (the metric of the background vacuum space), and therefore its determinant can be arbitrarily selected. In the case of a holonomic frame, that is, in EG, this limits the choice of Lagrangian coordinates by narrowing the EG symmetry subgroup H^g , associated with transitions between reference frames, to the group $SDiff$ of volume-preserving, or 'special' diffeomorphisms. It should be noted that the choice of Eulerian coordinates remains free, in particular, they can be dimensionless.

The symmetry group of UG is $\bar{S}_M^{vg} = SDiff \rtimes T_M^g$.

The gravitational field equations in UG in Lagrangian coordinates can be represented in the same form (66) as in EG, where we should fix v in expressions for the energy-momentum of the gravitational field \bar{t}_m^p (63) and superpotential \bar{B}_m^{ps} (65).

Narrowing down GL^g -symmetry of GRAF to its other subgroups (classes of allowed affine frames), we can obtain other locally equivalent variants of GR.

6. Discussion and Conclusions

In this work, the method of deformations of generalized gauge groups is applied to the study of GR symmetry. It is shown that all quantities which are conserved due to the invariance of the physical theories with respect to generalized gauge groups are quasilocal, that is, they have superpotentials.

GR is formulated in generalized reference frames, which are represented by (anholonomic in the general case) affine frame fields. The general principle of relativity is extended to the requirement of invariance of the theory with respect to

transitions between generalized reference frames, that is, with respect to the group GL^g of local linear transformations of affine frame fields.

GR is interpreted as the gauge theory of the gauge group of translations, and therefore is invariant under the space-time diffeomorphisms.

The consequence of the GL^g -invariance of the general relativity in an affine frame is the Palatini equation, which in the absence of torsion goes into the metricity condition, and vice versa, that is, is fulfilled identically in the Riemannian space.

The consequence of the T_M^g -invariance of GRAF is representation of the Einstein equation in superpotential form, that is, in the form of dynamic Maxwell equations (or Young-Mills equations).

At gauge translations, the energy-momentum of the gravitational field is transformed by the tensor law, and at transitions between relatively accelerated generalized reference frames - by the non-tensor law, which corresponds to the need to attract additional energy-momentum to provide such transitions. The groups GL^g and T_M^g are united into group $S_M^g = GL^g \times T_M^g$, which is their semidirect product and is the complete symmetry group of GRAF. Deformation of the group $S_M^g \rightarrow S_M^{gH}$ leads to renormalisation of energy-momentum of the gravitational field. Among the groups S_M^{gH} there is one DP - the group of parallel transports in the Riemannian space, action of which is agreed with the space geometrical structure. The energy-momentum of the gravitational field for DP is localizable.

Limiting the admissible generalized reference frames by imposing GL^g -gauge conditions on field variables of GRAF leads to various locally equivalent formulations of GR, such as EG, GROF or TEGR, DG, UG, and others. However, these theories can significantly be different by global solutions. So in [7] was suggested that "just like a conformally invariant theory of matter is insensitive to the cosmological singularity, a GL^g -invariant field theory will be insensitive to any singularity". This hope is based on a sufficiently broad gauge symmetry of such theory, which can provide the possibility of finding generalized reference frames in which singularities will be absent, just as singularity on the event horizon in the Schwarzschild solution disappears in the free-falling holonomic reference frame.

The choice of appropriate generalized d-dimensional reference frames in the framework of group-theory approach to unification of gravity with internal symmetry gauge interactions [24], [25] gives hope for getting rid of singularities in other theories of gauge interactions, too.

We are planning to consider the physical arguments regarding the choice of generalized reference frames in the future.

Acknowledgements

The author is grateful to Alisa Gryshchenko for her help with the prepare of article.

References

- [1] Aldrovandi R. and Pereira J., *An Introduction to Teleparallel Gravity*, Springer, Heidelberg 2013.
- [2] Blagojević M. and Hehl F., *Gauge Theories of Gravitation*, World Scientific, London 2013 [arXiv:1210.3775 [gr-qc]].
- [3] Cho Y., *Einstein Lagrangian as the Translational Yang - Mills Lagrangian*, Phys. Rev. D **14** (1976) 2521–2525.
- [4] Dirac P., *Long Range Forces and Broken Symmetries*, Proc. Roy. Soc. Lond. A **333** (1973) 403–418.
- [5] Duan Y. and Zhang J., *Conservation Law of Energy-momentum in General Relativity*, Acta Physica Sinica **11** (1963) 689–704 [arXiv:1803.09651 [gr-qc]].
- [6] Fontanini M., Huguet E. and Le Delliou M., *Teleparallel Gravity (TEGR) as a Gauge Theory: Translation or Cartan Connection?*, Phys. Rev. D **99** (2019) 064006 [arXiv:1811.03810 [gr-qc]].
- [7] Gielen S., de León Ardón R. and Percacci R., *Gravity with More or Less Gauging*, Class. Quant. Grav. **35** (2018) 195009 [arXiv:1805.11626 [gr-qc]].
- [8] Hehl F., von der Heyde P., Kerlick G. and Nester J., *General Relativity with Spin and Torsion: Foundation and Prospects*, Rev. Mod. Phys. **48** (1976) 393–416.
- [9] Kiriushcheva N. and Kuzmin S., *Translational Invariance of the Einstein-Cartan Action in Any Dimension*, Gen. Rel. Grav. **42** (2010) 2613–2631 [arXiv:0907.1999 [gr-qc]].
- [10] Landau L. and Lifshitz E., *The Classical Theory of Fields*, Pergamon Press, Oxford 1971.
- [11] Maluf J., *The Teleparallel Equivalent of General Relativity*, Annalen der Physik **525** (2013) 339–357 [arXiv:1303.3897 [gr-qc]].
- [12] Møller C., *Conservation Laws and Absolute Parallelism in General Relativity*, Mat. Fys. Skr. Dan. Vid. Selsk. **1** No. 10 (1961) 1–50.
- [13] Montesinos M., Gonzalez D., Celada D and Diaz B., *Reformulation of the Symmetries of First-order General Relativity*, Class. Quant. Grav. **34** (2017) 205002 [arXiv:1704.04248 [gr-qc]].

-
- [14] Noether E. and Tavel M., *Invariant Variation Problems*, Transport theory and statistical physics **1** (1971) 183–207 (English translation) [arXiv:physics/0503066 [physics.hist-ph]].
- [15] Obukhov Y. and Rubilar G., *Covariance Properties and Regularization of Conserved Currents in Tetrad Gravity*, Phys. Rev. D **73** (2006) 124017 [arXiv:gr-qc/0605045].
- [16] Pereira G. and Obukhov Y., *Gauge Structure of Teleparallel Gravity*, Universe **5** (2019) 139 [arXiv:1906.06287 [gr-qc]].
- [17] Rodichev V., *Theory of Gravity in an Orthogonal Repère*, Nauka, Moscow 1974 (in Russian).
- [18] Samokhvalov S., *Group-theoretical Description of Gauge Fields*, Theor. Math. Phys. **76** (1988) 709–717.
- [19] Samokhvalov S., *Infinite Deformed Groups and Their Geometrical and Physical Applications*, Bogolyubov Kyiv conference "Modern problems of mathematics and theoretical physics", Kyiv 2004 [arXiv:math/0508239 [math.DG]].
- [20] Samokhvalov S., *Group-theoretic Description of Riemannian Spaces*, Ukr. Math. J. **55** (2003) 1238–1248 [arXiv:0704.2967 [math.DG]].
- [21] Samokhvalov S., *General Reference Frames and Definition of Energy in the Theory of Gravity*, Math. mod. **35** (2016) 19–23 (in Ukrainian).
- [22] Samokhvalov S., *Group-theoretical Basis of the Holographic Principle*, Math. mod. **23** (2010) 7–11 (in Ukrainian).
- [23] Samokhvalov S. and Balakireva E., *Group-theoretic Matching of the Length and the Equality Principles in Geometry*, Rus. Math. (Iz. VUZ) **59** (2015) 26–37.
- [24] Samokhvalov S. and Vanyashin V., *Group Theory Approach to Unification of Gravity with Internal Symmetry Gauge Interactions. I. Canonical Electrodynamics*, Class. Quant. Grav. **8** (1991) 2277–2282 [arXiv:1802.08958 [gr-qc]].
- [25] Samokhvalov S., *Group Theory Approach to Unification of Gravity with Internal Symmetry Gauge Interactions. II. Relativity of Charges and Masses*, Probl. Nucl. Phys. Cosm. Rays **35** (1991) 50–58 [arXiv:1802.09842 [gr-qc]].
- [26] Szabados L., *Quasi-local Energy-momentum and Angular Momentum in General Relativity: A Review Article*, Living Rev. Relativity **7** (2004) 1–140.
- [27] Unruh W., *Unimodular Theory of Canonical Quantum Gravity*, Phys. Rev. D **40** (1989) 1048–1052.

Chair of Applied Mathematics

Dniprovsk State Technical University, UKRAINE

E-mail address: serh.samokhval@gmail.com