

# Revisit of Y-junctions for strings with currents: transonic elastic case

I. Yu. Rybak<sup>1, 2, \*</sup>

<sup>1</sup>*Centro de Astrofísica da Universidade do Porto, Rua das Estrelas, 4150-762 Porto, Portugal*

<sup>2</sup>*Instituto de Astrofísica e Ciências do Espaço, CAUP, Rua das Estrelas, 4150-762 Porto, Portugal*

(Dated: 18th February 2022)

We studied the formation of Y-junctions for transonic elastic strings. In particular, using the general solution for these strings, which is described by left- and right-moving modes, we obtained the dynamics of kinks and Y-junctions. Considering the linearized ansatz for straight strings, we constructed the parameter region space for which the formation of Y-junctions due to strings collisions is allowed.

## I. INTRODUCTION

Cosmic strings are hypothetical objects that were originally described by Tom Kibble [1]. They appear as a prediction of numerous models of early universe [2]. To highlight some of them it is worthwhile to mention brane inflation [3–8], supersymmetric grand unified theory [9–13] and theories of high energy particle physics [14–18].

Some types of cosmic strings allow the existence of bound states, named as Y-junctions. They might appear due to collisions of distinct strings that form tri-linear vertices. Y-junctions are common for non-Abelian strings [19], for Abelian-Higgs strings of the I type [20], for  $U(1) \times U(1)$  models with specific value of parameters [21] and for cosmic strings from brane inflation (cosmic superstrings) [5]. Using approximation that cosmic strings are infinitely thin, are described by Nambu-Goto action, it was demonstrated that kinematic constraints must be satisfied in order for the Y-junctions to be produced [22–24]. The result of kinematic constraints was confirmed by numerical simulations in a framework of field theory [25, 26]. The analytic description of cosmic strings via Nambu-Goto action also sheds light on dynamics of Y-junctions. In particular, one can estimate the average growth/reduction of string lengths for multi-tension cosmic string networks. This phenomenon is crucial for understanding the evolution of cosmic (super)string networks [27, 28].

Due to nontrivial interactions of fields that form a string core, cosmic strings might become superconducting [29]. This situation naturally arises for supersymmetric [30–34] and some non-Abelian strings [35, 36]. To obtain an effective description of superconducting cosmic strings, models for infinitely thin strings were developed [29, 37–41]. It was also suggested that some macroscopic properties can be captured by such current carrying Nambu-Goto strings. In particular, the barytropic cosmic string model, which also comes out from dimensional reduction [42, 43], provides an accurate depiction of “wiggly” (noisy) cosmic strings [44–47].

This study revisits the problem of Y-junctions for transonic elastic strings. We re-examine the exact solution for these strings [44, 47], obtain left-/right-moving modes, and in line with [22] we describe evolution of kinks and Y-junctions. In addition, we obtain kinematic conditions under which the production of Y-junction is possible.

The problem of Y-junctions for Nambu-Goto current carrying strings was initially studied in [48]. The authors developed a covariant formalism to investigate under which conditions the production of Y-junctions is possible. The result of paper [48] claims that for *magnetic* (space-like current) and *electric* (time-like current) current carrying strings the formation of Y-junction is impossible, unless the newly formed string is described by a more general equation of state. Detailed comparison of our result with work [48] can be found in appendix V.

## II. SOLUTION IN MINKOWSKI SPACE FOR TRANSONIC ELASTIC STRINGS

In this section we revisit the exact solution for transonic elastic strings, originally obtained in [44, 47], with the method developed in [49]. This approach allows us to show that only elastic and chiral strings lead to wave-like equations of motion.

We start consideration from the action

$$S = -\mu_0 \int f(\kappa) \sqrt{-\gamma} d\sigma d\tau, \quad (1)$$

where  $\mu_0$  is a constant defined by the symmetry breaking scale,  $\{\sigma, \tau\}$  are coordinates on the string worldsheet (Latin indexes “ $a$ – $d$ ” run over 0, 1) with induced metric

$$\gamma_{ab} \equiv x_{,a}^\mu x_{,b}^\nu \eta_{\mu\nu} \quad \text{and} \quad (2a)$$

$$\kappa \equiv \varphi_{,a} \varphi_{,b} \gamma^{ab}, \quad (2b)$$

$$\gamma \equiv \frac{1}{2} \varepsilon^{ac} \varepsilon^{bd} \gamma_{ab} \gamma_{cd}, \quad (2c)$$

$\varepsilon^{ac}$  is the Levi-Civita symbol,  $\eta^{\mu\nu}$  is Minkowski metric (Greek indexes run over space-time coordinates  $x^\mu$  from 0 to 3),  $x_{,a}^\mu \equiv \frac{\partial x^\mu}{\partial \sigma^a}$  and  $\varphi$  is a scalar function on the string worldsheet. The function  $f(\kappa)$  will be defined below.

\* Ivan.Rybak@astro.up.pt

The stress-energy tensor for the action (1) can be written as

$$T^{\mu\nu} \equiv -2 \frac{\delta S}{\delta g_{\mu\nu}} = \frac{\mu_0}{\sqrt{-g}} \int \delta^{(4)}(y - x(\sigma)) \sqrt{-\gamma} (U u^\mu u^\nu - T v^\mu v^\nu) d\sigma d\tau, \quad (3)$$

where  $u^\mu u_\mu = 1$  and  $v^\mu v_\mu = -1$  are orthonormal time-like and spacelike vectors. Mass per unity length  $U$  and tension  $T$  in (3) are given by expressions

$$\begin{aligned} U &= f - 2\kappa f'_\kappa \Theta[-\kappa f'_\kappa], \\ T &= f - 2\kappa f'_\kappa \Theta[\kappa f'_\kappa], \end{aligned} \quad (4)$$

where  $\Theta[\dots]$  is a Heaviside function and  $f'_\kappa = \frac{\partial f}{\partial \kappa}$  (for more details about the stress-energy tensor see section 4 in [50] or alternatively section 2 in [51]).

We can introduce the speed of “wiggles”  $c_E$  (propagation of transverse perturbations) and “woggles”  $c_L$  (propagation of longitudinal perturbations) according to [39, 47]

$$c_E^2 = \frac{T}{U}, \quad c_L^2 = -\frac{dT}{dU}. \quad (5)$$

In this way, for the standard Nambu-Goto string both propagations have the speed of light,  $c_E = c_L = 1$ . It is anticipated to have supersonic strings ( $c_E > c_L$ ) for most of regimes of superconducting strings [38, 52]. Meanwhile, the transonic model

$$c_L = c_E \leq 1 \quad (6)$$

can be considered as an effective description of wiggly strings [46, 47] and some particular limits of superconducting strings (see sections 5.8, 5.9 in [50]).

Using (4), the explicit form of (5) can be written as

$$\begin{aligned} c_E^2 &= \frac{f - 2\kappa f'_\kappa \Theta[\kappa f'_\kappa]}{f - 2\kappa f'_\kappa \Theta[-\kappa f'_\kappa]}, \\ c_L^2 &= -\frac{f'_\kappa - 2(f'_\kappa + \kappa f''_{\kappa\kappa})\Theta[\kappa f'_\kappa]}{f'_\kappa - 2(f'_\kappa + \kappa f''_{\kappa\kappa})\Theta[-\kappa f'_\kappa]}. \end{aligned} \quad (7)$$

Substituting (7) into condition (6) for transonic strings, one can obtain the equation for  $f(\kappa)$

$$(f'_\kappa)^2 + f f''_{\kappa\kappa} = 0 \Rightarrow f = \sqrt{c_1 \kappa + c_2}, \quad (8)$$

where  $c_1$  and  $c_2$  are constants of integration.

One can write down the equation of state for transonic strings using the expressions (4) together with (8)

$$UT = f(f - 2\kappa f'_\kappa) = c_2 = m^2, \quad (9)$$

where  $m$  is a mass dimensional constant.

We can define  $c_1 = \pm m^2$  and absorb  $m^2$  into the definition of  $\mu_0$ . These manipulations allow us to establish the function  $f(\kappa)$  for transonic elastic strings in the following form, as also presented in [44, 47],

$$f(\kappa) = \sqrt{1 - \kappa}, \quad \kappa \in (-\infty, 1], \quad UT = 1. \quad (10)$$

It is known that the transonic model has the general wave-like solution [44]. Let's use the method from [49] to demonstrate that there are only two types of strings, whose equations of motion can be reduced to the wave equation: *chiral* (see [40, 53]) and *transonic elastic* strings. We start consideration by writing down the equations of motion for the action (1) in Minkowski space [54]

$$\partial_a [\mathcal{T}^{ab} x^\mu_{,b}] = 0, \quad (11a)$$

$$\partial_a [\sqrt{-\gamma} \gamma^{ab} f'_\kappa \varphi_{,b}] = 0, \quad (11b)$$

where

$$\begin{aligned} \mathcal{T}^{ab} &= \sqrt{-\gamma} (\gamma^{ab} f - 2f'_\kappa \gamma^{ac} \gamma^{bd} \varphi_{,c} \varphi_{,d}) = \\ &= \sqrt{-\gamma} (\gamma^{ab} f + \theta^{ab}) \end{aligned} \quad (12)$$

(notice the change of the sign in (12) due to misprint in equation (6) of [54]).

Parametrization invariance of the string worldsheet allows us to make the transformation

$$\mathcal{T}^{ab} \mapsto \eta^{ab}, \quad (13)$$

if their determinants are equal [49]

$$\det \mathcal{T}^{ab} = \det \eta^{ab} = -1. \quad (14)$$

Let's expand the determinant of  $\mathcal{T}^{ab}$

$$\det \mathcal{T}^{ab} = -f^2 - f \text{Tr} \theta_c^a - \det \theta_c^a. \quad (15)$$

It is easy to check that  $\det \theta_c^a = 0$ , hence, we are left only with

$$\begin{aligned} \det \mathcal{T}^{ab} &= -f^2 + 2f f'_\kappa \text{Tr} [\gamma^{ac} \gamma^{bd} \varphi_{,c} \varphi_{,d}] = \\ &= -f(f - 2f'_\kappa \kappa) = -UT. \end{aligned} \quad (16)$$

It is seen from (16) that the transformation (13) is possible due to parametrization invariance only if the function  $f(\kappa)$  is defined as for transonic elastic strings (10), or  $f(\kappa)$  is defined as for chiral strings, where the current is a null vector  $\kappa \rightarrow 0$  [54].

The relation (16) together with (10) guarantees that the equations of motion for the string worldsheet (11a) has the general wave-like solution. Choosing the gauge where the worldsheet coordinate  $\tau$  coincides with physical time  $t$ , one can write down the solution for (11a) in the form of left- and right-moving modes

$$x^0 = \tau, \quad \mathbf{x} = \frac{1}{2} (\mathbf{a}(\sigma_+) + \mathbf{b}(\sigma_-)), \quad (17)$$

where  $\sigma_+ = \tau + \sigma$  and  $\sigma_- = \tau - \sigma$ .

Up to this point, we demonstrated how to obtain the result of [44] in a different manner. Let's study the equation of motion for the function  $\varphi$  (11b). To do so, we plug  $f f'_\kappa = \frac{1}{2}$  in (11b)

$$\begin{aligned} \partial_a [\sqrt{-\gamma} \gamma^{ab} f^2 f'_\kappa \varphi_{,b}] &= [\text{using (13)}] = \\ &= \partial_a [f (\eta^{ab} + 2f'_\kappa \sqrt{-\gamma} \gamma^{ac} \gamma^{bd} \varphi_{,c} \varphi_{,d}) f'_\kappa \varphi_{,b}] = \\ &= \partial_a [f f'_\kappa (\eta^{ab} + 2\kappa \sqrt{-\gamma} \gamma^{ab} f'_\kappa) \varphi_{,b}]. \end{aligned} \quad (18)$$

Transferring the right-hand side term with  $\sqrt{-\gamma}\gamma^{ab}$  to the left-hand side in (18), one obtains

$$\begin{aligned} \partial_a [\sqrt{-\gamma}\gamma^{ab} f'_\kappa f (f - 2\kappa f'_\kappa) \varphi_{,b}] = \\ = \partial_a [\sqrt{-\gamma}\gamma^{ab} f'_\kappa \varphi_{,b}] = \partial_a [f f'_\kappa \eta^{ab} \varphi_{,b}] = 0. \end{aligned} \quad (19)$$

Taking out the constant  $f f'_\kappa$  from the differentiation operation in (19), we derive the following equation

$$\partial_a [\eta^{ab} \varphi_{,b}] = 0, \quad (20)$$

which general solution is given by

$$\varphi = \frac{1}{2} (F(\sigma_+) + G(\sigma_-)). \quad (21)$$

The normalization of  $|\mathbf{a}'|$  and  $|\mathbf{b}'|$  are connected with values of the current as

$$\mathbf{a}'^2(\sigma_+) = 1 - F'^2(\sigma_+), \quad \mathbf{b}'^2(\sigma_-) = 1 - G'^2(\sigma_-) \quad (22)$$

for right- and left-moving modes.

Alternative treatment of elastic strings, as a Kaluza-Klein projection of standard Nambu-Goto strings in a space-time of 5-dimensions, can be found in [55]. In this approach the relation (22) can be seen as a normalization for unity of 4-dimensional vectors of left- and right-moving modes.

Using relations (22), one can write down the current (2b) as

$$\kappa = \frac{2F'G'}{1 + F'G' - \mathbf{a}' \cdot \mathbf{b}'}, \quad (23)$$

which is shown in figure 1 for different values of  $F'$  and  $G'$ .

It is seen that if the left(or right)-moving mode of the current is independent of  $\sigma_-$  (or  $\sigma_+$ ), the expression (23) goes to zero

$$\kappa = 0, \quad \text{if: } F' = 0, \quad (\text{or } G' = 0). \quad (24)$$

The situation (24) reproduces the chiral string properties, where only left(or right)-moving mode is allowed [40, 49, 53, 54].

### III. JUNCTIONS FOR TRANSONIC ELASTIC STRINGS

To study Y-junctions for transonic elastic strings we start with the action for three connected current carrying strings [48]

$$\begin{aligned} S = - \sum_{i=1}^3 \mu_i \int f(\kappa_i) \sqrt{-\gamma_i} \Theta(s_i(\tau) - \sigma_i) d\sigma_i d\tau + \\ + \sum_{i=1}^3 \int f_{\mu i} (x_i^\mu(s_i(\tau), \tau) - X^\mu(\tau)) d\tau + \\ + \sum_{i=1}^3 \int g_i (\varphi_i(s_i(\tau), \tau) - \Phi(\tau)) d\tau, \end{aligned} \quad (25)$$




Figure 1. The current  $\kappa$  defined by (23) for different values of scalar moving modes  $G'$ ,  $F'$  and for angles  $0, \pi$  between vectors  $\mathbf{a}'$  and  $\mathbf{b}'$ .

where the function  $f(\kappa)$  is given by (10),  $\mu_i$  are constants defined by the symmetry breaking scale,  $f_{\mu i}$ ,  $g_i$  are Lagrange multipliers for strings and currents, time functions  $X^\mu(\tau)$  and  $\Phi(\tau)$  define values for  $x_i^\mu$  and  $\varphi_i$  at the point where strings are connected, the index  $i = 1, 2, 3$  denotes each of the three strings (the summation over index  $i$  is carried out only when it is written explicitly).

Varying the action (25) with respect to  $x_i^\mu$  and  $\varphi_i$ , we obtain the equations of motion (11a) and (11b) for each type of strings. Using (16) and (20) the boundary terms from equations of motion, which are proportional to  $\delta(s_i(t) - \sigma_i)$ , can be expressed as

$$\begin{aligned} \mu_i \eta^{ab} x_{i,a}^\mu \lambda_{bi} = f_i^\mu, \\ 2\mu_i f_{\mu i} f'_{\kappa i} \eta^{ab} \varphi_{,a} \lambda_{bi} = g_i, \end{aligned} \quad (26)$$

where  $\lambda_{ai} = \{\dot{s}_i, -1\}$ .

The variation of the action (25) with respect to  $\mathcal{X}_i^\mu$  and  $\Phi$  gives us

$$\begin{aligned} \sum_{i=1}^3 f_i^\mu = 0, \\ \sum_{i=1}^3 g_i = 0, \end{aligned} \quad (27)$$

which can be rewritten using solutions (17) and (21) to-

gether with expressions (26) in the following way

$$\begin{aligned} \sum_{i=1}^3 \mu_i [\mathbf{a}'_i(1 + \dot{s}_i) - \mathbf{b}'_i(1 - \dot{s}_i)] &= 0, \\ \sum_{i=1}^3 \mu_i [F'_i(1 + \dot{s}_i) - G'_i(1 - \dot{s}_i)] &= 0. \end{aligned} \quad (28)$$

Finally, variation of the action (25) with respect to  $f_i^\mu$  and  $g_i$  provides us the following relations

$$\begin{aligned} x_i^\mu(s_i(\tau), \tau) &= X^\mu(\tau), \\ \varphi_i(s_i(\tau), \tau) &= \Phi(\tau). \end{aligned} \quad (29)$$

Differentiating (29), using the exact solutions (17) and (21) we obtain

$$\begin{aligned} (1 + \dot{s}_i)\mathbf{a}'_i + (1 - \dot{s}_i)\mathbf{b}'_i &= 2\dot{\mathbf{X}}(t), \\ F'_i(1 + \dot{s}_i) + G'_i(1 - \dot{s}_i) &= 2\dot{\Phi}(t). \end{aligned} \quad (30)$$

Manipulating vectors  $\mathbf{a}'_i$ ,  $\mathbf{b}'_i$  and using (28) with (30), it is possible to obtain the following equations

$$\begin{aligned} \mathbf{a}'_k(1 + \dot{s}_k) &= \frac{2}{\mu} \sum_{i=1}^3 (1 - \dot{s}_i)\mu_i \mathbf{b}'_i - (1 - \dot{s}_k)\mathbf{b}'_k, \\ F'_k(1 + \dot{s}_k) &= \frac{2}{\mu} \sum_{i=1}^3 (1 - \dot{s}_i)\mu_i G'_i - (1 - \dot{s}_k)G'_k \end{aligned} \quad (31)$$

and

$$\dot{\mathbf{X}} = \frac{1}{\mu} \sum_{i=1}^3 (1 - \dot{s}_i)\mu_i \mathbf{b}'_i. \quad (32)$$

Zero component of the vector equation in (26) provides energy conservation relation, which is identical to the standard Nambu-Goto scenario [22]

$$\sum_{i=1}^3 \mu_i \dot{s}_i = 0. \quad (33)$$

The relation (33) does not provide an additional constraint, but is a consequence of equations of motion. Hence, the relation (33) can be used as a check of numerical calculations that are carried out below.

We parametrize the string worldsheets in such way that modes  $\mathbf{a}'_i(\sigma_+)$ ,  $F'_i(\sigma_+)$  move outwards the string connection, while  $\mathbf{b}'_i(\sigma_-)$  and  $G'_i(\sigma_-)$  move towards the string connection. Such choice means that  $\mathbf{b}'_i(\sigma_-)$  and  $G'_i(\sigma_-)$  are initial values that define  $\mathbf{a}'_i(\sigma_+)$  and  $F'_i(\sigma_+)$  by equations (31). The first three equations for vectors  $\mathbf{a}'_i(\sigma_+)$  in (31) can be squared and using the normalization conditions (22) we eliminate  $\mathbf{a}'_i(\sigma_+)$ . Hence, we have the system of six independent algebraic equations (31) and six variables that can be found: three variables  $\dot{s}_i$  and three variables  $F'_i(\sigma_+)$ .

It is illustrative to compare values of  $\dot{s}_i$  for strings with currents and without. For this purpose we fix angles between  $\mathbf{b}'_i(\sigma_-)$ , define string constants  $\mu_i$  and evaluate the system of equations (31) for different values of  $G'_i(\sigma_-)$ . An example of such dependence is shown in figure 2.




Figure 2. Dynamics of Y-junctions represented by  $\dot{s}_i$  of strings with  $\mu_1 = 1$  (blue),  $\mu_2 = 1.2$  (red),  $\mu_3 = 1.4$  (green) and oriented with angles  $2\pi/3$  between them. Dashed lines show the values of  $\dot{s}_i$  depending on  $G'_1$ , dash-dotted on  $G'_2$ , solid on  $G'_3$ . Black dashed lines demonstrate no changes of  $\dot{s}_i$  when all  $G'_i$  increase simultaneously.

The description above demonstrates that dynamics of Y-junctions for transonic elastic strings can be described within Nambu-Goto approximation.

### A. Kinks for elastic strings

Having considered the Y-junctions, we can treat the formation and evolution of kinks for elastic strings. To do so we need simply change the sum in previous equations for 2 strings instead of 3. Let us consider the situation when parameters for strings are  $\mu_1 = \mu_2$ . Hence, from equations (30) and (31) one can deduce the following relations for two possible situations that satisfy conditions for the kink

$$\dot{s}_1 = -1 = -\dot{s}_2, F'_2 = G'_1, \quad (34a)$$

$$\text{or } \dot{s}_1 = 1 = -\dot{s}_2, F'_1 = G'_2. \quad (34b)$$

Illustrative example of two strings intercommutation, is shown on figure 3. After collision two kinks propagating in opposite direction are formed on each of strings. The kink  $\dot{s}_A$  corresponds to situation (34a) with  $F'_{sA} = G'_1$ , while another kink  $\dot{s}_B$  to (34b) with  $F'_{sB} = G'_2$ .

The velocities of kinks follow from equation (32), that

is

$$\dot{\mathbf{X}}_A^2 = (1 - G_1'^2), \quad (35a)$$

$$\dot{\mathbf{X}}_B^2 = (1 - G_2'^2), \quad (35b)$$

which are different from the speed of light if the correspondent current component is non-zero.




Figure 3. Upper panel demonstrates two cosmic strings before the collision. Arrows with  $G'$  and  $F'$  define left-right-moving modes of currents on cosmic strings. Bottom panel represents the situation when collided strings intercommute, exchange their moving modes.

It should be noted that if an elastic string collides with a standard Nambu-Goto (or chiral with  $G_1' = 0$ ) string, the intermediate growing section, between kinks  $\dot{s}_A$  and  $\dot{s}_B$ , is described by the chiral model.

In the same manner kinks appear when colliding strings form Y-junction: from the discontinuity of corresponding modes. The formation of kinks for elastic strings qualitatively is identical to the standard Nambu-Goto model considered in [23], except of the fact that the speed of kinks propagation is not equal to the speed of light, but given by (35a) and (35b).

#### IV. COLLISIONS OF TRANSONIC ELASTIC STRINGS

It is always possible to chose small region, where collided strings can be considered straight. We are going to study kinematic conditions for straight strings to produce a Y-junction.

We decompose the straight string solution as a linear combination of "bare" and current carrying parts [54]

$$\mathbf{x}_i = \mathbf{y}_i + \mathbf{z}_i, \quad (36)$$

where the "bare" part is given by

$$\begin{aligned} \mathbf{y}_{1,2} &= \{-\gamma_v^{-1}\sigma \cos \alpha; \mp \gamma_v^{-1}\sigma \sin \alpha; \pm v\tau\}, \\ \mathbf{y}_3 &= \{\gamma_u^{-1}\sigma \cos \theta; \gamma_u^{-1}\sigma \sin \theta; u\tau\}, \end{aligned} \quad (37)$$

while the current carrying part is described by

$$\mathbf{z}_i = -g_i\sigma_-(\dot{\mathbf{y}}_i - \mathbf{y}'_i) - f_i\sigma_+(\dot{\mathbf{y}}_i + \mathbf{y}'_i), \quad (38)$$

with  $\gamma_v^{-1} = \sqrt{1 - v^2}$ .

Constants  $f_i$  and  $g_i$  in (38) represent the current contribution for left- and right-moving modes.

From (36)-(38) one can find that

$$\begin{aligned} \mathbf{a}'_i &= (1 - 2f_i)(\dot{\mathbf{y}}_i + \mathbf{y}'_i), \quad |\mathbf{a}'_i|^2 = (1 - 2f_i)^2, \\ \mathbf{b}'_i &= (1 - 2g_i)(\dot{\mathbf{y}}_i - \mathbf{y}'_i), \quad |\mathbf{b}'_i|^2 = (1 - 2g_i)^2. \end{aligned} \quad (39)$$

Comparing constants  $f_i$  and  $g_i$  in (39) with (22), we establish the relations

$$\begin{aligned} f_i &= \frac{1}{2} \left( 1 - \sqrt{1 - F_i'^2} \right), \\ g_i &= \frac{1}{2} \left( 1 - \sqrt{1 - G_i'^2} \right). \end{aligned} \quad (40)$$

Parameters  $v$  and  $\alpha$  define orthogonal velocity and orientation of non-current carrying string [23], but for elastic (and chiral [49]) strings it is not the case. It happens due to presence of longitudinal component of velocity in considered parametrization. Hence, we will treat  $v$  and  $\alpha$  as some constant parameters combination of which provide string velocity and orientation. Alteration of parameters interpretation does not affect the validity of applied method.

In order to find out for which values of  $v$  and  $\alpha$ , the third string can be produced (which means that  $\dot{s}_3 > 0$ ), we need to derive the orientation and velocity of a newly created string, i.e.  $\theta$  and  $u$  parameters. To obtain these variables, we follow the procedure of [23], i.e. we write down the expression for  $\dot{\mathbf{X}}$ , given in (32), by substituting  $\sigma \rightarrow s_3(\tau)$  in (37) and (38)

$$\dot{\mathbf{X}} = \{T_1(\tau)\gamma_u^{-1} \cos \theta; T_1(\tau)\gamma_u^{-1} \sin \theta; T_2(\tau)u\}, \quad (41)$$

where  $T_1(\tau) = \dot{s}_3(\tau) + g_3(1 - \dot{s}_3(\tau)) - f_3(1 + \dot{s}_3(\tau))$  and  $T_2(\tau) = 1 - f_3(1 + \dot{s}_3(\tau)) + g_3(1 - \dot{s}_3(\tau))$ .

Combining (41) with (32) one can obtain the vector equation, from which  $\theta$  and  $u$  are determined via  $\mathbf{b}'_i$ .

To summarize, we have nine equations: six equations from (31) and three equations from (32). Therefore, we can derive eight variables  $F'_i, \dot{s}_i, u, \theta$  defining another eight variables  $\mu_i, G'_i, v, \alpha$ . The vector equality (32) with (41) does not provide three independent equations, but only two, similarly as in [23]. Having all this information, we can numerically solve this system of algebraic equations. As a result, one can obtain the region of values  $v$  and  $\alpha$  for which colliding strings give rise to Y-junctions ( $\dot{s}_3 > 0$ ), see figure 4 for symmetric string collision and figure 5 for asymmetric string collision.




Figure 4. Symmetric case. Range of parameters  $v$  and  $\alpha$ , which allow for colliding strings with  $\mu_1 = \mu_2 = 1$  to produce the Y-junction ( $\dot{s}_3 > 0$  corresponds to areas below lines) with  $\mu_3 = 1.2$ . The solid blue line corresponds to  $G'_1 = G'_2 = 0.7$ , red line to  $G'_3 = 0.99$ , while all others  $G'_i$  are zeros. Dashed black line represents the case when all  $G'_i = 0$ .




Figure 5. Asymmetric case. Range of parameters  $v$  and  $\alpha$ , which allow for colliding strings with  $\mu_1 = 1$  and  $\mu_2 = 1.2$  to produce the Y-junction ( $\dot{s}_3 > 0$  corresponds to areas below lines) with  $\mu_3 = 1.4$ . The solid blue line corresponds to  $G'_1 = 0.99$ , red line to  $G'_2 = 0.99$ , green line to  $G'_3 = 0.99$  while all others  $G'_i$  are zeros. Dashed black line represents the case when all  $G'_i = 0$ , while black solid when  $G'_1 = G'_2 = 0.65$ ,  $G'_3 = 0$ .

Production of Y-junction also leads to creation of kinks

$$\mathbf{K}_{1,2} = (1 - 2g_{1,2}) \{ \gamma_v^{-1} \cos \alpha; \pm \gamma_v^{-1} \sin \alpha; \pm v \} \tau \quad (42)$$

that propagate along collided strings, similarly as it happens for standard non-current Nambu-Goto strings [23].

It is important to note that for all strings the constants  $\mu_i$  were fixed, and we assumed that there is a relation between tensions of strings  $T_i$ . This assumption allows us

to eliminate one degree of freedom and treat  $G'_3$  as known value (it might be done through relation (23)). Possible bound between tensions of current carrying strings needs further investigation for particular models and goes beyond the scope of this paper.

## V. CONCLUSIONS

We revisited the exact solution for elastic transonic strings in Minkowski space [44, 47] with the method developed in [49]. The exact solution allowed us to consider left- and right-moving modes, which made it possible to treat the dynamics of Y-junctions in a similar manner as it was done in [23].

The system of equations (31) allowed us to obtain the rate of string lengths change  $\dot{s}_i$ , see figure 2, requiring the definition of incoming components of the current  $G'_i$ . The values of incoming current components  $G'_i$  should be determined by strings properties. Thus, in the case of cosmic superstrings the values of  $G'_i$  might be defined similarly to saturated BPS state (see [5, 56] for details), given by

$$\mu_{p,q} = \mu_F \sqrt{(p - qC_0)^2 + q^2/g_s^2}.$$

For superconducting and wiggly cosmic strings with Y-junctions, the definitions of  $G'_i$  should arise from the values of tensions and mass per unit lengths (4). The exact definition of  $G'_i$  for particular type of strings needs further investigation and goes beyond the scope of this paper.

We studied the kink dynamics for elastic strings in section III A. We obtained values of  $\dot{s}_i$  that are essential for existence of kink-like discontinuity. We also demonstrated an example of elastic strings intercommutation and determined the velocities of these kinks.

In section IV we found kinematic constraints that should be satisfied to give rise to a Y-junction for elastic strings. In particular, we obtained a range of parameters  $v$ ,  $\alpha$  and  $G'_i$  of collided strings (36) for which  $\dot{s}_3 > 0$ . The symmetric case of elastic strings collision is shown in figure 4 and asymmetric case is shown in figure 5.

## ACKNOWLEDGMENTS

This work was supported by FCT - Fundao para a Ciencia e a Tecnologia through national funds (PTDC/FIS-PAR/31938/2017) and by FEDER - Fundo Europeu de Desenvolvimento Regional through COMPETE2020 - Programa Operacional Competitividade e Internacionalizao (POCI-01-0145-FEDER-031938). We also would like to thank Juliane F. Oliveira, Lara G. Sousa, Ricardo C. Costa, Carlos J.A.P. Martins and Anastasios Avgoustidis for fruitful discussions and help in organizing the manuscript.

## APPENDIX: COMPARISON WITH [48]

In section IV we considered the formation of Y-junctions for transonic model of straight strings determined by the action (1) with  $f(\kappa)$  defined in (10). It was shown that for collision of transonic elastic straight strings there are kinematic conditions that should be satisfied to form a Y-junction. In our study we did not face overdetermined system of equations, in contrast to result in [48]. The general approach of [48] claims that for *magnetic* and *electric* superconducting strings the formation of Y-junction is impossible. On the other hand, the result in section IV states that formation of Y-junction is possible for a particular type of current, described by transonic elastic model. This appendix is intended to provide detailed comparison of our result with result obtained in [48]. For the sake of clarity we denote equations related to work [48] as (...\*).

### A. Comparison of equations

Let us write down equations of motion for Y-junction used in work [48] (above equations (16\*)-(17\*) in [48])

$$\partial_a \left( \sqrt{-\gamma_i} T_i^{ab} x_{i,b}^\mu \Theta(\tilde{s}_i(\tau_i) - \tilde{\sigma}_i) \right) = f_i^\mu \delta(\tilde{s}_i(\tau_i) - \tilde{\sigma}_i), \quad (43a)$$

$$\partial_a \left( \sqrt{-\gamma_i} z_i^a \Theta(\tilde{s}_i(\tau_i) - \tilde{\sigma}_i) \right) = g_i \delta(\tilde{s}_i(\tau_i) - \tilde{\sigma}_i), \quad (43b)$$

where  $z_i^a = \sqrt{\kappa_{0i}} c_i^a$ ,  $c_i^a = -2f_{\kappa_i}' \gamma_i^{ab} \varphi_{i,b}$ ,  $\sqrt{-\gamma_i} T_i^{ab} = \mathcal{T}_i^{ab}$ ,  $\kappa_{0i}$  is a constant multiplier and all derivatives are taken with respect to conformal gauge parameters  $\tau_i$  and  $\tilde{\sigma}_i$ , which are different for each string (in contrast to the gauge of the present study, where  $\tau$  is the same for all strings) and provide the relations

$$\partial_{\tau_i} x_i^\mu \partial_{\tilde{\sigma}_i} x_{i,\mu} = 0, \quad (\partial_{\tau_i} x_i^\mu)^2 = -(\partial_{\tilde{\sigma}_i} x_i^\mu)^2.$$

Equations (43) are parametrization invariant, namely one can chose any  $\tilde{\sigma}_i$  and  $\tau_i$ . If one choses the gauge  $(\tau, \sigma_i)$  of this work and uses expressions (13), (19) for elastic strings, defined by function (10), the system of equations (43) is reduced to

$$\partial_a \left( \eta^{ab} x_{i,b}^\mu \Theta(s_i(\tau) - \sigma_i) \right) = f_i^\mu \delta(s_i - \sigma_i), \quad (44a)$$

$$\partial_a \left( \eta^{ab} \varphi_{i,b} \Theta(s_i(\tau) - \sigma_i) \right) = g_i \delta(s_i - \sigma_i). \quad (44b)$$

Substituting exact solutions (17), (21) in (44) and using condition (27) one can see that boundary terms of equations (44) are identical to equations (28).

In case of conformal gauge  $(\tilde{\sigma}_i, \tau_i)$ , boundary terms of equations (43) have the form of equations (16\*)-(17\*) of reference [48], given by

$$\sum_i \sqrt{-\gamma_i} (T_i^{0b} \dot{\tilde{s}}_i - T_i^{1b}) x_{i,b}^\mu = 0, \quad (45a)$$

$$\sum_i \sqrt{-\gamma_i} (z_i^0 \dot{\tilde{s}}_i - z_i^1) = 0, \quad (45b)$$

Gauge of this work	Conformal gauge in [48]
Eq.(27) =	= Eq.(15*)
Eq.(28) =	= Eq.(14*)
Eq.(29) =	= Eqs.(16*)-(17*)

Table I. Correspondence between equations of the manuscript and work [48]

where  $\dot{\tilde{s}}_i \equiv \frac{d\tilde{s}_i}{d\tau_i}$ .

We demonstrated that equations (16\*)-(17\*) for Y-junction in [48] coincide with equations (28) of the main text. There is a full agreement between equations of [48] and equations of the manuscript, see table I for correspondence. Equations (25\*), (28\*) and (29\*) of [48], should be identical to (45) and to (28), but just written in a preferred rest-frame. Hence, the amount of equations in our study and work [48] are the same.

### B. Straight string solution and number of unknown variables

To understand where the disagreement with [48] comes from, we also need to count the number of unknown variables, since the number of equations in our study and in work [48] are the same.

To start, let us write down the solution for straight strings, which satisfies equations of motion (11) and can be considered as a linear term of Taylor expansion close to the point of strings collision, i.e. for gauge of this work

$$\begin{aligned} x_i^\mu(\sigma_i, \tau) &= A_i^\mu \sigma_i + B_i^\mu \tau + \mathcal{O}(\sigma_i^2, \tau^2), \\ \varphi_i(\sigma_i, \tau) &= C_i \sigma_i + D_i \tau + \mathcal{O}(\sigma_i^2, \tau^2) \\ &\text{or for conformal gauge} \\ x_i^\mu(\tilde{\sigma}_i, \tau_i) &= \tilde{A}^\mu \tilde{\sigma}_i + \tilde{B}^\mu \tau_i + \mathcal{O}(\tilde{\sigma}_i^2, \tau_i^2), \\ \varphi_i(\tilde{\sigma}_i, \tau_i) &= \tilde{C}_i \tilde{\sigma}_i + \tilde{D}_i \tau_i + \mathcal{O}(\tilde{\sigma}_i^2, \tau_i^2). \end{aligned} \quad (46)$$

where  $A_i^\mu$ ,  $B_i^\mu$ ,  $\tilde{A}_i^\mu$ ,  $\tilde{B}_i^\mu$ ,  $C_i$ ,  $D_i$ ,  $\tilde{C}_i$  and  $\tilde{D}_i$  are constants with possibly different physical meaning. The form of solution (46) provides left- and right-moving modes, similarly to (17) and (21).

Mass per unit length  $U$  and tension  $T$ , given by (4), are dynamical parameters that are constructed from left- and right-moving modes of (46) for the corresponding gauge. When  $U$ ,  $T$  (or current  $\kappa$  with action) are fixed one can still chose different left- and right-moving modes. This fact is well seen from expression (23), where the same value of the current  $\kappa$  can be constructed by different values of  $F'$  and  $G'$ .

Expressing all parameters, such as mass per unit length  $U$ , tension  $T$  and current  $\kappa$  by left- and right-moving modes, we can count number of unknown variables. All derivatives of outgoing modes, can be determined






Figure 6. Schematic picture of left- and right-moving modes of elastic transonic strings for the gauge described in the main text, section II. Upper panel shows left- and right-moving modes before the collision of strings, lower panel shows modes after. Moving modes with question symbol should be determined by equations for Y-junction (31)-(32).

through conditions (29), introducing  $\dot{\Phi}$  and  $\dot{\mathbf{X}}$ . The last variable can be substituted by expression with parameters  $\alpha$ ,  $u$  and  $\dot{s}_3$  of newly created string. Hence, there are 4-equations from (45) and energy conservation condition (which is automatically satisfied), 2-independent equations come from first expression of (29), providing 6 independent equations for six variables:  $u$ ,  $\alpha$ ,  $\dot{s}_i$  and  $\dot{\Phi}$ . Thus, expressing all parameters as left- and right-moving modes one obtains the system of equations, which has the same amount of equations and unknown variables (as it is mentioned in the end of section IV one also needs to define incoming mode  $G_3$ ).

### C. The difference with work [48]

The process of strings collision for elastic transonic strings is illustrated in figure 6. The upper panel demonstrates left-, right-moving modes of strings before the

collision, the lower - after. Modes denoted by question symbol should be obtained from Y-junction equations at  $J$ . Two kinks, illustrated in figure 6 as  $K_1$  and  $K_2$ , are formed after strings collision and, according to section III A, incoming modes propagate without modification through kinks  $K_1$  and  $K_2$  for special choice of parametrization, i.e. modes  $G_1$ ,  $b_1$ ,  $G_2$  and  $b_2$  can pass directly to Y-junction. Kinks for elastic transonic straight strings (36) are described by (42). Schematic figure 6 of strings collision with particular choice of parametrization is valid for elastic transonic strings as well as for standard Nambu-Goto strings [22, 23], and chiral strings [54].

For conformal gauge choice, which was used in [48], incoming modes for  $J - K_1$  and  $J - K_2$ , in general, are not the same as modes before kinks and should be determined by equations for kink discontinuity. As a result, to obtain similar parameter region space, which allows Y-junction formation, one should solve the system of equations that includes equations for kinks  $K_1$ ,  $K_2$  and junction  $J$  simultaneously. This fact demonstrates that the gauge choice, which was used in this work, simplifies equations allowing to consider incoming modes for Y-junction the same as modes before  $K_1$  and  $K_2$  kinks.

To summarize the comparison, it was demonstrated in section V A that equations for Y-junctions of this study and work [48] are in agreement, one system of equations can be transformed to the other, see table I. In section V B it was shown that any type of straight strings can be represented by the form (46) and split for left- and right-moving modes. Defining string parameters, such as  $T$ ,  $U$  and current through left- and right-moving modes, one obtains the system of equations that has the same amount of unknown variables and equations for Y-junction. Hence, equations of work [48] written via left- and right-moving modes (which do not require separated consideration of *electric* and *magnetic* types of current) are reduced to (31), (32) providing the same range of parameters that allows formation of Y-junction. In study [48] equations are not written in terms of left- and right-moving modes, but in terms of tension  $T$ , mass per unit length  $U$  and *electric* (or *magnetic*) current. Each value of  $U$  and  $T$  (as well as current) can be represented by different left- and right-moving modes. It means that there are more unknown variables in equations written via left- and right-moving modes than in equations written via  $U$  and  $T$ . Fixing mass per unit length  $U$  and tension  $T$ , as it was done in [48], one also fixes corresponding outgoing and incoming modes obtaining overdetermined system of equations, i.e. less unknown variables than equations for Y-junction.

- 
- [1] T. W. B. Kibble, Journal of Physics A: Mathematical and General **9**, 1387 (1976).
  - [2] T. Kibble, COSLAB (2004), arXiv:astro-ph/0410073v2 [astro-ph].
  - [3] C. P. Burgess, M. Majumdar, D. Nolte, F. Quevedo,

- G. Rajesh, and R.-J. Zhang, Journal of High Energy Physics **2001**, 047 (2001), arXiv:hep-th/0105204v3 [hep-th].
- [4] G. Dvali, R. Kallosh, and A. Van Proeyen, JHEP **2004**, 035 (2004), arXiv:hep-th/0312005v3 [hep-th].



- [5] E. J. Copeland, R. C. Myers, and J. Polchinski, JHEP **2004**, 013 (2004), arXiv:hep-th/0312067v5 [hep-th].
- [6] S. Sarangi and S.-H. H. Tye, Phys.Lett.B **536**, 185 (2002), arXiv:hep-th/0204074v1 [hep-th].
- [7] H. Firouzjahi and S.-H. H. Tye, JCAP **0503**, 009 (2005), arXiv:hep-th/0501099v3 [hep-th].
- [8] N. T. Jones, H. Stoica, and S.-H. H. Tye, Phys.Lett. **B**, 6 (2003), arXiv:hep-th/0303269v1 [hep-th].
- [9] R. Jeannerot, J. Rocher, and M. Sakellariadou, Phys.Rev.D **68**, 103514 (2003), arXiv:hep-ph/0308134v1 [hep-ph].
- [10] Y. Cui, S. P. Martin, D. E. Morrissey, and J. D. Wells, Phys.Rev.D **77**, 043528 (2008), arXiv:0709.0950v2 [hep-ph].
- [11] R. Jeannerot and M. Postma, JHEP **0412**, 043 (2004), arXiv:hep-ph/0411260 [astro-ph.CO].
- [12] A. Achúcarro, A. Celi, M. Esole, J. Van den Bergh, and V. P. A., JHEP **0601**, 102 (2006), arXiv:hep-th/0511001v2 [hep-th].
- [13] M. Majumdar and A. C. Davis, JHEP **2002**, 056 (2002), arXiv:hep-th/0202148v3 [hep-th].
- [14] R. H. Brandenberger, The Universe **1**, 6 (2013), arXiv:1401.4619v1 [astro-ph.CO].
- [15] M. Kawasaki, K. Saikawa, and T. Sekiguchi, Phys. Rev. D **91**, 065014 (2015), arXiv:1412.0789v3 [hep-ph].
- [16] M. Gorghetto, E. Hardy, and G. Villadoro, JHEP **2018** (2018), 10.1007/JHEP07(2018)151, arXiv:1806.04677 [hep-ph].
- [17] L. Fleury and G. D. Moore, Journal of Cosmology and Astroparticle Physics **2016**, 004 (2016), arXiv:1509.00026v1 [hep-ph].
- [18] B. Gripaio and O. Randal-Williams, Physics Letters B **782**, 94 (2018), arXiv:1610.05623v2 [hep-th].
- [19] D. Spergel and U.-L. Pen, The Astrophysical Journal **491**, L67 (1997), arXiv:astro-ph/9611198 [astro-ph].
- [20] L. Jacobs and C. Rebbi, Phys. Rev. B **19**, 4486 (1979).
- [21] P. M. Saffin, Journal of High Energy Physics **2005**, 011 (2005), arXiv:hep-th/0506138 [hep-th].
- [22] E. J. Copeland, T. W. B. Kibble, and D. A. Steer, Phys. Rev. Lett. **97**, 021602 (2006), arXiv:hep-th/0601153v3 [hep-th].
- [23] E. J. Copeland, T. W. B. Kibble, and D. A. Steer, Phys. Rev. D **75**, 065024 (2007), arXiv:hep-th/0611243v2 [hep-th].
- [24] E. J. Copeland, H. Firouzjahi, T. W. B. Kibble, and D. A. Steer, Phys. Rev. D **77**, 063521 (2008), arXiv:0712.0808v1 [hep-th].
- [25] P. Salmi, A. Achúcarro, E. J. Copeland, T. W. B. Kibble, R. de Putter, and D. A. Steer, Phys. Rev. D **77**, 041701(R) (2008), arXiv:0712.1204v2 [hep-th].
- [26] N. Bevis and P. M. Saffin, Phys. Rev. D **78**, 023503 (2008), arXiv:0804.0200v2 [hep-th].
- [27] A. Avgoustidis and E. P. S. Shellard, Phys. Rev. D **78**, 103510 (2008), arXiv:0705.3395v3 [astro-ph.CO].
- [28] I. Y. Rybak, A. Avgoustidis, and C. J. A. P. Martins, Phys. Rev. D **99**, 063516 (2019), arXiv:1812.04584v2 [astro-ph.CO].
- [29] E. Witten, Nuclear Physics B **249**, 557 (1985).
- [30] S. C. Davis, P. Bintruy, and A.-C. Davis, Physics Letters B **611**, 39 (2005), arXiv:hep-th/0501200v1 [hep-th].
- [31] S. C. Davis, A. C. Davis, and M. Trodden, Phys.Lett. **B**, 257 (1997), arXiv:hep-ph/9702360v1 [hep-ph].
- [32] S. C. Davis, A.-C. Davis, and M. Trodden, Phys. Rev. D **57**, 5184 (1998), arXiv:hep-ph/9711313v1 [hep-ph].
- [33] E. Allys, JCAP **1604**, 009 (2016), arXiv:1505.07888v3 [gr-qc].
- [34] M. Sakellariadou, Nuclear Physics B - Proceedings Supplements **192-193**, 68 (2009), theory and Particle Physics: The LHC Perspective and Beyond, arXiv:0902.0569v2 [hep-th].
- [35] A. E. Everett, Phys. Rev. Lett. **61**, 1807 (1988).
- [36] M. Hindmarsh, K. Rummukainen, and D. J. Weir, Phys.Rev.Lett. **117**, 251601 (2016), arXiv:arXiv:1607.00764v2 [hep-th].
- [37] B. Carter, Physics Letters B **228**, 466 (1989).
- [38] P. Peter, Phys. Rev. D **45**, 1091 (1992).
- [39] B. Carter and P. Peter, Phys. Rev. D **52**, R1744 (1995), arXiv:hep-ph/9411425v1 [hep-ph].
- [40] B. Carter and P. Peter, Phys.Lett. **B**, 41 (1999), arXiv:hep-th/9905025v1 [hep-th].
- [41] B. Carter, Ann. Phys. **9**, 247 (2000), arXiv:hep-th/0002162v1 [hep-th].
- [42] N. Nielsen, Nuclear Physics B **167**, 249 (1980).
- [43] N. Nielsen and P. Olesen, Nuclear Physics B **291**, 829 (1987).
- [44] B. Carter, Phys.Rev. **D41**, 3869 (1990).
- [45] A. Vilenkin, Phys. Rev. **D41**, 3038 (1990).
- [46] X. Martin, Phys. Rev. Lett. **74**, 3102 (1995).
- [47] B. Carter, Phys.Rev.Lett. **74**, 3098 (1995), arXiv:hep-th/9411231v1 [hep-th].
- [48] D. A. Steer, M. Lilley, D. Yamauchi, and T. Hiramatsu, Phys. Rev. D **97**, 023507 (2018), arXiv:1710.07475v1 [astro-ph.CO].
- [49] J. J. Blanco-Pillado, K. D. Olum, and A. Vilenkin, Phys. Rev. D **63**, 103513 (2001), arXiv:astro-ph/0004410v3 [astro-ph].
- [50] B. Carter, 2nd Mexican School on Gravitation and Mathematical Physics (1997), arXiv:hep-th/9705172 [hep-th].
- [51] I. Y. Rybak, A. Avgoustidis, and C. J. A. P. Martins, Phys. Rev. D **96**, 103535 (2017), arXiv:1709.01839v2 [astro-ph.CO].
- [52] P. Peter, Phys. Rev. D **47**, 3169 (1993).
- [53] A. C. Davis, T. W. B. Kibble, M. Pickles, and D. A. Steer, Phys. Rev. D **62**, 083516 (2000), arXiv:astro-ph/0005514v1 [astro-ph].
- [54] I. Y. Rybak, A. Avgoustidis, and C. J. A. P. Martins, Phys. Rev. D **98**, 063519 (2018), arXiv:1809.04033v1 [astro-ph.CO].
- [55] B. Carter and D. A. Steer, Phys. Rev. D **69**, 125002 (2004), arXiv:hep-th/0307161 [hep-th].
- [56] K. Dasgupta and S. Mukhi, Physics Letters B **423**, 261 (1998), arXiv:hep-th/9711094v1 [hep-th].