

Unbalanced $(p, 2)$ -fractional problems with critical growth

Deepak Kumar* and K. Sreenadh†

Department of Mathematics,
Indian Institute of Technology Delhi,
Hauz Khaz, New Delhi-110016, India.

Abstract

We study the existence, multiplicity and regularity results of non-negative solutions of following doubly nonlocal problem:

$$(P_\lambda) \begin{cases} (-\Delta)^{s_1} u + \beta(-\Delta)_p^{s_2} u = \lambda a(x)|u|^{q-2}u + \left(\int_\Omega \frac{|u(y)|^r}{|x-y|^\mu} dy \right) |u|^{r-2}u & \text{in } \Omega, \\ u = 0 & \text{in } \mathbb{R}^n \setminus \Omega, \end{cases}$$

where $\Omega \subset \mathbb{R}^n$ is a bounded domain with C^2 boundary $\partial\Omega$, $0 < s_2 < s_1 < 1$, $n > 2s_1$, $1 < q < p < 2$, $1 < r \leq 2_\mu^*$ with $2_\mu^* = \frac{2n-\mu}{n-2s_1}$, $\lambda, \beta > 0$ and $a \in L^{\frac{d}{d-q}}(\Omega)$, for some $q < d < 2_{s_1}^* := \frac{2n}{n-2s_1}$, is a sign changing function. We prove that each nonnegative weak solution of (P_λ) is bounded. Furthermore, we obtain some existence and multiplicity results using Nehari manifold method.

Key words: Non-local operator, Fractional $(p, 2)$ Laplacian, Choquard equation, Nehari manifold.

2010 Mathematics Subject Classification: 35J35, 35J60, 35R11.

1 Introduction

In this article, we are concerned with the regularity, existence and multiplicity results for solutions to $(p, 2)$ -fractional Choquard equation

$$(P_\lambda) \begin{cases} (-\Delta)^{s_1} u + \beta(-\Delta)_p^{s_2} u = \lambda a(x)|u|^{q-2}u + \left(\int_\Omega \frac{|u(y)|^r}{|x-y|^\mu} dy \right) |u|^{r-2}u & \text{in } \Omega \\ u = 0 & \text{in } \mathbb{R}^n \setminus \Omega, \end{cases}$$

where Ω is a bounded domain in \mathbb{R}^n with C^2 boundary $\partial\Omega$, $0 < s_2 < s_1 < 1$, $n > 2s_1$, $1 < q < p < 2$, $1 < r \leq 2_\mu^*$ with $2_\mu^* = \frac{2n-\mu}{n-2s_1}$, $\lambda, \beta > 0$, $0 < \mu < n$ and $a \in L^{\frac{d}{d-q}}(\Omega)$, where

*email: deepak.kr0894@gmail.com

†e-mail: sreenadh@maths.iitd.ac.in

$q < d < 2_{s_1}^*$ and $2_{s_1}^* = \frac{2n}{n-2s_1}$, is a sign changing function such that $a^+(x) := \max\{a(x), 0\} \not\equiv 0$. $(-\Delta)_m^s$ is the Fractional m -Laplace operator, defined as follows

$$(-\Delta)_m^s u(x) = -2 \lim_{\epsilon \rightarrow 0} \int_{\mathbb{R}^n \setminus B_\epsilon(x)} \frac{|u(y) - u(x)|^{m-2} (u(x) - u(y))}{|x - y|^{n+ms}} dy,$$

for $m > 1$ and $s \in (0, 1)$. The problems involving these kind of operators have their applications in obstacle problems, conservation laws, phase transition, image processing, anomalous diffusion, American options in finance. For more details, we refer to [11, 14, 26, 35] and the references therein.

In the local case, that is, when $s_1 = s_2 = 1$, and $\beta = 0$, the following equation with Choquard type nonlinearity

$$-\Delta u + V(x)u = (|x|^{-\mu} * |u|^p) |u|^{p-2} u, \text{ in } \mathbb{R}^n \quad (1.1)$$

has been studied extensively. In case of $n = 3, p = 2$ and $\mu = 1$, S. Pekar [34] describes (1.1) in quantum mechanics of a polaron at rest. Under the same assumptions, P. Choquard, in 1976, used (1.1), in modeling of an electron trapped in its own hole, in a certain approximation to Hartree-Fock theory of one component plasma [23]. Buffoni et al. [10] proved existence of at least one non-trivial solution of (1.1) when $p = 2, n = 3$ and potential V is sign changing periodic function with the assumption that 0 lies in the gap of the spectrum of the operator $-\Delta + V$. In [2], Alves et al. obtained the existence of a non-trivial solution via penalization method of the problem

$$-\Delta u + V(x)u = (|x|^{-\mu} * F(u)) f(u), \text{ in } \mathbb{R}^n,$$

where $0 < \mu < n, n = 3, V$ is a continuous real valued function and F is the primitive of f . Gao and Yang [18] studied the following Brezis-Nirenberg type problem

$$-\Delta u = \lambda u + \left(\int_{\Omega} \frac{|u(y)|^{2_\mu^*}}{|x - y|^\mu} dy \right) |u(x)|^{2_\mu^* - 2} u(x) \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega,$$

where $\Omega \subset \mathbb{R}^n$ is a bounded domain, $n \geq 3$ and $0 < \mu < n$. They proved existence, multiplicity and non-existence results with respect to the parameter λ .

In the nonlocal case D' Avenia et al. [15] considered

$$(-\Delta)^s u + \omega u = (|x|^{\alpha-n} * |u|^p) |u|^{p-2} u, \text{ in } \mathbb{R}^n,$$

where $\omega > 0, p > 1$ and $s \in (0, 1)$. In this work authors proved regularity, existence, non-existence, symmetry as well as decay properties of solution. Mukherjee and Sreenadh [29] obtained Brezis-Nirenberg type results for problem involving fractional Laplacian with Choquard type nonlinearity having critical growth. In [36], Pucci et al. studied problem involving critical Choquard nonlinearity with fractional p -Laplacian. For related work on this type of problems, we refer to [22, 27, 28, 39, 41] and the references therein.

Partial differential equations involving operator $-\Delta_p - \Delta_q$, known as the (p, q) -Laplacian, arises from important applications such as biophysics, plasma physics, reaction-diffusion (see [8, 13, 17, 25, 40]). Due to this, a lot of work has been done in last decade on (p, q) -Laplacian problems. Among them, Papageorgiou and Rădulescu [30] considered the problem

$$-\Delta_p u - \Delta u = f(x, u) \text{ in } \Omega, u = 0 \text{ on } \partial\Omega,$$

where Ω is a bounded domain and f is a Carathéodory function. Among other results, authors proved existence of four non-trivial solutions for the case $1 < p < 2$. Aizicovici et al. [1] proved existence of constant sign and nodal solutions for the problem

$$-\Delta_p u - \mu \Delta u = f(x, u) \text{ in } \Omega, u = 0 \text{ on } \partial\Omega,$$

where Ω is a bounded domain, $p > 2$ and $\mu > 0$. For general p and q , Yin and Yang [42] considered

$$-\Delta_p u - \Delta_q u = |u|^{p^*-2}u + \theta V(x)|u|^{r-2}u + \lambda f(x, u) \text{ in } \Omega, u = 0 \text{ on } \partial\Omega,$$

where $1 < r < q < p < n$ and $f(x, u)$ is a subcritical perturbation and they proved multiplicity of solutions using Lusternik-Schnirelman theory. Marano et. al [24] studied the problem with Carathéodory function having critical growth. Using critical point theory with truncation arguments and comparison principle authors also proved bifurcation type result.

As far as problems involving fractional (p, q) -Laplacian is concerned, there is not much literature available. Bhakta and Mukherjee [7] considered the problem

$$(Q_{\theta, \lambda}) \left\{ (-\Delta)_p^{s_1} u + (-\Delta)_q^{s_2} u = |u|^{p_{s_1}^*-2}u + \theta V(x)|u|^{r-2}u + \lambda f(x, u) \text{ in } \Omega, u = 0 \text{ in } \mathbb{R}^n \setminus \Omega, \right.$$

where $0 < s_2 < s_1 < 1$, $1 < r < q < p < \frac{n}{s_1}$, and V and f are some appropriate functions. Here they proved $(Q_{\theta, \lambda})$ has infinitely many weak solutions for some range of λ and θ . Moreover, for $V(x) \equiv 1$, $\lambda = 0$, and assuming $r > q$ and certain other conditions on n and r , they proved the existence of $cat_{\Omega}(\Omega)$ many solutions of $(Q_{\theta, \lambda})$ using Lusternik-Schnirelmann category theory. Goel et al. [21] studied the following fractional (p, q) -Laplacian problem

$$(-\Delta)_p^{s_1} u + \beta (-\Delta)_q^{s_2} u = \lambda a(x)|u|^{\delta-2}u + b(x)|u|^{r-2}u \text{ in } \Omega, u = 0 \text{ in } \mathbb{R}^n \setminus \Omega, \quad (1.2)$$

where $\Omega \subset \mathbb{R}^n$ is a bounded domain, $1 < \delta \leq q \leq p < r \leq p_{s_1}^* = \frac{np}{n - ps_1}$, $0 < s_2 < s_1 < 1$, $n > ps_1$, $\lambda, \beta > 0$, and a and b are sign changing functions. Using Nehari manifold method authors proved existence of at least two non-negative and non-trivial solutions in the subcritical case for all $\beta > 0$ and for some range of λ . For the critical case under some restriction on δ , they obtained multiplicity results in some range of β and λ . Furthermore, they proved weak solutions of (1.2) are in the space $L^\infty(\Omega) \cap C_{loc}^{0, \alpha}(\Omega)$, for some $\alpha \in (0, 1)$, when $2 \leq q \leq p < r < p_{s_1}^*$.

Inspired from all these works we study regularity, existence and multiplicity results of $(p, 2)$ fractional Laplacian problem with critical Choquard type non-linearity. Using Moser iteration technique we prove each weak solution of problem (P_λ) is bounded, in the case $\mu < 4s_1$. In this regard, we prove the following theorem.

Theorem 1.1 *Suppose $\mu < 4s_1$ and the function $a(x)$ is bounded in Ω . Let u be a non-negative solution of problem (P_λ) , then $u \in L^\infty(\Omega)$.*

Regarding the existence and multiplicity results, using the method of minimization over some suitable subset of the Nehari manifold we obtain existence of at least two non-trivial non-negative solutions for all $\beta > 0$ and λ in some range for the subcritical case. In the critical case, we prove the existence of solutions by identifying the first critical level (as defined in Lemma 5.2), below which the Palais-Smale sequences contain a convergent subsequence. We first prove multiplicity results for all $\beta > 0$ and for some range of λ but with some restriction on q . For this we estimate the fractional p -Laplacian norm of family of minimizers of S , the best constant of the embedding of the space X into $L^{2^*_{s_1}}(\Omega)$ (see Lemma 5.4). Later, we remove this restriction on q and prove multiplicity result for small λ and β . We show the following existence and multiplicity theorems for problem (P_λ) .

Theorem 1.2 *Let $r < 2^*_\mu$. Then, there exists $\lambda_0 > 0$ such that problem (P_λ) has at least two non-negative solutions for all $\lambda \in (0, \lambda_0)$ and $\beta > 0$.*

Theorem 1.3 *Let $r = 2^*_\mu$ and the function $a(x)$ be continuous in Ω . Then, there exist constants $\Lambda, \Lambda_0 > 0$ such that*

- (i) (P_λ) admits at least one non-negative solution for all $\lambda \in (0, \Lambda)$ and $\beta > 0$,
- (ii) (P_λ) admits at least two non-negative solutions for all $\lambda \in (0, \Lambda_0)$, $\beta > 0$ and
 - (I) for all $q > 0$, provided $1 < p < n/(n - s_1)$,
 - (II) for all $q \in (1, \frac{n(2-p)}{n-2s_1}) \cup (\frac{4n}{4n-np-4s_1}, p)$, provided $n/(n - s_1) \leq p < 2$.

Next, we remove this restriction on q , to obtain the existence of second solution in the critical case for all $1 < q < p < 2$ but for some range of λ and β . In this regard we state our theorem as follows.

Theorem 1.4 *Let $r = 2^*_\mu$, $1 < q < p < 2$ and the function $a(x)$ be continuous in Ω . Then, there exist constants $\Lambda, \Lambda_{00}, \beta_{00} > 0$ such that (P_λ) has at least two solutions for all $\lambda \in (0, \Lambda_{00})$ and $\beta \in (0, \beta_{00})$.*

We point out that the study of non-autonomous functionals characterized by the fact that the energy density changes its ellipticity and growth properties according to the point has been continued by Mingione *et al.* [4, 5, 6], Rădulescu *et al.* [3, 12, 31, 32, 37, 43], etc.

Some of the abstract methods used in this paper can be found in the recent monograph by Papageorgiou, Rădulescu and Repovš [33].

The paper is organized as follows: In section 2, we provide variational setting and regularity result. In section 3, we define Nehari manifold associated to problem (P_λ) and give fibering map analysis and some preliminary results. In section 4, we prove the existence and multiplicity results in the subcritical case. In section 5, we have the existence and multiplicity of solutions in the critical case.

2 Variational setting and regularity result

Let Ω be any open subset of \mathbb{R}^n , consider the function space, which were introduced in [38] for $p_i = 2$ and in [20] for general p ,

$$X_{p_i, s_i} := \left\{ u \in L^{p_i}(\mathbb{R}^n) : u = 0 \text{ a.e. in } \mathbb{R}^n \setminus \Omega, \int_Q \frac{|u(x) - u(y)|^{p_i}}{|x - y|^{n+p_i s_i}} dx dy < \infty \right\},$$

where $p_1 = 2$, $p_2 = p$, $0 < s_i < 1$ and $Q = \mathbb{R}^{2n} \setminus (\Omega^c \times \Omega^c)$, which is a reflexive Banach space when endowed with the norm

$$\|u\|_{X_{p_i, s_i}} := \left(\int_Q \frac{|u(x) - u(y)|^{p_i}}{|x - y|^{n+p_i s_i}} dx dy \right)^{\frac{1}{p_i}}. \quad (2.1)$$

Notice that the integral in (2.1) can be extended to \mathbb{R}^{2n} as $u = 0$ a.e. on $\mathbb{R}^n \setminus \Omega$.

For simplicity, we denote $X_1 := X_{2, s_1}$ and $X_2 := X_{p, s_2}$ and corresponding norms by $\|\cdot\|_{X_1}$ and $\|\cdot\|_{X_2}$, respectively. From [38], we have the continuous embedding of X_1 into $L^m(\Omega)$ for $1 \leq m \leq 2_{s_1}^*$, therefore we define

$$S_m = \inf_{u \in X_1 \setminus \{0\}} \frac{\|u\|_{X_1}^2}{\|u\|_m^2}.$$

For the sake of convenience, we denote $S_{2_{s_1}^*} = S$.

Regarding the spaces X_1 and X_2 , we have the following relation.

Lemma 2.1 *Let $1 < p \leq 2$ and $0 < s_2 < s_1 < 1$, then there exists a constant $C = C(|\Omega|, n, p, s_1, s_2) > 0$ such that*

$$\|u\|_{X_2} \leq C \|u\|_{X_1}, \quad \forall u \in X_1.$$

Proof. Proof follows from [21, Lemma 2.1]. □

The nonlocal nonlinear Choquard term present in the right hand side of (P_λ) is well defined due to the following result.

Theorem 2.2 *(Hardy-Littlewood-Sobolev inequality) Let $t, r > 1$ and $0 < \mu < n$ with $1/t + \mu/n + 1/r = 2$, $f \in L^t(\mathbb{R}^n)$ and $h \in L^r(\mathbb{R}^n)$. There exists a sharp constant $C(t, r, \mu, n) > 0$ independent of f, h , such that*

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{f(x)h(y)}{|x - y|^\mu} dx dy \leq C(t, r, \mu, n) |f|_t |h|_r.$$

In general, let $f = h = |u|^r$, then by Hardy-Littlewood-Sobolev inequality, we get

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(x)|^r |u(y)|^r}{|x-y|^\mu} dx dy \leq C(n, \mu, r) \|u\|_{tr}^{2r},$$

if $|u|^r \in L^t(\mathbb{R}^n)$ for some $t > 1$ satisfying $\frac{2}{t} + \frac{\mu}{n} = 2$. Thus, for $u \in H^{s_1}(\mathbb{R}^n)$, by Sobolev embedding theorems, we must have

$$\frac{2n - \mu}{n} \leq r \leq \frac{2n - \mu}{n - 2s_1}.$$

The term $2_\mu^* := (2n - \mu)/(n - 2s_1)$ is known as the upper critical exponent in the sense of Hardy-Littlewood-Sobolev inequality. In particular, when $r = \frac{2n - \mu}{n - 2s_1}$, for $u \in X_1$, we have

$$\left(\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(x)|^{2_\mu^*} |u(y)|^{2_\mu^*}}{|x-y|^\mu} dx dy \right)^{1/2_\mu^*} \leq C(n, \mu)^{\frac{1}{2_\mu^*}} \|u\|_{2_{s_1}^*}^2,$$

where $C(n, \mu)$ is a suitable constant and $2_{s_1}^* = 2n/(n - 2s_1)$. Let us define

$$S_H := \inf_{u \in H^{s_1}(\mathbb{R}^n) \setminus \{0\}} \frac{\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(x) - u(y)|^2}{|x-y|^{n+2s_1}} dx dy}{\left(\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(x)|^{2_\mu^*} |u(y)|^{2_\mu^*}}{|x-y|^\mu} dx dy \right)^{\frac{1}{2_\mu^*}}},$$

which is achieved if and only if u is of the form

$$C \left(\frac{t}{t^2 + |x - x_0|^2} \right)^{(n-2s_1)/2}, \text{ for } x \in \mathbb{R}^n,$$

for some $x_0 \in \mathbb{R}^n$, $C > 0$ and $t > 0$ (see [29]). Moreover,

$$S_H = \frac{S}{(C(n, \mu))^{1/2_\mu^*}}. \quad (2.2)$$

Lemma 2.3 ([29, Lemma 2.2]) *Define*

$$S_H(\Omega) := \inf_{u \in X_1 \setminus \{0\}} \frac{\int_Q \frac{|u(x) - u(y)|^2}{|x-y|^{n+2s_1}} dx dy}{\left(\int_\Omega \int_\Omega \frac{|u(x)|^{2_\mu^*} |u(y)|^{2_\mu^*}}{|x-y|^\mu} dx dy \right)^{1/2_\mu^*}}.$$

Then $S_H = S_H(\Omega)$ and $S_H(\Omega)$ is never achieved except when $\Omega = \mathbb{R}^n$.

Lemma 2.4 ([29, Lemma 3.4]) *Let*

$$\|u\|_{NL} = \left(\int_\Omega \int_\Omega \frac{|u(x)|^{2_\mu^*} |u(y)|^{2_\mu^*}}{|x-y|^\mu} dx dy \right)^{1/2 \cdot 2_\mu^*},$$

then $\|\cdot\|_{NL}$ defines a norm on $Y := \{u : \Omega \rightarrow \mathbb{R} : u \text{ is measurable, } \|u\|_{NL} < \infty\}$.

Notations: For simplicity, for $p_1 = 2$ and $p_2 = p$, we will use the following notations

$$\begin{aligned}\mathcal{K}_i(u, v) &= \int_Q \frac{|u(x) - u(y)|^{p_i-2} (u(x) - u(y))(v(x) - v(y))}{|x - y|^{n+p_i s_i}} dx dy, \quad \text{for all } u, v \in X_i, \\ \mathcal{A}_r(u, v) &= \int_\Omega \int_\Omega \frac{|u(y)|^r |u(x)|^{r-2} u(x) v(x)}{|x - y|^\mu} dx dy \quad \text{for all } u, v \in X_1\end{aligned}$$

and $\hat{r} = \frac{2n}{2n-\mu} r$.

Definition 2.5 A function $u \in X_1$ is said to be a solution of problem (P_λ) , if for all $v \in X_1$

$$\mathcal{K}_1(u, v) + \beta \mathcal{K}_2(u, v) - \lambda \int_\Omega a(x) |u|^{q-2} uv \, dx - \mathcal{A}_r(u, v) = 0.$$

The Euler functional $\mathcal{I}_\lambda : X_1 \rightarrow \mathbb{R}$ associated to the problem (P_λ) is defined as

$$\mathcal{I}_\lambda(u) = \frac{1}{2} \|u\|_{X_1}^2 + \frac{\beta}{p} \|u\|_{X_2}^p - \frac{\lambda}{q} \int_\Omega a(x) |u|^q dx - \frac{1}{2r} \int_\Omega \int_\Omega \frac{|u(x)|^r |u(y)|^r}{|x - y|^\mu} dx dy.$$

Now we present a regularity result, namely the L^∞ bound, for weak solutions of problem (P_λ) .

Proof of Theorem 1.1: Let u be a nonnegative solution of (P_λ) . For $\vartheta > 1$ and $T > 0$, define

$$\phi(t) = \phi_{T, \vartheta}(t) = \begin{cases} 0, & \text{if } t \leq 0 \\ t^\vartheta, & \text{if } 0 < t < T \\ \vartheta T^{\vartheta-1} t - (\vartheta - 1) T^\vartheta, & \text{if } t \geq T. \end{cases}$$

Then,

$$\phi'(t) = \begin{cases} 0, & \text{if } t \leq 0 \\ \vartheta t^{\vartheta-1}, & \text{if } 0 < t < T \\ \vartheta T^{\vartheta-1}, & \text{if } t \geq T. \end{cases}$$

As $u(x) \geq 0$, it is easy to observe that

$$\phi(u) \geq 0, \quad \phi'(u) \geq 0 \quad \text{and} \quad u\phi'(u) \leq \vartheta\phi(u).$$

Since ϕ is Lipschitz, we have $\phi(u) \in X_1$ and

$$\|\phi(u)\|_{X_1} = \left(\int_Q \frac{|\phi(u(x)) - \phi(u(y))|^2}{|x - y|^{n+2s_1}} dx dy \right)^{1/2} \leq K \|u\|_{X_1}.$$

Moreover, we have

$$S \|\phi(u)\|_{2_{s_1}^*}^2 \leq \|\phi(u)\|_{X_1}^2 = \int_\Omega \phi(u) (-\Delta)^{s_1} \phi(u). \quad (2.3)$$

Since ϕ is convex, we observe that $\phi(u)\phi'(u) \in X_1$ and

$$\int_{\Omega} \phi(u)(-\Delta)^{s_1} \phi(u) \leq \int_{\Omega} \phi(u)\phi'(u)(-\Delta)^{s_1} u. \quad (2.4)$$

Define $g : \mathbb{R} \rightarrow \mathbb{R}$ as $g(t) = \phi(t)\phi'(t)$, then g is increasing, and set $G(t) = \int_0^t g'(s)^{\frac{1}{p}} ds$ for $t \in \mathbb{R}$. Using the inequality (see [9, Lemma A.2])

$$|a - b|^{p-2}(a - b)(g(a) - g(b)) \geq |G(a) - G(b)|^p, \text{ for } a, b \in \mathbb{R},$$

we obtain

$$\begin{aligned} & \int_Q \frac{|u(x) - u(y)|^{p-2}(u(x) - u(y))(\phi(u(x))\phi'(u(x)) - \phi(u(y))\phi'(u(y)))}{|x - y|^{n+ps_2}} dx dy \\ & \geq \int_Q \frac{|G(u(x)) - G(u(y))|^p}{|x - y|^{n+ps_2}} dx dy \geq 0. \end{aligned} \quad (2.5)$$

Noting the fact that $1 < q < 2$, that is, $0 < q - 1 < 1$, there exists a constant $C > 0$ such that $\lambda|a(x)|u|^{q-2}u| \leq C(1 + |u|)$. Thus, from (2.3), (2.4) and (2.5), we deduce that

$$\begin{aligned} S\|\phi(u)\|_{2_{s_1}^*}^2 & \leq \|\phi(u)\|_{X_1}^2 = \int_{\Omega} \phi(u)(-\Delta)^{s_1} \phi(u) \\ & \leq \int_{\Omega} \phi(u)\phi'(u)(-\Delta)^{s_1} u \\ & \quad + \int_Q \frac{|u(x) - u(y)|^{p-2}(u(x) - u(y))(\phi(u(x))\phi'(u(x)) - \phi(u(y))\phi'(u(y)))}{|x - y|^{n+ps_2}} dx dy \\ & = \lambda \int_{\Omega} a(x)|u|^{q-2}u\phi(u)\phi'(u) dx + \int_{\Omega} \int_{\Omega} \frac{|u(y)|^r |u(x)|^{r-2} u(x)\phi(u)\phi'(u)}{|x - y|^{\mu}} dx dy \\ & \leq C \int_{\Omega} (1 + u)\phi(u)\phi'(u) + \int_{\Omega} \int_{\Omega} \frac{|u(y)|^r |u(x)|^{r-2} u(x)\phi(u)\phi'(u)}{|x - y|^{\mu}} dx dy. \end{aligned}$$

Using the relation $u\phi'(u) \leq \vartheta\phi(u)$ and $\phi'(u) \leq \vartheta(1 + \phi(u))$, we get

$$S\|\phi(u)\|_{2_{s_1}^*}^2 \leq C\vartheta \int_{\Omega} (\phi(u)^2 + \phi(u)) + \vartheta \int_{\Omega} \int_{\Omega} \frac{|u(y)|^r |u(x)|^{r-2} \phi(u)^2}{|x - y|^{\mu}} dx dy. \quad (2.6)$$

Now, using Hölder inequality, we see that

$$\int_{\Omega} \phi(u) \leq \left(\int_{\Omega} \phi(u)^2 \right)^{1/2} |\Omega|^{1/2} \leq \frac{1}{2} \int_{\Omega} \phi(u)^2 + \frac{1}{2} |\Omega|. \quad (2.7)$$

Next, to estimate the second term in (2.6), we follow the approach similar to [39, Proof of Theorem 1(2)]. Using Theorem 2.2, we have

$$\int_{\Omega} \int_{\Omega} \frac{|u(y)|^r |u(x)|^{r-2} \phi(u)^2}{|x - y|^{\mu}} dx dy \leq C(n, \mu) \| |u|^r \|_{\frac{2n}{2n-\mu}} \| |u|^{r-2} \phi(u)^2 \|_{\frac{2n}{2n-\mu}}. \quad (2.8)$$

By the embedding results of X_1 , we get $u \in L^{2^*_{s_1}}$, so we can assume $\|u\|_{2^*_{s_1}} \leq C$. Employing Hölder inequality, we get $\| |u|^r \|_{\frac{2n}{2n-\mu}} \leq C$. Next, for $m > 0$, we deduce that

$$\begin{aligned} \left(\int_{\Omega} (|u|^{r-2} \phi(u)^2)^{\frac{2n}{2n-\mu}} \right)^{\frac{2n-\mu}{2n}} &= \left[\left(\int_{\{u < m\}} + \int_{\{u \geq m\}} \right) (|u|^{r-2} \phi(u)^2)^{\frac{2n}{2n-\mu}} \right]^{\frac{2n-\mu}{2n}} \\ &\leq m^{r-2} \left(\int_{\{u < m\}} (\phi(u))^{\frac{2 \cdot 2n}{2n-\mu}} \right)^{\frac{2n-\mu}{2n}} \\ &\quad + \left(\int_{\{u \geq m\}} (|u|^{r-2} \phi(u)^2)^{\frac{2n}{2n-\mu}} \right)^{\frac{2n-\mu}{2n}}. \end{aligned}$$

With the help of Hölder inequality, we obtain

$$\left(\int_{\{u \geq m\}} (|u|^{r-2} \phi(u)^2)^{\frac{2n}{2n-\mu}} \right)^{\frac{2n-\mu}{2n}} \leq \left(\int_{\Omega} \phi(u)^{2^*_{s_1}} \right)^{\frac{2}{2^*_{s_1}}} \left(\int_{\{u \geq m\}} |u|^{\frac{(r-2) \cdot 2^*_{s_1}}{2^*_{s_1} - 2}} \right)^{\frac{4s_1 - \mu}{2n}}.$$

By the fact $r \leq 2^*_{\mu}$ and $\|u\|_{2^*_{s_1}} \leq C$, we can find $m > 0$ such that

$$\left(\int_{\{u \geq m\}} |u|^{\frac{(r-2) \cdot 2^*_{s_1}}{2^*_{s_1} - 2}} \right)^{\frac{4s_1 - \mu}{2n}} \leq \frac{S}{2 \cdot \vartheta C}.$$

Therefore, collecting these informations in (2.8), we get

$$\begin{aligned} \int_{\Omega} \int_{\Omega} \frac{|u(y)|^r |u(x)|^{r-2} \phi(u)^2}{|x-y|^{\mu}} dx dy &\leq C \left(m^{r-2} \left(\int_{\Omega} (\phi(u))^{\frac{4n}{2n-\mu}} \right)^{\frac{2n-\mu}{2n}} \right. \\ &\quad \left. + \frac{S}{2 \cdot \vartheta C} \left(\int_{\Omega} \phi(u)^{2^*_{s_1}} \right)^{\frac{2}{2^*_{s_1}}} \right). \end{aligned} \quad (2.9)$$

Hence, using (2.7) and (2.9) in (2.6), we obtain

$$\|\phi(u)\|_{2^*_{s_1}}^2 \leq 2 \cdot \vartheta C \left(1 + \int_{\Omega} \phi(u)^2 + \left(\int_{\Omega} |\phi(u)|^{\frac{4n}{2n-\mu}} \right)^{\frac{2n-\mu}{2n}} \right).$$

Taking $T \rightarrow \infty$, we get

$$\left(\int_{\Omega} u^{2^*_{s_1} \vartheta} \right)^{\frac{2}{2^*_{s_1}}} \leq C \vartheta \left(1 + \int_{\Omega} u^{2\vartheta} + \left(\int_{\Omega} u^{2\vartheta \frac{2^*_{s_1}}{2^*_{\mu}}} \right)^{\frac{2^*_{\mu}}{2^*_{s_1}}} \right),$$

with the help of Hölder inequality, we deduce that

$$\left(\int_{\Omega} u^{2^*_{s_1} \vartheta} \right)^{\frac{2}{2^*_{s_1}}} \leq C \vartheta \left[1 + (1 + |\Omega|^{1 - \frac{2^*_{\mu}}{2^*_{s_1}}}) \left(\int_{\Omega} u^{2\vartheta \frac{2^*_{s_1}}{2^*_{\mu}}} \right)^{\frac{2^*_{\mu}}{2^*_{s_1}}} \right] \leq C \vartheta \left[1 + \left(\int_{\Omega} u^{2\vartheta \frac{2^*_{s_1}}{2^*_{\mu}}} \right)^{\frac{2^*_{\mu}}{2^*_{s_1}}} \right],$$

therefore,

$$\left(\int_{\Omega} u^{2_{s_1}^* \vartheta} \right)^{1/2_{s_1}^* \vartheta} \leq (C\vartheta)^{\frac{1}{2\vartheta}} \left(1 + \left(\int_{\Omega} u^{2\vartheta \frac{2_{s_1}^*}{2_{\mu}^*}} \right)^{2_{\mu}^*/2_{s_1}^*} \right)^{1/2\vartheta}. \quad (2.10)$$

Now we consider the following cases:

Suppose there exists a sequence $\vartheta_k \rightarrow \infty$ such that $\int_{\Omega} u^{2\vartheta_k \frac{2_{s_1}^*}{2_{\mu}^*}} \leq 1$, which implies

$$\left(1 + \left(\int_{\Omega} u^{2\vartheta_k \frac{2_{s_1}^*}{2_{\mu}^*}} \right)^{2_{\mu}^*/2_{s_1}^*} \right)^{1/2\vartheta_k} \leq 2^{\frac{1}{2\vartheta_k}},$$

thus, from (2.10), we get $u \in L^{\infty}(\Omega)$. Otherwise, there exists $\vartheta_0 > 0$ such that

$$\int_{\Omega} u^{2\vartheta \frac{2_{s_1}^*}{2_{\mu}^*}} > 1, \text{ for all } \vartheta \geq \vartheta_0.$$

Therefore, from (2.10), we obtain

$$\|u\|_{2_{s_1}^* \vartheta} \leq (C\vartheta)^{\frac{1}{2\vartheta}} \|u\|_{2\vartheta \frac{2_{s_1}^*}{2_{\mu}^*}} \text{ for all } \vartheta \geq \vartheta_0. \quad (2.11)$$

Define a sequence $\{\vartheta_k\}$ such that $2_{s_1}^* \vartheta_k = 2\vartheta_{k+1} \frac{2_{s_1}^*}{2_{\mu}^*}$, that is, $2_{\mu}^* \vartheta_k = 2\vartheta_{k+1}$ for $k \geq 0$. By means of (2.10), it is easy to notice that $\|u\|_{2_{s_1}^* \vartheta_0} < \infty$. Thus, from (2.11), we get

$$\|u\|_{2_{s_1}^* \vartheta_1} \leq (C\vartheta_1)^{\frac{1}{2\vartheta_1}} \|u\|_{2\vartheta_1 \frac{2_{s_1}^*}{2_{\mu}^*}} = C^{2\frac{1}{2\vartheta_1}} \vartheta_1^{\frac{1}{2\vartheta_1}} \|u\|_{2_{s_1}^* \vartheta_0} < \infty,$$

and

$$\|u\|_{2_{s_1}^* \vartheta_2} \leq C^{2\frac{1}{2\vartheta_1} + \frac{1}{2\vartheta_2}} \vartheta_1^{\frac{1}{2\vartheta_1}} \vartheta_2^{\frac{1}{2\vartheta_2}} \|u\|_{2_{s_1}^* \vartheta_0},$$

similarly we obtain

$$\|u\|_{2_{s_1}^* \vartheta_k} \leq C^{\sum_{m=1}^k \frac{1}{2\vartheta_m}} \prod_{m=1}^k (\vartheta_m^{\frac{1}{2\vartheta_m}}) \|u\|_{2_{s_1}^* \vartheta_0}. \quad (2.12)$$

By ratio test we can see that $\sum_{m \in \mathbb{N}} \frac{1}{2\vartheta_m} < \infty$. Let $z_k = \prod_{m=1}^k \vartheta_m^{\frac{1}{2\vartheta_m}}$. Again by using ratio test, we get $\ln z_k = \sum_{m=1}^k \frac{\ln \vartheta_m}{2\vartheta_m}$ is convergent. Hence there exists a positive constant A such that $C^{\sum_{m=1}^k \frac{1}{2\vartheta_m}} \prod_{m=1}^k (\vartheta_m^{\frac{1}{2\vartheta_m}}) \leq A$ for all $k \geq 0$. Therefore, (2.12) implies

$$\|u\|_{2_{s_1}^* \vartheta_k} \leq A \|u\|_{2_{s_1}^* \vartheta_0} < \infty. \quad (2.13)$$

We claim that $u \in L^{\infty}(\Omega)$. Suppose not, then there exists $\varepsilon > 0$ and $M \subset \Omega$ with $|M| > 0$ such that

$$u(x) \geq A \|u\|_{2_{s_1}^* \vartheta_0} + \varepsilon \quad \text{a.e. } x \in M.$$

Thus,

$$\|u\|_{2_{s_1}^* \vartheta_k} \geq \left(\int_M |u(x)|^{2_{s_1}^* \vartheta_k} \right)^{\frac{1}{2_{s_1}^* \vartheta_k}} \geq (A\|u\|_{2_{s_1}^* \vartheta_0} + \varepsilon) |M|^{\frac{1}{2_{s_1}^* \vartheta_k}},$$

this implies that

$$\liminf_{k \rightarrow \infty} \|u\|_{2_{s_1}^* \vartheta_k} \geq (A\|u\|_{2_{s_1}^* \vartheta_0} + \varepsilon),$$

which is a contradiction to (2.13). Hence, $u \in L^\infty(\Omega)$. \square

3 Nehari manifold and fibering map analysis

Due to the fact that $1 < r$, we see that $\mathcal{I}_\lambda(tu) \rightarrow -\infty$, as $t \rightarrow \infty$ for $u(\neq 0) \in X_1$. Hence, the functional \mathcal{I}_λ is not bounded below on X_1 . Therefore, it is necessary to restrict \mathcal{I}_λ to a proper subset of X_1 on which it is bounded below. For this reason, we consider the Nehari set \mathcal{M}_λ associated to (P_λ) , which is defined as

$$\mathcal{M}_\lambda = \{u \in X_1 \setminus \{0\} : \langle \mathcal{I}'_\lambda(u), u \rangle = 0\},$$

where $\langle \cdot, \cdot \rangle$ is the duality between X_1 and its dual space. Obviously, \mathcal{M}_λ contains all the solution of (P_λ) . To study the critical points of the functional \mathcal{I}_λ , we define the fibering maps associated to it. For $u \in X_1$, define $\varphi_u : \mathbb{R}^+ \rightarrow \mathbb{R}$ as $\varphi_u(t) = \mathcal{I}_\lambda(tu)$, that is

$$\begin{aligned} \varphi_u(t) &= \frac{t^2}{2} \|u\|_{X_1}^2 + \beta \frac{t^p}{p} \|u\|_{X_2}^p - \frac{\lambda t^q}{q} \int_\Omega a(x) |u|^q dx - \frac{t^{2r}}{2r} \int_\Omega \int_\Omega \frac{|u(x)|^r |u(y)|^r}{|x-y|^\mu} dx dy, \\ \varphi'_u(t) &= t \|u\|_{X_1}^2 + \beta t^{p-1} \|u\|_{X_2}^p - \lambda t^{q-1} \int_\Omega a(x) |u|^q dx - t^{2r-1} \int_\Omega \int_\Omega \frac{|u(x)|^r |u(y)|^r}{|x-y|^\mu} dx dy, \end{aligned} \quad (3.1)$$

$$\begin{aligned} \varphi''_u(t) &= \|u\|_{X_1}^2 + \beta(p-1)t^{p-2} \|u\|_{X_2}^p - (q-1)\lambda t^{q-2} \int_\Omega a(x) |u|^q dx \\ &\quad - (2r-1)t^{2r-2} \int_\Omega \int_\Omega \frac{|u(x)|^r |u(y)|^r}{|x-y|^\mu} dx dy. \end{aligned} \quad (3.2)$$

It is clear that $tu \in \mathcal{M}_\lambda$ if and only if $\varphi'_u(t) = 0$ and in particular, $u \in \mathcal{M}_\lambda$ if and only if $\varphi'_u(1) = 0$. Hence, it is natural to split \mathcal{M}_λ into three parts corresponding to local minima, local maxima and points of inflection, namely

$$\mathcal{M}_\lambda^0 := \{u \in \mathcal{M}_\lambda : \varphi''_u(1) = 0\}, \text{ and } \mathcal{M}_\lambda^\pm := \{u \in \mathcal{M}_\lambda : \varphi''_u(1) \gtrless 0\}.$$

Define $\sigma_\lambda := \inf\{\mathcal{I}_\lambda(u) \mid u \in \mathcal{M}_\lambda\}$ and $\sigma_\lambda^\pm := \inf\{\mathcal{I}_\lambda(u) \mid u \in \mathcal{M}_\lambda^\pm\}$.

Lemma 3.1 \mathcal{I}_λ is coercive and bounded below on \mathcal{M}_λ .

Proof. Proof follows using Hölder inequality and Sobolev embedding results. \square

Lemma 3.2 *There exists $\lambda_0 > 0$ such that for all $\lambda \in (0, \lambda_0)$, we have $\mathcal{M}_\lambda^0 = \emptyset$.*

Proof. We distinguish the following cases:

Case 1: $u \in \mathcal{M}_\lambda$ such that $\int_\Omega a(x)|u|^q dx = 0$.

From (3.1), we have

$$\|u\|_{X_1}^2 + \beta\|u\|_{X_2}^p - \int_\Omega \int_\Omega \frac{|u(x)|^r |u(y)|^r}{|x-y|^\mu} dx dy = 0.$$

Therefore,

$$\begin{aligned} \phi_u''(1) &= \|u\|_{X_1}^2 + \beta(p-1)\|u\|_{X_2}^p - (2r-1) \int_\Omega \int_\Omega \frac{|u(x)|^r |u(y)|^r}{|x-y|^\mu} dx dy \\ &= (2-2r)\|u\|_{X_1}^2 + \beta(p-2r)\|u\|_{X_2}^p < 0, \end{aligned}$$

which implies $u \notin \mathcal{M}_\lambda^0$.

Case 2: $u \in \mathcal{M}_\lambda$ such that $\int_\Omega a(x)|u|^q dx \neq 0$.

If $u \in \mathcal{M}_\lambda^0$, then from (3.1) and (3.2), we have

$$(2-q)\|u\|_{X_1}^2 + \beta(p-q)\|u\|_{X_2}^p = (2r-q) \int_\Omega \int_\Omega \frac{|u(x)|^r |u(y)|^r}{|x-y|^\mu} dx dy \quad \text{and} \quad (3.3)$$

$$(2r-2)\|u\|_{X_1}^2 + \beta(2r-p)\|u\|_{X_2}^p = \lambda(2r-q) \int_\Omega a(x)|u|^q dx. \quad (3.4)$$

Define $E_\lambda : \mathcal{M}_\lambda \rightarrow \mathbb{R}$ as

$$E_\lambda(u) = \frac{(2r-2)\|u\|_{X_1}^2 + \beta(2r-p)\|u\|_{X_2}^p}{(2r-q)} - \lambda \int_\Omega a(x)|u|^q dx,$$

then from (3.4), $E_\lambda(u) = 0$ for all $u \in \mathcal{M}_\lambda^0$. Additionally, using Hölder inequality, we have

$$E_\lambda(u) \geq \|u\|_{X_1}^q \left[\left(\frac{2r-2}{2r-q} \right) \|u\|_{X_1}^{(2-q)} - \lambda \|a\|_{\frac{d}{d-q}} S_d^{\frac{-q}{2}} \right], \quad (3.5)$$

Now from (3.3) and Theorem 2.2, we get

$$\|u\|_{X_1} \geq \left(\frac{(2-q)S_{\hat{r}}^r}{(2r-q)C(n, \mu)} \right)^{\frac{1}{2r-2}}.$$

Using this in (3.5), we obtain

$$E_\lambda(u) \geq \|u\|_{X_1}^q \left(\left(\frac{2r-2}{2r-q} \right) \left(\frac{(2-q)S_{\hat{r}}^r}{(2r-q)C(n, \mu)} \right)^{\frac{2-q}{2r-2}} - \lambda \|a\|_{\frac{d}{d-q}} S_d^{\frac{-q}{2}} \right).$$

Set

$$\lambda_0 := \left(\frac{(2r-2)S_d^{\frac{q}{2}}}{(2r-q)\|a\|_{\frac{d}{d-q}}} \right) \left(\frac{(2-q)S_{\hat{r}}^r}{(2r-q)C(n, \mu)} \right)^{\frac{2-q}{2r-2}} > 0, \quad (3.6)$$

then from (3.5), for $\lambda \in (0, \lambda_0)$ we get $E_\lambda(u) > 0$, for all $u \in \mathcal{M}_\lambda^0$, which is a contradiction. Therefore, $\mathcal{M}_\lambda^0 = \emptyset$ for all $\lambda \in (0, \lambda_0)$. \square

Now, define $\psi_u : \mathbb{R}^+ \rightarrow \mathbb{R}$ by

$$\psi_u(t) = t^{2-q} \|u\|_{X_1}^2 + \beta t^{p-q} \|u\|_{X_2}^p - t^{2r-q} \int_{\Omega} \int_{\Omega} \frac{|u(x)|^r |u(y)|^r}{|x-y|^\mu} dx dy,$$

then

$$\psi'_u(t) = (2-q)t^{1-q} \|u\|_{X_1}^2 + \beta(p-q)t^{p-q-1} \|u\|_{X_2}^p - (2r-q)t^{2r-q-1} \int_{\Omega} \int_{\Omega} \frac{|u(x)|^r |u(y)|^r}{|x-y|^\mu} dx dy.$$

Then trivially, $tu \in \mathcal{M}_\lambda$ if and only if t is a solution of $\psi_u(t) = \lambda \int_{\Omega} a(x)|u|^q dx$ and if $tu \in \mathcal{M}_\lambda$, then $\varphi''_{tu}(1) = t^{q-1} \psi'_u(t)$. Moreover, we see $\psi_u(t) \rightarrow -\infty$ as $t \rightarrow \infty$, $\psi_u(t) > 0$ for t small enough and $\psi'_u(t) < 0$ for t large enough. Now based on the sign of $\int_{\Omega} a(x)|u|^q dx$, we will study the fibering map φ_u .

Lemma 3.3 *Let $u(\neq 0) \in X_1$ and $\lambda \in (0, \lambda_0)$.*

(i) *If $\int_{\Omega} a(x)|u|^q dx > 0$, then there exist unique $t_1 < t_{\max} < t_2$ such that $t_1 u \in \mathcal{M}_\lambda^+$ and $t_2 u \in \mathcal{M}_\lambda^-$. Moreover, $\mathcal{I}_\lambda(t_1 u) = \min_{0 \leq t \leq t_2} \mathcal{I}_\lambda(tu)$ and $\mathcal{I}_\lambda(t_2 u) = \max_{t \geq t_{\max}} \mathcal{I}_\lambda(tu)$.*

(ii) *If $\int_{\Omega} a(x)|u|^q dx < 0$, then there exists unique $t_2 > 0$ such that $t_2 u \in \mathcal{M}_\lambda^-$.*

Proof. (i) Let $u \in X_1$ such that $\int_{\Omega} a(x)|u|^q dx > 0$. We claim that there exists unique $t_{\max} > 0$ such that $\psi'_u(t_{\max}) = 0$. To prove this, it is sufficient to show the existence of unique t_{\max} such that $F_u(t_{\max}) = \beta(p-q)\|u\|_{X_2}^p$, where $F_u(t) := (2r-q)t^{2r-p} \int_{\Omega} \int_{\Omega} \frac{|u(x)|^r |u(y)|^r}{|x-y|^\mu} dx dy - (2-q)t^{2-p} \|u\|_{X_1}^2$. By the fact that $p < 2 < r$, we see that $F_u(t) < 0$ for t small enough, $F_u(t) \rightarrow \infty$ as $t \rightarrow \infty$. Hence, there exists unique $\hat{t} > 0$ such that $F_u(\hat{t}) = 0$. Moreover, there exists unique $\tilde{t} > 0$ such that $F'_u(\tilde{t}) = 0$. Therefore, there exists unique $t_{\max} > \hat{t} > 0$ such that $F_u(t_{\max}) = \beta(p-q)\|u\|_{X_2}^p$. Using these, we conclude that ψ_u is increasing in $(0, t_{\max})$, decreasing in (t_{\max}, ∞) . As a consequence,

$$\begin{aligned} (2-q)t_{\max}^2 \|u\|_{X_1}^2 &\leq (2-q)t_{\max}^2 \|u\|_{X_1}^2 + \beta(p-q)t_{\max}^p \|u\|_{X_2}^p \\ &= t_{\max}^{2r} (2r-q) \int_{\Omega} \int_{\Omega} \frac{|u(x)|^r |u(y)|^r}{|x-y|^\mu} dx dy \leq (2r-q)t_{\max}^{2r} C(n, \mu) S_{\hat{r}}^{-r} \|u\|_{X_1}^{2r}. \end{aligned}$$

Define

$$T_0 := \frac{1}{\|u\|_{X_1}} \left(\frac{(2-q)S_{\hat{r}}^r}{(2r-q)C(n, \mu)} \right)^{1/(2r-2)} \leq t_{\max}$$

then,

$$\begin{aligned} \psi_u(t_{\max}) &\geq \psi_u(T_0) \geq T_0^{2-q} \|u\|_{X_1}^2 - T_0^{2r-q} C(n, \mu) S_{\hat{r}}^{-r} \|u\|_{X_1}^{2r} \\ &= \|u\|_{X_1}^q \left(\frac{2r-2}{2r-q} \right) \left(\frac{(2-q)S_{\hat{r}}^r}{(2r-q)C(n, \mu)} \right)^{\frac{2-q}{2r-2}} \geq 0. \end{aligned}$$

Since $\lambda < \lambda_0$, then there exist $t_1 < t_{\max}$ and $t_2 > t_{\max}$ such that $\psi_u(t_1) = \psi_u(t_2) = \lambda \int_{\Omega} a(x)|u|^q dx$. That is, $t_1 u, t_2 u \in \mathcal{M}_\lambda$. Also $\psi'_u(t_1) > 0$, $\psi'_u(t_2) < 0$ implies $t_1 u \in \mathcal{M}_\lambda^+$ and

$t_2u \in \mathcal{M}_\lambda^-$. Since $\varphi'_u(t) = t^q(\psi_u(t) - \lambda \int_\Omega a(x)|u|^q dx)$, $\varphi'_u(t) < 0$ for all $t \in [0, t_1)$ and $\varphi'_u(t) > 0$ for all $t \in (t_1, t_2)$. So, $\mathcal{I}_\lambda(t_1u) = \min_{0 \leq t \leq t_2} \mathcal{I}_\lambda(tu)$. Also, $\varphi'_u(t) > 0$ for all $t \in [t_1, t_2)$, $\varphi'_u(t_2) = 0$ and $\varphi'_u(t) < 0$ for all $t \in (t_2, \infty)$ implies $\mathcal{I}_\lambda(t_2u) = \max_{t \geq t_{\max}} \mathcal{I}_\lambda(tu)$.

(ii) Let $u \in X_1$ be such that $\int_\Omega a(x)|u|^q dx < 0$. From (i), we have ψ_u is increasing in $(0, t_{\max})$, decreasing in (t_{\max}, ∞) and $\psi'_u(t_{\max}) = 0$. Since $\lambda \int_\Omega a(x)|u|^q dx < 0$ and $\psi_u(t_{\max}) > 0$, there exists unique $t_1 > 0$ such that $\psi_u(t_1) = \lambda \int_\Omega a(x)|u|^q dx$ and $\psi'_u(t_1) < 0$, which implies $t_1u \in \mathcal{M}_\lambda^-$, that is t_1u is a local maximum. \square

Lemma 3.4 *Let λ_0 be defined as in (3.6), then the following holds.*

(i) *There exists a constant $C_1 > 0$ such that $\sigma_\lambda \leq \sigma_\lambda^+ \leq -\frac{(2-q)(r-1)}{2qr} C_1 < 0$.*

(ii) $\inf\{\|u\| : u \in \mathcal{M}_\lambda^-\} > 0$.

Proof. To prove (i), let $u_0 \in X_1$ such that $\int_\Omega a(x)|u_0|^q dx > 0$. It implies there exists $t_0 = t_0(u_0) > 0$ such that $t_0u_0 \in \mathcal{M}_\lambda^+$ that is, $\varphi''_{t_0u_0}(1) > 0$. Therefore, we have

$$\begin{aligned} \mathcal{I}_\lambda(t_0u_0) &\leq \left(\frac{2-q}{q}\right) \left(\frac{1}{2r} - \frac{1}{2}\right) \|t_0u_0\|_{X_1}^2 + \beta \left(\frac{p-q}{q}\right) \left(\frac{1}{2r} - \frac{1}{p}\right) \|t_0u_0\|_{X_2}^p \\ &\leq \left(\frac{2-q}{q}\right) \left(\frac{1}{2r} - \frac{1}{2}\right) \|t_0u_0\|_{X_1}^2 \leq -\frac{(2-q)(r-1)}{2qr} C_1, \end{aligned}$$

where $C_1 = \|t_0u_0\|_{X_1}^2$. This implies $\sigma_\lambda^+ \leq -\frac{(2-q)(r-1)}{2qr} C_1 < 0$.

To prove (ii), let $u \in \mathcal{M}_\lambda^-$, then $\varphi''_u(1) < 0$ which implies that

$$\begin{aligned} (2-q)\|u\|_{X_1}^2 &\leq (2-q)\|u\|_{X_1}^2 + \beta(p-q)\|u\|_{X_2}^p \\ &< (2r-q) \int_\Omega \int_\Omega \frac{|u(x)|^r |u(y)|^r}{|x-y|^\mu} dx dy \\ &\leq (2r-q)C(n, \mu)S_{\tilde{r}}^{-r} \|u\|_{X_1}^{2r}, \end{aligned}$$

that is,

$$\|u\|_{X_1} > \left(\frac{(2-q)S_{\tilde{r}}^r}{(2r-q)C(n, \mu)} \right)^{1/(2r-2)}.$$

Hence the result follows. \square

Lemma 3.5 *Let $\lambda \in (0, \lambda_0)$, and $z \in \mathcal{M}_\lambda$, then there exists $\epsilon > 0$ and a differentiable function $\xi : \mathcal{B}(0, \epsilon) \subseteq X_1 \rightarrow \mathbb{R}^+$ such that $\xi(0) = 1$, the function $\xi(w)(z-w) \in \mathcal{M}_\lambda$ and*

$$\langle \xi'(0), w \rangle = \frac{2\mathcal{K}_1(z, w) + p\beta\mathcal{K}_2(z, w) - q\lambda \int_\Omega a(x)|z|^{q-2}zwdx - 2r\mathcal{A}_r(z, w)}{(2-q)\|z\|_{X_1}^2 + (p-q)\|z\|_{X_2}^p - (2r-q) \int_\Omega \int_\Omega \frac{|z(x)|^r |z(y)|^r}{|x-y|^\mu} dx dy},$$

for all $w \in X_1$.

Proof. For $z \in \mathcal{M}_\lambda$, define a function $\mathcal{H}_z : \mathbb{R} \times X_1 \rightarrow \mathbb{R}$ given by

$$\begin{aligned} \mathcal{H}_z(t, w) &:= \langle \mathcal{I}'_\lambda(t(z-w)), (t(z-w)) \rangle \\ &= t^2 \|z-w\|_{X_1}^2 + \beta t^p \|z-w\|_{X_2}^p - t^q \lambda \int_\Omega a(x) |z-w|^q dx \\ &\quad - t^{2r} \int_\Omega \int_\Omega \frac{|(z-w)(x)|^r |(z-w)(y)|^r}{|x-y|^\mu} dx dy. \end{aligned}$$

Then $\mathcal{H}_z(1, 0) = \langle \mathcal{I}'_\lambda(z), z \rangle = 0$ and by Lemma 3.2, we have $\frac{\partial}{\partial t} \mathcal{H}_z(1, 0) \neq 0$. Therefore, by implicit function theorem result follows (for details see [21, Lemma 3.5]). \square

Proposition 3.6 *Let $\lambda \in (0, \lambda_0)$, then there exists a minimizing sequence $\{u_k\} \subset \mathcal{M}_\lambda$ such that*

$$\mathcal{I}_\lambda(u_k) = \sigma_\lambda + o_k(1) \text{ and } \mathcal{I}'_\lambda(u_k) = o_k(1).$$

Proof. Using Lemma 3.1 and Ekeland variational principle [16], there exists a minimizing sequence $\{u_k\} \subset \mathcal{N}_\lambda$ such that

$$\mathcal{I}_\lambda(u_k) < \sigma_\lambda + \frac{1}{k}, \text{ and} \tag{3.7}$$

$$\mathcal{I}_\lambda(u_k) < \mathcal{I}_\lambda(v) + \frac{1}{k} \|v - u_k\|_{X_1} \text{ for each } v \in \mathcal{M}_\lambda.$$

By taking k large, using equation (3.7) and Lemma 3.4, we deduce that

$$\begin{aligned} \mathcal{I}_\lambda(u_k) &= \left(\frac{1}{2} - \frac{1}{2r} \right) \|u_k\|_{X_1}^2 + \beta \left(\frac{1}{p} - \frac{1}{2r} \right) \|u_k\|_{X_2}^p - \lambda \left(\frac{1}{q} - \frac{1}{2r} \right) \int_\Omega a(x) |u_k|^q dx \\ &< \sigma_\lambda + \frac{1}{k} < \sigma_\lambda^+ < 0, \end{aligned}$$

which gives us $u_k \neq 0$ for k large enough. Now, using Hölder's inequality, we get

$$\left(\frac{2qr(-\sigma_\lambda^+) S_d^{\frac{q}{2}}}{\lambda(2r-q) \|a\|_{\frac{d}{d-q}}} \right)^{1/q} \leq \|u_k\|_{X_1} \leq \left(\frac{\lambda(2r-q) \|a\|_{\frac{d}{d-q}}}{q(r-1) S_d^{\frac{q}{2}}} \right)^{1/(2-q)}.$$

Then, proof of the result $\mathcal{I}'_\lambda(u_k) \rightarrow 0$, as $k \rightarrow \infty$ follows exactly on the same line of [21, Proposition 4.1]. \square

4 Subcritical case (when $r < 2_\mu^*$)

In this section, we prove the existence of at least two non-negative solutions of problem (P_λ) .

Lemma 4.1 *The functional \mathcal{I}_λ satisfies $(PS)_c$ condition for all $c \in \mathbb{R}$. That is, if $\{u_k\} \subset X_1$ satisfies*

$$\mathcal{I}_\lambda(u_k) = c + o_k(1) \text{ and } \mathcal{I}'_\lambda(u_k) = o_k(1) \text{ in } X'_1, \tag{4.1}$$

then $\{u_k\}$ has a convergent subsequence in X_1 .

Proof. Let $\{u_k\} \subset X_1$ satisfies (4.1). Then it is easy to verify that sequence $\{u_k\}$ is bounded in X_1 . So up to subsequence $u_k \rightharpoonup u_0$ weakly in X_1 , $u_k \rightarrow u_0$ strongly in $L^\nu(\Omega)$, $1 \leq \nu < 2_{s_1}^*$ and $u_k(x) \rightarrow u_0(x)$ a.e. in Ω . Since $\langle \mathcal{I}'_\lambda(u_k) - \mathcal{I}'_\lambda(u_0), (u_k - u_0) \rangle \rightarrow 0$ as $k \rightarrow \infty$, we have

$$\begin{aligned} o_k(1) &= \langle \mathcal{I}'_\lambda(u_k) - \mathcal{I}'_\lambda(u_0), u_k - u_0 \rangle \\ &= \mathcal{K}_1(u_k, u_k - u_0) - \mathcal{K}_1(u_0, u_k - u_0) + \beta(\mathcal{K}_2(u_k, u_k - u_0) - \mathcal{K}_2(u_0, u_k - u_0)) \\ &\quad - \lambda \int_\Omega a(x) (|u_k(x)|^{q-2} u_k(x) - |u_0(x)|^{q-2} u_0(x)) (u_k(x) - u_0(x)) dx \\ &\quad - (\mathcal{A}_r(u_k, u_k - u_0) - \mathcal{A}_r(u_0, u_k - u_0)). \end{aligned} \tag{4.2}$$

Using Hölder inequality and the fact that $d < 2_{s_1}^*$, we obtain

$$\int_\Omega a(x) |u_k|^{q-2} u_k (u_k - u_0) dx \leq \|a\|_{\frac{d}{d-q}} \|u_k\|_d^{q-1} \|u_k - u_0\|_d \rightarrow 0 \text{ as } k \rightarrow \infty.$$

Again, using Hölder inequality, Hardy-Littlewood-Sobolev inequality with the fact $\hat{r} = \frac{2nr}{2n-\mu} < 2_{s_1}^*$, we deduce that

$$\begin{aligned} \mathcal{A}_r(u_k, u_k - u_0) &= \int_\Omega \int_\Omega \frac{|u_k(x)|^r |u_k(y)|^{r-2} u_k(y) (u_k(y) - u_0(y))}{|x-y|^\mu} dx dy \\ &\leq C(n, \mu) \|u_k\|_{\hat{r}}^{2r-1} \|u_k - u_0\|_{\hat{r}} \rightarrow 0, \text{ as } k \rightarrow \infty. \end{aligned}$$

Now, we claim that the sequence $\{u_k\}$ has a convergent subsequence.

Using the definition of \mathcal{K}_1 , it is easy to see that $\mathcal{K}_1(u_k, u_k - u_0) - \mathcal{K}_1(u_0, u_k - u_0) = \|u_k - u_0\|_{X_1}^2$. Furthermore, since we know that

$$|a - b|^\eta \leq C_\eta ((|a|^{\eta-2} a - |b|^{\eta-2} b)(a - b))^{\frac{\eta}{2}} (|a|^\eta + |b|^\eta)^{\frac{2-\eta}{2}} \text{ for } a, b \in \mathbb{R}^n, 1 < \eta \leq 2,$$

where C_η is some positive constant depending on η . Set $a = u_k(x) - u_k(y)$, $b = u_0(x) - u_0(y)$ and then using Hölder inequality, we deduce that

$$\begin{aligned} \|u_k - u_0\|_{X_2}^p &\leq C(\mathcal{K}_2(u_k, u_k - u_0) - \mathcal{K}_2(u_0, u_k - u_0))^{\frac{p}{2}} \\ &\quad \left(\int_Q \frac{|u_k(x) - u_k(y)|^p + |u_0(x) - u_0(y)|^p}{|x-y|^{n+ps_2}} \right)^{\frac{2-p}{2}} \end{aligned}$$

and boundedness of $\{u_k\}$ in X_2 (follows from boundedness in X_1 and Lemma 2.1), implies

$$\|u_k - u_0\|_{X_2}^p \leq C(\mathcal{K}_2(u_k, u_k - u_0) - \mathcal{K}_2(u_0, u_k - u_0))^{\frac{p}{2}}.$$

Collecting all these informations in (4.2), we obtain

$$o_k(1) = \langle \mathcal{I}'_\lambda(u_k) - \mathcal{I}'_\lambda(u_0), u_k - u_0 \rangle \geq \frac{1}{C} (\|u_k - u_0\|_{X_1}^2 + \beta \|u_k - u_0\|_{X_2}^2).$$

Hence, it concludes proof of the claim. \square

Proof of Theorem 1.2 : Using Proposition 3.6 and Lemma 4.1, there exist minimizing sequences $\{u_k\} \in \mathcal{M}_\lambda^+$, $\{v_k\} \in \mathcal{M}_\lambda^-$ and u_0 and $v_0 \in X_1$ such that $u_k \rightarrow u_0$ and $v_k \rightarrow v_0$ strongly in X_1 for $\lambda \in (0, \lambda_0)$. Therefore, for $\lambda \in (0, \lambda_0)$, u_0, v_0 are weak solutions of problem (P_λ) . By means of Lemma 3.4, we conclude that $u_0, v_0 \neq 0$, hence $u_0 \in \mathcal{M}_\lambda^+$ and $v_0 \in \mathcal{M}_\lambda^-$. Moreover, $\mathcal{I}_\lambda(u_0) = \sigma_\lambda^+$ and $\mathcal{I}_\lambda(v_0) = \sigma_\lambda^-$. Since $\mathcal{M}_\lambda^+ \cap \mathcal{M}_\lambda^- = \emptyset$, therefore u_0 and v_0 are distinct solutions.

Now we prove non-negativity of u_0 . If $u_0 \geq 0$, then we have a non negative solution of problem (P_λ) , which is also a minimizer for \mathcal{I}_λ in \mathcal{M}_λ^+ , otherwise we have $|u_0| \neq 0$, hence by fibering map analysis we get unique $t_1 > 0$ such that $t_1 u_0 \in \mathcal{M}_\lambda^+$. We note that $\psi_{|u_0|}(1) \leq \psi_{u_0}(1) = \lambda \int_\Omega a(x)|u_0|^q = \psi_{|u_0|}(t_1) \leq \psi_{u_0}(t_1)$ and $0 < \psi'_{u_0}(1)$, because of the fact $u_0 \in \mathcal{M}_\lambda^+$, which implies $t_1 \geq 1$. Thus,

$$\sigma_\lambda^+ \leq \varphi_{|u_0|}(t_1) \leq \varphi_{|u_0|}(1) \leq \varphi_{u_0}(1) = \sigma_\lambda^+.$$

Hence, $\mathcal{I}_\lambda(t_1|u_0) = \varphi_{|u_0|}(t_1) = \sigma_\lambda^+$ and $t_1|u_0| \in \mathcal{M}_\lambda^+$ that is, $t_1|u_0|$ is a nonnegative solution of problem (P_λ) in \mathcal{M}_λ^+ . \square

5 Critical case (when $r = 2_\mu^*$)

In this section we assume the function $a(x)$ is continuous in Ω and $a^+(x) = \max\{a(x), 0\} \neq 0$. Then without loss of generality we may assume there exists $\delta_1 > 0$ such that $m_a := \inf_{x \in B_{\delta_1}} a(x) > 0$.

Theorem 5.1 *Let $\{u_k\} \subset \mathcal{M}_\lambda$ be a $(PS)_c$ sequence for \mathcal{I}_λ such that $u_k \rightharpoonup u$ weakly in X_1 , then $\mathcal{I}'_\lambda(u) = 0$. Moreover, there exists a positive constant $C_0 = C_0(q, s_1, n, S, |\Omega|)$ such that $\mathcal{I}'_\lambda(u) \geq -C_0 \lambda^{\frac{2}{2-q}}$, where*

$$C_0 = \left(\frac{(2 \cdot 2_\mu^* - q)(2 - q)}{2 \cdot 2 \cdot 2_\mu^* q} \right) \left(\frac{2 \cdot 2_\mu^* - q}{2 \cdot 2_\mu^* - 2} \right)^{\frac{q}{2-q}} S^{\frac{-q}{2-q}} \|a\|_{\infty}^{\frac{2}{2-q}} |\Omega|^{\frac{2(2_{s_1}^* - q)}{2_{s_1}^*(2-q)}}. \quad (5.1)$$

Proof. Since $u_k \rightharpoonup u$ in X_1 , it implies $\{u_k\}$ is a bounded sequence in X_1 , and up to subsequence, $u_k \rightarrow u$ in $L^\nu(\Omega)$, $1 \leq \nu < 2_{s_1}^*$ and $u_k \rightarrow u$ a.e. in Ω . Now from the proof of [21, Theorem 4.3] it follows that

$$\mathcal{K}_i(u_k, v) \rightarrow \mathcal{K}_i(u, v) \text{ for } i = 1, 2, \text{ and } \int_\Omega a(x)(|u_k(x)|^{q-2}u_k(x) - |u(x)|^{q-2}u(x))v(x) dx \rightarrow 0$$

as $k \rightarrow \infty$, for all $v \in X_1$. From the continuous embedding of X_1 into $L^{2_{s_1}^*}$, we get $u_k \rightharpoonup u$ weakly in $L^{2_{s_1}^*}$, as $k \rightarrow \infty$. Therefore, $|u_k|^{2_\mu^*} \rightharpoonup |u|^{2_\mu^*}$ in $L^{2_{s_1}^*/2_\mu^*}(\Omega)$ and we know that Riesz potential defines a continuous linear map from $L^{2_{s_1}^*/2_\mu^*}(\Omega)$ to $L^{\frac{2n}{\mu}}(\Omega)$, thus we have

$$|x|^{-\mu} * |u_k|^{2_\mu^*} \rightharpoonup |x|^{-\mu} * |u|^{2_\mu^*} \text{ in } L^{\frac{2n}{\mu}}(\Omega).$$

Moreover, we have $|u_k|^{2^*_\mu-2}u_k \rightharpoonup |u|^{2^*_\mu-2}u$ in $L^{2^*_{s_1}/(2^*_\mu-1)}(\Omega)$. Combining all these facts, we obtain

$$\int_{\Omega} \int_{\Omega} \frac{|u_k(x)|^{2^*_\mu} |u_k(y)|^{2^*_\mu-2} u_k(y) \phi(y)}{|x-y|^\mu} dx dy \rightarrow \int_{\Omega} \int_{\Omega} \frac{|u(x)|^{2^*_\mu} |u(y)|^{2^*_\mu-2} u(y) \phi(y)}{|x-y|^\mu} dx dy$$

Therefore, we have

$$\begin{aligned} \langle \mathcal{I}'_\lambda(u_k) - \mathcal{I}'_\lambda(u), \phi \rangle &= \mathcal{K}_1(u_k, \phi) - \mathcal{K}_1(u, \phi) + \beta (\mathcal{K}_2(u_k, \phi) - \mathcal{K}_2(u, \phi)) \\ &\quad - \int_{\Omega} a(x) (|u_k(x)|^{q-2} u_k(x) - |u(x)|^{q-2} u(x)) \phi(x) dx \\ &\quad - (\mathcal{A}_{2^*_\mu}(u_k, \phi) - \mathcal{A}_{2^*_\mu}(u, \phi)) \rightarrow 0 \quad \text{for } \phi \in X_1. \end{aligned}$$

This implies $\mathcal{I}'_\lambda(u) = 0$. In particular, $\langle \mathcal{I}'_\lambda(u), u \rangle = 0$, that is

$$\begin{aligned} \mathcal{I}_\lambda(u) &= \left(\frac{1}{2} - \frac{1}{2 \cdot 2^*_\mu} \right) \|u\|_{X_1}^2 + \beta \left(\frac{1}{p} - \frac{1}{2 \cdot 2^*_\mu} \right) \|u\|_{X_2}^p - \lambda \left(\frac{1}{q} - \frac{1}{2 \cdot 2^*_\mu} \right) \int_{\Omega} a(x) |u|^q dx \\ &\geq \left(\frac{1}{2} - \frac{1}{2 \cdot 2^*_\mu} \right) \|u\|_{X_1}^2 - \lambda \left(\frac{1}{q} - \frac{1}{2 \cdot 2^*_\mu} \right) \int_{\Omega} a(x) |u|^q dx. \end{aligned} \quad (5.2)$$

By Hölder inequality, Sobolev embeddings and Young inequality, we obtain

$$\begin{aligned} \lambda \int_{\Omega} a(x) |u|^q dx &\leq \lambda \|a\|_{\infty} S^{-\frac{q}{2}} |\Omega|^{\frac{2^*_{s_1}-q}{2^*_{s_1}}} \|u\|_{X_1}^q \\ &= \left(\frac{2}{q} \left(\frac{1}{2} - \frac{1}{2 \cdot 2^*_\mu} \right) \left(\frac{1}{q} - \frac{1}{2 \cdot 2^*_\mu} \right)^{-1} \right)^{\frac{q}{2}} \|u\|_{X_1}^q \\ &\quad \lambda \left(\frac{2}{q} \left(\frac{1}{2} - \frac{1}{2 \cdot 2^*_\mu} \right) \left(\frac{1}{q} - \frac{1}{2 \cdot 2^*_\mu} \right)^{-1} \right)^{-\frac{q}{2}} \|a\|_{\infty} |\Omega|^{\frac{2^*_{s_1}-q}{2^*_{s_1}}} S^{-\frac{q}{2}} \quad (5.3) \\ &\leq \left(\frac{1}{2} - \frac{1}{2 \cdot 2^*_\mu} \right) \left(\frac{1}{q} - \frac{1}{2 \cdot 2^*_\mu} \right)^{-1} \|u\|_{X_1}^2 \\ &\quad + \left(\frac{2-q}{2} \right) \left(\frac{2 \cdot 2^*_\mu - q}{2 \cdot 2^*_\mu - 2} \right)^{\frac{q}{2-q}} S^{-\frac{q}{2-q}} \|a\|_{\infty}^{\frac{2}{2-q}} |\Omega|^{\frac{2(2^*_{s_1}-q)}{(2-q)2^*_{s_1}}} \lambda^{\frac{2}{2-q}}, \end{aligned}$$

Therefore, result follows from equations (5.2) and (5.3) with

$$C_0 = \left(\frac{1}{q} - \frac{1}{2 \cdot 2^*_\mu} \right) \left(\frac{2-q}{2} \right) \left(\frac{2 \cdot 2^*_\mu - q}{2 \cdot 2^*_\mu - 2} \right)^{\frac{q}{2-q}} S^{-\frac{q}{2-q}} \|a\|_{\infty}^{\frac{2}{2-q}} |\Omega|^{\frac{2(2^*_{s_1}-q)}{(2-q)2^*_{s_1}}}.$$

□

Lemma 5.2 (Palais-Smale range). \mathcal{I}_λ satisfies the $(PS)_c$ condition with $c \in (-\infty, c_\infty)$, where

$$c_\infty := \left(\frac{n-\mu+2s_1}{2(2n-\mu)} \right) \frac{S^{\frac{2n-\mu}{n-\mu+2s_1}}}{(C(n,\mu))^{\frac{n-2s_1}{n-\mu+2s_1}}} - C_0 \lambda^{\frac{2}{2-q}}$$

and C_0 is the positive constant defined in (5.1).

Proof. Let $\{u_k\}$ be a $(PS)_c$ sequence of \mathcal{I}_λ in X_1 . Then we have

$$\frac{1}{2}\|u_k\|_{X_1}^2 + \frac{\beta}{p}\|u_k\|_{X_2}^p - \frac{\lambda}{q} \int_{\Omega} a(x)|u_k|^q dx - \frac{1}{2.2_\mu^*}\|u_k\|_{NL}^{2.2_\mu^*} = c + o_k(1) \quad (5.4)$$

and

$$\|u_k\|_{X_1}^2 + \beta\|u_k\|_{X_2}^p - \lambda \int_{\Omega} a(x)|u_k|^q dx - \|u_k - u\|_{NL}^{2.2_\mu^*} = o_k(1). \quad (5.5)$$

As an easy consequence of this, we get $\{u_k\}$ is a bounded sequence in X_1 . Therefore, up to a subsequence, $u_k \rightharpoonup u$ in X_1 , for some $u \in X_1$ and by Lemma 5.1, we see that u is a critical point of \mathcal{I}_λ .

Claim: $u_k \rightarrow u$ strongly in X_1 .

Since, $u_k \rightarrow u$ strongly in $L^\nu(\Omega)$ for $1 \leq \nu < 2_{s_1}^*$, it implies $\int_{\Omega} a(x)|u_k|^q dx \rightarrow \int_{\Omega} a(x)|u|^q dx$.

Also, by Brezis -Leib Lemma, we have

$$\begin{aligned} \|u_k\|_{X_i}^{p_i} &= \|u_k - u\|_{X_i}^{p_i} + \|u\|_{X_i}^{p_i} + o_k(1), \quad 1 \leq i \leq 2, \quad p_1 = 2, \quad p_2 = p \quad \text{and} \\ \|u_k\|_{NL}^{2.2_\mu^*} &= \|u\|_{NL}^{2.2_\mu^*} + \|u_k - u\|_{NL}^{2.2_\mu^*} + o_k(1). \end{aligned} \quad (5.6)$$

Therefore, by using equations (5.4), (5.5) and (5.6), we get

$$\begin{aligned} \frac{1}{2}\|u_k - u\|_{X_1}^2 + \frac{\beta}{p}\|u_k - u\|_{X_2}^p - \frac{1}{2.2_\mu^*}\|u_k - u\|_{NL}^{2.2_\mu^*} &= c - \mathcal{I}_\lambda(u) + o_k(1) \\ \|u_k - u\|_{X_1}^2 + \beta\|u_k - u\|_{X_2}^p - \|u_k - u\|_{NL}^{2.2_\mu^*} &= o_k(1). \end{aligned} \quad (5.7)$$

Hence, let $\|u_k - u\|_{X_1}^2 + \beta\|u_k - u\|_{X_2}^p \rightarrow l$ and $\|u_k - u\|_{NL}^{2.2_\mu^*} \rightarrow l$, as $k \rightarrow \infty$. If $l = 0$, then claim is proved. So, we assume $l > 0$, then

$$l^{1/2_\mu^*} = \left(\lim_{k \rightarrow \infty} \|u_k - u\|_{NL}^{2.2_\mu^*} \right)^{1/2_\mu^*} \leq (C(n, \mu))^{1/2_\mu^*} \lim_{k \rightarrow \infty} (S^{-1}\|u_k - u\|_{X_1}^2) \leq (C(n, \mu))^{\frac{1}{2_\mu^*}} S^{-1}l.$$

This implies $l \geq C(n, \mu)^{\frac{-1}{2_\mu^*-1}} S^{\frac{2_\mu^*}{2_\mu^*-1}}$, that is, $l \geq \frac{S^{\frac{2n-\mu}{n-\mu+2s_1}}}{(C(n, \mu))^{\frac{n-2s_1}{n-\mu+2s_1}}}$. Now, from (5.7), we have

$$c - \mathcal{I}_\lambda(u) \geq \frac{1}{2}(\|u_k - u\|_{X_1}^2 + \beta\|u_k - u\|_{X_2}^p) - \frac{1}{2.2_\mu^*}\|u_k - u\|_{NL}^{2.2_\mu^*} = \left(\frac{n - \mu + 2s_1}{2(2n - \mu)} \right) l.$$

Therefore, with the help of Theorem 5.1, we get

$$c \geq \left(\frac{n - \mu + 2s_1}{2(2n - \mu)} \right) l + \mathcal{I}_\lambda(u) \geq \left(\frac{n - \mu + 2s_1}{2(2n - \mu)} \right) \frac{S^{\frac{2n-\mu}{n-\mu+2s_1}}}{(C(n, \mu))^{\frac{n-2s_1}{n-\mu+2s_1}}} - C_0 \lambda^{\frac{2}{2-q}} = c_\infty,$$

which is a contradiction. \square

Proof of Theorem 1.3 (i): Let $\gamma_0 > 0$ be such that for all $\lambda \in (0, \gamma_0)$, $c_\infty > 0$ holds. Set $\Lambda = \min\{\gamma_0, \lambda_0\} > 0$. By Proposition 3.6, for all $\lambda \in (0, \Lambda)$, there exists a minimizing sequence $u_k \in \mathcal{M}_\lambda$ such that $\mathcal{I}_\lambda(u_k) = \sigma_\lambda + o_k(1)$ and $\mathcal{I}'_\lambda(u_k) = o_k(1)$. Now using the fact $\sigma_\lambda \leq \sigma_\lambda^+ < 0$ and applying Lemma 5.2, there exists a $u_\lambda \in X_1$ such that $u_k \rightarrow u_\lambda$ in X_1 . By Theorem 5.1 and due to the fact $\sigma_\lambda < 0$, we get $\mathcal{I}'_\lambda(u_\lambda) = 0$ and $\int_\Omega a(x)|u_\lambda|^q > 0$ and hence $u_\lambda \neq 0$, thus $u_\lambda \in \mathcal{M}_\lambda$. Next, we will prove that $u_\lambda \in \mathcal{M}_\lambda^+$. Suppose on the contrary $u_\lambda \in \mathcal{M}_\lambda^-$, then by the fibering map analysis there exist $t_1 < t_2 = 1$ such that $t_1 u_\lambda \in \mathcal{M}_\lambda^+$ and $t_2 u_\lambda \in \mathcal{M}_\lambda^-$. Since ϕ_{u_λ} is increasing in $[t_1, t_2)$, it implies

$$\sigma_\lambda \leq \mathcal{I}_\lambda(t_1 u_\lambda) < \mathcal{I}_\lambda(t_2 u_\lambda) \leq \mathcal{I}_\lambda(u_\lambda) = \sigma_\lambda$$

for $t \in (t_1, 1)$, which is a contradiction. Hence $u_\lambda \in \mathcal{M}_\lambda^+$ and $\sigma_\lambda = \mathcal{I}_\lambda(u_\lambda) = \sigma_\lambda^+$. Moreover, by using same assertions and arguments as in proof of Theorem 1.2, we obtain that u_λ is nonnegative. \square

Consider the family of functions $U_\epsilon(x) = \epsilon^{-\frac{(n-2s_1)}{2}} u^*\left(\frac{x}{\epsilon}\right)$, where $u^*(x) = \bar{u}\left(\frac{x}{S^{2s_1}}\right)$, $\bar{u}(x) = \frac{\tilde{u}(x)}{\|\tilde{u}\|_{2_{s_1}^*}}$ and $\tilde{u}(x) = d(b^2 + |x|^2)^{-\frac{n-2s_1}{2}}$, $d \in \mathbb{R} \setminus \{0\}$ and $b > 0$. ([38]) U_ϵ satisfies

$$(-\Delta)^{s_1} U_\epsilon = |U_\epsilon|^{2_{s_1}^* - 2} U_\epsilon, \text{ in } \mathbb{R}^n, \text{ and } \|U_\epsilon\|_{X_1}^2 = \|U_\epsilon\|_{2_{s_1}^*}^{2_{s_1}^*} = S^{\frac{n}{2s_1}}. \quad (5.8)$$

Let $\eta \in C_c^\infty(\Omega)$ be such that $\eta = 1$ in $B_\delta(0)$, $\eta = 0$ in $B_{2\delta}^c(0)$, and $0 \leq \eta \leq 1$ in Ω . Set $u_\epsilon = \eta U_\epsilon$, then we have the following estimates

Lemma 5.3 (see [19, 38]) *The following hold true*

- (i) $\|u_\epsilon\|_{X_1}^2 \leq S^{\frac{n}{2s_1}} + O(\epsilon^{n-2s_1})$
- (ii) $\|u_\epsilon\|_{2_{s_1}^*}^{2_{s_1}^*} = S^{\frac{n}{2s_1}} + O(\epsilon^n)$
- (iii) $\|u_\epsilon\|_{NL}^2 \leq C(n, \mu) \frac{n}{2 \cdot 2_{s_1}^*} S_H^{\frac{n-2s_1}{2s_1}} + O(\epsilon^n)$
- (iv) $\|u_\epsilon\|_{NL}^2 \geq \left(C(n, \mu) \frac{n}{2s_1} S_H^{\frac{2n-\mu}{2s_1}} - O(\epsilon^n)\right) \frac{1}{2\mu}$.

Now, we will estimate the fractional p -Laplacian norm of the family of functions $\{u_\epsilon\}$.

Lemma 5.4 *Let $1 < p < 2$, then there exists $C > 0$ such that*

$$\|u_\epsilon\|_{X_2}^p \leq C \begin{cases} \epsilon^{\frac{n-2s_1}{2}p} & \text{if } 1 < p < n/(n-s_1) \\ \epsilon^{\frac{2-p}{2}n} & \text{if } n/(n-s_1) \leq p < 2. \end{cases} \quad (5.9)$$

Proof. Following the ideas of [38, Proposition 21], we define the following sets

$$\begin{aligned} D &:= \{(x, y) \in \mathbb{R}^{2n} : x \in B_\delta, y \in B_\delta^c, |x - y| > \delta/2\} \text{ and} \\ E &:= \{(x, y) \in \mathbb{R}^{2n} : x \in B_\delta, y \in B_\delta^c, |x - y| \leq \delta/2\}. \end{aligned}$$

Then, by the definition of u_ϵ , it is clear that

$$\begin{aligned} \|u_\epsilon\|_{X_2}^p &= \int_{\mathbb{R}^{2n}} \frac{|u_\epsilon(x) - u_\epsilon(y)|^p}{|x - y|^{n+ps_2}} dx dy = \int_{B_\delta} \int_{B_\delta} \frac{|U_\epsilon(x) - U_\epsilon(y)|^p}{|x - y|^{n+ps_2}} dx dy \\ &\quad + \left(2 \int_D + 2 \int_E + \int_{B_\delta^c} \int_{B_\delta^c} \right) \frac{|u_\epsilon(x) - u_\epsilon(y)|^p}{|x - y|^{n+ps_2}} dx dy. \end{aligned} \quad (5.10)$$

By [38, Claim 10 of Proposition 21], for $x, y \in B_\delta^c$, we have

$$|u_\epsilon(x) - u_\epsilon(y)| \leq C\epsilon^{(n-2s_1)/2} \min\{1, |x - y|\}.$$

Therefore,

$$I_1 = \int_{B_\delta^c} \int_{B_\delta^c} \frac{|u_\epsilon(x) - u_\epsilon(y)|^p}{|x - y|^{n+ps_2}} dx dy \leq C\epsilon^{\frac{n-2s_1}{2}p} \int_{B_{2\delta}} \int_{\mathbb{R}^n} \frac{\min\{1, |x - y|^p\}}{|x - y|^{n+ps_2}} dx dy = C\epsilon^{\frac{n-2s_1}{2}p}.$$

Next, by [38, (4.21)], for $x \in B_\delta$, $y \in B_\delta^c$ with $|x - y| \leq \delta/2$, we have

$$|u_\epsilon(x) - u_\epsilon(y)| \leq C\epsilon^{(n-2s_1)/2} |x - y|.$$

Then, proceeding similarly, we get

$$I_2 = \int_E \frac{|u_\epsilon(x) - u_\epsilon(y)|^p}{|x - y|^{n+ps_2}} dx dy \leq C\epsilon^{\frac{n-2s_1}{2}p}.$$

To evaluate $I_3 := \int_D \frac{|u_\epsilon(x) - u_\epsilon(y)|^p}{|x - y|^{n+ps_2}} dx dy$, we first note that due to the fact $1 < p < 2$, there exists $A_p > 0$ such that

$$|u_\epsilon(x) - u_\epsilon(y)|^p \leq |U_\epsilon(x) - U_\epsilon(y)|^p + |U_\epsilon(y) - u_\epsilon(y)|^p + A_p |U_\epsilon(x) - U_\epsilon(y)|^{p-1} |U_\epsilon(y) - u_\epsilon(y)|.$$

Now, using $|u_\epsilon(y)| \leq |U_\epsilon(y)| \leq C\epsilon^{(n-2s_1)/2}$ for all $y \in B_\delta^c$ ([38, (4.17)]), we obtain

$$\begin{aligned} \int_D \frac{|U_\epsilon(y) - u_\epsilon(y)|^p}{|x - y|^{n+ps_2}} dx dy &\leq 2^p \int_D \frac{|U_\epsilon(y)|^p}{|x - y|^{n+ps_2}} dx dy \leq C\epsilon^{\frac{n-2s_1}{2}p} \int_D \frac{1}{|x - y|^{n+ps_2}} dx dy \\ &= C\epsilon^{\frac{n-2s_1}{2}p}. \end{aligned} \quad (5.11)$$

Furthermore,

$$\int_D \frac{|U_\epsilon(x) - U_\epsilon(y)|^{p-1} |U_\epsilon(y) - u_\epsilon(y)|}{|x - y|^{n+ps_2}} dx dy \leq 2 \int_D \left(\frac{|U_\epsilon(x)|^{p-1} |U_\epsilon(y)|}{|x - y|^{n+ps_2}} + \frac{|U_\epsilon(y)|^p}{|x - y|^{n+ps_2}} \right) dx dy. \quad (5.12)$$

By [38, (4.17)] and definition of U_ϵ , we have

$$|U_\epsilon(x)|^{p-1} |U_\epsilon(y)| \leq C\epsilon^{\frac{n-2s_1}{2}} \left(b^2 + \left| \frac{x}{\epsilon S^{1/(2s_1)}} \right|^2 \right)^{-\frac{n-2s_1}{2}(p-1)}.$$

Therefore, for $\delta_\epsilon = \delta/\epsilon$, $\xi = \frac{x}{\epsilon S^{1/(2s_1)}}$ and $\zeta = |x - y|$, we deduce that

$$\begin{aligned} \int_D \frac{|U_\epsilon(x)|^{p-1}|U_\epsilon(y)|}{|x-y|^{n+ps_2}} dx dy &\leq C\epsilon^{\frac{n-2s_1}{2}} \epsilon^n \int_{\xi \in B_{\delta_\epsilon}} \int_{|\zeta| > \delta/2} (b^2 + |\xi|^2)^{-\frac{n-2s_1}{2}(p-1)} |\zeta|^{-(n+ps_2)} d\xi d\zeta \\ &\leq C\epsilon^{n+\frac{n-2s_1}{2}} \left[1 + \int_{\xi \in B_{\delta_\epsilon} \setminus B_1} (b^2 + |\xi|^2)^{-\frac{n-2s_1}{2}(p-1)} \right] \\ &\leq C\epsilon^{n+\frac{n-2s_1}{2}} (\epsilon^{(n-2s_1)(p-1)-n} + 1) \leq C\epsilon^{(n-2s_1)(p-1/2)}. \end{aligned}$$

Using this together with (5.11) and the fact that $(n - 2s_1)/2 \leq (n - 2s_1)(p - 1/2)$, (5.12) implies

$$\int_D \frac{|U_\epsilon(x) - U_\epsilon(y)|^{p-1} |U_\epsilon(y) - u_\epsilon(y)|}{|x-y|^{n+ps_2}} dx dy \leq C\epsilon^{\frac{n-2s_1}{2}p}.$$

Thus, from (5.10), we obtain

$$\|u_\epsilon\|_{X_2}^p = \int_{\mathbb{R}^{2n}} \frac{|u_\epsilon(x) - u_\epsilon(y)|^p}{|x-y|^{n+ps_2}} dx dy \leq \int_{\mathbb{R}^{2n}} \frac{|U_\epsilon(x) - U_\epsilon(y)|^p}{|x-y|^{n+ps_2}} dx dy + C\epsilon^{\frac{n-2s_1}{2}p}. \quad (5.13)$$

Now, it remains to evaluate only $\|U_\epsilon\|_{X_2}^p$. We recall $U_\epsilon(x) = \epsilon^{-\frac{n-2s_1}{2}} u^*\left(\frac{x}{\epsilon}\right)$ and $U_1(x) = u^*(x)$. Using suitable change of variables, we deduce that

$$\|U_\epsilon\|_{X_2}^p = \epsilon^{-\frac{n-2s_1}{2}p} \int_{\mathbb{R}^{2n}} \frac{|u^*\left(\frac{x}{\epsilon}\right) - u^*\left(\frac{y}{\epsilon}\right)|^p}{|x-y|^{n+ps_2}} dx dy = \epsilon^{-\frac{n-2s_1}{2}p} \int_{\mathbb{R}^{2n}} \frac{|u^*(z) - u^*(w)|^p}{|z-w|^{n+ps_2}} dz dw,$$

then, using Lemma 2.1 and (5.8), we get

$$\|U_\epsilon\|_{X_2}^p \leq C\epsilon^{-\frac{n-2s_1}{2}p} \|U_1\|_{X_1}^p \leq C\epsilon^{n(1-\frac{p}{2})}. \quad (5.14)$$

Hence, (5.13) and (5.14) give the required result of the Lemma. \square

Lemma 5.5 *There exists $\Lambda_0 > 0$ such that for every $\lambda \in (0, \Lambda_0)$ and $\beta > 0$, there holds $\sigma_\lambda^- < c_\infty$, where c_∞ is as defined in Lemma 5.2, in each of the following cases*

(I) *for all $q > 0$, if $1 < p < n/(n - s_1)$,*

(II) *for all $q \in (1, \frac{n(2-p)}{n-2s_1}) \cup (\frac{4n}{4n-np-4s_1}, p)$, if $n/(n - s_1) \leq p < 2$.*

Proof. To prove the lemma, we will show that there exists $\epsilon > 0$ such that $\sup_{t \geq 0} \mathcal{I}_\lambda(tu_\epsilon) < c_\infty$.

Let $\gamma_0 > 0$ be such that for all $\lambda \in (0, \gamma_0)$, $c_\infty > 0$ holds. Then, with the help of Lemma 2.1, we have

$$\begin{aligned} \mathcal{I}_\lambda(tu_\epsilon) &\leq \frac{t^2}{2} \|u_\epsilon\|_{X_1}^2 + \beta \frac{t^p}{p} \|u_\epsilon\|_{X_2}^p \\ &\leq \frac{t^2}{2} \|u_\epsilon\|_{X_1}^2 + C\beta \frac{t^p}{p} \|u_\epsilon\|_{X_1}^2 \leq C(t^2 + t^p). \end{aligned}$$

Therefore, there exists $t_0 > 0$ such that

$$\sup_{0 \leq t \leq t_0} \mathcal{I}_\lambda(tu_\epsilon) < c_\infty.$$

Let $\Theta(t) = \frac{t^2}{2} \|u_\epsilon\|_{X_1}^2 + \beta \frac{t^p}{p} \|u_\epsilon\|_{X_2}^p - \frac{t^{2.2_\mu^*}}{2.2_\mu^*} \|u_\epsilon\|_{NL}^{2.2_\mu^*}$, then we note that $\Theta(0) = 0$, $\Theta(t) > 0$ for t small enough, $\Theta(t) < 0$ for t large enough, and there exists $t_\epsilon > 0$ such that $\sup_{t \geq 0} \Theta(t) = \Theta(t_\epsilon)$, that is

$$0 = \Theta'(t_\epsilon) = t_\epsilon \|u_\epsilon\|_{X_1}^2 + \beta t_\epsilon^{p-1} \|u_\epsilon\|_{X_2}^p - t_\epsilon^{2.2_\mu^*-1} \|u_\epsilon\|_{NL}^{2.2_\mu^*}$$

which gives us

$$t_\epsilon^{2.2_\mu^*-p} = \frac{1}{\|u_\epsilon\|_{NL}^{2.2_\mu^*}} (t_\epsilon^{2-p} \|u_\epsilon\|_{X_1}^2 + \beta \|u_\epsilon\|_{X_2}^p) < C(1 + t_\epsilon^{2-p}).$$

Since $2.2_\mu^* > 2$, there exists $t_1 > 0$ such that $t_\epsilon \leq t_1$ for all $\epsilon > 0$. Also, for $\delta > 0$ such that $2\delta < \delta_1$, we have

$$\int_\Omega a(x) |u_\epsilon|^q dx = \int_{B_{2\delta}(0)} a(x) |u_\epsilon|^q dx \geq m_a \int_{B_{2\delta}(0)} |u_\epsilon|^q dx \geq m_a \int_{B_\delta(0)} |U_\epsilon|^q dx.$$

Therefore, we obtain

$$\begin{aligned} \sup_{t \geq t_0} \mathcal{I}_\lambda(tu_\epsilon) &\leq \sup_{t \geq 0} \Theta(t) - \frac{t_0^q m_a}{q} \lambda \int_{B_\delta(0)} |U_\epsilon|^q \\ &= \frac{t_\epsilon^2}{2} \|u_\epsilon\|_{X_1}^2 + \beta \frac{t_\epsilon^p}{p} \|u_\epsilon\|_{X_2}^p - \frac{t_\epsilon^{2.2_\mu^*}}{2.2_\mu^*} \|u_\epsilon\|_{NL}^{2.2_\mu^*} - \frac{t_0^q m_a}{q} \lambda \int_{B_\delta(0)} |U_\epsilon|^q \\ &\leq \sup_{t \geq 0} \left(\frac{t^2}{2} \|u_\epsilon\|_{X_1}^2 - \frac{t^{2.2_\mu^*}}{2.2_\mu^*} \|u_\epsilon\|_{NL}^{2.2_\mu^*} \right) + \beta \frac{t_\epsilon^p}{p} \|u_\epsilon\|_{X_2}^p - \frac{t_0^q m_a}{q} \lambda \int_{B_\delta(0)} |U_\epsilon|^q. \end{aligned} \quad (5.15)$$

Let $\Upsilon(t) = \frac{t^2}{2} \|u_\epsilon\|_{X_1}^2 - \frac{t^{2.2_\mu^*}}{2.2_\mu^*} \|u_\epsilon\|_{NL}^{2.2_\mu^*}$. It is easy to verify that Υ attains its maximum at $\hat{t} = \left(\frac{\|u_\epsilon\|_{X_1}^2}{\|u_\epsilon\|_{NL}^{2.2_\mu^*}} \right)^{1/(2_\mu^*-2)}$. Therefore,

$$\sup_{t \geq 0} \Upsilon(t) = \Upsilon(\hat{t}) = \frac{n - \mu + 2s_1}{2(2n - \mu)} \left(\frac{\|u_\epsilon\|_{X_1}^2}{\|u_\epsilon\|_{NL}^{2.2_\mu^*}} \right)^{2_\mu^*/(2_\mu^*-1)},$$

using the estimates of Lemma 5.3 and (2.2), we have

$$\begin{aligned} \sup_{t \geq 0} \Upsilon(t) &\leq \frac{n - \mu + 2s_1}{2(2n - \mu)} \left(\frac{(C(n, \mu)^{\frac{1}{2_\mu^*}} S_H)^{\frac{n}{2s_1}} + O(\epsilon^{n-2s_1})}{(C(n, \mu)^{\frac{n}{2s_1}} S_H^{\frac{2n-\mu}{2s_1}} - O(\epsilon^n))^{\frac{1}{2_\mu^*}}} \right)^{2_\mu^*/(2_\mu^*-1)} \\ &\leq \frac{n - \mu + 2s_1}{2(2n - \mu)} S_H^{\frac{2n-\mu}{n-\mu+2s_1}} + O(\epsilon^{n-2s_1}). \end{aligned} \quad (5.16)$$

For $\delta > 0$, sufficiently small such that $B_{2\delta}(0) \Subset \Omega$, $2\delta < \delta_1$ and $0 < \epsilon < \delta/2$, we have the following estimate ([19])

$$\int_{B_\delta(0)} |U_\epsilon|^q \geq C_3 \begin{cases} \epsilon^{n - \frac{(n-2s_1)}{2}q}, & \text{if } n < (n-2s_1)q \\ \epsilon^{\frac{n}{2}} |\ln \epsilon|, & \text{if } n = (n-2s_1)q \\ \epsilon^{\frac{(n-2s_1)}{2}q}, & \text{if } n > (n-2s_1)q. \end{cases} \quad (5.17)$$

Thus, using (5.9), (5.16) and (5.17) in (5.15), we get

$$\begin{aligned} \sup_{t \geq t_0} \mathcal{I}_\lambda(tu_\epsilon) &\leq \frac{n - \mu + 2s_1}{2(2n - \mu)} S_H^{\frac{2n-\mu}{n-\mu+2s_1}} + C_1 \epsilon^{n-2s_1} + C_4 \begin{cases} \epsilon^{\frac{n-2s_1}{2}p} & \text{if } 1 < p < n/(n-s_1) \\ \epsilon^{\frac{2-p}{2}n} & \text{if } n/(n-s_1) \leq p < 2. \end{cases} \\ &\quad - \lambda C_2 \begin{cases} \epsilon^{n - \frac{(n-2s_1)}{2}q}, & \text{if } n < (n-2s_1)q \\ \epsilon^{\frac{n}{2}} |\ln \epsilon|, & \text{if } n = (n-2s_1)q \\ \epsilon^{\frac{(n-2s_1)}{2}q}, & \text{if } n > (n-2s_1)q. \end{cases} \end{aligned} \quad (5.18)$$

Now, we consider the following cases.

Case(I): If $1 < p < n/(n-s_1)$.

In this case $(n-2s_1)p/2 < n(1-p/2)$ and $1 < q < p < n/(n-s_1) < n/(n-2s_1)$, therefore (5.18) implies

$$\sup_{t \geq t_0} \mathcal{I}_\lambda(tu_\epsilon) \leq \frac{1}{2} \frac{n - \mu + 2s_1}{2n - \mu} S_H^{\frac{2n-\mu}{n-\mu+2s_1}} + C_4 \epsilon^{\frac{n-2s_1}{2}p} - \lambda C_2 \epsilon^{\frac{(n-2s_1)}{2}q}.$$

And, therefore there exists $\gamma_1 > 0$ such that for all $\lambda \in (0, \gamma_1)$

$$C_4 \epsilon^{\frac{n-2s_1}{2}p} - \lambda C_2 \epsilon^{\frac{n-2s_1}{2}q} < -C_0 \lambda^{\frac{2}{2-q}}.$$

Case(II): If $n/(n-s_1) \leq p < 2$.

In this case, $(n-2s_1)p/2 \geq n(1-p/2)$, therefore (5.18) implies

$$\sup_{t \geq t_0} \mathcal{I}_\lambda(tu_\epsilon) \leq \frac{n - \mu + 2s_1}{2(2n - \mu)} S_H^{\frac{2n-\mu}{n-\mu+2s_1}} + C_4 \epsilon^{\frac{2-p}{2}n} - \lambda C_2 \begin{cases} \epsilon^{n - \frac{(n-2s_1)}{2}q}, & \text{if } n < (n-2s_1)q \\ \epsilon^{\frac{n}{2}} |\ln \epsilon|, & \text{if } n = (n-2s_1)q \\ \epsilon^{\frac{(n-2s_1)}{2}q}, & \text{if } n > (n-2s_1)q. \end{cases} \quad (5.19)$$

Subcase(a): If $n > (n-2s_1)q$.

In this case, we see that $\frac{(n-2s_1)}{2}q < n(1 - \frac{p}{2})$, if $q < \frac{n(2-p)}{n-2s_1}$. Then, proceeding similar to Case(I), there exists $\gamma_2 > 0$ such that for all $\lambda \in (0, \gamma_2)$

$$C_4 \epsilon^{\frac{n-2s_1}{2}p} - \lambda C_2 \epsilon^{\frac{n-2s_1}{2}q} < -C_0 \lambda^{\frac{2}{2-q}}.$$

Subcase(b): If $n < (n - 2s_1)q$.

Choosing $\epsilon = (\lambda^{\frac{2}{2-q}})^{\frac{2}{n(2-p)}} \leq \delta$, (5.19) yields

$$\sup_{t \geq t_0} \mathcal{I}_\lambda(tu_\epsilon) \leq \frac{n - \mu + 2s_1}{2(2n - \mu)} S_H^{\frac{2n-\mu}{n-\mu+2s_1}} + C_4 \lambda^{\frac{2}{2-q}} - \lambda C_2 \lambda^{\frac{4}{n(2-q)(2-p)}} \left(n - \frac{(n-2s_1)}{2} q \right).$$

Clearly,

$$1 + \frac{4}{n(2-q)(2-p)} \left(n - \frac{n-2s_1}{2} q \right) < \frac{2}{2-q}$$

if $q > 4n/(4n - np - 4s_1)$. Then, in this situation, there exists $\gamma_3 > 0$ such that for all $\lambda \in (0, \gamma_3)$,

$$C_4 \lambda^{\frac{2}{2-q}} - C_2 \lambda \lambda^{\frac{4}{n(2-q)(2-p)}} \left(n - \frac{n-2s_1}{2} q \right) < -C_0 \lambda^{\frac{2}{2-q}}.$$

Let $\Lambda_0 = \min\{\gamma_1, \gamma_2, \gamma_3, \Lambda, (\frac{\delta}{2})^{n(2-p)/2}\} > 0$, then for all $\lambda \in (0, \Lambda_0)$ and sufficiently small $\epsilon > 0$, we obtain

$$\sup_{t \geq 0} \mathcal{I}_\lambda(tu_\epsilon) < c_\infty.$$

Now choosing $\delta > 0$ sufficiently small, we see that $u_\epsilon \in X_1$ and using Lemma 3.3, there exists $\tilde{t} > 0$ such that $\tilde{t}u_\epsilon \in \mathcal{M}_\lambda^-$. Hence,

$$\sigma_\lambda^- \leq \mathcal{I}_\lambda(\tilde{t}u_\epsilon) \leq \sup_{t \geq 0} \mathcal{I}_\lambda(tu_\epsilon) < c_\infty.$$

This completes proof of the Lemma. \square

Proof of Theorem 1.3 (ii): With the help of Lemma 3.5 and Proposition 3.6, we get a minimizing sequence $\{v_k\} \subset \mathcal{M}_\lambda^-$, which is also $(PS)_{\sigma_\lambda^-}$ sequence. By Lemmas 5.5 and 5.2, there exists $v_\lambda \in X_1$ such that $v_k \rightarrow v_\lambda$ in X_1 . Using Theorem 5.1 and strong convergence, $v_\lambda \in \mathcal{M}_\lambda^-$ and $\mathcal{I}_\lambda(v_\lambda) = \sigma_\lambda^-$. Thus, v_λ is a weak solution of problem (P_λ) in \mathcal{M}_λ^- and since $u_\lambda \in \mathcal{M}_\lambda^+$, $u_\lambda \neq v_\lambda$. \square

Lemma 5.6 *There exists $\Lambda_{00}, \beta_{00} > 0$ such that for every $\lambda \in (0, \Lambda_{00})$ and $\beta \in (0, \beta_{00})$, there holds*

$$\sigma_\lambda^- < c_\infty.$$

Proof. Here we only give outline of the proof because arguments are similar to the previous lemma and [21, Lemma 4.7]. Let $\beta = \epsilon^\alpha$, where $\alpha > (n - 2s_1)$ and taking into account (5.16) and (5.17), from (5.15), it follows that

$$\sup_{t \geq t_0} \mathcal{I}_\lambda(tu_\epsilon) \leq \frac{n - \mu + 2s_1}{2(2n - \mu)} S_H^{\frac{2n-\mu}{n-\mu+2s_1}} + C_1 \epsilon^{n-2s_1} - \lambda C_2 \begin{cases} \epsilon^{n - \frac{(n-2s_1)}{2} q}, & \text{if } n < (n - 2s_1)q \\ \epsilon^{\frac{n}{2}} |\ln \epsilon|, & \text{if } n = (n - 2s_1)q \\ \epsilon^{\frac{(n-2s_1)}{2} q}, & \text{if } n > (n - 2s_1)q. \end{cases}$$

The case, when $n > (n - 2s_1)q$, follows exactly on the same lines of Case(I) of Lemma 5.5. For the other case, we set $\epsilon = (\lambda^{\frac{2}{2-q}})^{\frac{1}{n-2s_1}} \leq \delta$ and proceed similarly to [21, Lemma 4.7], to get the required result of the lemma for some $\Lambda_{00}, \beta_{00} > 0$. \square

Proof of Theorem 1.4: Proof follows exactly on the same lines of Proof of Theorem 1.3 (ii) by using Lemma 5.6 instead of Lemma 5.5. \square

References

- [1] S. Aizicovici, N.S. Papageorgiou and V. Staicu, *Nodal solutions for $(p, 2)$ -equations* Trans. Amer. Math. Soc. 367 (2015), no. 10, 7343-7372.
- [2] C.O. Alves, M.G. Figueiredo and M. Yang, *Existence of solutions for a nonlinear Choquard equation with potential vanishing at infinity*, Adv. Nonlinear Anal. 5 (2016), no. 4, 331-345.
- [3] A. Bahrouni, V.D. Rădulescu and D.D. Repovš, *Double phase transonic flow problems with variable growth: nonlinear patterns and stationary waves*, Nonlinearity 32 (2019), no. 7, 2481-2495.
- [4] P. Baroni, M. Colombo and G. Mingione, *Harnack inequalities for double phase functionals*, Nonlinear Anal. 121 (2015), 206-222.
- [5] P. Baroni, M. Colombo and G. Mingione, *Regularity for general functionals with double phase*, Calc. Var. Partial Differential Equations 57 (2018), Art 62.
- [6] L. Beck and G. Mingione, *Lipschitz bounds and non-uniform ellipticity*, Commun. Pure Appl. Math., to appear.
- [7] M. Bhakta and D. Mukherjee, *Multiplicity results for (p, q) fractional elliptic equations involving critical nonlinearities*, Adv. Differential Equ. 3/4(4) (2019), 185-228.
- [8] V. Bobkov and M. Tanaka, *On sign-changing solutions for (p, q) -Laplace equations with two parameters*, Adv. Nonlinear Anal. 8 (2019), no. 1, 101-129.
- [9] L. Brasco and E. Parini, *The second eigenvalue of the fractional p -Laplacian*, Adv. Calc. Var. 9 (2015), no. 4, 323-355.
- [10] B. Buffoni, L. Jeanjean and C.A. Stuart, *Existence of a nontrivial solution to a strongly indefinite semilinear equation*, Proc. Amer. Math. Soc. 119 (1993), no. 1, 179-186.
- [11] L. Caffarelli, *Nonlocal equations, drifts and games*, Nonlinear Partial Differential equations, Abel Symposia 7 (2012), 37-52.
- [12] M. Cencelj, V.D. Rădulescu and D.D. Repovš, *Double phase problems with variable growth*, Nonlinear Anal. 177 (2018), part A, 270-287.
- [13] L. Cherfils and Y. Ilyasov, *On the stationary solutions of generalized reaction diffusion equations with p & q -Laplacian*, Commun. Pure Appl. Anal. 4 (2005), 9-22.
- [14] E. Di Nezza, G. Palatucci and E. Valdinoci, *Hitchhikers guide to the fractional Sobolev spaces*, Bull. Sci. Math. 136 (2012), 521-573.
- [15] P. D'Avenia, G. Siciliano and M. Squassina, *Existence results for a doubly nonlocal equation*, São Paulo J. Math. Sci. 9 (2015), no. 2, 311-324.

- [16] I. Ekeland, *On the variational principle*, J. Math. Anal. Appl. 17 (1974), 324-353.
- [17] P. C. Fife, *Mathematical Aspects of Reacting and Diffusing Systems*, Lect. Notes in Biomath. 28, Springer, Berlin, 1979.
- [18] F. Gao, M. Yang, *On the Brezis-Nirenberg type critical problem for nonlinear Choquard equation*, Sci. China Math. 61 (2018), no. 7, 1219-1242.
- [19] J. Giacomoni, T. Mukherjee and K. Sreenadh, *Doubly nonlocal system with Hardy-Littlewood-Sobolev critical nonlinearity*, J. Math. Anal. Appl. 467 (2018), 638-672.
- [20] S. Goyal and K. Sreenadh, *Existence of multiple solutions of p -fractional Laplace operator with sign-changing weight function*, Adv. Nonlinear Anal. 4 (2015), no. 1, 3758.
- [21] D. Goel, D. Kumar and K. Sreenadh, *Regularity and multiplicity results for fractional (p, q) -Laplacian equations*, Commun. Contemp. Math. (2019), <https://doi.org/10.1142/S0219199719500652>.
- [22] D. Goel and K. Sreenadh, *Kirchhoff equations with Hardy-Littlewood-Sobolev critical nonlinearity*, Nonlinear Analysis 186 (2019), 162-186.
- [23] E. Lieb, *Existence and uniqueness of the minimizing solution of Choquard's nonlinear equation*, Stud. Appl. Math. 57 (1976/1977), 93-105.
- [24] S. A. Marano, G. Marino and N.S. Papageorgiou, *On a Dirichlet problem with (p, q) -Laplacian and parametric concave-convex nonlinearity*, J. Math. Anal. Appl. 475 (2019), 1093-1107.
- [25] S. Marano and S. Mosconi, *Some recent results on the Dirichlet problem for (p, q) -Laplacian equation*, Discrete Contin. Dyn. Syst. Ser. S 11 (2018), 279-291.
- [26] X. Mingqi, V.D. Rădulescu, B. Zhang, *Fractional Kirchhoff problems with critical Trudinger-Moser nonlinearity*, Calc. Var. Partial Differential Equations 58 (2019), no. 2, Art. 57, 27 pp.
- [27] X. Mingqi, V.D. Rădulescu, B. Zhang, *A critical fractional Choquard-Kirchhoff problem with magnetic field*, Comm. Contem. Math. 21 (2019), no. 4, 185004, 36 pp.
- [28] V. Moroz and J. V. Schaftingen, *A guide to the Choquard equation*, J. Fixed Point Theory Appl. 19 (2017), no. 1, 773-813.
- [29] T. Mukherjee and K. Sreenadh, *Fractional Choquard equation with critical nonlinearities*, NoDEA Nonlinear Differential Equations Appl. 24 (2017), 63.
- [30] N.S. Papageorgiou and V. D. Rădulescu, *Qualitative phenomena for some classes of quasilinear elliptic equations with multiple resonance*, Appl. Math. Optim. 69 (2014), 393-430.
- [31] N.S. Papageorgiou, V.D. Rădulescu and D.D. Repovš, *Double-phase problems with reaction of arbitrary growth*, Z. Angew. Math. Phys. 69 (2018), no. 4, Art. 108, 21 pp.
- [32] N.S. Papageorgiou, V.D. Rădulescu and D.D. Repovš, *Double-phase problems and a discontinuity property of the spectrum*, Proc. Amer. Math. Soc. 147 (2019), no. 7, 2899-2910.

- [33] N.S. Papageorgiou, V.D. Rădulescu and D.D. Repovš, *Nonlinear Analysis–Theory and Methods*, Springer Monographs in Mathematics, Springer Nature, Cham, 2019.
- [34] S. Pekar, *Untersuchung über die Elektronentheorie der Kristalle*, Akademie Verlag, Berlin, 1954.
- [35] H. Pham, *Optimal stopping, free boundary, and American option in a jump-diffusion model*, Appl. Math. Optim. 35 (1997), no. 2, 145-164.
- [36] P. Pucci, M. Xiang and B. Zhang, *Existence results for Schrödinger-Choquard-Kirchhoff equations involving the fractional p -Laplacian*, Adv. Calc. Var. 12 (2019), 253-275.
- [37] V.D. Rădulescu, *Isotropic and anisotropic double phase problems: old and new*, Opuscula Math. 39 (2019), 259-279.
- [38] R. Servadei and E. Valdinoci, *The Brezis-Nirenberg result for the fractional Laplacian*, Trans. Amer. Math. Soc. 367 (2015), no. 1, 67-102.
- [39] Y. Su and H. Chen, *Fractional Kirchhoff-type equation with Hardy-Littlewood-Sobolev critical exponent*, Computers and Mathematics with Applications (2019), <https://doi.org/10.1016/j.camwa.2019.03.052>.
- [40] H. Wilhelmsson, *Explosive instabilities of reaction-diffusion equations*, Phys. Rev. A, 36 (1987), 965-966.
- [41] M. Xiang, B.Zhang, V.D.Rădulescu, *Superlinear Schrödinger-Kirchhoff type problems involving the fractional p -Laplacian and critical exponent*, Adv. Nonlinear Anal. 9 (2020), no. 1, 690-709.
- [42] H. Yin and Z. Yang, *A class of $p - q$ -Laplacian type equation with concaveconvex nonlinearities in bounded domain*, J. Math. Anal. Appl. 382 (2011), 843-855.
- [43] Q. Zhang and V.D. Rădulescu, *Double phase anisotropic variational problems and combined effects of reaction and absorption terms*, J. Math. Pures Appl. 118 (2018), 159203.