

Graceful Labeling For bipartite graceful Graphs and related Graphs^{*}

Chunfeng Liu^a, Xiuying Wang^b, Guozhu Liu^{b*}

(^aCollege of Science, Liaoning University of Technology, Liaoning, Jinzhou, 121013)

(^bCollege of Information Science and Technology, Qingdao University of Science and Technology, Shandong, Qingdao 266000)

Abstract: The concept of graceful labels was proposed by Rosa, scholars began to study graceful labels of various graphs and obtained relevant results. In this paper, let G is a bipartite graceful graph, we proved that $S(t \bullet G)$, $P_n^t(tn \bullet G)$, $C(t \bullet G)$ and G^* are graceful labelings.

Key words: Graceful labeling, Bipartite graceful Graphs, Open star of graphs, Cycle of a graph, Path union of a graph

1. INTRODUCTION AND LEMMEL

The graceful labeling was introduced by A Rosa [1] in 1967. Golomb [2] proved that the complete bipartite graph is graceful. Barrientos [3] proved that union of complete bipartite graphs is also graceful. Vaidya [4] introduced a star of cycle. Kaneria and Makadia [5] proved that star of a cycle is graceful. Kaneria [6] proved that join sum of path union and star are graceful graphs. The detail survey of graph labeling refer Gallian [7].

Let G is a bipartite graceful graph, In this paper, we proved that $S(t \bullet G)$, $P_n^t(tn \bullet G)$, $C(t \bullet G)$ and G^* are graceful graphs. We assume G is a simple undirected, finite graph, with vertices and edges. For all terminology and notations we follow Harary [8]. We shall give brief summary of definitions which are useful in this paper.

Definition 1.1 A function f is called graceful labeling of a (p, q) -graph G if $f: V(G) \rightarrow \{0, 1, 2, \dots, q\}$ is injective and induced function $f^*: E(G) \rightarrow \{0, 1, 2, \dots, q\}$ defined as $f(e) = |f(u) - f(v)|$ is bijective for every edge $e = uv$. A graph G is called graceful graph if it admits a graceful labeling.

Definition 1.2 Let (p, q) -graph G is a bipartite graph with the bipartition (X, Y) , f is a graceful labeling of G , if $\max\{f(x) | x \in X\} < \min\{f(y) | y \in Y\}$, then f called is a bipartite graceful labeling of G . if G admits a bipartite graceful labeling, then call G is a bipartite graceful graph.

Definition 1.3 Let G_1, G_2, \dots, G_n ($n \geq 2$) be n graph, Then the graph obtained by adding an edge from G_i to G_{i+1} ($1 \leq i \leq n-1$) is called path union of graph G_1, G_2, \dots, G_n . We shall denote such graph by $P(G_1, G_2, \dots, G_n)$.

If $G_1 = G_2 = \dots = G_n = G$, we shall denote by $P(nt \bullet G)$.

Definition 1.4^[10] A graph obtained by replacing each vertex of $K_{1,t}$ except the apex vertex by the graphs G_1, G_2, \dots, G_t is known as open star of graphs. We shall denote such graph by $S(G_1, G_2, \dots, G_t)$.

If we replace each vertices of $K_{1,n}$ except the apex vertex by a graph G . i.e. $G_1 = G_2 = \dots = G_t = G$, such open star of a graph, we shall denote by $S(t \bullet G)$.

Definition 1.5^[10] A graph is obtained by replacing each edge of $K_{1,t}$ by a path of P_n length n on $n+1$ vertices is called one point union for t copies of path P_n . We shall denote such graph G by P_n^t .

Definition 1.6^[10] A graph obtained by replacing each vertices of except the central vertex by the graphs G_1, G_2, \dots, G_m is known as one point union for path of graphs. We shall denote such graph G by $P_n^t(G_1, G_2, \dots, G_m)$, where P_n^t is the one point union of t copies of path P_n . If we replace each vertices of P_n^t except the central vertex by a graph G , i.e. $G = G_1 = G_2 = \dots = G_m$ such one point union of path graph, we shall denote it by $P_n^t(tn, G)$.

Definition 1.7^[11] For a cycle C_n , each vertices of C_n is replace by connected graphs G_1, G_2, \dots, G_n is known as *cycle of graphs* and we shall denote it by $C(G_1, G_2, \dots, G_n)$. If we replace each vertices by graph G , i.e. $G = G_1 = G_2 = \dots = G_n$, such cycle of a graph G , we shall denote it by $C(n \bullet G)$.

^{*}This work was Supported by the National Natural Science Foundation of China (61773107, 61603168)

*Corresponding Author at College of Information Science and Technology, Qingdao University of Science and Technology, Shandong, Qingdao 266000, E-mail: lgz_0228@163.com

Definition 1.8^[6] A graph obtained by replacing each vertex of star $K_{1,n}$ by a graph G of n vertices is called star of G and it is Denoted by G^\star . The graph G which replaced at the center of $K_{1,n}$ we call the central copy of G^\star .

Let G is a bipartite graceful graph, In this paper, We also proved that $P(n \cdot G), S(t \cdot G) P_n^l(tn \cdot G), C(t \cdot G)$ and G^\star are graceful graphs.

Obviously, there are following conclusions:

Lemma 1.1 P_n is bipartite graceful graph.

Lemma 1.2 $C_n (n \equiv 0 \pmod{4})$ is bipartite graceful graph.

Lemma 1.3 $K_{m,n} (m, n \in \mathbb{N})$ is bipartite graceful graph.

Lemma 1.4 $P_m \times P_n (m, n \in \mathbb{N})$ is bipartite graceful graph.

2 MAIN RESULTS

Theorem 2.1 Let G is bipartite graceful graph and $G_i (i=1, 2, \dots, t)$ is t copies of graph G , Then $P(n \cdot G)$ is graceful.

Proof: Let (p_0, q_0) -graph G is a bipartite graph with the bipartition (V_1, V_2) , and $V_1 = \{u_i | i \in [1, m]\}$, $V_2 = \{v_j | j \in [1, r]\}$, f_0 is a bipartite graceful labeling of G , $f_0(u_i) < f_0(u_{i+1}), i \in [1, m-1]$, $f_0(v_j) < f_0(v_{j+1}), j \in [1, r-1]$, and $f_0(u_m) < f_0(v_1)$. Let $u_{l,i} (i \in [1, m]), v_{l,j} (j \in [1, r])$ be the vertices of G which is l^{th} copy of the path union of n copies of G . We shall join $v_{l,1}$ with $u_{l,1}$.

We shall define labeling function of $P(n \cdot G) f$ as follows:

$f: P(n \cdot G) \rightarrow [0, q]$, Where $q = n(q_0 + 1) - 1$ defined by,

$$f(u_{1,i}) = f_0(u_i), \quad i \in [1, m];$$

$$f(v_{1,j}) = q - q_0 + f_0(v_j), \quad j \in [1, r];$$

$$f(u_{l,i}) = f(u_{l-1,m}) + f_0(u_i) + 1, \quad i \in [1, m], l \in [2, n];$$

$$f(v_{l,j}) = f(v_{l-1,1}) + f_0(v_j) - q_0 - 1, \quad j \in [1, r], l \in [2, n].$$

Above labeling pattern give rise graceful labeling to the path union graph $P(n \cdot G)$ of n copies of G .




Figure 1. Graph P_{14} and its bipartite graceful labeling




Figure 2. Graph $P(5 \cdot P_{14})$ and its bipartite graceful labeling

By theorem 2.1 and lemma 1.1-1.4, respectively, have following consequences.

Corollary 2.1.1 $P(t \cdot P_n)$ is graceful.

Corollary 2.1.2^[13] $P(t \cdot C_n)$ is graceful graph, where $n \equiv 0 \pmod{4}$.

Corollary 2.1.3 $P(t \cdot K_{m,n})$ is graceful.

Corollary 2.1.4^[12] $P(t \cdot P_m \times P_n)$ is graceful.

Theorem 2.2 Let G is bipartite graceful graph and $G_i(i= 1,2,\dots, t)$ is be t copies of graph G , Then $S(t \cdot G)$ is graceful.

Proof: Let (p_0, q_0) -graph G is a bipartite graph with the bipartition (V_1, V_2) , and $V_1 = \{u_i | i \in [1, m]\}$, $V_2 = \{v_j | j \in [1, n]\}$, f_0 is a bipartite graceful labeling of G , $f_0(u_i) < f_0(u_{i+1})$, $i \in [1, m-1]$, $f_0(v_j) < f_0(v_{j+1})$, $j \in [1, n-1]$, and $f_0(u_m) < f_0(v_1)$. Let $S(t \cdot G)$ be a graph obtained by replacing each vertices of $K_{1,t}$, except the apex vertex by the graph G . Let v_0 is the apex vertex of $K_{1,t}$. i.e. it is central vertex of the graph $S(t \cdot G)$. Let $u_{li}(1 \leq i \leq m)$, $v_{lj}(1 \leq j \leq n)$ be the vertices of l^{th} copy of $G^{(l)}$ of G in $S(t \cdot G)$, $\forall l \in [1, t]$.

We shall join v_{lm} with the vertex v_0 by an edge to form the open star of graphs $S(t \cdot G)$, $\forall l \in [1, t]$. where $q = t(q_0 + 1)$. We define labeling function $f: V(G) \rightarrow [1, q]$ as follows:

$$\begin{aligned} f(v_0) &= 0; \\ f(u_{1,i}) &= f_0(u_i) + 1, & i \in [1, m]; \\ f(v_{1,j}) &= q - q_0 + f_0(v_j), & j \in [1, n]; \\ f(u_{2,i}) &= f(u_{1,i}) + q - q_0 - 1, & i \in [1, m]; \\ f(v_{2,j}) &= f(v_{1,j}) - q + q_0 + 1, & j \in [1, n]; \\ f(u_{l,i}) &= f(u_{l-2,i}) - (-1)^l (q_0 + 1), & i \in [1, m], l \in [3, t]; \\ f(v_{l,j}) &= f(v_{l-2,j}) + (-1)^l (q_0 + 1), & j \in [1, n], l \in [3, t] \end{aligned}$$

Above labeling pattern give rise graceful labeling to the graph and so it is a graceful graph.



(a) Graph Q and its bipartite graceful labeling (b) Graph $K_{4,3}$ and its bipartite graceful labeling
Figure 3




Figure 4. A graph obtained by open star of Q and its graceful labeling .




Figure 5. A graph obtained by open star of $K_{4,3}$ and its graceful labeling .

By theorem 2.2 and lemma 1.1–1.4, there are following consequences respectively.

Corollary 2.2.1 $S(t \cdot P_n)$ is graceful.

Corollary 2.2.2^[10] $S(t \cdot C_n)$ is graceful graph, where $n \equiv 0 \pmod{4}$.

Corollary 2.2.3^[9] $S(t \cdot K_{m,n})$ is graceful.

Corollary 2.2.4^[9] $S(t \cdot P_m \times P_n)$ is graceful.

Theorem 2.3 Let G is bipartite graceful graph and $G_i (i= 1, 2, \dots, t)$ is i th copies of graph G , Then $P_n^t(tn \cdot G)$ is graceful.

Proof Let (p_0, q_0) -graph G is a bipartite graph with the bipartition (V_1, V_2) , and $V_1 = \{u_i \mid i \in [1, m]\}$, $V_2 = \{v_j \mid j \in [1, r]\}$, f_0 is a bipartite graceful labeling of G , $f_0(u_i) < f_0(u_{i+1})$, $i \in [1, m-1]$, $f_0(v_j) < f_0(v_{j+1})$, $j \in [1, r-1]$, and $f_0(u_m) < f_0(v_1)$. Let be a graph obtained by replacing each vertices of P_n^t except the central vertex by the graph G i.e. $P_n^t(tn \cdot G)$ is the graph obtained by replacing each vertices of $K_{1,t}$ except the apex vertex by the path union of n copies of the graph G . Let u_0 be the central vertex for the graph $P_n^t(tn \cdot G)$. Let $u_{s,l,i} (i \in [1, m])$, $v_{s,l,j} (j \in [1, r])$ be the vertices of G which is l th copy of the path union of n copies of G lies in s th branch of the graph $P_n^t(tn \cdot G)$, $l \in [1, n]$ and $s \in [1, t]$.

First we shall join $v_{s,l,m}$ with the vertex $u_{s,l+1,1}$ by an edge to form the path union of n copies of G for s th branch of $P_n^t(tn \cdot G)$, $l \in [1, n-1]$ and $s \in [1, t]$. Now we shall join $u_{s,1,1}$ with the vertex u_0 by an edge to form the one point union for path of graphs $P_n^t(tn \cdot G)$, $s \in [1, t]$. We shall define labeling function f for the first copy (branch) of the path union of n copies of G as follows:

$f: P_n^t(tn \cdot G) \rightarrow [0, q]$, Where $q = n(q_0 + 1) - 1$ defined by,

$$\begin{aligned} f(u_{1,1,i}) &= f_0(u_i), & i \in [1, m]; \\ f(v_{1,1,j}) &= q - q_0 + f_0(v_j), & j \in [1, r]; \\ f(u_{1,l,i}) &= f(u_{1,l-1,m}) + f_0(u_i) + 1, & i \in [1, m], l \in [2, n]; \\ f(v_{1,l,j}) &= f(v_{1,l-1,1}) + f_0(v_j) - q_0 - 1, & j \in [1, r], l \in [2, n]. \end{aligned}$$

Above labeling pattern give rise graceful labeling to the path union of n copies of G , which lies in first branch of $P_n^t(tn \cdot G)$. Now we shall define labeling $g: V(P_n^t(tn \cdot G)) \rightarrow [0, Q]$, where $Q = |E(P_n^t(tn \cdot G))| = tn(q_0 + 1)$ as follows:

$$\begin{aligned} g(u_0) &= 0; \\ g(u_{1,l,i}) &= f(u_{1,l,i}) + 1, & i \in [1, m], l \in [1, n]; \\ g(v_{1,l,j}) &= f(v_{1,l,j}) + Q - q, & j \in [1, r], l \in [1, n]; \\ g(u_{2,l,i}) &= g(u_{1,l,i}) + Q - q - 1, & i \in [1, m], l \in [1, n]; \\ g(v_{2,l,j}) &= g(v_{1,l,j}) - Q + q + 1, & j \in [1, r], l \in [1, n]; \\ g(u_{s,l,i}) &= g(u_{s-2,l,i}) - (-1)^s(q+1), & i \in [1, m], l \in [1, n], s \in [3, t]; \\ g(v_{s,l,j}) &= g(v_{s-2,l,j}) + (-1)^s(q+1), & j \in [1, r], l \in [1, n], s \in [3, t]. \end{aligned}$$

Above labeling pattern give rise graceful labeling to the graph $P_n^t(tn \cdot G)$ and so it is a graceful graph.



Figure 6 . A graph Q and its bipartite graceful labeling




Figure 7 . The path union 5 copies of Q and its graceful labeling




Figure 8 . $P_3^3(3 \times 5 \cdot Q)$ and its graceful labeling

By theorem 2.3 and lemma 1.1-1.4, respectively, have following consequences.

Corollary 2.3.1 $P_n^t(tn \cdot P_n)$ is graceful.

Corollary 2.3.2^[10] $P_n^t(tn \cdot K_{1,m})$ is graceful.

Corollary 2.3.3^[10] $P_n^t(tn \cdot C_m)$ is graceful, where $m \equiv 0 \pmod{4}$.

Corollary 2.3.4^[9] $P_n^t(tn \cdot K_{m,r})$ is graceful.

Corollary 2.3.5^[10] $P_n^t(tn \cdot P_r \times P_s)$ is graceful.

Theorem 2.4 Cycle $C(t \cdot G)$ ($t \equiv 0 \pmod{2}$) of bipartite graceful graph G is graceful graphs.

Proof Let Let (p_0, q_0) -graph G is a bipartite graph with the bipartition (V_1, V_2) , and $V_1 = \{u_i | i \in [1, m]\}$, $V_2 = \{v_j | j \in [1, n]\}$, f_0 is a bipartite graceful labeling of G , $f_0(u_i) < f_0(u_{i+1})$, $i \in [1, m-1]$, $f_0(v_j) < f_0(v_{j+1})$, $j \in [1, n-1]$, and $f_0(u_m) < f_0(v_1)$. $C(t \cdot G)$ be a graph which contains t copies of the bipartite graceful graph G , where $t \equiv 0 \pmod{2}$. Let $u_{i,j}$ ($i \in [1, m]$) and $v_{i,j}$ ($j \in [1, n]$) be vertices of i^{th} copy of G , $i \in [1, t]$. Now join $v_{i,1}$ with $u_{i+1,1}$ ($i \in [1, (t/2)-1]$), join $v_{i,2,1}$ with $v_{(t/2)+1,1}$, join $u_{i,1}$ with $v_{i+1,1}$ ($i \in [(t/2)+1, t-1]$) and join $u_{t,1}$ with $u_{1,1}$ by an edge to form cycle of graphs $C(t \cdot G)$.

We define the labeling function $f: V(C(tG)) \rightarrow [0, q]$, where $q = t(q_0 + 1)$ as follows:

$$\begin{aligned} f(u_{1,j}) &= f_0(u_j), & j \in [1, m]; \\ f(v_{1,j}) &= q - f_0(v_n) + f_0(v_j), & j \in [1, n]; \\ f(u_{l,i}) &= f(u_{l-1,m}) + 1 + f_0(u_j), & j \in [1, m], l \in [2, t/2]; \\ f(v_{l,i}) &= f(v_{l-1,1}) - 1 - f_0(v_n) + f_0(v_j), & j \in [1, n], l \in [2, t/2]; \\ f(u_{(t/2)+1,i}) &= f(v_{t/2,1}) - 2 - f_0(u_m) + f_0(u_j), & j \in [1, m]; \end{aligned}$$

$$f(v_{(t/2+1,j)})=f(u_{t/2,m})+1-f_0(v_1)+f_0(v_j), \quad j \in [1,n];$$

$$f(u_{l,i})=f(u_{l-1,1})-1-f_0(u_m)+f_0(u_i), \quad j \in [1,m], l \in [(t/2)+2, t];$$

$$f(v_{l,j})=f(v_{l-1,n})+1-f_0(v_1)+f_0(v_j), \quad j \in [1,n], l \in [(t/2)+2, t].$$

Above labeling pattern give rise graceful labeling to the graph $C(tG)$ and so it is a graceful graph.




Figure 9 . G_0 and its bipartite graceful labeling




Figure 10. Cycle $C(6 \cdot G_0)$ and its graceful labeling

By theorem 2.4 and lemma 1.1-1.4, respectively, have following consequences.

Corollary 2.4.1 Cycle of path graph $C(t \cdot P_n)$ is graceful.

Corollary 2.4.2^[14] Cycle of C_{4m} graphs $C(t \cdot C_{4m})$, $t \equiv 0 \pmod{2}$ is graceful graphs.

Corollary 2.4.3^[11] Cycle of complete bipartite graphs $C(t \cdot K_{m,n})$, $t \equiv 0 \pmod{2}$, $m, n \in \mathbb{N}$ is graceful graphs.

Corollary 2.4.5^[12] Cycle of grid graph $C(t \cdot P_r \times P_s)$ is graceful.

Theorem 2.5. G^\star the star of a bipartite graceful graph G , is graceful graphs.

Proof Let Let (p_0, q_0) -graph G is a bipartite graph with the bipartition (V_1, V_2) , and $V_1 = \{u_i | i \in [1, m]\}$, $V_2 = \{v_j | j \in [1, n]\}$, f_0 is a bipartite graceful labeling of G , $f_0(u_i) < f_0(u_{i+1})$, $i \in [1, m-1]$,

$f_0(v_j) < f_0(v_{j+1})$, $j \in [1, n-1]$, and $f_0(u_m) < f_0(v_1)$. Let $u_{0,i}$ ($1 \leq i \leq m$), $v_{0,j}$ ($1 \leq j \leq n$) be vertices of central copy of G and $u_{l,i}$ ($1 \leq i \leq m$), $v_{l,j}$ ($1 \leq j \leq n$) be vertices of other copies of G , $\forall l = 1, 2, \dots, m+n$. We define labeling function $f: V \rightarrow \{0, 1, \dots, q\}$, where $q = (m+n+1)q_0 + m+n$ as follows:

$$f(u_{0,i})=f_0(u_i), \quad i \in [1, m];$$

$$f(v_{0,j})=q-f_0(v_n)+f_0(v_j) \quad j \in [1, n];$$

$$f(u_{1,i})=f(v_{0,1})-1-f_0(u_m)+f_0(u_i), \quad i \in [1, m];$$

$$f(v_{1,j})=f_0(v_j), \quad j \in [1, n];$$

$$f(u_{l,i})=f(u_{l-2,i})+q_0+1, \quad i \in [1, m], l \in [1, m+n] \text{ and } l \text{ even};$$

$$f(v_{l,j})=f(v_{l-2,j})-q_0-1, \quad j \in [1, n], l \in [1, m+n] \text{ and } l \text{ even};$$

$$f(u_{l,i})=f(u_{l-2,i})-q_0-1, \quad i \in [1, m], l \in [1, m+n] \text{ and } l \text{ odd};$$

$$f(v_{l,j})=f(v_{l-2,j})+q_0+1, \quad j \in [1, n], l \in [1, m+n] \text{ and } l \text{ odd}.$$

Above defined labeling function f give rise edge labels $1, 2, \dots, q_0, q_0+2, q_0+3, \dots, 2q_0+1, 2q_0+3, 2q_0+4, \dots, 3q_0+2, \dots, (q_0+1)(m+n)+1, \dots, q$ to all the copies of G . To make G^* as graceful graph $q_0+1, 2(q_0+1), \dots, (m+n)(q_0+1)$ edge labels to be require.

Now we see that the difference of vertex labels for the central copy $G^{(0)}$ with its other copies $G^{(i)}, 1 \leq i \leq m+n$ is precisely following sequence.

$$\begin{aligned}
 & (m+n)(q_0+1) \\
 & \quad q_0+1 \\
 & (m+n-1)(q_0+1) \\
 & \quad 2(q_0+1) \\
 & \quad \dots \\
 & (m+n)(q_0+1)/2, \text{ when } m+n \equiv 0 \pmod{2} \\
 & \quad \text{or} \\
 & (m+n-1)(q_0+1)/2, \text{ when } m+n \equiv 1 \pmod{2} \\
 & \quad (m+n+1)(q_0+1)/2
 \end{aligned}$$

Using above sequence we can produce required edge labels by joining corresponding vertex of $G^{(0)}$ with its other copies of $G^{(i)}, \forall i = 1, 2, \dots, m+n$. Thus G^* admits a graceful labeling and so it is a graceful graph.




Figure 11 . G and its bipartite graceful labeling




Figure 12 . G^* and its bipartite graceful labeling

By theorem 2.5 and lemma 1.1-1.4, respectively, have following consequences.

Corollary 2.5.1 P_n^* of path graph P_n is graceful

Corollary 2.5.2 C_{4m}^* of C_{4m} graphs is graceful graphs.

Corollary 2.5.3^[6] $K_{m,n}^*$ of complete bipartite graphs is graceful graphs.

Corollary 2.5.4^[12] $(P_r \times P_s)^*$ of grid graph $P_r \times P_s$ is graceful.

REFERENCES

- [1] A. Rosa, On certain valuation of the vertices of a graph, Theory of Graphs, (Rome, July 1966), Goden and Breach, N.Y. and Paris, 1967, 329- 335.
- [2] S. W. Golomb, How to number a graph, in Graph Theory And Computing, R. C. Read, ed., Academic Press, New York, (1972), 23- 37.

- [3] C. Barrientos, The gracefulness of unions of cycles and complete bipartite graphs, *J. Combin. Math. Combin. Comput.*, 52(2005),69-78.
- [4] S. K. Vaidya, G. V. Ghodasara, Sweta Srivastav and V.J.Kaneria, Cordial and 3 equitable labeling of star of a cycle, *Mathematics Today*, 24(2008), 54-64.
- [5] V. J. Kaneria, H. M. Makadia, Some Graceful Graphs, *J. of Math. Research*, 4(1), (2012), 54-57.
- [6] V. J. Kaneria, H. M. Makadia, M.M.Jariya and Meera Meghpara, Graceful Labeling for Complete Bipartite graphs, *Applied Mathematical Sciences*,8(103), 2014, 5099 – 5104.
- [7] J. A. Gallian, A Dynamic Survey of Graph Labeling, *The Electronics Journal of Combinatorics*, (2017), # DS6.
- [8] F. Harary, *Graph theory* Addison Wesley, Massachusetts, 1972.
- [9] V.J.Kaneria,Meera Meghpara,H M Makadia, Graceful labeling for Open star of graphs,*International Journal of Mathematics and statistics Invention*,2(9),2014,19-23.
- [10] V.J.Kaneria,Meera Meghpara, Graceful labeling for one point union path of graphs,*International Journal of Mathematics and its Application*, 3(1),2015,49-54.
- [11] V.J.Kaneria,H.M.Makadia,Meera Meghpara, Gracefulness cycle of cycles and cycle of complete bipartite graphs, *International Journal of Mathematics Trends technology*,12(1),2014,19-26.
- [12] V.J.Kaneria,H.M.Makadia,Meera Meghpara, Graceful labeling for grid related graphs, *International Journal of Mathematics and soft computing*, 5(1),2015,111-117.
- [13] V.J.Kaneria,H.M.Makadia,Meera Meghpara,Some Graceful graphs, *International Journal of Mathematics and soft computing*, 4(2),2014,165-172.
- [14] V. J. Kaneria, H. M. Makadia, M. M. Jariya, Graceful Labeling for Cycle of Graphs, *International Journal of Mathematics Research*, 6(2), 2014, 173–178.