

# AN INVERSE PROBLEM FOR MOORE–GIBSON–THOMPSON EQUATION ARISING IN HIGH INTENSITY ULTRASOUND

RODRIGO LECAROS, ALBERTO MERCADO, AND SEBASTIÁN ZAMORANO

ABSTRACT. In this article we study the inverse problem of recovering a space-dependent coefficient of the Moore–Gibson–Thompson (MGT) equation, from knowledge of the trace of the solution on some open subset of the boundary. We obtain the Lipschitz stability for this inverse problem, and we design a convergent algorithm for the reconstruction of the unknown coefficient. The techniques used are based on Carleman inequalities for wave equations and properties of the MGT equation.

## 1. INTRODUCTION AND MAIN RESULTS

In this paper, we are interested in an inverse problem for the Moore–Gibson–Thompson (MGT) equation. It consists in recovering an unknown time-independent coefficient from a single measurement of some trace of the solution at the boundary.

Let  $\Omega \subseteq \mathbb{R}^N$  be a nonempty bounded open set (for  $N = 2$  or  $N = 3$ ), with a smooth boundary  $\Gamma$ , and let  $T > 0$ . We consider the MGT equation

$$(1) \quad \begin{cases} \tau u_{ttt} + \alpha u_{tt} - c^2 \Delta u - b \Delta u_t = f, \Omega \times (0, T) \\ u = 0, \Gamma \times (0, T) \\ u(\cdot, 0) = u_0, u_t(\cdot, 0) = u_1, u_{tt}(\cdot, 0) = u_2, \Omega, \end{cases}$$

where  $c \in \mathbb{R}$  and  $\tau, b > 0$  are given constants. We study the inverse problem of recovering the unknown space-dependent coefficient  $\alpha = \alpha(x)$  from partial knowledge of some trace of the solution  $u$  at the boundary, namely,

$$\frac{\partial u}{\partial n} \text{ on } \Gamma_0 \times (0, T),$$

where  $\Gamma_0 \subset \Gamma$  is an relatively open subset, called the observation region, and  $n$  is the outward unit normal vector on  $\Gamma$ . We will often write  $u(\alpha)$  to denote the dependence of  $u$  on the coefficient  $\alpha$ . Concerning the well posedness of the system, it is known (see Theorem 2.1) that, given a coefficient  $\alpha \in L^\infty(\Omega)$  and data satisfying

$$(2) \quad (u_0, u_1, u_2) \in H^2(\Omega) \cap H_0^1(\Omega) \times H_0^1(\Omega) \times L^2(\Omega), \quad f \in L^2(0, T; L^2(\Omega)),$$

---

*Key words and phrases.* Carleman inequalities, Bukhgeim–Klibanov method, hidden regularity, Moore–Gibson–Thompson equation.

The work of the first author is partially supported by FONDECYT (Chile) Grant NO: 1171712. R. Lecaros was partially supported by FONDECYT(Chile) Grant NO: 11180874. The first and second author was partially supported by BASAL Project, CMM - U. de Chile. The third author is supported by the FONDECYT(Chile) Postdoctoral Grant NO: 3180322.

the system (1) admits a unique weak solution  $(u, u_t, u_{tt})$  satisfying

$$(u, u_t, u_{tt}) \in C([0, T]; H^2(\Omega) \cap H_0^1(\Omega) \times H_0^1(\Omega) \times L^2(\Omega)).$$

More precisely, in this paper we study the following properties of the stated inverse problem:

- **Uniqueness:**

$$\frac{\partial u(\alpha_1)}{\partial n} = \frac{\partial u(\alpha_2)}{\partial n} \text{ on } \Gamma_0 \times (0, T) \text{ implies } \alpha_1 = \alpha_2 \text{ in } \Omega.$$

- **Stability:**

$$\|\alpha_1 - \alpha_2\|_{X(\Omega)} \leq C \left\| \frac{\partial u(\alpha_1)}{\partial n} - \frac{\partial u(\alpha_2)}{\partial n} \right\|_{Y(\Gamma_0)},$$

for some appropriate spaces  $X(\Omega)$  and  $Y(\Gamma_0)$ .

- **Reconstruction:** Design an algorithm to recover the coefficient  $\alpha$  from the knowledge of  $\frac{\partial u(\alpha)}{\partial n}$  on  $\Gamma_0$ .

The first part of this work is concerned with the uniqueness and stability issues of the inverse problem. We will obtain a stability result, which directly implies a uniqueness one, under certain conditions for  $\alpha$ ,  $\Gamma_0$  and the time  $T$ . To prove the stability, we use the Bukheim–Klibanov method, which is based on the idea of applications of the so-called Carleman estimates to prove the result for the inverse problem.

In order to state the precise result that we obtain, we consider the following set of admissible coefficients:

$$(3) \quad \mathcal{A}_M = \left\{ \alpha \in L^\infty(\Omega), \quad \frac{c^2}{b} \leq \alpha(x) \leq M \quad \forall x \in \overline{\Omega} \right\},$$

and the geometrical assumptions, sometimes referred to as the Gamma-condition of Lions or the multiplier condition::

$$(4) \quad \exists x_0 \notin \Omega \text{ such that } \Gamma_0 \supset \{x \in \Gamma : (x - x_0) \cdot n \geq 0\},$$

and

$$(5) \quad T > \sup_{x \in \Omega} |x - x_0|.$$

Our main result concerning the stability is the following.

**Theorem 1.1.** *For  $\Gamma_0 \subset \Gamma$ ,  $M > 0$  and  $T > 0$  satisfying (4)–(5), suppose there exists  $\eta > 0$  such that*

$$(6) \quad |u_2| \geq \eta > 0 \quad \text{a.e. in } \Omega,$$

and  $\alpha_2 \in \mathcal{A}_M$  is such that the unique solution  $u(\alpha_2)$  of (1) satisfies

$$(7) \quad u(\alpha_2) \in H^3(0, T; L^\infty(\Omega)).$$

Then there exists a constant  $C > 0$  such that

$$(8) \quad C^{-1} \|\alpha_1 - \alpha_2\|_{L^2(\Omega)}^2 \leq \left\| \frac{\partial u(\alpha_1)}{\partial n} - \frac{\partial u(\alpha_2)}{\partial n} \right\|_{H^2(0,T;L^2(\Gamma_0))}^2 \leq C \|\alpha_1 - \alpha_2\|_{L^2(\Omega)}^2$$

for all  $\alpha_1 \in \mathcal{A}_M$ .

Let us mention some comments about Theorem 1.1.

**Remark 1.2.** *The hypothesis  $u(\alpha_2) \in H^3(0, T; L^\infty(\Omega))$  in Theorem 1.1 is satisfied if more regularity is imposed on the data. For instance, taking  $m > \frac{N}{2} + 1$ , it is enough to take  $(u_0, u_1, u_2) \in (H^{m+2}(\Omega) \times H^{m+1}(\Omega) \times H^m(\Omega))$ ,  $f \equiv 0$ , and  $\alpha_2 \in H^{m-1}(\Omega)$ , plus appropriate boundary compatibility conditions. Indeed, from Theorem 2.1 (Section 2), we obtain*

$$u = u(\alpha_2) \in C([0, T]; H^{m+2}(\Omega)) \cap C^1([0, T]; H^{m+1}(\Omega)) \cap C^2([0, T]; H^m(\Omega)).$$

*Using the fact that the Sobolev space  $H^s(\Omega)$  with  $s > \frac{N}{2}$  is an algebra, we have that  $u_{tt}$  solves a similar equation than (1) with initial condition  $(u_{tt}(\cdot, 0), u_{ttt}(\cdot, 0), u_{tttt}(\cdot, 0)) \in (H^m(\Omega) \times H^{m-1}(\Omega) \times H^{m-2}(\Omega))$ . Using again Theorem 2.1, we deduce that  $u_{tt} \in C([0, T]; H^m(\Omega)) \cap C^1([0, T]; H^{m-1}(\Omega)) \cap C^2([0, T]; H^{m-2}(\Omega))$ . Hence*

$$u_{ttt} \in C([0, T]; H^{m-1}(\Omega)) \cap C^1([0, T]; H^{m-2}(\Omega)),$$

*and using Sobolev's embedding theorem, we get that  $u_{ttt} \in L^2(0, T; L^\infty(\Omega))$ .*

**Remark 1.3.** *The inverse problem studied in this paper was previously considered by Liu and Triggiani [19, Theorem 15.5]. They considered  $\alpha \in H^m(\Omega)$  and initial data  $(u_0, u_1, u_2) \in (H^{m+2}(\Omega) \times H^{m+1}(\Omega) \times H^m(\Omega))$  with  $m > \frac{N}{2} + 2$ . By using Carleman estimates for a general hyperbolic equation, the authors proved global uniqueness of any damping coefficient  $\alpha$  with boundary measurement given by*

$$\frac{c^2}{b} \frac{\partial u}{\partial n} + \frac{\partial u_t}{\partial n}, \quad \text{on } \Gamma_0 \times [0, T],$$

*and the initial data is supposed to satisfy (6) and*

$$(9) \quad \frac{c^2}{b} u_0(x) + u_1(x) = 0, \quad x \in \Omega.$$

*In this paper, using an appropriate Carleman inequality and the method of Bukhgeim-Klibanov, we obtain stability around any regular state, under hypothesis  $m > \frac{N}{2} + 1$  and without the additional assumption (9).*

**Remark 1.4.** *The hypotheses (4) and (5) on  $\Gamma_0$  and  $T$  typically arises in the study of stability or observability inequalities for the wave equation, see [11] where the multiplier method is used, or [10, 28] where some observability inequalities are obtained from Carleman estimates. These hypotheses provide a particular case of the geometric control condition stated in [1].*

**Remark 1.5.** *The assumption of the positiveness for  $u_2$  appearing in Theorem 1.1 is classical when applying the Bukhgeim-Klibanov method and Carleman estimates for inverse problems with only one boundary measurement, see [2, 18, 25].*

As we mentioned before, in order to study and proved the stated inverse problem, we use global Carleman estimates and the method of Bukhgeim-Klibanov, introduced in [8]. To state our Carleman estimates precisely, we shall need the following notations.

Assume that  $\Gamma_0$  satisfies (4) for some  $x_0 \in \mathbb{R}^N \setminus \overline{\Omega}$ . For  $\lambda > 0$ , we define the weight functions  $\phi$  and  $\varphi_\lambda$  as follows

$$\varphi_\lambda(x, t) = e^{\lambda\phi(x, t)},$$

where

$$\phi(x, t) = |x - x_0|^2 - \beta t^2 + M_0, \quad 0 < \beta < 1,$$

and  $M_0$  is chosen such that

$$\forall (x, t) \in \Omega \times (-T, T), \quad \phi(x, t) \geq 1.$$

To prove our main Theorem 1.1, we shall use the following Carleman estimate.

**Theorem 1.6.** *Suppose that  $\Gamma_0$  and  $T$  satisfies (4), (5). Let  $M > 0$  and  $\alpha \in \mathcal{A}_M$ . Let  $\beta \in (0, 1)$  such that*

$$(10) \quad \beta T > \sup_{x \in \Omega} \|x - x_0\|.$$

*Then, there exists  $s_0 > 0$ ,  $\lambda > 0$  and a positive constant  $C$  such that*

$$(11) \quad \begin{aligned} & \sqrt{s} \int_{\Omega} e^{2s\varphi_\lambda(\cdot, 0)} |y_{tt}(\cdot, 0)|^2 dx + s\lambda c^4 \int_0^T \int_{\Omega} e^{2s\varphi_\lambda} \varphi_\lambda (|y_t|^2 + |\nabla y|^2) dx dt \\ & \quad + s^3 \lambda^3 c^4 \int_0^T \int_{\Omega} e^{2s\varphi_\lambda} \varphi_\lambda^3 |y|^2 dx dt + s\lambda \int_0^T \int_{\Omega} e^{2s\varphi_\lambda} \varphi_\lambda (|y_{tt}|^2 + |\nabla y_t|^2) dx dt \\ & \quad + s^3 \lambda^3 \int_0^T \int_{\Omega} e^{2s\varphi_\lambda} \varphi_\lambda^3 |y_t|^2 dx dt \leq C \int_0^T \int_{\Omega} e^{2s\varphi_\lambda} |\tau y_{ttt} + \alpha y_{tt} - c^2 \Delta y - b \Delta y_t|^2 dx dt \\ & \quad \quad \quad + Cs\lambda \int_0^T \int_{\Gamma_0} e^{2s\varphi_\lambda} (|\nabla y_t \cdot n|^2 + c^4 |\nabla y \cdot n|^2) d\sigma dt, \end{aligned}$$

*for all  $s \geq s_0$  and for all  $y \in L^2(0, T; H_0^1(\Omega))$  satisfying  $Ly := y_{ttt} + \alpha y_{tt} - c^2 \Delta y - b \Delta y_t \in L^2(\Omega \times (0, T))$ ,  $y(\cdot, 0) = y_t(\cdot, 0) = 0$  in  $\Omega$ , and  $y_{tt}(\cdot, 0) \in L^2(\Omega)$ .*

Let us mention that Carleman inequalities like (11) is a very technical but powerful mathematical tool. In this work, we established this estimate with an alternative methodology. The classical way to get inequalities as (11) is to consider an operator of the form  $P_0(w) := e^{s\varphi_\lambda} L(e^{-s\varphi_\lambda} w)$ , with  $w := e^{s\varphi_\lambda} y$  and then decompose this operator into the symmetric part and the anti-symmetric part. Then, after several integrations by parts, the use of the weighted functions, and energy estimates, the Carleman inequality is obtained. However, as we mention before, instead of developing a Carleman estimate for the MGT system, we get (11) using in an appropriate way the well-known Carleman estimate for the wave operator. The most important reason for obtaining an inequality like (11), similar to a Carleman estimate, but not following the classical technique, is due that the MGT equation has poor control properties as was proved in [20]. The authors in [20] showed that

the null controllability is impossible to get in a cylindrical subset  $\omega \times (0, T) \subset \Omega \times (0, T)$ . This immediately implies that also with a boundary control the null controllability is false. Therefore, if we prove a Carleman estimate for MGT equation, this would tell us that the equation can be controllable through a boundary control, obtaining a contradiction with the result in [20].

A similar idea used to obtain an observability inequality was considered in [27], where a Carleman estimate was obtained to prove the unique continuation property for a linearized Benjamin–Bona–Mahony equation, from the use of Carleman inequality for the Laplace operator.

These Bukhgeim-Klibanov method and Carleman estimate have been widely used for obtaining stability of coefficients with one-measurement observations. Concerning inverse problems for wave equations with boundary observations, in [24] is studied the problem of recovering a source term of the equation, [26] deals with the problem of recovering a coefficient of the zero-order term, and [6] concerns the recovering of the main coefficient. A complete list of references using this method is too long to be given here, but we mention [7] for a complete account of classic and recent results.

The second part of this work is focused on giving a constructive and iterative algorithm which allow us to find the coefficient  $\alpha$  from the knowledge of the additional data  $\frac{\partial u}{\partial n}$  on the observation domain  $\Gamma_0$ . For that, we will analysis the following functional

$$(12) \quad J[\mu, g](y) = \frac{1}{2s} \int_0^T \int_{\Omega} q^2(s, x, t) |\tau y_{ttt} + \alpha y_{tt} - c^2 \Delta y - b \Delta y_t - g|^2 dx dt \\ + \frac{1}{2} \int_0^T \int_{\Gamma_0} q^2(s, x, t) \left( \left| \frac{\partial y}{\partial n} - \mu \right|^2 + \left| \frac{\partial y_t}{\partial n} - \mu_t \right|^2 \right) d\sigma dt,$$

where  $\alpha \in \mathcal{A}_M$ ,  $g \in L^2(\Omega \times (0, T))$ ,  $\mu \in H^1(0, T; L^2(\Gamma_0))$ , and  $q$  is an appropriate exponential function depending on the parameter  $s > 0$ , which is given explicitly in the proof of Theorem 1.1, and on the trajectories  $y \in L^2(0, T; H_0^1(\Omega))$ , with  $Ly := y_{ttt} + \alpha y_{tt} - c^2 \Delta y - b \Delta y_t \in L^2(\Omega \times (0, T))$ ,  $y(\cdot, 0) = y_t(\cdot, 0) = 0$  in  $\Omega$ , and  $y_{tt}(\cdot, 0) \in L^2(\Omega)$ .

We will show that this functional admits a unique minimizer (see Theorem 4.2) on an appropriate admissible set. Moreover, we will show how the minimum of (12) depends on  $g \in L^2(\Omega \times (0, T))$  (see Theorem 4.3). We refer to Section 4 for details. Using this results, we prove the convergence of the iterative algorithm presented below, which is adapted from [3], where it was introduced an algorithm for recovering zero-order terms in the wave equation. We can also mention the works of Beilina and Klibanov [4, 5], where the authors studied the reconstruction of a coefficient in a hyperbolic equation using the Carleman weight.

---

**Algorithm:**

---

- (1) **Initialization:**  $\alpha^0 = \frac{c^2}{b}$ .
- (2) **Iteration:** From  $k$  to  $k + 1$

Step 1 - Given  $\alpha^k$  we consider  $\mu^k = \partial_t \left( \frac{\partial u(\alpha^k)}{\partial n} - \frac{\partial u(\alpha)}{\partial n} \right)$  and  $\mu_t^k = \partial_t \left( \frac{\partial u_t(\alpha^k)}{\partial n} - \frac{\partial u_t(\alpha)}{\partial n} \right)$  on  $\Gamma_0 \times (0, T)$ , where  $u(\alpha^k)$  and  $u(\alpha)$  are the solution of the problems

$$(13) \quad \begin{cases} \tau u_{ttt} + \alpha^k u_{tt} - c^2 \Delta u - b \Delta u_t = f, \Omega \times (0, T) \\ u = g, \Gamma \times (0, T) \\ u(\cdot, 0) = u_0, u_t(\cdot, 0) = u_1, u_{tt}(\cdot, 0) = u_2, \Omega \end{cases}$$

and

$$(14) \quad \begin{cases} \tau u_{ttt} + \alpha u_{tt} - c^2 \Delta u - b \Delta u_t = f, \Omega \times (0, T) \\ u = g, \Gamma \times (0, T) \\ u(\cdot, 0) = u_0, u_t(\cdot, 0) = u_1, u_{tt}(\cdot, 0) = u_2, \Omega. \end{cases}$$

Step 2 - Minimize the functional  $J[\mu^k, 0]$  on the admissible trajectories  $y$  (see Section 4 for details).

Step 3 - Let  $y^{*,k}$  the minimizer of  $J[\mu^k, 0]$  and

$$(15) \quad \tilde{\alpha}^{k+1} = \alpha^k + \frac{y_{tt}^{*,k}(\cdot, 0)}{u_2}.$$

Step 4 - Finally, consider  $\alpha^{k+1} = T(\tilde{\alpha}^{k+1})$ , where

$$(16) \quad T(\alpha) = \begin{cases} M & \text{if } \alpha > M \\ \alpha & \text{if } \frac{c^2}{b} \leq \alpha \leq M \\ \frac{c^2}{b} & \text{if } \alpha < \frac{c^2}{b}. \end{cases}$$

This function  $T$  is to guarantee at each step that  $\alpha^k$  belongs to the admissible set  $\mathcal{A}_M$ .

Let us finally conclude this section giving some references about the MGT equation considering in this work. The MGT equation

$$\tau u_{ttt} + \alpha(x) u_{tt} - c^2 \Delta u - b \Delta u_t = 0,$$

arises, e.g., as a linearization of a model for wave propagation in viscous thermally relaxing fluids. In that cases, the space-dependent coefficient  $\alpha$  depended on a viscosity of the fluid [14]. This third order in time equation (linear and nonlinear) was previously analysed by several authors from a different point of view. We can mentioned, among others, the works [13, 15, 16, 19, 21, 22, 20] for a variety of problems related to this equation.

In particular, one interesting and different characteristics of this equation from a familiar second order systems ( $\tau = 0, \alpha > 0$ ), is that the structural damping  $b$  plays an crucial role for the well-posedness: in [15] it is proved that, if  $b = 0$  and  $\alpha$  a positive constant, there does not exist an infinitesimal generator of the semigroup, in contrast with second order

equations, where the structural damping does not affect the well-posedness of the equation. The parameter  $\gamma := \alpha - \frac{\tau c^2}{b}$  gives relevant information regarding the stability of the system. If  $\gamma > 0$ , the group associated to the equation is exponentially stable, and for  $\gamma = 0$ , the group is conservative, see for instance [21]. On the other hand, Conejero, Lizama and Rodenas [9] proved that the one-dimensional equation exhibits a chaotic behavior if  $\gamma < 0$ . Also, for the case in which  $\alpha$  is given by a function depending on space and time, the well posedness and the exponential decay was proved by Kaltenbacher and Lasiecka in [14].

The remaining of this paper is organized as follows. In section 2 we present some basic results about the well posedness of the MGT equation which are needed for the inverse problem. Besides, we prove the hidden regularity for the MGT equation. Also, we present Carleman estimates for the wave equation. In section 3 we prove the main result of our work, namely Theorems 1.1 and 1.6. Finally, in section 4 we focus on the algorithm for the reconstruction of coefficient  $\alpha$  and we prove the convergence of this Algorithm.

## 2. PRELIMINARIES

In this section we present the well posedness framework for studying our problem. Furthermore, we state and prove some estimates for the Laplacian of a solution of (1) and a hidden regularity estimate for the solution of the MGT equation. Also, we recall the results concerning to the Carleman estimates for the wave equation. This estimation play an important role in our approach.

**2.1. Well-posedness for Moore-Gibson-Thompson equation.** Let us now introduce the basic properties for the equation under studying. We first review some results given by Kaltenbacher and Lasiecka [14], in the context of semigroup theory. Let  $\Omega$  be a bounded and smooth domain of  $\mathbb{R}^N$ ,  $N = 2$  or  $3$ , and consider the equation

$$(17) \quad \begin{cases} \tau u_{ttt} + \alpha u_{tt} - c^2 \Delta u - b \Delta u_t = f, \Omega \times (0, T) \\ u = 0, \Gamma \times (0, T) \\ u(\cdot, 0) = u_0, u_t(\cdot, 0) = u_1, u_{tt}(\cdot, 0) = u_2, \Omega \end{cases}$$

where  $\alpha = \alpha(x)$  is a variable coefficient depending on a viscosity of the fluid,  $c$  is the speed of the sound,  $\tau$  is a positive constant accounting for relaxation,  $b = \delta + \tau c^2$  where  $\delta$  is the diffusivity of sound. Without loss of generality, we normalize  $\tau$  and set  $\tau = 1$ . We denote by  $\Gamma$  the boundary of  $\Omega$ , and  $T$  is a strictly positive number. We recall that  $n$  is the outward unit normal vector on  $\Gamma$ .

The homogeneous part of equation (17) can be rewritten as a first order abstract system

$$\begin{cases} U_t(t) = AU(t), & t > 0, \\ U(0) = U_0 = (u_0, u_1, u_2), \end{cases}$$

where the operator  $A$  is defined by

$$(18) \quad A = \begin{pmatrix} 0 & I & 0 \\ 0 & 0 & I \\ c^2 \Delta & b \Delta & -\alpha I \end{pmatrix}$$

The natural domain of  $A$  is given by  $D(A) = (H^2(\Omega) \cap H_0^1(\Omega)) \times (H^2(\Omega) \cap H_0^1(\Omega)) \times H_0^1(\Omega)$ .

Let us consider

$$(19) \quad \alpha(x) \in \mathcal{A}_M$$

for some  $M > 0$ .

**Theorem 2.1** ([14], Theorem 2.2). *Let  $b > 0$  and  $\alpha \in L^\infty(\Omega)$ . Then the operator  $A$  defined by (18) generates a strongly continuous group on the state space*

$$H = (H^2(\Omega) \cap H_0^1(\Omega)) \times H_0^1(\Omega) \times L^2(\Omega).$$

*That is, for each  $(u_0, u_1, u_2) \in H$  and  $f \in L^2(0, T; L^2(\Omega))$ , there exists a unique solution  $U = (u, u_t, u_{tt}) \in C([0, T]; H)$  of (17).*

**2.2. Bound of Laplacian of the solutions.** From now, throughout the article, we define

$$(20) \quad \gamma(x) := \alpha(x) - \frac{c^2}{b},$$

and then the set of admissible coefficients is given by

$$(21) \quad \mathcal{B}_M := \{\gamma \in L^\infty(\Omega), \quad 0 \leq \gamma(x) \leq M, \quad \forall x \in \overline{\Omega}\}.$$

We also define the energy

$$(22) \quad E_e(y) := \frac{b}{2} \|\nabla y\|_{L^2(\Omega)}^2 + \frac{1}{2} \|y_t\|_{L^2(\Omega)}^2.$$

In order to prove our main results, some technical estimations are necessary. One of them is the following:

**Theorem 2.2.** *Let  $b > 0$  and  $M > 0$  such that  $\alpha \in \mathcal{A}_M$ . Then there exists  $C > 0$  such that the total energy*

$$(23) \quad \overline{E}(t) := E_e(u_t(t)) + E_e(u(t)),$$

*satisfies*

$$\overline{E}(t) \leq C \left( \overline{E}(0) + \|f\|_{L^2(0, T; L^2(\Omega))}^2 \right), \quad t \in [0, T],$$

*for every  $(u_0, u_1, u_2) \in (H^2(\Omega) \cap H_0^1(\Omega)) \times H_0^1(\Omega) \times L^2(\Omega)$  and  $f \in L^2(0, T; L^2(\Omega))$ , where  $u$  be the unique solution of (17).*

*Proof.* Without loss of generality, we assume that  $b = 1$ . Then, the equation

$$u_{ttt} + \alpha(x)u_{tt} - c^2\Delta u - b\Delta u_t = f$$

can be write as follows (recall the definition of  $\gamma$  in (20))

$$(24) \quad Lu := L_0 u_t + c^2 L_0 u + \gamma(x) u_{tt} = f,$$

where the operator  $L_0$  is the wave operator given by

$$(25) \quad L_0 := \partial_t^2 - \Delta.$$

Let us multiply the equation (24) by  $u_{tt}(t) + c^2 u_t(t) \in L^2(\Omega)$  and after integrating on  $\Omega$ , we deduce that



$$\frac{d}{dt}E_e(u_t + c^2u) + \int_{\Omega} \gamma u_{tt}(u_{tt} + c^2u_t) = \int_{\Omega} f(u_{tt} + c^2u_t),$$

thus, we have

$$\frac{d}{dt}E_e(u_t + c^2u) + \frac{c^2}{2} \frac{d}{dt} \|\gamma^{1/2}u_t\|_{L^2(\Omega)}^2 \leq \frac{1}{2} \|f\|_{L^2(\Omega)}^2 + E_e(u_t + c^2u).$$

And using Gronwall's inequality, there exists a constant  $C > 0$ , such that

$$(26) \quad E_e(u_t + c^2u) + \frac{c^2}{2} \|\gamma^{1/2}u_t\|_{L^2(\Omega)}^2 \leq C(\|f\|_{L^2(0,T;L^2(\Omega))}^2 + \overline{E}(0)), \quad \forall t \in [0, T].$$

On other side, a directly computation give us

$$E_e(u_t + c^2u) = E_e(u_t) + c^4E_e(u) + 2c^2 \frac{d}{dt}E_e(u),$$

and replacing the above identities in (26), we have

$$2c^2 \frac{d}{dt}E_e(u) \leq C(\|f\|_{L^2(0,T;L^2(\Omega))}^2 + \overline{E}(0)).$$

Hence, integrating we obtain that, there exists a constant  $C > 0$ , such that

$$(27) \quad E_e(u) \leq C(\|f\|_{L^2(0,T;L^2(\Omega))}^2 + \overline{E}(0)), \quad \forall t \in [0, T].$$

Finally, if we take  $\varepsilon < 1$ , we observe that

$$2c^2 \frac{d}{dt}E_e(u) = c^2 \int_{\Omega} (u_{tt}u_t + \nabla u \cdot \nabla u_t) \leq \varepsilon^2 E_e(u_t) + \frac{c^4}{\varepsilon^2} E_e(u),$$

replacing the above identities in (26) and using (27), we obtain that, there exists a constant  $C > 0$ , such that,

$$(28) \quad E_e(u_t) \leq C(\|f\|_{L^2(0,T;L^2(\Omega))}^2 + \overline{E}(0)), \quad \forall t \in [0, T].$$

combined (27) and (28) we conclude the proof.  $\square$

**Theorem 2.3.** *Let  $b > 0$  and  $M > 0$  such that  $\alpha \in \mathcal{A}_M$ . Let  $(u, u_t, u_{tt})$  be the unique solution of (17) with data  $(u_0, u_1, u_2) \in (H^2(\Omega) \cap H_0^1(\Omega)) \times H_0^1(\Omega) \times L^2(\Omega)$  and  $f \in L^2(0, T; L^2(\Omega))$ . Then, the term  $\Delta u(t)$  can be bounded as follows*

$$\|\Delta u(t)\|_{L^2(\Omega)}^2 \leq C \left( \|f\|_{L^2(0,T;L^2(\Omega))}^2 + \overline{E}(0) + \|\Delta u_0\|_{L^2(\Omega)}^2 \right), \quad \forall t \in [0, T].$$

*Proof.* Since the term  $u_{tt}(t), \Delta u(t) \in L^2(\Omega)$ , let us multiply the equation (24) by  $L_0u$  and after integrating on  $\Omega$ , we deduce that

$$(29) \quad \frac{d}{dt} \|L_0(u(t))\|_{L^2(\Omega)}^2 + 2c^2 \|L_0(u(t))\|_{L^2(\Omega)}^2 = 2 \langle f(t) - \gamma u_{tt}(t), L_0(u(t)) \rangle_{L^2(\Omega)}.$$

Using Hölder inequality, the right hand side of (29) is majorized by

$$2\langle f(t) - \gamma u_{tt}(t), L_0(u(t)) \rangle_{L^2(\Omega)} \leq 2\|f(t) - \gamma u_{tt}(t)\|_{L^2(\Omega)} \|L_0(u(t))\|_{L^2(\Omega)}.$$

Now, for every  $\varepsilon > 0$

$$(30) \quad 2\|f(t) - \gamma u_{tt}(t)\|_{L^2(\Omega)} \|L_0(u(t))\|_{L^2(\Omega)} \leq \frac{1}{\varepsilon} \|f(t) - \gamma u_{tt}(t)\|_{L^2(\Omega)}^2 + \varepsilon \|L_0(u(t))\|_{L^2(\Omega)}^2.$$

Therefore, taking  $\varepsilon = c^2 > 0$  in (30) and replacing it in (29) we obtain

$$\begin{aligned} \frac{d}{dt} \|L_0(u(t))\|_{L^2(\Omega)}^2 + 2c^2 \|L_0(u(t))\|_{L^2(\Omega)}^2 - c^2 \|L_0(u(t))\|_{L^2(\Omega)}^2 \\ \leq \frac{1}{c^2} \|f(t) - \gamma u_{tt}(t)\|_{L^2(\Omega)}^2, \end{aligned}$$

which implies

$$(31) \quad \frac{d}{dt} \|L_0(u(t))\|_{L^2(\Omega)}^2 \leq \frac{1}{c^2} \|f(t) - \gamma u_{tt}(t)\|_{L^2(\Omega)}^2.$$

Integrating (31) from 0 to  $t > 0$ , we obtain that

$$\|L_0(u(t))\|_{L^2(\Omega)}^2 \Big|_0^t \leq \frac{1}{c^2} \int_0^t \|f(\tau) - \gamma u_{tt}(\tau)\|_{L^2(\Omega)}^2 d\tau.$$

Then, we have

$$\|L_0(u(t))\|_{L^2(\Omega)}^2 - \|L_0(u(t))\|_{L^2(\Omega)}^2 \Big|_{t=0} \leq \frac{2}{c^2} \int_0^t \|f(\tau)\|_{L^2(\Omega)}^2 d\tau + \frac{2}{c^2} \int_0^t \|\gamma u_{tt}(\tau)\|_{L^2(\Omega)}^2 d\tau,$$

and then using Theorem 2.2 we obtain that, there exists a constant  $C > 0$ , such that

$$\|\Delta u(t)\|_{L^2(\Omega)}^2 \leq C \left( \|f\|_{L^2(0,T;L^2(\Omega))}^2 + \overline{E}(0) + \|\Delta u_0\|_{L^2(\Omega)}^2 \right), \quad \forall t \in [0, T].$$

□

**2.3. Hidden regularity.** We can observe that the inverse problem considered in this paper needs that the normal derivative of the solution can be defined on the boundary. It is well know that the wave equation satisfies certain extra regularity called *hidden regularity* [17]. it is natural to expect an analogous result for the Moore–Gibson–Thompson equation, due its hyperbolic nature [15]. In the following result, using the multiplier method, we obtain a hidden regularity for the solutions of this equation.

**Theorem 2.4.** *The unique solution  $(u, u_t, u_{tt}) \in C([0, T]; (H^2(\Omega) \cap H_0^1(\Omega)) \times H_0^1(\Omega) \times L^2(\Omega))$  of (17) satisfies*

$$(32) \quad \frac{\partial u}{\partial n} \in H^1(0, T; L^2(\Gamma)).$$

Moreover, the normal derivative satisfies

$$(33) \quad \left\| \frac{\partial u}{\partial n} \right\|_{H^1(0,T;L^2(\Gamma))}^2 \leq C (\|u_0\|_{H^2(\Omega) \cap H_0^1(\Omega)}^2 + \|u_1\|_{H_0^1(\Omega)}^2 + \|u_2\|_{L^2(\Omega)}^2 + \|f\|_{L^2(0,T;L^2(\Omega))}^2).$$

Consequently, the mapping

$$(f, u_0, u_1, u_2) \mapsto \frac{\partial u}{\partial n}$$

is linear continuous from  $L^2(0, T; L^2(\Omega)) \times (H^2(\Omega) \cap H_0^1(\Omega)) \times H_0^1(\Omega) \times L^2(\Omega)$  into  $H^1(0, T; L^2(\Gamma))$ .

*Proof.* We use the multiplier method for the proof. Let  $m \in W^{1,\infty}(\Omega; \mathbb{R}^N)$  and let us multiply  $L_0(u)$  by  $m \nabla u$  and  $L_0(u_t)$  by  $m \nabla u_t$ . Using the summation convention for repeated index, we obtain, respectively

$$(34) \quad \int_0^T \int_{\Omega} L_0(u_t) m \nabla u_t dx dt = \frac{1}{2} \int_0^T \int_{\Omega} \operatorname{div}(m) |u_{tt}|^2 dx dt + \int_{\Omega} u_{tt} m \nabla u_t \Big|_0^T dx \\ + \int_0^T \int_{\Omega} \frac{\partial u_t}{\partial x_i} \frac{\partial m_j}{\partial x_i} \frac{\partial u_t}{\partial x_j} dx dt - \frac{1}{2} \int_0^T \int_{\Omega} \operatorname{div}(m) |\nabla u_t|^2 dx dt - \frac{1}{2} \int_0^T \int_{\partial \Omega} |\nabla u_t \cdot n|^2 (m \cdot n) d\sigma dt.$$

and

$$(35) \quad \int_0^T \int_{\Omega} L_0(u) m \nabla u dx dt = \frac{1}{2} \int_0^T \int_{\Omega} \operatorname{div}(m) |u_t|^2 dx dt + \int_{\Omega} u_t m \nabla u \Big|_0^T dx \\ + \int_0^T \int_{\Omega} \frac{\partial u}{\partial x_i} \frac{\partial m_j}{\partial x_i} \frac{\partial u}{\partial x_j} dx dt - \frac{1}{2} \int_0^T \int_{\Omega} \operatorname{div}(m) |\nabla u|^2 dx dt - \frac{1}{2} \int_0^T \int_{\partial \Omega} |\nabla u \cdot n|^2 (m \cdot n) d\sigma dt.$$

Now, taking the multiplier  $m$  as a lifting of the outward unit normal  $n$ , so that  $m \cdot n = 1$ , on  $\Gamma$  and using that  $(u, u_t, u_{tt}) \in C([0, T]; (H^2(\Omega) \cap H_0^1(\Omega)) \times H_0^1(\Omega) \times L^2(\Omega))$  we obtain

$$(36) \quad \int_0^T \int_{\partial \Omega} |\nabla u \cdot n|^2 d\sigma dt + \int_0^T \int_{\partial \Omega} |\nabla u_t \cdot n|^2 d\sigma dt \\ = - \int_0^T \int_{\Omega} (f - c^2 L_0(u) - \gamma u_{tt}) m \nabla u_t dx dt + \frac{1}{2} \int_0^T \int_{\Omega} \operatorname{div}(m) |u_{tt}|^2 dx dt \\ + \int_{\Omega} u_{tt} m \nabla u_t \Big|_0^T dx + \int_0^T \int_{\Omega} \frac{\partial u_t}{\partial x_i} \frac{\partial m_j}{\partial x_i} \frac{\partial u_t}{\partial x_j} dx dt - \frac{1}{2} \int_0^T \int_{\Omega} \operatorname{div}(m) |\nabla u_t|^2 dx dt \\ - \int_0^T \int_{\Omega} L_0(u) m \nabla u dx dt + \frac{1}{2} \int_0^T \int_{\Omega} \operatorname{div}(m) |u_t|^2 dx dt + \int_{\Omega} u_t m \nabla u \Big|_0^T dx \\ + \int_0^T \int_{\Omega} \frac{\partial u}{\partial x_i} \frac{\partial m_j}{\partial x_i} \frac{\partial u}{\partial x_j} dx dt - \frac{1}{2} \int_0^T \int_{\Omega} \operatorname{div}(m) |\nabla u|^2 dx dt \\ \leq C \left( \|f\|_{L^2(0,T;L^2(\Omega))}^2 + \overline{E}(0) + \|\Delta u_0\|_{L^2(\Omega)}^2 \right)$$

Concerning the estimate (33) is direct from the previous computation, Theorem 2.3 and the continuous dependence of the solution with respect to the data. The proof is finished.  $\square$

**2.4. Carleman Inequalities for wave equation.** Now, we provide global Carleman estimates corresponding to the wave operator. Here, we rely on the work of Puel [23], Imanuvilov [12], and also the work of Baudouin et al. [3].

Consider the wave equation

$$(37) \quad \begin{cases} y_{tt} - \Delta y = f, \Omega \times (0, T) \\ y = g, \Gamma \times (0, T) \\ y(\cdot, 0) = y_0, y_t(\cdot, 0) = y_1, \Omega. \end{cases}$$

We know a measure of  $\frac{\partial y}{\partial n}$  on a subset of the boundary, namely

$$\frac{\partial y}{\partial n} \text{ on } \Gamma_0 \times (0, T).$$

Consider the wave operator  $L_0$  defined by

$$L_0 = \partial_t^2 - \Delta$$

and  $v \in L^2(-T, T; L^2(\Omega))$  such that

$$(38) \quad L_0 v \in L^2(-T, T; L^2(\Omega)), \quad v = 0 \text{ on } \Gamma \times (-T, T),$$

$$(39) \quad v(-T) = v(T) = v_t(-T) = v_t(T) = v_{tt}(-T) = v_{tt}(T) = 0 \text{ in } \Omega.$$

Next, for  $x_0 \in \mathbb{R}^N \setminus \overline{\Omega}$  and  $\lambda > 0$ , we define the weight functions  $\phi$  and  $\varphi_\lambda$  as follows

$$(40) \quad \varphi_\lambda(x, t) = e^{\lambda\phi(x, t)},$$

where

$$(41) \quad \phi(x, t) = |x - x_0|^2 - \beta t^2 + M_0, \quad 0 < \beta < 1,$$

and  $M_0$  is chosen such that

$$(42) \quad \forall (x, t) \in \Omega \times (-T, T), \quad \phi(x, t) \geq 1.$$

For a parameter  $s > 0$ , let us consider  $P_0 w := e^{s\varphi_\lambda} L_0(e^{-s\varphi_\lambda} w)$ , with  $w := e^{s\varphi_\lambda} v$  and the decomposition

$$P_0 w = P_1 w + P_2 w + R_0 w,$$

where

$$P_1 w := w_{tt} - \Delta w + s^2 \lambda^2 \varphi_\lambda^2 (\phi_t^2 - |\nabla \phi|^2) w,$$

$$P_2 w := (M_1 - 1) s \lambda \varphi_\lambda (\phi_{tt} - \Delta \phi) w - s \lambda^2 \varphi_\lambda (\phi_t^2 - |\nabla \phi|^2) w - 2s \lambda \varphi_\lambda (\phi_t w_t - \nabla \phi \nabla w),$$

where  $M_1 \in (0, 1)$  is appropriately chosen and

$$R_0 w := -M_1 s \lambda \varphi_\lambda (\phi_{tt} - \Delta \phi) w.$$

Finally, for  $x_0 \in \mathbb{R}^N$ , we define

$$(43) \quad \Gamma_{x_0} := \{x \in \Gamma, n(x) \cdot (x - x_0) > 0\}.$$

**Theorem 2.5** ([23], Theorem 2.0.1). *Let us suppose that there exists  $x_0 \in \mathbb{R}^N \setminus \overline{\Omega}$  such that  $\Gamma_0 \supset \Gamma_{x_0}$ , and consider  $v \in L^2(-T, T; L^2(\Omega))$  such that fulfills conditions (38) and (39). Then there exists  $\lambda_0 > 0$  and  $s_0 > 0$  and there exists a constant  $C = C(s_0, \lambda_0, \Omega, \beta, x_0)$ , we have, for every  $\lambda \geq \lambda_0$  and  $s \geq s_0$ ,*

$$(44) \quad s\lambda \int_{-T}^T \int_{\Omega} e^{2s\varphi_\lambda} \varphi_\lambda (|v_t|^2 + |\nabla v|^2) dxdt + s^3 \lambda^3 \int_{-T}^T \int_{\Omega} e^{2s\varphi_\lambda} \varphi_\lambda^3 |v|^2 dxdt \\ + \int_{-T}^T \int_{\Omega} (|P_1 w|^2 + |P_2 w|^2) dxdt \leq C \int_{-T}^T \int_{\Omega} e^{2s\varphi_\lambda} |L_0 v|^2 dxdt + C s \lambda \int_{-T}^T \int_{\Gamma_0} e^{2s\varphi_\lambda} \left| \frac{\partial v}{\partial n} \right|^2 d\sigma dt.$$

The following result is a different way to established Carleman estimates for wave equation, which is easier to use in the context of inverse problems.

**Theorem 2.6** ([3], Theorem 2.10). *Let us suppose that there exists  $x_0 \in \mathbb{R}^N \setminus \overline{\Omega}$  such that  $\Gamma_0 \supset \Gamma_{x_0}$  and the following time condition*

$$(45) \quad T > \sup_{x \in \Omega} \|x - x_0\|.$$

*Besides, define the weight functions  $\phi$  as in (41) with  $\beta \in (0, 1)$  such that*

$$(46) \quad \beta T > \sup_{x \in \Omega} \|x - x_0\|.$$

*Then, there exists  $s_0 > 0$ ,  $\lambda > 0$  and a positive constant  $C > 0$ , such that for all  $s \geq s_0$*

$$(47) \quad \sqrt{s} \int_{\Omega} e^{2s\varphi_\lambda(\cdot, 0)} |v_t(\cdot, 0)|^2 dx + s\lambda \int_{-T}^T \int_{\Omega} e^{2s\varphi_\lambda} \varphi_\lambda (|v_t|^2 + |\nabla v|^2) dxdt \\ + s^3 \lambda^3 \int_{-T}^T \int_{\Omega} e^{2s\varphi_\lambda} \varphi_\lambda^3 |v|^2 dxdt \leq C \int_{-T}^T \int_{\Omega} e^{2s\varphi_\lambda} |L_0 v|^2 dxdt + C s \lambda \int_{-T}^T \int_{\Gamma_0} e^{2s\varphi_\lambda} \left| \frac{\partial v}{\partial n} \right|^2 d\sigma dt,$$

*for all  $v \in L^2(-T, T; H_0^1(\Omega))$  satisfying  $(v_{tt} - \Delta v) \in L^2(\Omega \times (-T, T))$ ,  $(\nabla v \cdot n) \in L^2(\Gamma_0 \times (-T, T))$  and  $v(\cdot, 0) = 0$  in  $\Omega$ .*

**Remark 2.7.** *We observe that when the time  $T$  is large enough (see (45)), the conditions on the function  $v$  at time  $\pm T$  can be removed of the hypotheses of Theorem 2.5. However, it is necessary to add the null assumption on the initial condition.*

### 3. PROOF OF MAIN RESULTS

In this section we prove our main results, that is, Theorem 1.1 and Theorem 1.6. First, we obtain the Carleman estimate given in Theorem 1.6 and then we apply this inequality to solve our inverse problem.

We use the following notation for the energy of the wave operator  $L_0$

$$(48) \quad W(y) := s\lambda \int_0^T \int_{\Omega} e^{2s\varphi_\lambda} \varphi_\lambda (|y_t|^2 + |\nabla y|^2) dxdt + s^3 \lambda^3 \int_0^T \int_{\Omega} e^{2s\varphi_\lambda} \varphi_\lambda^3 |y|^2 dxdt,$$

with  $\varphi_\lambda$  is given by (40). And also we recall below the operator  $L$  defined in Section 2

$$Ly := L_0 y_t + c^2 L_0 y + \gamma y_{tt}.$$



and, taking into account that  $v = v_t = 0$  at  $t = T$  and  $t = -T$ , we have

$$\begin{aligned} \int_{-T}^T \int_{\Omega} \widehat{c}^2 e^{s\varphi_\lambda} \partial_t |L_0 v|^2 dx dt &= \int_0^T \int_{\Omega} c^2 e^{s\varphi_\lambda} \partial_t |L_0 v|^2 dx dt - \int_{-T}^0 \int_{\Omega} c^2 e^{s\varphi_\lambda} \partial_t |L_0 v|^2 dx dt \\ &= -2c^2 \int_{\Omega} |L_0 v(\cdot, 0)|^2 e^{s\varphi_\lambda(\cdot, 0)} dx - \int_0^T \int_{\Omega} s c^2 |L_0 v|^2 (\partial_t \varphi_\lambda) e^{s\varphi_\lambda} dx dt \\ &\quad + \int_{-T}^0 \int_{\Omega} s c^2 |L_0 v|^2 (\partial_t \varphi_\lambda) e^{s\varphi_\lambda} dx dt. \end{aligned}$$

Also, from the definition of the weight function, we have

$$\begin{cases} \partial_t \varphi_\lambda < 0, & t \in (0, T), \\ \partial_t \varphi_\lambda > 0, & t \in (-T, 0), \end{cases}$$

and then

$$(53) \quad \int_{-T}^T \int_{\Omega} \widehat{c}^2 e^{s\varphi_\lambda} \partial_t |L_0 v|^2 dx dt \geq -2c^2 \int_{\Omega} |L_0 v(\cdot, 0)|^2 e^{s\varphi_\lambda(\cdot, 0)} dx.$$

From (52) and (53), using that  $v(\cdot, 0) = 0$ , we deduce that

$$(54) \quad \|L_0 v_t\|_w^2 + c^4 \|L_0 v\|_w^2 - 2c^2 \int_{\Omega} |y_2(x)|^2 e^{s\varphi_\lambda(\cdot, 0)} dx \leq \|Pv\|_w^2 + \|\widehat{\gamma} v_{tt}\|_w^2.$$

On the other hand, from (10), we can chose  $\delta > 0$  small enough such that

$$\phi(x, t) < M_0 < \phi(x, 0)$$

for all  $x \in \Omega$  and  $t \in I_\delta := [-T, -T + \delta] \cup [T - \delta, T]$ . Hence, taking into account that  $\phi(x, t) \leq \phi(x, 0)$  for all  $x \in \Omega$  and that the derivatives of  $\theta$  vanish on  $[-T + \delta, T - \delta]$ , we get

$$(55) \quad \begin{aligned} \int_{-T}^T \int_{\Omega} e^{2s\varphi_\lambda} |Pv|^2 dx dt &\leq C \int_0^T \int_{\Omega} e^{2s\varphi_\lambda(\cdot, 0)} |f|^2 dx dt + C e^{2se^{\lambda M_0}} \int_{T-\delta}^{\delta} \overline{E}(t) dt \\ &\quad + C e^{2se^{\lambda M_0}} \int_{T-\delta}^{\delta} \int_{\Omega} |\Delta y|^2 dx dt, \end{aligned}$$

where  $\overline{E}$  is the energy defined in (23) of  $y$ . Using Theorems 2.2 and 2.3 we get that

$$(56) \quad \|Pv\|_w^2 \leq C \int_0^T \int_{\Omega} e^{2s\varphi_\lambda(\cdot, 0)} |f|^2 dx dt + C \int_{\Omega} e^{2s\varphi_\lambda(\cdot, 0)} |y_2|^2 dx dt,$$

which together with (54) gives

$$(57) \quad \|L_0 v_t\|_w^2 + c^4 \|L_0 v\|_w^2 \leq C \int_0^T \int_{\Omega} e^{2s\varphi_\lambda(\cdot, 0)} |f|^2 dx dt + C \int_{\Omega} e^{2s\varphi_\lambda(\cdot, 0)} |y_2|^2 dx + \|\widehat{\gamma} v_{tt}\|_w^2.$$

Since  $\widehat{\gamma} \in L^\infty(\Omega \times (-T, T))$ , from (57) we obtain that  $L_0 v$  and  $L_0 v_t$  belongs to  $L^2(\Omega \times (-T, T))$ . Therefore, using the hidden regularity for the wave equation, we have that

$\frac{\partial v}{\partial n} \in H^1(-T, T; L^2(\Gamma_0))$ . Then, we can apply the Carleman estimates (47) for the wave equation to each term  $L_0 v$  and  $L_0 v_t$ . Namely, we have

$$(58) \quad s\lambda \int_{-T}^T \int_{\Omega} e^{2s\varphi_\lambda} \varphi_\lambda (|v_t|^2 + |\nabla v|^2) dx dt + s^3 \lambda^3 \int_{-T}^T \int_{\Omega} e^{2s\varphi_\lambda} \varphi_\lambda^3 |v|^2 dx dt \\ \leq C \int_{-T}^T \int_{\Omega} e^{2s\varphi_\lambda} |L_0 v|^2 dx dt + Cs\lambda \int_{-T}^T \int_{\Gamma_0} e^{2s\varphi_\lambda} \left| \frac{\partial v}{\partial n} \right|^2 d\sigma dt,$$

where we use the fact that  $v_t(\cdot, 0) = 0$ , and

$$(59) \quad \sqrt{s} \int_{\Omega} e^{2s\varphi_\lambda(\cdot, 0)} |y_2|^2 dx + s\lambda \int_{-T}^T \int_{\Omega} e^{2s\varphi_\lambda} \varphi_\lambda (|v_{tt}|^2 + |\nabla v_t|^2) dx dt \\ + s^3 \lambda^3 \int_{-T}^T \int_{\Omega} e^{2s\varphi_\lambda} \varphi_\lambda^3 |v_t|^2 dx dt \leq C \int_{-T}^T \int_{\Omega} e^{2s\varphi_\lambda} |L_0 v_t|^2 dx dt + Cs\lambda \int_{-T}^T \int_{\Gamma_0} e^{2s\varphi_\lambda} \left| \frac{\partial v_t}{\partial n} \right|^2 d\sigma dt.$$

Thus, from (57), (58) and (59) we obtain

$$(60) \quad \sqrt{s} \int_{\Omega} e^{2s\varphi_\lambda(\cdot, 0)} |y_2|^2 dx + c^4 W(v) + W(v_t) \leq C \int_0^T \int_{\Omega} e^{2s\varphi_\lambda(\cdot, 0)} |f|^2 dx dt \\ + \|\widehat{\gamma} v_{tt}\|_w^2 + C \int_{\Omega} |v_{tt}(\cdot, 0)|^2 e^{s\varphi_\lambda(\cdot, 0)} dx + s\lambda C \int_{-T}^T \int_{\Gamma_0} e^{2s\varphi_\lambda} \left( \left| \frac{\partial v_t}{\partial n} \right|^2 + c^4 \left| \frac{\partial v}{\partial n} \right|^2 \right) d\sigma dt.$$

Then, there exists  $s_0 > 0$  and  $\lambda$  such that for every  $s \geq s_0$  we absorb the second and third term in the right hand side of (60) which implies

$$\sqrt{s} \int_{\Omega} e^{2s\varphi_\lambda(\cdot, 0)} |v_{tt}(\cdot, 0)|^2 dx + c^4 W(v) + W(v_t) \leq C \int_0^T \int_{\Omega} e^{2s\varphi_\lambda(\cdot, 0)} |f|^2 dx dt \\ + s\lambda C \int_{-T}^T \int_{\Gamma_0} e^{2s\varphi_\lambda} \left( \left| \frac{\partial v_t}{\partial n} \right|^2 + c^4 \left| \frac{\partial v}{\partial n} \right|^2 \right) d\sigma dt.$$

Finally, without loss of generality, we can take  $M_0 > 0$  and  $C > 1$  in definition (41) such that  $\phi(x, 0) \leq C\phi(x, t)$  for all  $x \in \Omega$  and  $t \in [0, T]$ . Then we have  $\varphi_\lambda(x, 0) \leq C_1 \varphi_\lambda(x, t)$  for some  $C_1 = C_1(\lambda)$  independent of  $(x, t) \in \Omega \times [0, T]$ , from where we conclude the desired estimates (11).  $\square$

With the previous Carleman inequality, we can prove the main result of this article.

**Proof of Theorem 1.1.** Using the notation settled in the the previous section (see (20) and (24)), we write the MGT equation in the following way.

$$(61) \quad \begin{cases} L_0 u_t + c^2 L_0 u + \gamma u_{tt} = f, \Omega \times (0, T) \\ u = h, \Gamma \times (0, T) \\ u(\cdot, 0) = u_0, u_t(\cdot, 0) = u_1, u_{tt}(\cdot, 0) = u_2, \Omega. \end{cases}$$

Hence, we will prove a stability estimate for coefficient  $\gamma = \gamma(x)$  in equation (61).



Let us denote by  $u^k$  the solution of equation (61) with coefficient  $\gamma_k$ , for  $k = 1, 2$ , which existence is guaranteed by Theorem 2.1. Hence  $z := u^1 - u^2$  solves the following system.

$$(62) \quad \begin{cases} L_0 z_t + c^2 L_0 z + \gamma_1(x) z_{tt} = (\gamma_2 - \gamma_1) R(x, t), \Omega \times (0, T) \\ z = 0, \Gamma \times (0, T) \\ z(\cdot, 0) = z_t(\cdot, 0) = z_{tt}(\cdot, 0) = 0, \Omega \end{cases}$$

where  $R = \partial_t^2 u^2$ . Then  $y := \partial_t z$  satisfies

$$(63) \quad \begin{cases} L_0 y_t + c^2 L_0 y + \gamma_1(x) y_{tt} = (\gamma_2 - \gamma_1) \partial_t R(x, t), \Omega \times (0, T) \\ y = 0, \Gamma \times (0, T) \\ y(\cdot, 0) = y_t(\cdot, 0) = 0, y_{tt}(\cdot, 0) = (\gamma_2 - \gamma_1) R(x, 0), \Omega \end{cases}$$

Since  $\gamma_2 - \gamma_1$  belongs, in particular, to  $L^2(\Omega)$  and  $R \in H^1(0, T; L^\infty(\Omega))$ , by Theorem 2.1, we obtain that the Cauchy problem (63) is well-posed and admits a unique solution

$$(y, y_t, y_{tt}) \in C([0, T]; (H^2(\Omega) \cap H_0^1(\Omega)) \times H_0^1(\Omega) \times L^2(\Omega)).$$

Moreover, from Theorem 2.4 the normal derivative  $\frac{\partial y}{\partial n}$  belongs to  $H^1(0, T; L^2(\Gamma))$  and satisfy

$$\left\| \frac{\partial y}{\partial n} \right\|_{H^1(0, T; L^2(\Gamma))}^2 \leq C \|\gamma_2 - \gamma_1\|_{L^2(\Omega)}^2 (\|R(\cdot, 0)\|_{L^\infty(\Omega)}^2 + \|\partial_t R\|_{L^2(0, T; L^\infty(\Omega))}^2).$$

This last estimate gives at the end  $\frac{\partial z}{\partial n}$  belongs to  $H^2(0, T; L^2(\Gamma_0))$  and proves the second inequality of (8).

Next, we apply Theorem 1.6 to  $y$ . From system (63) we have

$$\int_0^T \int_\Omega e^{2s\varphi_\lambda} |Ly|^2 dx dt \leq C (\|\gamma_1\|_{L^\infty(\Omega)}, \|\partial_t R\|_{L^2(0, T; L^\infty(\Omega))}) \int_\Omega e^{2s\varphi_\lambda(\cdot, 0)} |\gamma_2 - \gamma_1|^2 dx.$$

Thus, from (11)

$$\begin{aligned} \sqrt{s} \int_\Omega e^{2s\varphi_\lambda(\cdot, 0)} |\gamma_2 - \gamma_1|^2 |R(x, 0)|^2 dx &\leq C \int_\Omega e^{2s\varphi_\lambda(\cdot, 0)} |\gamma_2 - \gamma_1|^2 dx \\ &\quad + Cs\lambda \int_0^T \int_{\Gamma_0} e^{2s\varphi_\lambda} \left( \left| \frac{\partial y_t}{\partial n} \right|^2 + c^4 \left| \frac{\partial y}{\partial n} \right|^2 \right) d\sigma dt, \end{aligned}$$

which implies, using that  $|R(x, 0)| = |u_2| \geq \eta > 0$  a.e in  $\Omega$ ,

$$\begin{aligned} \eta^2 \sqrt{s} \int_\Omega e^{2s\varphi_\lambda(\cdot, 0)} |\gamma_2 - \gamma_1|^2 dx &\leq C \int_\Omega e^{2s\varphi_\lambda(\cdot, 0)} |\gamma_2 - \gamma_1|^2 dx \\ &\quad + Cs\lambda \int_0^T \int_{\Gamma_0} e^{2s\varphi_\lambda} \left( \left| \frac{\partial y_t}{\partial n} \right|^2 + c^4 \left| \frac{\partial y}{\partial n} \right|^2 \right) d\sigma dt. \end{aligned}$$

Therefore, taking  $s$  large enough we absorb the first term in the right hand side and we have

$$\eta^2 \int_{\Omega} e^{2s\varphi_{\lambda}(\cdot, 0)} |\gamma_2 - \gamma_1|^2 dx \leq C\sqrt{s}\lambda \int_0^T \int_{\Gamma_0} e^{2s\varphi_{\lambda}} \left( \left| \frac{\partial y_t}{\partial n} \right|^2 + c^4 \left| \frac{\partial y}{\partial n} \right|^2 \right) d\sigma dt,$$

which is the first inequality of (8) and the proof is finished.  $\square$

#### 4. RECONSTRUCTION OF THE COEFFICIENT

In this section we shall propose a reconstruction algorithm for the unknown parameter  $\gamma$ , from measurements of the normal derivative of the solution  $u(\gamma)$  of the MGT equation (61). This algorithm is an extension of the work of Baudouin, Buhan and Ervedoza [3], in which they propose a reconstruction algorithm for the potential of the wave equation.

By Theorem 1.1, we know that the knowledge of  $\frac{\partial u}{\partial n}$  on  $\Gamma_0 \times (0, T)$  is enough to identify the parameter  $\gamma$ .

Let  $\gamma \in \mathcal{B}_M$ . Let  $g \in L^2(\Omega \times (0, T))$  and  $\mu \in H^1(0, T; L^2(\Gamma_0))$ . Given  $\varphi_{\lambda}$  defined in (40) with  $\lambda > 0$  given by Theorem 1.6, we define the functional

$$(64) \quad J[\mu, g](y) = \frac{1}{2s} \int_0^T \int_{\Omega} e^{2s\varphi_{\lambda}} |Ly - g|^2 dx dt + \frac{1}{2} \int_0^T \int_{\Gamma_0} e^{2s\varphi_{\lambda}} \left( \left| \frac{\partial y}{\partial n} - \mu \right|^2 + \left| \frac{\partial y_t}{\partial n} - \mu_t \right|^2 \right) d\sigma dt,$$

defined in the space

$$(65) \quad \mathcal{V} = \{y \in L^2(0, T; H_0^1(\Omega)) \text{ with } Ly \in L^2(\Omega \times (0, T)), y(\cdot, 0) = y_t(\cdot, 0) = 0 \text{ and } y_{tt}(\cdot, 0) \in L^2(\Omega)\},$$

with the family of semi-norms

$$(66) \quad \|y\|_{\mathcal{V}, s}^2 := \frac{1}{s} \int_0^T \int_{\Omega} e^{2s\varphi_{\lambda}} |Ly|^2 dx dt + \int_0^T \int_{\Gamma_0} e^{2s\varphi_{\lambda}} \left( \left| \frac{\partial y}{\partial n} \right|^2 + \left| \frac{\partial y_t}{\partial n} \right|^2 \right) d\sigma dt.$$

A few remarks about this semi-norms (for more details see [3, Section 4]):

**Remark 4.1.** (1) *Since the weighted functions  $e^{s\varphi_{\lambda}}$  are bounded from below and from above by a positive constants depending on  $s$ , the semi-norm (66) is equivalent to*

$$\|y\|_{\mathcal{V}}^2 := \int_0^T \int_{\Omega} |Ly|^2 dx dt + \int_0^T \int_{\Gamma_0} \left( \left| \frac{\partial y}{\partial n} \right|^2 + \left| \frac{\partial y_t}{\partial n} \right|^2 \right) d\sigma dt,$$

*in the sense that there exists a constant  $C = C(s)$ , such that for all  $y \in \mathcal{V}$*

$$\frac{1}{C} \|y\|_{\mathcal{V}}^2 \leq \|y\|_{\mathcal{V}, s}^2 \leq C \|y\|_{\mathcal{V}}^2.$$

- (2) By Theorem 1.6, there exists  $s_0 > 0$  such that for every  $s \geq s_0$  the semi-norm (66) is actually a norm. Hence, from (1) we have that  $\|\cdot\|_{\mathcal{V},s}$  is a norm for all  $s > 0$ . In the rest of the paper, we will omit the subscript  $s$  in the notation.

The first result concerning the reconstruction of  $\gamma$ , is to guarantee that the functional  $J[\mu, g]$  reaches the minimum. Moreover, we have the following uniqueness result.

**Theorem 4.2.** *Assume the same hypotheses of Theorem 1.6 and assume that  $g \in L^2(\Omega \times (0, T))$  and  $\mu \in H^1(0, T; L^2(\Gamma_0))$ . Then, for all  $s > 0$  and  $\gamma \in \mathcal{B}_M$ , the functional  $J[\mu, g]$  defined by (64) is continuous, strictly convex and coercive on  $\mathcal{V}$ . Besides, the minimizer  $y^*$  satisfies*

$$\|y^*\|_{\mathcal{V}}^2 \leq \frac{4}{s} \int_0^T \int_{\Omega} e^{2s\varphi\lambda} |g|^2 dxdt + 4 \int_0^T \int_{\Gamma_0} e^{2s\varphi\lambda} (|\mu|^2 + |\mu_t|^2) d\sigma dt.$$

*Proof.* The continuity and convexity is immediately. Let us see the coercivity.

$$\begin{aligned} J[\mu, g](y) &= \frac{1}{2s} \int_0^T \int_{\Omega} e^{2s\varphi\lambda} |Ly - g|^2 dxdt + \frac{1}{2} \int_0^T \int_{\Gamma_0} e^{2s\varphi\lambda} \left( \left| \frac{\partial y}{\partial n} - \mu \right|^2 + \left| \frac{\partial y_t}{\partial n} - \mu_t \right|^2 \right) d\sigma dt \\ &= \frac{1}{2s} \int_0^T \int_{\Omega} e^{2s\varphi\lambda} |Ly|^2 dxdt + \frac{1}{2s} \int_0^T \int_{\Omega} e^{2s\varphi\lambda} |g|^2 dxdt - \frac{1}{s} \int_0^T \int_{\Omega} e^{2s\varphi\lambda} g Ly dxdt \\ &\quad + \frac{1}{2} \int_0^T \int_{\Gamma_0} e^{2s\varphi\lambda} \left( \left| \frac{\partial y}{\partial n} \right|^2 + \left| \frac{\partial y_t}{\partial n} \right|^2 \right) d\sigma dt - \int_0^T \int_{\Gamma_0} e^{2s\varphi\lambda} \left( \mu \frac{\partial y}{\partial n} + \mu_t \frac{\partial y_t}{\partial n} \right) \\ &\quad + \frac{1}{2} \int_0^T \int_{\Gamma_0} e^{2s\varphi\lambda} (|\mu|^2 + |\mu_t|^2) d\sigma dt. \end{aligned}$$

Using the fact that  $2ab \leq 2a^2 + \frac{b^2}{2}$ , we deduce

$$\begin{aligned} J[\mu, g](y) &\geq \frac{1}{4s} \int_0^T \int_{\Omega} e^{2s\varphi\lambda} |Ly|^2 dxdt - \frac{1}{s} \int_0^T \int_{\Omega} e^{2s\varphi\lambda} |g|^2 dxdt \\ &\quad + \frac{1}{4} \int_0^T \int_{\Gamma_0} e^{2s\varphi\lambda} \left( \left| \frac{\partial y}{\partial n} \right|^2 + \left| \frac{\partial y_t}{\partial n} \right|^2 \right) d\sigma dt - \int_0^T \int_{\Gamma_0} e^{2s\varphi\lambda} (|\mu|^2 + |\mu_t|^2) d\sigma dt \\ &= \frac{1}{4} \|y\|_{\mathcal{V}}^2 - \frac{1}{s} \int_0^T \int_{\Omega} e^{2s\varphi\lambda} |g|^2 dxdt - \int_0^T \int_{\Gamma_0} e^{2s\varphi\lambda} (|\mu|^2 + |\mu_t|^2) d\sigma dt \end{aligned}$$

Therefore, the functional  $J[\mu, g]$  admits a unique minimizer  $y^*$  in  $\mathcal{V}$ .

Now, let us prove the estimates on the minimizer. First, we develop the functional  $J[\mu, g](y^*)$ :

$$\begin{aligned} J[\mu, g](y^*) &= \frac{1}{2s} \int_0^T \int_{\Omega} e^{2s\varphi\lambda} |Ly^* - g|^2 dxdt + \frac{1}{2} \int_0^T \int_{\Gamma_0} e^{2s\varphi\lambda} \left( \left| \frac{\partial y^*}{\partial n} - \mu \right|^2 + \left| \frac{\partial y_t^*}{\partial n} - \mu_t \right|^2 \right) d\sigma dt \\ &= \frac{1}{2s} \int_0^T \int_{\Omega} e^{2s\varphi\lambda} |Ly^*|^2 dxdt + \frac{1}{2} \int_0^T \int_{\Gamma_0} e^{2s\varphi\lambda} \left( \left| \frac{\partial y^*}{\partial n} \right|^2 + \left| \frac{\partial y_t^*}{\partial n} \right|^2 \right) d\sigma dt \\ &\quad + \frac{1}{2s} \int_0^T \int_{\Omega} e^{2s\varphi\lambda} |g|^2 dxdt + \frac{1}{2} \int_0^T \int_{\Gamma_0} e^{2s\varphi\lambda} (|\mu|^2 + |\mu_t|^2) d\sigma dt \\ &\quad - \frac{1}{s} \int_0^T \int_{\Omega} e^{2s\varphi\lambda} g Ly^* dxdt - \int_0^T \int_{\Gamma_0} e^{2s\varphi\lambda} \left( \mu \frac{\partial y^*}{\partial n} + \mu_t \frac{\partial y_t^*}{\partial n} \right) d\sigma dt. \end{aligned}$$

Next, since  $y^*$  is the minimizer, we have that  $J[\mu, g](y^*) \leq J[\mu, g](0)$ , which implies in particular

$$\begin{aligned} &\frac{1}{2s} \int_0^T \int_{\Omega} e^{2s\varphi\lambda} |Ly^*|^2 dxdt + \frac{1}{2} \int_0^T \int_{\Gamma_0} e^{2s\varphi\lambda} \left( \left| \frac{\partial y^*}{\partial n} \right|^2 + \left| \frac{\partial y_t^*}{\partial n} \right|^2 \right) d\sigma dt \\ &\leq \frac{1}{s} \int_0^T \int_{\Omega} e^{2s\varphi\lambda} g Ly^* dxdt + \int_0^T \int_{\Gamma_0} e^{2s\varphi\lambda} \left( \mu \frac{\partial y^*}{\partial n} + \mu_t \frac{\partial y_t^*}{\partial n} \right) d\sigma dt \end{aligned}$$

Therefore, using that  $2ab \leq 2a^2 + \frac{b^2}{2}$  and the definition of the norm  $\|\cdot\|_{\mathcal{V}}$ , we deduce

$$\frac{1}{4} \|y^*\|_{\mathcal{V}}^2 \leq \frac{1}{s} \int_0^T \int_{\Omega} e^{2s\varphi\lambda} |g|^2 dxdt + \int_0^T \int_{\Gamma_0} e^{2s\varphi\lambda} (|\mu|^2 + |\mu_t|^2) d\sigma dt.$$

□

Secondly, the following Theorem gives a relationship between the unique minimizer of  $J[\mu, g]$  and the source term  $g$ . This is, together with the Theorem 1.6, an essential result for the proof of convergence of our algorithm of reconstruction.

**Theorem 4.3.** *Assume the same hypotheses of Theorem 1.6 and assume that  $\mu \in H^1(0, T; L^2(\Gamma_0))$  and  $g^1, g^2 \in L^2(\Omega \times (0, T))$ . Let  $y^{*,i}$  be the unique minimizer of the functional  $J[\mu, g^i]$ , for  $i = 1, 2$ . Then, there exists  $s_0 > 0$  and a constant  $C > 0$  such that for all  $s \geq s_0$*

$$(67) \quad \sqrt{s} \int_{\Omega} e^{2s\varphi\lambda(\cdot, 0)} |y_{tt}^{*,1}(\cdot, 0) - y_{tt}^{*,2}(\cdot, 0)|^2 dx \leq C \int_0^T \int_{\Omega} e^{2s\varphi\lambda} |g^1 - g^2|^2 dxdt.$$

*Proof.* Since  $y^{*,i}$  is the unique minimizer of  $J[\mu, g^i]$ , for  $i = 1, 2$ , we have that for all  $y \in \mathcal{V}$

$$(68) \quad \begin{aligned} &\frac{1}{s} \int_0^T \int_{\Omega} e^{2s\varphi\lambda} (Ly^{*,1} - g^1) Ly dxdt \\ &\quad + \int_0^T \int_{\Gamma_0} e^{2s\varphi\lambda} \left[ \left( \frac{\partial y^{*,1}}{\partial n} - \mu \right) \frac{\partial y}{\partial n} + \left( \frac{\partial y_t^{*,1}}{\partial n} - \mu_t \right) \frac{\partial y_t}{\partial n} \right] d\sigma dt = 0, \end{aligned}$$

and

$$(69) \quad \frac{1}{s} \int_0^T \int_{\Omega} e^{2s\varphi\lambda} (Ly^{*,2} - g^2) Ly dx dt + \int_0^T \int_{\Gamma_0} e^{2s\varphi\lambda} \left[ \left( \frac{\partial y^{*,2}}{\partial n} - \mu \right) \frac{\partial y}{\partial n} + \left( \frac{\partial y_t^{*,2}}{\partial n} - \mu_t \right) \frac{\partial y_t}{\partial n} \right] d\sigma dt = 0.$$

Subtracting (68) and (69), for  $y = y^{*,1} - y^{*,2}$ , we deduce that

$$\begin{aligned} \frac{1}{s} \int_0^T \int_{\Omega} e^{2s\varphi\lambda} |Ly|^2 dx dt + \int_0^T \int_{\Gamma_0} e^{2s\varphi\lambda} \left( \left| \frac{\partial y}{\partial n} \right|^2 + \left| \frac{\partial y_t}{\partial n} \right|^2 \right) d\sigma dt \\ = \frac{1}{s} \int_0^T \int_{\Omega} e^{2s\varphi\lambda} (g^1 - g^2) Ly dx dt, \end{aligned}$$

Then, applying again  $2ab \leq 2a^2 + \frac{b^2}{2}$  we obtain

$$(70) \quad \frac{1}{2} \int_0^T \int_{\Omega} e^{2s\varphi\lambda} |Ly|^2 dx dt + s \int_0^T \int_{\Gamma_0} e^{2s\varphi\lambda} \left( \left| \frac{\partial y}{\partial n} \right|^2 + \left| \frac{\partial y_t}{\partial n} \right|^2 \right) d\sigma dt \leq 2 \int_0^T \int_{\Omega} e^{2s\varphi\lambda} |g^1 - g^2|^2 dx dt,$$

Finally, by the estimate (11) of Theorem 1.6 we obtain the desired result.  $\square$

Under the new writing of the equation, we present our algorithm and the convergence result of this.

---

### Algorithm:

---

(1) **Initialization:**  $\gamma^0 = 0$ .

(2) **Iteration:** From  $k$  to  $k + 1$

Step 1 - Given  $\gamma^k$  we consider  $\mu^k = \partial_t \left( \frac{\partial u(\gamma^k)}{\partial n} - \frac{\partial u(\gamma)}{\partial n} \right)$  and  $\mu_t^k = \partial_t \left( \frac{\partial u_t(\gamma^k)}{\partial n} - \frac{\partial u_t(\gamma)}{\partial n} \right)$  on  $\Gamma_0 \times (0, T)$ , where  $u(\gamma^k)$  and  $u(\gamma)$  are the solution of the problems

$$(71) \quad \begin{cases} L_0 u_t + c^2 L_0 u + \gamma^k(x) u_{tt} = f, \Omega \times (0, T) \\ u = g, \Gamma \times (0, T) \\ u(\cdot, 0) = u_0, u_t(\cdot, 0) = u_1, u_{tt}(\cdot, 0) = u_2, \Omega \end{cases}$$

and

$$(72) \quad \begin{cases} L_0 u_t + c^2 L_0 u + \gamma(x) u_{tt} = f, \Omega \times (0, T) \\ u = g, \Gamma \times (0, T) \\ u(\cdot, 0) = u_0, u_t(\cdot, 0) = u_1, u_{tt}(\cdot, 0) = u_2. \Omega \end{cases}$$

Step 2 - Minimize the functional  $J[\mu^k, 0]$  on the admissible trajectories  $y \in \mathcal{V}$ :

$$J[\mu^k, 0](y) = \frac{1}{2s} \int_0^T \int_{\Omega} e^{2s\varphi\lambda} |L_0 y_t + c^2 L_0 y + \gamma^k(x) y_{tt}|^2 dx dt \\ + \frac{1}{2} \int_0^T \int_{\Gamma_0} e^{2s\varphi\lambda} \left( \left| \frac{\partial y}{\partial n} - \mu^k \right|^2 + \left| \frac{\partial y_t}{\partial n} - \mu_t^k \right|^2 \right) d\sigma dt$$

Step 3 - Let  $y^{*,k}$  the minimizer of  $J[\mu^k, 0]$  and

$$(73) \quad \tilde{\gamma}^{k+1} = \gamma^k + \frac{y_{tt}^{*,k}(\cdot, 0)}{u_2}.$$

Step 4 - Finally, consider  $\gamma^{k+1} = T(\tilde{\gamma}^{k+1})$ , where

$$(74) \quad T(\gamma) = \begin{cases} M & \text{if } \gamma(x) > M \\ \gamma & \text{if } 0 \leq \gamma(x) \leq M \\ 0 & \text{if } \gamma(x) < 0. \end{cases}$$

Therefore, under the previous Theorem 4.3, we can prove the convergence of this algorithm:

**Theorem 4.4.** *Assume the same hypotheses of Theorem 1.6, and the following assumption of  $u(\gamma)$  :*

$$(75) \quad u(\gamma) \in H^3(0, T; L^\infty(\Omega)) \text{ and } |u_2| \geq \eta > 0.$$

Then, there exists a constant  $C > 0$  and  $s_0 > 0$  such that for all  $s \geq s_0$  and  $k \in \mathbb{N}$

$$(76) \quad \int_{\Omega} e^{2s\varphi\lambda(\cdot, 0)} (\gamma^{k+1} - \gamma)^2 dx \leq \frac{C}{\sqrt{s}} \int_{\Omega} e^{2s\varphi\lambda(\cdot, 0)} (\gamma^k - \gamma)^2 dx.$$

*Proof.* We consider  $y^k = \partial_t(u(\gamma^k) - u(\gamma))$ , which is the solution of

$$(77) \quad \begin{cases} L_0 y_t^k + c^2 L_0 y^k + \gamma(x)^k y_{tt}^k = (\gamma - \gamma^k) \partial_t R(x, t), \Omega \times (0, T) \\ y^k = 0, \Gamma \times (0, T) \\ y^k(\cdot, 0) = 0, y_t^k(\cdot, 0) = 0, y_{tt}^k(\cdot, 0) = (\gamma - \gamma^k) R(x, t), \Omega \end{cases}$$

where  $R(x, t) = \partial_t^2 u(\gamma)$ . Thus,

$$(78) \quad \mu^k = \frac{\partial y^k}{\partial n}, \quad \mu_t^k = \frac{\partial y_t^k}{\partial n}.$$

We observe that  $y^k$  belongs to  $\mathcal{V}$ . Therefore, by (78), the solution  $y^k$  of (77) satisfy the Euler–Lagrange equations associated to the functional  $J[\mu^k, g^k]$ , where  $g^k = (\gamma - \gamma^k) \partial_t R(x, t)$ . Since  $J[\mu^k, g^k]$  admits a unique minimizer,  $y^k$  corresponds to minimum of  $J[\mu^k, g^k]$ .

Let  $y^{*,k}$  be the minimizer of  $J[\mu^k, 0]$ . From Theorem 4.3 we obtain that

$$(79) \quad \sqrt{s} \int_{\Omega} e^{2s\varphi_{\lambda}(\cdot,0)} |y_{tt}^{*,k}(\cdot, 0) - y_{tt}^k(\cdot, 0)|^2 dx \leq C \int_0^T \int_{\Omega} e^{2s\varphi_{\lambda}} |(\gamma - \gamma^k) \partial_t R(x, t)|^2 dx dt.$$

From (73) and (77)

$$y_{tt}^{*,k}(\cdot, 0) = (\tilde{\gamma}^{k+1} - \gamma^k) u_2, \quad y_{tt}^k(\cdot, 0) = (\gamma - \gamma^k) u_2.$$

This implies that, using that  $|u_2| \geq \eta > 0$

$$(80) \quad \eta^2 \sqrt{s} \int_{\Omega} e^{2s\varphi_{\lambda}(\cdot,0)} (\tilde{\gamma}^{k+1} - \gamma)^2 dx \leq C \int_0^T \int_{\Omega} e^{2s\varphi_{\lambda}} |(\gamma - \gamma^k) \partial_t R(x, t)|^2 dx dt.$$

Since the function  $T$  defined in (74) is Lipschitz continuous and satisfy  $T(\gamma) = \gamma$ , we obtain

$$(81) \quad |\tilde{\gamma}^{k+1} - \gamma| \geq |T(\tilde{\gamma}^{k+1}) - T(\gamma)| = |\gamma^{k+1} - \gamma|.$$

On the other hand, since  $\phi(\cdot, t)$  is decreasing in  $t \in (0, T)$  and  $\partial_t R(x, t) \in L^2(0, T; L^{\infty}(\Omega))$ , we conclude

$$\int_{\Omega} e^{2s\varphi_{\lambda}(\cdot,0)} (\tilde{\gamma}^{k+1} - \gamma)^2 dx \leq \frac{C}{\sqrt{s}} \frac{\|\partial_t R(x, t)\|_{L^2(0,T;L^{\infty}(\Omega))}}{\eta^2} \int_{\Omega} e^{2s\varphi_{\lambda}(\cdot,0)} |(\gamma - \gamma^k)| dx dt.$$

□

## REFERENCES

- [1] C. Bardos, G. Lebeau, and J. Rauch. Sharp sufficient conditions for the observation, control, and stabilization of waves from the boundary. *SIAM Journal on Control and Optimization*, 30(5):1024–1065, 1992. 3
- [2] L. Baudouin. Lipschitz stability in an inverse problem for the wave equation. Preprint, available at: <http://hail.archives-ouvertes.fr/hal-00598876/fr/>. 2010. 3
- [3] L. Baudouin, M. De Buhan, and S. Ervedoza. Global Carleman estimates for waves and applications. *Communications in Partial Differential Equations*, 38(5):823–859, 2013. 5, 12, 13, 18
- [4] L. Beilina and M. V. Klibanov. *Approximate global convergence and adaptivity for coefficient inverse problems*. Springer Science & Business Media, 2012. 5
- [5] L. Beilina and M. V. Klibanov. Globally strongly convex cost functional for a coefficient inverse problem. *Nonlinear analysis: real world applications*, 22:272–288, 2015. 5
- [6] M. Bellassoued. Uniqueness and stability in determining the speed of propagation of second-order hyperbolic equation with variable coefficients. *Appl. Anal.*, 83(10):983–1014, 2004. 5
- [7] M. Bellassoued and M. Yamamoto. *Carleman estimates and applications to inverse problems for hyperbolic systems*. Springer Monographs in Mathematics. Springer, Tokyo, 2017. 5
- [8] A. L. Bukhgeim and M. V. Klibanov. Uniqueness in the large of a class of multidimensional inverse problems. *Dokl. Akad. Nauk SSSR*, 260(2):269–272, 1981. 4
- [9] J. A. Conejero, C. Lizama, and F. Rodenas. Chaotic behaviour of the solutions of the Moore-Gibson-Thompson equation. *Applied Mathematics & Information Sciences*, 9(5):2233, 2015. 7
- [10] A. Fursikov and O. Y. Imanuvilov. *Controllability of Evolution Equations. Lecture Notes Series*, volume Vol. 34. Seoul: Seoul National University Research Institute of Mathematics Global Analysis Research Center. 1996. 3
- [11] L. F. Ho. Observabilité frontière de l'équation des ondes. *Comptes rendus de l'Académie des sciences. Série 1, Mathématique*, 302(12):443–446, 1986. 3

- [12] O. Y. Imanuvilov. On Carleman estimates for hyperbolic equations. *Asymptotic Analysis*, 32(3, 4):185–220, 2002. 12
- [13] B. Kaltenbacher. Mathematics of nonlinear acoustics. *Evolution Equations & Control Theory*, 4(4), 2015. 6
- [14] B. Kaltenbacher and I. Lasiecka. Exponential decay for low and higher energies in the third order linear Moore-Gibson-Thompson equation with variable viscosity. *Palest. J. Math*, 1(1):1–10, 2012. 6, 7, 8
- [15] B. Kaltenbacher, I. Lasiecka, and R. Marchand. Wellposedness and exponential decay rates for the Moore-Gibson-Thompson equation arising in high intensity ultrasound. *Control and Cybernetics*, 40:971–988, 2011. 6, 10
- [16] B. Kaltenbacher, I. Lasiecka, and M. K. Pospieszalska. Well-posedness and exponential decay of the energy in the nonlinear Jordan–Moore–Gibson–Thompson equation arising in high intensity ultrasound. *Mathematical Models and Methods in Applied Sciences*, 22(11):1250035, 2012. 6
- [17] J. L. Lions. *Contrôlabilité exacte perturbations et stabilisation de systèmes distribués. Tome 1, Contrôlabilité exacte.*, volume 8. Recherches en Mathématiques Appliquées, Masson, 1988. 10
- [18] S. Liu and R. Triggiani. Global uniqueness and stability in determining the damping and potential coefficients of an inverse hyperbolic problem. *Nonlinear Analysis: Real World Applications*, 12(3):1562–1590, 2011. 3
- [19] S. Liu and R. Triggiani. Inverse problem for a linearized Jordan–Moore–Gibson–Thompson equation. In *New Prospects in Direct, Inverse and Control Problems for Evolution Equations*, pages 305–351. Springer, 2014. 3, 6
- [20] C. Lizama and S. Zamorano. Controllability results for the moore–gibson–thompson equation arising in nonlinear acoustics. *Journal of Differential Equations*, 266(12):7813–7843, 2019. 4, 5, 6
- [21] R. Marchand, T. McDevitt, and R. Triggiani. An abstract semigroup approach to the third-order Moore–Gibson–Thompson partial differential equation arising in high-intensity ultrasound: structural decomposition, spectral analysis, exponential stability. *Mathematical Methods in the Applied Sciences*, 35(15):1896–1929, 2012. 6, 7
- [22] M. Pellicer and J. d. Solà-Morales Rubió. Optimal scalar products in the moore-gibson-thompson equation. *Evolution Equations and Control Theory*, 8(1):203–220, 2019. 6
- [23] J. Puel. Global Carleman inequalities for the wave equations and applications to controllability and inverse problems, in “control of solids and structures: Mathematical modelling and engineering applications”. *Udine (Italy), June, 2004*. 12, 13
- [24] J.-P. Puel and M. Yamamoto. On a global estimate in a linear inverse hyperbolic problem. *Inverse Problems*, 12(6):995–1002, 1996. 5
- [25] M. Yamamoto. Uniqueness and stability in multidimensional hyperbolic inverse problems. *Journal de mathématiques pures et appliquées*, 78(1):65–98, 1999. 3
- [26] M. Yamamoto. Uniqueness and stability in multidimensional hyperbolic inverse problems. *J. Math. Pures Appl. (9)*, 78(1):65–98, 1999. 5
- [27] M. Yamamoto. One unique continuation for a linearized benjaminbonamahony equation. *Journal of Inverse and Ill-posed Problems jiiip*, 11(5):537–543, 2003. 5
- [28] X. Zhang. Explicit observability inequalities for the wave equation with lower order terms by means of Carleman inequalities. *SIAM Journal on Control and Optimization*, 39(3):812–834, 2000. 3



UNIVERSIDAD TÉCNICA FEDERICO SANTA MARÍA, DEPARTAMENTO DE MATEMÁTICA, CASILLA 110-V, VALPARAÍSO, CHILE.

*E-mail address:* `rodrigo.lecaros@usm.cl`

UNIVERSIDAD TÉCNICA FEDERICO SANTA MARÍA, DEPARTAMENTO DE MATEMÁTICA, CASILLA 110-V, VALPARAÍSO, CHILE AND INSTITUT DE MATHÉMATIQUES DE TOULOUSE UMR 5219, CNRS, TOULOUSE, FRANCE.

*E-mail address:* `alberto.mercado@usm.cl`

UNIVERSIDAD DE SANTIAGO DE CHILE, FACULTAD DE CIENCIA, DEPARTAMENTO DE MATEMÁTICA Y CIENCIA DE LA COMPUTACIÓN, CASILLA 307, CORREO 2, SANTIAGO, CHILE.

*E-mail address:* `sebastian.zamorano@usach.cl`