

Generalized Dobrushin Ergodicity Coefficient and Ergodicities of Non-homogeneous Markov Chains

FARRUKH MUKHAMEDOV

Department of Mathematical Sciences, College of Science,
United Arab Emirates University 15551, Al-Ain,
United Arab Emirates.
e-mail: far75m@gmail.com; farrukh.m@uaeu.ac.ae

AHMED AL-RAWASHDEH

Department of Mathematical Sciences, College of Science,
United Arab Emirates University 15551, Al-Ain,
United Arab Emirates.
e-mail: aalrawashdeh@uaeu.ac.ae

Abstract

In our earlier paper, a generalized Dobrushin ergodicity coefficient of Markov operators (acting on abstract state spaces) with respect to a projection P , has been introduced and studied. It turned out that the introduced coefficient was more effective than the usual ergodicity coefficient. In the present work, by means of a left consistent Markov projections and the generalized Dobrushin's ergodicity coefficient, we investigate uniform and weak P -ergodicities of non-homogeneous discrete Markov chains (NDMC) on abstract state spaces. It is easy to show that uniform P -ergodicity implies a weak one, but in general the reverse is not true. Therefore, some conditions are provided together with weak P -ergodicity of NDMC which imply its uniform P -ergodicity. Furthermore, necessary and sufficient conditions are found by means of the Doeblin's condition for the weak P -ergodicity of NDMC. The weak P -ergodicity is also investigated in terms of perturbations. Several perturbative results are obtained which allow us to produce nontrivial examples of uniform and weak P -ergodic NDMC. Moreover, some category results are also obtained. We stress that all obtained results have potential applications in the classical and non-commutative probabilities.

MSC: 47A35; 60J10, 28D05

Key words: uniform P -ergodic; weak P -ergodic; Markov operator; projection; ergodicity coefficient; non-homogeneous discrete Markov chain;

1. INTRODUCTION

The present work is a continuation of the paper [34] where we have introduced a generalized Dobrushin ergodicity coefficient $\delta_P(T)$ of Markov operators (acting on abstract state spaces) with respect to a projection P , and studied its properties. It turns out that the introduced coefficient was more effective than the usual Dobrushin's ergodicity coefficient [7]. Indeed, uniform stability of the trajectories of Markov operator to some projection have been investigated by means of the generalized Dobrushin's ergodicity coefficient, while usual ergodicity coefficient is not applicable in that situation. We stress that in the literature, much attention is paid to homogeneous Markov processes (see [3, 4, 5, 11, 23, 38]). However, a limited number of papers (see [14, 31, 32, 33, 45])

were devoted to investigations of ergodic properties of nonhomogeneous Markov processes in the abstract scheme. In those papers the limiting operators are considered as one-dimensional projections. Therefore, one of the aims of the present paper is to investigate stabilities of nonhomogeneous discrete Markov chains (NDMC) to some projection in the abstract framework. We notice that this abstract scheme contains both classical and quantum settings as particular cases [1, 9] which implies that the obtained results will be new in both cases. In this abstract setting, certain limiting behaviors of Markov operators were investigated in [3, 10, 15, 38].

It is stressed that due to the nonhomogeneity of Markov processes, the investigations of limiting behavior of such processes become very complicated. The Dobrushin's ergodicity coefficient was effectively used in the investigation of ergodicity of nonhomogeneous Markov chains when their limit is a one-dimensional projection (see [7, 16, 19, 20, 23, 35, 36, 37, 39, 42, 44]). If the limit of the chain is a more general projection then the usual coefficient is not effective. Therefore, using the generalized coefficient, we are going to investigate uniform and weak ergodicities of NDMC in the abstract state spaces. In [17, 18] a very simple case of NDMC (on finite dimensional spaces) has been investigated when the limit is a projection. We point out that all obtained results will be new for classical and quantum Markov chains.

As it is mentioned that our purpose is to investigate the stability (in uniform and weak topologies) of NDMC acting on abstract state spaces. Here, by an abstract state space it is meant an ordered Banach space, where the norm has an additivity property on the cone of positive elements. Examples of these spaces include all classical L^1 -spaces and the space of density operators acting on some Hilbert spaces [1, 24]. Moreover, any Banach space can be embedded into some abstract spaces (see Appendix, Example A.3). In the present paper, we are going to study the asymptotic stabilities of NDMC in terms of generalized Dobrushin's ergodicity coefficient. We notice that the Dobrushin coefficient (which extends $\delta(P)$ to abstract state spaces) has been introduced and studied in [15, 32, 33], for Markov operators acting on abstract state spaces.

Let us briefly describe the organization of the paper. In Section 2, we provide preliminary definitions and results on properties of the generalized Dobrushin's ergodicity coefficient. Moreover, the uniform and weak P -ergodicities of NDMC are defined. In Section 3, we introduce sequences of left consistent Markov projections which will be used in the forthcoming sections. We note that if the chain is homogeneous then the uniform and weak P -ergodicities coincide [34], but in the non-homogeneous setting the situation is much more complicated [35, 39, 42]. Therefore, in Section 4, certain relations between these notions are going to be investigated. The results of this section are known facts related to NDMC [35, 32] when the limiting projection is one-dimensional. In Section 5, necessary and sufficient conditions are established by means of the Doeblin's condition for the weak P -ergodicity of NDMC. These extend the results of [8, 32, 33] to an abstract scheme. Note that in [8] similar conditions were found for classical nonhomogeneous Markov processes to satisfy weak ergodicity. In Section 6, we study the weak P -ergodicity in terms of perturbations. Several perturbation results are obtained which allow us to produce nontrivial examples of uniform and weak P -ergodic NDMC. Finally, in Section 7 some category results are also obtained. Namely, in the set of all NDMC (consistent with a left decreasing sequence of projections $\{P_n\}$) $\mathfrak{S}_{\{P_n\}}(X)$, we introduce a metric, according to the set $\mathfrak{S}_{\{P_n\}}^w(X)$ of all weak ergodic w.r.t. $\{P_n\}$ NDMC is a G_δ -dense subset of $\mathfrak{S}_{\{P_n\}}(X)$. Some similar kinds of results have been proved in [36] when $X = \ell^1$. We stress that all obtained results have potential applications in the classical and non-commutative probabilities.

2. PRELIMINARIES

In this section, we recall some necessary definitions and results about abstract state spaces.

Let X be an ordered vector space with a cone $X_+ = \{x \in X : x \geq 0\}$. A subset \mathcal{K} is called a *base* for X , if $\mathcal{K} = \{x \in X_+ : f(x) = 1\}$ for some strictly positive (i.e. $f(x) > 0$ for $x > 0$) linear functional f on X . An ordered vector space X with generating cone X_+ (i.e. $X = X_+ - X_+$) and a fixed base \mathcal{K} , defined by a functional f , is called an *ordered vector space with a base* [1]. Let U be the convex hull of the set $\mathcal{K} \cup (-\mathcal{K})$, and let

$$\|x\|_{\mathcal{K}} = \inf\{\lambda \in \mathbb{R}_+ : x \in \lambda U\}.$$

Then one can see that $\|\cdot\|_{\mathcal{K}}$ is a seminorm on X . Moreover, one has $\mathcal{K} = \{x \in X_+ : \|x\|_{\mathcal{K}} = 1\}$, $f(x) = \|x\|_{\mathcal{K}}$ for $x \in X_+$. Assume that the seminorm becomes a norm and X is a complete space w.r.t. this norm and X_+ is closed subset, then (X, X_+, \mathcal{K}, f) is called *abstract state space*. In this case, \mathcal{K} is a closed face of the unit ball of X , and U contains the open unit ball of X . If the set U is *radially compact* [1], i.e. $\ell \cap U$ is a closed and bounded segment for every line ℓ through the origin of X , then $\|\cdot\|_{\mathcal{K}}$ is a norm. The radial compactness is equivalent to the coincidence of U with the closed unit ball of X . In this case, X is called a *strong abstract state space*. In the sequel, for the sake of simplicity, instead of $\|\cdot\|_{\mathcal{K}}$, the standard notation $\|\cdot\|$ is used. For a better understanding of the difference between a strong abstract state space and a more general class of base norm spaces, the reader is referred to [43].

A positive cone X_+ of an ordered Banach space X is said to be λ -generating if, given $x \in X$, we can find $y, z \in X_+$ such that $x = y - z$ and $\|y\| + \|z\| \leq \lambda\|x\|$. The norm on X is called *regular* (respectively, *strongly regular*) if, given x in the open (respectively, closed) unit ball of X , y can be found in the closed unit ball with $y \geq x$ and $y \geq -x$. The norm is said to be additive on X_+ if $\|x + y\| = \|x\| + \|y\|$ for all $x, y \in X_+$. If X_+ is 1-generating, then X can be shown to be strongly regular. Similarly, if X_+ is λ -generating for all $\lambda > 1$, then X is regular [43]. The following results are well-known.

Theorem 2.1. [46, p.90] *Let X be an ordered Banach space with closed positive cone X_+ . Then the following statements are equivalent:*

- (i) X is an abstract state space;
- (ii) X is regular, and the norm is additive on X_+ ;
- (iii) X_+ is λ -generating for all $\lambda > 1$, and the norm is additive on X_+ .

Theorem 2.2. [43] *Let X be an ordered Banach space with closed positive cone X_+ . Then the following statements are equivalent:*

- (i) X is a strong abstract state space;
- (ii) X is strongly regular, and the norm is additive on X_+ ;
- (iii) X_+ is 1-generating and the norm is additive on X_+ .

In this paper, we consider a general abstract state space for which the convex hull of the base \mathcal{K} and $-\mathcal{K}$ is not assumed to be radially compact (in our previous papers [12, 13, 32, 33] this condition was essential). This consideration has an important advantage: whenever X is an ordered Banach space with a generating cone X_+ whose norm is additive on X_+ , then X admits an equivalent norm that coincides with the original norm on X_+ and renders X that base norm space. Hence, to apply the results of the paper one would only have to check if the norm is additive on X_+ .

Let (X, X_+, \mathcal{K}, f) be an abstract state space. A linear operator $T : X \rightarrow X$ is called *positive*, if $Tx \geq 0$ whenever $x \geq 0$. A positive linear operator $T : X \rightarrow X$ is said to be

Markov, if $T(\mathcal{K}) \subset \mathcal{K}$. It is clear that $\|T\| = 1$, and its adjoint mapping $T^* : X^* \rightarrow X^*$ acts in ordered Banach space X^* with unit f , and moreover, one has $T^*f = f$.

Recall that a family of Markov operators $\{T^{m,n} : X \rightarrow X\}$ ($m \leq n$, $m, n \in \mathbb{N}$) is called a *non-homogeneous discrete Markov chain (NDMC)* if

$$T^{m,n} = T^{k,n}T^{m,k-1}$$

for every $m \leq k \leq n$. Due to this property, to any NDMC $\{T^{m,n}\}$ one can associate a sequence $\{T_n\}_{n=1}^\infty$, (where $T_n = T^{n,n+1}$) of Markov operators. Conversely, any given a sequence of Markov operators $\{T_n\}_{n=1}^\infty$ on X and for $k < n$, by putting

$$T^{k,n} := T_n T_{n-1} \dots T_{k+1}.$$

we also can define a NDMC $\{T^{k,n}\}$. This chain is generated by $\{T_n\}$, such a sequence $\{T_n\}$ is called *generating sequence* of the NDMC. Therefore, NDMC $\{T^{k,n}\}$ can be identified with its generating sequence. In the last section 7, we will use this identification.

Recall that if for a given NDMC $\{T^{k,m}\}$ one has $T^{k,m} = (T^{0,1})^{m-k}$, then such a chain becomes *homogeneous*. In what follows, by $\{T^n\}$ we denote a homogeneous Markov chain, where $T := T^{0,1}$. Equivalently, any NDMC is homogeneous, if its generating sequence is stationary, i.e. $T_n = T_1$ for all $n \in \mathbb{N}$.

Let (X, X_+, \mathcal{K}, f) be an abstract state space and let $\{T^n\}$ be a homogeneous Markov chain on X . Consider a projection operator $P : X \rightarrow X$ (i.e. $P^2 = P$). According to [34] $\{T^n\}$ is called *uniformly P-ergodic* if

$$\lim_{n \rightarrow \infty} \|T^n - P\| = 0.$$

From this definition we immediately find that P must be a Markov projection.

Analogously, we say that a NDMC $\{T^{m,n}\}$ is called *uniformly P-ergodic* if for every $m \geq 0$ one has

$$\lim_{n \rightarrow \infty} \|T^{m,n} - P\| = 0.$$

We note that if $P = T_y$, for some $y \in X_+$, where $T_y(x) = f(x)y$, then the uniform P -ergodicity coincides with uniform ergodicity or uniform asymptotical stability considered in [32, 33].

In [34], we have introduced a generalized notion of the Dobrushin's ergodicity coefficient as follows:

Let (X, X_+, \mathcal{K}, f) be an abstract state space and let $T : X \rightarrow X$ be a linear bounded operator and P be a non-trivial projection operator on X . Then we define

$$(1) \quad \delta_P(T) = \sup_{x \in N_P, x \neq 0} \frac{\|Tx\|}{\|x\|},$$

where

$$(2) \quad N_P = \{x \in X : Px = 0\}.$$

If $P = I$, we put $\delta_P(T) = 1$. The quantity $\delta_P(T)$ is called the *generalized Dobrushin ergodicity coefficient of T with respect to P* .

We notice that if $X = \mathbb{R}^n$, then there are some formulas to calculate this coefficient (see [17, 18]).

In the following remarks, let us have a brief comparison between the coefficients $\delta_P(T)$ and $\delta(T)$. It is noticed that $\delta(T)$ has been introduced and investigated in [32, 33].

Remark 2.3. Let $y_0 \in \mathcal{K}$ and consider the projection $Px = f(x)y_0$. Then one can see that N_P coincides with

$$N = \{x \in X; f(x) = 0\},$$

and in this case $\delta_P(T) = \delta(T)$. Hence, $\delta_P(T)$ indeed is a generalization of $\delta(T)$.

Remark 2.4. Let P be a Markov projection on X . Then, for any Markov operator $T : X \rightarrow X$

$$\delta_P(T) \leq \delta(T).$$

Using this coefficient, we define weak P -ergodicity of NDMC. Namely, a NDMC $\{T^{m,n}\}$ is called *weakly P -ergodic* if for every $m \geq 0$ one has

$$\lim_{n \rightarrow \infty} \delta_P(T^{m,n}) = 0.$$

We point out that the relations between uniform and weak P -ergodicities will be discussed in Section 4.

Let us denote by $\Sigma(X)$ the set of all Markov operators defined on X , and by $\Sigma_P(X)$ we denote the set of all Markov operators T on X with $PT = TP$.

We recall certain properties of $\delta_P(T)$, which are given in the following theorem:

Theorem 2.5. [34] Let (X, X_+, \mathcal{K}, f) be an abstract state space, P be a projection on X and let $T, S \in \Sigma(X)$. Then:

- (i) $0 \leq \delta_P(T) \leq 1$;
 - (ii) $|\delta_P(T) - \delta_P(S)| \leq \delta_P(T - S) \leq \|T - S\|$;
 - (iii) if $P \in \Sigma(X)$, one has
- (3)
$$\delta_P(T) \leq \frac{\lambda}{2} \sup\{\|Tu - Tv\|; u, v \in \mathcal{K} \text{ with } u - v \in N_P\}.$$
- (iv) if $H : X \rightarrow X$ is a bounded linear operator such that $HP = PH$, then
- $$\delta_P(TH) \leq \delta_P(T)\|H\|;$$
- (v) if $H : X \rightarrow X$ is a bounded linear operator such that $PH = 0$, then
- $$\|TH\| \leq \delta_P(T)\|H\|;$$
- (vi) if $S \in \Sigma_P(X)$, then
- $$\delta_P(TS) \leq \delta_P(T)\delta_P(S).$$

We stress that the condition $PS = SP$ in (vi) can be weakened as follows:

Proposition 2.6. Let (X, X_+, \mathcal{K}, f) be an abstract state space, P be a projection on X and let T_1 and T_2 be operators on X . If $T_2(N_P) \subseteq N_P$, then

$$\delta_P(T_1T_2) \leq \delta_P(T_1)\delta_P(T_2).$$

Proof. For all $x \in N_P$ we have $T_2x \in N_P$, so

$$\begin{aligned} \|T_1(T_2x)\| &\leq \delta_P(T_1)\|T_2x\| \\ &\leq \delta_P(T)\delta_P(T_2)\|x\|, \end{aligned}$$

which implies

$$\frac{\|T_1T_2x\|}{\|x\|} \leq \delta_P(T_1)\delta_P(T_2), \quad \forall x \in N_P,$$

then

$$\delta_P(T_1T_2) \leq \delta_P(T_1)\delta_P(T_2),$$

and hence the result follows. \square

Lemma 2.7. *Let (X, X_+, \mathcal{K}, f) be an abstract state space, P be a projection on X and let T be an operator on X . Then $T(N_P) \subseteq N_P$ if and only if $PT = PTP$.*

Proof. Assume that $PT = PTP$. If $x \in N_P$, then $P(Tx) = PTP(x) = 0$, so we get $Tx \in N_P$. Conversely, suppose that $T(N_P) \subseteq N_P$ and $x \in N_P$ and as $N_P = (I - P)X$, so $x = (y - Py)$, for some $y \in X$. Therefore,

$$0 = PTx = PT(y - Py) = PTy - PTPy,$$

which implies that $PT = PTP$. \square

Remark 2.8. *It is easy to check that if $PT = P$ or $PT = TP$, then $PT = PTP$.*

In what follows, we need the following auxiliary fact.

Lemma 2.9. [34] *Let (X, X_+, \mathcal{K}, f) be an abstract state space and let P be a Markov projection. Then for every $x \in N_P$ there exist $u, v \in \mathcal{K}$ with $u - v \in N_P$ such that*

$$x = \alpha(x)(u - v),$$

where $\alpha(x) \in \mathbb{R}_+$ and $\alpha(x) \leq \frac{\lambda}{2}\|x\|$.

3. LEFT CONSISTENT PROJECTIONS

In this section, we are going to study a relation between projections of X .

Definition 3.1. *Let P and Q be projections on X . We say that P is left consistent by Q , and denoted by $P \leq^\ell Q$, if $PQ = P$.*

Proposition 3.2. *The relation \leq^ℓ has the following properties:*

- (i) \leq^ℓ is reflexive;
- (ii) \leq^ℓ is transitive.

Proof. (i) is obvious. To prove (ii), assume that $P_1 \leq^\ell P_2$ and $P_2 \leq^\ell P_3$, which implies that $P_1P_2 = P_1$ and $P_2P_3 = P_2$. Then

$$P_1P_3 = P_1P_2P_3 = P_1P_2 = P_1,$$

and hence $P_1 \leq^\ell P_3$. \square

In what follows, one needs the following property of $\delta_P(T)$ with the relation \leq^ℓ .

Proposition 3.3. *Let $T : X \rightarrow X$ be a linear bounded operator. If P and Q are two projections on X such that $P \leq^\ell Q$, then $\delta_Q(T) \leq \delta_P(T)$.*

Proof. Assume that $P \leq^\ell Q$. Then for every $x \in N_Q$ we get $Px = PQx = 0$, therefore $N_Q \subseteq N_P$ which yields the desired inequality. \square

A sequence $\{P_n\}_{n=1}^\infty$ of projections of X is called *left decreasing*, if $P_{n+1} \leq^\ell P_n$, for all $n \in \mathbb{N}$, i.e. P_{n+1} is left consistent by P_n .

Example 3.4. *Let (X, X_+, \mathcal{K}, f) be an abstract state space and let $\{z_n\}$ be a sequence in \mathcal{K} such that $z_n \rightarrow z$. Construct the one dimensional projection $P_n := T_{z_n}$, i.e. $P_n x = f(x)z_n$. Then $\{P_n\}$ is a left decreasing sequence of projections, indeed*

$$\begin{aligned} P_{n+1}P_n x &= P_{n+1}(f(x)z_n) \\ &= f(x)P_{n+1}(z_n) \\ &= f(x)f(z_n)z_{n+1} \\ &= f(x)z_{n+1} \\ &= P_{n+1}(x). \end{aligned}$$

Moreover $P_n \rightarrow P = T_z$

In particular, we consider the following example:

Example 3.5. Consider the space ℓ_1 , and recall that $\mathcal{K} = \{x \in \ell_1; \sum_{n=1}^{\infty} x_n = 1, x_n \geq 0\}$. For all $n \in \mathbb{N}$, define

$$z_n = \left(\frac{1}{2}, \frac{1}{2^2}, \frac{1}{2^3}, \dots, \frac{1}{2^n}, \frac{1}{2^n}, 0, 0, \dots, 0 \right).$$

Then it is clear that $z_n \in \mathcal{K}$ and $z_n \rightarrow z = (\frac{1}{2}, \frac{1}{2^2}, \frac{1}{2^3}, \dots)$, as $\|z_n - z\| = \sum_{k=n+1}^{\infty} \frac{1}{2^k} \rightarrow 0$. Due to the previous example, the projections $P_n = T_{z_n}$ form a left decreasing sequence.

Next result gives some important properties of left consistent sequences of projections.

Lemma 3.6. Let $\{P_n\}_{n=1}^{\infty}$ be a sequence of projections of X . Then the following statements hold:

- (i) if $\{P_n\}_{n=1}^{\infty}$ is left decreasing, then $P_m \leq^\ell P_k$, for all $m \geq k$;
- (ii) if $\{P_n\}_{n=1}^{\infty}$ is left decreasing and $P_n \rightarrow P$ in norm, as $n \rightarrow \infty$, then P is a projection and $P \leq^\ell P_k$, for all $k \in \mathbb{N}$.

Proof. (i) Assume that $\{P_n\}_{n=1}^{\infty}$ is a left decreasing sequence. Then for all $m \geq k$, we have

$$\begin{aligned} P_m P_k &= P_m P_{m-1} P_k \\ &= P_m P_{m-1} P_{m-2} P_k \\ &\vdots \\ &= P_m P_{m-1} P_{m-2} \dots P_{k+1} P_k \\ &= P_m P_{m-1} P_{m-2} \dots P_{k+1} \\ &\vdots \\ &= P_m P_{m-1} \\ &= P_m. \end{aligned}$$

(ii) is obvious. □

4. UNIFORM P -ERGODICITY AND WEAK P -ERGODICITY OF NDMC

In this section, we discuss some relations between weak and uniform P -ergodicities of NDMC. The following result show that weak P -ergodicity is indeed weaker than the uniform P -ergodicity.

Proposition 4.1. Let X be an abstract state space. Then every uniformly P -ergodic NDMC is weakly P -ergodic.

Proof. By Theorem 2.5(iii), we have

$$\begin{aligned} \delta_P(T^{k,n}) &\leq \frac{\lambda}{2} \sup \|T^{k,n}u - T^{k,n}v\| \quad (u, v \in \mathcal{K}, \text{ and } Pu = Pv) \\ &= \frac{\lambda}{2} \sup \|T^{k,n}u - Pu + Pv - T^{k,n}v\| \\ &\leq \frac{\lambda}{2} (\sup \|T^{k,n}u - Pu\| + \sup \|T^{k,n}v - Pv\|) \\ &\leq \lambda \|T^{k,n} - P\| \rightarrow 0, \text{ as } n \rightarrow \infty \end{aligned}$$

and hence $\delta_P(T^{k,n}) \rightarrow 0$ as $n \rightarrow \infty$, which completes the proof. \square

Before discussing the reverse direction, let us consider some examples of uniform P -ergodic NDMC.

Example 4.2. Let (X, X_+, \mathcal{K}, f) be an abstract state space. Let $\{\tilde{T}_n\}_{n=1}^\infty$ be a sequence of Markov operators such that $\bigcap_{n=1}^\infty \text{Fix}(\tilde{T}_n) \neq \emptyset$ (here $\text{Fix}(T) = \{x \in \mathcal{K} : Tx = x\}$). For a given $a \in (0, 1)$ and $z_0 \in \mathcal{K}$ with $z_0 \in \bigcap_{n=1}^\infty \text{Fix}(\tilde{T}_n)$, define the operators

$$T_n = aT_{z_0} + (1-a)\tilde{T}_n.$$

One can see that

$$\begin{aligned} T_n T_m &= (aT_{z_0} + (1-a)\tilde{T}_n)(aT_{z_0} + (1-a)\tilde{T}_m) \\ &= a^2 T_{z_0} + a(1-a)T_{z_0}\tilde{T}_m + a(1-a)\tilde{T}_n T_{z_0} + (1-a)^2 \tilde{T}_n \tilde{T}_m \\ &= a^2 T_{z_0} + a(1-a)T_{z_0} + a(1-a)T_{z_0} + (1-a)^2 \tilde{T}_n \tilde{T}_m \\ &= (1 - (1-a)^2)T_{z_0} + (1-a)^2 \tilde{T}_n \tilde{T}_m. \end{aligned}$$

Hence, for all $k < n$

$$T^{k,n} = T_n T_{n-1} \dots T_{k+1} = (1 - (1-a)^{n-k})T_{z_0} + (1-a)^{n-k} \tilde{T}_n \dots \tilde{T}_{k+1}.$$

Therefore,

$$\|T^{k,n} - T_{z_0}\| \leq (1-a)^{n-k} \|T_{z_0} - \tilde{T}_n \dots \tilde{T}_{k+1}\| \leq 2(1-a)^{n-k} \rightarrow 0,$$

as $n \rightarrow \infty$, which proves that $\{T^{k,n}\}$ is uniformly P -ergodic, where $P = T_{z_0}$.

It is interesting to find some conditions which together with weak P -ergodicity of NDMC imply its uniform P -ergodicity.

Theorem 4.3. Let (X, X_+, \mathcal{K}, f) be an abstract state space and $\{T_n\}$ be a generating sequence of NDMC. Let $\{P_n\}$ be a sequence of projections of X such that

- (i) $T_n P_n = P_n T_n = P_n, \forall n \in \mathbb{N}$,
- (ii) $\{P_n\}$ is left decreasing sequence of projections,
- (iii) $\sum_{n=k}^\infty \|P_{n+1} - P_n\| \rightarrow 0$, as $k \rightarrow \infty$.

If $\{T^{k,n}\}$ is weakly P -ergodic, then it is uniformly P -ergodic.

Proof. From the hypotheses (iii) and using the standard argument, one finds that $P_n \rightarrow P$ (in norm), where P is a projection. Due to (ii) Lemma 3.6 we have $P \leq^\ell P_k$, for all $k \in \mathbb{N}$. As

$$\|T^{m,n} - P\| \leq \|T^{m,n} - P_n\| + \|P_n - P\|$$

for $m < n$, it is enough to prove that $\|T^{m,n} - P_n\| \rightarrow 0$, as $n \rightarrow \infty$.

According to (i), one has

$$\begin{aligned} T^{k,n} P_{k+1} &= T^{k+1,n} T_{k+1} P_{k+1} \\ &= T^{k+1,n} P_{k+1} \\ &= T^{k+1,n} (P_{k+1} - P_{k+2}) + T^{k+1,n} P_{k+2} \\ &\vdots \\ &= \sum_{l=k+1}^{n-1} T^{l,n} (P_l - P_{l+1}) + P_n. \end{aligned} \tag{4}$$

For any $\varepsilon > 0$, and by (iii), there is $k_0 \in \mathbb{N}$ (without loss of generality we may assume that $k_0 > m$) such that

$$(5) \quad \sum_{l=k_0}^{\infty} \|P_l - P_{l+1}\| < \varepsilon.$$

Therefore, from (4) and (5), we have

$$(6) \quad \|T^{k_0,n}P_{k_0+1} - P_n\| \leq \sum_{l=k_0}^{n-1} \|T^{l+1,n}(P_l - P_{l+1})\| \leq \sum_{l=k_0}^{\infty} \|P_l - P_{l+1}\| < \varepsilon.$$

For all $n \in \mathbb{N}$, and by (i), one finds

$$(7) \quad PT_n = PP_nT_n = PP_n = P,$$

which implies that $PT^{m,k_0-1} = P$, for all $m < k_0$, and hence $P(T^{m,k_0-1} - P_{k_0+1}) = 0$. So, (v) of Theorem (2.5) implies

$$(8) \quad \|T^{k_0,n}(T^{m,k_0-1} - P_{k_0+1})\| \leq \delta_P(T^{k_0,n})\|T^{m,k_0-1} - P_{k_0+1}\| \leq 2\delta_P(T^{k_0,n}).$$

Due to the weak P -ergodicity of $\{T^{k,n}\}$, there is $N_0 \in \mathbb{N}$ such that $\delta_P(T^{k_0,n}) < \varepsilon$ for all $n \geq N_0$.

From (6) and (8), we obtain

$$\begin{aligned} \|T^{m,n} - P_n\| &\leq \|T^{k_0,n}T^{m,k_0-1} - T^{k_0,n}P_{k_0+1}\| + \|T^{k_0,n}P_{k_0+1} - P_n\| \\ &= \|T^{k_0,n}(T^{m,k_0-1} - P_{k_0+1})\| + \|T^{k_0,n}P_{k_0+1} - P_n\| \\ &\leq 2\delta_P(T^{k_0,n}) + \varepsilon \\ &\leq 3\varepsilon, \quad \text{for all } n \geq N_0 \end{aligned}$$

hence, we deduce that $\|T^{m,n} - P\| \rightarrow 0$, as $n \rightarrow \infty$ which completes the proof. \square

Remark 4.4. We note that if the generating sequence $\{T_n\}$ is stationary (i.e. $T_n = T, \forall n \in \mathbb{N}$), then the corresponding NDMC reduces to a homogeneous chain. If for some projection P with $PT = TP = P$ and $P_n = P$, for all $n \in \mathbb{N}$, then the conditions of Theorem 4.3 are satisfied, hence we obtain the equivalence of the weak and uniform P -ergodicities for homogeneous Markov chain $\{T^n\}$. This result has been proven in [34] (namely by Corollary 4.7 and Proposition 4.10). However, it is known that even in the case of L^1 -spaces, the weak P -ergodicity does not imply uniform P -ergodicity for NDMC (see [35]).

Now, let us recall the following lemma which will be used to prove Theorem 4.6:

Lemma 4.5. [33] Let $\{a_{j,n}\}$ be a sequence of real numbers such that

$$\begin{aligned} 0 &\leq a_{j,n} \leq 1, \quad \text{for all } j, n \in \mathbb{N}, \\ a_{j,n} &\leq a_{j,m}a_{m+1,n} \quad \text{for all } j \leq m \leq n. \end{aligned}$$

If there is a constant $K > 0$ such that

$$\sum_{j=1}^n a_{j,n} \leq K \quad \text{for all } n \in \mathbb{N},$$

then for each j one has $a_{j,n} \rightarrow 0$ as $n \rightarrow \infty$.

Theorem 4.6. *Let (X, X_+, \mathcal{K}, f) be an abstract state space and $\{T_n\}$ be a generating sequence of the NDMC $\{T^{m,n}\}$. Let $\{P_n\}$ be a left decreasing sequence of projections of X such that $T_n P_n = P_n T_n = P_n$, for all $n \in \mathbb{N}$, and $P_n \rightarrow P$ in norm. If there exists a constant $C > 0$ with*

$$(9) \quad \sum_{l=1}^{n-1} \delta_P(T^{l,n}) \leq C, \quad \forall n \in \mathbb{N},$$

then $\{T^{k,n}\}$ is uniformly P -ergodic.

Proof. For $j, n \in \mathbb{N}$, put $a_{j,n} = \delta_P(T^{j,n})$, and using Equation (7), we have $PT_n = T_n P$. Then by Lemma 4.5 and the hypothesis (9), we find that $\{T^{k,n}\}$ is weakly P -ergodic.

As $P_n \rightarrow P$, for any $\varepsilon > 0$, there is $N_1 \in \mathbb{N}$ such that

$$(10) \quad \|P_n - P\| < \varepsilon \quad \text{for all } n \geq N_1.$$

One can see that

$$(11) \quad \|T^{m,n} - P\| \leq \|T^{m,n} - P_n\| + \|P_n - P\|.$$

Due to (10) it is enough to estimate $\|T^{m,n} - P_{n+1}\|$. By (7), one gets the following:

$$\begin{aligned} \|T^{m,n} - P_{n+1}\| &= \|T_n T^{m,n-1} - P_n\| \\ &\leq \|T_n T^{m,n-1} - T_n P_{n-1}\| + \|T_n P_{n-1} - P_n\| \\ &= \|T_{n+1} T_n T^{m,n-2} - T_n T_{n-1} P_{n-1}\| + \|T_n P_{n-1} - T_n P_n\| \\ &\leq \|T^{n-1,n} T^{m,n-2} - T^{n-1,n} P_{n-1}\| + \delta_P(T_n) \|P_{n-1} - P_n\| \\ &\leq \|T^{n-1,n} T^{m,n-2} - T^{n-1,n} P_{n-2}\| + \|T^{n-1,n} P_{n-2} - T^{n-1,n} P_{n-1}\| \\ &\quad + \delta_P(T_n) \|P_{n-1} - P_n\| \\ &\leq \|T^{n-2,n} T^{m,n-3} - T^{n-2,n} P_{n-2}\| + \delta_P(T^{n-1,n}) \|P_{n-2} - P_{n-1}\| \\ &\quad + \delta_P(T_n) \|P_{n-1} - P_n\| \\ &\quad \vdots \\ &\leq \|T_m - P_m\| \delta_P(T^{m+1,n}) + \sum_{\ell=m+1}^{n-1} \|P_{\ell-1} - P_\ell\| \delta_P(T^{\ell,n}). \end{aligned}$$

Due to $\delta_P(T^{m+1,n}) \rightarrow 0$, there exists an $N_2 \in \mathbb{N}$ such that $\delta_P(T^{m+1,n}) < \varepsilon$, for all $n \geq N_2$.

On other hand, from (10) we infer that $\|P_{n-1} - P_n\| < 2\varepsilon$ for all $n \geq N_1$. Without loss of generality, we may assume that $N_1 > m$. Hence,

$$\begin{aligned} \sum_{\ell=m+1}^{n-1} \|P_{\ell-1} - P_\ell\| \delta_P(T^{\ell,n}) &= \underbrace{\sum_{\ell=m+1}^{N_1} \|P_{\ell-1} - P_\ell\| \delta_P(T^{\ell,n})}_{I_1} \\ &\quad + \underbrace{\sum_{j=N_1+1}^{n-1} \|P_{j-1} - P_j\| \delta_P(T^{j,n})}_{I_2} \end{aligned} \quad (12)$$

Let us estimate I_1 and I_2 , separately.

We start with I_2 . From (9) we easily find

$$(13) \quad I_2 \leq 2\varepsilon \sum_{j=N_1+1}^{n-1} \delta_P(T^{j,n}) \leq 2\varepsilon C.$$

Now consider I_1 . For any $\ell \in \{m+1, \dots, N_1\}$ one has $\|P_{\ell-1} - P_\ell\| \leq 2$. The weakly P -ergodicity of $T^{m,n}$ implies the existence of an $N_3 \in \mathbb{N}$ such that $\delta_P(T^{N_1,n}) < \varepsilon$ for all $n \geq N_3$. Due to Equation (7) and by (vi) Theorem 2.5, one can see that

$$\delta_P(T^{\ell,n}) \leq \delta_P(T^{N_1,n}) < \varepsilon.$$

Therefore,

$$(14) \quad I_1 = \sum_{\ell=m+1}^{N_1} \|P_{\ell-1} - P_\ell\| \delta_P(T^{\ell,n}) \leq 2 \sum_{\ell=m+1}^{N_1} \delta_P(T^{N_1,n}) \leq 2(N_1 - m)\varepsilon.$$

Hence, from (10)-(14) we obtain

$$\|T^{m,n} - P\| \leq (3 + C + 2(N_1 - m))\varepsilon$$

for all $n \geq \max\{N_1, N_2, N_3\}$, which proves the assertion. \square

Theorem 4.7. *Let (X, X_+, \mathcal{K}, f) be an abstract state space and $\{T_n\}$ be a generating sequence of the NDMC. Let $\{P_n\}$ be a left decreasing sequence of projections of X such that $T_n P_n = P_n T_n = P_n, \forall n \in \mathbb{N}$. Assume that $\{T^{k,n}\}$ is uniformly P -ergodic for a given projection P and suppose there exist $k_n \in \mathbb{N}$ and $\gamma_n \in [0, 1)$ such that*

$$\delta_{P_n}(T_n^{k_n}) \leq \gamma_n \quad \text{with} \quad \sup_n \frac{k_n}{1 - \gamma_n} < \infty.$$

Then P_n converges to P in norm.

Proof. As $\{T^{m,n}\}$ is uniformly P -ergodic, for each $m \in \mathbb{N}$, we have $\|T^{m,n} - P\| \rightarrow 0$ as $n \rightarrow \infty$. Therefore, without loss of generality, we may assume $m = 1$. Define

$$(15) \quad E_n = T^{1,n} - T^{1,n-1}, \quad D_n = P_n - T^{1,n-1}.$$

One can see that

$$\begin{aligned} \|P_n - P\| &\leq \|P_n - T^{1,n-1}\| + \|T^{1,n-1} - P\| \\ &= \|D_n\| + \|T^{1,n-1} - P\| \end{aligned}$$

so it is enough to show that $\|D_n\| \rightarrow 0$ as $n \rightarrow \infty$, since $\|T^{1,n-1} - P\| \rightarrow 0$. Notice that

$$T_n D_n = T_n P_n - T_n T^{1,n-1} = P_n - T^{1,n} = D_n - E_n,$$

which yields

$$T_n(T_n D_n + E_n) = T_n(D_n - E_n + E_n) = T_n D_n = D_n - E_n.$$

Then

$$(16) \quad D_n = E_n + T_n^2 D_n + T_n E_n$$

Now iterating (16) N times, one gets

$$\begin{aligned}
 D_n &= E_n + T_n E_n + T_n^2 E_n + T_n^3 E_n + T_n^4 D_n \\
 &\vdots \\
 (17) \quad &= E_n + \sum_{j=1}^{2^N-1} T_n^j E_n + T_n^{2^N} D_n.
 \end{aligned}$$

As $\{P_n\}$ is left decreasing sequence and for every $n \in \mathbb{N}$, $P_n T_n = P_n$,

$$\begin{aligned}
 P_n T^{1,n} &= P_n T_n T_{n-1} \dots T_2 \\
 &= P_n T_{n-1} \dots T_2 \\
 &= P_n P_{n-1} T_{n-1} \dots T_2 \\
 &= P_n P_{n-1} T_{n-2} \dots T_2 \\
 &= P_n.
 \end{aligned}$$

Therefore,

$$P_n E_n = P_n (T^{1,n} - T^{1,n-1}) = P_n T^{1,n} - P_n T^{1,n-1} = P_n - P_n = 0.$$

Also, we have

$$P_n D_n = P_n - P_n T^{1,n-1} = 0.$$

Hence, by (17) and using (v) of Theorem 2.5, we obtain

$$\begin{aligned}
 \|D_n\| &\leq \|E_n\| + \sum_{j=1}^{2^N-1} \|T_n^j E_n\| + \|T_n^{2^N} D_n\| \\
 &\leq \|E_n\| + \sum_{j=1}^{2^N-1} \delta_{P_n}(T_n^j) \|E_n\| + \delta_{P_n}(T_n^{2^N}) \|D_n\| \\
 (18) \quad &\leq \|E_n\| \left(1 + \sum_{j=1}^{2^N-1} \delta_{P_n}(T_n^j) \right) + 2\delta_{P_n}(T_n^{2^N}).
 \end{aligned}$$

By $\delta_{P_n}(T_n^{k_n}) \leq \gamma_n < 1$, it follows that $\delta_{P_n}(T_n^{2^N}) < \varepsilon$ for a sufficiently large N . Moreover,

$$\begin{aligned}
 \sum_{j=1}^{2^N-1} \delta_{P_n}(T_n^j) &\leq \sum_{j=1}^{\infty} \delta_{P_n}(T_n^j) \\
 &= \sum_{j=1}^{k_n} \delta_{P_n}(T_n^j) + \sum_{j=k_n+1}^{2k_n} \delta_{P_n}(T_n^j) + \dots \\
 &\leq k_n + k_n \gamma_n + k_n \gamma_n^2 + \dots \\
 &= \frac{k_n}{1 - \gamma_n} \leq K,
 \end{aligned}$$

Then, using (18), we deduce

$$\|D_n\| \leq \|E_n\|(1 + K) + 2\varepsilon.$$

Now, according to $\|E_n\| \rightarrow 0$ as $n \rightarrow \infty$, one finds $\|D_n\| \rightarrow 0$, which completes the proof. \square

Remark 4.8. *We stress that these types of results are even new in the case of classical L^1 -spaces. Moreover, if one considers abstract state spaces associated with C^* -algebras, we get totally new sort of results which open new insight into the field of non-commutative probability.*

5. WEAK P -ERGODICITY AND THE DOEBLIN CONDITION

In this section, we are going to investigate the weak P -ergodicity of NDMC by means of an analogue of Doeblin's condition on abstract state spaces.

Let (X, X_+, \mathcal{K}, f) be an abstract state space and let P be a projection on X . Now we are going to provide an analogue of Doeblin's condition for NDMC associated with P [8, 31], as follows:

Condition \mathfrak{D}_P . We say that a NDMC $\{T^{k,n}\}$ defined on X satisfies *condition \mathfrak{D}_P* , if for every $k \in \mathbb{N}$, there exist $\lambda_k \in [0, 1]$ and $n_k \in \mathbb{N}$, and for every $x, y \in \mathcal{K}$, with $x - y \in N_P$, one can find $z_k^{xy} \in \mathcal{K}$, and $\varphi_{x,y}^k \in X_+$ with $\sup \|\varphi_{x,y}^k\| \leq \frac{\lambda_k}{2}$ such that

$$(19) \quad T^{k,k+n_k}x + \varphi_{x,y}^k \geq \lambda_k z_k^{xy} \quad \text{and} \quad T^{k,k+n_k}y + \varphi_{x,y}^k \geq \lambda_k z_k^{xy}.$$

The next result is the main result of this section.

Theorem 5.1. *Let (X, X_+, \mathcal{K}, f) be an abstract state space, P be a Markov projection on X , and $\{T_n\}$ be a generating sequence of the NDMC $\{T^{k,n}\}$. Assume that $T_n P = P T_n$, for all $n \in \mathbb{N}$. Then the following conditions are equivalent:*

- (i) *the chain $\{T^{k,n}\}$ is weakly P -ergodic;*
- (ii) *the chain $\{T^{k,n}\}$ satisfies the condition \mathfrak{D}_P with $\sum_{k=1}^{\infty} \lambda_k$ diverges;*

- (iii) *for each $k \in \mathbb{N}$ there exist $\mu_k \in [0, 1]$ and a number $n_k \in \mathbb{N}$ such that*

$$\delta_P(T^{k,k+n_k}) \leq \mu_k$$

with $\sum_{k=1}^{\infty} (1 - \mu_k)$ diverges.

Proof. (i) \Rightarrow (ii): Assume that $\delta_P(T^{k,n}) \rightarrow 0$ as $n \rightarrow \infty$. From the definition of δ_P , we have

$$\sup\{\|T^{k,n}x\|; \|x\| \neq 0 \text{ with } x \in N_P\} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

In particular, one finds

$$\sup\{\|T^{k,n}x - T^{k,n}y\|; |x - y| \neq 0 \text{ with } x - y \in N_P\} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Now fix $y_0 \in \mathcal{K}$. Then there exists $n_k \in \mathbb{N}$ such that for all $x \neq y$ with $x - y_0 \in N_P, y - y_0 \in N_P$ we have

$$\|T^{k,k+n_k}x - T^{k,k+n_k}y_0\| < \frac{1}{4}, \quad \|T^{k,k+n_k}y - T^{k,k+n_k}y_0\| < \frac{1}{4}.$$

By means of the decomposition

$$\begin{aligned} T^{k,k+n_k}x - T^{k,k+n_k}y_0 &= (T^{k,k+n_k}x - T^{k,k+n_k}y_0)_+ - (T^{k,k+n_k}x - T^{k,k+n_k}y_0)_- \\ T^{k,k+n_k}y - T^{k,k+n_k}y_0 &= (T^{k,k+n_k}y - T^{k,k+n_k}y_0)_+ - (T^{k,k+n_k}y - T^{k,k+n_k}y_0)_- \end{aligned}$$

let us define

$$\tilde{\varphi}_u^{(k)} := (T^{k,k+n_k}u - T^{k,k+n_k}y_0)_-, \text{ where } u \in \{x, y\}.$$

Put

$$\varphi_{xy}^{(k)} := \tilde{\varphi}_x^{(k)} + \tilde{\varphi}_y^{(k)}.$$

Then, for all $x, y \in \mathcal{K}$, with $x - y \in N_P$, we have $\|\varphi_{xy}^{(k)}\| \leq \frac{1}{2}$. Therefore,

$$\begin{aligned} T^{k,k+n_k}u + \varphi_{xy}^{(k)} &\geq T^{k,k+n_k}u + \tilde{\varphi}_u^{(k)}, \quad u \in \{x, y\} \\ &= T^{k,k+n_k}y_0 + T^{k,k+n_k}u - T^{k,k+n_k}y_0 + \tilde{\varphi}_u^{(k)} \\ &\geq T^{k,k+n_k}y_0. \end{aligned}$$

By defining $z_k^{xy} := T^{k,k+n_k}y_0$ and $\lambda_k = 1$, the condition \mathfrak{D}_P is obtained.

(ii) \Rightarrow (iii): For each $k \in \mathbb{N}$, and for every $x, y \in \mathcal{K}$ with $x - y \in N_P$, by the condition \mathfrak{D}_P , we have

$$T^{k,k+n_k}x + \varphi_{x,y}^k \geq \lambda_k z_k^{xy}, \quad T^{k,k+n_k}y + \varphi_{x,y}^k \geq \lambda_k z_k^{xy},$$

with $\|\varphi_{x,y}^k\| \leq \frac{\lambda_k}{2}$, $\forall x, y \in \mathcal{K}$. Then

$$\begin{aligned} \|T^{k,k+n_k}x + \varphi_{x,y}^k - \lambda_k z_k^{xy}\| &= f(T^{k,k+n_k}x + \varphi_{x,y}^k - \lambda_k z_k^{xy}) \\ &= 1 - (\lambda_k - f(\varphi_{x,y}^k)) \\ &\leq 1 - \frac{\lambda_k}{2}. \end{aligned}$$

Similarly, $\|T^{k,k+n_k}y + \varphi_{x,y}^k - \lambda_k z_k^{xy}\| \leq 1 - \frac{\lambda_k}{2}$. Let $c = \lambda_k - f(\varphi_{x,y}^k)$ and

$$\begin{aligned} x_1 &:= \frac{1}{1-c}(T^{k,k+n_k}x + \varphi_{x,y}^k - \lambda_k z_k^{xy}) \\ y_1 &:= \frac{1}{1-c}(T^{k,k+n_k}y + \varphi_{x,y}^k - \lambda_k z_k^{xy}). \end{aligned}$$

Then $x_1, y_1 \in \mathcal{K}$ and

$$\|T^{k,k+n_k}x - T^{k,k+n_k}y\| = (1-c)\|x_1 - y_1\| \leq 2(1-c),$$

which implies that

$$\delta_P(T^{k,k+n_k}) \leq 1 - \frac{\lambda_k}{2},$$

which proves (ii) by taking $\mu_k = 1 - \frac{\lambda_k}{2}$.

(iii) \Rightarrow (i): Given $k \in \mathbb{N}$, then there exists $\mu_k \in [0, 1)$ and $n_k \in \mathbb{N}$ such that

$$\delta_P(T^{k,k+n_k}) \leq \mu_k.$$

Let $l_1 = k + n_k$, by (ii), one finds $n_{l_1} \in \mathbb{N}$ and $\mu_{l_1} \in [0, 1)$ such that $\delta_P(T^{l_1, l_1+n_{l_1}}) \leq \mu_{l_1}$. Continuing in the same argument, there exists a sequence $\{l_j\}_{j=0}^\infty$, with $l_0 = k$ and $\mu_{l_j} \in [0, 1)$ such that

$$\delta_P(T^{l_j, l_j+n_{l_j}}) \leq \mu_{l_j}.$$

For a large n , we define L_n as

$$L_n := \max\{j; l_j + n_{l_j} \leq n\}.$$

Due to the hypothesis of the theorem, we have $T^{k,n}P = PT^{k,n}$, and then, from (vi) Theorem 2.5 one finds

$$\begin{aligned}\delta_P(T^{k,n}) &= \delta_P(T^{l_{L_n},n}T^{l_{L-1},l_L}, \dots T^{l_0,l_1}) \\ &\leq \prod_{j=1}^{L_n} \delta_P(T^{l_{L_n-j},l_{L_n-j+1}}) \\ &\leq \prod_{j=1}^{L_n} \mu_{l_j} \rightarrow 0 \text{ as } n \rightarrow \infty,\end{aligned}$$

since $\sum_{k=1}^{\infty} (1 - \mu_k)$ diverges, hence (i) holds. This completes the proof. \square

Remark 5.2. We point out that the condition $T_n P = P T_n$, $n \in \mathbb{N}$ is only used to establish the implication (iii) \Rightarrow (i). Hence, we conclude that the implications (i) \Rightarrow (ii) \Rightarrow (iii) are true without the stated condition.

Remark 5.3. It is also worth to mention that an analogous kind of results to Theorem 5.1 have been established in [18, 35, 37] in the setting of $X = \ell_1$ and P is a one dimensional projection.

As an application of the previous theorem, we have the following result.

Theorem 5.4. Let (X, X_+, \mathcal{K}, f) be an abstract state space, P be a Markov projection on X , and for every $n \in \mathbb{N}$ let $T_n \in \Sigma_P(X)$. Assume that $\|T_n - P\| < \epsilon_n$, where $\epsilon_n \rightarrow 0$. Then the NDMC $\{T^{m,n}\}$ is uniformly P -ergodic.

Proof. For every $n \in \mathbb{N}$, let $Q_n = T_n - P$. Then, $\|Q_n\| < \epsilon_n$ and $Q_n P = P Q_n$. For every $n \in \mathbb{N}$, one finds

$$\begin{aligned}\delta_P(T_n) &= \sup_{x \in N_P, \|x\|=1} \|T_n(x)\| \\ &= \sup_{x \in N_P, \|x\|=1} \|Q_n x + P x\| \\ &= \sup_{x \in N_P, \|x\|=1} \|Q_n x\| \\ &= \delta_P(Q_n) \\ &\leq \|Q_n\| < \epsilon_n.\end{aligned}$$

Also, we have

$$\begin{aligned}P T_n &= P(Q_n + P) \\ &= P^2 Q_n + P \\ &= P Q_n P + P^3 \\ &= P(Q_n + P)P \\ &= P T_n P.\end{aligned}$$

Now let us choose $n_k \in \mathbb{N}$ such that $\epsilon_{k+n_k} \leq \frac{1}{2^k}$. Then by Proposition 2.7,

$$\begin{aligned}\delta_P(T^{k,k+n_k}) &= \delta_P(T_{k+n_k} T_{k+n_k-1} \dots T_{k+1}) \\ &= \delta_P(T_{k+n_k}) \delta_P(T_{k+n_k-1}) \dots \delta_P(T_{k+1}) \\ &= \delta_P(T_{k+n_k}) \\ &\leq \epsilon_{k+n_k} \\ &\leq \frac{1}{2^k}.\end{aligned}$$

Letting $\mu_k = \frac{1}{2^k}$, and using Theorem 5.1, we infer that $\{T^{k,n}\}$ is weakly P -ergodic. Now, for every $n \in \mathbb{N}$, put $P_n = P$ and then the three conditions (i), (ii) and (iii) of Theorem 4.3 are satisfied which yields that $\{T^{m,n}\}$ is uniformly P -ergodic. \square

Now, we are going to construct a left decreasing sequence of projections $\{P_n\}$ on X which converges to a projection P and to construct a sequence $\{T_n\}$ of uniformly P_n -ergodic Markov operators on X such that the generated NDMC $\{T^{k,n}\}$ is uniformly P -ergodic.

Example 5.5. Consider the space ℓ_1 , the subspaces $\mathcal{A} = \{x \in \ell_1; x_{2n} = 0\}$ and the operator $P : \ell_1 \rightarrow \mathcal{A}$ defined by

$$P_1(x) = (x_1 + x_2, 0, x_3 + x_4, 0, x_5 + x_6, 0 \dots).$$

Then P is a projection on \mathcal{A} . Let $Q_1 : \ell_1 \rightarrow \ell_1$ be the operator defined by

$$Q_1(x) = \left(\frac{-x_2}{2}, \frac{x_2}{2}, \frac{-x_4}{2}, \frac{x_4}{2}, \frac{-x_6}{2}, \frac{x_6}{2}, \dots \right).$$

It is clear that $Q_1^n \rightarrow 0$, as $n \rightarrow \infty$ so for some $n_0 \in \mathbb{N}$, we have $\|Q_1^{n_0}\| < 1$. Also, $P_1 Q_1 = Q_1 P_1 = 0$. Then by Theorem 5.2 of [34], we have that the operator $T_1 = P_1 + Q_1$ is uniformly P_1 -ergodic. Similarly, define the operators P_2 and Q_2 on ℓ_1 by

$$P_2(x) = (0, 0, x_3 + x_4, 0, x_5 + x_6, 0, x_7 + x_8, 0 \dots), \text{ and } Q_2(x) = \frac{1}{2} Q_1(x).$$

One can see that $P_2 Q_2 = Q_2 P_2 = 0$ and $P_2 \leq^l P_1$, with $T_2 = P_2 + Q_2$ is uniformly P_2 -ergodic. Also, define the operators P_3 and Q_3 on ℓ_1 by

$$P_3(x) = (0, 0, 0, 0, x_5 + x_6, 0, x_7 + x_8, 0 \dots), \text{ and } Q_3(x) = \frac{1}{3} Q_1(x),$$

and having $P_3 Q_3 = Q_3 P_3 = 0$ and $P_3 \leq^l P_2$, with $T_3 = P_3 + Q_3$ is uniformly P_3 -ergodic. Fixing some $N_0 \in \mathbb{N}$ and using the same argument, for every n , $1 \leq n \leq N_0$, put

$$P_n(x) = (0, 0, \dots, 0, \overbrace{x_{2n-1} + x_{2n}}^{(2n-1)^{th}-place}, 0, x_{2n+1} + x_{2n+2}, 0 \dots),$$

and for $n > N_0$, put $P_n = P_{N_0}$. Then $\{P_n\}_{n=1}^\infty$ is a left decreasing sequence of projections on \mathcal{A} , which converges to $P = P_{N_0}$. For every $n \in \mathbb{N}$, define $Q_n = \frac{1}{n} Q_1$. It is clear that $Q_n^m \rightarrow 0$, as $m \rightarrow \infty$ and $\|Q_n\| < \frac{1}{n}$. Therefore, using Theorem 5.2 of [34], the operator $T_n = P_n + Q_n$ is uniformly P_n -ergodic, $\forall n \in \mathbb{N}$.

Considering the NHDC $\{T^{k,n}\}$ generated by T_n , noting $T_n \rightarrow P$, as $n \rightarrow \infty$, we have $\|T_n - P\| < \frac{1}{n}$, and hence by Theorem 5.4 we deduce that $\{T^{k,n}\}$ is uniformly P -ergodic.

6. WEAKLY P -ERGODICITY AND ITS APPLICATION TO PERTURBATIONS

Let (X, X_+, \mathcal{K}, f) be an abstract state space, T be a Markov operator on X and let P be a projection on X . In [34], we have proved that the homogenous Markov chain $\{T^n\}$ is uniformly P -ergodic if and only if $TP = P$ and $T = P + Q$, where Q is an operator on X such that $PQ = QP = 0$ and $\|Q^{n_0}\| < 1$, for some $n_0 \in \mathbb{N}$. Moreover,

$$\delta_P(T) \leq \|Q\| \leq 2\delta_P(T).$$

Let $\{P_n\}$ be a sequence of projections of X . Then a NDMC $\{T^{m,n}\}$ is called *weakly ergodic w.r.t $\{P_n\}$* if for every $m > 0$ one has

$$\lim_{n \rightarrow \infty} \delta_{P_{m+1}}(T^{m,n}) = 0.$$

In this section, we are going to establish similar kind of results for NDMC. Before, to formulate the main result of this section, one needs some auxiliary facts.

Lemma 6.1. *Let (X, X_+, \mathcal{K}, f) be an abstract state space and $\{T_n\}$ be a generating sequence of NDMC $\{T^{k,n}\}$. Let $\{P_n\}$ be a left decreasing sequence of projections of X such that $P_n T_n = P_n, \forall n \in \mathbb{N}$. If $\{T^{m,n}\}$ is weakly ergodic w.r.t $\{P_n\}$, then*

$$\delta_{P_l}(T^{m,n}) \rightarrow 0, \text{ as } n \rightarrow \infty, \text{ for all } l \geq m+1.$$

Proof. The condition $P_n T_n = P_n$ ($\forall n \in \mathbb{N}$) together with the left consistency of $\{P_n\}$, one finds $P_k T_n = P_k$, for all $k \geq n$. Moreover,

$$\begin{aligned} P_{m+l+1} T^{m,m+l-1} &= P_{m+l+1} (T_{m+l-1} T_{m+l-2} \dots T_{m+1}) \\ &= P_{m+l+1} T_{m+l-2} \dots T_{m+1} \\ &= P_{m+l+1}. \end{aligned}$$

Therefore, by Remark 2.8 and Proposition 2.7, we have

$$\begin{aligned} \delta_{P_{m+l+1}}(T^{m,n}) &= \delta_{P_{m+l+1}}(T^{m+l,n} T^{m,m+l-1}) \\ &\leq \delta_{P_{m+l+1}}(T^{m+l,n}) \delta_{P_{m+l+1}}(T^{m,m+l-1}) \\ &\leq \delta_{P_{m+l+1}}(T^{m+l,n}) \rightarrow 0 \text{ as } n \rightarrow \infty \end{aligned}$$

hence the lemma is proved. \square

Lemma 6.2. *Let (X, X_+, \mathcal{K}, f) be an abstract state space and T be an operator on X . If $\{P_n\}$ is a sequence of projections such that $P_n \rightarrow P$ in norm, then $\delta_{P_n}(T) \rightarrow \delta_P(T)$.*

Proof. Given $\epsilon > 0$, let $y = x - Px, \forall n \in \mathbb{N}$, let $y_n = x - P_n x$ with $\|y_n\| \leq 1$. Then $y \in N_P, y_n \in N_{P_n}$ and $\|y\| \leq 1$. As T is continuous, there exists $N_0 \in \mathbb{N}$ such that

$$\|Ty_n - Ty\| < \epsilon; \forall n > N_0.$$

Therefore, $\forall n > N_0$ we have

$$\|Ty_n\| < \epsilon + \|Ty\| \leq \delta_P(T) + \epsilon$$

and then $\delta_{P_n}(T) \leq \delta_P(T) + \epsilon$, which implies that

$$|\delta_{P_n}(T) - \delta_P(T)| \leq \epsilon; \forall n \geq N_0$$

this yields the required assertion. \square

Now, we are ready to formulate the main result of this section.

Theorem 6.3. *Let (X, X_+, \mathcal{K}, f) be an abstract state space, $\{P_n\}$ be a left decreasing sequence of Markov projections of X and $\{T_n\}$ be a generation sequence of Markov operators of NDMC $\{T^{m,n}\}$. Let $T_n \in \Sigma_{P_n}(X)$, for all $n \in \mathbb{N}$ with $T_n P_n = P_n$ and assume that $T_n = P_n + Q_n$. Then the following statements are equivalent:*

- (i) *for every $k \in \mathbb{N}$, $\{T^{m,n}\}$ is weakly P_k -ergodic;*
- (ii) *$\{T^{m,n}\}$ is weakly ergodic w.r.t $\{P_n\}$;*
- (iii) *for every $m \in \mathbb{N}$ one has*

$$\|Q_{m+n} \dots Q_{m+1}\| \rightarrow 0, \text{ as } n \rightarrow \infty.$$

Moreover,

$$\delta_{P_{m+1}}(T^{m,m+n}) \leq \|Q_{m+n} \dots Q_{m+1}\| \leq 2\delta_{P_{m+2}}(T^{m+1,m+n}).$$

Moreover, if $P_n \rightarrow P$ (in norm), then the above statements are equivalent to

- (iv) *$\{T^{m,n}\}$ is weakly P -ergodic.*

Proof. (i) \Rightarrow (ii): Obvious as by the assumption $\{T^{m,n}\}$ is weakly P_{m+1} -ergodic.
(ii) \Rightarrow (i): Let us consider the following two cases: (a) For $0 < k \leq m+1$, we have $P_{m+1} \leq^\ell P_k$ and by Proposition 3.3, we get

$$\delta_{P_k}(T^{m,n}) \leq \delta_{P_{m+1}}(T^{m,n}),$$

which tends to zero by the assumption. (b) For $k > m+1$, and using Lemma 6.1, we have that

$$\delta_{P_k}(T^{m,n}) \rightarrow 0, \text{ as } n \rightarrow \infty,$$

hence we deduced (i).

To establish the implication (ii) \Rightarrow (iii), let us first prove the following equality

$$(20) \quad Q_{m+n} \dots Q_{m+1} = T^{m,m+n} - T^{m+1,m+n} P_{m+1}.$$

by induction on n .

For $n = 2$, by noticing that

$$P_{m+2}T_{m+1} = P_{m+2}P_{m+1}T_{m+1} = P_{m+2}P_{m+1} = P_{m+2}.$$

we find

$$\begin{aligned} Q_{m+2}Q_{m+1} &= (T_{m+2} - P_{m+2})(T_{m+1} - P_{m+1}) \\ &= T_{m+2}T_{m+1} - T_{m+2}P_{m+1} - P_{m+2}T_{m+1} + P_{m+2}P_{m+1} \\ &= T_{m+2}T_{m+1} - T_{m+2}P_{m+1} \\ &= T^{m,m+2} - T^{m+1,m+2}P_{m+1}. \end{aligned}$$

Now assume the statement is true for $k = n$ and let us establish it for $k = n+1$:
First notice that

$$\begin{aligned} P_{m+n+1}T^{m,m+n} &= P_{m+n+1}T_{m+n} \dots T_{m+1} \\ &= P_{m+n+1}P_{m+n}T_{m+n} \dots T_{m+1} \\ &= P_{m+n+1}P_{m+n}T_{m+n-1} \dots T_{m+1} \\ &= P_{m+n+1}T_{m+n-1} \dots T_{m+1} \\ &\vdots \\ &= P_{m+n+1}. \end{aligned}$$

Similarly,

$$P_{m+n+1}T^{m+1,m+n}P_{m+1} = P_{m+n+1}.$$

Therefore,

$$\begin{aligned} Q_{m+n+1} \dots Q_{m+1} &= Q_{m+n+1}(Q_{m+n} \dots Q_{m+1}) \\ &= Q_{m+n+1}Q_{m+n}(T^{m,m+n} - T^{m+1,m+n}P_{m+1}) \\ &= T_{m+n+1}T^{m,m+n} - T_{m+n+1}T^{m+1,m+n}P_{m+1} \\ &\quad - P_{m+n+1}T^{m,m+n} + P_{m+n+1}T^{m+1,m+n}P_{m+1} \\ &= T^{m,m+n+1} - T^{m+1,m+n+1}P_{m+1}, \end{aligned}$$

which yields Equation (20).

(ii) \Rightarrow (iii): By (20), one finds

$$\begin{aligned}
\|Q_{m+n} \cdots Q_{m+1}\| &= \|T^{m,m+n} - T^{m+1,m+n}P_{m+1}\| \\
&= \|T^{m+1,m+n}T_{m+1} - T^{m+1,m+n}P_{m+1}\| \\
&= \|T^{m+1,m+n}(T_{m+1} - P_{m+1})\| \\
&\leq \delta_{P_{m+1}}(T^{m+1,m+n})\|T_{m+1} - P_{m+1}\| \\
&\leq 2\delta_{P_{m+1}}(T^{m+1,m+n}) \text{ (as both are Markov operators)} \\
&\leq 2\delta_{P_{m+2}}(T^{m+1,m+n}) \text{ (using Proposition 3.3),}
\end{aligned}$$

which tends to 0 by (ii).

(iii) \Rightarrow (ii): Again using (20), we have

$$T^{m,m+n} = \prod_{j=0}^{n-1} Q_{m+n-j} + T^{m+1,m+n}P_{m+1},$$

then

$$\begin{aligned}
\delta_{P_{m+1}}(T^{m,m+n}) &= \sup_{x \in N_{P_{m+1}}} \left\| \prod_{j=0}^{n-1} Q_{m+n-j}(x) + T^{m+1,m+n}P_{m+1}(x) \right\| \\
&= \sup_{x \in N_{P_{m+1}}} \left\| \prod_{j=0}^{n-1} Q_{m+n-j}(x) \right\| \\
&= \delta_{P_{m+1}}\left(\prod_{j=0}^{n-1} Q_{m+n-j}\right) \\
&\leq \left\| \prod_{j=0}^{n-1} Q_{m+n-j} \right\| \rightarrow 0, \text{ as } n \rightarrow \infty \text{ by (iii).}
\end{aligned}$$

(iv) \Rightarrow (ii): As $\{P_n\}$ is a left decreasing sequence and $P_n \rightarrow P$, by (ii) of Lemma 3.6, one concludes that P is a projections and $P \leq^\ell P_n$ for all $n \in \mathbb{N}$. Therefore, due to Proposition 3.3 one gets

$$\delta_{P_{m+1}}(T^{m,n}) \leq \delta_P(T^{m,n}) \rightarrow 0, \text{ as } n \rightarrow \infty.$$

(ii) \Rightarrow (iv): Assume that $\delta_{P_{m+1}}(T^{m,n}) \rightarrow 0$ as $n \rightarrow \infty$, then by Lemma 6.1, for all $l \geq m+1$ we have that

$$\delta_{P_l}(T^{m,n}) \rightarrow 0, \text{ as } n \rightarrow \infty.$$

Letting $l \rightarrow \infty$ and using Lemma 6.2, one gets

$$\delta_P(T^{m,n}) \rightarrow 0, \text{ as } n \rightarrow \infty$$

which proves (iv), and hence the proof is completed. \square

Example 6.4. Consider the space ℓ_1 and for every $n \in \mathbb{N}$ let

$$e_n = (\underbrace{0, 0, \dots, 1}_n, 0, \dots).$$

Construct the one dimensional projections $P_n := T_{e_n}$. Then $\{P_n\}$ is a left decreasing sequence of projections (see Example 3.4). Take any $r \in (0, 1/2)$, and for every $n \in \mathbb{N}$, define $T_n := P_n + Q_n$, where $Q_n = r(I - P_n)$. Therefore,

$$\|Q_{m+n} \cdots Q_{m+1}\| = r^n \|(I - P_{m+n}) \cdots (I - P_{m+1})\| = (2r)^n \rightarrow 0, \text{ as } n \rightarrow \infty.$$

Hence by Theorem 6.3, the generated chain $\{T^{m,n}\}$ is weakly ergodic w.r.t. $\{P_n\}$.

Next, we will discuss a perturbations of weakly and uniformly P -ergodic chains. We need the following auxiliary fact.

Lemma 6.5. *Let (X, X_+, \mathcal{K}, f) be an abstract state space and let $\{T_n\}, \{S_n\}$ be two generating sequences of NDMCs $\{T^{k,n}\}, \{S^{k,n}\}$ on X , respectively. Assume that*

$$\sum_{n=1}^{\infty} \|T_n - S_n\| < \infty,$$

then for any $\epsilon > 0$, there exists $m_0 \in \mathbb{N}$ such that

$$\|T^{m,n} - S^{m,n}\| < \epsilon; \quad \forall m \geq m_0, \quad \forall n > m.$$

Proof. For every $n \in \mathbb{N}$, let $R_n = T_n - S_n$, and put $r_n = \|R_n\|$. Then

$$\begin{aligned} T^{m,n} &= \prod_{j=0}^{n-m-1} T_{n-j} \\ &= \prod_{j=0}^{n-m-1} (S_{n-j} + R_{n-j}) \\ &= \prod_{j=0}^{n-m-1} S_{n-j} + R_{m,n} = S^{m,n} + R_{m,n}, \end{aligned}$$

where $R_{m,n}$ contains all possible products of S_i and R_i and keeping in mind $\|S_i\| = 1$ for all i , one has

$$\begin{aligned} \|R_{m,n}\| &\leq \sum_i r_i + \sum_{i,j} r_i r_j + \sum_{i,j,k} r_i r_j r_k + \cdots + \prod_{i=m+1}^n r_i \\ &= \prod_{i=m+1}^n (1 + r_i) - 1. \end{aligned}$$

As $\sum_i r_i < \infty$, the product $\prod_{i=1}^n (1 + r_i)$ converges. Hence, for any $\epsilon > 0$, there exists $m_0 \in \mathbb{N}$ such that $\|R_{m,n}\| < \epsilon, \forall m \geq m_0$ and $\forall n > m$, which completes the proof. \square

Next result is about perturbations of weakly and uniformly P -ergodicities of NDMC.

Theorem 6.6. *Let (X, X_+, \mathcal{K}, f) be an abstract state space, $\{P_n\}$ be a left decreasing sequence of Markov projections of X which converges to P (in norm). Let $\{T_n\}, \{S_n\}$ be two generating sequences of NDMCs $\{T^{k,n}\}, \{S^{k,n}\}$ on X , respectively. Assume that $\forall n \in \mathbb{N}, T_n, S_n \in \Sigma_{P_n}(X)$ with $T_n P_n = P_n, S_n P_n = P_n$ and $\sum_{n=1}^{\infty} \|T_n - S_n\| < \infty$. Then the following statements hold:*

- (i) $\{T^{m,n}\}$ is uniformly P -ergodic if and only if $\{S^{m,n}\}$ is uniformly P -ergodic;
- (ii) $\{T^{m,n}\}$ is weakly P -ergodic if and only if $\{S^{m,n}\}$ is weakly P -ergodic.

Proof. (i). Given any $\epsilon > 0$. As $\{T^{k,n}\}$ is uniformly P -ergodic, there is $N_0 \in \mathbb{N}$ such that

$$\|T^{k,n} - P\| < \epsilon, \quad n \geq N_0.$$

On the other hand, by Proposition 6.5, there exists $m_0 \in \mathbb{N}$ such that

$$(21) \quad \|T^{k,n} - S^{k,n}\| < \epsilon, \quad \forall k \geq m_0, \quad \forall n > k.$$

Therefore, if $k \geq m_0$, then one finds

$$(22) \quad \|S^{k,n} - P\| \leq \|T^{k,n} - S^{k,n}\| + \|T^{k,n} - P\| < 2\varepsilon \quad \forall n > N_0.$$

Now, if $0 \leq k \leq m_0 - 1$, then by noticing $PS^{k,n} = P$ and (21), we find

$$\begin{aligned} \|S^{k,n} - P\| &= \|S^{m_0,n} S^{k,m_0-1} - PS^{k,m_0-1}\| \\ &= \|(S^{m_0,n} - P)S^{k,m_0-1}\| \\ &\leq \|S^{m_0,n} - P\| < 2\varepsilon \quad \forall n > N_0. \end{aligned}$$

This shows that $\{S^{m,n}\}$ is uniformly P -ergodic. The reverse can be proved by the same argument.

(ii). Assume that $\{T^{m,n}\}$ is weakly P -ergodic. For every $n \in \mathbb{N}$, let us denote $Q_n = T_n - P_n$ and $\tilde{Q}_n = S_n - P_n$. By Theorem 6.3, the weak P -ergodicity of $\{T^{m,n}\}$ implies

$$(23) \quad \|Q_{m+n} \dots Q_{m+1}\| \rightarrow 0, \text{ as } n \rightarrow \infty; \quad \forall m \in \mathbb{N}.$$

Let us establish

$$(24) \quad \|\tilde{Q}_{m+n} \dots \tilde{Q}_{m+1}\| \rightarrow 0, \text{ as } n \rightarrow \infty; \quad \forall m \in \mathbb{N}.$$

Using (20), we have

$$\begin{aligned} Q_{m+n} \dots Q_{m+1} &= T^{m,m+n} - T^{m+1,m+n} P_{m+1}, \\ \tilde{Q}_{m+n} \dots \tilde{Q}_{m+1} &= S^{m,m+n} - S^{m+1,m+n} P_{m+1}. \end{aligned}$$

According to the weak P -ergodicity of $\{T^{k,n}\}$, there is $N_1 \in \mathbb{N}$ such that

$$\delta_P(T^{m_0,m+n}) < \varepsilon, \quad \forall n \geq N_1.$$

Hence, by (21) one finds

$$\begin{aligned} \|Q_{m+n} \dots Q_{m+1} - \tilde{Q}_{m+n} \dots \tilde{Q}_{m+1}\| &\leq \|T^{m,m+n} - S^{m,m+n}\| \\ &\quad + \|T^{m+1,m+n} - S^{m+1,m+n}\| \\ &\leq \|T^{m_0,m+n} T^{m,m_0-1} - T^{m_0,m+n} S^{m,m_0-1}\| \\ &\quad + \|S^{m_0,m+n} S^{m,m_0-1} - T^{m_0,m+n} S^{m,m_0-1}\| \\ &\quad + \|T^{m_0,m+n} T^{m+1,m_0-1} - T^{m_0,m+n} S^{m+1,m_0-1}\| \\ &\quad + \|S^{m_0,m+n} S^{m+1,m_0-1} - T^{m_0,m+n} S^{m+1,m_0-1}\| \\ &\leq \|T^{m_0,m+n} (T^{m,m_0-1} - S^{m,m_0-1})\| \\ &\quad + \|(S^{m_0,m+n} - T^{m_0,m+n}) S^{m,m_0-1}\| \\ &\quad + \|T^{m_0,m+n} (T^{m+1,m_0-1} - S^{m+1,m_0-1})\| \\ &\quad + \|(S^{m_0,m+n} - T^{m_0,m+n}) S^{m+1,m_0-1}\| \\ &\leq 4\delta_P(T^{m_0,m+n}) + 2\|S^{m_0,m+n} - T^{m_0,m+n}\| \\ &< 6\varepsilon, \quad \text{for all } n > \max\{N_0, N_1\}. \end{aligned}$$

Consequently, by (23) we arrive at (24), which yields that $\{S^{m,n}\}$ is weakly P -ergodic. \square

Remark 6.7. We stress that if one considers, as a particular case, the left consistence sequence $\{P_n\}$ defined in Example 3.4, then the results of Theorem 6.6 is even new in the non-commutative setting. When $X = L^1$ and P is a one dimensional projection, in the mentioned setting, an analogue of Theorem 6.6 is known as the classical result [35].

Let us provide some examples as application of the proved theorem.

Example 6.8. Let (X, X_+, \mathcal{K}, f) be an abstract state space and $\{P_n\}$ be a left decreasing sequence of Markov projections of X which converges to P (in norm). Assume that $\{T^{k,n}\}$ be weakly (resp. uniformly) P -ergodic NDMC which is generated by a sequence of Markov operators $\{T_n\}_{n=1}^\infty$ such that $T_n P_n = P_n T_n = P_n, \forall n \in \mathbb{N}$. Let $\{\lambda_n\} \subset (0, 1)$ such that

$$(25) \quad \sum_{n=1}^{\infty} (1 - \lambda_n) < \infty.$$

Let us define a sequence $\{S_n\}$ of Markov operators by

$$S_n = (1 - \lambda_n)P_n + \lambda_n T_n.$$

It is clear that $S_n P_n = P_n S_n = P_n$, and we have

$$\|S_n - T_n\| = |1 - \lambda_n| \|P_n - T_n\| \leq 2(1 - \lambda_n)$$

which due to (25) implies

$$\sum_{n=1}^{\infty} \|S_n - T_n\| < \infty.$$

Hence, by Theorem 6.6 we conclude that the NDMC $\{S^{k,n}\}$ is weakly (resp. uniformly) P -ergodic.

Now, let us provide a concrete example of uniformly P -ergodic NDMC.

Example 6.9. Let (X, X_+, \mathcal{K}, f) be an abstract state space, and let $\{P_n\}$ be a left decreasing sequence of Markov projections of X which converges to P (in norm). Assume that $\{\tilde{T}_n\}_{n=1}^\infty$ is a sequence of Markov operators such that $\tilde{T}_n P_n = P_n \tilde{T}_n = P_n$. Let $\{\alpha_n\}_{n=1}^\infty \subseteq (0, 1)$ be such that

$$(26) \quad \sum_{n=1}^{\infty} (1 - \alpha_n) \text{ diverges.}$$

Let us define the sequence $\{T_n\}$ by

$$T_n = (1 - \alpha_n)P + \alpha_n \tilde{T}_n.$$

Then, we have

$$\begin{aligned} T_n T_m &= ((1 - \alpha_n)P + \alpha_n \tilde{T}_n)((1 - \alpha_m)P + \alpha_m \tilde{T}_m) \\ &= (1 - \alpha_n)(1 - \alpha_m)P + \alpha_m(1 - \alpha_n)P \tilde{T}_m + \alpha_n(1 - \alpha_m)\tilde{T}_m P + \alpha_n \alpha_m \tilde{T}_n \tilde{T}_m \\ &= (1 - \alpha_n \alpha_m)P + \alpha_n \alpha_m \tilde{T}_n \tilde{T}_m. \end{aligned}$$

Hence $\forall k < n$

$$T^{k,n} = T_n T_{n-1} \dots T_{k+1} = \left(1 - \prod_{j=k+1}^n \alpha_j\right)P + \left(\prod_{j=k+1}^n \alpha_j\right)\tilde{T}_n \dots \tilde{T}_{k+1}.$$

Therefore, due to (26), one finds

$$\|T^{k,n} - P\| \leq \left(\prod_{j=k+1}^n \alpha_j \right) \|P - \tilde{T}_n \dots \tilde{T}_{k+1}\| \leq 2 \prod_{j=k+1}^n \alpha_j \rightarrow 0,$$

as $n \rightarrow \infty$, which shows that $\{T^{k,n}\}$ is uniformly P -ergodic.

Assume that a sequence $\{\beta_n\}_{n=1}^\infty \subseteq (0, 1)$ satisfies

$$(27) \quad \sum_{n=1}^{\infty} |\beta_n - \alpha_n| < \infty.$$

It is obvious that this together with (26) yields that

$$\sum_{n=1}^{\infty} (1 - \beta_n) \text{ diverges.}$$

Due to $P_n \rightarrow P$, we suppose (if it is needed by taking subsequence) that

$$(28) \quad \|P_n - P\| \leq \frac{1}{(1 - \beta_n)n^{1+\gamma}}, \text{ for some } \gamma > 0.$$

Now, let us define the sequence $\{S_n\}$ as follows:

$$S_n := (1 - \beta_n)P_n + \beta_n \tilde{T}_n.$$

It is clear that $\forall n \in \mathbb{N}$, the operator S_n is Markov and $S_n P_n = P_n S_n = P_n$. Moreover,

$$\begin{aligned} S_n - T_n &= (P_n - P) + (\alpha_n P - \beta_n P_n) + (\beta_n - \alpha_n) \tilde{T}_n \\ &= (P_n - P) + (\alpha_n P - \beta_n P) + (\beta_n P - \beta_n P_n) + (\beta_n - \alpha_n) \tilde{T}_n \\ &= (1 - \beta_n)(P_n - P) + (\alpha_n - \beta_n)(P - T_n) \end{aligned}$$

and due to (27), (28), we infer that

$$\begin{aligned} \sum_{n=1}^{\infty} \|S_n - T_n\| &\leq \sum_{n=1}^{\infty} (1 - \beta_n) \|P_n - P\| + \sum_{n=1}^{\infty} |\alpha_n - \beta_n| \|P - T_n\| \\ &\leq \sum_{n=1}^{\infty} \frac{1}{n^{1+\gamma}} + 2 \sum_{n=1}^{\infty} |\alpha_n - \beta_n| < \infty. \end{aligned}$$

Hence, by Theorem 6.6 we conclude that the NDMC $\{S^{k,n}\}$ associated with $\{S_n\}$ is uniformly P -ergodic.

The above theorem can be generalized as follows:

Theorem 6.10. Let (X, X_+, \mathcal{K}, f) be an abstract state space $\{P_n\}$ and $\{\bar{P}_n\}$ be left decreasing sequences of Markov projections of X such that $\|P_n - \bar{P}_n\| \rightarrow 0$, and $P_n \rightarrow P$ as $n \rightarrow \infty$. Suppose that the generating sequences $\{T_n\}_{n=1}^\infty \subseteq \Sigma_{P_n}(X)$, $\{S_n\}_{n=1}^\infty \subseteq \Sigma_{\bar{P}_n}(X)$ such that $T_n P_n = P_n$, $S_n \bar{P}_n = \bar{P}_n$ and $\sum_{n=1}^\infty \|T_n - S_n\| < \infty$. Then $\{T^{m,n}\}$ is weakly P -ergodic if and only if $\{S^{m,n}\}$ is weakly P -ergodic.

Proof. As $\|P_n - \bar{P}_n\| \rightarrow 0$ and $P_n \rightarrow P$, we have $\bar{P}_n \rightarrow P$. For every $n \in \mathbb{N}$, let us denote $Q_n = T_n - P_n$ and $\tilde{Q}_n = S_n - \bar{P}_n$. Using (20), we have

$$\begin{aligned} Q_{m+n} \dots Q_{m+1} &= T^{m,m+n} - T^{m+1,m+n} P_{m+1}, \\ \tilde{Q}_{m+n} \dots \tilde{Q}_{m+1} &= S^{m,m+n} - S^{m+1,m+n} \bar{P}_{m+1}. \end{aligned}$$

Now, given any $\epsilon > 0$, by Proposition 6.5, there exists $m_0 \in \mathbb{N}$ such that

$$\|T^{m,n} - S^{m,n}\| < \epsilon; \quad \forall m \geq m_0, \quad \forall n > m.$$

Also, there exists $m_1 \in \mathbb{N}$ such that

$$\|P_m - \bar{P}_m\| < \epsilon, \quad \forall m \geq m_1.$$

Therefore,

$$\begin{aligned} \|Q_{m+n} \cdots Q_{m+1} - \tilde{Q}_{m+n} \cdots \tilde{Q}_{m+1}\| &\leq \|T^{m,m+n} - S^{m,m+n}\| \\ &\quad + \|T^{m+1,m+n} P_{m+1} - S^{m+1,m+n} \bar{P}_{m+1}\| \\ &\leq \|T^{m,m+n} - S^{m,m+n}\| \\ &\quad + \|T^{m+1,m+n} - S^{m+1,m+n}\| \\ &\quad + \|P_{m+1} - \bar{P}_{m+1}\|. \end{aligned}$$

Repeating the argument of the proof of Theorem 6.6, we obtain the desired assertion. \square

7. BAIR CATEGORY RESULTS

In this section, we are going to prove Baire category results to describe the size of uniformly P -ergodic and weakly ergodic w.r.t. $\{P_n\}$ NDMC's.

In the sequel, we will identify the generating sequence $\mathcal{T} = \{T_n\}_{n=1}^\infty$ of Markov operators with the corresponding NDMC $\{T^{k,n}\}$.

Let (X, X_+, \mathcal{K}, f) be an abstract state space, and let $\{P_n\}$ be a left decreasing sequence of Markov projections of X . Let us define

$$\begin{aligned} \mathfrak{S}_{\{P_n\}}(X) &:= \{\mathcal{T} = \{T_n\}_{n=1}^\infty : T_n \in \Sigma_{P_n}(X) \text{ and } T_n P_n = P_n, \forall n \in \mathbb{N}\}, \\ \mathfrak{S}_{\{P_n\}}^w(X) &:= \{\mathcal{T} \in \mathfrak{S}_{\{P_n\}}(X) : \{T^{m,n}\} \text{ is weakly ergodic w.r.t. } \{P_n\}\}. \end{aligned}$$

If P is a any projection of X , then we define:

$$\begin{aligned} \mathfrak{S}_P(X) &:= \{\mathcal{T} = \{T_n\}_{n=1}^\infty : T_n \in \Sigma_P(X) \text{ and } T_n P = P, \forall n \in \mathbb{N}\}, \\ \mathfrak{S}_P^u(X) &:= \{\mathcal{T} \in \mathfrak{S}_P(X) : \{T^{m,n}\} \text{ is uniformly } P\text{-ergodic}\}. \end{aligned}$$

The sets $\mathfrak{S}_{\{P_n\}}(X)$ and $\mathfrak{S}_P(X)$ are endowed with the following metric

$$d(\mathcal{T}, \mathcal{S}) := \sum_{n=1}^{\infty} \frac{\|T_n - S_n\|}{2^n}, \quad \mathcal{T} = \{T_n\}, \mathcal{S} = \{S_n\}.$$

Then let us prove the following density theorems.

Theorem 7.1. *Let (X, X_+, \mathcal{K}, f) be an abstract state space, and let P be a Markov projection of X . Then $\mathfrak{S}_P^u(X)$ is a dense subset of $\mathfrak{S}_P(X)$.*

Proof. Let $\epsilon > 0$ and let $\mathcal{T} \in \mathfrak{S}_P(X)$. For each $n \in \mathbb{N}$, put $S_n = \frac{\epsilon}{2}P + (1 - \frac{\epsilon}{2})T_n$. Clearly, $\{S_n\}$ is a sequence of Markov operators on X satisfying: $S_n P = P S_n = P$, and $\mathcal{S} = \{S_n\} \in \mathfrak{S}_P(X)$. Moreover, one has

$$d(\mathcal{S}, \mathcal{T}) = \sum_{n=1}^{\infty} \frac{\|S_n - T_n\|}{2^n} = \frac{\epsilon}{2} \sum_{n=1}^{\infty} \frac{\|P - T_n\|}{2^n} < \epsilon.$$

Now, we claim that the generated NHMC $\{S^{k,n}\}$ is uniformly P -ergodic. Notice that

$$S^{k,n} = S_n \cdots S_{k+1} = \left(1 - \left(1 - \frac{\epsilon}{2}\right)^{n-k}\right)P + \left(1 - \frac{\epsilon}{2}\right)^{n-k} T_n \cdots T_{k+1}.$$

Hence, we have

$$\begin{aligned}\|S^{k,n} - P\| &\leq \left(1 - \frac{\epsilon}{2}\right)^{n-k} \|P - T_n \dots T_{k+1}\| \\ &\leq 2 \left(1 - \frac{\epsilon}{2}\right)^{n-k} \rightarrow 0,\end{aligned}$$

as $n \rightarrow \infty$. So, we arrive at $\mathcal{S} \in \mathfrak{S}_P^u(X)$ which completes the proof. \square

Theorem 7.2. *Let (X, X_+, \mathcal{K}, f) be an abstract state space, and let $\{P_n\}$ be a left decreasing sequence of Markov projections of X . Then $\mathfrak{S}_{\{P_n\}}^w(X)$ is a dense G_δ -subset of $\mathfrak{S}_{\{P_n\}}(X)$.*

Proof. Let $\epsilon > 0$ and let $\mathcal{T} = \{T_n\} \in \mathfrak{S}_{\{P_n\}}(X)$. For each $n \in \mathbb{N}$, we define

$$S_n = \frac{\epsilon}{2}P_n + (1 - \frac{\epsilon}{2})T_n.$$

As in Example 6.8, $\{S_n\}$ is a sequence of Markov operators on X satisfying: $S_n P_n = P_n S_n = P_n$, and $\mathcal{S} = \{S_n\} \in \mathfrak{S}_{\{P_n\}}(X)$. Moreover, one has

$$d(\mathcal{S}, \mathcal{T}) = \sum_{n=1}^{\infty} \frac{\|S_n - T_n\|}{2^n} = \frac{\epsilon}{2} \sum_{n=1}^{\infty} \frac{\|P_n - T_n\|}{2^n} < \epsilon.$$

For every x and y with $x - y \in N_{P_n}$, (for all $n \in \mathbb{N}$) we have

$$\begin{aligned}\|S_n x - S_n y\| &= \left\| \frac{\epsilon}{2}P_n x + (1 - \frac{\epsilon}{2})T_n x - \frac{\epsilon}{2}P_n y - (1 - \frac{\epsilon}{2})T_n y \right\| \\ &= (1 - \frac{\epsilon}{2})\|T_n(x - y)\| \\ (29) \quad &\leq (1 - \frac{\epsilon}{2})\|(x - y)\|.\end{aligned}$$

For all $m, n \in \mathbb{N}$ with $m < n$, one has $P_n(S^{m,n-1}) = P_n$ which, for all x, y with $x - y \in N_{P_{m+1}}$, implies $S^{m,n-1}x - S^{m,n-1}y \in N_{P_n}$. Indeed,

$$P_n(S^{m,n-1}x - S^{m,n-1}y) = P_n(x - y) = P_n P_{m+1}(x - y) = 0.$$

Therefore, iterating (29), we obtain

$$\begin{aligned}\|S^{m,n}x - S^{m,n}y\| &= \|S_n(S^{m,n-1}x - S^{m,n-1}y)\| \\ &\leq (1 - \frac{\epsilon}{2})\|S^{m,n-1}x - S^{m,n-1}y\| \\ &\vdots \\ &\leq (1 - \frac{\epsilon}{2})^{n-m}\|(x - y)\|,\end{aligned}$$

which yields to

$$\delta_{P_{m+1}}(S^{m,n}) \leq (1 - \frac{\epsilon}{2})^{n-m} \rightarrow 0, \text{ as } n \rightarrow \infty,$$

and this proves that $\{S^{m,n}\}$ is weakly ergodic w.r.t. $\{P_n\}$. Hence, $\mathfrak{S}_{\{P_n\}}^w(X)$ is a dense subset of $\mathfrak{S}_{\{P_n\}}(X)$.

Now for every $m \in \mathbb{N}$, we have

$$\begin{aligned}\delta_{P_{m+1}}(T^{m,n+1}) &= \delta_{P_{m+1}}(T_{n+1}T^{m,n}) \\ &\leq \delta_{P_{m+1}}(T_{n+1})\delta_{P_{m+1}}(T^{m,n}) \\ &\leq \delta_{P_{m+1}}(T^{m,n}),\end{aligned}$$

which proves that $\{\delta_{P_{m+1}}(T^{m,n})\}_{n=1}^\infty$ is a non-increasing sequence. Therefore,

$$(30) \quad \mathfrak{S}_{\{P_n\}}^w(X) = \bigcap_{m=1}^\infty \bigcap_{k=1}^\infty \bigcup_{n=m+1}^\infty \left\{ \mathcal{T} \in \mathfrak{S}_{\{P_n\}}(X); \delta_{P_{m+1}}(T^{m,n}) < \frac{1}{k} \right\}.$$

For all $m < n$, define the mapping $\Phi_{m,n} : \mathfrak{S}_{\{P_n\}}(X) \rightarrow [0, 1]$ by

$$\Phi_{m,n}(\mathcal{T}) = \delta_{P_{m+1}}(T^{m,n}).$$

One can see that $\Phi_{m,n}$ is continuous, indeed

$$\begin{aligned} |\Phi_{m,n}(\mathcal{T}) - \Phi_{m,n}(\mathcal{S})| &= |\delta_{P_{m+1}}(T^{m,n}) - \delta_{P_{m+1}}(S^{m,n})| \\ &\leq \|T^{m,n} - S^{m,n}\| \\ &= \|T_n T_{n-1} \dots T_{m+1} - S_n S_{n-1} \dots S_{m+1}\| \\ &\leq \sum_{k=m+1}^n \|T_k - S_k\| \\ &\leq 2^n \sum_{k=1}^\infty \frac{\|T_k - S_k\|}{2^k} \\ &= 2^n d(\mathcal{T}, \mathcal{S}). \end{aligned}$$

Hence, for every $k \in \mathbb{N}$ the set $\Phi_{m,n}^{-1}([0, \frac{1}{k}])$ is an open subset of $\mathfrak{S}_{\{P_n\}}(X)$, and from (30), we infer that $\mathfrak{S}_{\{P_n\}}^w(X)$ is a G_δ -subset of $\mathfrak{S}_{\{P_n\}}(X)$. This completes the proof. \square

Remark 7.3. We point out that the residual properties of homogeneous Markov chains on C^* -algebras or von Neumann algebras were intensively studied in [4, 5, 32]. In the case $X = L^1(\mu)$, these results were investigated in [2, 21]. An analogue of Theorem 7.2 has been proved in [21, 36] when $X = \ell^1$. Therefore, our results are even new in the case of $X = L^1(\mu)$.

APPENDIX A. EXAMPLES OF ABSTRACT STATE SPACES

Let us provide some examples of abstract state spaces.

Example A.1. Let M be a von Neumann algebra. Let $M_{h,*}$ be the Hermitian part of the predual space M_* of M . As a base \mathcal{K} we define the set of normal states of M . Then $(M_{h,*}, M_{*,+}, \mathcal{K}, \mathbf{1})$ is a strong abstract state spaces, where $M_{*,+}$ is the set of all positive functionals taken from M_* , and $\mathbf{1}$ is the unit in M . In particular, if $M = L^\infty(E, \mu)$, then $M_* = L^1(E, \mu)$ is an abstract state space.

Example A.2. Let A be a real ordered linear space and, as before, let A_+ denote the set of positive elements of A . An element $e \in A_+$ is called order unit if for every $a \in A$ there exists a number $\lambda \in \mathbb{R}_+$ such that $-\lambda e \leq a \leq \lambda e$. If the order is Archimedean, then the mapping $a \rightarrow \|a\|_e = \inf\{\lambda > 0 : -\lambda e \leq a \leq \lambda e\}$ is a norm. If A is a Banach space with respect to this norm, the pair (A, e) is called an order-unit space with the order unit e . An element $\rho \in A^*$ is called positive if $\rho(a) \geq 0$ for all $a \in A_+$. By A_+^* we denote the set of all positive functionals. A positive linear functional is called a state if $\rho(e) = 1$. The set of all states is denoted by $S(A)$. Then it is well-known that $(A^*, A_+^*, S(A), e)$ is a strong abstract state space [1]. In particular, if \mathfrak{A}_{sa} is the self-adjoint part of an unital C^* -algebra, \mathfrak{A}_{sa} becomes order-unit spaces, hence $(\mathfrak{A}_{sa}^*, \mathfrak{A}_{sa,+}^*, S(\mathfrak{A}_{sa}), \mathbf{1})$ is a strong abstract state space.

Example A.3. Let X be a Banach space over \mathbb{R} . Consider a new Banach space $\mathcal{X} = \mathbb{R} \oplus X$ with a norm $\|(\alpha, x)\| = \max\{|\alpha|, \|x\|\}$. Define a cone $\mathcal{X}_+ = \{(\alpha, x) : \|x\| \leq \alpha, \alpha \in \mathbb{R}_+\}$ and a positive functional $f(\alpha, x) = \alpha$. Then one can define a base $\mathcal{K} = \{(\alpha, x) \in \mathcal{X} : f(\alpha, x) = 1\}$. Clearly, we have $\mathcal{K} = \{(1, x) : \|x\| \leq 1\}$. Then $(\mathcal{X}, \mathcal{X}_+, \mathcal{K}, f)$ is an abstract state space [24]. Moreover, X can be isometrically embedded into \mathcal{X} . Using this construction one can study several interesting examples of abstract state spaces.

Example A.4. Let A be the disc algebra, i.e. the sup-normed space of complex-valued functions which are continuous on the closed unit disc, and analytic on the open unit disc. Let $X = \{f \in A : f(1) \in \mathbb{R}\}$. Then X is a real Banach space with the following positive cone $X_+ = \{f \in X : f(1) = \|f\|\} = \{f \in X : f(1) \geq \|f\|\}$. The space X is an abstract state space, but not strong one (see [43] for details).

APPENDIX B. EXAMPLES OF MARKOV OPERATORS

Let us consider several examples of Markov operators.

Example B.1. Let $X = L^1(E, \mu)$ be the classical L^1 -space. Then any transition probability $P(x, A)$ defines a Markov operator T on X , whose dual T^* acts on $L^\infty(E, \mu)$ as follows

$$(T^*f)(x) = \int f(y)P(x, dy), \quad f \in L^\infty.$$

Example B.2. Let M be a von Neumann algebra, and consider $(M_{h,*}, M_{*,+}, \mathcal{K}, \mathbf{1})$ as in Example A.1. Let $\Phi : M \rightarrow M$ be a positive, unital ($\Phi(\mathbf{1}) = \mathbf{1}$) linear mapping. Then the operator given by $(Tf)(x) = f(\Phi(x))$, where $f \in M_{h,*}, x \in M$, is a Markov operator.

Example B.3. Let $X = C[0, 1]$ be the space of real-valued continuous functions on $[0, 1]$. Denote

$$X_+ = \{x \in X : \max_{0 \leq t \leq 1} |x(t) - x(1)| \leq 2x(1)\}.$$

Then X_+ is a generating cone for X , and $f(x) = x(1)$ is a strictly positive linear functional. Then $\mathcal{K} = \{x \in X_+ : f(x) = 1\}$ is a base corresponding to f . One can check that the base norm $\|x\|$ is equivalent to the usual one $\|x\|_\infty = \max_{0 \leq t \leq 1} |x(t)|$. Due to closedness of X_+ we conclude that (X, X_+, \mathcal{K}, f) is an abstract state space. Let us define a mapping T on X as follows:

$$(Tx)(t) = tx(t).$$

It is clear that T is a Markov operator on X .

Example B.4. Let X be a Banach space over \mathbb{R} . Consider the abstract state space $(\mathcal{X}, \mathcal{X}_+, \tilde{\mathcal{K}}, f)$ constructed in Example A.3. Let $T : X \rightarrow X$ be a linear bounded operator with $\|T\| \leq 1$. Then the operator $\mathcal{T} : \mathcal{X} \rightarrow \mathcal{X}$ defined by $\mathcal{T}(\alpha, x) = (\alpha, Tx)$ is a Markov operator.

Example B.5. Let A be the disc algebra, and let X be the abstract state space as in Example A.4. A mapping T given by $Tf(z) = zf(z)$ is clearly a Markov operator on X .

REFERENCES

1. E.M. Alfsen, *Compact convex sets and boundary integrals*, Springer-Verlag, Berlin, 1971.
2. W. Bartoszek, Norm residuality of ergodic operators, *Bull. Pol. Ac. Sci. Math* **29**(1981), 165–167.
3. W. Bartoszek, Asymptotic properties of iterates of stochastic operators on (AL) Banach lattices, *Anal. Polon. Math.* **52**(1990), 165–173.
4. W. Bartoszek, B. Kuna, On residualities in the set of Markov operators on C_1 , *Proc. Amer. Math. Soc.* **133** (2005), 2119–2129.
5. W. Bartoszek, B. Kuna, Strong mixing Markov semigroups on C_1 are meager, *Colloq. Math.* **105**(2006), 311–317.
6. W. Bartoszek, M. Pulka, On mixing in the class of quadratic stochastic operators, *Nonlin. Anal.: Theor. Methods* **86** (2013), 95–113.
7. R. L. Dobrushin, Central limit theorem for nonstationary Markov chains. I,II, *Theor. Probab. Appl.* **1**(1956), 65–80; 329–383.
8. C.C.Y. Dorea, A.G.C. Pereira, A note on a variation of Doeblin’s condition for uniform ergodicity of Markov chains, *Acta Math. Hungar.* **110**(2006), 287–292.
9. E. Yu. Emelyanov, *Non-spectral asymptotic analysis of one-parameter operator semigroups*, Birkhäuser Verlag, Basel, 2007.
10. E. Yu. Emel’yanov, M.P.H. Wolff, Asymptotic behavior of Markov semigroups on non-commutative L_1 -spaces, In book: *Quantum probability and infinite dimensional analysis* (Burg, 2001), 77–83, QP-PQ: Quantum Probab. White Noise Anal., 15, World Sci. Publishing, River Edge, NJ, 2003.
11. E. Yu. Emel’yanov, M.P.H. Wolff, Positive operators on Banach spaces ordered by strongly normal cones, *Positivity* **7**(2003), 3–22.
12. N. Erkursun-Ozcan, F. Mukhamedov, Uniform ergodicities and perturbation bounds of Markov chains on ordered Banach spaces, *J. Phys.: Conf. Ser.* **819**(2017), 012015.
13. N. Erkursun-Ozcan, F. Mukhamedov, Uniform ergodicities and perturbation bounds of Markov chains on ordered Banach spaces, *Queast. Math.* **41** (2018), no. 6, 863–876.
14. N. Erkursun-Ozcan, F. Mukhamedov, Uniform ergodicities of Lotz - Rübiger nets of Markov operators on ordered Banach spaces, *Results Math.* **73** (2018), no. 1, 35.
15. S. Gaubert, Z. Qu, Dobrushin’s ergodicity coefficient for Markov operators on cones and beyond, *Integ. Eqs. Operator Theor.* **81**(2014), 127–150.
16. J. Hajnal, Weak ergodicity in nonhomogeneous Markov chains, *Proc. Cambridge Phil. Soc.* **54**(1958) 233–246.
17. D.J. Hartfiel. Coefficients of ergodicity for imprimitive matrices, *Commun. Statis. Stochastic Models* **15**(1999), 81–88.
18. D.J. Hartfiel, U.G. Rothblum, Convergence of inhomogeneous products of matrices and coefficients of ergodicity, *Lin. Alg. Appl.* **277**(1998), 1–9.
19. H. Hennion, L. Harve, *Limit theorems for Markov chains and stochastic properties of dynamical systems by quasi-compactness*, Lec. Notes Math. **1766** (2001), Springer-Verlag, Berlin.
20. I.C.F. Ipsen, T.M. Salee, Ergodicity coefficients defined by vector norms, *SIAM J. Matrix Anal. Appl.* **32**(2011), 153–200.
21. A. Iwanik, Baire category of mixing for stochastic operators, *Rend. Circ. Mat. Palermo, Serie II* **28** (1992), 201–217.
22. J. Jachymski, Convergence of iterates of linear operators and the Kelisky- Rivlin type theorems, *Studia Math.*, **195** (2009), 99–113.
23. R.Jajte, *Strong limit theorems in non-commutative probability*, Lecture Notes in Math. vol. 1110, Berlin-Heidelberg: Springer 1984.
24. G. Jameson, *Ordered linear spaces*, Lect. Notes Math. V. 141, Springer-Verlag, Berlin, 1970.
25. J. Johnson, D. Isaacson, Conditions for strong ergodicity using intensity matrices, *J. Appl. Probab.* **25**(1988) 34–42.
26. I. Kontoyiannis, S.P. Meyn, Geometric ergodicity and the spectral gap of non-reversible Markov chains, *Probab. Theory Relat. Fields* **154**(2012), 327–339.
27. U. Krengel, *Ergodic Theorems*, Walter de Gruyter, Berlin-New York, 1985.
28. R. W. Madsen, D. L. Isaacson, Strongly ergodic behavior for non-stationary Markov processes, *Ann. Probab.* **1** (1973), 329–335.
29. S.P. Meyn, R.L. Tweedie, *Markov chains and stochastic stability*, Springer-Verlag, 1996.
30. F. Mukhamedov, Dobrushin ergodicity coefficient and ergodicity of noncommutative Markov chains, *J. Math. Anal. Appl.* **408** (2013), 364–373.

31. F. Mukhamedov, On L_1 -Weak Ergodicity of nonhomogeneous discrete Markov processes and its applications, *Rev. Mat. Comput.* **26**(2013), 799–813.
32. F. Mukhamedov, Ergodic properties of nonhomogeneous Markov chains defined on ordered Banach spaces with a base, *Acta. Math. Hungar.* **147** (2015), 294–323.
33. F. Mukhamedov, Uniform stability and weak ergodicity of nonhomogeneous Markov chains defined on ordered Banach spaces with a base, *Positivity*, **20**(2016), 135–153.
34. F. Mukhamedov A. Al-Rawashdeh, On generalized Dobrushin ergodicity coefficient and uniform Ergodicites of Markov Operators, *Positivity*, DOI 10.1007/s11117-019-00713-0 (2019).
35. A. Paz, Ergodic theorems for infinite probabilistic tables, *Ann. Math. Statist.* **41**(1970), 539–550.
36. M. Pulka, On the mixing property and the ergodic principle for nonhomogeneous Markov chains, *Linear Alg. Appl.* **434** (2011), 1475–1488.
37. A. Rhodius, On ergodicity coefficients of infinite stochastic matrices, *Zeit. Anal. Anwen.* **19**(2000), 873–887.
38. T.A. Sarymsakov, N.P. Zimakov, Ergodic principle for Markov semi-groups in ordered normal spaces with basis, *Dokl. Akad. Nauk. SSSR* **289** (1986), 554–558.
39. E. Seneta, On the historical development of the theory of finite inhomogeneous Markov chains, *Proc. Cambridge Philos. Soc.*, **74** (1973), 507–513
40. E. Seneta, *Non-negative matrices and Markov chains*, Springer, Berlin, 2006.
41. O. Szehr, M.M. Wolf, Perturbation bounds for quantum Markov processes and their fixed points, *J. Math. Phys.* **54**(2013), 032203.
42. Ch. P. Tan, On the weak ergodicity of nonhomogeneous Markov chains, *Statis. & Probab. Lett.* **26**(1996), 293–295.
43. D. Yost, A base norm space whose cone is not 1- generating, *Glasgow Math. J.* **25** (1984), 35–36.
44. A. I. Zeifman, D. L. Isaacson, On strong ergodicity for nonhomogeneous continuous-time Markov chains, *Stochast. Process. Appl.* **50**(1994), 263–273.
45. J. J. Vardy, B. A. Watson, Markov processes on Riesz spaces, *Positivity* **16** (2012), 373–391.
46. Y. C. Wong, K. F. Ng, *Partially ordered topological vector spaces*, Clarendon Press, 1973.