

The multi-degree of coverings on Lie groups

Haibao Duan*

Institute of Mathematics, Chinese Academy of Sciences,
Beijing 100190, P.R. China

Siye Wu

Department of Mathematics, National Tsing Hua University,
Hsinchu 30013, Taiwan

March 13, 2024

Abstract

We associate to each covering map of simple Lie groups a sequence of integers, called *the multi-degree of the covering*; develop a unified method to evaluate the sequence; and apply the results to solve two extension problems of mathematical physics.

2010 Mathematical Subject Classification: 57T10; 55R20; 81T13.

Key words and phrases: cohomology of Lie groups; Serre spectral sequence; Schubert calculus; WZW models; Gauge theory

Emails: dhb@math.ac.cn; swu@math.nthu.edu.tw

1 Main results

Covering maps on Lie groups are essential to extend the constructions and calculations of various physical models associated with simply connected Lie groups, to that associated with non-simply connected Lie groups. Naturally, topological invariants of the coverings are useful to formulate, or to solve, the relevant extension problems. In this paper we introduce for each covering map of Lie groups an invariant, called *the multi-degree of the covering*; extend Schubert calculus to evaluate the invariant; and apply the results to two outstanding topological problems arising from the studies of the Wess–Zumino–Witten models and the topological Gauge theories [6, 16, 25]. The main tool in our approach is *the Chow rings* of Lie groups, introduced by Grothendieck [19] in 1958.

To clarify the concept of *multi-degree* of a covering on Lie groups we recall the classical results of Hopf and Chevalley on the real cohomology of Lie groups. The Lie groups G under consideration will be compact and connected. For a maximal torus T on G the dimension $n := \dim T$ is an invariant of G , called *the rank of G* . The multiplication $\mu : G \times G \rightarrow G$ on G defines the *co-product* on the real cohomology algebra $H^*(G; \mathbb{R})$

*Supported by NSFC 11131008; 11661131004

$$(1.1) \quad \mu^* : H^*(G; \mathbb{R}) \rightarrow H^*(G \times G; \mathbb{R}) \cong H^*(G; \mathbb{R}) \otimes H^*(G; \mathbb{R}),$$

where the isomorphism is given by the Kunneth formula. An element $y \in H^*(G; \mathbb{R})$ is called *primitive* if the relation $\mu^*(y) = y \otimes 1 + 1 \otimes y$ is satisfied. Since the linear combinations of primitive elements are also primitive such elements form a subspace $P(G; \mathbb{R})$ of the algebra $H^*(G; \mathbb{R})$. Hopf [18] has shown that

Theorem 1.1. *For a Lie group G with rank n we have $\dim P(G; \mathbb{R}) = n$.*

Moreover, if $\{y_1, \dots, y_n\}$ is a basis of the space $P(G; \mathbb{R})$, then the real cohomology algebra $H^(G; \mathbb{R})$ is the exterior algebra generated by y_1, \dots, y_n :*

$$(1.2) \quad H^*(G; \mathbb{R}) = \Lambda_{\mathbb{R}}(y_1, \dots, y_n), \deg y_i \equiv 1 \pmod{2}. \square$$

By (1.2) the sequence $I_G = \{r_1, \dots, r_n\}$ of integers defined by

$$r_i := \frac{1}{2}(\deg y_i + 1), 1 \leq i \leq n,$$

is also an invariant of the group G , which has been shown by Chevalley [4],[27, (3.2)Theorem] to be the degree sequence of *the basic Weyl invariants of G* . We may assume that the sequence I_G is ordered by $r_1 \leq \dots \leq r_n$. In particular, if the group G is simple one has $2 = r_1 < \dots < r_n$ (see in Table 1).

In general, for a CW-complex X the torsion subgroup $\tau(X)$ of the integral cohomology $H^*(X)$ is an ideal and therefore, defines the (graded) quotient ring $\mathcal{F}(X) := H^*(X)/\tau(X)$. In view of the obvious additive decomposition

$$H^*(X) = \mathcal{F}(X) \oplus \tau(X)$$

we may call $\mathcal{F}(X)$ *the torsion free part* of the integral cohomology $H^*(X)$. Furthermore, if $f : X \rightarrow Y$ is a continuous map between two CW-complexes, the induce ring map f^* on the cohomologies preserves the torsion ideals. Therefore, passing to the quotients yields the (graded) ring map $f^\# : \mathcal{F}(Y) \rightarrow \mathcal{F}(X)$. In particular, the multiplication μ on G furnishes the ring $\mathcal{F}(G)$ with a co-product

$$\mu^\# : \mathcal{F}(G) \rightarrow \mathcal{F}(G) \otimes \mathcal{F}(G).$$

The following result may be seen as an integral refinement of Theorem 1.1.

Theorem A. *For a Lie group G with $I_G = \{r_1, \dots, r_n\}$, there exist n elements $x_1, \dots, x_n \in \mathcal{F}(G)$, $\deg x_i = 2r_i - 1$, so that*

- i) $\mathcal{F}(G) = \Lambda(x_1, \dots, x_n)$ (i.e. the exterior ring over \mathbb{Z})
- ii) $\mu^\#(x_i) = x_i \otimes 1 + 1 \otimes x_i, 1 \leq i \leq n$.

In addition,, if the group G is simple, then the generators x_1, \dots, x_n satisfying i) and ii) are unique up to the sign \pm convention.

Let G be a simple Lie group with non-trivial center $\mathcal{Z}(G)$. The quotient $c : G \rightarrow PG := G/\mathcal{Z}(G)$ is a covering map of Lie groups, and induces a ring map on the torsion free parts of the cohomologies

$$(1.3) \quad c^\# : \mathcal{F}(PG) = \Lambda(y_1, \dots, y_n) \rightarrow \mathcal{F}(G) = \Lambda(x_1, \dots, x_n),$$

where $\deg y_i = \deg x_i$ by the well known fact $I_G = I_{PG}$ of invariant theory. Granted with Theorem A we shall show that

Theorem B. *There exists a unique sequence $\{a_1, \dots, a_n\}$ of n positive integers, so that*

- i) *the map $c^\#$ is given by $c^\#(y_i) = a_i \cdot x_i$, $1 \leq i \leq n$;*
- ii) *the product $a_1 \cdot \dots \cdot a_n$ is the order $|\mathcal{Z}(G)|$ of the center.*

The unique sequence $\{a_1, \dots, a_n\}$ obtained by Theorem B, written $\mathcal{D}(G, PG)$ and called *the multi-degree* of the covering c , will be the main concern of this work. To be precise we tabulate all the simply connected simple Lie groups G with non-trivial centers $\mathcal{Z}(G)$, together with the degree sequences I_G of their basic Weyl invariants, using the table below.

Table 1. The simply connected simple Lie groups with non-trivial centers

G	$\text{rank } G$	I_G	$\mathcal{Z}(G)$
$SU(n)$	$n - 1$	$\{2, 3, \dots, n\}$	\mathbb{Z}_n
$Sp(n)$	n	$\{2, 4, \dots, 2n\}$	\mathbb{Z}_2
$Spin(2n + 1)$	n	$\{2, 4, \dots, 2n\}$	\mathbb{Z}_2
$Spin(2n), n \equiv 1 \pmod{2}$	n	$\{2, 4, \dots, 2(n - 1)\} \amalg \{n\}$	\mathbb{Z}_4
$Spin(2n), n \equiv 0 \pmod{2}$	n	$\{2, 4, \dots, 2(n - 1)\} \amalg \{n\}$	$\mathbb{Z}_2 \oplus \mathbb{Z}_2$
E_6	6	$\{2, 5, 6, 8, 9, 12\}$	\mathbb{Z}_3
E_7	7	$\{2, 6, 8, 10, 12, 14, 18\}$	\mathbb{Z}_2

Let $J(PG)$ be the subring of $H^*(PG)$ generated multiplicatively by the second cohomology group $H^2(PG)$, together with the multiplicative unit $1 \in H^0(PG)$. By computing with the Chow rings [19] of the adjoint Lie groups PG we shall show that

Theorem C. *For each simply connected Lie group G in Table 1 we have*

$$(1.4) \quad H^2(PG) = \mathcal{Z}(G).$$

Moreover, there exist generators

$$\begin{aligned} \omega &\in H^2(PG) \text{ for } G \neq Spin(2n) \text{ with } n \equiv 0 \pmod{2}; \text{ or} \\ \omega_1, \omega_2 &\in H^2(PG) = \mathbb{Z}_2 \oplus \mathbb{Z}_2 \text{ for } G = Spin(2n) \text{ with } n \equiv 0 \pmod{2} \end{aligned}$$

so that the ring $J(PG)$ has the following presentation:

$$\begin{aligned} \text{i) } J(PSU(n)) &= \frac{\mathbb{Z}[\omega]}{\langle b_{n,r}\omega^r, 1 \leq r \leq n \rangle}, \text{ where } b_{n,r} = g.c.d.\{\binom{n}{1}, \dots, \binom{n}{r}\}; \\ \text{ii) } J(PSp(n)) &= \frac{\mathbb{Z}[\omega]}{\langle 2\omega, \omega^{2r+1} \rangle}, \text{ where } n = 2^r(2k + 1); \\ \text{iii) } J(PSpin(2n + 1)) &= \frac{\mathbb{Z}[\omega]}{\langle 2\omega, \omega^{2^s+1} \rangle}, \text{ where } 2^s \leq n < 2^{s+1}; \\ \text{iv) } J(PSpin(2n)) &= \begin{cases} \frac{\mathbb{Z}[\omega]}{\langle 4\omega, 2\omega^2, \omega^{2^r+1} \rangle} & \text{if } n \equiv 1 \pmod{2}; \\ \frac{\mathbb{Z}[\omega_1, \omega_2]}{\langle 2\omega_1, 2\omega_2, \omega_1^{2^t}, \omega_2^{2^r+1} \rangle} & \text{if } n \equiv 0 \pmod{2}, \end{cases} \\ &\text{where } n = 2^t(2k + 1), 2^r < n \leq 2^{r+1}; \\ \text{v) } J(PE_6) &= \frac{\mathbb{Z}[\omega]}{\langle 3\omega, \omega^9 \rangle}; \quad \text{vi) } J(PE_7) = \frac{\mathbb{Z}[\omega]}{\langle 2\omega, \omega^2 \rangle}. \end{aligned}$$

Our main result implies that, the multi-degree sequence $\mathcal{D}(G, PG)$ is determined entirely by the structure of the ring $J(PG)$ given by Theorem C (see Theorem 4.4).

Theorem D. *For a Lie group G in Table 1 the multi-degree $\mathcal{D}(G, PG)$ is*

G	$\mathcal{D}(G, PG)$
$SU(n)$	$\{a_1, \dots, a_{n-1}\}, a_k = \frac{b_{n,k}}{b_{n,k+1}},$
$Sp(n), n = 2^t(2b+1)$	$\{1, \dots, 1, 2_{(2^t)}, 1, \dots, 1\}$
$Spin(2n+1), 2^s \leq n < 2^{s+1}$	$\{1, \dots, 1, 2_{(2^s)}, 1, \dots, 1\}$
$Spin(2n), 2^s < n = 2b+1 \leq 2^{s+1}$	$\{2, 1, \dots, 1, 2_{(2^s)}, 1, \dots, 1\}$
$Spin(2n), 2^s < n = 2^t(2b+1) \leq 2^{s+1}, t \geq 1$	$\{1, \dots, 1, 2_{(2^{t-1})}, 1, \dots, 1, 2_{(2^s)}, 1, \dots, 1\}$
E_6 or E_7	$\{1, 1, 1, 1, 3, 1\}$ or $\{2, 1, 1, 1, 1, 1, 1\}$

where the notion $2_{(h)}$ stands for $a_h = 2$, and where if $G = SU(n)$ and the integer n has the prime factorization $n = p_1^{r_1} \dots p_t^{r_t}$, then (see [10, Lemma 3.1])

$$(1.5) \quad a_k = \begin{cases} p_i & \text{if } k = p_i^s \text{ with } 1 \leq i \leq t \text{ and } 1 \leq s \leq r_i; \\ 1 & \text{otherwise.} \end{cases}$$

Theorem D is a natural extension of [10, Theorem 1.1], where the multi-degree of the universal covering $SU(n) \rightarrow PSU(n)$ on the projective unitary group $PSU(n)$ has been computed. On the other hand, E.B. Dynkin [14, 15] raised the problem to determine induce actions of Lie group homomorphisms $G \rightarrow G'$ on the cohomologies, which is essentially solved by Theorem D for the cases of universal coverings of the adjoint Lie groups. Nevertheless, the present paper is motivated by two topics from mathematical physics. To start with we note by Theorem D that

Corollary 1.2. *For a Lie group G in Table 1 the leading term a_1 of the sequence $\mathcal{D}(G, PG)$ is either 1 or 2, where $a_1 = 2$ occurs if and only if G is isomorphic to one of the following groups*

- i) $SU(n)$ with $n \equiv 0 \pmod{2}$;
- ii) $Sp(n)$ with $n \equiv 1 \pmod{2}$;
- iii) $Spin(2^t(2b+1))$ with $t = 1, 2$;
- iv) E_7 . \square

In the study on the Wess–Zumino–Witten models with simple Lie groups Felder, Gawedzki and Kupiainen obtained the short exact sequence

$$(1.6) \quad 0 \rightarrow H_3(G)(= \mathbb{Z}) \xrightarrow{c_*} H_3(PG) \rightarrow \mathcal{Z}(G) \rightarrow 0,$$

and investigated its extension problem [16, Appendix 1], where G is simply connected and simple. This problem was emphasized by Dijkgraaf and Witten in the work [6] on the topological Gauge theories, and by Mathai and Rosenberg [25] in the study on the relationship between Langlands duality and T–duality for compact Lie groups. Since the map c_* in (1.6) is Kronecker dual to the map c^* on $H^3(PG)$ we get from Corollary 1.2 the following result.

Corollary 1.3. *For the Lie groups G given in Table 1 we have*

$$H_3(PG) = \mathbb{Z} \oplus \mathcal{Z}(G)$$

with the following exceptions

- i) $H_3(PSU(2m)) = \mathbb{Z} \oplus \mathbb{Z}_m$, where $m \in \mathbb{Z}$;
- ii) $H_3(PSp(2b+1)) = H_3(PE_7) = \mathbb{Z}$, where $b \in \mathbb{Z}$;
- iii) $H_3(PSpin(2^t(2b+1))) = \mathbb{Z} \oplus \mathbb{Z}_2$, where $t = 1, 2$, $b \in \mathbb{Z}$. \square

Remark 1.4. In the inspiring work [25] Mathai and Rosenberg have computed the leading terms a_1 of the sequences $\mathcal{D}(G, PG)$. In the cases of $G = SU(n)$ with n divisible by 4 and $G = Spin(2(2b+1))$, our results in Corollary 1.2 are different with the ones stated in [25, Theorem 1, (1), (3)]. We note that these are precisely the cases where the integral cohomologies $H^*(G)$ have torsion elements of the order 4. Therefore, working with the \mathbb{Z}_2 algebra $H^*(G; \mathbb{Z}_2)$ alone may not suffice to decide a_1 . \square

Let B_G be the classifying space of a simple Lie group G . By the naturality of the cohomology suspension $\tau_G : H^r(B_G) \rightarrow H^{r-1}(G)$ [17, p.22] in the universal G -bundle

$$G \hookrightarrow E_G \rightarrow B_G$$

the covering $c : G \rightarrow PG$ induces the commutative diagram

$$(1.7) \quad \begin{array}{ccc} H^4(B_{PG}) & \xrightarrow{\tau_{PG}} & H^3(PG) \\ Bc^* \downarrow & & c^* \downarrow \\ H^4(B_G) & \xrightarrow{\cong \tau_G} & H^3(G) \end{array} \quad (\text{compare with [6, (4.14)]})$$

where, as being pointed out in [6] that, the four vertices groups in (1.7) are all isomorphic to \mathbb{Z} , and where the suspension τ_G at the bottom is an isomorphism when G is simply connected. Let ω and $\tilde{\omega}$ denote the generators of respectively $H^4(B_{PG})$ and $H^4(B_G)$, ξ and $\tilde{\xi}$ denote the generators of respectively $H^3(PG)$ and $H^3(G)$. In [6, Section 4.3] Dijkgraaf and Witten raised the interesting problem to decide the pair (α, β) of integers characterized by the relations

$$Bc^*(\omega) = \alpha \cdot \tilde{\omega}, \tau_{PG}(\omega) = \beta \cdot \xi,$$

where the pair (α, β) has shown to be $(\frac{2n}{g.c.d.\{2, n-1\}}, n)$ for the case $G = SU(n)$.

On the other hand, by the commutativity of the diagram (1.7) one has $\alpha = a_1 \cdot \beta$, where a_1 is the leading term of the sequence $\mathcal{D}(G, PG)$. Therefore, combining results of Corollary 1.2 with Dijkgraaf–Witten’s formula [6, (4.18)] evaluating the integer α we obtain the following results.

Corollary 1.5. *For a Lie group G in Table 1 the pair (α, β) of integers is given by the table:*

G	(α, β)
$SU(n)$	$(\frac{2n}{g.c.d.\{2, n-1\}}, n)$
$Spin(2n+1)$	$(2, 2)$
$Sp(n)$, $n = 2^t(2b+1)$	$(8, 4), (4, 4), (2, 2)$ or $(1, 1)$ for $t = 0, 1, 2$ or ≥ 3
$Spin(2n)$, $n = 2^t(2b+1)$	$(8, 4), (4, 4)$ or $(2, 2)$ for $t = 0, 1$ or ≥ 2
E_6	$(3, 3)$
E_7	$(4, 2)$

The paper is arranged as follows. In Section 2 we develop fundamental properties of the Serre spectral sequence of the torus fibration $\pi : G \rightarrow G/T$, where T is a maximal torus on G . The results are applied in Section 3 to show Theorems A, B and D. Section 4 is devoted to an exact sequence associated to the cyclic coverings of Lie groups, by which the proof of Theorem D is reduced to computing with the Chow rings of Lie groups.

In this paper the cohomologies and spectral sequences are over the ring \mathbb{Z} of integers, unless otherwise stated. Given a subset S of a ring A the notion $\langle S \rangle$ stands for the ideal generated by S , while $A/\langle S \rangle$ denotes the quotient ring. In addition, the elements in a graded ring or algebra are assumed to be homogeneous.

Remark 1.6. The problem of computing the cohomology of Lie groups was raised by E. Cartan in 1929, which was solved only for the cohomologies with field coefficients, see Reeder [27] and Kač [21] for accounts about the history. As for the task of the present work general results on the integral cohomology of Lie groups, such as Theorems A and B, are requested.

In his problem 15 Hilbert asked for a rigorous foundation of Schubert calculus. Van der Waerden [28] and A. Weil [29, p.331] attributed the problem to the determination of the cohomology rings of the flag manifolds G/T . The proofs of Theorems C and D illustrate how Schubert calculus could be extended to computing with the integral cohomologies of Lie groups [13, Remark 6.3]. \square

2 The integral cohomology of Lie groups

For a Lie group G with a maximal torus T consider the torus fibration

$$(2.1) \quad T \rightarrow G \xrightarrow{\pi} G/T.$$

on the group G . The Borel transgression in π is the composition

$$\tau = (\pi^*)^{-1} \circ \delta : H^1(T) \xrightarrow{\delta} H^2(G, T) \xrightarrow[\cong]{(\pi^*)^{-1}} H^2(G/T) \quad ([9]),$$

where δ is the connecting homomorphism in the cohomological exact sequence of the pair (G, T) , and where the map π^* from $H^2(G/T)$ to $H^2(G, T)$ is always an isomorphism. By the Leray-Serre theorem [24, p.135] we have that

Lemma 2.1. *The second page of the Serre spectral sequence $\{E_r^{*,*}(G), d_r\}$ of the fibration π is the Koszul complex*

$$(2.2) \quad E_2^{*,*}(G) = H^*(G/T) \otimes H^*(T) \quad (\text{see [24, p.259]})$$

on which the differential d_2 is determined by the transgression τ as

$$\begin{aligned} \text{i)} \quad & d_2(x \otimes 1) = 0, \quad d_2(1 \otimes t) = \tau(t) \otimes 1, \quad t \in H^1(T), \\ \text{ii)} \quad & d_2(z \cdot z') = d_2(z) \cdot z' + (-1)^{\deg z} z \cdot d_2(z'), \quad z, z' \in E_2^{*,*}(G). \end{aligned}$$

In particular, if $\dim G/T = m$ and $\dim T = n$, then

$$(2.3) \quad E_2^{m,n}(G) = H^m(G) \otimes H^n(T) = E_r^{m,n}(G) = H^{n+m}(G) = \mathbb{Z}, \quad r \geq 2;$$

$$(2.4) \quad E_3^{*,0}(G) = H^*(G/T) / \langle \text{Im } \tau \rangle;$$

Proof. The base manifold G/T of π is the complete flag manifold associated to the Lie group G , hence is simply connected. Formula (2.2), together with properties i) and ii), are standard [24, p.259].

By formula (2.2) we have that

$$E_2^{m,n}(G) = H^m(G/T) \otimes H^n(T) = \mathbb{Z}$$

and that $E_2^{s,t}(G) = 0$ if either $s > m$ or $t > n$. It implies that any differential d_r that acts or lands on the group $E_r^{m,n}(G)$ must be trivial, confirming the isomorphisms in (2.3).

Finally, by formula i) the differential

$$d_2 : E_2^{*,1}(G) = H^*(G/T) \otimes H^1(T) \rightarrow E_2^{*,0}(G) = H^*(G/T)$$

is $d_2(x \otimes t) = x \cup \tau(t)$, $t \in H^1(T)$. We get formula (2.4) from

$$E_3^{*,0}(G) = E_2^{*,0}(G) / \text{Im } d_2 = H^*(G/T) / \langle \text{Im } \tau \rangle. \square$$

The Koszul complex $\{E_2^{*,*}(G), d_2\}$ has now been well understood by the following works:

- a) the base manifold G/T is a flag variety which has a canonical decomposition into the Schubert cells on G/T [2, 5];
- b) presentation of the ring $H^*(G/T)$ by special Schubert classes on G/T has been completed by Duan and Zhao in [11];
- c) with respect to the Schubert basis on $H^2(G/T)$ formula of the transgression τ is available in [9, Theorem 2.5].

Combining these results explicit construction of the ring $E_3^{*,*}(G)$ has been carried out in [8, 12], among which we shall only need the following two results.

In [19] Grothendieck defined the *Chow ring of a Lie group G* to be the subring $\mathcal{A}^*(G) := \pi^*(H^*(G/T))$ of $H^*(G)$, where he has also shown

$$\mathcal{A}^*(G) = H^*(G/T) / \langle \text{Im } \tau \rangle \quad ([19, \text{p.21, Rem.2}]).$$

Comparing this with formula (2.4) we obtain that

Lemma 2.2. *The map π in (2.1) induces an isomorphism*

$$(2.5) \quad \mathcal{A}^*(G) = E_3^{*,0}(G).$$

In particular, for the Lie groups G given in Table 1, the rings $\mathcal{A}^(G)$ and $\mathcal{A}^*(PG)$ admit the following presentations (in terms of generators–relations):*

$$(2.6) \quad \mathcal{A}^*(SU(n)) = \mathbb{Z}, \mathcal{A}^*(PSU(n)) = \frac{\mathbb{Z}[\omega]}{\langle b_{n,r}\omega^r, 1 \leq r \leq n \rangle}, b_{n,r} = g.c.d.\{\binom{n}{1}, \dots, \binom{n}{r}\};$$

$$(2.7) \quad \mathcal{A}^*(Sp(n)) = \mathbb{Z}, \mathcal{A}^*(PSp(n)) = \frac{\mathbb{Z}[\omega]}{\langle 2\omega, \omega^{2^r+1} \rangle}, \text{ where } n = 2^r(2k+1);$$

$$\begin{aligned}
(2.8) \quad \mathcal{A}^*(Spin(2n+1)) &= \frac{\mathbb{Z}[x_3, x_5, \dots, x_{2[\frac{n+1}{2}]-1}]}{\langle 2x_{2i-1}, x_{2i-1}^{k_i}; 2 \leq i \leq [\frac{n+1}{2}] \rangle}, \\
\mathcal{A}^*(PSpin(2n+1)) &= \frac{\mathbb{Z}[x_1, x_3, x_5, \dots, x_{2[\frac{n+1}{2}]-1}]}{\langle 2x_{2i-1}, x_{2i-1}^{k_i}; 1 \leq i \leq [\frac{n+1}{2}] \rangle}; \\
(2.9) \quad \mathcal{A}^*(Spin(2n)) &= \frac{\mathbb{Z}[x_3, x_5, \dots, x_{2[\frac{n}{2}]-1}]}{\langle 2x_{2i-1}, x_{2i-1}^{k_i}; 2 \leq i \leq [\frac{n}{2}] \rangle}; \\
\mathcal{A}^*(PSpin(2n)) &= \begin{cases} \frac{\mathbb{Z}[x_1, x_3, x_5, \dots, x_{2[\frac{n}{2}]-1}]}{\langle 4x_1, 2x_1^2, x_1^{k_1}, 2x_{2i-1}, x_{2i-1}^{k_i}; 2 \leq i \leq [\frac{n}{2}] \rangle} & \text{if } n \equiv 1 \pmod{2}, \\ \frac{\mathbb{Z}[\omega, x_1, x_3, x_5, \dots, x_{2[\frac{n}{2}]-1}]}{\langle 2\omega, \omega^{2^h}, 2x_{2i-1}, x_{2i-1}^{k_i}; 1 \leq i \leq [\frac{n}{2}] \rangle} & \text{if } n = 2^h(2b+1), h \geq 1; \end{cases} \\
(2.10) \quad \mathcal{A}^*(E_6) &= \frac{\mathbb{Z}[x_3, x_4]}{\langle 2x_3, 3x_4, x_3^2, x_4^3 \rangle}, \\
\mathcal{A}^*(PE_6) &= \frac{\mathbb{Z}[\omega, x_3, x_4]}{\langle 3\omega, 2x_3, 3x_4, (x_3 + \omega^3)^2, \omega^9, x_4^3 \rangle}; \\
(2.11) \quad \mathcal{A}^*(E_7) &= \frac{\mathbb{Z}[x_3, x_4, x_5, x_9]}{\langle 2x_3, 3x_4, 2x_5, 2x_9, x_3^2, x_4^3, x_5^2, x_9^2 \rangle}, \\
\mathcal{A}^*(PE_7) &= \frac{\mathbb{Z}[\omega, x_3, x_4, x_5, x_9]}{\langle 2\omega, \omega^2, 2x_3, 3x_4, 2x_5, 2x_9, x_3^2, x_4^3, x_5^2, x_9^2 \rangle},
\end{aligned}$$

where the generators ω, x_i ($\deg \omega = 2, \deg x_k = 2k$) are the π^* -images of certain Schubert classes on G/T specified in [11], and where the power k_i 's appearing in the denominator of the quotient (2.7) or (2.8) are respectively

$$k_i = 2^{\lfloor \ln \frac{n}{2} \rfloor + 1} \text{ or } k_i = 2^{\lfloor \ln \frac{n-1}{2} \rfloor + 1}.$$

Proof. Combining the results mentioned in b) and c) above it is straightforward to evaluate the Chow rings $\mathcal{A}(G)$ and $\mathcal{A}^*(PG)$ using formula (2.4). As examples we show the results for $G = PSU(n)$ and $PSpin(2n)$, and refer the remaining cases to [11, Corollary 6.2; formula (6.13)] and [23, p.i].

For $G = SU(n)$ the ring $H^*(G/T)$ has the presentation

$$H^*(G/T) = \frac{\mathbb{Z}[t_1, t_2, \dots, t_n]}{\langle e_1, \dots, e_n \rangle} \text{ (by Borel),}$$

where $\deg t_i = 2$, e_r is the r^{th} elementary symmetric polynomial in the t_1, \dots, t_n . On the other hand for the group $PSU(n)$ we have

$$\text{Im } \tau = \{t_1 - t_2, t_2 - t_3, \dots, t_{n-1} - t_n\} \text{ by [9, Corollary 3.2].}$$

From (2.4) we get

$$\mathcal{A}^*(PSU(n)) = \frac{\mathbb{Z}[t_1, t_2, \dots, t_n]}{\langle e_1, \dots, e_n \rangle} \big|_{t_i=t_j} = \frac{\mathbb{Z}[\omega]}{\langle \binom{n}{r} \omega^r, 1 \leq r \leq n \rangle},$$

where $\omega := t_1 = \dots = t_n$. It implies that the order of the power ω^r in the ring $\mathcal{A}^*(PSU(n))$ is precisely $b_{n,r} = g.c.d. \{ \binom{n}{1}, \dots, \binom{n}{r} \}$, showing (2.6).

For $G = Spin(2n)$ we have by Marlin [23, p.20, Theorem 3] that

$$H^*(Spin(2n)/T) = \frac{\mathbb{Z}[t_1, t_2, \dots, t_n, x_1, x_2, \dots, x_{n-1}]}{\langle \delta_1, \dots, \delta_n, q_1, \dots, q_{n-1} \rangle}, \deg t_i = 2, \deg x_r = 2r,$$

where, with $x_k = 0$ for $k \geq n$ being understood,

$$\begin{aligned}\delta_r &= 2x_r - e_r(t_1, \dots, t_n), \quad 1 \leq r \leq n-1; \quad \delta_n = e_n(t_1, \dots, t_n), \\ q_r &= x_{2r} + 2 \sum_{1 \leq i \leq r-1} (-1)^i x_i x_{2r-i} + (-1)^r x_r^2, \quad 1 \leq r \leq n-1,\end{aligned}$$

and where $e_r(t_1, \dots, t_n)$ is the r^{th} elementary symmetric polynomials in t_1, \dots, t_n . On the other hand, for the group $PSpin(2n)$ we have

$$\text{Im } \tau = \{t_1 - t_2, \dots, t_{n-1} - t_n, 2t_n\} \text{ by [9, Corollary 3.2].}$$

The second formula in (2.9) is verified by

$$\begin{aligned}\mathcal{A}^*(PSpin(2n)) &= H^*(Spin(2n)/T) / \langle \text{Im } \tau \rangle \\ &= H^*(Spin(2n)/T) |_{t_1=\dots=t_n, 2t_n=0},\end{aligned}$$

where $x_1 = \frac{1}{2}e_1$ (by the relation δ_1), and where $\omega := t_1 = \dots = t_n$. \square

With the product inherited from that on $E_2^{*,*}(G)$ the third page $E_3^{*,*}(G)$ is a bi-graded ring [30, P.668]. In particular, $E_3^{*,1}(G)$ can be considered as a module over the subring $\mathcal{A}^*(G) = E_3^{*,0}(G) \subset E_3^{*,*}(G)$.

Lemma 2.3. *For each compact Lie group G with $I_G = \{r_1, \dots, r_n\}$ there exist n elements $\rho_k \in E_3^{2r_k-2,1}(G)$, $1 \leq k \leq n$, so that*

- i) *as a $\mathcal{A}^*(G)$ -module $E_3^{*,1}(G)$ is spanned additively by $\{\rho_1, \dots, \rho_n\}$;*
- ii) *the product $\rho_1 \cdots \rho_n$ generates the group $E_3^{m,n}(G) = \mathbb{Z}$ (see (2.3)).* \square

Example 2.4. A classical result of Leray [9, Example 1.4] states that for the cohomology with real coefficients one has isomorphism of algebras

$$H^*(G; \mathbb{R}) = E_3^{*,*}(G; \mathbb{R}) = \Lambda_{\mathbb{R}}(y_1, \dots, y_n)$$

where the generators y_1, \dots, y_n is a basis of the subspace $E_3^{*,1}(G; \mathbb{R})$ constructed from the basic Weyl invariants of the group G . The elements $\rho_1, \dots, \rho_n \in E_3^{*,1}(G)$ asserted by Lemma 2.3 may be seen as the integral lifts of the classes y_1, \dots, y_n , and will be called a *basis* of the $\mathcal{A}^*(G)$ -module $E_3^{*,1}(G)$ (for the latter convenience). Note that $E_3^{*,1}(G)$ may fail to be a free module over the ring $\mathcal{A}^*(G)$, e.g. see [9, Example 1.4] for the case $G = PSU(8)$.

Without the loss of generalities we can assume in Lemma 2.3 that the Lie group G is simple. In this case a basis $\{\rho_1, \dots, \rho_n\}$ of the module $E_3^{*,1}(G)$ has been constructed uniformly for all simply connected G in [12], and for the non-simply connected G in [8, 9]. In this paper knowing the degrees of these elements is sufficient for our purpose.

It is worth to mention that, in the course to compute the integral cohomology of the spinor group $Spin(n)$ Pittie [26] has constructed a basis $\{\rho_1, \rho_2, \dots\}$ of the module $E_3^{*,1}(G)$ for the classical groups $G = U(n), SO(n)$ and $Spin(n)$. \square

To see the implication of Lemma 2.3 let F^p be the filtration on the integral cohomology $H^*(G)$ defined by the map π . That is ([24, P.146])

$$0 = F^{r+1}(H^r(G)) \subseteq F^r(H^r(G)) \subseteq \dots \subseteq F^0(H^r(G)) = H^r(G)$$

with

$$E_{\infty}^{p,q}(G) = F^p(H^{p+q}(G))/F^{p+1}(H^{p+q}(G)).$$

The fact $H^{2k+1}(G/T) = 0$ due to Bott and Samelson [3] implies that

$$\text{a) } E_r^{2k+1,*} = 0 \text{ for } k \geq 0; \quad \text{b) } E_3^{4s,2} = E_4^{4s,2} = \dots = E_{\infty}^{4s,2}.$$

From $F^{2s+1}(H^{2s+1}(G)) = F^{2s+2}(H^{2s+1}(G)) = 0$ by a) one finds that

$$E_{\infty}^{2s,1}(G) = F^{2s}(H^{2s+1}(G)) \subset H^{2s+1}(G).$$

Combining this with the routine relation $d_r(E_r^{*,1}) = 0$ for $r \geq 3$ yields the composition

$$(2.12) \quad \kappa : E_3^{*,1}(G) \rightarrow E_4^{*,1}(G) \rightarrow \dots \rightarrow E_{\infty}^{*,1}(G) \subset H^*(G)$$

that interprets elements of $E_3^{*,1}$ directly as cohomology classes of G . Carrying on the results of Lemma 2.3 we obtain the following characterization of the integral cohomology $H^*(G)$ in term of its free part and torsion ideal.

Lemma 2.5. *The integral cohomology of G has the additive presentation*

$$(2.13) \quad H^*(G) = \Delta(\xi_1, \dots, \xi_n) \oplus \tau(G) \text{ with } \xi_i := \kappa(\rho_i) \in H^*(G),$$

where $\Delta(\xi_1, \dots, \xi_n)$ denotes the free \mathbb{Z} -module with the basis

$$\Phi := \{1, \xi_I = \bigcup_{i \in I} \xi_i \in H^*(G) \mid I \subseteq \{1, \dots, n\}\}.$$

Proof. Let $\{E_r^{*,*}(G; \mathbb{R}), d_r\}$ be the Serre spectral sequence of π with real coefficients. According to Leray [22] the algebra $E_3^{*,*}(G; \mathbb{R})$ is generated multiplicatively by its subspace $E_3^{*,1}(G; \mathbb{R})$, while the map κ in (2.12) induces

- i) an isomorphism $E_3^{*,1}(G; \mathbb{R}) \cong P(G; \mathbb{R})$ of vector spaces, and
- ii) an isomorphism $E_3^{*,*}(G; \mathbb{R}) \cong H^*(G; \mathbb{R})$ of algebras,

respectively. Assume that $\{\rho_1, \dots, \rho_n\}$ is a basis of the $\mathcal{A}^*(G)$ -module $E_3^{*,1}(G)$, and that $\dim G/T = m$, $\dim T = n$. In addition to $\xi_i := \kappa(\rho_i)$ we put

$$\xi_i^R := \xi_i \otimes 1 \in H^*(G; \mathbb{R}) = H^*(G) \otimes \mathbb{R}, \quad 1 \leq i \leq n.$$

Since $\mathcal{A}^*(G) \otimes \mathbb{R} = \mathbb{R}$ the space $E_3^{*,1}(G; \mathbb{R}) = E_3^{*,1}(G) \otimes \mathbb{R}$ has the basis $\{\rho_1 \otimes 1, \dots, \rho_n \otimes 1\}$ by i) of Lemma 2.3. By the isomorphisms i) and ii) above

$$H^*(G; \mathbb{R}) = \Lambda_{\mathbb{R}}(\xi_1^R, \dots, \xi_n^R) \text{ with } \xi_i^R \in P(G; \mathbb{R}),$$

implying that the set Φ of 2^n monomials is linearly independent in $H^*(G)$. It remains to show that the set Φ spans a direct summand of $H^*(G)$.

Assume, on the contrary, that there exist a monomial $\xi_I \in \Phi$, an integral class $\varsigma \in H^*(G)$, as well as some integer $a > 1$, so that a relation of the form $\xi_I = a \cdot \varsigma$ holds in $H^*(G)$. Multiplying both sides by ξ_J with J the complement of $I \subseteq \{1, \dots, n\}$ yields that

$$\xi_1 \cup \dots \cup \xi_n = (-1)^r a \cdot (\varsigma \cup \xi_J) \text{ (for some } r \in \{0, 1\}).$$

However, in view of the identification (2.3) the map κ in (2.12) transforms the generator $\rho_1 \dots \rho_n$ of $E_3^{m,n}(G) = \mathbb{Z}$ to the generator $\xi_1 \cup \dots \cup \xi_n$ of $H^{n+m}(G) = \mathbb{Z}$ (by $\xi_i := \kappa(\rho_i)$). The proof is completed by this contradiction. \square

3 Proofs of Theorems A, B and C

Let $\{\rho_1, \dots, \rho_n\}$ be a basis of the $\mathcal{A}^*(G)$ -module $E_3^{*,1}(G)$. With $\xi_i = \kappa(\rho_i)$ and $\xi_i^R = \xi_i \otimes 1$ we have by the proof of Lemma 2.5 that

$$(3.1) \quad H^*(G; \mathbb{R}) = \Lambda_{\mathbb{R}}(\xi_1^R, \dots, \xi_n^R), \text{ where } \xi_i^R \in P(G; \mathbb{R}).$$

Proof of Theorem A. Let $q : H^*(G) \rightarrow \mathcal{F}(G) = H^*(G)/\tau(G)$ be the quotient map and put $x_i := q(\xi_i)$. Then $\mathcal{F}(G) = \Delta(x_1, \dots, x_n)$ by (2.13), where

$$\deg x_i = \deg \xi_i = 2r_i - 1, 1 \leq i \leq n, I_G = \{r_1, \dots, r_n\}.$$

In addition, with $\deg \xi_i \equiv 1 \pmod{2}$ we get $x_i^2 = 0$ from $\xi_i^2 \in \tau(G)$, showing

$$(3.2) \quad \mathcal{F}(G) = \Lambda(x_1, \dots, x_n) \text{ (i.e. the formula i) of Theorem A).}$$

For the co-product μ^* on $H^*(G)$ we can assume, in general, that

$$\mu^*(\xi_i) = \xi_i \otimes 1 + 1 \otimes \xi_i + z_i + d_i, 1 \leq i \leq n,$$

where $z_i = \sum a_j \otimes b_j \in \mathcal{F}^+(G) \times \mathcal{F}^+(G)$ is a mixed term, $d_i \in \tau(G \times G)$. With $\xi_i^R \in P(G; \mathbb{R})$ by (3.1) there must be $z_i = 0$. Thus, applying $q \times q$ to this equality verifying the assertion ii) of Theorem A

$$\mu^\#(x_i) = x_i \otimes 1 + 1 \otimes x_i.$$

Finally, suppose that the group G is simple, and that in addition to (3.2) one has $\mathcal{F}(G) = \Lambda(x'_1, \dots, x'_n)$ with

$$\deg x_i = \deg x'_i \text{ and } \mu^\#(x'_i) = x'_i \otimes 1 + 1 \otimes x'_i, 1 \leq i \leq n.$$

Let $D \subset \mathcal{F}(G)$ be the subring consists of the decomposable elements in the positive degrees. Since both $\{x'_1, \dots, x'_n\}$ and $\{x_1, \dots, x_n\}$ are basis of the free \mathbb{Z} -module $\mathcal{F}(G)/D$, and since $2 = r_1 < \dots < r_n$, the transition functions on $\mathcal{F}(G)$ between these two sets of generators take the forms

$$x'_i = \pm x_i + d_i \text{ with } d_i \in D, 1 \leq i \leq n.$$

Applying the co-product $\mu^\#$ to both sides, and using the relations $\mu^\#(y) = y \otimes 1 + 1 \otimes y$ for $y = x_i$ or x'_i to simplify the resulting equation, yields that

$$d_i \otimes 1 + 1 \otimes d_i = \mu^\#(d_i).$$

With $d_i \in D$ this implies $d_i = 0$, completing the proof of Theorem A. \square

Proof of Theorem B. Let G be a simple Lie group with maximal torus T and non-trivial center $\mathcal{Z}(G)$. Since the covering $c : G \rightarrow PG = G/\mathcal{Z}(G)$ carries T to the maximal torus $T' := c(T)$ on PG , it can be viewed as a bundle map over the identity of G/T

$$\begin{array}{ccc} T & \xrightarrow{c'} & T' \\ \cap & & \cap \\ G & \xrightarrow{c} & PG \quad (\text{note that } G/T = PG/T'), \\ \pi \downarrow & & \pi' \downarrow \\ G/T & = & PG/T' \end{array}$$

hence induces a map $c^* : E_2^{*,*}(PG) \rightarrow E_2^{*,*}(G)$ of Koszul complexes. Assume by Lemma 2.3 that $\{\rho'_1, \dots, \rho'_n\}$ is a basis of the $\mathcal{A}(PG)$ -module $E_3^{*,1}(PG)$, and that $\{\rho_1, \dots, \rho_n\}$ is a basis of the $\mathcal{A}^*(G)$ -module $E_3^{*,1}(G)$. Then, by the proof of Theorem A

$$\begin{aligned}\mathcal{F}(PG) &= \Lambda(y_1, \dots, y_n) \text{ with } y_i := q \circ \kappa(\rho'_i) \in \mathcal{F}(PG), \\ \mathcal{F}(G) &= \Lambda(x_1, \dots, x_n) \text{ with } x_i := q \circ \kappa(\rho_i) \in \mathcal{F}(G).\end{aligned}$$

On the other hand, for the degree reason $\deg \rho_i = \deg \rho'_i$ we can assume by i) and Lemma 2.4 that

$$(3.3) \quad c^*(\rho'_i) = a_i \rho_i + b_{i-1} \rho_{i-1} + \dots + b_1 \rho_1, \quad a_i \in \mathbb{Z}, \quad b_j \in \mathcal{A}^+(G),$$

where $\mathcal{A}^+(G)$ denotes the subring of $\mathcal{A}^*(G)$ consisting of elements in the positive degrees. Since the ring $\mathcal{A}^+(G)$ is always finite (3.3) implies that $c^\#(y_i) = a_i \cdot x_i$ on the quotient $\mathcal{F}(G)$, where we can assume $a_i \geq 0$ by modifying the sign of the generator x_i whenever is necessary, to get the assertion i) of Theorem B.

Finally, applying c^* to the generator $\rho'_1 \cdots \rho'_n$ of $E_3^{m,n}(PG) = H^{m+n}(PG) = \mathbb{Z}$ (see Lemma 2.3 and (2.3)), and noting that the partial sum $b_{i-1} \rho_{i-1} + \dots + b_1 \rho_1$ in (3.3) is of finite order, we get

$$(3.4) \quad c^*(\rho'_1 \cdots \rho'_n) = (a_1 \cdots a_n) \rho_1 \cdots \rho_n \text{ on } E_3^{m,n}(G) = H^{m+n}(G) = \mathbb{Z}.$$

Since the mapping degree of c is the order $|\mathcal{Z}(G)|$ of the center, and since the product $\rho_1 \cdots \rho_n$ generates the group $E_3^{m,n}(G) = H^{m+n}(G) = \mathbb{Z}$ by Lemma 2.3, we get from (3.4) that $a_1 \cdots a_n = |\mathcal{Z}(G)|$, completing the proof of Theorem B. \square

Proof of Theorem C. Since the second homotopy group of PG is trivial, the homotopy exact sequence of π' contains the free resolution of the group $\pi_1(PG)$

$$0 \rightarrow \pi_2(G/T) \rightarrow \pi_1(T') \xrightarrow{j^*} \pi_1(PG)(= \mathcal{Z}(G)) \rightarrow 0,$$

where j is the inclusion of the maximal torus. Applying the co-functor $Hom(\cdot, \mathbb{Z})$ to this sequence, and using the Hurewicz isomorphisms

$$\pi_2(G/T) = H_2(G/T), \quad \pi_1(T') = H_1(T'), \quad \pi_1(PG) = H_1(PG)$$

to substitute the relevant groups, one obtains the exact sequence

$$(3.6) \quad 0 \rightarrow H^1(PG) \xrightarrow{j^*} H^1(T') \xrightarrow{\tau'} H^2(G/T) \xrightarrow{\pi'^*} Tor H^2(PG) = \mathcal{Z}(G) \rightarrow 0$$

in cohomologies, where τ' is the transgression in π' . It follows that

$$H^2(G/T)/\text{Im } \tau' \cong H^2(PG) = Tor H^2(PG) = \mathcal{Z}(G),$$

where the first equality follows from $H^2(PG) \otimes \mathbb{R} = 0$ by Theorem 1.1. With $\mathcal{A}^2(PG) = H^2(G/T)/\text{Im } \tau'$ this implies that the inclusion $\mathcal{A}(PG) \subseteq H^*(PG)$ restricts to an isomorphism in degree 2. Therefore, the subring $J(PG)$ of $H^*(PG)$ generated by $H^2(PG)$ agrees with the subring of $\mathcal{A}^*(PG)$ generated by $\mathcal{A}^2(PG)$. In particular, one reads out $J(PG)$ from the formulae of the ring $\mathcal{A}^*(PG)$ presented in Lemma 2.2, showing Theorem C. \square

4 An exact sequence for cyclic coverings

As a covering $c : G \rightarrow G'$ on Lie groups is a group homomorphism we have $\ker c \subseteq \mathcal{Z}(G)$. The covering c is called *cyclic* if $\ker c$ is a cyclic subgroup. We note that

- a) if G is one of the simply connected Lie groups in Table 1 with $G \neq Spin(2n)$, then the covering $G \rightarrow PG$ is always cyclic;
- b) if $G = Spin(2n)$ the covering $G \rightarrow PG$ can be decomposed into the composition $Spin(2n) \xrightarrow{c_1} SO(2n) \xrightarrow{c_2} PSpin(2n)$ of two cyclic ones, both with order 2.

In addition, for a cyclic covering $G \rightarrow G'$ between simple Lie groups, the multi-degree $\mathcal{D}(G, G')$ is also defined by the proof of Theorem B. Summarizing, to show Theorem D it suffices to compute the invariant $\mathcal{D}(G, G')$ for the cyclic coverings. In this section we establish an exact sequence, that reduces this task to the calculation with the Chow ring $\mathcal{A}^*(G')$ of the group G' .

Assume that $c : G \rightarrow G'$ is a cyclic covering on simple Lie groups. The *central extension* of c is the principle $U(1)$ -fibration over the group G'

$$(4.1) \quad 0 \rightarrow U(1) \rightarrow \tilde{G} := G \times_{\ker c} U(1) \xrightarrow{C} G' \rightarrow 0,$$

where $\ker c$ acts on the cycle $U(1)$ as the anti-clockwise rotation through the angle $2\pi/|\ker c|$, and where \tilde{G} is furnished with the obvious group structure. Moreover, fix once for all a maximal torus T on G , $\dim T = n$. Then both

$$\tilde{T} := T \times_{\ker c} U(1) \subset \tilde{G} \text{ and } T' := c(T) \subset G'$$

are respectively maximal torus of the corresponding Lie groups, while the $U(1)$ -fibration C can be regarded as a bundle map between two torus fibrations:

$$(4.2) \quad \begin{array}{ccccc} U(1) & \hookrightarrow & \tilde{T} & \xrightarrow{C'} & T' \\ \parallel & & \cap & & \cap \\ U(1) & \hookrightarrow & \tilde{G} & \xrightarrow{C} & G' \quad (\text{see (2.1)}). \\ & & \tilde{\pi} \downarrow & & \pi' \downarrow \\ & & G/T & = & G'/T \end{array}$$

Since both \tilde{T} and T' are torus groups, the restriction C' of C on \tilde{T} is splittable, implying that the group $H^1(\tilde{T})$ admits a basis $\{t_0, t_1, \dots, t_n\}$ so that the induced map C'^* carries the ring $H^*(T')$ isomorphically to the subring $\Lambda(t_1, \dots, t_n)$ of $H^*(\tilde{T}) = \Lambda(t_0, t_1, \dots, t_n)$. It follows that

Lemma 4.1. *For a cyclic covering $c : G \rightarrow G'$ of simple Lie groups we have*

$$\begin{aligned} E_2^{*,*}(\tilde{G}) &= H^*(G/T) \otimes \Lambda(t_0, t_1, \dots, t_n); \\ E_2^{*,*}(G') &= H^*(G/T) \otimes \Lambda(t_1, \dots, t_n), \end{aligned}$$

on which C induces a map $C^* : E_2^{*,*}(G') \rightarrow E_2^{*,*}(\tilde{G})$ of Koszul complexes with

- i) $C^*(z \otimes t) = z \otimes t$, where $z \in H^*(G/T)$, $t \in H^*(T')$;
- ii) the transgression τ' of π' is the restriction of the transgression $\tilde{\tau}$ of $\tilde{\pi}$ to the subgroup $H^1(T') \subset H^1(\tilde{T})$.
- iii) the Euler class of the $U(1)$ -fibration C over G' is
$$\omega = [\tilde{\tau}(t_0) \otimes 1] \in E_3^{2,0}(G') = H^2(G'),$$

where $[x]$ denotes the cohomology class of a d'_2 -cocycle $x \in E_2^{2,k}(G')$, and where the identification $E_3^{2,0}(G') = H^2(G')$ follows from the proof of Theorem C. \square

By Lemma 4.1 the map C^* fits into the short exact sequence

$$(4.3) \quad 0 \rightarrow E_2^{*,k}(G') \xrightarrow{C^*} E_2^{*,k}(\tilde{G}) \xrightarrow{\theta} E_2^{*,k-1}(G') \rightarrow 0,$$

of Koszul complexes, where the map θ is evaluated by the simple rule:

$$(4.4) \quad \theta(x \otimes y) = x \otimes y_2 \text{ if } y = y_1 + t_0 \cdot y_2 \text{ with } y_1, y_2 \in H^*(T').$$

Moreover, with $\text{rank} H^2(G/T) = n$ and $\dim \tilde{T} = n + 1$, the transgression $\tilde{\tau}$ satisfies that $\ker \tilde{\tau} = \mathbb{Z} \subseteq H^1(\tilde{T})$. Taking a generator $s \in \ker \tilde{\tau}$ and noting that $1 \otimes s \in E_2^{0,1}(\tilde{G})$ is \tilde{d}_2 -closed by i) of Lemma 2.1, we get the cohomology class

$$\rho_0 = [1 \otimes s] \in E_3^{0,1}(\tilde{G}).$$

Theorem 4.2. *For a cyclic covering $c : G \rightarrow G'$ one has the exact sequence*

$$(4.5) \quad 0 \rightarrow E_3^{*,*}(G') / \langle \omega \rangle \xrightarrow{C^*} E_3^{*,*}(\tilde{G}) \xrightarrow{\theta} E_3^{*,*}(G') \xrightarrow{\omega} \omega \cdot E_3^{*,*}(G') \rightarrow 0,$$

in which

$$(4.6) \quad E_3^{*,*}(\tilde{G}) = E_3^{*,*}(G) \otimes \Lambda(\rho_0),$$

where ω is the Euler class of the central extension C of c , and where for any $z \in E_3^{*,*}(G')$, $y \in E_3^{*,*}(\tilde{G})$,

- i) $\theta(C^*(z) \cdot y) = z \cdot \theta(y)$;
- ii) $\theta(\rho_0) = |\ker c| \in E_3^{0,0}(G') = \mathbb{Z}$;
- iii) $\omega(z) = \omega \cdot z$;
- iv) $C^*(z) \equiv c^*(z) \otimes 1 \pmod{\rho_0}$ (with respect to (4.6)).

Proof. The short exact sequence (4.3) of Koszul complexes yields the long exact sequence in cohomologies

$$\cdots \rightarrow E_3^{*,r}(G') \xrightarrow{C^*} E_3^{*,r}(\tilde{G}) \xrightarrow{\bar{\theta}} E_3^{*,r-1}(G') \xrightarrow{\omega} E_3^{*,r-1}(G') \xrightarrow{C^*} \cdots$$

for which properties i), ii) and iii) follows easily from Lemma 4.1. Since $\ker C^* = \omega \cdot E_3^{*,*}(G') \subset E_3^{*,*}(G')$ (by the exactness) we obtain (4.5) from

$$\text{Im } \omega = \omega \cdot E_3^{*,*}(G') \text{ and } \text{coker } \omega = E_3^{*,*}(G') / \text{Im } \omega = E_3^{*,*}(G') / \langle \omega \rangle.$$

It remains to establish the decomposition (4.6), together with the relation iv).

In addition to (4.2) the group \tilde{G} also fits into the fibrations

$$(4.7) \quad \begin{array}{ccccccc} 0 & \rightarrow & T & \hookrightarrow & \tilde{T} & \xrightarrow{g'} & U(1) \rightarrow 0 \\ & & \cap & & \cap & & \parallel \\ & & G & \xrightarrow{i} & \tilde{G} & \xrightarrow{g} & U(1) \rightarrow 0 \\ & & \pi \downarrow & & \tilde{\pi} \downarrow & & \\ & & G/T & = & \tilde{G}/T & & \end{array}$$

where g is the quotient of the projection $G \times U(1) \rightarrow U(1)$ by $\ker c$; g' is the restriction of g on \tilde{T} ; and where the upper two rows are short exact sequences of Lie groups. It follows from Lemma 2.1 that $E_2^{*,*}(\tilde{G}) = E_2^{*,*}(G) \otimes \Lambda(s)$. With $\rho_0 = [1 \otimes s]$ by our convention we obtain (4.6) from the Künnth formula. Finally, the relation iv) is transparent, since the inclusion i in (4.7) identifies G with the normal subgroup $q^{-1}(1)$ of \tilde{G} that satisfies the relation $C \circ i = c$. \square

Recall that a $U(1)$ -fibration $E \rightarrow X$ over a CW-complex X is classified by its Euler class $\omega \in H^2(X)$. The following result, fairly transparent in the context of [1], provides a geometric interpretation of the generators of the rings $J(PG)$ given in Theorem C.

Lemma 4.3. *Let G be a simply connected simple Lie group given in Table 1.*

i) if $G \neq Spin(2^t(2b+1))$ with $t \geq 2$, the Euler class of the central extension of the cyclic covering $c : G \rightarrow PG$ is the generator $\omega \in J(PG)$;

ii) if $G = Spin(2n)$ with $n \equiv 0 \pmod{2}$, the Euler class of the central extension of the 2 sheets covering $c : SO(2n) \rightarrow PSpin(2n)$ is the generator $\omega_1 \in J(PSpin(2n))$. \square

It is crucial to note in (4.5) that, with respect to the bi-gradation on $E_3^{s,t}$ imposed by the "base degrees s " and "fiber degrees t ", the map C^* preserves both the base and fiber degrees; the map θ preserves the base degrees, but reduces the fiber degrees by 1; and that the map ω increases the base degrees by 2, and preserves the fiber degrees. In particular, for each $k \geq 0$ one has by (4.5) the exact sequences with four terms

$$(4.8) \quad 0 \rightarrow E_3^{2k,1}(PG)/\langle \omega \rangle \xrightarrow{C^*} E_3^{2k,1}(\tilde{G}) \xrightarrow{\theta} \mathcal{A}^{2k}(PG) \xrightarrow{\omega} \omega \cdot \mathcal{A}^{2k}(PG) \rightarrow 0$$

where the groups $E_3^{2k,0}(PG) = \mathcal{A}^{2k}(PG)$ and $E_3^{2k,0}(\tilde{G}) = \mathcal{A}^{2k}(G)$ have been decided by Lemma 2.2. Furthermore, if $\{\rho'_1, \dots, \rho'_n\}$ is a basis of the $\mathcal{A}^*(PG)$ -module $E_3^{*,1}(PG)$, and $\{\rho_1, \dots, \rho_n\}$ is a basis of the $\mathcal{A}^*(G)$ -module $E_3^{*,1}(G)$, then, with $\mathcal{A}(PG)/\langle \omega \rangle = \mathcal{A}(G)$ by Lemma 2.2,

$$(4.9) \quad \begin{aligned} E_3^{*,1}(\tilde{G}) &\text{ is a } \mathcal{A}^*(G)\text{-module with basis } \{\rho_0, \rho_1, \dots, \rho_n\} \text{ by (4.6);} \\ E_3^{*,1}(PG)/\langle \omega \rangle &\text{ is a } \mathcal{A}^*(G)\text{-module with basis } \{\rho'_1, \dots, \rho'_n\}. \end{aligned}$$

Proof of Theorem D. Let G be a simply connected Lie group given in Table 1, and assume that $D(G, PG) = \{a_1, \dots, a_n\}$. By ii) of Theorem B we have

$$(4.10) \quad a_1 \cdots a_n = |\mathcal{Z}(G)|.$$

In addition, by the naturality of the map κ in (2.12) with respect to bundle maps, we have the commutative diagram

$$(4.11) \quad \begin{array}{ccccccc} E_3^{2k,1}(PG) & \xrightarrow{\kappa} & H^*(PG) & \xrightarrow{q} & \mathcal{F}(PG) \\ C^* \downarrow & & C^* \downarrow & & C^\# \downarrow \\ E_3^{2k,1}(\tilde{G}) & \xrightarrow{\kappa} & H^*(\tilde{G}) & \xrightarrow{q} & \mathcal{F}(\tilde{G}) \end{array}$$

allowing us to calculate the eigenvalues a_1, \dots, a_n of $c^\#$ (see i) of Theorem B) by computing with the action of C^* on $E_3^{2k,1}(PG)$. Granted with the relations (4.8), (4.9) and (4.10), together with the diagram (4.11), the proof of Theorem D will be given in the following order

$$G = SU(n), Sp(n), E_7, E_6, Spin(2n+1) \text{ and } Spin(2n),$$

where $\mathbb{Z}_m\{x\}$ (resp. $\mathbb{Z}\{x\}$) denotes the cyclic group of order m (resp. of order ∞) with generator x .

Case 1. $G = SU(n)$. For each $1 \leq k \leq n-1$ we have by (4.8) the exact sequence

$$0 \rightarrow \mathbb{Z}\{\rho'_k\} \xrightarrow{a_k} \mathbb{Z}\{\rho_k\} \rightarrow \mathbb{Z}_{b_{n,k}}\{\omega^k\} \xrightarrow{\omega} \omega \cdot \mathbb{Z}_{b_{n,k}}\{\omega^k\} \rightarrow 0,$$

where $A^{2k}(PG) = \mathbb{Z}_{b_{n,k}}\{\omega^k\}$ by (2.6), and where

$$E_3^{2k,1}(PG)/\langle \omega \rangle = \mathbb{Z}\{\rho'_k\}, E_3^{2k,1}(\tilde{G}) = \mathbb{Z}\{\rho_k\} \text{ by (4.9).}$$

As the order of the power ω^r is precisely $b_{n,r}$, we get by the exactness that $\ker \omega = \mathbb{Z}_{b_{n,k}/b_{n,k+1}}\{\omega\}$, showing $a_k = \frac{b_{n,k}}{b_{n,k+1}}$, as that stated in Theorem D.

Case 2. $G = Sp(n)$ with $n = 2^r(2b+1)$. Taking in (4.8) that $k = 2^{r+1} - 1$ we get the exact sequence

$$0 \rightarrow \mathbb{Z}\{\rho'_{2^r}\} \xrightarrow{a_{2^r}} \mathbb{Z}\{\rho_{2^r}\} \xrightarrow{\theta} \mathbb{Z}_2\{\omega^{2^{r+1}-1}\} \xrightarrow{\omega} \omega \cdot \mathbb{Z}_2\{\omega^{2^{r+1}-1}\} = 0,$$

where $A^{2k}(PG) = \mathbb{Z}_2\{\omega^k\}$ and $\omega \cdot \mathbb{Z}_2\{\omega^k\} = 0$ by (2.7), and where

$$E_3^{2k,1}(\tilde{G}) = \mathbb{Z}\{\rho_{2^r}\}, E_3^{2k,1}(PG)/\langle \omega \rangle = \mathbb{Z}\{\rho'_{2^r}\} \text{ by (4.9).}$$

This shows that $a_{2^r} = 2$. Moreover, with $|\mathcal{Z}(G)| = 2$ the relation (4.10) forces $a_i = 1$ for $i \neq 2^r$, verifying Theorem D for $G = Sp(n)$.

Case 3. $G = E_7$. Taking $k = 1$ in (4.8) we obtain the short exact sequence

$$0 \rightarrow \mathbb{Z}\{\rho'_1\} \xrightarrow{a_1} \mathbb{Z}\{\rho_1\} \xrightarrow{\theta} \mathbb{Z}_2\{\omega\} \xrightarrow{\omega} \omega \cdot \mathbb{Z}_2\{\omega\} = 0$$

where $A^2(PG) = \ker \omega = \mathbb{Z}_2\{\omega\}$ by (2.11), and where

$$E_3^{2,1}(\tilde{G}) = \mathbb{Z}\{\rho_1\}, E_3^{2,1}(PG)/\langle \omega \rangle = \mathbb{Z}\{\rho'_1\} \text{ by (4.9).}$$

It shows that $a_1 = 2$. Consequently, $a_2 = \dots = a_7 = 1$ by (4.10).

Case 4. $G = E_6$. Taking $k = 8$ in (4.8) we get the short exact sequence

$$0 \rightarrow E_3^{16,1}(PG)/\langle \omega \rangle \xrightarrow{C^*} E_3^{16,1}(\tilde{G}) \xrightarrow{\theta} \mathbb{Z}_3\{\omega^8\} \rightarrow 0$$

where $A^{16}(PG) = \mathbb{Z}_3\{\omega^8\} \oplus \mathbb{Z}_3\{x_4^2\}$ with $\ker \omega = \mathbb{Z}_3\{\omega^8\}$ by (2.10), and where

$$E_3^{16,1}(\tilde{G}) = \mathbb{Z}\{\rho_5\} \oplus x_3 \cdot \mathbb{Z}\{\rho_3\} \oplus x_4 \cdot \mathbb{Z}\{\rho_2\},$$

$$E_3^{16,1}(PG)/\langle \omega \rangle = \mathbb{Z}\{\rho'_5\} \oplus x_3 \cdot \mathbb{Z}\{\rho'_3\} \oplus x_4 \cdot \mathbb{Z}\{\rho'_2\} \text{ by (4.9).}$$

Moreover, by the relation i) of Theorem 4.2, together $\text{Im } \theta = \mathbb{Z}_3\{\omega^8\}$,

$$\theta(x_3 \cdot \rho_3) = x_3 \cdot \theta(\rho_3) = 0; \theta(x_4 \cdot \rho_2) = x_4 \cdot \theta(\rho_2) = 0.$$

These imply, by the exactness, that $\theta(\rho_5) = \omega^8$. Since $\theta(\rho_5)$ is of order 3 there must be $C^*(\rho'_5) = 3\rho_5$, showing $a_5 = 3$. As result $a_i = 1$ for $i \neq 5$ by (4.10).

Case 5. $G = Spin(2n+1)$, $2^s \leq n < 2^{s+1}$. By the formula (2.8) of $\mathcal{A}^*(PG)$

$$\ker\{A^{2k}(PG) \xrightarrow{\omega} \omega \cdot A^{2k}(PG)\} = \begin{cases} 0 & \text{if } 0 \leq k < 2^{s+1} - 1; \\ \mathbb{Z}_2\{\omega^{2^{s+1}-1}\} & \text{if } k = 2^{s+1} - 1, \end{cases}$$

where $\omega = x_1$. By (4.8) we have the exact sequences

$$(4.12) \quad 0 \rightarrow E_3^{2k,1}(PG)/\langle \omega \rangle \xrightarrow{C^*} E_3^{2k,1}(\tilde{G}) \xrightarrow{\theta} 0 \text{ for } 1 \leq k < 2^{s+1} - 1;$$

$$(4.13) \quad 0 \rightarrow E_3^{2(2^{s+1}-1),1}(PG)/\langle \omega \rangle \xrightarrow{C^*} E_3^{2(2^{s+1}-1),1}(\tilde{G}) \xrightarrow{\theta} \mathbb{Z}_2\{\omega^{2^{s+1}-1}\} \rightarrow 0.$$

In particular, by (4.12)

$$(4.14) \quad \theta(\rho_i) = 0, \quad 1 < i < 2^s.$$

Moreover, by (4.9) the groups $E_3^{2(2^{s+1}-1),1}(\tilde{G})$ and $E_3^{2(2^{s+1}-1),1}(PG)/\langle \omega \rangle$ in (4.13) are spanned respectively by the following elements

$$\begin{aligned} &\rho_{2^s}, a_i \cdot \rho_i, \quad 1 \leq i < 2^s, \quad a_i \in \mathcal{A}^{4(2^s-i)}(G) \text{ and} \\ &\rho'_{2^s}, a_i \cdot \rho'_i, \quad 1 \leq i < 2^s, \quad a_i \in \mathcal{A}^{4(2^s-i)}(G), \end{aligned}$$

with respect to them, by i) of Lemma 4.2 and by (4.14),

$$\theta(a_i \cdot \rho_i) = a_i \cdot \theta(\rho_i) = 0, \quad 1 \leq i < 2^s,$$

By the exactness of the sequence (4.13) we get

$$\theta(\rho_{2^s}) = \omega^{2^{s+1}-1}, \text{ hence } C^*(\rho'_{2^s}) = 2\rho_{2^s},$$

showing $a_{2^s} = 2$. With $\mathcal{Z}(G) = \mathbb{Z}_2$ we get $a_i = 1$ for $i \neq 2^s$ by (4.10).

Case 6. $G = Spin(2n)$, $2^s < n = 2b+1 \leq 2^{s+1}$. According to (2.9) we have

$$\mathcal{A}^*(PG) = \frac{\mathbb{Z}[\omega, x_3, x_5, \dots, x_{2[\frac{n}{2}-1]}]}{\langle 4\omega, 2\omega^2, \omega^{2^{s+1}}, 2x_{2i-1}, x_{2i-1}^{k_i}; 2 \leq i \leq [\frac{n}{2}] \rangle},$$

where $\omega = x_1$. It implies that

$$\ker\{A^{2k}(PG) \xrightarrow{\omega} \omega \cdot A^{2k}(PG)\} = \begin{cases} \mathbb{Z}_2\{2\omega\} & \text{if } k = 1; \\ \mathbb{Z}_2\{\omega^{2^{s+1}-1}\} & \text{if } k = 2^{s+1} - 1. \end{cases}$$

By (4.8) we have the exact sequences

$$0 \rightarrow \mathbb{Z}\{\rho'_1\} \xrightarrow{C^*} \mathbb{Z}\{\rho_1\} \xrightarrow{\theta} \mathbb{Z}_2\{2\omega\} \rightarrow 0 \text{ and}$$

$$0 \rightarrow E_3^{2(2^{s+1}-1),1}(PG)/\langle \omega \rangle \xrightarrow{C^*} E_3^{2(2^{s+1}-1),1}(\tilde{G}) \xrightarrow{\theta} \mathbb{Z}_2\{\omega^{2^{s+1}-1}\} \rightarrow 0$$

implying respectively that $a_1 = 2$ and $a_{2^s} = 2$ (by an argument similar to the previous case). With $\mathcal{Z}(G) = \mathbb{Z}_4$ we get $a_i = 1$ for $i \neq 1$ or 2^s by (4.10).

Case 7. $G = Spin(2n)$, $2^s < n = 2^t(2b+1) \leq 2^{s+1}$, $t \geq 1$. Decompose the covering $c : G \rightarrow PG$ into the composition of two cyclic coverings of order 2

$$c = c_2 \circ c_1 : Spin(2n) \xrightarrow{c_1} SO(2n) \xrightarrow{c_2} PSpin(2n),$$

where, in addition to formulae (2.9) of $\mathcal{A}^*(G)$ for $G = Spin(2n)$, $PSpin(2n)$,

$$\mathcal{A}^*(SO(2n)) = \frac{\mathbb{Z}[x_1, x_3, x_5, \dots, x_{2[\frac{n}{2}-1]}]}{\langle 2x_{2i-1}, x_{2i-1}^{h_i}; 1 \leq i \leq [\frac{n}{2}] \rangle}, \deg x_k = 2k, h_i = 2^{\lfloor \ln \frac{n-1}{i} \rfloor + 1}.$$

by Marlin [23]. The same calculation as that in Case 5 shows that

$$\mathcal{D}(Spin(2n), SO(2n)) = \{1, \dots, 1, 2_{(2^s)}, 1, \dots, 1\},$$

$$\mathcal{D}(SO(2n), PSpin(2n)) = \{1, \dots, 1, 2_{(2^{t-1})}, 1, \dots, 1\}.$$

For $G = Spin(2n)$ Theorem D is verified by the multiplicative property of multi-degree with respect to the composition of coverings. \square

For a cyclic covering $c : G \rightarrow G'$ on simple Lie groups let $J(\omega) \subseteq H^*(G')$ be the subring generated by the Euler class ω of the central extension $\tilde{G} \rightarrow G'$, and let b_r be the order of the power ω^r in $J(\omega)$. Define the characteristic of the ring $J(\omega)$ to be the sequence $Ch(J(\omega)) := \{b_1, b_2, \dots\}$ of integers. It is clear that

- i) b_{r+1} divides b_r for $r \geq 1$; and
- ii) $J(\omega) = J(PG)$ for $G \neq Spin(2n)$ with $n \equiv 1 \pmod{2}$

by Theorem C. For examples, we get from ii) that

$$Ch(J(\omega)) = \{3, \dots, 3, 1\} \text{ for the three sheets covering } E_6 \rightarrow PE_6$$

$$Ch(J(\omega)) = \{2, 1, \dots, 1\} \text{ for the two sheets covering } E_7 \rightarrow PE_7.$$

Since the proof of Theorem D goes through (essentially) all the possible cyclic coverings between simple Lie groups, it implies the following result that is independent of the types of simple Lie groups.

Theorem 4.4. *For any cyclic covering $c : G \rightarrow G'$ between simple Lie groups the two invariants $\mathcal{D}(G, G')$ and $Ch(J(\omega))$ are related by the equalities*

$$a_k = b_{r_k} / b_{r_k+1}, 1 \leq k \leq n, \text{ where } I_G = \{r_1, \dots, r_n\}. \square$$

Remarks 4.5. The passage from a cyclic covering $c : G \rightarrow G'$ to its central extension $\tilde{G} \rightarrow G'$ is an useful construction in geometry. As examples, for the cyclic covering $SU(n) \rightarrow PSU(n)$ of order n we have $\tilde{G} = U(n)$, the unitary group of rank n ; for the cyclic covering $Spin(n) \rightarrow SO(n)$ of order 2 we get the spin^c group $\tilde{G} = Spin^c(n)$.

In the context of Schubert calculus the integral cohomologies of all simply connected simple Lie groups G have been constructed in [12]. In [7, 8] the exact sequence (4.5) has been applied to extend the construction to the integral cohomology of the adjoint Lie groups PG . \square

References

- [1] A. Borel, F. Hirzebruch, Characteristic classes and homogeneous space I, Amer. J. Math., vol. 80 (1958), 458-538.
- [2] I. N. Bernstein, I. M. Gel'fand, S. I. Gel'fand, Schubert cells and cohomology of the spaces G/P , Russian Math. Surveys 28 (1973), 1-26.
- [3] R. Bott and H. Samelson, The cohomology ring of G/T , Nat. Acad. Sci. 41(1955), 490-492.
- [4] C. Chevalley, The Betti numbers of the exceptional simple Lie groups, Proceedings of the International Congress of Mathematicians, Cambridge, Mass., Vol. 2 (1950), 21-24.
- [5] C. Chevalley, La théorie des groupes algébriques, Proc. 1958 ICM, Cambridge Univ. Press, 1960, 53-68.
- [6] P. Dijkgraaf, E. Witten, Topological gauge theories and group cohomology, commun. Math. Phys. 129(1990), p.393-429.
- [7] H. Duan, The integral cohomology of the projective unitary group $PU(n)$, math.AT. arXiv:1710.09222.
- [8] H. Duan, Schubert calculus and cohomology of Lie groups, Part II. compact Lie groups. math.AT. arXiv:1502.00410.
- [9] H. Duan, On the Borel transgression in the fibration $G \rightarrow G/T$, Homology, Homotopy and Applications 20(1) (2018), 79-86.
- [10] H. Duan and X. Lin, Topology of unitary groups and the prime orders of binomial coefficients, Sci. China Math. 60 (2017), no.9, 1543-1548.
- [11] H. Duan, Xuezhi Zhao, Schubert presentation of the integral cohomology ring of the flag manifolds G/T , LMS J. Comput. Math. Vol.18, no.1(2015), 489-506.
- [12] H. Duan, Xuezhi Zhao, Schubert calculus and cohomology of Lie groups. Part I. 1-connected Lie groups, math.AT (math.AG). arXiv:0711.2541.
- [13] H. Duan, Xuezhi Zhao, On Schubert's Problem of Characteristics, arXiv:1912.10745 (to appear in Springer proceedings in Mathematics & Statistics).
- [14] E.B. Dynkin, Homologies of compact Lie groups. Amer. Math. Soc. Transl. (2) 12(1959), 251-300.
- [15] E.B. Dynkin, Topological characteristics of homomorphisms of compact Lie groups. Amer. Math. Soc. Transl. (2) 12 (1959), 301-342.
- [16] G. Felder, K.Gawedzki, A. Spectra, Spectra of Wess-Zumino-Witten models with arbitrary simple groups, commun. Math. Phys. 117(1988), p.127-158.
- [17] J. R. Harper, Secondary cohomology operations, Graduate Studies in Mathematics 49., Amer. Math. Soc., 2002.

- [18] H. Hopf, Über die topologie der gruppenmannigfaltigkeiten und ihre verallgemeinerungen, Ann. of Math. 42(1941), 22–52.
- [19] A. Grothendieck, Torsion homologique et sections rationnelles, Sem. C. Chevalley, ENS 1958, exposé 5, Secrétariat Math. IHP, Paris, 1958.
- [20] G. Felder, K. Gawedzki, A. Spectra, Spectra of Wess–Zumino–Witten models with arbitrary simple groups, commun. Math. Phys. 117(1988), p.127–158.
- [21] V.G. Kač, Torsion in cohomology of compact Lie groups and Chow rings of reductive algebraic groups, Invent. Math. 80(1985), no. 1, 69–79.
- [22] J. Leray, Sur l’homologie des groupes de Lie, des espaces homogènes et des espaces fibré principaux, Colloque de topologie Algébrique, Bruxelles (1950), 101–115.
- [23] R. Marlin, Anneaux de Chow des groupes algébriques $SU(n)$, $Sp(n)$, $SO(n)$, $Spin(n)$, G_2 , F_4 , C. R. Acad. Sci. Paris, A 279(1974), 119–122.
- [24] J. McCleary, A user’s guide to spectral sequences, Second edition. Cambridge Studies in Advanced Mathematics, 58, 2001.
- [25] V. Mathai and J. Rosenberg, Group dualities, T–dualities, and twisted K–Theory, J. London Mathematical Society, (2) 97 (2018), 1–23.
- [26] H. Pittie, The integral homology and cohomology rings of $SO(n)$ and $Spin(n)$, J. Pure Appl. Algebra 73(1991), 105–153.
- [27] M. Reeder, On the cohomology of compact Lie groups. Enseign. Math. (2) 41 (1995), no. 3–4, 181–200.
- [28] B. L. van der Waerden, Topologische Begründung des Kalküls der abzählenden Geometrie, Math. Ann. 102 (1930), no. 1, 337–362.
- [29] A. Weil, Foundations of algebraic geometry, American Mathematical Society, Providence, R.I. 1962.
- [30] G.W. Whitehead, Elements of homotopy theory, Graduate Texts in Mathematics, 61. Springer-Verlag, New York-Berlin, 1978.