

Fermi liquid theory for nonlinear transport through a multilevel Anderson impurity

Yoshimichi Teratani

Department of Physics, Osaka City University, Sumiyoshi-ku, Osaka 558-8585, Japan

Rui Sakano

The Institute for Solid State Physics, the University of Tokyo, Kashiwa, Chiba 277-8581, Japan

Akira Oguri

Department of Physics, Osaka City University, Sumiyoshi-ku, Osaka 558-8585, Japan and
Nambu Yoichiro Institute of Theoretical and Experimental Physics, Osaka City University, Osaka 558-8585, Japan
(Dated: March 13, 2024)

We present a microscopic Fermi-liquid view on the low-energy transport through an Anderson impurity with N discrete levels, at arbitrary electron filling N_d . It is applied to nonequilibrium current fluctuations, for which the two-quasiparticle collision integral and the three-body correlations that determine the quasiparticle energy shift play important roles. Using the numerical renormalization group up to $N = 6$, we find that for strong interactions the three-body fluctuations are determined by a single parameter other than the Kondo energy scale in a wide filling range $1 \lesssim N_d \lesssim N - 1$. It significantly affects the current noise for $N > 2$ and the behavior of noise in magnetic fields.

Introduction.— Highly correlated low-energy states of the Kondo systems show fascinating universal behavior [1], which can be described by a Fermi liquid (FL) theory in zero dimension [2–6]. FL behaviors have been observed for the nonlinear current through quantum dots [7, 8] and also the current noise [9–12] which is now one of the most important probes to explore quantum states. Furthermore, in addition to the spin, internal degrees of freedom such as *orbital*, *flavor*, etc., bring an interesting variety in the Kondo effect, occurring in a carbon nanotube [12, 13] and novel quantum systems such as ultracold atomic gases [14] and quark matters [15].

Transport properties of the local FL have successfully been described by the renormalized quasiparticles and their collisions due to the residual interaction, especially at the *symmetric point* where both the particle-hole (PH) and time-reversal (TR) symmetries are present [16–19]. These symmetries are broken in real systems by external fields, such as a gate voltage and a magnetic field. In this case, a single quasiparticle captures the quadratic dependence on frequency ω , temperature T , and bias voltage V not only through the well-investigated *damping rate* but also through the *energy shift*. It has recently been clarified that the quadratic *energy shift* is determined by the three-body correlations between the impurity electrons [20–24]. It shows that the three-body correlations are essential parameters for describing the FL transport.

Despite its importance, the current noise [25–31] has been still less elucidated away from the *symmetric point*. A major milestone was achieved by Mora *et al* [20], who have extended Nozières phenomenological FL theory [3] to give the formula of the nonlinear noise for a PH asymmetric single-orbital Anderson model at zero magnetic field. Further investigation, however, is required to clarify the physics of nonequilibrium fluctuations in the Kondo systems with various internal degrees of freedom.

In this letter, we give a microscopic view on the low-energy transport through a multilevel Anderson impurity for a wide range of electron fillings N_d . It is described in terms of *five* FL parameters, which can be calculated using the numerical renormalization group (NRG) [2] up to $N = 6$. We find that for strong interactions the three-body correlations for N degenerate levels are determined by a single parameter over a wide filling range $1 \lesssim N_d \lesssim N - 1$, which includes the intermediate valence regions. We also provide a current-noise formula for the FL, taking into account all the two-quasiparticle collision processes [32, 33]. It satisfies a Ward identity [4–6] for the Keldysh vertex function, and resolves an essential problem of the current conservation of the correlated electrons under a nonequilibrium condition [16, 25]. We also calculate the nonlinear noise using the NRG, and demonstrate that the internal degrees of freedom give a wide variety to the filling dependence. We also examine the effect of a magnetic field that breaks the TR symmetry, and show that the noise of a spin-1/2 quantum dot exhibits a universal Kondo scaling behavior.

Model.— We consider an N -level Anderson impurity coupled to two leads on the left (L) and right (R):

$$\mathcal{H} = \sum_{\sigma=1}^N \epsilon_{d\sigma} n_{d\sigma} + \sum_{\lambda=L,R} \sum_{\sigma=1}^N v_{\lambda} \left(\psi_{\lambda\sigma}^{\dagger} d_{\sigma} + d_{\sigma}^{\dagger} \psi_{\lambda\sigma} \right) + \sum_{\lambda=L,R} \sum_{\sigma=1}^N \int_{-D}^D d\epsilon \epsilon c_{\epsilon\lambda\sigma}^{\dagger} c_{\epsilon\lambda\sigma} + \frac{U}{2} \sum_{\sigma \neq \sigma'} n_{d\sigma} n_{d\sigma'}. \quad (1)$$

d_{σ}^{\dagger} creates an impurity electron with energy $\epsilon_{d\sigma}$, $n_{d\sigma} \equiv d_{\sigma}^{\dagger} d_{\sigma}$, and U the Coulomb repulsion. Conduction electrons are normalized as $\{c_{\epsilon\lambda\sigma}, c_{\epsilon'\lambda'\sigma'}^{\dagger}\} = \delta_{\lambda\lambda'} \delta_{\sigma\sigma'} \delta(\epsilon - \epsilon')$. The coupling v_{λ} between $\psi_{\lambda\sigma} \equiv \int_{-D}^D d\epsilon \sqrt{\rho_c} c_{\epsilon\lambda\sigma}$ and d_{σ}^{\dagger} yields a resonance of the width $\Delta \equiv \Gamma_L + \Gamma_R$, with $\Gamma_{\lambda} = \pi \rho_c v_{\lambda}^2$, $\rho_c = 1/(2D)$, and D the half band width.

TABLE I. Coefficients C 's introduced in Eq. (5). W 's and Θ 's represent the two- and three-body contributions, respectively.

$C_S = \frac{\pi^2}{192} \left[W_S - \cos 2\delta \left\{ \Theta_I + 3(N-1)\Theta_{II} \right\} \right]$	$W_S \equiv \cos 4\delta + \left[4 + 5 \cos 4\delta + \frac{3}{2}(1 - \cos 4\delta)(N-2) \right] (N-1)(R-1)^2$
$C_V = \frac{\pi^2}{64} \left[W_V + \Theta_I + 3(N-1)\Theta_{II} \right]$	$W_V \equiv - \left[1 + 5(N-1)(R-1)^2 \right] \cos 2\delta$
$C_T = \frac{\pi^2}{48} \left[W_T + \Theta_I + (N-1)\Theta_{II} \right]$	$W_T \equiv - \left[1 + 2(N-1)(R-1)^2 \right] \cos 2\delta$
$C_\kappa^{\text{QD}} = \frac{7\pi^2}{80} \left[W_\kappa^{\text{QD}} + \Theta_I + \frac{5}{21}(N-1)\Theta_{II} \right]$	$W_\kappa^{\text{QD}} \equiv \frac{10-11 \cos 2\delta}{21} - \frac{6}{7}(N-1)(R-1)^2 \cos 2\delta$

In this work, we study the nonlinear current noise [25]

$$S_{\text{noise}}^{\text{QD}} = \int_{-\infty}^{\infty} dt \left\langle \delta \hat{J}(t) \delta \hat{J}(0) + \delta \hat{J}(0) \delta \hat{J}(t) \right\rangle_V. \quad (2)$$

Here, $\delta \hat{J}(t) \equiv \hat{J}(t) - \langle \hat{J}(0) \rangle_V$ is the current fluctuation operator through the quantum dot [34], and $\langle \cdots \rangle_V$ is the Keldysh steady-state average defined at finite bias voltages $eV \equiv \mu_L - \mu_R$ with μ_λ the chemical potential for $\lambda = L, R$ [35]. The average current $J \equiv \langle \hat{J}(0) \rangle_V$ is given by [16],

$$J = \frac{e}{h} \sum_{\sigma} \int_{-\infty}^{\infty} d\omega \left[f_L(\omega) - f_R(\omega) \right] \mathcal{T}_{\sigma}(\omega). \quad (3)$$

Here, $f_{\lambda}(\omega) \equiv [e^{(\omega - \mu_{\lambda})/T} + 1]^{-1}$ the Fermi function, $\mathcal{T}_{\sigma}(\omega) \equiv -\frac{4\Gamma_L\Gamma_R}{\Gamma_L + \Gamma_R} \text{Im} G_{\sigma}^r(\omega)$ the transmission probability, and $G_{\sigma}^r(\omega) = [\omega - \epsilon_{d\sigma} + i\Delta - \Sigma_{\sigma}^r(\omega)]^{-1}$ the retarded Green's function with $\Sigma_{\sigma}^r(\omega)$ the self-energy. From this $\mathcal{T}_{\sigma}(\omega)$, we can also deduce the thermal conductivity κ_{QD} [36] for the heat current $J_Q = -\kappa_{\text{QD}} \delta T$, induced by the temperature difference δT between the two leads [37].

Fermi-liquid parameters.— We investigate low-energy transport up to next leading order. To this end, we ex-

pand $\Sigma_{\sigma}^r(\omega)$ up to terms of order ω^2 , T^2 , and $(eV)^2$ for general N , extending the latest FL description for spin 1/2 case [22, 23]. The expansion coefficients play an important role as the FL parameters.

The phase shift $\delta_{\sigma} \equiv \cot^{-1}(\epsilon_{d\sigma}^*/\Delta)$ is a parameter of primary importance, with $\epsilon_{d\sigma}^* \equiv \epsilon_{d\sigma} + \Sigma_{\sigma}^r(0)|_{T=eV=0}$ the effective impurity level. It determines the occupation number $\langle n_{d\sigma} \rangle = \delta_{\sigma}/\pi$, and the density of states $\rho_{d\sigma} \equiv \sin^2 \delta_{\sigma}/(\pi\Delta)$. The renormalization factor is given by the first derivative $z_{\sigma} \equiv [1 - \frac{\partial \Sigma_{\sigma}^r(\omega)}{\partial \omega}|_{\omega=0}]^{-1}$, defined at $T = eV = 0$. It is also related to the static susceptibility $\chi_{\sigma_1\sigma_2} \equiv \int_0^{1/T} d\tau \langle \delta n_{d\sigma_1}(\tau) \delta n_{d\sigma_2} \rangle$, as $\chi_{\sigma\sigma} \xrightarrow{T \rightarrow 0} \rho_{d\sigma}/z_{\sigma}$, with $\delta n_{d\sigma} \equiv n_{d\sigma} - \langle n_{d\sigma} \rangle$ [4–6]. The second derivative is a complex number, the imaginary part of which corresponds to the single-quasiparticle *damping rate* of order ω^2 , T^2 , and $(eV)^2$ [16, 17]. The real part corresponds to the quadratic *energy shift* that is determined by the nonlinear susceptibility defined at equilibrium [22, 23]:

$$\chi_{\sigma_1\sigma_2\sigma_3}^{[3]} \equiv - \int_0^{1/T} d\tau_3 \int_0^{1/T} d\tau_2 \langle T_{\tau} \delta n_{d\sigma_3}(\tau_3) \delta n_{d\sigma_2}(\tau_2) \delta n_{d\sigma_1} \rangle,$$

with T_{τ} the imaginary-time ordering operator. It can also be written as $\chi_{\sigma_1\sigma_2\sigma_3}^{[3]} = \partial \chi_{\sigma_1\sigma_2}/\partial \epsilon_{d\sigma_3}$, and contributes to the transport when the PH or TR symmetry is broken.

SU(N) symmetric case.— In the case at which the N impurity levels are degenerate $\epsilon_{d\sigma} \equiv \epsilon_d$, the linear susceptibility $\chi_{\sigma\sigma'}$ has only two independent components. The diagonal element determines the energy scale $T^* \equiv 1/(4\chi_{\sigma\sigma})$, by which the T -linear specific heat is scaled as $\mathcal{C}_{\text{imp}} = \frac{N\pi^2}{12}(T/T^*)$. It can also be identified as the Kondo temperature in the strong-coupling limit. The other one is the off-diagonal element $\chi_{\sigma\sigma'}$ for $\sigma \neq \sigma'$, which is related to the Wilson ratio $R \equiv 1 - \chi_{\sigma\sigma'}/\chi_{\sigma\sigma}$ [38]. Similarly, the nonlinear susceptibility has three independent components for $N \geq 3$: the diagonal element $\chi_{\sigma\sigma\sigma}^{[3]}$ and two off-diagonal ones, which can also be expressed in the following form for $\sigma \neq \sigma' \neq \sigma'' \neq \sigma$,

$$\begin{aligned} -(N-1)\chi_{\sigma\sigma'\sigma'}^{[3]} &= \chi_{\sigma\sigma\sigma}^{[3]} - \frac{\partial \chi_{\sigma\sigma}}{\partial \epsilon_d}, \\ \frac{(N-1)(N-2)}{2}\chi_{\sigma\sigma'\sigma''}^{[3]} &= \chi_{\sigma\sigma\sigma}^{[3]} - \frac{\partial \chi_{\sigma\sigma}}{\partial \epsilon_d} + \frac{N-1}{2} \frac{\partial \chi_{\sigma\sigma'}}{\partial \epsilon_d}. \end{aligned} \quad (4)$$

In this work, we obtain the low-energy expansion of $S_{\text{noise}}^{\text{QD}}$, J and κ_{QD} up to next leading order, specifically for symmetric junctions $\Gamma_L = \Gamma_R$ and $\mu_L = -\mu_R = eV/2$:

$$\begin{aligned} S_{\text{noise}}^{\text{QD}} &= \frac{2Ne^2|eV|}{h} \left[\frac{\sin^2 2\delta}{4} + C_S \left(\frac{eV}{T^*} \right)^2 + \cdots \right], \\ \frac{dJ}{dV} &= \frac{Ne^2}{h} \left[\sin^2 \delta - C_T \left(\frac{\pi T}{T^*} \right)^2 - C_V \left(\frac{eV}{T^*} \right)^2 + \cdots \right], \\ \kappa_{\text{QD}} &= \frac{N\pi^2 T}{3h} \left[\sin^2 \delta - C_{\kappa}^{\text{QD}} \left(\frac{\pi T}{T^*} \right)^2 + \cdots \right]. \end{aligned} \quad (5)$$

The explicit expressions of the coefficients C_S , C_V , C_T and C_{κ}^{QD} are listed in table I. Each of these C 's consists of two parts, denoted as W and Θ . The W -part represents two-body contributions which can be described in terms of R and δ . The Θ -part represents dimensionless three-

body contributions:

$$\Theta_I \equiv \frac{\sin 2\delta}{2\pi} \frac{\chi_{\sigma\sigma\sigma}^{[3]}}{\chi_{\sigma\sigma}^2}, \quad \Theta_{II} \equiv \frac{\sin 2\delta}{2\pi} \frac{\chi_{\sigma\sigma'\sigma'}^{[3]}}{\chi_{\sigma\sigma}^2}. \quad (6)$$

Therefore, the low-energy transport of the $SU(N)$ Fermi liquid are determined completely by *five* parameters: δ , T^* , R , Θ_I , and Θ_{II} . These FL parameters can also be deduced experimentally through measurements of the coefficients C 's. We note that another parameter for three different levels, $\Theta_{III} \equiv \frac{\sin 2\delta}{2\pi} \frac{\chi_{\sigma\sigma'\sigma''}^{[3]}}{\chi_{\sigma\sigma}^2}$, does *not* affect C 's for symmetric junctions. Nevertheless, it contributes to the transport for $N \geq 3$ when the tunneling couplings or the chemical potentials are asymmetric.

The nonlinear noise of the Fermi liquid is determined not only by a single-quasiparticle excitation but also by two-quasiparticle collisions described by the Keldysh vertex corrections [25]. In this work, we calculate the vertex function up to order eV [35], extending the diagrammatic approach of Yamada-Yosida [4–6]. Consequently, the collision contributions C_S^{coll} and the single-quasiparticle ones C_S^{qp} yield the nonlinear noise $C_S = C_S^{\text{qp}} + C_S^{\text{coll}}$:

$$C_S^{\text{coll}} = \left[\frac{7 + 5 \cos 4\delta}{2} + \frac{3}{2} (1 - \cos 4\delta) (N - 2) \right] \frac{\tilde{K}^2}{N - 1}$$

with $\tilde{K} \equiv (N - 1)(R - 1)$. The second term in the bracket emerges through the collisions specific to multilevel impurities for $N > 2$, and it vanishes in the $SU(2)$ symmetric case or the PH symmetric case at which $\delta = \pi/2$.

Filling dependence of the FL state.— How does the FL state evolve as the number of levels N and their position ϵ_d vary? As the electron configuration $N_d \equiv \sum_{\sigma} \langle n_{d\sigma} \rangle$ continuously varies with ϵ_d , a different class of the Kondo and valence-fluctuation states emerge for multilevel systems $N > 2$. To our knowledge, however, the behavior of three-body correlations Θ 's that determine the nonlinear transport has not been explored so much, whereas the two-body correlations have been well investigated for $N = 4$ [30, 39, 40]. In this work, we calculate the FL parameters for $N = 4, 6$ with the NRG, using the interleaved algorithm particularly for $N = 6$ [41]. To be specific, we choose the Coulomb interaction to be much larger than the hybridization energy scale: $U/(\pi\Delta) = 5.0$. The results are plotted vs $\xi_d \equiv \epsilon_d + (N - 1)U/2$ in Fig. 1 for (left panels) $N = 4$ and (right panels) $N = 6$.

The top panels of Fig. 1 show the two-body correlations, relating to $\langle n_{d\sigma} \rangle$, $\chi_{\sigma\sigma}$, and $\chi_{\sigma\sigma'}$. We see that $\sin^2 \delta$, which determines $\mathcal{T}_{\sigma}(0)$ at $T = eV = 0$, shows a flat Kondo ridge of the unitary limit $\delta \simeq \pi/2$ near the PH symmetric point $|\xi_d| \lesssim U/2$ where the occupation number is almost locked at $N_d \simeq N/2$. The other Kondo ridges also emerge at ξ_d where N_d approaches an integer: $\xi_d \simeq \pm U$ for $N = 4$, and also $\xi_d \simeq \pm U, \pm 2U$ for $N = 6$.

The renormalization factor z , which determines the energy scale $T^* = z\pi\Delta/(4\sin^2 \delta)$, is also shown in the top




FIG. 1. Fermi-liquid parameters for $SU(N)$ Anderson model are plotted vs $\xi_d \equiv \epsilon_d + (N - 1)U/2$ for $U/(\pi\Delta) = 5.0$, $N = 4$ (left panels) and $N = 6$ (right panels). Top panels: $\sin^2 \delta$, renormalization factor z , and $\tilde{K} \equiv (N - 1)(R - 1)$. Middle panels: Θ_I , $-\tilde{\Theta}_{II} \equiv -(N - 1)\Theta_{II}$, and $\tilde{\Theta}_{III} \equiv \frac{(N - 1)(N - 2)}{2}\Theta_{III}$. Bottom panels: $\frac{48}{\pi^2}C_T$, $\frac{64}{\pi^2}C_V$, $\frac{80}{7\pi^2}C_K^{\text{QD}}$, and $\frac{192}{\pi^2}C_S$.

panels. It is significantly suppressed over a wide range of gate voltages $|\xi_d| \lesssim \frac{N-1}{2}U$, and appears as a broad valley. This valley becomes shallow as N increases, and vanishes in the large N limit [40]. Inside the valley, z has minimums at $\xi_d \simeq \frac{N-2M}{2}U$ for $M = 1, 2, \dots, N-1$, where the occupation number approaches an integer $N_d = M$. At these minimums, the low-energy states can be described by the $SU(N)$ Kondo model in the strong-coupling limit $U \gg \Delta$. We find that z is also suppressed at local maximums corresponding to the intermediate valence states, for both $N = 4$ and 6. In the top panels, the rescaled Wilson ratio \tilde{K} is also shown. It is almost saturated to the universal value $\tilde{K} = 1$ and its derivative becomes very small $\partial\tilde{K}/\partial\epsilon_d \sim 0$ in the whole region of the broad valley $1 \lesssim N_d \lesssim N - 1$. It reveals the fact that not only the charge susceptibility $\chi_c \equiv -\langle n_{d\sigma} \rangle / \partial\epsilon_d = \chi_{\sigma\sigma}(1 - \tilde{K})$ but its derivative $\partial\chi_c/\partial\epsilon_d$ is suppressed in this region.

The three-body correlation Θ_I is plotted in the middle panels of Fig. 1, together with the other two rescaled ones: $-\tilde{\Theta}_{II} \equiv -(N - 1)\Theta_{II}$, and $\tilde{\Theta}_{III} \equiv \frac{(N - 1)(N - 2)}{2}\Theta_{III}$. These Θ 's also show plateau structures due to the Kondo effect at the values of ξ_d corresponding to integer N_d , and almost vanish at $|\xi_d| \lesssim \frac{U}{2}$. We find that these three parameters Θ_I , $-\tilde{\Theta}_{II}$, and $\tilde{\Theta}_{III}$ approach each other very closely over a wide gate-voltage range $|\xi_d| \lesssim \frac{N-2}{2}U$, at




FIG. 2. Nonlinear current-current correlations. Left panel: $F_K \equiv \frac{C_S}{C_V/3}$ vs ξ_d/U for $SU(N)$ symmetric case for $N = 4$ (\bullet, \blacklozenge) and $N = 6$ (\circ, \diamond), for $U/(\pi\Delta) = 1/3$ (diamonds) and $U/(\pi\Delta) = 5$ (circles). Right panel: \overline{C}_S^b vs b/T_K for $N = 2$ at half filling $\epsilon_d = -U/2$, for $U/(\pi\Delta) = 2.5, 3.0, 3.5, 4.0$, with b the magnetic field and T_K the Kondo temperature at $b = 0$.

which $1 \lesssim N_d \lesssim N - 1$. This indicates that contributions of the diagonal element $\chi_{\sigma\sigma\sigma}^{[3]}$ dominate the terms in the right-hand side of Eq. (4); i.e. $\chi_{\sigma\sigma\sigma}^{[3]}$ becomes much greater than $\frac{\partial\chi_{\sigma\sigma}}{\partial\epsilon_d}$ and $\frac{N-1}{2} \frac{\partial\chi_{\sigma\sigma'}}{\partial\epsilon_d}$. It also reveals the fact that not only $\partial\chi_c/\partial\epsilon_d$ but also $\partial\chi_s/\partial\epsilon_d$, the derivative of the spin susceptibility $\chi_s \propto \chi_{\sigma\sigma} - \chi_{\sigma\sigma'}$, becomes much smaller than $(T^*)^{-2}$. Thus, for large U , the FL properties are characterized by *three* parameters δ , T^* and Θ_I over the wide filling range $1 \lesssim N_d \lesssim N - 1$. Outside this region, the Θ 's approach the noninteracting values: $\Theta_I \rightarrow -2$, and the other two vanish as $|\xi_d| \rightarrow \infty$.

Nonequilibrium FL fluctuations.— We show in the following how the transport coefficients evolve as N_d varies continuously. The NRG results are also plotted in Fig. 1. The difference between the C 's near half filling $|\xi_d| \lesssim \frac{U}{2}$ is caused by the two-body contributions W 's as the Θ 's almost vanish. In particular, the T^2 conductance C_T is determined by W_T over the wide filling range $1 \lesssim N_d \lesssim N - 1$ as the three-body contributions almost cancel out $\Theta_I + (N-1)\Theta_{II} \approx 0$, reflecting the suppression of $\partial\chi_c/\partial\epsilon_d$ and $\partial\chi_s/\partial\epsilon_d$ mentioned above. For the thermal conductivity, the three-body contributions become negative in this region, $\Theta_I + \frac{5}{21}(N-1)\Theta_{II} \approx -\frac{16}{21}\tilde{\Theta}_{II}$, but otherwise C_κ^{QD} shows a similar ξ_d dependence to that of C_T . The three-body contributions on the $(eV)^2$ conductance C_V are given by $\Theta_V \equiv \Theta_I + 3(N-1)\Theta_{II}$, which takes a value $\Theta_V \approx 2\tilde{\Theta}_{II}$ in the same filling range. Thus, C_V is significantly enhanced at $N_d \simeq 1$ and $N - 1$ where $-\tilde{\Theta}_{II}$ shows a deep valley. It pushes the tail of the C_V curve outside than that of the C_T in the valence fluctuation region towards the empty or fully-occupied limit.

The $|eV|^3$ current noise also exhibits the Kondo plateau structures as shown in Fig. 1. For C_S , the three-body contributions enter through Θ_V with a sinusoidal factor: $-\Theta_V \cos 2\delta \approx \frac{\sin 4\delta}{2\pi} \frac{\chi_{\sigma\sigma\sigma}^{[3]}}{\chi_{\sigma\sigma}^2}$ over the range of $1 \lesssim N_d \lesssim N - 1$. In the valence fluctuation regions mentioned above, C_S has a minimum caused by the higher-harmonic $\sin 4\delta$ and $\cos 4\delta$ contributions. We also find

that C_S approaches zero almost simultaneously with C_V at $|\xi_d| \simeq 1.4U$ for $N = 4$, and at $|\xi_d| \simeq 2.4U$ for $N = 6$. This proximity of the zero points affects the behavior of an extended Fano factor F_K , defined as the ratio of order $(eV)^3$ current noise to the nonlinear current [20, 29, 35]:

$$F_K \equiv \lim_{|eV| \rightarrow 0} \frac{S_{\text{noise}}^{\text{QD}} - \frac{2Ne^2|eV|}{h} \frac{\sin^2 2\delta}{4}}{-2|e| \left(J - \frac{Ne^2|V|}{h} \sin^2 \delta \right)} = \frac{C_S}{C_V/3}. \quad (7)$$

This formula for the $SU(N)$ Anderson model includes the result of Mora *et al*, obtained for $N = 2$ at zero magnetic field [20], as a special case. In the strong coupling limit at integer N_d , it also agrees with another noise formula of Mora *et al* for the $SU(N)$ Kondo model [29].

The Fano factor for $N = 4, 6$ is plotted vs ξ_d for two different values of U in the left panel of Fig. 2. It reaches the local maximum $F_K \rightarrow (N-1+9\tilde{K}^2)/(N-1+5\tilde{K}^2)$ at $\xi_d = 0$ [30], and has positive plateaus for large U at integer N_d . In the limit of $|\xi_d| \rightarrow \infty$, the ratio becomes negative and takes the noninteracting value $F_K \rightarrow -1$. By definition, F_K changes sign at the zero points of C_S . It also diverges at the zero point of C_V , where the nonlinear component of J changes direction from backward to forward. Such a singularity already exists for $U = 0$ at $|\xi_d| = \Delta/\sqrt{3}$. For large U , F_K diverges near $|\xi_d| \simeq \frac{N-1}{2}U$ in the valence fluctuation region towards the empty or fully-occupied limit. We can see that sign of the coefficient C_S at the singular points becomes positive for large U , whereas it is negative for small U . Sign change occurs, for both $N = 4$ and 6 , at a finite U between the two examined cases $U/(\pi\Delta) = 1/3$ and 5 . It is associated with the large enhancement of three-body contributions Θ_V occurring in the Kondo regime at $N_d \simeq 1$ and $N - 1$ for $N > 2$. In contrast, the NRG calculations examined so far indicate that sign is always negative in the $SU(2)$ case for any $U \geq 0$ [20, 35]. The main difference is that in the $SU(2)$ case the three-body correlations evolve in the valence fluctuation region where electron correlations become less important and the two-body contributions W_S dominate C_S near the singular point.

Magnetic-field dependence.— We next consider effects of a magnetic field b that breaks the $SU(N)$ and TR symmetries: specifically at half filling for $N = 2$, where $\epsilon_{d\uparrow} = -\frac{U}{2} - b$, $\epsilon_{d\downarrow} = -\frac{U}{2} + b$, and the electron filling is fixed at $\langle n_{d\uparrow} \rangle + \langle n_{d\downarrow} \rangle = 1$. In this case, the transport coefficients can be described also by five FL parameters: magnetization $m_d \equiv \langle n_{d\uparrow} \rangle - \langle n_{d\downarrow} \rangle$, susceptibilities $\chi_{\uparrow\uparrow} = \chi_{\downarrow\downarrow}$ and $\chi_{\uparrow\downarrow} = \chi_{\downarrow\uparrow}$, and three-body correlations $\chi_{\uparrow\uparrow\uparrow}^{[3]} = -\chi_{\downarrow\downarrow\downarrow}^{[3]}$ and $\chi_{\uparrow\uparrow\downarrow}^{[3]} = -\chi_{\uparrow\downarrow\downarrow}^{[3]}$. The nonlinear current for this case has previously been studied [21–23]. However, behavior of its fluctuations has not been clarified so far.

Here, we examine the current noise at $T = 0$ [42]:

$$S_{\text{noise}}^{\text{QD}} = \frac{4e^2|eV|}{h} \left[\frac{\sin^2(\pi m_d)}{4} + \overline{C}_S^b \left(\frac{eV}{T_K} \right)^2 + \dots \right].$$

Note that the second term is scaled by $T_K \equiv T^*|_{b=0}$, the Kondo temperature defined at zero field. Thus, the coefficient \overline{C}_S^b includes all effects of b , which enter through the FL parameters. In the right panel of Fig. 2, NRG results for \overline{C}_S^b are plotted as a function of b/T_K for several different values of U . We find that the nonlinear noise exhibits a universal behavior for $U/(\pi\Delta) \gtrsim 2.0$ in a similar way that the nonlinear current shows [22, 23]. It decreases rapidly as b increases for small fields, changes sign at $b \approx 0.36T_K$, takes a minimum at $b \approx 0.5T_K$, and then approaches zero at $b \gtrsim T_K$. We note that order T^3 thermal conductivity also exhibits the scaling behavior [35]. These observations reflect the fact that the three-body fluctuations show the universal scaling behavior in the Kondo regime without the TR symmetry.

Conclusion.— Nonlinear transport through the $SU(N)$ Anderson impurity has been described in a unified way with five FL parameters. We have demonstrated how the FL state evolves as electron filling N_d varies, using the NRG up to $N = 6$. For strong interactions U , not only charge fluctuations but also the derivatives of charge and spin susceptibilities are suppressed over a wide filling range $1 \lesssim N_d \lesssim N - 1$. It reduces the number of variable FL parameters from five to three, and causes the Kondo plateau structures emerging for all the coefficients C 's. In particular, the three-body contributions on C_V are significantly enhanced at $N_d \simeq 1$ and $N - 1$ for $N > 2$. It also affects the behavior of nonlinear Fano factor F_K in the valence fluctuation region. We have also shown that the nonlinear current noise exhibits the universal magnetic-field scaling in the Kondo regime. The FL parameters can also be deduced from experiments and can be used to predict behaviors of unmeasured observables.

We would like to thank K. Kobayashi, T. Hata, M. Ferrier, R. Deblock, and A. C. Hewson for valuable discussions. This work was supported by JSPS KAKENHI Grand Numbers JP18J10205, JP18K03495, JP26220711, and JST CREST Grant No. JPMJCR1876.

[1] A. C. Hewson, *The Kondo Problem to Heavy Fermions* (Cambridge University Press, 1993).
[2] K. G. Wilson, Rev. Mod. Phys. **47**, 773 (1975).
[3] P. Nozières, J. Low Temp. Phys. **17**, 31 (1974).
[4] K. Yamada, Prog. Theor. Phys. **54**, 316 (1975).
[5] H. Shiba, Prog. Theor. Phys. **54**, 967 (1975).
[6] A. Yoshimori, Prog. Theor. Phys. **55**, 67 (1976).
[7] M. Grobis, I. G. Rau, R. M. Potok, H. Shtrikman, and D. Goldhaber-Gordon, Phys. Rev. Lett. **100**, 246601 (2008).
[8] G. D. Scott, Z. K. Keane, J. W. Ciszek, J. M. Tour, and D. Natelson, Phys. Rev. B **79**, 165413 (2009).
[9] O. Zarchin, M. Zaffalon, M. Heiblum, D. Mahalu, and V. Umansky, Phys. Rev. B **77**, 241303(R) (2008).
[10] T. Delattre, C. Feuillet-Palma, L. G. Herrmann,

P. Morfin, J.-M. Berroir, G. FShu ve, B. PlaXun ais, D. C. Glatthli, M.-S. Choi, C. Mora, and T. Kontos, Nature Physics **5**, 208 (2009).
[11] Y. Yamauchi, K. Sekiguchi, K. Chida, T. Arakawa, S. Nakamura, K. Kobayashi, T. Ono, T. Fujii, and R. Sakano, Phys. Rev. Lett. **106**, 176601 (2011).
[12] M. Ferrier, T. Arakawa, T. Hata, R. Fujiwara, R. Delagrangé, R. Weil, R. Deblock, R. Sakano, A. Oguri, and K. Kobayashi, Nat. Phys. **12**, 230 (2016).
[13] E. A. Laird, F. Kuemmeth, G. A. Steele, K. Grove-Rasmussen, J. Nygård, K. Flensberg, and L. P. Kouwenhoven, Rev. Mod. Phys. **87**, 703 (2015).
[14] K. Ono, J. Kobayashi, Y. Amano, K. Sato, and Y. Takahashi, Phys. Rev. A **99**, 032707 (2019).
[15] S. Yasui and S. Ozaki, Phys. Rev. D **96**, 114027 (2017).
[16] S. Hershfield, J. H. Davies, and J. W. Wilkins, Phys. Rev. B **46**, 7046 (1992).
[17] A. Oguri, Phys. Rev. B **64**, 153305 (2001).
[18] A. A. Aligia, J. Phys.: Condens. Matter **24**, 015306 (2012).
[19] E. Muñoz, C. J. Bolech, and S. Kirchner, Phys. Rev. Lett. **110**, 016601 (2013).
[20] C. Mora, C. P. Moca, J. von Delft, and G. Zaránd, Phys. Rev. B **92**, 075120 (2015).
[21] M. Filippone, C. P. Moca, A. Weichselbaum, J. von Delft, and C. Mora, Phys. Rev. B **98**, 075404 (2018).
[22] A. Oguri and A. C. Hewson, Phys. Rev. Lett. **120**, 126802 (2018).
[23] A. Oguri and A. C. Hewson, Phys. Rev. B **97**, 035435 (2018).
[24] A. Oguri and A. C. Hewson, Phys. Rev. B **98**, 079905(E) (2018).
[25] S. Hershfield, Phys. Rev. B **46**, 7061 (1992).
[26] A. O. Gogolin and A. Komnik, Phys. Rev. Lett. **97**, 016602 (2006).
[27] E. Sela, Y. Oreg, F. von Oppen, and J. Koch, Phys. Rev. Lett. **97**, 086601 (2006).
[28] E. Sela and J. Malecki, Phys. Rev. B **80**, 233103 (2009).
[29] C. Mora, P. Vitushinsky, X. Leyronas, A. A. Clerk, and K. Le Hur, Phys. Rev. B **80**, 155322 (2009).
[30] R. Sakano, T. Fujii, and A. Oguri, Phys. Rev. B **83**, 075440 (2011).
[31] D. B. Karki, C. Mora, J. von Delft, and M. N. Kiselev, Phys. Rev. B **97**, 195403 (2018).
[32] L. P. Pitaevskii and E. M. Lifshitz, *Physical Kinetics* (Butterworth-Heinemann, 1981).
[33] H. Haug and A.-P. Jauho, *Quantum Kinetics in Transport and Optics of Semiconductors* (Springer, 2008).
[34] $\hat{J} \equiv (\Gamma_L \hat{J}_R + \Gamma_R \hat{J}_L)/(\Gamma_L + \Gamma_R)$, with \hat{J} the current flowing between the dot and lead on λ ($= L, R$) side.
[35] See supplemental material for details.
[36] $\kappa_{\text{QD}}^{\text{QD}} \equiv \frac{1}{\hbar T} [\sum_{\sigma} \mathcal{L}_{2,\sigma}^{\text{QD}} - (\sum_{\sigma} \mathcal{L}_{1,\sigma}^{\text{QD}})^2 / (\sum_{\sigma} \mathcal{L}_{0,\sigma}^{\text{QD}})]$, and $\mathcal{L}_{n,\sigma}^{\text{QD}} = \int_{-\infty}^{\infty} d\omega \omega^n (-\frac{\partial f}{\partial \omega}) \mathcal{T}_{\sigma}(\omega)$ for $n = 0, 1, 2$.
[37] T. A. Costi and V. Zlatić, Phys. Rev. B **81**, 235127 (2010).
[38] A. C. Hewson, J. Phys.: Condens. Matter **13**, 10011 (2001).
[39] Y. Nishikawa, D. J. G. Crow, and A. C. Hewson, Phys. Rev. B **82**, 245109 (2010).
[40] A. Oguri, Phys. Rev. B **85**, 155404 (2012).
[41] K. M. Stadler, A. K. Mitchell, J. von Delft, and A. Weichselbaum, Phys. Rev. B **93**, 235101 (2016).
[42] $\overline{C}_S^b = \frac{\pi^2}{192} [\cos(2\pi m_d) + \{4 + 5 \cos(2\pi m_d)\}(R - 1)^2 + \Theta_V^b \cos(\pi m_d)] (\frac{T_K}{T^*})^2$, and $\Theta_V^b \equiv -\frac{\sin(\pi m_d)}{2\pi} \frac{\chi_{\uparrow\uparrow\uparrow}^{[3]} + 3\chi_{\uparrow\uparrow\downarrow}^{[3]}}{\chi_{\uparrow\uparrow}^2}$.

Fermi liquid theory for nonlinear transport through a multilevel Anderson impurity (Supplemental Material)

Yoshimichi Teratani¹, Rui Sakano², and Akira Oguri^{1,3}

¹*Department of Physics, Osaka City University, Sumiyoshi-ku, Osaka 558-8585, Japan*

²*The Institute for Solid State Physics, the University of Tokyo, Kashiwa, Chiba 277-8581, Japan*

³*NITEP, Osaka City University, Sumiyoshi-ku, Osaka 558-8585, Japan*

I. DERIVATION OF THE TRANSPORT COEFFICIENTS C_T , C_V AND C_κ^{QD}

We describe here outline of the derivation of the coefficients C 's, listed in table I in the main text. The steady current J through the quantum dots has been calculated using the formula given in Eq. (3) with the transmission probability, defined by

$$\mathcal{T}_\sigma(\omega) = \frac{-4\Gamma_L\Gamma_R}{\Gamma_L + \Gamma_R} \text{Im} G_\sigma^r(\omega), \quad G_\sigma^r(\omega) = \frac{1}{\omega - \epsilon_{d\sigma} + i\Delta - \Sigma_\sigma^r(\omega)}. \quad (8)$$

We have also calculated the thermal conductivity κ_{QD} , which can be expressed in the following form at $eV = 0$,

$$\kappa_{\text{QD}} = \frac{1}{2\pi\hbar T} \left[\sum_\sigma \mathcal{L}_{2,\sigma}^{\text{QD}} - \frac{\left(\sum_\sigma \mathcal{L}_{1,\sigma}^{\text{QD}}\right)^2}{\sum_\sigma \mathcal{L}_{0,\sigma}^{\text{QD}}} \right], \quad \mathcal{L}_{n,\sigma}^{\text{QD}} \equiv \int_{-\infty}^{\infty} d\omega \omega^n \mathcal{T}_\sigma(\omega) \left(-\frac{\partial f(\omega)}{\partial \omega} \right). \quad (9)$$

The coefficients for the charge and heat currents, C_T , C_V , and C_κ^{QD} , can be deduced from the low-energy expansion of the retarded self-energy $\Sigma_\sigma^r(\omega)$ obtained up to terms of order ω^2 , T^2 , and $(eV)^2$. We note that, in order to determine also the thermopower of quantum dots $S_{\text{TP}}^{\text{QD}}$ up to next leading order, additional terms of order ω^3 and ωT^2 of the self-energy are necessary. This is because the leading term of $S_{\text{TP}}^{\text{QD}}$ already includes the derivative $\rho'_{d\sigma} \equiv \frac{\partial \rho_{d\sigma}(\omega)}{\partial \omega} \Big|_{\omega=0}$ which describes a variation from the ground state:

$$S_{\text{TP}}^{\text{QD}} \equiv \frac{-1}{|e|T} \frac{\sum_\sigma \mathcal{L}_{1,\sigma}^{\text{QD}}}{\sum_\sigma \mathcal{L}_{0,\sigma}^{\text{QD}}} = -\frac{\pi^2}{3|e|} \frac{\sum_\sigma \rho'_{d\sigma}}{\sum_\sigma \rho_{d\sigma}} T + O(T^3), \quad \rho'_{d\sigma} = \frac{\chi_{\sigma\sigma} \sin 2\delta_\sigma}{\Delta}. \quad (10)$$

The expansion coefficients of $\Sigma_\sigma^r(\omega)$ can be expressed in terms of the linear and nonlinear susceptibilities,

$$\text{Im} \Sigma_\sigma^r(\omega) = -\frac{\pi(N-1)}{2} \frac{\chi_{\sigma\sigma'}}{\rho_{d\sigma}} \left[\omega^2 + \frac{3}{4}(eV)^2 + (\pi T)^2 \right] + \dots, \quad (\sigma' \neq \sigma), \quad (11)$$

$$\epsilon_d + \text{Re} \Sigma_\sigma^r(\omega) = \Delta \cot \delta + (1 - \tilde{\chi}_{\sigma\sigma})\omega + \frac{1}{2} \frac{\partial \tilde{\chi}_{\sigma\sigma}}{\partial \epsilon_{d\sigma}} \omega^2 + \frac{N-1}{6} \frac{\chi_{\sigma\sigma'}^{[3]}}{\rho_{d\sigma}} \left[\frac{3}{4}(eV)^2 + (\pi T)^2 \right] + \dots, \quad (12)$$

for $\Gamma_L = \Gamma_R = \Delta/2$ and $\mu_L = -\mu_R = eV/2$ in the $\text{SU}(N)$ symmetric case. Note that $\tilde{\chi}_{\sigma\sigma''} \equiv \delta_{\sigma\sigma''} + \frac{\partial \Sigma_\sigma^r(0)}{\partial \epsilon_{d\sigma''}} \Big|_{T=eV=0}$, $\chi_{\sigma\sigma''} = \rho_{d\sigma} \tilde{\chi}_{\sigma\sigma''}$, and $\rho_{d\sigma} = \sin^2 \delta / (\pi \Delta)$ for $\sigma = 1, 2, 3, \dots, N$. These results are obtained by extending further the latest version of Fermi-liquid description [A. Oguri and A. C. Hewson, Phys. Rev. B **97**, 035435] to the multilevel cases $N > 2$. At $T = eV = 0$, the self-energy satisfies the Ward identity of the following form, which yields the Fermi-liquid relations between the expansion coefficients,

$$\left(\delta_{\sigma\sigma'} \frac{\partial}{\partial \omega} + \frac{\partial}{\partial \epsilon_{d\sigma'}} \right) \Sigma_{\sigma\sigma'}^{--}(\omega) = -\Gamma_{\sigma\sigma';\sigma'\sigma}^{--;--}(\omega, 0; 0, \omega) \rho_{d\sigma'}. \quad (13)$$

Here, $\Sigma_{\sigma\sigma'}^{--}(\omega) = \text{Re} \Sigma_\sigma^r(\omega) + i \text{sgn}(\omega) \text{Im} \Sigma_\sigma^r(\omega)$, and $\Gamma_{\sigma\sigma';\sigma'\sigma}^{--;--}(\omega, \omega'; \omega', \omega)$ is the causal vertex function at $T = eV = 0$. It has been determined up to linear order terms with respect to ω and ω' ,

$$\Gamma_{\sigma\sigma';\sigma\sigma}^{--;--}(\omega, \omega'; \omega', \omega) \rho_{d\sigma}^2 = i\pi \sum_{\sigma'(\neq \sigma)} \chi_{\sigma\sigma'}^2 |\omega - \omega'| + \dots, \quad (\sigma' \neq \sigma), \quad (14)$$

$$\Gamma_{\sigma\sigma';\sigma'\sigma}^{--;--}(\omega, \omega'; \omega', \omega) \rho_{d\sigma} \rho_{d\sigma'} = -\chi_{\sigma\sigma'} + \rho_{d\sigma} \frac{\partial \tilde{\chi}_{\sigma\sigma'}}{\partial \epsilon_{d\sigma}} \omega + \rho_{d\sigma'} \frac{\partial \tilde{\chi}_{\sigma'\sigma}}{\partial \epsilon_{d\sigma'}} \omega' + i\pi \chi_{\sigma\sigma'}^2 (|\omega - \omega'| - |\omega + \omega'|) + \dots. \quad (15)$$

The Ward identity itself follows from the current conservation between the dot and leads:

$$\frac{\partial}{\partial t}(e n_{d\sigma}) + \hat{J}_{R,\sigma} - \hat{J}_{L,\sigma} = 0, \quad \hat{J}_{L,\sigma} \equiv i e v_L (\psi_{L\sigma}^\dagger d_\sigma - d_\sigma^\dagger \psi_{L\sigma}), \quad \hat{J}_{R,\sigma} \equiv -i e v_R (\psi_{R\sigma}^\dagger d_\sigma - d_\sigma^\dagger \psi_{R\sigma}). \quad (16)$$

II. FERMI-LIQUID CORRECTIONS FOR NONLINEAR CURRENT NOISE $S_{\text{noise}}^{\text{QD}}$

In contrast to the average current J and thermal conductivity κ_{QD} , the current noise depends also on the two-quasiparticle collisions which correspond to the vertex corrections for the current-current correlation function $\mathcal{K}_{\sigma'\sigma}^{\nu'\nu}$:

$$S_{\text{noise}}^{\text{QD}} = \int_{-\infty}^{\infty} dt \sum_{\sigma\sigma'} i \left[\mathcal{K}_{\sigma'\sigma}^{+-}(t, 0) + \mathcal{K}_{\sigma'\sigma}^{-+}(t, 0) \right], \quad \delta \hat{J}_\sigma(t) \equiv \hat{J}_\sigma(t) - \langle \hat{J}_\sigma(t) \rangle_V, \quad (17)$$

$$\mathcal{K}_{\sigma'\sigma}^{+-}(t, 0) \equiv -i \langle \delta \hat{J}_{\sigma'}(t) \delta \hat{J}_\sigma(0) \rangle_V, \quad \mathcal{K}_{\sigma'\sigma}^{-+}(t, 0) \equiv -i \langle \delta \hat{J}_\sigma(0) \delta \hat{J}_{\sigma'}(t) \rangle_V. \quad (18)$$

Here, $\hat{J}_\sigma \equiv (\Gamma_L \hat{J}_{R,\sigma} + \Gamma_R \hat{J}_{L,\sigma}) / (\Gamma_L + \Gamma_R)$ is a symmetrized current operator. In this work, we have expanded $S_{\text{noise}}^{\text{QD}}$ up to terms of order $(eV)^3$, using the diagrammatic representation illustrated in Fig. 3. To this end, all the components of Keldysh Green's function $G_{\sigma}^{\nu'\nu}(\omega)$ have been deduced up to order ω^2 and $(eV)^2$, and the Keldysh vertex function $\Gamma_{\sigma\sigma';\sigma'\sigma}^{\nu_1\nu_2;\nu_3\nu_4}(\omega, \omega'; \omega'\omega)$ have been calculated up to linear order in ω, ω' and eV . We have checked that the result satisfies the nonequilibrium Ward identity [A. Oguri, Y. Tetatani, and S. Sakano, unpublished],

$$\left(\delta_{\sigma\sigma'} \frac{\partial}{\partial \omega} + \frac{\partial}{\partial \epsilon_{d\sigma'}} \right) \Sigma_{\sigma}^{\nu_4\nu_1}(\omega) = - \int_{-\infty}^{\infty} \frac{d\omega'}{2\pi} \sum_{\nu_2\nu_3} \Gamma_{\sigma\sigma';\sigma'\sigma}^{\nu_1\nu_2;\nu_3\nu_4}(\omega, \omega'; \omega'\omega) 2\Delta G_{\sigma'}^r(\omega') G_{\sigma'}^a(\omega') \left(-\frac{\partial f_{\text{eff}}(\omega')}{\partial \omega'} \right). \quad (19)$$

Here, $f_{\text{eff}}(\omega) = \{\Gamma_L f_L(\omega) + \Gamma_R f_R(\omega)\} / (\Gamma_L + \Gamma_R)$, and $\Sigma_{\sigma}^{\nu_4\nu_1}(\omega)$ is the Keldysh self-energy. The expansion coefficient for order $(eV)^3$ current noise is given in table I. It can be separated into two parts $C_S = C_S^{\text{qp}} + C_S^{\text{coll}}$, as mentioned in the main text. Here, C_S^{qp} and C_S^{coll} represent respectively the contributions of the bubble diagram and that of the vertex corrections shown in Fig. 3.

The nonlinear Fano factor, $F_K = \frac{C_S}{C_V/3}$, can be expressed in the following form for the $\text{SU}(N)$ Anderson impurity at arbitrary electron fillings,

$$F_K = \frac{\cos 4\delta + \left[4 + 5 \cos 4\delta + \frac{3}{2}(1 - \cos 4\delta)(N - 2) \right] \frac{\tilde{K}^2}{N - 1} - \cos 2\delta \left[\Theta_{\text{I}} + 3(N - 1)\Theta_{\text{II}} \right]}{- \left[1 + 5 \frac{\tilde{K}^2}{N - 1} \right] \cos 2\delta + \Theta_{\text{I}} + 3(N - 1)\Theta_{\text{II}}}. \quad (20)$$

For $N = 2$, it reproduces the previous result, obtained by Mora *et al* [Eq. (11) of Phys. Rev. B **92**, 075120 (2015)]: their notation and our one correspond to each other such that $\alpha_{\sigma}^{(1)}/\pi = \chi_{\sigma\sigma}$, $\phi_{\sigma\sigma'}^{(1)}/\pi = -\chi_{\sigma\sigma'}$, $\alpha_{\sigma}^{(2)}/\pi = -\frac{1}{2}\chi_{\sigma\sigma\sigma}^{[3]}$, and $\phi_{\sigma\sigma'}^{(2)}/\pi = 2\chi_{\sigma\sigma'\sigma'}^{[3]}$, for $\sigma' \neq \sigma$. In Fig. 4, we have also plotted F_K and other Fermi-liquid parameters for the $\text{SU}(2)$ symmetric case as functions of ϵ_d for comparisons with those for $N = 4$ and 6 shown in the main text.



FIG. 3. The Feynman diagrams for the correlation function $\int_{-\infty}^{\infty} dt \mathcal{K}_{\sigma'\sigma}^{\nu'\nu}(t, 0)$. The solid lines denote the Keldysh Green's functions $G_{\sigma}^{\nu'\nu}(\omega)$. The shaded region in the diagram on the right represents the Keldysh vertex function $\Gamma_{\sigma\sigma';\sigma'\sigma}^{\nu_1\nu_2;\nu_3\nu_4}(\omega, \omega'; \omega'\omega)$. The superscripts ν, ν' and ν_i ($i = 1, 2, 3, 4$) specify the branches of Keldysh time-loop contour. We are using the notation in which $\nu = -$ and $+$ represent the forward and return paths, respectively.




FIG. 4. Fermi-liquid parameters for the SU(2) symmetric case are plotted vs $\xi_d \equiv \epsilon_d + U/2$ for $U/(\pi\Delta) = 5.0$. (a): $\sin^2 \delta$, renormalization factor z , and $\tilde{K} \equiv (N-1)(R-1)$. (b): three-body correlators Θ_I , and $-\tilde{\Theta}_{II} \equiv -(N-1)\Theta_{II}$. (c): $\frac{48}{\pi^2}C_T$, $\frac{64}{\pi^2}C_V$, $\frac{192}{\pi^2}C_S$, and $\frac{80}{7\pi^2}C_K^{\text{QD}}$. (d): nonlinear Fano factor $F_K = \frac{C_S}{C_V/3}$ is plotted also for $U/(\pi\Delta) = 1/3, 5$.

Equation (20) also reproduces the previous result in the particle-hole symmetric case, at which $\delta = \pi/2$ and the three-body contributions vanish $\Theta_I = \Theta_{II} = 0$ [Sakano *et al* Phys. Rev. B **83**, 075440 (2011)] :

$$F_K \xrightarrow{\epsilon_d \rightarrow -(N-1)U/2} \frac{1 + \frac{9\tilde{K}^2}{N-1}}{1 + \frac{5\tilde{K}^2}{N-1}} \xrightarrow{U \rightarrow \infty} \frac{1 + \frac{9}{N-1}}{1 + \frac{5}{N-1}}. \quad (21)$$

In the strong coupling limit $U \rightarrow \infty$, the occupation number $N_d \equiv \sum_{\sigma} \langle n_{d\sigma} \rangle$ becomes integer $M = 1, 2, \dots, N-1$ at $\epsilon_d = -(M-1/2)U$ and the phase shift is locked at $\delta = \pi M/N$. The charge and spin susceptibilities satisfy the stationary conditions in this case, and thus Eq. (20) can be rewritten as

$$F_K \rightarrow \frac{1 + \sin^2\left(\frac{2\pi M}{N}\right) + \frac{9-13\sin^2\left(\frac{2\pi M}{N}\right)}{N-1} + 2\Theta_I \cos\left(\frac{2\pi M}{N}\right)}{-\left[1 + \frac{5}{N-1}\right] \cos\left(\frac{2\pi M}{N}\right) - 2\Theta_I}. \quad (22)$$

This expression is consistent with the corresponding noise formula for the SU(N) Kondo model, obtained by Mora *et al* [Eq. (51) of Phys. Rev. B **80**, 155322 (2009), after inserting some parenthesis for correcting minor typos].

III. NRG CALCULATIONS

NRG calculations for the SU(N) Anderson model for $N = 2, 4, 6$ have been carried out, dividing N channels into $N/2$ pairs and exploiting the SU(2) spin and U(1) charge symmetries for each of the pairs, i.e. using $\prod_{k=1}^{N/2} \{\text{SU}(2) \otimes \text{U}(1)\}_k$ symmetries. The discretization parameter Λ and the number of retained low-lying excited states N_{trunc} are chosen such that $(\Lambda, N_{\text{trunc}}) = (2, 4000)$ for $N = 2$, $(6, 10000)$ for $N = 4$, and $(20, 30000)$ for $N = 6$. We have also exploited methods of Stadler's *et al* [Phys. Rev. B **93**, 235101 (2017)] for $N = 6$. The truncation is performed at each step after adding states from each pair of the channels, using Olivera's \mathcal{Z} -trick [Phys. Rev. B **49**, 11986 (1994)] and choosing different \mathcal{Z} values for different pairs: $\mathcal{Z}_i = 1/2 + i/N$ for the i -th pair ($i = 1, 2, \dots, N/2$).

TABLE II. The coefficients C 's at finite magnetic fields b for $N = 2$ at half filling $\epsilon_d = -\frac{U}{2}$.
See also, A. Oguri and A. C. Hewson, Phys. Rev. B **98**, 079905 (E).

$C_S^b = \frac{\pi^2}{192} [W_S^b + (\Theta_I^b + 3\Theta_{II}^b) \cos(\pi m_d)]$	$W_S^b \equiv \cos(2\pi m_d) + [4 + 5 \cos(2\pi m_d)](R-1)^2$
$C_V^b = \frac{\pi^2}{64} [W_V^b + \Theta_I^b + 3\Theta_{II}^b]$	$W_V^b \equiv [1 + 5(R-1)^2] \cos(\pi m_d)$
$C_T^b = \frac{\pi^2}{48} [W_T^b + \Theta_I^b + \Theta_{II}^b]$	$W_T^b \equiv [1 + 2(R-1)^2] \cos(\pi m_d)$
$C_{\kappa,b}^{\text{QD}} = \frac{7\pi^2}{80} [W_{\kappa,b}^{\text{QD}} + \Theta_I^b + \frac{5}{2}\Theta_{II}^b]$	$W_{\kappa,b}^{\text{QD}} \equiv [1 + \frac{6}{7}(R-1)^2] \cos(\pi m_d)$

IV. MAGNETIC-FIELD DEPENDENCE OF CURRENT NOISE FOR $N = 2$

We describe here supplemental information about the nonlinear current noise at finite magnetic field b , specifically for $N = 2$ at half filling, where the impurity level is given by $\epsilon_{d\sigma} \equiv -\frac{U}{2} - \text{sgn}(\sigma)b$ with $\text{sgn}(\uparrow) = +1$ and $\text{sgn}(\downarrow) = -1$. In this case, the phase shift takes the form $\delta_\sigma = \pi\{1 + \text{sgn}(\sigma)m_d\}/2$ with $m_d \equiv \langle n_{d\uparrow} \rangle - \langle n_{d\downarrow} \rangle$, and the other correlation functions have symmetry properties: $\chi_{\uparrow\uparrow} = \chi_{\downarrow\downarrow}$, $\chi_{\uparrow\downarrow} = \chi_{\downarrow\uparrow}$, $\chi_{\uparrow\downarrow}^{[3]} = -\chi_{\downarrow\uparrow}^{[3]}$, and $\chi_{\downarrow\downarrow}^{[3]} = -\chi_{\uparrow\uparrow}^{[3]}$. Thus, the transport coefficients up to the next leading order can be described by five parameters, for instance, m_d , $T^* = 1/(4\chi_{\uparrow\uparrow})$, $R = 1 - \chi_{\uparrow\downarrow}/\chi_{\uparrow\uparrow}$ and the following 3-body correlation functions,

$$\Theta_I^b \equiv -\frac{\sin(\pi m_d)}{2\pi} \frac{\chi_{\uparrow\uparrow\uparrow}^{[3]}}{\chi_{\uparrow\uparrow}^2}, \quad \Theta_{II}^b \equiv -\frac{\sin(\pi m_d)}{2\pi} \frac{\chi_{\uparrow\downarrow\downarrow}^{[3]}}{\chi_{\uparrow\uparrow}^2}. \quad (23)$$

The low-energy expansion of the current noise $S_{\text{noise}}^{\text{QD}}$, conductance dJ/dV , and thermal conductivity κ_{QD} for this case can be written in the following form, with the coefficients C 's listed in table II,

$$\begin{aligned} S_{\text{noise}}^{\text{QD}} &= 2 \frac{2e^2}{h} |eV| \left[\frac{\sin^2(\pi m_d)}{4} + C_S^b \left(\frac{eV}{T^*} \right)^2 + \dots \right], \quad (24) \\ \frac{dJ}{dV} &= \frac{2e^2}{h} \left[\cos^2 \left(\frac{\pi m_d}{2} \right) - C_T^b \left(\frac{\pi T}{T^*} \right)^2 - C_V^b \left(\frac{eV}{T^*} \right)^2 + \dots \right], \\ \kappa_{\text{QD}} &= \frac{2\pi^2 T}{3h} \left[\cos^2 \left(\frac{\pi m_d}{2} \right) - C_{\kappa,b}^{\text{QD}} \left(\frac{\pi T}{T^*} \right)^2 + \dots \right]. \end{aligned}$$




FIG. 5. $F_K^b = \frac{\overline{C}_S^b}{\overline{C}_V^{b/3}}$ and $\overline{C}_{\kappa,b}^{\text{QD}} \equiv (T_K/T^*)^2 C_{\kappa,b}^{\text{QD}}$ for $N = 2$ are plotted vs b/T_K at half filling for $U/(\pi\Delta) = 2.5, 3.0, 3.5, 4.0$: T_K varies with U , and $T_K \xrightarrow{U \rightarrow 0} \pi\Delta/4$.

In order to see the magnetic field dependences in the Kondo regime, it is preferable to rescale the next leading $(eV)^2$ and T^2 contributions by the Kondo temperature defined at zero field $T_K = \lim_{b \rightarrow 0} T^*$. This is because all effects of b are absorbed into the coefficients redefined such that $\overline{C}_V^b \equiv (T_K/T^*)^2 C_V^b$, $\overline{C}_S^b \equiv (T_K/T^*)^2 C_S^b$, and $\overline{C}_{\kappa,b}^{\text{QD}} \equiv (T_K/T^*)^2 C_{\kappa,b}^{\text{QD}}$. We have presented the NRG results for the nonlinear current noise \overline{C}_S^b in the main text; \overline{C}_V^b was examined previously [A. Oguri and A. C. Hewson, Phys. Rev. Lett. **120**, 126802 (2018)]. In Fig. 5, $F_K^b \equiv \frac{\overline{C}_S^b}{\overline{C}_V^{b/3}}$ and $\overline{C}_{\kappa,b}^{\text{QD}}$ are plotted as functions of b/T_K for several different values of U . The nonlinear Fano factor F_K^b shows the Kondo scaling behavior for strong interactions $U/(\pi\Delta) \gtrsim 2.0$. The universal curve of F_K^b deviates significantly from the curve for $U = 0$ keeping its qualitative characteristics unchanged. We also find that the thermal conductivity $\overline{C}_{\kappa,b}^{\text{QD}}$ exhibits the universal scaling behavior.