ISOTRIVIALITY OF SMOOTH FAMILIES OF VARIETIES OF GENERAL TYPE

CHUANHAO WEI AND LEI WU

ABSTRACT. In this paper, we proved that a log smooth family of log general type klt pairs with a special (in the sense of Campana) quasi-projective base is isotrivial. As a consequence, we proved the generalized Kebekus-Kovács conjecture [WW18, Conjecture 1.1], for smooth families of general type varieties as well as log smooth families of log canonical pairs of log general type, assuming the existence of relative good minimal models.

1. INTRODUCTION

We start with $f_V : (U, D^U) \to V$, a log smooth family of projective log pairs of log general type, over a smooth quasi-projective variety V, with the coefficients of D^U are in [0, 1). Here, log smoothness of f_V means that f_V and each stratum of D^U is smooth over V (see §3.1). Due to [BCHM10], f_V has a relative canonical model, denoted by f_V^c . The relative canonical model is a stable family by the invariance of log plurigenera [HMX18, Theorem 4.2], and hence it induces a moduli map $\mu : V \to \mathfrak{M}$, where \mathfrak{M} is the coarse moduli space of the corresponding stable family (with a fixed set of coefficients) as in [KP16, §6]. Note that \mathfrak{M} is a projective variety, [KP16, Theorem 1.1].

Following [KP16, Definition 6.16], we denote the variation of f_V by

$$\operatorname{Var}(f_V) = \operatorname{Var}(f_V^c) := \dim(\mu(V)).$$

We say that f_V is of maximal variation if $\operatorname{Var}(f_V) = \dim V$ and that f_V is isotrivial if $\operatorname{Var}(f_V) = 0$. Since fibers of f_V are of log general type, the above definition is compatible with the variation defined in [Vie83] and [Kaw85]. We write the log Kodaira dimension of V by $\bar{\kappa}(V) = \kappa(Y, E^Y)$, where the latter is defined to be the Iitaka dimension of the log canonical sheaf $\omega_Y(E^Y)$ and Y is a projective compactification of V with $E^Y = Y \setminus U$ a normal crossing divisor. We call (Y, E^Y) a log smooth compactification of V. Note that $\bar{\kappa}(V)$ does not depend on the choice of (Y, E^Y) and it is a birational invariant, that is, $\bar{\kappa}(V)$ remains the same, if one replaces V by a smooth birational model. In this paper, we prove

Theorem 1.0.1. With notations above, we have:

- (1) if $\bar{\kappa}(V) > -\infty$, then $\bar{\kappa}(V) \ge \operatorname{Var}(f_V)$,
- (2) if $\bar{\kappa}(V) = -\infty$, then dim $V > \operatorname{Var}(f_V)$.

When D^U is empty, we particularly have:

Theorem 1.0.2. Let $f_V : U \to V$ be a smooth family of projective varieties of general type. Then,

(1) if $\bar{\kappa}(V) > -\infty$, then $\bar{\kappa}(V) \ge \operatorname{Var}(f_V)$,

(2) if $\bar{\kappa}(V) = -\infty$, then dim $V > \operatorname{Var}(f_V)$.

When f_V is a smooth family of canonical polarized varieties, the above statement is conjectured by Kebekus and Kovács [KK08, Conjecture 1.6], which is a natural extension of Viehweg's hyperbolicity conjecture [Vie01]. Meanwhile, Campana made a related conjecture for smooth families of canonical polarized varieties [Cam07, Conjecture 12.19] (see also [JK11a, Conjecture 1.4]), the Isotriviality Conjecture, which implies the Kebekus-Kovács conjecture. The Kebekus-Kovács conjecture for smooth families of canonical polarized varieties is proved by Taji [Taj16] by proving the Isotriviality Conjecture of Campana.

In the special case that $\operatorname{Var}(f_V) = \dim V$, Campana's isotriviality conjecture is also known as Viehweg's hyperbolicity conjecture. For smooth families of canonically polarized varieties, it was proved by Campana and Păun [CP15]. Their result has been extended to the case for smooth families of general type varieties by Popa and Schnell using Hodge modules [PS17, Theorem A]. They [PS17, §4.3] implicitly gave an extension of the Kebekus-Kovács conjecture for families with geometric generic fibers admitting good minimal models (particularly for smooth families of general type varieties). They proved in *loc. cit.* a special case, with additionally assuming an abundance-type conjecture of Campana–Peternell. Based on the Hodge-module construction of Popa and Schnell, we further extended Viehweg's hyperbolicity to smooth families of log general type pairs (see [WW18, Theorem A(1)]). We then extended the Kebekus-Kovács conjecture to the case for smooth families of log general type pairs (see [WW18, Conjecture 1.1]).

Theorem 1.0.1 gives a confirmative answer to the generalized Kebekus-Kovács conjecture. It is worth mentioning that the generalized Kebekus-Kovács conjecture even for smooth families of general type varieties was not known before.

There are three reasons why we consider Theorem 1.0.1 (the pair version) instead of Theorem 1.0.2 (the non-pair version):

(1) The statement of Theorem 1.0.2 is about the log Kodaira dimension of the base, where the log Kodaira dimension is defined by embedding the base into a pair. (2) To prove Theorem 1.0.2, we first need to compactify f_V

$$U \longleftrightarrow (X, D^X)$$
$$\downarrow^{f_V} \qquad \qquad \downarrow^f$$
$$V \longleftrightarrow (Y, D^Y),$$

and then do stable reductions (see §4), where f is a morphism of log smooth pairs. (3) The proof of Theorem 1.0.2 needs to use Campana's fibrations, but they are naturally defined as morphisms of pairs (see §2.1).

Remark 1.0.3. Actually, we can consider the case that f_V is a log smooth family of log smooth pairs of log general type with the coefficients of D^U in [0, 1], if we assume that f_V admits a relative good minimal model over a Zariski open subset $V_0 \subseteq V$. Then, we have the smooth family $f_V^{\epsilon} : (U, D_{\epsilon}^U := (1 - \epsilon)D^U) \to V$ of log smooth pairs of log general type with the coefficients of D_{ϵ}^U in [0, 1) and $\operatorname{Var}(f_V^{\epsilon}) \geq \operatorname{Var}(f_V)$ by [WW18, Lemma 3.1]. See [WW18, §3] for the definition of variation in the log canonical case. The inequality $\operatorname{Var}(f_V^{\epsilon}) \geq \operatorname{Var}(f_V)$ implies that Theorem 1.0.1 still holds for smooth families of log canonical pairs of log general type, assuming the existence of relative good minimal models. In order to prove Theorem 1.0.1, we prove Campana's Isotriviality Conjecture for log smooth families of log general type pairs, Theorem 1.0.4 below. We continue to use the notations in Theorem 1.0.1. We now fix a log smooth compactification (Y, E^Y) of V, such that the moduli map $\mu : V \to \mathfrak{M}$ extends to $\bar{\mu} : Y \to \mathfrak{M}$. Considering the Stein factorization of $\bar{\mu}$, we get a morphism $s : Y \to M$ with M a normal variety so that $\bar{\mu}$ factors through s and s has generically connected fibers. In particular, $s : (Y, E^Y) \to M$ is a fibration of (Y, E^Y) over M with dim $M = \operatorname{Var}(f_V)$ (see Definition 2.1.1). Campana defined the fiberations of general type and speciality in the category of geometric orbifolds; see Definition 2.1.3 for a simplified version in our setting.

Theorem 1.0.4. With notations as above, the fibration s is of general type. As a consequence, if V is a special quasi-projective variety, then f_V is birationally isotrivial, i.e. $Var(f_V) = 0$.

The first statement of Theorem 1.0.4 implies Theorem 1.0.1, thanks to the additivity of the log Kodaira dimension for fibrations of general type, [Cam11, Theorem 6.3].

When f_V is a smooth family of canonically polarized variety, the second statement of Theorem 1.0.4 is the Isotriviality Conjecture of Campana as we mentioned above. In the case that dim $V \leq 3$, it is proved by Jabbusch and Kebekus [JK11a]. Their proof uses Campana's theory of geometric orbifolds and the so-called Viehweg-Zuo sheaves constructed in [VZ02] as well as a refine result on Viehweg-Zuo sheaves in [JK11b], which roughly asserts that Viehweg-Zuo sheaves factor through the moduli. Based on the strategy of Jabbusch and Kebekus and the method of Campana and Păun in proving Viehweg's hyperbolicity conjecture, Taji [Taj16] proved the isotriviality conjecture for smooth families of canonically polarized varieties in general.

By using Hodge modules, Popa and Schnell [PS17] constructed the Viehweg-Zuo sheaves for smooth families of general type varieties with maximal variation (or more precisely for smooth families admitting relative good minimal models). We further constructed Viehweg-Zuo sheaves for families of pairs with maximal variation in [WW18]. A key step to prove Theorem 1.0.4 is to construct the Viehweg-Zuo sheaves for families of (log) general type varieties (pairs) with arbitrary variation and prove that they factor through the "moduli" in the way analoguous to the result of Jabbusch and Kebekus.

Definition 1.0.5. In regard to a log smooth compactification (Y, E) of V and an extension $\bar{\mu}: Y \to \mathfrak{M}$ of the moduli map μ , we define the subsheaf

$$\mathcal{B}_{(Y,E^Y)} \subseteq \Omega_Y(\log E^Y)$$

by the saturation of the image of the natural morphism $\bar{\mu}^* \Omega_{\mathfrak{M}} \to \Omega_Y(\log E^Y)$ in $\Omega_Y(\log E^Y)$, where $\Omega_{\mathfrak{M}}$ is the sheaf of Kähler differentials (over \mathbb{C}) and $\Omega_Y(\log E^Y)$ the sheaf of log 1-forms with log poles along E^Y .

The above definition follows [JK11b, Notation 1.2]. However, they only considered the case when \mathfrak{M} is the moduli space of canonically polarised manifolds in *loc. cit.* Note that, since $\bar{\mu}$ factors through $s: Y \to M$, and $M \to \mathfrak{M}$ is quasi-finite, we have that $\mathcal{B}_{(Y,E^Y)}$ is also the saturation of the image of $s^*\Omega_M \to \Omega_Y(\log E^Y)$, and the rank of $\mathcal{B}_{(Y,E^Y)}$ is equal to the dimension of M, which is also the variation of f_V . Now, we state the main technical result of this paper. **Theorem 1.0.6.** Let $f_V : (U, D^U) \to V$ be a log smooth family of projective log pairs of log general type, over a smooth quasi-projective variety V, with the coefficients of D^U in [0,1). If $\operatorname{Var}(f_V) > 0$, then after replacing V by a further log resolution, we have a log smooth compactification (Y, E^Y) of V, such that there exists an invertible subsheaf $\mathcal{A} \subset \operatorname{Sym}^{[n]} \mathcal{B}_{(Y,E^Y)}$, for some positive integer n, with $\kappa(\mathcal{A}) \geq \operatorname{Var}(f_V)$.

We call the invertible sheaf \mathcal{A} the refined Viehweg-Zuo Sheaf of f_V on (Y, E^Y) . The proof of Theorem 1.0.6 is built upon the refinement of the stable reduction used in [WW18] (see Section 5 for details). Another key input of its proof is the use of Hodge modules. Roughly speaking, we essentially need Saito's decomposition theorem for pure Hodge modules to compare Viehweg-Zuo sheaves before and after base-changes, see Theorem 3.5.1 in Section 3 for details.

Structure of the paper. In Section 2, we recall some of Campana's definitions and results about orbifold fibrations, and concludes the proof of Theorem 1.0.4 using Theorem 1.0.6. In Section 3, we fixed the notations and show some useful results using Saito's theory of Hodge modules. In Section 4, we prove results related to stable reductions, and using them to make the geometric constructions that are needed in Section 5 to construct the refined Viehweg-Zuo Sheaf in our setting.

Acknowledgement. We would like to thank Christian Schnell for useful discussions during the preparation of the paper. The first author also gets some inspiration from a workshop held in Shanghai Center for Mathematical Sciences.

2. BIRATIONALLY EQUIVALENT FIBRATIONS

2.1. Birationally equivalent fibration in the sense of Campana. We now recall Campana's birational equivalence of fibrations. We mainly restrict ourselves to the setting of Theorem 1.0.4 for our application while Campana works more generally in the category of geometric orbifolds in [Cam11]. See also [JK11a] for a more approachable introduction.

Definition 2.1.1. We say that $s: (Y, E^Y) \to M$ is a fibration of a log pair (Y, E^Y) over M, if s is a dominate projective morphism with generically connected fibers, and W a normal variety. For simplicity we always assume that Y is a smooth quasiprojective variety. Given two fibrations $s: (Y, E^Y) \to M$ and $s': (Y', E^{Y'}) \to M'$, we say that s' is dominant over s, if we have the following commutative diagram

$$(Y, E^Y) \xleftarrow{u} (Y', E^{Y'})$$
$$\downarrow^s \qquad \qquad \downarrow^{s'}$$
$$M \xleftarrow{v} M',$$

with both u and v are birational, and $u_*E^{Y'} = E^Y$. We say two fibrations s and s' are birationally equivalent if they both can be dominated by a third fibration $s'': (Y'', E^{Y''}) \to M''$.

Using the recipe in [Cam11, Definition 3.2] (see also [JK11a, Construction and Definition 5.3]), we obtain the C-base (M, Δ^s) associated to the fibration s. We say that a fibration $s : (Y, E^Y) \to (M, \Delta^s)$ is a *neat* model if (Y, E^Y) and (M, Δ^s) are log smooth, and for all of the divisors F with codim $s(F) \ge 2$, $F \subset E^Y$, [JK11a,

Assumption 5.4]. The name of neat model is adopted from [Taj16, Definition 4.1], which serves a similar purpose as "strictement nette et haute" model in [Cam11], in the general case. By definition, starting from an arbitrary fibration $s : (Y, E^Y) \to M$, the associated fibration $s : (Y, E^Y) \to (M, \Delta^s)$ over the associated \mathcal{C} -base (M, Δ^s) is not always neat even when the pairs (Y, E^Y) and (M, Δ^s) are log smooth. We recall the following result in [Taj16], which is essentially proved in [JK11a, Section 10].

Proposition 2.1.2. [Taj16, Proposition 4.2] Every fibration $s : (Y, E^Y) \to (M, \Delta^s)$ is dominated by a neat model.

Definition 2.1.3. ([Cam11, Definition 4.10, 4.16 and 4.17]) Let $s : (Y, E^Y) \to M$ be a fibration with (Y, E^Y) log smooth and M projective. Let $\tilde{s} : (\tilde{Y}, E^{\tilde{Y}}) \to (\tilde{M}, \Delta)$ be neat model dominant over s, with (\tilde{M}, Δ) the induced C-base of \tilde{s} . We define the canonical dimension of s by $\kappa(\tilde{M}, \Delta^{\tilde{s}})$. It does not depend on the choice of the neat model \tilde{s} dominant over s (see [Cam11, Corollaire 4.11]). Then we define:

- (1) a fibration s is of general type if its canonical dimension is the same as the dimension of the base,
- (2) (Y, E^Y) is special if there exists no fibration $s : (Y, E^Y) \to M$ of general type with dim(M) > 0.

We also say the smooth quasi-projective variety $V = Y \setminus E^Y$ is special if (Y, E^Y) is so. One can easily check that the specialty of V does not depend on the choice of the compactification (Y, E^Y) and that specialty is a birational invariant, that is, V is special if and only if V' is so, where V' is a smooth quasi-projective variety properly birational to V.

2.2. Using the refined Viehweg-Zuo sheaf to prove the main theorem. We use Theorem 1.0.6 to give a proof of Theorem 1.0.4. The proof follows the strategy in [JK11a], which is also used by Taji in proving the Isotriviality conjecture of Campana [Taj16, Theorem 1.5].

Proof of Theorem 1.0.4. When f_V is birationally isotrivial, the first statement of Theorem 1.0.4 is obvious. We hence assume $\operatorname{Var}(f_V) > 0$. Using Proposition 2.1.2, we fix a neat model dominating $s, \tilde{s} : (\tilde{Y}, E^{\tilde{Y}}) \to (\tilde{M}, \Delta^{\tilde{s}})$, . Let $\mu : (\tilde{Y}, E^{\tilde{Y}}) \to (Y, E^Y)$ be the induced birational morphism. Without loss of generality, we further assume that $E^{\tilde{Y}} \supset \mu^{-1}E^Y$ by adding more components to $E^{\tilde{Y}}$ (by doing this the \mathcal{C} -base (\tilde{M}, Δ) stays the same).

Recall that $\mathcal{B}_{(Y,E^Y)}$ is defined to be the saturation of the image of the natural morphism $\tilde{s}^* : \Omega_M \to \Omega_Y(\log E^Y)$. We similarly define $\mathcal{B}_{(\tilde{Y},E^{\tilde{Y}})}$ to be the saturation of the image of the natural morphism $\tilde{s}^* : \Omega_{\tilde{M}} \to \Omega_{\tilde{Y}}(\log E^{\tilde{Y}})$. By [JK11a, Proposition 3.3], we have that $\operatorname{Sym}^{[n]}\mathcal{B}_{(\tilde{Y},E^{\tilde{Y}})}$ is also the saturation of the image of the composed natural morphisms

 $\mu^* \operatorname{Sym}^{[n]} \mathcal{B}_{(Y, E^Y)} \to \mu^* \operatorname{Sym}^{[n]} \Omega_Y(\log E^Y) \to \operatorname{Sym}^{[n]} \Omega_{\tilde{Y}}(\log E^{\tilde{Y}}).$

Hence, the refined Viehweg-Zuo sheaf $\mathcal{A} \subset \operatorname{Sym}^{[n]}\mathcal{B}_{(Y,E^Y)}$ on (Y, E^Y) lifted to $\mu^*\mathcal{A} \subset \operatorname{Sym}^{[n]}\mathcal{B}_{(\tilde{Y},E^{\tilde{Y}})}$. Hence, to make notations simple, we can assume that $s: (Y, E^Y) \to (M, \Delta^s)$ itself is neat.

By definition, to prove the fibration $s : (Y, E^Y) \to M$ is of general type, one only needs to show that (M, Δ^s) is of log general type. Since $\mathcal{B}_{(Y, E^Y)}$ is saturated, $\operatorname{Sym}^{[n]}\mathcal{B}_{(Y, E^Y)} \subseteq \operatorname{Sym}^{[n]}\Omega_Y(\log E^Y)$ is also a saturated subsheaf by [JK11a, Proposition 3.3]. Due to [JK11a, Proposition 5.7], see [JK11a, Notation 4.1] for the notations, we have a natural isomorphism

$$\iota: \operatorname{Sym}_{\mathcal{C}}^{[n]}\Omega_M(\log \Delta^s) \to s_* \operatorname{Sym}^{[n]}\mathcal{B}_{(Y,E^Y)}$$

for all *n*. Using Theorem 1.0.6, we have the refined Viehweg-Zuo sheaf $\mathcal{A} \subset \operatorname{Sym}^{[n]}\mathcal{B}_{(Y,E^Y)}$, and we can further assume that \mathcal{A} is saturated. Let \mathcal{A}_M be the saturation of $s_*\mathcal{A}$ in $\operatorname{Sym}_{\mathcal{C}}^{[n]}\Omega_M(\log \Delta^s)$, and in particular, it is a line bundle with $\kappa_{\mathcal{C}}(\mathcal{A}_M) = \kappa(\mathcal{A}) = \dim M$, by [JK11a, Proposition 5.7, Corollary 5.8]. By applying [Taj16, Theorem 5.2], we get that (M, Δ^s) is of log general type. \Box

3. Construction of Higgs sheaves

In this section, we use Seito's theory of Hodge modules to show some results about Hodge bundles that will be used to construct the Viehweg-Zuo Sheaves. Some constructions are inspired by [PS17].

3.1. Notations and remarks on log smooth morphism. For a log pair (X, D^X) , we mean that X is a normal variety with D^X a \mathbb{Q} -divisor and the log canonical divisor $K_X + D^X$ is \mathbb{Q} -Cartier. We also write $\omega_X(D^X)$ the \mathbb{Q} -line bundle given by the \mathbb{Q} -Cartier divisor $K_X + D^X$. We follow the terminology of singularities of pairs as in [KM98, §2.3].

We say that the pair (X, D^X) is log smooth if X is smooth and the support of D^X , denoted by D^X_{red} , is normal crossing. We denote $\Omega_X(\log D^X) := \Omega_X(\log D^X_{\text{red}})$, the sheaf of log 1-forms with logarithmic poles along D^X_{red} . Notice that we have used $\Omega_X(\log D^X)$ to denote the sheaf of \mathcal{C} -log forms in the sense of Campana in the proof of Theorem 1.0.4. However, Campana's \mathcal{C} -sheaves only make their appearance in the proof of Theorem 1.0.4 in Section 2 but not in the rest part of this paper.

Definition 3.1.1. We say that $f: (X, D^X) \to (Y, D^Y)$ is a morphism of log smooth pairs, if both (X, D^X) and (Y, D^Y) are log smooth, and $f^{-1}(D^Y) := f^*(D^Y)_{\text{red}} \subset D^X_{\text{red}}$. We say that f is strict if $f^{-1}(D^Y) = D^X_{\text{red}}$.

We say that f (as a morphism of log smooth pairs) is log smooth if we further have that f is dominant as a morphism of schemes and the cokernal of the log differential map

$$df: f^*\Omega_Y(\log D^Y) \to \Omega_X(\log D^X)$$

is locally free. In the case that D^Y is empty, we have the following well-known result. For completeness, we give a brief proof here.

Lemma 3.1.2. $f : (X, D^X) \to Y$ being log smooth is equivalent to each stratum (including X) of (X, D^X) being smooth over Y.


Proof. Fix one component D_1^X of D^X . Consider the following commutative diagram, and define \mathcal{A} and \mathcal{B} with all horizontal and vertical complexes forming short

exact sequences

$$\begin{array}{cccc} f^*\Omega_Y(-D_1^X) & \longrightarrow & f^*\Omega_Y & \longrightarrow & (f^*\Omega_Y)|_{D_1^X} \\ & \downarrow & & \downarrow & & \downarrow \\ \Omega_X(\log D^X)(-D_1^X) & \longrightarrow & \Omega_X(\log D^X - D_1^X) & \longrightarrow & \Omega_{D_1^X}(\log(D^X - D_1^X)|_{D_1^X}) \\ & \downarrow & & \downarrow & & \downarrow \\ \Omega_f(-D_1^X) & \longrightarrow & \mathcal{A} & \longrightarrow & \mathcal{B}. \end{array}$$

Working locally on X, if each stratum of (X, D^X) being smooth over Y, then by induction on the number of components of D^X , we can assume that the right two vertical complexes split, and hence the left complex splits which implies that f is log smooth.

On the other hand, consider the following commutative diagram, with all horizontal and vertical complexes forming short exact sequences



If f is log smooth, then locally on X the middle vertical short exact complex splits, which implies the left one splits. This implies the right vertical complex in the first commutative diagram splits. Hence, by induction, each stratum of (X, D^X) is smooth over Y.

In the case that the morphism $f: (X, D^X) \to (Y, D^Y)$ is dominant, we further denote by D_h^X , the horizontal part of D^X , which means it contains all components that are dominant over Y, and by $D_v^X := D^X - D_h^X$, the vertical part of D^X . In this case, we are always making the following assumption in this pape:

(Assumption. 0) D^Y , the boundary divisor of the log smooth base, and D_v^X , the vertical boundary divisor, are reduced, that is, their coefficients are 1, and the coefficients of D_h^X , the horizontal boundary divisors are always in (0, 1].

Notice that (Assumption. 0) is enough for the proof of Theorem 1.0.1 as the log smooth morphism f_V has no boundary on the base and D^U is dominant over V.

Moreover we write by $\lceil f \rceil : (X, \lceil D^X \rceil) \to (Y, \lceil D^Y \rceil = D^Y)$ the morphism with rounding up the pairs. Hence f is log smooth if and only if $\lceil f \rceil$ is.

When we write $\operatorname{Sym}^{i} \mathscr{F}$ (or $\operatorname{Sym}^{[i]} \mathscr{F}$ to be constent with notations from references), we always consider the reflexive hull of the *i*-th symmetric power of the coherent sheaf \mathscr{F} ; similarly, for det \mathscr{F} , we also consider the reflexive hull of the determinant of \mathscr{F} . Actually, due to the following important remark, taking the reflexive hull is also not necessary, since it only modifies the sheaf over a closed subset of codimension ≥ 2 . The next remark will be frequently used in later sections.

Remark 3.1.3. To prove Theorem 1.0.6 we only need to prove it over a *big open* subset of Y, a Zariski open subset with its complement of codimension ≥ 2 , so,

we can feel free to ignore a *small closed subset* of Y, a Zariski closed subset of codimension ≥ 2 . In particular, we can always assume the divisors on Y are smooth. Furthermore, if we have the morphism of log smooth pairs, $f: (X, D^X) \to (Y, D^Y)$, is log smooth outside of the vertical boundary, i.e. f is log smooth over $Y \setminus D^Y$, then by ignoring a small closed subset of Y, we have that f itself is log smooth by the following lemma.

Lemma 3.1.4. Fix a morphism of log smooth pairs, $f : (X, D^X) \to (Y, D^Y)$. Assume that $f|_{X \setminus f^{-1}D^Y}$ is log smooth over $Y \setminus D^Y$, then it is log smooth over a big open subset of Y.

Proof. Since we can ignore all components in D^X that maps to a small closed subset of Y, we only need to show that, for any closed point $p \in E$, where E is any smooth stratum of D^X that dominates a smooth component F of D^Y , the cokernal of the natural morphism $df : f^*\Omega_Y(\log D^Y) \to \Omega_X(\log D^X)$ is locally free, on an open neighborhood U of p. However, restricted on U, the local sections of $\Omega_X(\log D^X)|_U$ are generated by the sections from $\Omega_E(\log D^E)$ and $\{dx_1/x_1, ..., dx_k/x_k\}$, where $D^E := D_h^X|_E$, (noting that E is contained in D_v^X ,) and x_i are local regular functions that defines E. Similarly, locally around f(p), local sections of $\Omega_Y(\log D^Y)$ are generated by the local sections of Ω_F and dy/y, where y is a local function that defines F. However, on U, we have $f^*y = u \Pi_i x_i^{a_i}$, where u is a local unit, and a_i 's are positive integers. Hence we have, $df(f^*dy/y) = \sum_i a_i dx_i/x_i$, which is a local section of $\Omega_X(\log D^X)|_U$ without zero locus. Now we are only left to show that the cokernal of the natural morphism $f|_E^*\Omega_F \to \Omega_E(\log D^E)$ is locally free, over a Zariski open subset of F, which is true by generic smoothness.

Definition 3.1.5. Given a log smooth pair (X, D), we denote

$$\mathcal{T}_{(X,D)} := \Omega_X (\log D)^{\vee}$$
$$\mathscr{A}_{(X,D)}^{\bullet} := \bigoplus_i \operatorname{Sym}^i \mathcal{T}_{(X,D)}$$

that is, $\mathcal{T}_{(X,D)}$ is the sheaf of (algebraic) vector fields with logarithmic zeroes along D^X and $\mathscr{A}^{\bullet}_{(X,D)}$ is the associated graded algebra of symmetric powers. The sheaves $\mathcal{T}_{(X,D)}$ and \mathscr{O}_X generate a subalgebra $\mathscr{D}_{X,D}$ of \mathscr{D}_X , the sheaf of (algebraic) differential operators. We call $\mathscr{D}_{(X,D)}$ the sheaf of (algebraic) log differential operators. The order filtration F_{\bullet} on \mathscr{D}_X induces the order filtration on $\mathscr{D}_{(X,D)}$. With this filtration, we have a canonical isomorphism

$$\mathscr{A}^{\bullet}_{(X,D)} = \mathrm{Gr}^{F}_{\bullet} \mathscr{D}_{(X,D)}$$

where $\operatorname{Gr}^F_{\bullet} \mathscr{D}_{(X,D)}$ denotes the associated graded algebra.

Definition 3.1.6. Given a log smooth morphism of log smooth pairs $f : (X, D^X) \to (Y, D^Y)$,

$$\Omega_f := \operatorname{coker}(f^*\Omega_Y(\log D^Y) \to \Omega_X(\log D^X)),$$

$$\Omega_f^i := \wedge^i \Omega_f,$$

$$\operatorname{DR}_f := [\mathscr{O}_X \to \Omega_f^1 \to \dots \to \Omega_f^k],$$

with cohomology degrees at $-k, ..., 0, k = \dim X - \dim Y$. We call DR_f the relative log de Rham complex of f.

We set $F_p\Omega_f^i = \Omega_f^i$ for $p \ge 0$ and $F_p\Omega_f^i = 0$ for p < 0. Then we have the induced filtration on the complex DR_f :

$$F_{\bullet}\mathrm{DR}_f := [F_{\bullet-k}\mathscr{O}_X \to F_{\bullet-k+1}\Omega^1_f \to \dots \to F_{\bullet}\Omega^k_f].$$

The associated graded complexes are $\operatorname{Gr}_i^F \operatorname{DR}_f = \Omega_f^{k-i}[i]$ for all *i*.

Taking the push-forward functor Rf_* on $(DR_f, F_{\bullet}DR_f)$, we then have the relative log Hodge-to-de Rham spectral squence. See Theorem 3.3.1 for further disussions.

Since D^X and D^Y are possibly \mathbb{Q} -divisors, we write the \mathbb{Q} -line bundle by

$$\omega_f := \omega_X(D^X) \otimes f^*(\omega_Y^{-1}(-D^Y)).$$

Using $\lceil f \rceil$, we particularly have $\omega_{\lceil f \rceil} = \Omega_f^k$. When D^X and D^Y are empty, $\omega_f = \omega_X \otimes f^*(\omega_Y^{-1})$, the relative canonical sheaf of f.

Everything defined as above except ω_f , only depends on the the information of [f]. We still use the sub-index f to make notations simpler.

3.2. Direct image of graded \mathscr{A}^{\bullet} -modules. Given a morphism of log smooth pairs $f: (X, D^X) \to (Y, D^Y)$, fixing a line bundle \mathcal{L} on X, we denote

$$\mathcal{M}^{i}_{f,\mathcal{L},\bullet} := \mathbf{R}^{i} f_{*}(\mathcal{L}^{-1} \otimes \omega_{X}(\lceil D^{X} \rceil) \otimes_{\mathscr{A}^{\bullet}_{(X,D^{X})}}^{\mathbf{L}} f^{*}(\mathscr{A}^{\bullet}_{(Y,D^{Y})} \otimes \omega_{Y}^{-1}(-\lceil D^{Y} \rceil)))$$

for each *i*. By definition, $\mathcal{M}_{f,\mathcal{L},\bullet}^i$ is a graded $\mathscr{A}_{(Y,D^Y)}^{\bullet}$ -modules. Hence it possesses a Higgs-type morphism

$$\theta^{i}_{f,\mathcal{L}}: \mathcal{M}^{i}_{f,\mathcal{L},\bullet} \to \mathcal{M}^{i}_{f,\mathcal{L},\bullet+1} \otimes \Omega_{(Y,D^{Y})}.$$

Denote $\mathcal{K}^i_{f,\mathcal{L},\bullet} := \operatorname{Ker}(\theta^i_{f,\mathcal{L}})$. By repeatedly composing $\theta^i_{f,\mathcal{L}} \otimes \operatorname{Id}$ with itself k times, we have the induced morphism

$$\theta_{f,\mathcal{L},\bullet}^{i,k}:\mathcal{M}_{f,\mathcal{L},\bullet}^{i}\to\mathcal{M}_{f,\mathcal{L},\bullet+k}^{i}\otimes\mathrm{Sym}^{k}\Omega_{(Y,D^{Y})}.$$

The following proposition is the pair analogue of [WW18, Lemma 6.2].

Proposition 3.2.1. If f is log smooth, we have the following quasi-isomorphism of graded $\mathscr{A}^{\bullet}_{(Y,D^Y)}$ -modules

$$\mathcal{M}_{f,\mathcal{L},\bullet}^{i} \stackrel{q.i.}{\simeq} \mathbf{R}^{i} f_{*}(\mathcal{L}^{-1} \otimes \operatorname{Gr}_{\bullet} \operatorname{DR}_{f}).$$

In particular, we have that $\mathcal{M}^{i}_{f,\mathcal{L},\bullet}$ are coherent over \mathscr{O}_{Y} .

Proof. First, we have the relative log Spencer complex

$$\mathscr{A}^{\bullet}_{(X,D^X)} \otimes \mathcal{T}^k_f \to \mathscr{A}^{\bullet}_{(X,D^X)} \otimes \mathcal{T}^{k-1}_f \to \dots \to \mathscr{A}^{\bullet}_{(X,D^X)},$$

given by

$$P \otimes v_1 \wedge \dots \wedge v_p \mapsto \sum_i P v_i \otimes v_1 \wedge \dots \wedge \hat{v}_i \wedge \dots \wedge v_p,$$

for v_i sections of \mathcal{T}_f , where \mathcal{T}_f^i is the \mathscr{O}_X -dual of Ω_f^i . We locally choose a free basis of \mathcal{T}_f^k , denoted by (ξ_1, \ldots, ξ_k) . Then by construction, the relative log Spencer complex locally is the Koszul complex of $\mathscr{A}_{(X,D^X)}^{\bullet}$, with actions given by multiplications of ξ_i , for all *i*. Since *f* is log smooth, we know that (ξ_1, \ldots, ξ_k) is a regular sequence in $\mathscr{A}_{(X,D^X)}^{\bullet}$ and the 0-th cohomology sheaf is $f^*\mathscr{A}_{(Y,D^Y)}^{\bullet}$. Therefore, the relative

log Spencer complex is a locally free resolution of $f^*\mathscr{A}^{\bullet}_{(Y,D^Y)}$. Using the dual-pair $(\Omega^i_f, \mathcal{T}^i_f)$, we then see that $\mathcal{M}_{f,\mathcal{L},\bullet}$ is quasi-isomorphic to

$$\mathbf{R}^i f_*(\mathcal{L}^{-1} \otimes \operatorname{Gr}_{\bullet} \operatorname{DR}_f).$$

Simplification of Notations. To simplify notations, we write

$$\mathcal{M}_{f,\mathcal{L}} = \mathbf{R}^0 f_*(\mathcal{L}^{-1} \otimes \operatorname{Gr}_{\bullet} \operatorname{DR}_f),$$

for the rest of this paper. In the case that ${\cal L}$ is trivial, we omit ${\cal L}$ from the lower-index, that is, we write

$$\mathcal{M}_f^i = \mathbf{R}^i f_*(\mathrm{Gr}_{\bullet}\mathrm{DR}_f) \text{ and } \mathcal{M}_f = \mathcal{M}_f^0.$$

Similarly, when \mathcal{L} is trivial, $\mathcal{K}_{f,\mathcal{L},\bullet}^0$ (resp. $\theta_{f,\mathcal{L},\bullet}^{0,k}$) is simplified to $\mathcal{K}_{f,\bullet}$ (resp. $\theta_{f,\bullet}^k$) or \mathcal{K}_{\bullet} (resp. θ_{\bullet}^k) when there is no ambiguity of f.

3.3. Direct image of filtered log \mathscr{D} -modules and canonical extensions of variation of mixed Hodge structures. We now discuss variations of mixed Hodge structures (VMHS) associated to log smooth morphisms. Let us refer to [Kas86] for the definition of VMHS and admissible VMHS. In contrast to the definition in *loc. cit.*, we assume the Hodge filtrations and the weight filtrations are both increasing filtrations, to be consistent with the good filtration for \mathscr{D} -modules underlying Hodge modules.

The following theorem is well-known; see for instance [FF12] and [Kaw14]. We give it an alternative proof by using Saito's mixed Hodge modules.

Theorem 3.3.1. Suppose that $f : (X, D^X) \to (Y, D^Y)$ is a projective log smooth morphism between log smooth pairs with D^Y smooth (but not necessarily irreducible). We then have:

- (1) The relative log Hodge-to-de Rham spectral sequence degenerate at E_1 .
- (2) We have an admissible VMHS V underlying the local system $\mathbf{R}^{i} f_{*}^{o} \mathbb{Q}_{U^{X}}$, where f^{o} denotes the morphism $f^{o} : U^{X} \coloneqq X \setminus D^{X} \to U^{Y} \coloneqq Y \setminus D^{Y}$. Moreover, if f is strict, then V is a variation of Hodge structure (VHS).
- (3) The associated graded module (with respect to the Hodge filtration) of the upper-canonical extension $V^{\geq 0}$, that is, the logarithmic extension of V with eigenvalues of the monodromy of $\mathbf{R}^i f^o_* \mathbb{Q}_{U^X}$ along D^Y in [0,1)), is \mathcal{M}^i_f .

Proof. We first deal with the case that D^Y is empty. We consider the mixed Hodge module M underlying $Rj_*\mathbb{Q}_{U^X}[n]$, where $j: U^X \hookrightarrow X$. The underlying filtered \mathscr{D}_X -module of M is $\mathscr{O}_X(*D^X)$ with

$$\mathscr{O}_X(*D^X) = \bigcup_{k \in \mathbb{Z}} \mathscr{O}_X(kD^X)$$

and the filtration

$$F_p \mathscr{O}_X(*D^X) = \begin{cases} \mathscr{O}_X((p+1)D^X), & q \ge 0\\ 0, & q < 0 \end{cases}$$

By [BPW17, Proposition 2.3], we obtained a filtered quasi-isomorphism

(3.3.1)
$$f_+(\mathscr{O}_X(*D^X), F_{\bullet}) \simeq Rf_*(\mathrm{DR}_f, F_{\bullet}DR_f)$$

where f_+ denotes the filtered \mathscr{D} -module direct image functor. By the strictness of the Hodge filtration in [Sai88, 3.3.17] and [Sai90, 2.15], the filtered complex

 $f_+(\mathscr{O}_X(*D^X), F_{\bullet})$ is strict or equivalently the relative log Hodge-to-de Rham spectral sequence degenerate at E_1 by the filtered quasi-isomorphism (3.3.1). Hence Part (1) follows in this case.

Since $R^i f_* DR_f$ is \mathscr{O} -coherent and hence it is locally free over \mathscr{O}_Y as

$$R^i f_* \mathrm{DR}_f \simeq \mathcal{H}^i f_+ \mathscr{O}_X(*D^X)$$

are both \mathscr{D}_Y -modules. Hence, the underlying perverse sheaf $\mathbf{R}^i f^o_* \mathbb{Q}_{U^X}[n]$ is locally constant (upto a shift). By [Sai90, 3.27], we hence obtain Part (2) in this case.

If D^Y is smooth but not empty, then one can use the double-strictness of direct image functor for mixed Hodge modules of normal crossing type (see [Sai90, §3] and also [Wei17]). More precisely, one applies for instance [Wei17, Theorem 15] and Part (1), (2) and (3) follow in general.

We also need the following weak negativity of Kodaira-Spencer kernals proved in [PW16].

Theorem 3.3.2. [PW16, Theorem 4.8] In the situation of Theorem 3.3.1, if f is strict, then \mathcal{K}_p^{\vee} is weakly positive for each $p \in \mathbb{Z}$, where \bullet^{\vee} denotes the \mathcal{O} -dual of \bullet .

3.4. Adding extra boundary divisors. We assume that $f: (X, D^X) \to (Y, D^Y)$ is a log smooth morphism of log smooth pairs. If we add a divisor D' to D^Y (assuming D' is not supported on D^Y) and obtain another normal crossing divisor D'^Y over a big open subset of Y (by getting rid of the singular locus of D' and the intersection of D' and D^Y), then we define D'^X by $D'^X_h = D^X_h$ and $D'^X_v = f^{-1}D'^Y$, and denote $f': (X, D'^X) \to (Y, D'^Y)$ the new log smooth morphism (by Lemma 3.1.4). Since f is smooth away from D^Y , by a local computation one easily obtains that

(3.4.1)
$$\Omega_f \simeq \Omega_{f'},$$

over a big open subset of Y, which further implies

(3.4.2)
$$\mathcal{M}^i_{f,\mathcal{L}} \simeq \mathcal{M}^i_{f',\mathcal{L}},$$

as graded $\mathscr{A}_{(Y,D^Y)}$ -modules over a big open subset of Y.

More generally, we have the following.

Proposition 3.4.1. Fix a projective log smooth morphism $f : (X, D^X) \to (Y, D^Y)$, and a morphism of log smooth pairs $\eta : (Y', D^{Y'}) \to (Y, D^Y)$, with the underlying morphism of schemes $Y' \to Y$ being smooth. Let $X' = X \times_Y Y'$, $D^{Y'}$ be a SNC divisor containing $\eta^{-1}D^Y$ and $f' : (X', D^{X'}) \to (Y', D^{Y'})$ be the induced morphism of log smooth pairs with $D^{X'} = f'^{-1}D^{Y'} \cup \overline{\eta}^{-1}D^X$ in the following diagram:

$$\begin{array}{ccc} (X', D^{X'}) & \stackrel{\overline{\eta}}{\longrightarrow} (X, D^X) \\ & \downarrow^{f'} & \downarrow^{f} \\ (Y', D^{Y'}) & \stackrel{\eta}{\longrightarrow} (Y, D^Y) \end{array}$$

Then, over a big open subset of Y', we have an natural isomorphism of graded $\mathscr{A}^{\bullet}_{(Y',DY')}$ -modules

$$\eta^* \mathcal{M}_f \simeq \mathcal{M}_{f'}.$$

Proof. We first note that, using the identity (3.4.2), we only need to show the case when $D^{Y'} = \eta^{-1}D^Y$. In this case we also have $D^{X'} = \bar{\eta}^{-1}D^X$. The required statement then follows from Proposition 3.2.1 and the smooth base-change.

Proposition 3.4.2. Assume that $f: (X, D^X) \to (Y, D^Y)$ is a projective log smooth morphism with $D^X = f^{-1}D^Y$ (that is, f is strict), and $\psi: Y' \to Y$ is a finite morphism branched over D^Y with $\psi^{-1}D^Y$ normal crossing. Let $(X', D^{X'})$ be a log resolution of the normalization of the main component of $X \times_Y Y'$ with $D^{Y'}$ a SNC divisor containing $\psi^{-1}D^Y$ and $D^{X'} = f'^{-1}D^{Y'}$ (we can assume that $D^{X'}$ is also normal crossing after taking further log resolutions), and $f': (X', D^{X'}) \to$ $(Y', D^{Y'})$ be the induced projective morphism of log smooth pairs in the following diagram

$$(X', D^{X'}) \xrightarrow{\psi} (X, D^X)$$
$$\downarrow^{f'} \qquad \qquad \downarrow^f$$
$$(Y', D^{Y'}) \xrightarrow{\psi} (Y, D^Y)$$

Then, over a big open subset of Y', we have a natural inclusion of $\mathscr{A}^{\bullet}_{(Y',DY')}$ -modules

$$\psi^* \mathcal{M}_f \to \mathcal{M}_{f'}.$$

Proof. Since over $Y' \setminus D^{Y'} \psi$ is étale, we have $\psi^* \mathcal{M}_f$ is identical with $\mathcal{M}_{f'}$ over $Y' \setminus D^{Y'}$, by Proposition 3.4.2. Then by Theorem 3.3.1 (3), we further have that \mathcal{M}_f (resp. $\mathcal{M}_{f'}$) is the associated graded module of the upper-canonical extension $V_f^{\geq 0}$ (resp. $V_{f'}^{\geq 0}$) of the VHS of the 0-th cohomology of the smooth fibers of f (resp. f'). By [PW16, Proposition 4.4], we see that

$$\psi^* V_f^{\ge 0} \subseteq V_{f'}^{\ge 0}.$$

Then we apply [PW16, Corollary 4.7] for each filtrant of $V_f^{\geq 0}$ and $V_{f'}^{\geq 0}$ in their Hodge filtrations and obtain the required inclusion.

3.5. Birationally invariant Hodge modules.

Theorem 3.5.1. Given a projective dominant morphism $f : X \to Y$ with X and Y being smooth, there exists a pure Hodge module M_f on Y, which corresponds to a generically defined VHS (by using the equivalence in [Sai90, 3.21]), such that for any projective morphism $f' : X' \to Y$ with X' being smooth and birational to X, M_f is isomorphic to a direct summand of $\mathcal{H}^0 f'_+ \mathbb{Q}^H_{X'}$, and its lowest filtered piece in the Hodge filtration is isomorphic to $f_*\omega_f = f'_*\omega_{f'}$, where $\mathbb{Q}^H_{X'}$ is the trivial pure Hodge module on X' and f'_+ denotes the (derived) direct image functor for Hodge modules.

Proof. We first fix \tilde{X} , a common resolution of X and X'.

$$X \xleftarrow{\phi_f} \tilde{X} \xrightarrow{\psi} X'$$

Note that $f'_*\omega_{f'} = f_*\omega_f = \tilde{f}_*\omega_{\tilde{f}}$ as X and X' are both smooth. Now we define M_f the smallest sub-Hodge module of $\mathcal{H}^0\tilde{f}_+\mathbb{Q}^H_{\tilde{X}}$ that contains $\tilde{f}_*\omega_{\tilde{f}}$. More precisely,

due to the semi-simplicity of pure Hodge modules (see [Sai88, §5]), we define \mathcal{M} the sub-pure Hodge module of $\mathcal{H}^0 \tilde{f}_+ \mathbb{Q}_{\tilde{X}}^H$ that consists of all the simple sub-pure Hodge modules that intersect $\tilde{f}_* \omega_{\tilde{f}}$ non-trivially. Thanks to the decomposition theorem of the direct image of pure Hodge modules under projective morphism [Sai88, Théorème 1 and Corollaire 3] (see also [Sch14, §16]) and the equivalence in [Sai90, 3.21], we have that both $\mathcal{H}^0 f_+ \mathbb{Q}_X^H$ and $\mathcal{H}^0 f'_+ \mathbb{Q}_{X'}^H$ are direct summands of $\mathcal{H}^0 \tilde{f}_+ \mathbb{Q}_{\tilde{X}}^H$, and all three pure Hodge modules share the same lowest filtered piece. As a consequence, we conclude that M_f satisfies the conditions.

Proposition 3.5.2. Fix two projective morphisms of log smooth pairs

$$f_i: (X_i, D^{X_i}) \to (Y, D^Y), \ i = 1, 2,$$

with D^Y a smooth divisor (but not necessarily irreducible) and X_1 and X_2 birational. Assume that $D^{X_i} = f_i^{-1}D^Y$ (in particular, D^{X_i} are SNC divisors) and f_i are smooth over $Y \setminus D^Y$, for i = 1, 2. Recalling the notations in §3.2, we can find a graded $\mathscr{A}^{\bullet}_{(Y,D^Y)}$ -module $\hat{\mathcal{M}}$, which is a direct summand of both \mathcal{M}_{f_i} for i = 1, 2. Furthermore, all of $\hat{\mathcal{M}}, \mathcal{M}_{f_i}$ share the same lowest graded piece.

Proof. Recall that \mathcal{M}_{f_i} are isomorphic to the associated graded modules of the upper-canonical extension along D^Y of the VHS given by the restriction of $\mathcal{H}^0 f_{i+} \mathbb{Q}^H_{X_i}$ on $Y \setminus D^Y$ (see Theorem 3.3.1(2) and (3)). By the previous theorem, we get a pure Hodge module M_{f_1} that corresponds to a VHS on $Y \setminus D^Y$, called V. Now we set $\hat{\mathcal{M}}$ to be the associated graded module of the canonical extension along D^Y of V, with the real part of the eigenvalues of the residues in [0, 1). \Box

3.6. Construction of Higgs sheaves from a section. Suppose that we have the following diagram of projective morphisms of log smooth pairs:

$$(Z, D^Z) \xrightarrow{\pi} (X, D^X)$$

$$\xrightarrow{g} \qquad \qquad \downarrow^f$$

$$(Y, D^Y),$$

with π generically finite, and both f and g being log smooth. By forgetting the horizontal part of D^Z respect to g, we denote $g^0: (Z, D_v^Z) \to (Y, D^Y)$. Note that, by Lemma 3.1.2 and Lemma 3.1.4, g^0 is log smooth at least over a big open subset of Y. For simplicity, in the rest of this subsection, we assume g^0 is log smooth by removal a small closed subset of Y.

We have a natural morphism $\pi^* \Omega_X(\log D^X) \to \Omega_Z(\log D^Z)$, which induces a natural graded morphism of graded complexes

(3.6.1)
$$\pi^* \operatorname{Gr}_{\bullet} \operatorname{DR}_f \to \operatorname{Gr}_{\bullet} \operatorname{DR}_q.$$

For invertible sheaves \mathcal{L} and \mathcal{A} on X and Y respectively, we assume the following two assumptions:

(Assumption. 1) $H^0(Z, \pi^* \mathcal{L} \otimes g^* \mathcal{A}^{-1}) \neq 0.$

(Assumption. 2) the support of the effective Cartier divisor (θ) contains D_h^Z .

By (Assumption. 1), fixing one non-trivial section

$$\theta \in H^0(Z, \pi^*\mathcal{L} \otimes g^*\mathcal{A}^{-1}) \neq 0,$$

we have an induced inclusion

$$\tau^* \mathcal{L}^{-1} \to g^* \mathcal{A}^{-1}.$$

We then have an induced graded morphism

(3.6.2)
$$\pi^*(\operatorname{Gr}_{\bullet}\operatorname{DR}_f \otimes \mathcal{L}^{-1}) \to \operatorname{Gr}_{\bullet}\operatorname{DR}_g \otimes g^*\mathcal{A}^{-1}.$$

as the composition of the natural morphisms

$$\pi^*(\mathrm{Gr}_{\bullet}\mathrm{DR}_f \otimes_{\mathscr{O}} \mathcal{L}^{-1}) \to \mathrm{Gr}_{\bullet}\mathrm{DR}_g \otimes_{\mathscr{O}} \pi^* \mathcal{L}^{-1} \to \mathrm{Gr}_{\bullet}\mathrm{DR}_g \otimes_{\mathscr{O}} g^* \mathcal{A}^{-1}.$$

By (Assumption. 2), the morphism (3.6.2) naturally factors through

$$\pi^*(\operatorname{Gr}_{\bullet}\operatorname{DR}_f \otimes \mathcal{L}^{-1}) \to \operatorname{Gr}_{\bullet}\operatorname{DR}_g(-D_h^Z) \otimes g^*(\mathcal{A}^{-1}).$$

Note that the natural inclusion $\Omega_Z(\log D^Z)(-D_h^Z) \to \Omega_Z(\log D_v^Z)$ induces a natural inclusion of complexes:

$$\operatorname{Gr}_{\bullet}\operatorname{DR}_g(-D_h^Z) \to \operatorname{Gr}_{\bullet}\operatorname{DR}_{g^0}.$$

We then have an induced morphism

$$\pi^*(\mathrm{Gr}_{\bullet}\mathrm{DR}_f\otimes\mathcal{L}^{-1})\to\mathrm{Gr}_{\bullet}\mathrm{DR}_{g^0}\otimes g^*\mathcal{A}^{-1}.$$

This further induces a morphism

$$\eta_{\theta}: \mathcal{M}_{f,\mathcal{L}} \to \mathcal{M}_{g^0,g^*\mathcal{A}},$$

as graded $\mathscr{A}^{\bullet}_{(Y,D^Y)}$ -modules by Proposition 3.2.1. Furthermore, the lowest graded piece of η_{θ} :

$$g_*\pi^*(\omega_f\otimes\mathcal{L}^{-1})\to g_*\omega_{g^0}\otimes\mathcal{A}^{-1})$$

is induced by the morphism $\pi^*(\omega_f \otimes \mathcal{L}^{-1}) \to \omega_{g^0} \otimes g^* \mathcal{A}^{-1}$. Since $\pi^*(\omega_f \otimes \mathcal{L}^{-1}) \to \omega_{g^0} \otimes g^* \mathcal{A}^{-1}$ is induced by θ and hence injective, by the left exactness of g_* , we see that

$$g_*\pi^*(\omega_f \otimes \mathcal{L}^{-1}) \to g_*\omega_{g^0} \otimes \mathcal{A}^{-1},$$

is also injective.

4. Geometric construction

4.1. Stable reduction for families with arbitrary variation. In [WW18, §4], we introduced stable reduction for log smooth families with maximal variation. In this section, we discuss stable reduction for log smooth families with arbitrary variation.

Suppose that $f_V: (U, D^U) \to V$ is a log smooth family of projective log smooth pairs of log general type, with (U, D^U) being klt and V smooth. Consider the relative canonical model

$$f_{V,c}\colon (U_c, D^{U_c}) \to V.$$

We write v the volume of $K_{U_{c,y}} + D_y^{U_c}$ for $y \in V$ (by invariance of pluri-genera, v is constant over V). Let I be a finite coefficient set I closed under addition and containing the coefficients of D^{U_c} . We consider the coarse moduli space, denoted by \mathfrak{M} , for stable log varieties of a fixed dimension, volume v and the coefficient set I. Let us refer to [KP16, §6] for the construction of \mathfrak{M} and related properties. It is proved in *loc. cit.* that \mathfrak{M} is a projective reduced scheme and the corresponding moduli stack is an Deligne-Mumford stack. The relative canonical model $f_{V,c}$ (, which is a stable family thanks to invariance of pluri-genera in the log smooth case [HMX18, Theorem 4.2],) induces a moduli map $V \longrightarrow \mathfrak{M}$. Since \mathfrak{M} is projective,

we take a projective compactification Y of V so that $E^Y = Y \setminus V$ is normal crossing and the moduli map extends to a projective map $Y \to \mathfrak{M}$ (we probably need to replace f_V by a birational model).

Proposition 4.1.1. Under the above construction, replacing Y by a further resolution and removing a small closed subset, we can construct the following commutative diagram

(4.1.1)

$$\begin{array}{cccc} (X_c, D^{X_c}) & \xleftarrow{\psi_c} & (X'_{1,c}, D^{X'_{1,c}}) & \xrightarrow{\rho} & (X'_{2,c}, D^{X'_{2,c}}) & \xrightarrow{\eta_c} & (X^{\sharp}_c, D^{X^{\sharp}_c}) \\ & \downarrow^{f_c} & \Box_1 & & & & \\ & Y & \xleftarrow{\eta} & & & & & \\ & Y & \xleftarrow{\eta} & & & & & & \\ \end{array}$$

such that

(0) $(X_c^{\bullet}, D^{X_c^{\bullet}})$ are all klt pairs with $\bullet = \emptyset_{,1}, \emptyset_{,2}, \sharp$;

(1) f_c^{\sharp} is a stable family with maximal variation;

- (2) the square \Box_1 is Cartesian over V;
- (3) the square \Box_2 is Cartesian;
- (4) η is a dominant smooth morphism, not necessarily proper;
- (5) ψ is finite and flat;
- (6) for all sufficiently positive and divisible m, we have isomorphisms

$$f'_{2,c*}\omega^m_{X'_{2,c}/Y'}(mD^{X'_{2,c}}) \simeq \eta^* f^{\sharp}_* \omega^m_{X^{\sharp}_c/Y^{\sharp}}(mD^{X^{\sharp}_c});$$

$$\det f'_{2,c*}\omega^m_{X'_c-/Y'}(mD^{X'_{2,c}}) \simeq \eta^* \det f^{\sharp}_* \omega^m_{X^{\sharp}_c/Y^{\sharp}}(mD^{X^{\sharp}_c});$$

(7) for sufficiently positive and divisible N_m (depending on m), we can find a line bundle \mathcal{A}_m on Y such that

$$\psi^* \mathcal{A}_m \simeq (\det f'_{2,c*} \omega^m_{X'_{2,c}} / Y' (mD^{X'_{2,c}}))^{N_m}$$

Proof. By [LMB00, Theorem 16.6], \mathfrak{M} has a finite covering $S \to \mathfrak{M}$ which is induced by a stable family over S in the moduli stack of \mathfrak{M} . We then take Y' to be a desingularization of the main component of $Y \times_{\mathfrak{M}} S$ (the component dominant Y'). Then we got a generic finite map $\psi \colon Y' \to Y$. We then take the Stein factorization of the induced morphism $Y' \to S$ and obtain a fibration $\eta: Y' \to Y^{\sharp}$. Thanks to Raynaud-Gruson flattening theorem ([RG71, Théorèm 5.2.2]), after replacing Y^{\sharp} by a further resolution, we can assume that $\eta: Y' \to Y^{\sharp}$ is a flat fibration, with Y^{\sharp} being projective and smooth. Note that Y' might have more than 1 components, and we then replace Y' by the main component. Now we take a Kawamata covering $Y^{\sharp'} \to Y^{\sharp}$ so that there exists a Y'', a desingularization of the main component of $Y' \times_{Y^{\sharp}} Y^{\sharp'}$ satisfying that $Y'' \to Y^{\sharp'}$ is semistable in codimension 1; semistable means that the morphism is flat with reduced fibres, see for instance [AK00] for a stronger result. In particular, we can remove a small closed subset of Y'' to make the morphism be smooth. Replace Y' by Y'' and Y^{\sharp} by $Y^{\sharp'}$. To finish the construction of the bases, we remove the largest reduced divisor F on Y' such that $\operatorname{codim} \psi(F) \geq 2$, and replace Y by the image of Y' (with F removal) under ψ . Note that we only removed a small subset of Y.

Now we start to construct the X^{\bullet} level. We first take a klt log smooth pair (X, D^X) with a projective morphism $f_Y : (X, D^X) \to Y$ so that $f_Y|_V = f_V$. Since

general fibers of f are of log general type, by [BCHM10, Theorem 1.2] we take f_c to be the relative log canonical model of f_Y . In particular, (X, D^{X_c}) is a klt pair. We then set $X'_{1,c}$, the main component of the normalization of $X_c \times_Y Y'$, with boundary divisor $D^{X'_{1,c}}$ given by $\omega_{X'_{1,c}}(D^{X'_{1,c}}) = \psi_c^* \omega_{X_c}(D^{X_c})$. Since ψ is finite, so is ψ_c , hence $(X'_{1,c}, D^{X'_{1,c}})$ is klt by [KM98, Proposition 5.20]. The induced morphism $Y^{\sharp} \to S$ is generic finite, so it induces a stable family $f_c^{\sharp} : (X_c^{\sharp}, D^{X_c^{\sharp}}) \to Y^{\sharp}$. $f'_{2,c} : (X'_{2,c}, D^{X'_{2,c}}) \to Y'$ is defined as the stable family induced by f_c^{\sharp} under the smooth base change η . Note that by construction, we have that $f'_{1,c}$ and $f'_{2,c}$ coincide on $\psi^{-1}V$, hence we can blowup $(X'_{1,c}, D^{X'_{1,c}})$ to make ρ a morphism, without changing $f'_{1,c}$ over $\psi^{-1}V$. Now the morphisms satisfy (1) to (5), and (6) follows by the flat base change.

To prove (7), we use the argument similar to the proof of [VZ03, Corollary ix)]. Note that, over an open set $V_0 \subset Y$, since $f'_{2,c}$ can be induced by f_c using a flat base change ψ , we can canonically identifying $\psi^* \det f_* \omega^m_{X_c/Y}$ and $\det f'_{2,c*} \omega^m_{X'_{2,c}/Y'}(mD^{X'_{2,c}})$ over $V'_0 := \psi^{-1}V_0$, and set *B* the divisor support on $Y' \setminus V'_0$, satisfying

$$\psi^* \det f_* \omega^m_{X_c/Y}(mD^{X_c}) = \det f'_{2,c*} \omega^m_{X'_2,c/Y'}(mD^{X'_{2,c}})(B),$$

that is also canonically defined. Now we only need to show that B is the pullback of some \mathbb{Q} -divisor on Y, so only need to show that, for any two components B_1 and B_2 , if they have a same image under ψ , then they share a same coefficient. Now we only need to show that over a general point p of $\psi(B_1)$. Take a local curve Qthat only intersect with $\psi(B)$ at p. Now f_Q , the restriction of f_c onto Q, is stable over $Q \setminus p$, so we can find a cyclic cover $\pi : \tilde{Q} \to Q$ that is totally ramified at p, such that the induced new family \tilde{f}_Q over \tilde{Q} is stable and compatible with the base change of f_Q over $Q \setminus p$. Denote Q' any irreducible component of $\psi^{-1}Q$, and let $\tilde{Q}' = \tilde{Q} \times_Q Q'$. Although we can induce two stable families over \tilde{Q}' , but since they are the same over general point, hence they are the same family, by the properness of the moduli functor, and we denote the family by $f_{\tilde{Q}'}$. Note that $f_{\tilde{Q}}, f_{Q'}$ and $f_{\tilde{Q}'}$ are all compatible with flat base change, so we only need to verify the proposition on f_Q and $f_{\tilde{Q}}$. This is true, since π is totally ramified at p.

Since f_c^{\sharp} is a stable family with maximal variation, we have det $f_{c*}^{\sharp} \omega_{X_c^{\sharp}/Y^{\sharp}}^m (mD^{X_c^{\sharp}})$ is big by [KP16, Theorem 7.1]. We now fix a pair of m and N_m as in the previous proposition. To simplify notation, we set

$$\begin{aligned} \mathcal{A}^{\sharp} &:= (\det f_{c*}^{\sharp} \omega_{X_{c}^{\sharp}/Y^{\sharp}}^{m} (mD^{X_{c}^{\sharp}}))^{N_{m}} \\ \mathcal{A}' &:= (\det f_{2,c*}' \omega_{X_{2,c}'/Y'}^{m} (mD^{X_{2,c}'}))^{N_{m}} \\ \mathcal{A} &:= \mathcal{A}_{m}. \end{aligned}$$

By Part (6) and (7) in Proposition 4.1.1, we have

(4.1.2)
$$\psi^* \mathcal{A} = \mathcal{A}' = \eta^* \mathcal{A}^{\sharp} \text{ and } \kappa(\mathcal{A}) = \kappa(\mathcal{A}') = \kappa(\mathcal{A}^{\sharp}) = \operatorname{Var}(f_V).$$

4.2. Birational refinement of the stable reduction. In this subsection, we refine the stable reduction in Proposition 4.1.1, which will be used to construct Viehweg-Zuo sheaf in later section. We first show the following lemma.

Lemma 4.2.1. Assume that we have two log smooth projective morphisms

$$f_i: (X_i, D^{X_i}) \to (Y, D^Y), with \ i = 1, 2,$$

whose generic fibers share the same birational equivalent model, i.e. we have log resolutions over ξ , $\phi_{i,\xi} : (X_{\xi}, D^{X_{\xi}}) \to (X_{i,\xi}, D^{X_{i,\xi}})$, satisfying

$$\phi_{i,\xi}^*\omega_{X_{\xi}}(D^{X_{i,\xi}}) \simeq \omega_{X_{\xi}}(D^{X_{\xi}}),$$

where ξ is the generic point of Y. After adding components to D^Y if necessary (cf. §3.4), forgetting the information over a small closed subset of Y, and replacing $D_v^{X_i}$ by $f^{-1}(D^Y)$, we have $f_{1*}\omega_{f_1}^m = f_{2*}\omega_{f_2}^m$.

Proof. According to the assumption, we can have a common log resolution, hence the following commutative diagram, with all pairs are log smooth,

so that $D^X|_{X_{\xi}} = D^{X_{\xi}}$. We can assume that every irreducible component of D^X is dominant over Y, i.e. the vertical part of D^X over Y is zero. After we extend D^Y and forget a small closed subset of Y, we can assume that f is smooth away from D^Y without changing log smoothness of f_1 and f_2 . We then replace the vertical parts of D^{X_1} , D^X and D^{X_2} by $f_1^{-1}(D^Y)$, $f^{-1}(D^Y)$ and $f_2^{-1}(D^Y)$ respectively. Using Lemma 3.1.4, we can assume that all vertical morphisms are projective log smooth morphisms.

To prove the statement, by symmetry we only need to show that $\phi_{1*}(\omega_X(D^X))^m = (\omega_{X_1}(D^{X_1}))^m$. Consider a Cartier divisor B with

$$\mathscr{O}(B) = (\omega_X(D^X))^m \otimes \phi_1^*(\omega_{X_1}(D^{X_1}))^{-m},$$

which is an ϕ_1 -exceptional divisor. Since f is log smooth, any irreducible ϕ_1 -exceptional divisor E on X maps to D^Y or dominates Y. By the assumption of f_i over the generic point of Y, E cannot dominant Y. Then E maps to D^Y and E is f-exceptional. Hence the coefficient of E in D^X is 1 by construction (see (Assumption. 0) in §3.1). Since (X_1, D^{X_1}) is log canonical (by (Assumption. 0) in §3.1), the coefficient of E in B is positive, which concludes the proof.

Proposition 4.2.2. Assume that we have a commutative diagram (4.1.1) as in Proposition 4.1.1, we can always find a log resolution $\pi : (X, D^X) \to (X_c, D^{X_c})$, so that we can have the following commutative diagram, allowing forgetting a small closed subset of Y, Y' and Y^{\sharp} .

(4.2.1)

such that

(1) all the log pairs are log smooth;

(2) f is a projective compactification of f_V ;

CHUANHAO WEI AND LEI WU

- (3) all the downwards morphisms are log smooth, with vertical boundaries are reduced and the coefficients of horizontal boundaries are in (0, 1);
- (4) the square \Box_1 is Cartesian over $Y \setminus D^Y$ and ψ is étale over $Y \setminus D^Y$;
- (5) the square \Box_2 is Cartesian;
- (6) all ϕ_{\bullet} are birational, with $\bullet = 1, 2;$
- (7) $\bar{\psi}$ factors through the natural morphism $\psi_0 : X'_0 \to X$, where X'_0 is the normalization of the main component of $X \times_Y Y'$, and $\phi_0 : X'_1 \to X'_0$ is birational, and denote $f'_0 : X'_0 \to Y'$ the induced morphism;
- (8) for all m sufficiently positive and divisible, we have isomorphisms

$$f_{0*}'\psi_0^*\omega_f^m \simeq f_{1*}'\bar{\psi}^*\omega_f^m \simeq f_{1*}'\omega_{f_1'}^m \simeq f_{*}'\omega_{f'}^m \simeq f_{2*}'\omega_{f_2'}^m \simeq \eta^*f_{*}^{\sharp}\omega_f^m \sharp$$

(9) denoting $\lceil f^{\bullet} \rceil : (X^{\bullet}, \lceil D^{X^{\bullet}} \rceil) \to (Y^{\bullet}, D^{Y^{\bullet}})$, with $\bullet = \emptyset, '_{1}, '_{2}, ^{\sharp}$, i.e. only considering the boundary with reduced structure, for all m sufficiently positive and divisible, we have isomorphisms


$$f'_{0*}\psi^*_0\omega^m_{\lceil f\rceil} \simeq f'_{1*}\bar{\psi}^*\omega^m_{\lceil f\rceil} \simeq f'_{1*}\omega^m_{\lceil f'_1\rceil} \simeq f'_*\omega^m_{\lceil f'_1\rceil} \simeq f'_{2*}\omega^m_{\lceil f'_2\rceil} \simeq \eta^*f^\sharp_*\omega^m_{\lceil f^\sharp_1\rceil}.$$

Proof. From Proposition 4.1.1, we first replace $(X_c^{\bullet}, D^{X_c^{\bullet}})$ by their log resolutions $(X^{\bullet}, D^{X^{\bullet}})$, with $\bullet = \emptyset_{,1,2}^{\prime}, \sharp$, and the horizontal part of the boundary divisor $D^{X^{\bullet}}$ is defined by taking the positive and horizontal part of $\omega_{X^{\bullet}}^{-1} \otimes \pi^{\bullet*} \omega_{X_c^{\bullet}}(D^{X_c^{\bullet}})$, where π^{\bullet} are the corresponding log resolutions. Since $(X_c^{\bullet}, D^{X_c^{\bullet}})$ are klt, so are (X, D^X) . Moreover, by the construction of f_c in Proposition 4.1.1, we can assume $f: (X, D^X) \to Y$ is a projective compactification of the initial log smooth family $f_V: (V, D^U) \to V$. In particular, we can keep track the initial boundary divisor $E^Y = Y \setminus V$ because ultimately we are interested in the log Kodaira dimension of V.

We do not consider the vertical parts of $D^{X^{\bullet}}$ at this stage, but they will be specified after we fixed the boundary divisors on Y^{\bullet} . Since base changing by an étale or smooth morphism will keep the log smoothness, we can keep \Box_1 and \Box_2 are Cartesian over the assigned locus, and make $\bar{\psi}$ and $\bar{\eta}$ are morphisms of log smooth pairs. By the construction of $f'_{1,c}$ in Proposition 4.1.1, we have (7).

Set D^Y and $D^{Y^{\sharp}}$ be divisors on Y and Y^{\sharp} respectively, so that f and f^{\sharp} are log smooth over the base outside of the boundary (cf. Lemma 3.1.2). By expanding D^Y , we can assume that ψ is étale over $Y \setminus D^Y$. Set $D^{Y'}$ a divisor on Y', so that it contains both $\psi^{-1}D^Y$ and $\eta^{-1}D^{Y^{\sharp}}$, and both f'_1 and f'_2 are log smooth over $Y \setminus D^{Y'}$. By Lemma 3.1.4, also Remark 3.1.3, we can assume that f'_1 and f'_2 are log smooth, by removing a small subset of Y'. Hence, we can apply the previous lemma, by further expending $D^{Y'}$, constructing f' as in the proof of the lemma, and replacing all vertical boundaries by the inverse image $D^{Y'}$, we have the identities of the third term to the fifth term of (8). Then, we replace D^Y and $D^{Y^{\sharp}}$, by $\psi_*D^{Y'}$ and $\eta_*D^{Y'}$. Now we expanding $D^{X^{\bullet}}$ by setting $D^{X^{\bullet}_v} = f^{\bullet -1}D^{Y^{\bullet}}$. Meanwhile we take further log resolutions of $(X^{\bullet}, D^{X^{\bullet}})$ if needed, to keep them being log smooth, but do not change the locus that are already log smooth, so that we can keep the Cartesian condition. At last, remove some small subsets of Y, Y', and Y^{\sharp} , we have all the downwards morphisms are log smooth.

18



The first identity in (8) is due to the projection formula. Note that ψ is étale outside of the boundary, so we have $\psi^* \omega_Y(D^Y) \simeq \omega_{Y'}(D^{Y'})$. Set $D^{X'_0}$ by $\bar{\psi}_0^* \omega_X(D^X) = \omega_{X'_0}(D^{X'_0})$. Since \Box_1 is Cartesian over $Y \setminus D^Y$, so we have that the ramification locus of $\bar{\psi}_0$ is contained in D_v^X , which is reduced. This implies that $D_v^{X'_0}$ is also reduced and $\phi_{0*}(D^{X'_1}) = D^{X'_0}$. Hence the support of $(\bar{\psi}^* \omega_X(D^X))^{-1} \otimes \omega_{X'_1}(D^{X'_1})$ is contained in the exceptional divisor respect to ϕ_0 , hence also contained in $D_v^{X'_1}$, and it is an effective \mathbb{Q} -divisor, due to (X, D^X) is log canonical, and $D_v^{X'_1}$ is reduced. This implies that $f'_{1*}\bar{\psi}^*\omega_X^m(mD^X) \simeq f'_{1*}\omega_{X'_1}^m(mD^{X'_1})$, which concludes the second identity.

The last identity of (8) can be deduced by flat base change, combining the fact that we have η is smooth and $X'_2 = X^{\sharp} \times_{Y^{\sharp}} Y'$, so $\bar{\eta}^* D^{X^{\sharp}} = D^{X'_2}$ and $\eta^* D^{Y^{\sharp}} = D^{Y'}$.

Since we only need that all log pairs are log canonical in proving all identities in (8), we get (9).

4.3. Viehweg's fiber product trick and construction of cyclic covering. We continue assuming that we are in the situation of Proposition 4.1.1. We then have three line bundles \mathcal{A}^{\bullet} on Y^{\bullet} respectively, with $\bullet = \emptyset$, ' or \sharp , satisfying

(4.3.1)
$$\psi^* \mathcal{A} = \mathcal{A}' = \eta^* \mathcal{A}^\sharp$$

as in (4.1.2)

Recall that $\mathcal{A}^{\sharp} = (\det f_{c*}^{\sharp} \omega_{X_c^{\sharp}/Y^{\sharp}}^{m} (mD^{X^{\sharp}}))^{N_m}$, and \mathcal{A}^{\sharp} is a big line bundle. Hence, we can assume that $\mathcal{A}^{\sharp N}(-D^{Y^{\sharp}})$ has a global section, for some positive integer N, where $D^{Y^{\sharp}}$ is the divisor as in (4.2.1). Since f_c is stable, we see that $f_{c*}^{\sharp} \omega_{X^{\sharp}/Y^{\sharp}}^{m} (mD^{X_c^{\sharp}})$ is reflexive. Then, we have a natural inclusion

$$\det f^{\sharp}_{c*}\omega^{m}_{X^{\sharp}_{c}/Y^{\sharp}}(mD^{X^{\sharp}_{c}}) \to (f^{\sharp}_{c*}\omega^{m}_{X^{\sharp}_{c}/Y^{\sharp}}(mD^{X^{\sharp}_{c}}))^{\otimes r_{0}},$$

where r_0 is the rank of $f_{c*}^{\sharp} \omega_{X_c^{\sharp}/Y^{\sharp}}^m (mD^{X_c^{\sharp}})$. These give us a sequence of inclusions

$$(\mathcal{A}^{\sharp}(D^{Y^{\sharp}}))^{m} \to \mathcal{A}^{\sharp m(N+1)} \to (f_{c*}^{\sharp} \omega_{X_{c}^{\sharp}/Y^{\sharp}}^{m}(mD^{X_{c}^{\sharp}}))^{\otimes r},$$

where $r = m \cdot N_m(N+1)$. In particular, we have

$$H^{0}(Y^{\sharp}, (f^{\sharp}_{c*}\omega^{m}_{X^{\sharp}_{c}/Y^{\sharp}}(mD^{X^{\sharp}_{c}}))^{\otimes r} \otimes (\mathcal{A}^{\sharp}(D^{Y^{\sharp}}))^{-m}) \neq 0.$$

We recall Viehweg's fiber product trick (see [Vie83] and also [WW18, Section 4] for the log case). Suppose that we have a morphism $f : X \to Y$ with X and Y both smooth. We then define

$$X_Y^r := X \times_Y X \times_Y \cdots \times_Y X,$$

the r-th fiber product, and

$$f^r: X^r_Y \to Y$$

the induced morphism. If D^X is a Q-Cartier divisor on X, we write

$$(D^X)_Y^r = \sum_{i=1}^r p_i^* D^X,$$

where $p_i \colon X_Y^r \longrightarrow X$ is the *i*-th projection; X_Y^r and $(D^X)_Y^r$ are also denoted by X^r and D^r respectively, if f is obvious from the context.

We first apply the fiber product trick on f_c^{\sharp} , and have

$$(f_c^{\sharp})^r : ((X_c^{\sharp})^r, (D^{X_c^{\sharp}})^r) \to Y^{\sharp}.$$

By [WW18, Lemma 4.2], we have that $(f_c^{\sharp})^r$ is also a stable family. Then by the flat base change, we have

$$(f_{c*}^{\sharp}\omega_{X_c^{\sharp}/Y^{\sharp}}^{m}(mD^{X_c^{\sharp}}))^r \otimes (\mathcal{A}^{\sharp}(D^{Y^{\sharp}}))^{-m} \simeq (f_c^{\sharp})_*^r \omega_{(X_c^{\sharp})^r/Y^{\sharp}}^{m}(m(D^{X_c^{\sharp}})^r) \otimes (\mathcal{A}^{\sharp}(D^{Y^{\sharp}}))^{-m}$$

Therefore,

$$H^{0}(Y^{\sharp}, (f_{c}^{\sharp})_{*}^{r} \omega_{(X_{c}^{\sharp})^{r}/Y^{\sharp}}^{m} (m(D^{X_{c}^{\sharp}})^{r}) \otimes (\mathcal{A}^{\sharp}(D^{Y^{\sharp}}))^{-m}) \neq 0.$$

Note that, by the construction of (4.2.1), $D^{Y^{\sharp}}$ does not depend on r.

Now we apply the fiber product trick on the log smooth morphism f^{\sharp} in (4.2.1) and obtain

$$(f^{\sharp})^r : ((X^{\sharp})^r, (D^{X^{\sharp}})^r) \to Y^{\sharp}.$$

We then replace $(X^{\sharp})^r$ by a log resolution of the main component of the normalization of $(X^{\sharp})^r$ and $(f^{\sharp})^r$ by the induced morphism. Note that these operations do not change $(f^{\sharp})^r$ over $Y^{\sharp} \setminus D^{Y^{\sharp}}$. Hence, we replace $(D^{X^{\sharp}})^r$ by the closure of $(D^{X^{\sharp}})^r|_{((f^{\sharp})^r)^{-1}(Y^{\sharp}\setminus D^{\sharp})}$ inside the new $(X^{\sharp})^r$. We can assume that $(D^{X^{\sharp}})^r$ is normal crossing by taking further log resolutions. We then add $((f^{\sharp})^r)^{-1}(D^{Y^{\sharp}})$ to $(D^{X^{\sharp}})^r$ and obtain a morphism of log smooth pairs

$$(f^{\sharp})^r : ((X^{\sharp})^r, (D^{X^{\sharp}})^r) \to (Y^{\sharp}, D^{Y^{\sharp}}).$$

By Lemma 3.1.4, $(f^{\sharp})^r$ is log smooth after getting rid of a small closed subset of Y^{\sharp} .

In the diagram (4.2.1), we replace f^{\sharp} by the new induced morphism $(f^{\sharp})^r : (X^{\sharp}, D^{X^{\sharp}}) \to (Y, D^{Y^{\sharp}})$, and replace f_c^{\sharp} by $(f_c^{\sharp})^r$. Then, by a local computation, we have that

$$f^{\sharp}_{*}\omega^{m}_{f^{\sharp}} \otimes \mathcal{A}^{\sharp-m} \supset f^{\sharp}_{c*}\omega^{m}_{X^{\sharp}_{c}/Y^{\sharp}}(mD^{X^{\sharp}_{c}}) \otimes (\mathcal{A}^{\sharp}(D^{Y^{\sharp}}))^{-m}$$

In particular, $f_*^{\sharp} \omega_{f^{\sharp}}^m \otimes \mathcal{A}^{\sharp-m}$ has non-trivial global sections.

We similarly apply the fiber product trick on all vertical morphisms in (4.2.1), and then, replace all $(X^{\bullet}, D^{X^{\bullet}})$ by $(X_{Y^{\bullet}}^{\bullet \times r}, D_{Y^{\bullet}}^{X^{\bullet} \times r})$, as we did in the $\bullet = \sharp$ case. We end up with a new commutative diagram as (4.2.1), satisfying all properties in Proposition 4.2.2. Hence, we have that for any sufficient positive and divisible m,

$$(f_{0*}'\psi_0^*(\omega_f^m \otimes f^*\mathcal{A}^{-m})) \simeq (f_{1*}'\omega_{f_1'}^m \otimes \mathcal{A}'^{-m}) \simeq (f_{2*}'\omega_{f_2'}^m \otimes \mathcal{A}'^{-m}) \simeq \eta^*(f_*^{\sharp}\omega_{f^{\sharp}}^m \otimes \mathcal{A}^{\sharp-m})$$

and they all have non-trivial global sections.

20

Since ψ_0 is finite, we can find l > 0, such that $(\omega_f \otimes f^* \mathcal{A}^{-1})^{ml}$, has non-trivial global sections, thanks to Lemma 5.1.4. From now on, we replace ml by m, and denote $\mathcal{A}'_1 := \mathcal{A}'_2 := \mathcal{A}'$ on Y', hence we can find four sections

$$\bar{s}^{\bullet} \in H^0(X^{\bullet}, (\omega_f \cdot \otimes f^{\bullet *}(\mathcal{A}^{\bullet})^{-1})^m),$$

with $\bullet =_{1,2}', \sharp$ or \emptyset , and satisfying that, on $f_1'^{-1}(Y' \setminus D^{Y'})$ and $f_2'^{-1}(Y' \setminus D^{Y'})$, $\bar{\psi}^*\bar{s} = \bar{s}_1'$ and $\bar{s}_2' = \bar{\eta}^*\bar{s}^{\sharp}$, respectively.

Moreover, since all coefficients of $D_h^{X^{\bullet}}$ are in (0,1), we can find four sections

(4.3.2)
$$s^{\bullet} \in H^0(X^{\bullet}, (\omega_{\lceil f^{\bullet} \rceil} \otimes f^{\bullet*}(\mathcal{A}^{\bullet})^{-1})^{nm}),$$

for some integral n > 0, with $\bullet =_{1,2}', \sharp$ or \emptyset , and satisfying that, on $f_1'^{-1}(Y' \setminus D^{Y'})$ and $f_2'^{-1}(Y' \setminus D^{Y'}), \bar{\psi}^* s = s_1'$ and $s_2' = \bar{\eta}^* s^{\sharp}$, and the support of the divisor $(s^{\bullet} = 0)$ contains the support of $D_h^{X^{\bullet}}$ respectively, by the definition of $\omega_{\lceil f^{\bullet} \rceil}$.

We now take the *nm*-th cyclic coverings given by s^{\bullet} . Then we have generic finite morphisms $\pi^{\bullet} : (Z^{\bullet}, D^{Z^{\bullet}}) \to (X^{\bullet}, D^{X^{\bullet}})$, with $\bullet = \emptyset'_{,1}, {}'_{,2}, {}^{\sharp}$, where Z^{\bullet} is a log resolution of the main component of the normalization of the cyclic covering given by s^{\bullet} and $D^{Z^{\bullet}}$ is given by $\pi^{\bullet-1}D^{X^{\bullet}}$. They induce the following commutative diagram:

(4.3.3)

where Z' is a common log resolution of Z'_1 and Z'_2 , and $D^{Z'}$ is given by the union of the inverse image of $D^{Z'_1}$ and $D^{Z'_2}$ (so $D^{Z'}$ is a reduced divisor). We also have \Box_1 and \Box_2 are Cartesian over $Y \setminus D^Y$ and $Y^{\sharp} \setminus D^{Y^{\sharp}}$. Furthermore, we have sections

$$\theta^{\bullet} \in H^0(\pi^{\bullet*}\omega_{\lceil f^{\bullet}\rceil} \otimes g_0^{\bullet*}(\mathcal{A}^{\bullet})^{-1})$$

on $(Z^{\bullet}, D^{Z^{\bullet}})$, with $\bullet = \emptyset_{1,2}'$ and \sharp , such that their zero locus contains $D_{h}^{Z^{\bullet}}$.

4.4. Extra boundaries and summary of the geometric construction. We make the final modification of the diagram (4.3.3), by adding more boundaries and summarize the geometric construction.

We continue assuming that we are in the situation of the diagram (4.3.3). Suppose that $f_V : (U, D^U) \to V$ is the log smooth family from where Proposition 4.1.1 starts. Then the morphism $f_V^r : (U^r, (D_U)^r) \to V$ remains log smooth. Then by the construction of (4.3.3), we know that

$$\begin{array}{ccc} U^r & \stackrel{\subseteq}{\longrightarrow} & X \\ & & \downarrow^{f_V^r} & & \downarrow^f \\ V & \stackrel{\subseteq}{\longrightarrow} & Y, \end{array}$$

where f is the first vertical morphism (forgetting log structure) in (4.2.1) in Proposition 4.2.2 after taking the fiber product tricks as in §4.3. We now set $E^Y := Y \setminus V$, E_h^X the closure of $(D^U)^r$ in X and $E_v^X := f^{-1}E^Y$ and $E^X = E_h^X + E_h^X$. Remeber

that V is important, since we are interested in the log Kodaira dimension of it. We then obtain the following commutative diagram:

(4.4.1)

$$(X, E^{X}) \xleftarrow{id} (X, D^{X}) \downarrow_{f_{0}} \downarrow_{f} \downarrow_{f} \downarrow_{g_{0}} \downarrow_{f} \downarrow_{f} \downarrow_{g} \downarrow_{g} \downarrow_{f} \downarrow_{f} \downarrow_{g} \downarrow_{g} \downarrow_{f} \downarrow_{f} \downarrow_{g} \downarrow_{g} \downarrow_{f} \downarrow_{g} \downarrow_{$$

where f_0 is the same as f as scheme-morphisms but with different log structures.

We now make the final modification of the diagram (4.3.3) as follows. We add extra boundary divisors on Y^{\bullet} so that all the down morphisms are log smooth over $Y^{\bullet} \setminus D^{Y^{\bullet}}$, and both ψ and η are morphisms of pairs, with ψ being finite with ramification locus contained in D^Y , and η being smooth. Then we add all vertical divisors onto those $D^{Z^{\bullet}}$, and make further resolutions if needed, to make all down arrows are morphisms of log smooth pairs, but keeping \Box_1 and \Box_2 are still Cartesian over $(Y \setminus D^Y)$ and $Y^{\sharp} \setminus D^{Y^{\sharp}}$. We then use g^{\bullet} to denote the new down morphisms respectively. After ignoring small closed subsets on those Y^{\bullet} respectively, we can have that all of those g^{\bullet} are log smooth, and $D^{Y^{\bullet}}$ are smooth divisors on Y^{\bullet} . In summary, we end up with the following commutative diagram:

$$(4.4.2)$$

$$(Z, D_v^Z) \longleftarrow (Z'_1, D_v^{Z'^1}) \longleftarrow (Z', D_v^{Z'}) \longrightarrow (Z'_2, D_v^{Z'_2}) \longrightarrow (Z^{\sharp}, D_v^{Z^{\sharp}})$$

$$\downarrow^g \Box_1 \qquad \qquad \downarrow^{g'} \downarrow^{g'} \qquad \qquad \Box_2 \qquad \downarrow^{g^{\sharp}} \qquad \qquad \Box_2 \qquad \downarrow^{g^{\sharp}} \qquad \qquad (Y, D^Y) \longleftarrow (Y', D^{Y'}) \xrightarrow{\eta} \qquad (Y^{\sharp}, D^{Y^{\sharp}}).$$

5. Construction of the refined Viehweg-Zuo sheaf

5.1. Construction of the Viehweg-Zuo sheaf. The goal of this section is the construction of the so-called Viehweg-Zuo sheaf in our setting, as indicated in the following theorem.

We assume that we are in the situation of the diagram (4.4.2) constructed in the previous section.

Theorem 5.1.1. There exists an invertible subsheaf $\mathcal{A} \subset \Omega_Y^{\otimes n}(\log E^Y)$, for some positive integer n, with $\kappa(\mathcal{A}) \geq \operatorname{Var}(f_V)$.

To prove the above theorem, we also use the terminology and notations established in §3.1 and §3.2. We apply Proposition 3.5.2 for g'_1 and g' in the diagram (4.4.2), and obtain graded $\mathscr{A}^{\bullet}_{(Y',D^{Y'})}$ -modules $\hat{\mathcal{M}}$ and $\mathcal{M}_{g'_1}$. We then prove the following lemma.

Lemma 5.1.2. We have the following sequence of morphisms, as graded $\mathscr{A}^{\bullet}_{(Y',D^{Y'})}$ -modules:

(5.1.1)
$$\psi^* \mathcal{M}_{f_0,\mathcal{L}} \xrightarrow{\simeq} \psi^* \mathcal{M}_{f,\mathcal{L}} \to \psi^* \mathcal{M}_{g,g^*\mathcal{A}} \to \mathcal{M}_{g'_1,g'_1^*\mathcal{A}'} \to \hat{\mathcal{M}}_{\mathcal{A}'},$$

away from a small closed subset of Y, where $\mathcal{L} = \omega_{\lceil f \rceil} = \omega_{\lceil f_0 \rceil}$, \mathcal{A}^{\bullet} are defined in (4.1.2), and $\hat{\mathcal{M}}_{\mathcal{A}'} := \hat{\mathcal{M}} \otimes_{\mathscr{O}_{Y'}} \mathcal{A}'^{-1}$. Moreover, the lowest graded pieces of those morphisms are inclusions. In particular, the composition of those morphisms is not trivial.

Proof. We construct (5.1.1) one by one from the left to the right. By (3.4.2), we obtain the first isomorphism. To obtain the second morphism, we need the construction in §3.6. More precisely, by (4.1.2) and the construction of Z in the §4.3, we see that (Assumption. 1) and (Assumption. 2) in §3.6 are fulfilled. Taking $\theta = s$ there in §3.6, we see that the second morphism is induced by η_s . We apply projection formula and see that

$$\mathcal{M}_{g,g^*\mathcal{A}} = \mathcal{M}_g \otimes_{\mathscr{O}} \mathcal{A} \text{ and } \mathcal{M}_{g'_1,g'^*\mathcal{A}'} = \mathcal{M}_{g'_1} \otimes_{\mathscr{O}} \mathcal{A}'.$$

We then obtain the third morphism by Proposition 3.4.2. By Proposition 3.5.2, we know that $\hat{\mathcal{M}}$ is a direct summand of $\mathcal{M}_{g'_1}$. Hence, the last morphism is defined to be the natural projection

$$\mathcal{M}_{g_1',g_1'^*\mathcal{A}'} \to \mathcal{M}_{\mathcal{A}'}.$$

The second morphism on the lowest graded pieces is injective by construction. The third morphism is injective. The last morphism on the lowest graded pieces is identical thanks to Proposition 3.5.2 again. Therefore, the lowest graded pieces of those morphisms are inclusions.

Denote by $\mathcal{N}_{g'}$, the image of $\psi^* \mathcal{M}_{f_0,\mathcal{L}}$ in $\hat{\mathcal{M}}_{\mathcal{A}'}$, under the morphisms in (5.1.1). Note that it is naturally a sub- $\mathscr{A}^{\bullet}_{(Y',DY')}$ -module of $\hat{\mathcal{M}}_{\mathcal{A}'}$. However, since $\mathcal{M}_{f_0,\mathcal{L}}$ is a $\mathscr{A}^{\bullet}_{(Y,EY)}$ -module, we have the following commutative diagram,

Consequently, the Higgs-type morphism $\mathcal{N}_{g'} \to \mathcal{N}_{g'} \otimes \Omega_{Y'}(\log D^{Y'})$ factorizes through

(5.1.2)
$$\delta: \mathcal{N}_{g'} \to \mathcal{N}_{g'} \otimes \psi^* \Omega_Y(\log E^Y)$$

We denote by $C_{g'}$ the kernel of δ and denoted by

$$\delta^k : \mathcal{N}_{q'} \to \mathcal{N}_{q'} \otimes \psi^* \operatorname{Sym}^k \Omega_Y(\log E^Y),$$

the k-th iterated morphism of δ .

Proposition 5.1.3. For any positive integer m, we can find another positive integer N_m , such that $\psi^* \mathcal{A}^{-m} \otimes \operatorname{Sym}^{N_m} \mathcal{C}_{a'}^{\vee}$ is generically globally generated.

Proof. We first focus on the morphism f^{\sharp} in (4.2.1) in Proposition 4.2.2 after taking the fiber product tricks as in §4.3. We apply Theorem 3.3.2 and get that $\mathcal{K}_{g^{\sharp}}^{\vee}$ is weakly positive. Since \mathcal{A}^{\sharp} is big, we hence have that

$$\mathcal{K}_{q^{\sharp},q^{\sharp*}\mathcal{A}^{\sharp}}^{ee}\simeq\mathcal{K}_{q^{\sharp}}^{ee}\otimes\mathcal{A}^{\sharp}$$

is also big. By the definition of bigness (see for instance [KP16, Definition 4.7(2)]), we then have that for any integer m, there exists an integer N_m , such that

$$\mathcal{A}^{\sharp-m} \otimes \operatorname{Sym}^{N_m} \mathcal{K}_{q^{\sharp},q^{\sharp*}\mathcal{A}^{\sharp}}^{\vee}$$

is generically globally generated. Since η is smooth, we have $\eta^* \mathcal{M}_{g^{\sharp}} \simeq \mathcal{M}_{g'_2}$ and hence $\eta^* \mathcal{K}_{g^{\sharp}} \simeq \mathcal{K}_{g'_2}$. Since $\eta^* \mathcal{A}^{\sharp} \simeq \mathcal{A}' \simeq \psi^* \mathcal{A}$ (see (4.1.2)), we hence have an isomorphism

$$\eta^*(\mathcal{A}^{\sharp-m}\otimes \operatorname{Sym}^{N_m}\mathcal{K}_{g^{\sharp},g^{\sharp*}\mathcal{A}^{\sharp}}^{\vee}) \to \psi^*\mathcal{A}^{-m}\otimes \operatorname{Sym}^{N_m}\mathcal{K}_{g'_2,g'_2*\mathcal{A}'}^{\vee}$$

Therefore, $\psi^* \mathcal{A}^{-m} \otimes \operatorname{Sym}^{N_m} \mathcal{K}^{\vee}_{g'_2, g'_2^* \mathcal{A}'}$ is also generically globally generated.

By applying Proposition 3.5.2 for g' and g'_2 , we see that $\hat{\mathcal{M}}_{\mathcal{A}'}$ is a direct summand of $\mathcal{M}_{g'_2,g'^{2*}\mathcal{A}'}$. We have a natural inclusion $\mathcal{C}_{g'} \to \mathcal{K}_{g'_2,g^*_2\mathcal{A}'}$, which induces a morphism

$$\psi^* \mathcal{A}^{-m} \otimes \operatorname{Sym}^{N_m} \mathcal{K}_{g'_2, g'^*_2 \mathcal{A}'}^{\vee} \to \psi^* \mathcal{A}^{-m} \otimes \operatorname{Sym}^{N_m} \mathcal{C}_{g'}^{\vee},$$

which is generically surjective. Hence we can conclude the proof.

The following technical lemma is needed in the proof of Theorem 5.1.1. It is well-know by experts, but we still give a proof here.

Lemma 5.1.4. Given a quasi-coherent sheaf \mathcal{F} on a variety X, and a finite covering $f: X' \to X$, assuming that $f^*\mathcal{F}$ has a non-trivial global section, then there exists a positive integer n, such that $H^0(X, \mathcal{F}^{\otimes n}) \neq 0$.

Proof. Let $\overline{f}: \overline{X} \to X$ be the Galois closure of f, with Galois group G of order n. Pick any non-trivial global section s in $H^0(\overline{X}, \overline{f^*}\mathcal{F})$. Then $\bigotimes_{g \in G} g \cdot s$ gives a G-invariant global section of $H^0(\overline{X}, \overline{f^*}\mathcal{F}^{\otimes n})$, and hence induces a global section of $\mathcal{F}^{\otimes n}$.

Proof of Theorem 5.1.1. By construction, the lowest graded piece of $\mathcal{M}_{f_0,\mathcal{L}}$ is $f_*\mathcal{O}_X$, which has a non-trivial global section (generating \mathcal{O}_Y), and denote by t its image in $\mathcal{N}_{g'}$. By the inclusions obtained in Lemma 5.1.2, we have that t is not a trivial section. Let k be the largest number, such that δ^k does not map t to 0. By Proposition 5.1.3, we have that $k \geq 1$. Hence δ^k maps t to $\mathcal{C}_{g'} \otimes \psi^* \operatorname{Sym}^k \Omega_Y(\log E^Y)$, which induces a non-trivial morphism

$$\mathcal{C}_{q'}^{\vee} \to \psi^* \operatorname{Sym}^k \Omega_Y(\log E^Y).$$

Since $\psi^* \mathcal{A}^{-1} \otimes \operatorname{Sym}^m \mathcal{C}_{g'}^{\vee}$ is generically globally generated, for some m, we can find a non-trivial morphism

$$\psi^* \mathcal{A} \to \psi^* \operatorname{Sym}^{mk} \Omega_Y(\log E^Y).$$

Since ψ is finite, we can find a positive integer n, such that we have a non-trivial morphism

$$\mathcal{A}^n \to \operatorname{Sym}^{nmk}\Omega_Y(\log E^Y)$$

thanks to Lemma 5.1.4. Since \mathscr{A} is a line bundle, the above non-trivial morphism is an inclusion. We conclude the proof, by replacing \mathcal{A} by \mathcal{A}^n .

5.2. Refinement of Viehweg-Zuo sheaves. We first denote

$$\mathcal{Q}_{(Y,D^Y)} := \ker(\mathcal{T}_Y(-\log D^Y) \to \mathcal{B}_{(Y,D^Y)}^{\vee}).$$

For a graded $\mathscr{A}^{\bullet}_{(Y,D^Y)}$ -module \mathcal{M} , to show that the Higgs-type morphism

$$\mathcal{M} \to \mathcal{M} \otimes \Omega_Y(\log D^Y)$$

factors through

$$\mathcal{M} \to \mathcal{M} \otimes \mathcal{B}_{(Y,D^Y)}$$

we only need to show that $\mathcal{Q}_{(Y,D^Y)}$ acts on \mathcal{M} trivially. If we further have \mathcal{M} being \mathcal{O}_Y -torsion-free, we only need to show that it acts trivially, over a dense open subset of Y.

Proof of Theorem 1.0.6. We first show that the higgs morphism (5.1.2)

$$\delta: \mathcal{N}_{q'} \to \mathcal{N}_{q'} \otimes \psi^* \Omega_Y(\log E^Y).$$

factors through

$$\delta_0: \mathcal{N}_{g'} \to \mathcal{N}_{g'} \otimes \psi^* \mathcal{B}_{(Y, E^Y)}.$$

Using the two stable families $f'_{2,c}$ and f^{\sharp}_{c} constructed in Proposition 4.1.1, over Y'and Y^{\sharp} , respectively, we have the induced moduli maps $\mu^{\bullet}: Y^{\bullet} \to \mathfrak{M}$, with $\bullet =', \sharp$. Hence, following Definition 1.0.5, we can similarly define $\mathcal{B}_{(Y',D^{Y'})}$ and $\mathcal{B}_{(Y^{\sharp},D^{Y^{\sharp}})}$, as well as $\mathcal{Q}_{(Y',D^{Y'})}$ and $\mathcal{Q}_{(Y^{\sharp},D^{Y^{\sharp}})}$ as above.

Note that, since ψ is finite, over a dense open subset of Y', we have $\psi^* \mathcal{Q}_{(Y,E^Y)} = \mathcal{Q}_{(Y',D^{Y'})}$. Hence, we only need to show that $\mathcal{Q}_{(Y',D^{Y'})}$ acts on $\mathcal{N}_{g'}$ trivially, over some dense open subset of Y'.

The isomorphism $\eta^* \mathcal{M}_{g^{\sharp}, g^{\sharp*} \mathcal{A}^{\sharp}} \simeq \mathcal{M}_{g'_2, g'^*_2 \mathcal{A}'}$ implies that the kernel of the natural morphism $d\eta^* : \mathcal{T}_{Y'}(-\log D^{Y'}) \to \eta^* \mathcal{T}_{Y^{\sharp}}(-\log D^{Y^{\sharp}})$ acts trivially on $\mathcal{M}_{g'_2, g'^*_2 \mathcal{A}'}$, hence on $\mathcal{N}_{g'}$. Since f^{\sharp} is of maximal variation, we have $\Omega_{Y^{\sharp}}(\log D^{Y^{\sharp}}) = \mathcal{B}_{(Y^{\sharp}, D^{Y^{\sharp}})}$, and that the inclusion $\eta^* \Omega_{Y^{\sharp}}(\log D^{Y^{\sharp}}) \subset \mathcal{B}_{(Y', D^{Y'})}$ actually is an isomorphism, over a dense open subset of Y'. By taking the dual, we have that, over a dense open subset, $\mathcal{Q}_{(Y', D^{Y'})}$ is isomorphic to the kernel of $d\eta^*$, and hence it acts trivially on $\mathcal{N}_{g'}$.

Furthermore, we have that $C_{g'}$, the kernel of δ , is also the kernel of δ_0 , since the induced morphism

$$\mathcal{N}_{q'} \otimes \psi^* \mathcal{B}_{(Y, E^Y)} \to \mathcal{N}_{q'} \otimes \psi^* \Omega_Y(\log E^Y)$$

is an inclusion, thanks to the fact that $\mathcal{N}_{g'} \otimes \psi^* \mathcal{B}_{(Y, E^Y)}$ is torsion-free, as ψ is flat. Now, we can apply the argument in the proof of Theorem 5.1.1, by replacing

 $\Omega_Y(\log E^Y)$ by $\mathcal{B}_{(Y,E^Y)}$, and δ by δ^0 , from which, the proof is accomplished. \Box

References

- [AK00] D. Abramovich and K. Karu. Weak semistable reduction in characteristic 0. Invent. Math., 139(2):241–273, 2000.
- [BCHM10] Caucher Birkar, Paolo Cascini, Christopher D. Hacon, and James McKernan. Existence of minimal models for varieties of log general type. J. Amer. Math. Soc., 23(2):405–468, 2010.
- [BPW17] Bo Berndtsson, Mihai Păun, and Xu Wang. Algebraic fiber spaces and curvature of higher direct images. 2017. arxiv.org/abs/1704.02279.
- [Cam07] Frédéric Campana. Orbifoldes spéciales et classification biméromorphe des variétés kählériennes compactes. arXiv preprint arXiv:0705.0737, 2007.
- [Cam11] Frédéric Campana. Orbifoldes géométriques spéciales et classification biméromorphe des variétés kählériennes compactes. J. Inst. Math. Jussieu, 10(4):809–934, 2011.
- [CP15] Frédéric Campana and Mihai Păun. Orbifold generic semi-positivity: an application to families of canonically polarized manifolds. Ann. Inst. Fourier (Grenoble), 65(2):835– 861, 2015.
- [FF12] Osamu Fujino and Taro Fujisawa. Variations of mixed hodge structure and semipositivity theorems. arXiv preprint arXiv:1203.6697, 2012.
- [HMX18] Christopher D. Hacon, James McKernan, and Chenyang Xu. Boundedness of moduli of varieties of general type. J. Eur. Math. Soc. (JEMS), 20(4):865–901, 2018.

- [JK11a] Kelly Jabbusch and Stefan Kebekus. Families over special base manifolds and a conjecture of Campana. Math. Z., 269(3-4):847–878, 2011.
- [JK11b] Kelly Jabbusch and Stefan Kebekus. Positive sheaves of differentials coming from coarse moduli spaces. Ann. Inst. Fourier (Grenoble), 61(6):2277–2290 (2012), 2011.
- [Kas86] Masaki Kashiwara. A study of variation of mixed hodge structure. Publications of the Research Institute for Mathematical Sciences, 22(5):991–1024, 1986.
- [Kaw85] Yujiro Kawamata. Minimal models and the Kodaira dimension of algebraic fiber spaces. J. Reine Angew. Math., 363:1–46, 1985.
- [Kaw14] Yujiro Kawamata. Hodge theory on generalized normal crossing varieties. Proceedings of the Edinburgh Mathematical Society, 57(1):175–189, 2014.
- [KK08] Stefan Kebekus and Sándor Kovács. Families of canonically polarized varieties over surfac. Invent. Math., 172(3):657–682, 2008. DOI:10.1016/j.aim.2008.01.005.
- [KM98] János Kollár and Shigefumi Mori. Birational geometry of algebraic varieties, volume 134 of Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 1998. With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original.
- [KP16] Sándor Kovács and Zsolt Patakfalvi. Projectivity of the moduli space of stable logvarieties and subadditivity of log-kodaira dimension. Journal of the American Mathematical Society, 2016.
- [LMB00] Gérard Laumon and Laurent Moret-Bailly. Champs algébraiques, volume 39 of Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]. Springer-Verlag, Berlin, 2000.
- [PS17] Mihnea Popa and Christian Schnell. Viehweg's hyperbolicity conjecture for families with maximal variation. *Invent. Math.*, 208(3):677–713, 2017.
- [PW16] Mihnea Popa and Lei Wu. Weak positivity for Hodge modules. Math. Res. Lett., 23(4):1139–1155, 2016.
- [RG71] Michel Raynaud and Laurent Gruson. Critères de platitude et de projectivité. Techniques de "platification" d'un module. Invent. Math., 13:1–89, 1971.
- [Sai88] Morihiko Saito. Modules de Hodge polarisables. Publ. Res. Inst. Math. Sci., 24(6):849– 995 (1989), 1988.
- [Sai90] Morihiko Saito. Mixed Hodge modules. Publ. Res. Inst. Math. Sci., 26(2):221–333, 1990.
- [Sch14] Christian Schnell. An overview of morihiko saito's theory of mixed hodge modules. arXiv preprint arXiv:1405.3096, 2014.
- [Taj16] Behrouz Taji. The isotriviality of smooth families of canonically polarized manifolds over a special quasi-projective base. *Compos. Math.*, 152(7):1421–1434, 2016.
- [Vie83] Eckart Viehweg. Weak positivity and the additivity of the Kodaira dimension for certain fibre spaces. In Algebraic varieties and analytic varieties (Tokyo, 1981), volume 1 of Adv. Stud. Pure Math., pages 329–353. North-Holland, Amsterdam, 1983.
- [Vie01] Eckart Viehweg. Positivity of direct image sheaves and applications to families of higher dimensional manifolds. In School on Vanishing Theorems and Effective Results in Algebraic Geometry (Trieste, 2000), volume 6 of ICTP Lect. Notes, pages 249–284. Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2001.
- [VZ02] Eckart Viehweg and Kang Zuo. Base spaces of non-isotrivial families of smooth minimal models. In Complex geometry (Göttingen, 2000), pages 279–328. Springer, Berlin, 2002.
- [VZ03] Eckart Viehweg and Kang Zuo. On the Brody hyperbolicity of moduli spaces for canonically polarized manifolds. Duke Math. J., 118(1):103–150, 2003.
- [Wei17] Chuanhao Wei. Logarithmic comparison theorems in mixed hodge modules. arXiv preprint arXiv:1710.07407, 2017.
- [WW18] Chuanhao Wei and Lei Wu. Hyperbolicity for log smooth families with maximal variation. 2018. preprint arXiv:1811.07466.

Department of Mathematics, Stony Brook University, 100 Nicolls Rd, Stony Brook, NY 11794, USA

E-mail address: chuanhao.wei@stonybrook.edu

Department of Mathematics, University of Utah, 155 S 1400 E, Salt Lake City, UT 84112, USA

E-mail address: lwu@math.utah.edu