

# On lower bound of relaxation time for massless fluid in presence of magnetic field

Sarthak Satapathy<sup>1,\*</sup>, Jayanta Dey<sup>1</sup>, Prashant Murmu<sup>2</sup> Sabyasachi Ghosh<sup>1\*</sup>

<sup>1</sup>*Indian Institute of Technology Bhilai, GEC Campus,  
Sejbahar, Raipur 492015, Chhattisgarh, India and*

<sup>2</sup>*Indian School of Mines Dhanbad 826004, Jharkhand, India*

Present work has gone through analytic calculation of shear viscosity and entropy density of massless relativistic fluid, facing external magnetic field  $B$  and then explored its fluid property.

Before going to finite magnetic field picture, let us remind the expression of shear viscosity for bosonic/fermionic fluid in absence of magnetic field [1],

$$\eta = \frac{\beta}{15} \int \frac{d^3 \mathbf{p}}{(2\pi)^3} \frac{\mathbf{p}^4}{\omega^2} \tau_c f(1 - af), \quad (1)$$

where  $f = 1/(e^{\beta\omega} + a)$ , with  $a = \pm 1$  for fermionic/bosonic fluid,  $\omega = \sqrt{\mathbf{p}^2 + m^2}$  is energy of massive constituent of medium, and most important quantity is the relaxation time  $\tau_c$ , which proportionally control the strength of shear viscosity.

Now, when we come the magnetic field picture, an additional time scale  $\tau_B = \omega/(eB)$  will come into the picture and in terms of these two time scale  $\tau_c$  and  $\tau_B$ , we will get an effective time scale

$$\tau_{\text{eff}} = \tau_c / \left\{ 1 + \left( \frac{\tau_c}{\tau_B} \right)^2 \right\} \quad (2)$$

which will be entered in place of  $\tau_c$  of Eq. (1) for finite magnetic field picture. Hence we will get shear viscosity of bosonic/fermionic fluid in presence of magnetic field [2] as

$$\eta_2 = \frac{\beta}{15} \int \frac{d^3 \mathbf{p}}{(2\pi)^3} \frac{\mathbf{p}^4}{\omega^2} \tau_{\text{eff}} f(1 - af), \quad (3)$$

where the subscript 2 stands for one of the component of shear viscosities among 5 components, appeared due to magnetic field.

For massless case, Eq. (1) becomes

$$\begin{aligned} \eta &= \frac{4 \tau_c}{5\pi^2} \zeta(4) T^4 = \frac{4\pi^2 \tau_c}{450} T^4 \quad \text{for Boson} \\ &= \left( \frac{7}{8} \right) \frac{4 \tau_c}{5\pi^2} \zeta(4) T^4 = \frac{7\pi^2 \tau_c}{900} T^4 \quad \text{for Fermion} \end{aligned} \quad (4)$$

and Eq. (3) becomes

$$\begin{aligned} \eta_2 &= \frac{\eta(B=0)}{1 + (\tau_c/\tau_B)^2} = \frac{\frac{4\pi^2 \tau_c}{450} T^4}{1 + (\tau_c/\tau_B)^2} \quad \text{for Boson} \\ &= \frac{\eta(B=0)}{1 + (\tau_c/\tau_B)^2} = \frac{\frac{7\pi^2 \tau_c}{900} T^4}{1 + (\tau_c/\tau_B)^2} \quad \text{for Fermion} . \end{aligned} \quad (5)$$

For calculation simplification, we have considered average energy in  $\tau_B$

$$\begin{aligned} \tau_B &= \left\{ \frac{\zeta(4)}{\zeta(3)} \right\} \frac{3T}{eB} \quad \text{for Boson} \\ &= \left\{ \frac{7\zeta(4)}{2\zeta(3)} \right\} \frac{3T}{eB} \quad \text{for Fermion} \end{aligned} \quad (6)$$

Now, let us come to the perfect fluid aspects of fermionic/bosonic fluid. Fluidity of the medium is measured by the shear viscosity to entropy density ratio  $\eta/s$ , where entropy density  $s$  of medium can be expressed as

$$s = \int \frac{d^3 \mathbf{p}}{(2\pi)^3} \left( \omega + \frac{\mathbf{p}^2}{3\omega} \right) f, \quad (7)$$

whose massless limit is

$$\begin{aligned} s &= \frac{4}{\pi^2} \zeta(4) T^3 = \frac{4\pi^2}{90} T^3 \quad \text{for Boson} \\ &= \left( \frac{7}{8} \right) \frac{4}{\pi^2} \zeta(4) T^3 = \frac{7\pi^2}{180} T^3 \quad \text{for Fermion} . \end{aligned} \quad (8)$$

\*Electronic address: sarthaks@iitbhlai.ac.in

In classical case, we can imagine a perfect fluid, having  $\eta/s = 0$  but in quantum case, we get a lower bound of  $\eta/s$ , which is also known as KSS bound [3], and it is  $1/(4\pi)$ . For massless medium, using  $\eta$  from Eq. (4) and  $s$  from Eq. (8), we can get  $\eta/s = \tau_c T/5$ , which is interestingly same for bosonic or fermionic fluid, although their individual  $\eta$  and  $s$  expressions are different. If at  $\tau_c = \tau_c^0$ , lower bound of  $\eta/s$  for massless medium is achieved, then we get

$$\begin{aligned} \frac{\eta}{s} &= \frac{\tau_c^0 T}{5} = \frac{1}{4\pi} \\ \Rightarrow \tau_c^0 &= \frac{5}{4\pi T}, \end{aligned} \quad (9)$$

which is quite standard well-known results. In this context, the present article provide similar type of analytic expression of relaxation time as a function  $T$  and  $B$  for massless fluid in presence of magnetic field. Using normal viscosity component  $\eta_2$  from Eq. (5) and  $s$  from Eq. (8), we will get  $\eta_2/s$ , which will be now different for Maxwell-Boltzmann (MB), Bose-Einstein (BE) and Fermi-Dirac (FD) distribution cases as it contains  $\tau_B$ , whose average values are different for different cases. By restricting  $\eta_2/s = 1/4\pi$ , we can get quadratic equation of  $\tau_c$ :

$$\begin{aligned} \tau_c^2 - 4\pi\tau_B^2 \frac{\eta(B=0)}{s} + \tau_B^2 &= 0 \\ \Rightarrow \tau_c^2 - \left(\frac{4\pi T \tau_B^2}{5}\right) \tau_c + \tau_B^2 &= 0 \\ \Rightarrow \tau_c^2 - \left(\frac{\tau_B^2}{\tau_c^0}\right) \tau_c + \tau_B^2 &= 0. \end{aligned} \quad (10)$$

The solution of above equation is

$$\tau_c = \tau_c^{2\pm} = \frac{\tau_B^2}{2\tau_c^0} \left[ 1 \pm \sqrt{1 - 4\left(\frac{\tau_c^0}{\tau_B}\right)^2} \right]. \quad (11)$$

So far from our best knowledge, we are first time addressing an *analytic* expressions of  $\tau_c(T, B)$  [2], where massless bosonic/fermionic matter in presence of magnetic field reach the KSS bound. To get a physical solution of Eq. (11), we need

$$\begin{aligned} 1 - 4\left(\frac{\tau_c^0}{\tau_B}\right)^2 &\geq 0 \\ \Rightarrow \tau_B &\geq 2\tau_c^0. \end{aligned} \quad (12)$$

Using MB relation  $\tau_B = \frac{3T}{eB}$  in above inequality, we have

$$\begin{aligned} \frac{3T}{eB} &\geq 2\frac{5}{4\pi T} \\ T &\geq \left(\frac{5eB}{6\pi}\right)^{1/2}. \end{aligned} \quad (13)$$

Corresponding FD and BE relations from Eq. (6) will give

$$\begin{aligned} T &\geq \left[\left(\frac{\zeta(3)}{\zeta(4)}\right)\frac{5eB}{6\pi}\right]^{1/2} \text{ for BE} \\ T &\geq \left[\left(\frac{2\zeta(3)}{7\zeta(4)}\right)\frac{5eB}{6\pi}\right]^{1/2} \text{ for FD.} \end{aligned} \quad (14)$$

Hence, Eqs. (13) and (14) can be identified as upper allowed domain, where KSS bound can be achieved while lower domain is forbidden zone if we believe that  $\eta_2/s$  never goes below  $\frac{1}{4\pi}$ . Within the allowed  $T$ - $B$  zone, we will get two positive values of relaxation time  $\tau_c^\pm$ , where KSS bound can be achieved. It can be understandable mathematically as follows. For smaller values of  $\tau_c$  or  $\tau_c/\tau_B \ll 1$ ,  $\eta_2/s \propto \tau_c$ , while for larger values of  $\tau_c$  or  $\tau_c/\tau_B \gg 1$ ,  $\eta_2/s \propto 1/\tau_c$ . Due to this two opposite trends of  $\tau_c$  dependence, we are getting a lower and upper values of  $\tau_c$ , where  $\eta_2/s$  reach the lower bound  $1/(4\pi)$ . The detail analysis of this investigation is documented in Ref. [2].

## References

- [1] S. Gavin, *Transport Coefficients In Ultrarelativistic Heavy Ion Collisions*, Nucl. Phys. **A 435**, 826 (1985).
- [2] J. Dey, S. Satapathy, P. Murmu, S. Ghosh, *Shear viscosity and electrical conductivity of relativistic fluid in presence of magnetic field: a massless case*, arXiv:1907.11164 [hep-ph]
- [3] P. Kovtun, D.T. Son, A.O. Starinets, *Viscosity in strongly interacting quantum field theories from black hole physics*, Phys. Rev. Lett. 94 (2005) 111601.