

A FLOW METHOD FOR THE DUAL ORLICZ-MINKOWSKI PROBLEM

YANNAN LIU JIAN LU

ABSTRACT. In this paper the dual Orlicz-Minkowski problem, a generalization of the L_p dual Minkowski problem, is studied. By studying a flow involving the Gauss curvature and support function, we obtain a new existence result of solutions to this problem for smooth measures.

1. INTRODUCTION

As important developments of the classical Brunn-Minkowski theory in convex geometry, the Orlicz-Brunn-Minkowski theory and the dual Orlicz-Brunn-Minkowski theory are developing rapidly in recent years. They have attracted great attention from many researchers, see for example [11, 20, 22, 27, 29, 30, 32, 33, 34, 36, 43, 45, 51, 53, 55, 56, 58, 60, 63, 66, 69] and references therein. In the recent groundbreaking work [33], Huang, Lutwak, Yang and Zhang introduced dual curvature measures and found their associated variational formulas for the first time. These new curvature measures were proved to connect two different known measures, namely cone-volume measure and Alexandrov integral curvature. Then the dual curvature measures were extended into the L_p case in [57]. The so called L_p dual curvature measures partially unified various measures in the L_p Brunn-Minkowski theory and the dual Brunn-Minkowski theory. Very recently, the L_p dual curvature measures were developed to the Orlicz case [21, 23], which unified more curvature measures and were named *dual Orlicz curvature measures*. These curvature measures are of central importance in the dual Orlicz-Brunn-Minkowski theory, and the corresponding Minkowski problems are called the *dual Orlicz-Minkowski problems*.

Let $\varphi : (0, +\infty) \rightarrow (0, +\infty)$ and $G : \mathbb{R}^n \setminus 0 \rightarrow (0, +\infty)$ be two given continuous functions. For a convex body $K \subset \mathbb{R}^n$ with the origin $0 \in K$, the dual Orlicz curvature measure of K is defined as

$$\tilde{C}_{\varphi, G}(K, E) := \frac{1}{n} \int_{\alpha_K^*(E)} \frac{\varphi(h_K(\alpha_K(u))) G(\rho_K(u)u) \rho_K^n(u)}{h_K(\alpha_K(u))} du$$

2010 *Mathematics Subject Classification.* 35J96, 52A20, 53A07, 53C44.

Key words and phrases. Monge-Ampère equation, dual Orlicz-Minkowski problem, Gauss curvature flow, Existence of solutions.

The authors were supported by Natural Science Foundation of China (11871432). The first author was also supported in part by Beijing Natural Science Foundation (1172005).

for each Borel set $E \subset S^{n-1}$, where α_K^* is the reverse radial Gauss image, α_K is the radial Gauss map, h_K is the support function and ρ_K is the radial function. See the next section for more details. For convenience, we here have used a slightly different notation in the definition of $\tilde{C}_{\varphi,G}(K, \cdot)$ from that of [21, 23]. The *dual Orlicz-Minkowski problem* asks what are the necessary and sufficient conditions for a Borel measure μ on the unit sphere S^{n-1} to be a multiple of the dual Orlicz curvature measure of a convex body K . Namely, this problem is to find a convex body $K \subset \mathbb{R}^n$ such that

$$(1) \quad c \, d\tilde{C}_{\varphi,G}(K, \cdot) = d\mu \text{ on } S^{n-1}$$

for some positive constant c .

When the Radon-Nikodym derivative of μ with respect to the spherical measure on S^{n-1} exists, namely $d\mu = \frac{1}{n}f \, dx$ for a non-negative integrable function f , the equation (1) is reduced into

$$(2) \quad c \, \varphi(h_K) G(\bar{\nabla} h_K) \det(\nabla^2 h_K + h_K I) = f \text{ on } S^{n-1},$$

where ∇ is the covariant derivative with respect to an orthonormal frame on S^{n-1} , I is the unit matrix of order $n-1$, and $\bar{\nabla} h_K(x) := \nabla h_K(x) + h_K(x)x$ is the point on ∂K whose outer unit normal vector is $x \in S^{n-1}$. This is a Monge-Ampère type equation. When both φ and G are constant functions, Eq. (1) or Eq. (2) is just the classical Minkowski problem.

When $\varphi(s) = s^{1-p}$ and $G(y) = |y|^{q-n}$, the dual Orlicz curvature measure $\tilde{C}_{\varphi,G}(K, \cdot)$ is reduced into the (p, q) -th dual curvature measure $\tilde{C}_{p,q}(K, \cdot)$ defined in [57]. Correspondingly, equation (1) or equation (2) will become into

$$(3) \quad d\tilde{C}_{p,q}(K, \cdot) = d\mu \text{ on } S^{n-1}$$

or

$$(4) \quad h_K^{1-p} |\bar{\nabla} h_K|^{q-n} \det(\nabla^2 h_K + h_K I) = f \text{ on } S^{n-1}$$

respectively. Here the constant c can be merged into h_K because of the homogeneity of $\tilde{C}_{p,q}$. Eq. (3) or Eq. (4) is just the L_p dual Minkowski problem. Many important Minkowski type problems are special cases of the L_p dual Minkowski problem. When $q = n$, Eq. (3) is the L_p Minkowski problem, which includes the logarithmic Minkowski problem ($p = 0$) and the centroaffine Minkowski problem ($p = -n$). The L_p Minkowski problem has been extensively studied, see e.g. [5, 7, 8, 15, 17, 28, 32, 36, 39, 40, 41, 47, 48, 49, 50, 52, 54, 67, 68] and Schneider's book [59], and corresponding references therein. When $p = 0$, Eq. (3) is the dual Minkowski problem, which was first introduced by [33] and then studied by [6, 14, 29, 31, 42, 64, 65]. Note that Eq. (3) also contains the Alexandrov problem [1] ($p = 0$ and $q = 0$), which is the prescribed Alexandrov integral curvature problem.

After the L_p dual Minkowski problem was first proposed in [57], several existence results of solutions were established in [4, 12, 35]. Huang and Zhao [35] proved the existence of solutions for general measure μ when $p > 0$ and $q < 0$, for even μ when

$pq > 0$ and $p \neq q$, and for smooth f when $p > q$. Chen, Huang and Zhao [12] studied the smooth case when f is even and $pq \geq 0$. Böröczky and Fodor [4] proved the existence of solutions for general measure μ when $p > 1$ and $q > 0$. Their existence results for the non-even case can be stated as the following two theorems. For a general measure μ , we have

Theorem A. (1) [35, Theorem 1.2]. *Let $p > 0$, $q < 0$, and μ be a non-zero finite Borel measure on \mathbb{S}^{n-1} . There exists a convex body $K \subset \mathbb{R}^n$ containing the origin in its interior such that Eq. (3) holds if and only if μ is not concentrated on any closed hemisphere.*

(2) [4, Theorem 1.2]. *Let $p > 1$, $q > 0$, and μ be a finite Borel measure on \mathbb{S}^{n-1} that is not concentrated on any closed hemisphere. Then there exists a convex body $K \subset \mathbb{R}^n$ containing the origin such that Eq. (3) holds when $p \neq q$ or holds up to a positive constant when $p = q$.*

And for a function f , there is

Theorem B ([35, Theorem 1.5]). *Suppose $p > q$. For any given positive function $f \in C^\alpha(\mathbb{S}^{n-1})$ with $\alpha \in (0, 1)$, there exists a unique solution $h_K \in C^{2,\alpha}(\mathbb{S}^{n-1})$ to Eq. (4). If f is smooth, then the solution is also smooth.*

In this paper we mainly consider the existence of solutions to the dual Orlicz-Minkowski problem. Our purpose is to extend Theorems A and B to Eq. (1) or Eq. (2). Very recently, Gardner et al. provided a sufficient condition for the existence of solutions to Eq. (1), which contained Theorem A (1) as a special case, see [21, Theorem 6.4]. In the sequel [23], they obtained another existence result about Eq. (1), which contained Theorem A (2) as a special case, see [23, Theorem 6.3]. To the best of our knowledge, there is no other existence result about Eq. (1) or Eq. (2).

In this paper we extend Theorem B to the dual Orlicz-Minkowski problem. Namely we obtain the following

Theorem 1. *Suppose $\alpha \in (0, 1)$, $\varphi \in C^\alpha(0, +\infty)$, and $G \in C^1(\mathbb{R}^n \setminus \{0\})$. For any given positive function $f \in C^\alpha(\mathbb{S}^{n-1})$ satisfying*

$$(5) \quad \limsup_{s \rightarrow +\infty} [\varphi(s) \max_{|y|=s} G(y) s^{n-1}] < f < \liminf_{s \rightarrow 0^+} [\varphi(s) \min_{|y|=s} G(y) s^{n-1}],$$

there exists a solution $h_K \in C^{2,\alpha}(\mathbb{S}^{n-1})$ to Eq. (2) with $c = 1$. If φ , G and f are smooth, then the solution is also smooth.

One can see that Theorem 1 applies to the situation when $\varphi(s) = s^{1-p}$, $G(y) = |y|^{q-n}$, $p > q$, and $f > 0$, which recovers Theorem B. In the dual Orlicz-Minkowski problem, the constant c in Eq. (1) or Eq. (2) is not known and can not be set to 1 in general, see [21, 23]. The equation of the Orlicz-Minkowski problem also contains a constant c which is not equal to 1 in general, see [26, 30, 38]. Note that in our Theorem 1, we can require $c = 1$ in Eq. (2). This is an advantage of the flow method used in our proof, compared to the usual variational method.

Theorem B was proved by the continuity method in [35], for which the uniqueness of solution to Eq. (4) is crucial. However solutions to the dual Orlicz-Minkowski problem may be not unique in general. One can easily construct examples when f is a constant function in Eq. (2). We may need to impose strong restrictions on φ and G to obtain the uniqueness, see our discussion Theorem 3 for $G(y) = G(|y|)$ and see also [23, Theorem 6.5]. Due to the lack of uniqueness of Eq. (2) in Theorem 1, the continuity method is no longer applicable to prove Theorem 1. Instead, we use a flow method involving the Gauss curvature and support function.

Let M_0 be a smooth, closed, uniformly convex hypersurface in Euclidean space \mathbb{R}^n enclosing the origin. Suppose that M_0 is given by a smooth embedding $X_0 : \mathbb{S}^{n-1} \rightarrow \mathbb{R}^n$. In this paper we study the long-time existence and convergence of a family of hypersurfaces $\{M_t\}$ given by smooth maps $X(\cdot, t) : \mathbb{S}^{n-1} \rightarrow \mathbb{R}^n$ satisfying the following flow equation:

$$(6) \quad \begin{aligned} \frac{\partial X}{\partial t}(x, t) &= -f(\nu)\mathcal{K} \frac{\langle X, \nu \rangle}{G(X)\varphi(\langle X, \nu \rangle)}\nu + X(x, t), \\ X(x, 0) &= X_0(x), \end{aligned}$$

where $\langle \cdot, \cdot \rangle$ is the standard inner product in \mathbb{R}^n , \mathcal{K} is the Gauss curvature of the hypersurface M_t at X , and ν is the unit outer normal vector of M_t at X .

The geometric flow generated by the Gauss curvature was first studied by Firey [18] to describe the shape of a tumbling stone. Since then, various Gauss curvature flows have been extensively studied, see e.g. [2, 3, 9, 10, 13, 16, 19, 24, 25, 37, 61, 62] and references therein. Our flow (6) is inspired by [46] in which a flow was used to study the dual Minkowski problem. However flow (6) is more complicated than that in [46], since it involves two more functions φ and G . Note G is a function of X other than $|X|$, which needs more efforts to deal with when estimating the principal curvatures of M_t .

In this paper, we obtain the long-time existence and convergence of the flow (6).

Theorem 2. *Let M_0 be a smooth closed uniformly convex hypersurface in Euclidean space \mathbb{R}^n enclosing the origin. Assume f, φ, G are smooth and satisfy the condition (5). Then flow (6) has a smooth solution M_t for all time $t > 0$, and a subsequence of M_t converges in C^∞ to a positive, smooth, uniformly convex solution to Eq. (2) with $c = 1$.*

This paper is organized as follows. In section 2, we give some basic knowledge about convex bodies and flow (6). In section 3, the long-time existence of flow (6) will be proved. The main difficulties in deriving long-time existence are to obtain uniform positive upper and lower bounds for support function and principal curvatures along the flow. By choosing proper auxiliary functions, the bounds of principal curvatures are obtained after delicate computations. Then long-time existence follows by standard arguments. In section 4, by considering a related geometric functional, which is decreasing along the flow, we can prove that the smooth solution of flow (6)

can converge for a subsequence to a smooth solution of Eq. (2) with $c = 1$. This completes the proof of Theorem 2. Then Theorem 1 follows by a simple approximation. At last, we provide a uniqueness result about (2) in a special case.

2. PRELIMINARIES

2.1. Convex body. Let \mathbb{R}^n be the n -dimensional Euclidean space, and \mathbb{S}^{n-1} be the unit sphere in \mathbb{R}^n . Assume M is a smooth closed uniformly convex hypersurface in \mathbb{R}^n . Without loss of generality, we may assume that M encloses the origin. The support function h of M is defined as

$$h(x) := \max_{y \in M} \langle y, x \rangle, \quad x \in \mathbb{S}^{n-1}.$$

And the radial function ρ of M is given by

$$\rho(u) := \max \{ \lambda > 0 : \lambda u \in M \}, \quad u \in \mathbb{S}^{n-1}.$$

We see that $\rho(u)u \in M$. Denote by ν_M the Gauss map of M . Then the radial Gauss map α_M is defined as

$$\alpha_M(u) := \nu_M(\rho(u)u) \text{ for } u \in \mathbb{S}^{n-1}.$$

And the reverse radial Gauss map α_M^* is given by

$$\alpha_M^*(x) := \{ u \in \mathbb{S}^{n-1} : \rho(u)u \in \nu_M^{-1}(x) \}.$$

For any Borel set $E \subset \mathbb{S}^{n-1}$ and any bounded integrable function $g : \mathbb{S}^{n-1} \rightarrow \mathbb{R}$, we have

$$(7) \quad \int_{\alpha_M^*(E)} g(u) du = \int_E g(\alpha_M^*(x)) h(x) \rho(\alpha_M^*(x))^{-n} \det(\nabla^2 h + hI)(x) dx.$$

Note that

$$\rho(u)u = (\bar{\nabla} h)(\alpha_M(u)), \quad u \in \mathbb{S}^{n-1}.$$

Then we easily see that equations (1) and (2) are equivalent. One can consult [33] for more details.

2.2. Curvature flow. Suppose that M is parametrized by the inverse Gauss map $X : \mathbb{S}^{n-1} \rightarrow M$, namely $X(x) = \nu_M^{-1}(x)$. The support function h of M can be computed by

$$(8) \quad h(x) = \langle x, X(x) \rangle, \quad x \in \mathbb{S}^{n-1}.$$

Here one can see x is the outer normal of M at $X(x)$. Let e_{ij} be the standard metric of the sphere \mathbb{S}^{n-1} and ∇ be the gradient on \mathbb{S}^{n-1} . Differentiating (8), we have

$$\nabla_i h = \langle \nabla_i x, X(x) \rangle + \langle x, \nabla_i X(x) \rangle.$$

Since $\nabla_i X(x)$ is tangent to M at $X(x)$, we have

$$\nabla_i h = \langle \nabla_i x, X(x) \rangle.$$

It follows that

$$(9) \quad X(x) = \nabla h + hx.$$

By differentiating (8) twice, the second fundamental form A_{ij} of M can be computed in terms of the support function, see for example [61],

$$(10) \quad A_{ij} = \nabla_{ij}h + he_{ij},$$

where $\nabla_{ij} = \nabla_i \nabla_j$ denotes the second order covariant derivative with respect to e_{ij} . The induced metric matrix g_{ij} of M can be derived by Weingarten's formula,

$$(11) \quad e_{ij} = \langle \nabla_i x, \nabla_j x \rangle = A_{ik} A_{lj} g^{kl}.$$

The principal radii of curvature are the eigenvalues of the matrix $b_{ij} = A^{ik} g_{jk}$. When considering a smooth local orthonormal frame on \mathbb{S}^{n-1} , by virtue of (10) and (11), we have

$$(12) \quad b_{ij} = A_{ij} = \nabla_{ij}h + h\delta_{ij}.$$

We will use b^{ij} to denote the inverse matrix of b_{ij} . The Gauss curvature of $X(x) \in M$ is given by

$$\mathcal{K}(x) = (\det(\nabla_{ij}h + h\delta_{ij}))^{-1}.$$

From the evolution equation of $X(x, t)$ in (6), we derive the evolution equation of the corresponding support function $h(x, t)$:

$$(13) \quad \frac{\partial h(x, t)}{\partial t} = -f(x)\mathcal{K}(x, t) \frac{h}{G(\nabla h + hx)} \frac{1}{\varphi(h)} + h(x, t), \quad x \in \mathbb{S}^{n-1}.$$

A self-similar solution of this flow is just equation (2).

Note that M can be represented by its radial function, namely

$$M = \{\rho(u)u : u \in \mathbb{S}^{n-1}\}.$$

It is easy to see that the outer normal vector $\nu = x(u)$ at $\rho(u)u \in M$ is given by

$$\nu = \frac{\rho(u)u - \nabla \rho}{\sqrt{\rho^2 + |\nabla \rho|^2}}.$$

And the induced metric and the second fundamental form of M can also be computed according to the radial function, see for example [24].

Denote the radial function of M_t by $\rho(u, t)$. From (9), u and x are related by

$$(14) \quad \rho(u)u = \nabla h(x) + h(x)x.$$

Let $x = x(u, t)$, by (14), we have

$$\log \rho(u, t) = \log h(x, t) - \log x \cdot u.$$

Differentiating the above identity, we have

$$\begin{aligned}
\frac{1}{\rho(u, t)} \frac{\partial \rho(u, t)}{\partial t} &= \frac{1}{h(x, t)} \left(\nabla h \cdot \frac{\partial x(u, t)}{\partial t} + \frac{\partial h(x, t)}{\partial t} \right) - \frac{1}{u \cdot x} u \cdot \frac{\partial x(u, t)}{\partial t} \\
&= \frac{1}{h(x, t)} \frac{\partial h(x, t)}{\partial t} + \frac{1}{h(x, t)} (\nabla h - \rho(u, t) u) \cdot \frac{\partial x(u, t)}{\partial t} \\
&= \frac{1}{h(x, t)} \frac{\partial h(x, t)}{\partial t}.
\end{aligned}$$

The evolution equation of radial function then follows from (13),

$$(15) \quad \frac{\partial \rho(u, t)}{\partial t} = -f \mathcal{K}(u, t) \frac{\rho}{G(\rho, u)} \frac{1}{\varphi(h)} + \rho(u, t),$$

where $\mathcal{K}(u, t)$ denotes the Gauss curvature at $\rho(u, t)u \in M_t$ and f takes value at the normal vector $x(u, t)$. Here and after, we sometimes also write $G(X) = G(\rho, u)$ for convenience.

3. LONG-TIME EXISTENCE OF THE FLOW

In this section, we will give a priori estimates about support function and obtain the long-time existence of the flow (6).

We begin with the estimate of the upper and lower bounds of h .

Lemma 1. *Let h be a smooth solution of (13) on $\mathbb{S}^{n-1} \times [0, T)$, and f, φ, G are smooth functions satisfying (5), then*

$$\frac{1}{C} \leq h(x, t) \leq C,$$

where C is a positive constant depending on $\max_{\mathbb{S}^{n-1}} h(x, 0)$, $\min_{\mathbb{S}^{n-1}} h(x, 0)$, $\max_{\mathbb{S}^{n-1}} f(x)$, $\min_{\mathbb{S}^{n-1}} f(x)$, $\limsup_{s \rightarrow +\infty} [\varphi(s) \max_{|y|=s} G(y) s^{n-1}]$ and $\liminf_{s \rightarrow 0^+} [\varphi(s) \max_{|y|=s} G(y) s^{n-1}]$.

Proof. Suppose $\max_{\mathbb{S}^{n-1}} h(x, t)$ is attained at $x^0 \in \mathbb{S}^{n-1}$, then at x^0 , we have

$$\nabla h = 0, \quad \nabla^2 h \leq 0, \quad \rho = h,$$

and

$$\nabla^2 h + hI \leq hI.$$

So at x^0 ,

$$\begin{aligned}
\frac{\partial h}{\partial t} &= -f(x) \mathcal{K}(x) \frac{h}{G(\rho, u)} \frac{1}{\varphi(h)} + h \\
&\leq -f(x) \frac{1}{G(h, u) h^{n-1}} \frac{h}{\varphi(h)} + h \\
&= \frac{1}{G(h, u) h^{n-2} \varphi(h)} (G(h, u) h^{n-1} \varphi(h) - f(x))
\end{aligned}$$

Let $\overline{A} = \limsup_{s \rightarrow +\infty} [\varphi(s) \max_{|y|=s} G(y) s^{n-1}]$. By (5), $\varepsilon = \frac{1}{2}(\min_{\mathbb{S}^{n-1}} f(x) - \overline{A})$ is positive and there exists a positive constant $C_1 > 0$ such that

$$G(h, u)h^{n-1}\varphi(h) < \overline{A} + \varepsilon$$

for $h > C_1$. It follows that

$$G(h, u)h^{n-1}\varphi(h) - f(x) < \overline{A} + \varepsilon - \min_{\mathbb{S}^{n-1}} f(x) < 0,$$

from which we have

$$\frac{\partial h}{\partial t} < 0$$

at maximal points. So

$$h \leq \max\{C_1, \max_{\mathbb{S}^{n-1}} h(x, 0)\}.$$

Similarly, we can estimate $\min_{\mathbb{S}^{n-1}} h(x, t)$. Suppose $\min_{\mathbb{S}^{n-1}} h(x, t)$ is attained at x^1 , then at this point,

$$\nabla h = 0, \quad \nabla^2 h \geq 0, \quad \rho = h,$$

and

$$\nabla^2 h + hI \geq hI.$$

And at x^1 ,

$$\begin{aligned} \frac{\partial h}{\partial t} &= -f(x)\mathcal{K}(x)\frac{h}{G(\rho, u)}\frac{1}{\varphi(h)} + h \\ &\geq -f(x)\frac{1}{G(h, u)h^{n-1}}\frac{h}{\varphi(h)} + h \\ &= \frac{1}{G(h, u)h^{n-2}\varphi(h)}(G(h, u)h^{n-1}\varphi(h) - f(x)). \end{aligned}$$

Let $\underline{A} = \liminf_{s \rightarrow 0^+} [\varphi(s) \max_{|y|=s} G(y) s^{n-1}]$. By (5), $\varepsilon = \frac{1}{2}(\underline{A} - \max_{\mathbb{S}^{n-1}} f(x))$ is positive and there exists some positive constant $C_2 > 0$ such that

$$G(h, u)h^{n-1}\varphi(h) > \underline{A} - \varepsilon$$

for $h < C_2$. It follows by (5)

$$G(h, u)h^{n-1}\varphi(h) - f(x) > \underline{A} - \varepsilon - \max_{\mathbb{S}^{n-1}} f(x) > 0,$$

which shows that

$$\frac{\partial h}{\partial t} > 0$$

at minimal points. Hence

$$h \geq \min\{C_2, \min_{\mathbb{S}^{n-1}} h(x, 0)\}.$$

The proof of the lemma is completed. □

Corollary 1. *Let h be a smooth solution of (13) on $\mathbb{S}^{n-1} \times [0, T)$, and f, φ, G are smooth functions satisfying (5). Then we have*

$$(16) \quad |\nabla h(x, t)| \leq C, \quad \forall (x, t) \in \mathbb{S}^{n-1} \times [0, T),$$

where C is the same positive constant as in Lemma 1.

Proof. By (14), we know that

$$\min_{\mathbb{S}^{n-1}} h(x, t) \leq \rho(u, t) \leq \max_{\mathbb{S}^{n-1}} h(x, t),$$

and

$$\rho^2 = h^2 + |\nabla h|^2.$$

These facts together with Lemma 1 imply the result. \square

Next we will establish the uniform upper and lower bounds for the principal curvatures of the flow (6). We first derive an upper bound for the Gauss curvature. For convenience, we write

$$F = f(x)\mathcal{K}(x)\frac{h}{G(\nabla h + hx)}\frac{1}{\varphi(h)}.$$

In this section, we take a local orthonormal frame $\{e_1, \dots, e_{n-1}\}$ on \mathbb{S}^{n-1} such that the standard metric on \mathbb{S}^{n-1} is $\{\delta_{ij}\}$. The local coordinate of a point x in \mathbb{S}^{n-1} is denoted by $x = (x_1, \dots, x_{n-1})$. Double indices always mean to sum from 1 to $n-1$.

By Lemma 1 and Corollary 1, if h is a smooth solution of (13) on $\mathbb{S}^{n-1} \times [0, T)$, and f, φ, G are smooth functions satisfying (5), then along the flow for $[0, T)$, $\nabla h + hx$ and h are smooth functions whose ranges are within some bounded domain $\Omega_{[0, T)}$ and bounded interval $I_{[0, T)}$ respectively. Here $\Omega_{[0, T)}$ and $I_{[0, T)}$ depend only on the upper and lower bounds of h on $[0, T)$.

Lemma 2. *Let h be a smooth solution of (13) on $\mathbb{S}^{n-1} \times [0, T)$, and f, φ, G are smooth functions satisfying (5). Then on $\mathbb{S}^{n-1} \times [0, T)$,*

$$\mathcal{K}(x, t) \leq C,$$

where C is a positive constant depending on $\|f\|_{C^0(\mathbb{S}^{n-1})}$, $\|\varphi\|_{C^1(I_{[0, T)})}$, $\|G\|_{C^1(\Omega_{[0, T)})}$ and $\|h\|_{C^1(\mathbb{S}^{n-1} \times [0, T))}$.

Proof. Consider the following auxiliary function

$$Q(x, t) = \frac{1}{h - \varepsilon_0}(F - h),$$

where ε_0 is a positive constant satisfying

$$\varepsilon_0 < \min_{\mathbb{S}^{n-1} \times [0, T)} h(x, t).$$

Recall that $F = \frac{f(x)\mathcal{K}(x)h}{G(\nabla h + hx)\varphi(h)}$, then by Lemma 1 and Corollary 1, the upper bound of $\mathcal{K}(x, t)$ follows from that of $Q(x, t)$. Hence we only need to derive the upper bound of $Q(x, t)$. First we compute the evolution equation of $Q(x, t)$. Note that

$$\begin{aligned}\nabla_i Q &= \frac{F_i - h_i}{h - \varepsilon_0} - \frac{F - h}{(h - \varepsilon_0)^2} h_i, \\ \nabla_{ij} Q &= \frac{F_{ij} - h_{ij}}{h - \varepsilon_0} - \frac{(F_i - h_i)h_j}{(h - \varepsilon_0)^2} - \frac{(F_j - h_j)h_i + (F - h)h_{ij}}{(h - \varepsilon_0)^2} + 2\frac{(F - h)h_i h_j}{(h - \varepsilon_0)^3} \\ &= \frac{F_{ij} - h_{ij}}{h - \varepsilon_0} - \frac{(F - h)h_{ij}}{(h - \varepsilon_0)^2} - \frac{\nabla_i Q h_j}{h - \varepsilon_0} - \frac{\nabla_j Q h_i}{h - \varepsilon_0},\end{aligned}$$

and

$$\frac{\partial Q}{\partial t} = \frac{F_t - h_t}{h - \varepsilon_0} + \frac{h_t^2}{(h - \varepsilon_0)^2} = \frac{F_t}{h - \varepsilon_0} + Q + Q^2.$$

The evolution equation of $Q(x, t)$ is given by

$$\begin{aligned}\frac{\partial Q}{\partial t} - Fb^{ij}\nabla_{ij}Q &= \frac{1}{h - \varepsilon_0}(F_t - Fb^{ij}F_{ij}) + Q + Q^2 \\ &\quad + (Q + 1)\frac{Fb^{ij}h_{ij}}{h - \varepsilon_0} + \frac{\nabla_i Q Fb^{ij}h_j}{h - \varepsilon_0} + \frac{\nabla_j Q Fb^{ij}h_i}{h - \varepsilon_0}.\end{aligned}$$

Now, we need to compute the evolution equation of F .

$$\begin{aligned}F_t &= \frac{hf(x)}{G(\nabla h + hx)\varphi(h)} \frac{\partial \mathcal{K}}{\partial t} + \mathcal{K}(x, t)f(x) \frac{\partial}{\partial t} \left(\frac{h}{G(\nabla h + hx)\varphi(h)} \right) \\ &= -Fb^{ij}(h_{ij} + \delta_{ij}h)_t + \mathcal{K}(x, t)f(x) \frac{\partial}{\partial t} \left(\frac{h}{G(\nabla h + hx)\varphi(h)} \right) \\ &= -Fb^{ij}(h_t)_{ij} - Fb^{ij}\delta_{ij}h_t + \mathcal{K}(x, t)f(x) \frac{\partial}{\partial t} \left(\frac{h}{G(\nabla h + hx)\varphi(h)} \right) \\ &= -Fb^{ij}(-F + h)_{ij} - Fb^{ij}\delta_{ij}h_t + \mathcal{K}(x, t)f(x) \frac{\partial}{\partial t} \left(\frac{h}{G(\nabla h + hx)\varphi(h)} \right) \\ &= Fb^{ij}F_{ij} - Fb^{ij}b_{ij} + F^2b^{ij}\delta_{ij} + \mathcal{K}(x, t)f(x) \frac{\partial}{\partial t} \left(\frac{h}{G(\nabla h + hx)\varphi(h)} \right),\end{aligned}$$

where we have used the fact

$$(17) \quad \frac{\partial \mathcal{K}}{\partial b_{ij}} = -\mathcal{K}b^{ij}.$$

Therefore we have

$$\frac{\partial F}{\partial t} - Fb^{ij}\nabla_{ij}F = -F(n - 1) + F^2b^{ij}\delta_{ij} + \mathcal{K}(x, t)f(x) \frac{\partial}{\partial t} \left(\frac{h}{G(\nabla h + hx)\varphi(h)} \right).$$

At a spatial maximal point of $Q(x, t)$, if we take an orthonormal frame such that b_{ij} is diagonal, we have

$$\begin{aligned}
& \frac{\partial Q}{\partial t} - b^{ii} F \nabla_{ii} Q \\
& \leq \frac{1}{h - \varepsilon_0} (F_t - b^{ii} F \nabla_{ii} F) + Q + Q^2 + \frac{F b^{ii} h_{ii}}{h - \varepsilon_0} + \frac{(F - h) F b^{ii} h_{ii}}{(h - \varepsilon_0)^2} \\
& = \frac{1}{h - \varepsilon_0} (-F b^{ii} b_{ii} + F^2 b^{ii} \delta_{ii} + \mathcal{K}(x, t) f(x) \frac{\partial}{\partial t} (\frac{h}{G(\nabla h + hx) \varphi(h)})) + Q + Q^2 \\
& \quad + \frac{F b^{ii} (b_{ii} - h \delta_{ii})}{h - \varepsilon_0} + \frac{Q F b^{ii} (b_{ii} - h \delta_{ii})}{h - \varepsilon_0} \\
& = \frac{F^2}{h - \varepsilon_0} \sum_i b^{ii} + Q + Q^2 + \frac{1}{h - \varepsilon_0} \mathcal{K}(x, t) f(x) \frac{\partial}{\partial t} (\frac{h}{G(\nabla h + hx) \varphi(h)}) \\
& \quad - \frac{h F}{h - \varepsilon_0} \sum_i b^{ii} + \frac{Q F (n - 1)}{h - \varepsilon_0} - \frac{Q F h}{h - \varepsilon_0} \sum_i b^{ii} \\
& \leq F Q (1 - \frac{h}{h - \varepsilon_0}) \sum_i b^{ii} + C_1 Q + C_2 Q^2 \\
& \quad + \frac{1}{h - \varepsilon_0} \mathcal{K}(x, t) f(x) \frac{\partial}{\partial t} (\frac{h}{G(\nabla h + hx) \varphi(h)}).
\end{aligned}$$

From the evolution equation of (6), we have

$$\begin{aligned}
\frac{\partial G}{\partial t} &= \langle \bar{\nabla} G, \partial_t X \rangle \\
&= \langle \bar{\nabla} G, -F \nu + X \rangle \\
&= -F \langle \bar{\nabla} G, \nu \rangle + \langle \bar{\nabla} G, X \rangle,
\end{aligned}$$

where $\bar{\nabla}$ is the standard metric on \mathbb{R}^n .

Now, we have

$$\begin{aligned}
\frac{\partial}{\partial t} (\frac{h}{G(\nabla h + hx) \varphi(h)}) &= \frac{1}{G \varphi} h_t - \frac{h}{G^2 \varphi^2} (\varphi \frac{\partial G}{\partial t} + G \frac{\partial \varphi}{\partial t}) \\
&= \frac{1}{G \varphi} h_t - \frac{h \varphi'}{G \varphi^2} h_t - \frac{h}{G^2 \varphi} \frac{\partial G}{\partial t} \\
&\leq c Q (h - \varepsilon_0) + c,
\end{aligned}$$

where φ' denotes $\frac{\partial \varphi(s)}{\partial s}$, and c is a positive constant depending on $\|\varphi\|_{C^1(I_{[0, T]})}$, $\|G\|_{C^1(\Omega_{[0, T]})}$ and $\|h\|_{C^1(\mathbb{S}^{n-1} \times [0, T])}$.

For a Q large enough, by Lemma 1 and Corollary 1, it is easy to see

$$\frac{1}{C_0} \mathcal{K} \leq Q \leq C_0 \mathcal{K},$$

and

$$\sum_i b^{ii} \geq (n-1)\mathcal{K}(x)^{\frac{1}{n-1}}.$$

Hence if we take ε_0 as a positive constant satisfying $\varepsilon_0 < \min_{S^{n-1} \times (0, +\infty)} h(x, t)$, then for Q large,

$$\frac{\partial Q}{\partial t} \leq C_1 Q^2 (C_2 - \varepsilon_0 Q^{\frac{1}{n-1}}) < 0.$$

Now the upper bound of $\mathcal{K}(x, t)$ follows. \square

Next we will estimate the lower bound of principal curvatures. It is equivalent to estimate the upper bound of the eigenvalues of matrix $\{b_{ij}\}$ w.r.t. $\{\delta_{ij}\}$.

Lemma 3. *Let h be a smooth solution of (13) on $\mathbb{S}^{n-1} \times [0, T)$, and f, φ, G are smooth functions satisfying (5), then the principal curvature κ_i for $i = 1, \dots, n-1$ satisfies*

$$\kappa_i \geq C,$$

where C is a positive constant depending on $\|f\|_{C^2(\mathbb{S}^{n-1})}$, $\|\varphi\|_{C^2(I_{[0, T)})}$, $\|G\|_{C^2(\Omega_{[0, T)})}$ and $\|h\|_{C^1(\mathbb{S}^{n-1} \times [0, T))}$.

Proof. Consider the auxiliary function

$$w(x, t) = \log \lambda_{\max}(b_{ij}) - A \log h + B |\nabla h|^2,$$

where A, B are constants to be determined. λ_{\max} is the maximal eigenvalue of b_{ij} .

For any fixed t , we assume that $\max_{\mathbb{S}^{n-1}} w(x, t)$ is attained at $p \in \mathbb{S}^{n-1}$. At p , we take an orthogonal frame such that $b_{ij}(p, t)$ is diagonal and $\lambda_{\max}(p, t) = b_{11}(p, t)$.

Now we can write $w(x, t)$ as

$$w(x, t) = \log b_{11} - A \log h + B |\nabla h|^2.$$

First, we compute the evolution equation of w . Note that

$$\begin{aligned} \frac{\partial \log b_{11}}{\partial t} - b^{ii} F \nabla_{ii} \log b_{11} &= b^{11} \left(\frac{\partial b_{11}}{\partial t} - F b^{ii} \nabla_{ii} b_{11} \right) + F b^{ii} (b^{11})^2 (\nabla_i b_{11})^2, \\ \frac{\partial \log h}{\partial t} - b^{ii} F \nabla_{ii} \log h &= \frac{1}{h} \left(\frac{\partial h}{\partial t} - F b^{ii} \nabla_{ii} h \right) + \frac{F b^{ii} h_i^2}{h^2}, \\ \frac{\partial |\nabla h|^2}{\partial t} - b^{ii} F \nabla_{ii} |\nabla h|^2 &= 2 h_k \left(\frac{\partial h_k}{\partial t} - F b^{ii} \nabla_{ii} h_k \right) - 2 F b^{ii} h_{ii}^2, \end{aligned}$$

we have

$$\begin{aligned} \frac{\partial w}{\partial t} - F b^{ii} \nabla_{ii} w &= b^{11} \left(\frac{\partial b_{11}}{\partial t} - F b^{ii} \nabla_{ii} b_{11} \right) + F b^{ii} (b^{11})^2 (\nabla_i b_{11})^2 \\ &\quad - \frac{A}{h} \left(\frac{\partial h}{\partial t} - F b^{ii} \nabla_{ii} h \right) - \frac{F b^{ii} h_i^2}{h^2} \\ &\quad + 2 B h_k \left(\frac{\partial h_k}{\partial t} - F b^{ii} \nabla_{ii} h_k \right) - 2 B F b^{ii} h_{ii}^2. \end{aligned}$$

Let

$$M = \log \frac{hf(x)}{G(\nabla h + hx)\varphi(h)},$$

then

$$\log F = \log \mathcal{K} + M.$$

Differentiating the above equation, by (17) we have

$$\frac{\nabla_k F}{F} = \frac{1}{\mathcal{K}} \frac{\partial \mathcal{K}}{\partial b_{ij}} \nabla_k b_{ij} + \nabla_k M = -b^{ij} \nabla_k b_{ij} + M_k,$$

and

$$\frac{\nabla_{kl} F}{F} - \frac{\nabla_k F \nabla_l F}{F^2} = -b^{ij} \nabla_{kl} b_{ij} + b^{ii} b^{jj} \nabla_k b_{ij} \nabla_l b_{ij} + \nabla_{lk} M.$$

From the above results and the evolution equation of h , we have

$$\begin{aligned} \frac{\partial h}{\partial t} - F b^{ii} \nabla_{ii} h &= -F + h - F b^{ii} (b_{ii} - \delta_{ii} h) \\ &= -F n + h + F h \sum_i b^{ii}, \end{aligned}$$

and

$$\begin{aligned} \frac{\partial h_k}{\partial t} - F b^{ii} \nabla_{ii} h_k &= -F_k + h_k - F b^{ii} \nabla_k b_{ii} + F h_k \sum_i b^{ii} \\ &= -M_k F + h_k + F h_k \sum_i b^{ii}. \end{aligned}$$

On the other hand, from the evolution equation of h , we have

$$\begin{aligned} \frac{\partial h_{kl}}{\partial t} &= -\nabla_{kl} F + h_{kl} \\ &= -\frac{\nabla_k F \nabla_l F}{F} + F b^{ij} \nabla_{kl} b_{ij} - F b^{ii} b^{jj} \nabla_k b_{ij} \nabla_l b_{ij} - F \nabla_{lk} M + h_{kl}. \end{aligned}$$

By the Gauss equation, see [61] for details,

$$\nabla_{kl} h_{ij} = \nabla_{ij} h_{kl} + 2\delta_{kl} h_{ij} - 2\delta_{ij} h_{kl} + \delta_{kj} h_{il} - \delta_{li} h_{kj},$$

or

$$\nabla_{kl} b_{ij} = \nabla_{ij} b_{kl} + \delta_{kl} h_{ij} - \delta_{ij} h_{kl} + \delta_{kj} h_{il} - \delta_{li} h_{kj}.$$

Then

$$\begin{aligned} \frac{\partial h_{kl}}{\partial t} &= F b^{ij} \nabla_{ij} h_{kl} + 2\delta_{kl} F b^{ij} h_{ij} - F b^{ij} \delta_{ij} h_{kl} + F b^{ik} h_{il} - F b^{jl} h_{kj} \\ &\quad - \frac{\nabla_k F \nabla_l F}{F} - F b^{ii} b^{jj} \nabla_k b_{ij} \nabla_l b_{ij} - F \nabla_{lk} M + h_{kl}. \end{aligned}$$

Hence

$$\begin{aligned}
\frac{\partial b_{kl}}{\partial t} &= Fb^{ij}\nabla_{ij}b_{kl} + \delta_{kl}Fb^{ij}(b_{ij} - h\delta_{ij}) - Fb^{ij}\delta_{ij}(b_{kl} - h\delta_{kl}) \\
&\quad + Fb^{ik}h_{il} - Fb^{jl}h_{kj} + b_{kl} - h\delta_{kl} + (-F + h)\delta_{kl} \\
&\quad - \frac{\nabla_k F \nabla_l F}{F} - Fb^{ii}b^{jj}\nabla_k b_{ij}\nabla_l b_{ij} - F\nabla_{lk}M \\
&= Fb^{ij}\nabla_{ij}b_{kl} + \delta_{kl}F(n-2) - Fb^{ij}\delta_{ij}b_{kl} \\
&\quad + Fb^{ik}h_{il} - Fb^{jl}h_{kj} + b_{kl} \\
&\quad - \frac{\nabla_k F \nabla_l F}{F} - Fb^{ii}b^{jj}\nabla_k b_{ij}\nabla_l b_{ij} - F\nabla_{lk}M.
\end{aligned}$$

When $k = l = 1$, we have

$$\begin{aligned}
\frac{\partial b_{11}}{\partial t} &= Fb^{ii}\nabla_{ii}(b_{11}) + F(n-2) - F\sum_i b^{ii}b_{11} + b_{11} \\
&\quad - \frac{F_1^2}{F} - Fb^{ii}b^{jj}\nabla_1(b_{ij})^2 - FM_{11}.
\end{aligned}$$

By the above results, we get

$$\begin{aligned}
&\frac{\partial w}{\partial t} - Fb^{ii}\nabla_{ii}w \\
&\leq b^{11}F(n-2) + 1 - A - b^{11}FM_{11} + \frac{AFn}{h} - AF\sum_i b^{ii} \\
&\quad - 2BFh_kM_k + 2B|\nabla h|^2 + 2BF|\nabla h|^2\sum_i b^{ii} - 2BF\sum_i b_{ii} + 4BFh(n-1) \\
&= b^{11}F(n-2) - b^{11}FM_{11} - 2BFh_kM_k - (A - 2B|\nabla h|^2)F\sum_i b^{ii} \\
&\quad - (A - 1 - 2B|\nabla h|^2) - 2BF\sum_i b_{ii} + 4BFh(n-1) + \frac{AFn}{h}.
\end{aligned}$$

If we let A satisfy

$$A \geq 2B \max_{\mathbb{S}^{n-1} \times [0, T)} |\nabla h|^2 + 1,$$

then

$$\begin{aligned}
(18) \quad \frac{\partial w}{\partial t} - Fb^{ii}\nabla_{ii}w &\leq b^{11}F(n-2) - b^{11}FM_{11} - 2BFh_kM_k \\
&\quad - 2BF\sum_i b_{ii} + 4BFh(n-1) + \frac{AFn}{h}.
\end{aligned}$$

Now, we estimate $-b^{11}FM_{11} - 2BFh_kM_k$. Since

$$\begin{aligned} M &= \log \frac{hf(x)}{G(\nabla h + hx)\varphi(h)} \\ &= \log f(x) + \log h(x) - \log G(\nabla h + hx) - \log \varphi(h), \end{aligned}$$

then

$$\begin{aligned} \nabla_k M &= \frac{f_k}{f} + \frac{h_k}{h} - \frac{1}{G} \langle \bar{\nabla} G, \nabla_k X \rangle - \frac{\varphi'}{\varphi} h_k \\ &= \frac{f_k}{f} + \frac{h_k}{h} - \frac{\varphi'}{\varphi} h_k - \frac{1}{G} \langle \bar{\nabla} G, e_i \rangle b_{ki}. \end{aligned}$$

And

$$\begin{aligned} \nabla_{11} M &= \frac{f_{11}}{f} - \frac{f_1^2}{f^2} + \frac{h_{11}}{h} - \frac{h_1^2}{h^2} - \frac{\varphi'' h_1^2 + \varphi' h_{11}}{\varphi} + \frac{(\varphi')^2 h_1^2}{\varphi^2} \\ &\quad + \frac{1}{G^2} \langle \bar{\nabla} G, e_i \rangle \langle \bar{\nabla} G, e_j \rangle b_{1i} b_{1j} - \frac{1}{G} \langle \nabla_1 \bar{\nabla} G, e_i \rangle b_{1i} \\ &\quad + \frac{1}{G} \langle \bar{\nabla} G, x \rangle b_{11} - \frac{1}{G} \langle \bar{\nabla} G, e_i \rangle b_{11i}, \end{aligned}$$

where $\varphi'' = \frac{\partial^2 \varphi(h)}{\partial h^2}$ and we have used the Gauss formula on \mathbb{S}^{n-1}

$$\nabla_{ij} x = -\delta_{ij} x.$$

From above computations, we obtain

$$\begin{aligned} -2Bh_k M_k &= -2Bh_k \left(\frac{f_k}{f} + \frac{h_k}{h} - \frac{1}{G} \langle \bar{\nabla} G, e_i \rangle b_{ki} - \frac{\varphi'}{\varphi} h_k \right) \\ &\leq 2B \left(\frac{|\nabla f| |\nabla h|}{f} + |\nabla h|^2 \frac{\varphi'}{\varphi} \right) + 2B \frac{h_k}{G} \langle \bar{\nabla} G, e_k \rangle b_{kk} \\ &\leq c_1 B + 2B \frac{h_k}{G} \langle \bar{\nabla} G, e_k \rangle b_{kk}, \end{aligned}$$

where c_1 is positive constants depending on $\|\varphi\|_{C^1(I_{[0,T]})}$, $\|f\|_{C^1(\mathbb{S}^{n-1})}$ and $\|h\|_{C^1(\mathbb{S}^{n-1} \times [0,T])}$.

We also have

$$\begin{aligned} -b^{11} M_{11} &= -b^{11} \left(\frac{f_{11}}{f} - \frac{f_1^2}{f^2} - \frac{h_1^2}{h^2} - \frac{\varphi'' h_1^2}{\varphi} + \frac{(\varphi')^2 h_1^2}{\varphi^2} \right) + b^{11} \frac{\varphi'(b_{11} - h)}{\varphi} - b^{11} \frac{b_{11} - h}{h} \\ &\quad - \frac{1}{G^2} \langle \bar{\nabla} G, e_1 \rangle^2 b_{11} + \frac{1}{G} \langle \nabla_1 \bar{\nabla} G, e_i \rangle - \frac{1}{G} \langle \bar{\nabla} G, x \rangle + \frac{1}{G} \langle \bar{\nabla} G, e_i \rangle b^{11} b_{11i} \\ &\leq c_2 b^{11} + c_3 + b_{11} \frac{1}{G} \langle \bar{\nabla}^2 G, e_1 \rangle, e_1 \rangle + \frac{1}{G} \langle \bar{\nabla} G, e_i \rangle b^{11} b_{11i} \\ &\leq c_2 b^{11} + c_3 + c_4 b_{11} + \frac{1}{G} \langle \bar{\nabla} G, e_i \rangle b^{11} b_{11i}, \end{aligned}$$

where c_2 is positive constant depending on $\|\varphi\|_{C^2(I_{[0,T]})}$, $\|f\|_{C^2(\mathbb{S}^{n-1})}$ and $\|h\|_{C^1(\mathbb{S}^{n-1} \times [0,T])}$ and positive constants c_3 and c_4 depend on $\|G\|_{C^2(\Omega_{[0,T]})}$.

From the definition of w ,

$$\begin{aligned} b^{11}\nabla_l b_{11} &= \nabla_l w + \frac{A}{h}h_l - 2Bh_{lk}h_k \\ &= \nabla_l w + \frac{A}{h}h_l - 2Bb_{lk}h_k + 2Bhh_l, \end{aligned}$$

At point p , we have

$$-b^{11}M_{11} \leq c_1B + c_2b^{11} + c_4b_{11} + c_5 - 2B\frac{h_k}{G}\langle \bar{\nabla}G, e_k \rangle b_{kk},$$

where c_5 depends on $\|G\|_{C^2(\Omega_{[0,T]})}$ and $\|h\|_{C^1(\mathbb{S}^{n-1} \times [0,T])}$.

Now, we have proved that

$$-b^{11}FM_{11} - 2BFh_kM_k \leq F(c_1B + c_2b^{11} + c_4b_{11} + c_5).$$

Hence, from (18) we have

$$\frac{\partial w}{\partial t} - Fb^{ii}\nabla_{ii}w \leq F(c_1B + c_2b^{11} + c_4b_{11} + c_5) - 2BF\sum_i b_{ii} + 4BFh(n-1) + \frac{AFn}{h}.$$

If we take

$$B > c_4,$$

then for b_{ii} large enough, there is

$$\begin{aligned} \frac{\partial w}{\partial t} - Fb^{ii}\nabla_{ii}w &\leq F(c_1B + c_2b^{11} + c_5) - BF\sum_i b_{ii} + 4BFh(n-1) + \frac{AFn}{h} \\ &< 0, \end{aligned}$$

which implies that

$$\frac{\partial w}{\partial t} \leq 0.$$

Now recalling Lemma 1 and Corollary 1, one can easily see the conclusion of this lemma holds. \square

Now we have proved that the principal curvatures of M_t have uniform upper and lower bounds, this together with Lemma 1 and Corollary 1 implies that the evolution equation (13) is uniformly parabolic on any finite time interval. Thus, the result of [44] and the standard parabolic theory show that the smooth solution of (13) exists for all time. Namely flow (6) has a long-time solution. And by these estimates again, a subsequence of M_t converges in C^∞ to a positive, smooth, uniformly convex hypersurface M_∞ in \mathbb{R}^n . Now to complete the proof of Theorem 2, it remains to check M_∞ satisfies Eq. (2) with $c = 1$.

4. EXISTENCE OF THE SOLUTIONS TO THE EQUATION

In this section, we first complete the proof of Theorem 2, namely we will prove the support function h_∞ of M_∞ satisfies the following equation:

$$(19) \quad \det(\nabla^2 h + hI)G(\nabla h + hx)\varphi(h) = f(x), \quad x \in \mathbb{S}^{n-1}.$$

To achieve this, we define a functional, which is non-increasing along the flow.

Define

$$\begin{aligned} \tilde{G}(r, u) &= \int^r G(s, u)s^{n-1}ds, \\ \tilde{\varphi}(t) &= \int \frac{1}{\varphi(s)}ds. \end{aligned}$$

It can be checked that flow (6) is the gradient flow of the functional given by

$$J(M) = \int_{S^{n-1}} \tilde{\varphi}(h)f(x)dx - \int_{S^{n-1}} \tilde{G}(\rho, u)du.$$

Here h and ρ are the support function and radial function of M respectively.

Now, we show that the functional $J(M_t)$ is non-increasing along the flow (6).

Lemma 4. *The functional $J(M_t)$ is non-increasing along the flow (6). That is*

$$\frac{\partial J}{\partial t} \leq 0,$$

and the equality holds if and only if M_t satisfies the elliptic equation (19).

Proof. We begin from the equation

$$\frac{\partial J}{\partial t} = \int_{S^{n-1}} \frac{1}{\varphi(h)}f(x)\frac{\partial h}{\partial t}dx - \int_{S^{n-1}} G(\rho, u)\rho^{n-1}\frac{\partial \rho}{\partial t}du.$$

Since

$$\frac{1}{\rho(u, t)}\frac{\partial \rho}{\partial t} = \frac{1}{h(x, t)}\frac{\partial h}{\partial t},$$

then by the fact

$$\rho^n du = \frac{h(x)}{\mathcal{K}(x)}dx,$$

we have

$$\begin{aligned} \frac{\partial J}{\partial t} &= \int_{S^{n-1}} \left(\frac{f(x)}{\varphi(h)} - \frac{G(\rho, u)}{\mathcal{K}(x)} \right) \frac{\partial h}{\partial t} dx \\ &= - \int_{S^{n-1}} \frac{G(\rho, u)}{\mathcal{K}(x)} h \left(\frac{f(x)}{\varphi(h)} \frac{\mathcal{K}(x)}{G(\rho, u)} - 1 \right)^2 dx \\ &\leq 0. \end{aligned}$$

The equality holds if and only if

$$\frac{f(x)}{\varphi(h)} \frac{\mathcal{K}(x)}{G(\rho, u)} = 1,$$

which is just the equation (19). □

From Lemma 1, Corollary 1 and the definition of $J(M_t)$, there exists a positive constant C which is independent of t , such that

$$(20) \quad |J(M_t)| \leq C, \quad \forall t > 0.$$

By the proof of Lemma 4, we have

$$(21) \quad J(M_0) - J(M_t) = \int_0^t dt \int_{S^{n-1}} \frac{G(X_t)}{\mathcal{K}(x)} h(x, t) \left(\frac{f(x)}{\varphi(h)} \frac{\mathcal{K}(x)}{G(X_t)} - 1 \right)^2 dx.$$

Combining (20) and (21), and recalling Lemmas 2 and 3, we have that

$$\int_0^\infty dt \int_{S^{n-1}} \left(\frac{f(x)}{\varphi(h)} \frac{\mathcal{K}(x)}{G(X_t)} - 1 \right)^2 dx < \infty.$$

This implies that there exists a subsequence of times $t_j \rightarrow \infty$ such that

$$\int_{S^{n-1}} \left(\frac{f(x)}{\varphi(h)} \frac{\mathcal{K}(x)}{G(X_{t_j})} - 1 \right)^2 dx \rightarrow 0 \quad \text{as } t_j \rightarrow \infty.$$

Passing to the limit, we obtain

$$\int_{S^{n-1}} \left(\frac{f(x)}{\varphi(h_\infty)} \frac{\mathcal{K}(x)}{G(X_\infty)} - 1 \right)^2 dx = 0,$$

where h_∞ is the support function of M_∞ , and $X_\infty = \overline{\nabla} h_\infty$. This implies that

$$\frac{f(x)}{\varphi(h_\infty)} \frac{\mathcal{K}(x)}{G(\overline{\nabla} h_\infty)} = 1, \quad \forall x \in \mathbb{S}^{n-1},$$

which is just equation (19). The proof of Theorem 2 is now completed.

At the same time, for Theorem 1, we have showed that for smooth φ, G, f , there exists a smooth solution h to (19). We now begin to prove Theorem 1 for general case. Note that

Lemma 5. *For any positive solution h to Eq. (2) with $c = 1$, we have*

$$C_1 \leq h \leq C_2 \quad \text{on } \mathbb{S}^{n-1},$$

where positive constants C_1 and C_2 depend only on the bounds of $f(x)$ and $\varphi(s) \max_{|y|=s} G(y) s^{n-1}$.

Proof. Assume $\max_{\mathbb{S}^{n-1}} h = h(\bar{x}) = M$. From its equation

$$\varphi(h) G(\overline{\nabla} h) \det(\nabla^2 h + hI) = f \quad \text{on } \mathbb{S}^{n-1},$$

we have

$$\begin{aligned} f(\bar{x}) &\leq \varphi(h(\bar{x})) G(h(\bar{x}) \bar{x}) h(\bar{x})^{n-1} \\ &\leq \varphi(M) \max_{|y|=M} G(y) M^{n-1}, \end{aligned}$$

which together with the assumption (5) implies that $M \leq C$ for a constant C depending only on f and $\varphi(s) \max_{|y|=s} G(y)s^{n-1}$. Similarly, considering the minimum point of h , we obtain the lower bound of h . \square

Now by a standard approximation and the regularity theory of Monge-Ampère equation, one can easily see Theorem 1 holds.

As said in the Introduction, for general φ and G , there is no uniqueness for solutions to Eq. (2). We here provide a special uniqueness result to end this paper. A similar result can be found in [23, Theorem 6.5]. Here we provide a different proof which is inspired by [17, Proposition 2.1].

Theorem 3. *Assume $G(y) = G(|y|)$. If whenever*

$$(22) \quad \varphi(\lambda s_1)G(\lambda s_2) \leq \varphi(s_1)G(s_2)\lambda^{1-n}$$

holds for some positive s_1, s_2 and λ , there must be $\lambda \geq 1$. Then the solution to the following equation

$$(23) \quad \det(\nabla^2 h + hI)G(\bar{\nabla} h)\varphi(h) = f(x)$$

is unique.

Proof. Let h_1 and h_2 be two solutions of (23), we first want to prove that

$$(24) \quad \max \frac{h_1}{h_2} \leq 1.$$

Suppose (24) is not true, namely $\max \frac{h_1}{h_2} > 1$. We want to derive a contradiction. Assume $\frac{h_1}{h_2}$ attains its maximum at $p_0 \in \mathbb{S}^{n-1}$. Then $h_1(p_0) > h_2(p_0)$. Let $L = \log \frac{h_1}{h_2}$, then at p_0 ,

$$0 = \nabla L = \frac{\nabla h_1}{h_1} - \frac{\nabla h_2}{h_2},$$

and

$$\begin{aligned} 0 &\geq \nabla^2 L \\ &= \frac{\nabla^2 h_1}{h_1} - \frac{\nabla h_1 \otimes \nabla h_1}{h_1^2} - \frac{\nabla^2 h_2}{h_2} + \frac{\nabla h_2 \otimes \nabla h_2}{h_2^2} \\ &= \frac{\nabla^2 h_1}{h_1} - \frac{\nabla^2 h_2}{h_2}. \end{aligned}$$

By equation (23), we have at p_0

$$\begin{aligned}
1 &= \frac{\det(\nabla^2 h_2 + h_2 I) G(|\nabla h_2 + h_2 x|) \varphi(h_2)}{\det(\nabla^2 h_1 + h_1 I) G(|\nabla h_1 + h_1 x|) \varphi(h_1)} \\
&= \frac{h_2^{n-1} \det(\frac{\nabla^2 h_2}{h_2} + I) G(h_2(\sqrt{|\frac{\nabla h_2}{h_2}|^2 + 1})) \varphi(h_2)}{h_1^{n-1} \det(\frac{\nabla^2 h_1}{h_1} + I) G(h_1(\sqrt{|\frac{\nabla h_1}{h_1}|^2 + 1})) \varphi(h_1)} \\
&\geq \frac{h_2^{n-1} G(h_2(\sqrt{|\frac{\nabla h_1}{h_1}|^2 + 1})) \varphi(h_2)}{h_1^{n-1} G(h_1(\sqrt{|\frac{\nabla h_1}{h_1}|^2 + 1})) \varphi(h_1)}.
\end{aligned}$$

Write $h_2(p_0) = \delta h_1(p_0)$, and $s = h_1(\sqrt{|\frac{\nabla h_1}{h_1}|^2 + 1})(p_0)$. Then the above inequality reads

$$\frac{\delta^{n-1} G(\delta s) \varphi(h_2)}{G(s) \varphi(h_1)} \leq 1,$$

namely

$$G(\delta s) \varphi(\delta h_1) \leq G(s) \varphi(h_1) \delta^{1-n}.$$

By our assumption (22), we have $\delta \geq 1$. Namely $h_2(p_0) \geq h_1(p_0)$. This is a contradiction. Thus (24) holds.

Interchanging h_1 and h_2 , (24) implies

$$\max \frac{h_2}{h_1} \leq 1.$$

Combining it with (24), we have $h_1 \equiv h_2$. The proof is completed. \square

REFERENCES

- [1] A. ALEXANDROFF, *Existence and uniqueness of a convex surface with a given integral curvature*, C. R. (Doklady) Acad. Sci. URSS (N.S.), 35 (1942), pp. 131–134.
- [2] B. ANDREWS, *Monotone quantities and unique limits for evolving convex hypersurfaces*, Internat. Math. Res. Notices, (1997), pp. 1001–1031.
- [3] B. ANDREWS, P. GUAN, AND L. NI, *Flow by powers of the Gauss curvature*, Adv. Math., 299 (2016), pp. 174–201.
- [4] K. J. BÖRÖCZKY AND F. FODOR, *The L_p dual Minkowski problem for $p > 1$ and $q > 0$* , J. Differential Equations, 266 (2019), pp. 7980–8033.
- [5] K. J. BÖRÖCZKY, P. HEGEDŰS, AND G. ZHU, *On the discrete logarithmic Minkowski problem*, Int. Math. Res. Not. IMRN, (2016), pp. 1807–1838.
- [6] K. J. BÖRÖCZKY, M. HENK, AND H. POLLEHN, *Subspace concentration of dual curvature measures of symmetric convex bodies*, J. Differential Geom., 109 (2018), pp. 411–429.
- [7] K. J. BÖRÖCZKY, E. LUTWAK, D. YANG, AND G. ZHANG, *The logarithmic Minkowski problem*, J. Amer. Math. Soc., 26 (2013), pp. 831–852.
- [8] K. J. BÖRÖCZKY AND H. T. TRINH, *The planar L_p -Minkowski problem for $0 < p < 1$* , Adv. in Appl. Math., 87 (2017), pp. 58–81.
- [9] S. BRENDLE, K. CHOI, AND P. DASKALOPOULOS, *Asymptotic behavior of flows by powers of the Gaussian curvature*, Acta Math., 219 (2017), pp. 1–16.

- [10] P. BRYAN, M. N. IVAKI, AND J. SCHEUER, *A unified flow approach to smooth, even L_p -Minkowski problems*, Anal. PDE, 12 (2019), pp. 259–280.
- [11] U. CAGLAR AND D. YE, *Affine isoperimetric inequalities in the functional Orlicz-Brunn-Minkowski theory*, Adv. in Appl. Math., 81 (2016), pp. 78–114.
- [12] C. CHEN, Y. HUANG, AND Y. ZHAO, *Smooth solutions to the L_p dual Minkowski problem*, Math. Ann., 373 (2019), pp. 953–976.
- [13] H. CHEN AND Q.-R. LI, *The L_p dual Minkowski problem and related parabolic flows*. Preprint.
- [14] S. CHEN AND Q.-R. LI, *On the planar dual Minkowski problem*, Adv. Math., 333 (2018), pp. 87–117.
- [15] S. CHEN, Q.-R. LI, AND G. ZHU, *The logarithmic Minkowski problem for non-symmetric measures*, Trans. Amer. Math. Soc., 371 (2019), pp. 2623–2641.
- [16] K.-S. CHOU AND X.-J. WANG, *A logarithmic Gauss curvature flow and the Minkowski problem*, Ann. Inst. H. Poincaré Anal. Non Linéaire, 17 (2000), pp. 733–751.
- [17] ———, *The L_p -Minkowski problem and the Minkowski problem in centroaffine geometry*, Adv. Math., 205 (2006), pp. 33–83.
- [18] W. J. FIREY, *Shapes of worn stones*, Mathematika, 21 (1974), pp. 1–11.
- [19] M. E. GAGE AND Y. LI, *Evolving plane curves by curvature in relative geometries. II*, Duke Math. J., 75 (1994), pp. 79–98.
- [20] R. J. GARDNER, D. HUG, AND W. WEIL, *The Orlicz-Brunn-Minkowski theory: a general framework, additions, and inequalities*, J. Differential Geom., 97 (2014), pp. 427–476.
- [21] R. J. GARDNER, D. HUG, W. WEIL, S. XING, AND D. YE, *General volumes in the Orlicz-Brunn-Minkowski theory and a related Minkowski problem I*, Calc. Var. Partial Differential Equations, 58 (2019), pp. Art. 12, 35 pp.
- [22] R. J. GARDNER, D. HUG, W. WEIL, AND D. YE, *The dual Orlicz-Brunn-Minkowski theory*, J. Math. Anal. Appl., 430 (2015), pp. 810–829.
- [23] R. J. GARDNER, D. HUG, S. XING, AND D. YE, *General volumes in the Orlicz-Brunn-Minkowski theory and a related Minkowski problem II*. arXiv: 1809.09753.
- [24] C. GERHARDT, *Non-scale-invariant inverse curvature flows in Euclidean space*, Calc. Var. Partial Differential Equations, 49 (2014), pp. 471–489.
- [25] P. GUAN AND L. NI, *Entropy and a convergence theorem for Gauss curvature flow in high dimension*, J. Eur. Math. Soc. (JEMS), 19 (2017), pp. 3735–3761.
- [26] C. HABERL, E. LUTWAK, D. YANG, AND G. ZHANG, *The even Orlicz Minkowski problem*, Adv. Math., 224 (2010), pp. 2485–2510.
- [27] C. HABERL, F. E. SCHUSTER, AND J. XIAO, *An asymmetric affine Pólya-Szegő principle*, Math. Ann., 352 (2012), pp. 517–542.
- [28] Y. HE, Q.-R. LI, AND X.-J. WANG, *Multiple solutions of the L_p -Minkowski problem*, Calc. Var. Partial Differential Equations, 55 (2016), pp. Art. 117, 13 pp.
- [29] M. HENK AND H. POLLEHN, *Necessary subspace concentration conditions for the even dual Minkowski problem*, Adv. Math., 323 (2018), pp. 114–141.
- [30] Q. HUANG AND B. HE, *On the Orlicz Minkowski problem for polytopes*, Discrete Comput. Geom., 48 (2012), pp. 281–297.
- [31] Y. HUANG AND Y. JIANG, *Variational characterization for the planar dual Minkowski problem*. Journal of Functional Analysis, 277 (2019), pp. 22092236.
- [32] Y. HUANG, J. LIU, AND L. XU, *On the uniqueness of L_p -Minkowski problems: the constant p -curvature case in \mathbb{R}^3* , Adv. Math., 281 (2015), pp. 906–927.
- [33] Y. HUANG, E. LUTWAK, D. YANG, AND G. ZHANG, *Geometric measures in the dual Brunn-Minkowski theory and their associated Minkowski problems*, Acta Math., 216 (2016), pp. 325–388.

- [34] ———, *The L_p -Aleksandrov problem for L_p -integral curvature*, J. Differential Geom., 110 (2018), pp. 1–29.
- [35] Y. HUANG AND Y. ZHAO, *On the L_p dual Minkowski problem*, Adv. Math., 332 (2018), pp. 57–84.
- [36] D. HUG, E. LUTWAK, D. YANG, AND G. ZHANG, *On the L_p Minkowski problem for polytopes*, Discrete Comput. Geom., 33 (2005), pp. 699–715.
- [37] M. N. IVAKI, *Deforming a hypersurface by Gauss curvature and support function*, J. Funct. Anal., 271 (2016), pp. 2133–2165.
- [38] H. JIAN AND J. LU, *Existence of solutions to the Orlicz-Minkowski problem*, Adv. Math., 344 (2019), pp. 262–288.
- [39] H. JIAN, J. LU, AND X.-J. WANG, *Nonuniqueness of solutions to the L_p -Minkowski problem*, Adv. Math., 281 (2015), pp. 845–856.
- [40] ———, *A priori estimates and existence of solutions to the prescribed centroaffine curvature problem*, J. Funct. Anal., 274 (2018), pp. 826–862.
- [41] H. JIAN, J. LU, AND G. ZHU, *Mirror symmetric solutions to the centro-affine Minkowski problem*, Calc. Var. Partial Differential Equations, 55 (2016), pp. Art. 41, 22 pp.
- [42] Y. JIANG AND Y. WU, *On the 2-dimensional dual Minkowski problem*, J. Differential Equations, 263 (2017), pp. 3230–3243.
- [43] H. KONE, *Valuations on Orlicz spaces and L^ϕ -star sets*, Adv. in Appl. Math., 52 (2014), pp. 82–98.
- [44] N. V. KRYLOV AND M. V. SAFONOV, *A property of the solutions of parabolic equations with measurable coefficients*, Izv. Akad. Nauk SSSR Ser. Mat., 44 (1980), pp. 161–175, 239.
- [45] A.-J. LI, *The generalization of Minkowski problems for polytopes*, Geom. Dedicata, 168 (2014), pp. 245–264.
- [46] Q.-R. LI, W. SHENG, AND W. XU-JIA, *Flow by Gauss curvature to the Aleksandrov and dual Minkowski problems*, J. Eur. Math. Soc. (2018, in press).
- [47] J. LU, *Nonexistence of maximizers for the functional of the centroaffine Minkowski problem*, Sci. China Math., 61 (2018), pp. 511–516.
- [48] ———, *A remark on rotationally symmetric solutions to the centroaffine Minkowski problem*, J. Differential Equations, 266 (2019), pp. 4394–4431.
- [49] J. LU AND H. JIAN, *Topological degree method for the rotationally symmetric L_p -Minkowski problem*, Discrete Contin. Dyn. Syst., 36 (2016), pp. 971–980.
- [50] J. LU AND X.-J. WANG, *Rotationally symmetric solutions to the L_p -Minkowski problem*, J. Differential Equations, 254 (2013), pp. 983–1005.
- [51] M. LUDWIG, *General affine surface areas*, Adv. Math., 224 (2010), pp. 2346–2360.
- [52] E. LUTWAK, *The Brunn-Minkowski-Firey theory. I. Mixed volumes and the Minkowski problem*, J. Differential Geom., 38 (1993), pp. 131–150.
- [53] E. LUTWAK, D. YANG, AND G. ZHANG, *L_p affine isoperimetric inequalities*, J. Differential Geom., 56 (2000), pp. 111–132.
- [54] ———, *On the L_p -Minkowski problem*, Trans. Amer. Math. Soc., 356 (2004), pp. 4359–4370.
- [55] ———, *L_p John ellipsoids*, Proc. London Math. Soc. (3), 90 (2005), pp. 497–520.
- [56] ———, *Orlicz projection bodies*, Adv. Math., 223 (2010), pp. 220–242.
- [57] ———, *L_p dual curvature measures*, Adv. Math., 329 (2018), pp. 85–132.
- [58] T. MESIKEPP, *M-addition*, J. Math. Anal. Appl., 443 (2016), pp. 146–177.
- [59] R. SCHNEIDER, *Convex bodies: the Brunn-Minkowski theory*, vol. 151 of Encyclopedia of Mathematics and its Applications, Cambridge University Press, Cambridge, expanded ed., 2014.
- [60] Y. SUN AND Y. LONG, *The planar Orlicz Minkowski problem in the L^1 -sense*, Adv. Math., 281 (2015), pp. 1364–1383.
- [61] J. URBAS, *An expansion of convex hypersurfaces*, J. Differential Geom., 33 (1991), pp. 91–125.

- [62] X. J. WANG, *Existence of convex hypersurfaces with prescribed Gauss-Kronecker curvature*, Trans. Amer. Math. Soc., 348 (1996), pp. 4501–4524.
- [63] D. XI, H. JIN, AND G. LENG, *The Orlicz Brunn-Minkowski inequality*, Adv. Math., 260 (2014), pp. 350–374.
- [64] Y. ZHAO, *The dual Minkowski problem for negative indices*, Calc. Var. Partial Differential Equations, 56 (2017), p. 56:18.
- [65] ———, *Existence of solutions to the even dual Minkowski problem*, J. Differential Geom., 110 (2018), pp. 543–572.
- [66] B. ZHU, J. ZHOU, AND W. XU, *Dual Orlicz-Brunn-Minkowski theory*, Adv. Math., 264 (2014), pp. 700–725.
- [67] G. ZHU, *The logarithmic Minkowski problem for polytopes*, Adv. Math., 262 (2014), pp. 909–931.
- [68] ———, *The L_p Minkowski problem for polytopes for $0 < p < 1$* , J. Funct. Anal., 269 (2015), pp. 1070–1094.
- [69] D. ZOU AND G. XIONG, *Orlicz-John ellipsoids*, Adv. Math., 265 (2014), pp. 132–168.

YANNAN LIU: SCHOOL OF MATHEMATICS AND STATISTICS, BEIJING TECHNOLOGY AND BUSINESS UNIVERSITY, BEIJING 100048, P.R. CHINA
E-mail address: liuyn@th.btbu.edu.cn

JIAN LU: SOUTH CHINA RESEARCH CENTER FOR APPLIED MATHEMATICS AND INTERDISCIPLINARY STUDIES, SOUTH CHINA NORMAL UNIVERSITY, GUANGZHOU 510631, P.R. CHINA
E-mail address: jianlu@m.scnu.edu.cn
E-mail address: lj-tshu04@163.com