

From Nesterov’s Estimate Sequence to Riemannian Acceleration

Kwangjun Ahn

KJAHN@MIT.EDU

Suvrit Sra

SUVRIT@MIT.EDU

Department of Electrical Engineering and Computer Science, Massachusetts Institute of Technology

Abstract

We propose the first global accelerated gradient method for Riemannian manifolds. Toward establishing our result we revisit Nesterov’s estimate sequence technique and develop an alternative analysis for it that may also be of independent interest. Then, we extend this analysis to the Riemannian setting, localizing the key difficulty due to non-Euclidean structure into a certain “metric distortion.” We control this distortion by developing a novel geometric inequality, which permits us to propose and analyze a Riemannian counterpart to Nesterov’s accelerated gradient method.

1. Introduction

First-order methods enjoy a well-developed theory for convex problems, while also demonstrating practical success in current machine learning tasks involving *non-convex* problems. But non-convex problems are in general intractable, so the corresponding theory of first-order methods either limits itself to local (stationarity) results, or relies on special structure that permits sharper global analysis.

A promising instance of such special structure arises via non-Euclidean geometry. Specifically, via the concept of *geodesic convexity* (*g-convexity*), which defines convexity along geodesics in metric spaces (Gromov, 1978; Burago et al., 2001; Bridson and Haefliger, 2013). Problems that are non-convex under a Euclidean view can sometimes be transformed into *g-convex* problems in a non-Euclidean space, potentially uncovering their *tractability*. This viewpoint has proved fruitful in several applications (e.g. see (Zhang and Sra, 2016, Section 1.1)), as well as towards tackling some theoretical questions in computer science, mathematics, and physics—see (Bürgisser et al., 2019; Goyal and Shetty, 2019) and references therein, as well as §1.2 of this paper.

Paralleling this viewpoint, several works have sought to analyze first-order methods in non-Euclidean settings, primarily in Riemannian manifolds (Udriste, 1994; Absil et al., 2009) and CAT(0) spaces (Bacák, 2014). Earlier studies focus on *asymptotic* analysis, while Zhang and Sra (2016) obtain the first *non-asymptotic* global iteration complexity analysis for Riemannian (stochastic) gradient descent methods, assuming *g-convexity*. Subsequently, iteration-complexity results were established for Riemannian proximal-point methods (Bento et al., 2017), Frank-Wolfe schemes (Weber and Sra, 2019), variance reduced methods (Zhang et al., 2016; Kasai et al., 2016; Zhang et al., 2018; Zhou et al., 2019), trust-region methods (Agarwal et al., 2018), among others.

Despite this progress, a landmark achievement of Euclidean convex optimization remains elusive in the Riemannian setting: that is, obtaining an analogue of Nesterov’s accelerated gradient method (Nesterov, 1983). This gap motivates the central question of our paper:

Is it possible to develop accelerated gradient methods for Riemannian manifolds?

This natural question, however, is believed to be highly non-trivial, especially because Nesterov’s analysis heavily relies on the linear structure of Euclidean space. In particular, recent efforts were able to answer this question only *partially*. See §1.2 for details.

1.1. Overview of our main results

In this paper, we take a major step toward answering this question by developing the *first global accelerated first-order method* for Riemannian manifolds (formal statement in Theorem 4.1):

Theorem 1.1 (Informal) *Let f be L -smooth and μ -strongly convex in a geodesic sense. Then, there exists a computationally tractable optimization algorithm satisfying*

$$f(x_t) - f(x_*) = O((1 - \xi_1)(1 - \xi_2) \cdots (1 - \xi_t)),$$

*where $\{\xi_t\}$ satisfies (i) $\{\xi_t\}_{t \geq 1} > \mu/L$ (**strictly faster than gradient descent**); and (ii) $\exists \lambda \in (0, 1)$ such that $\forall \epsilon > 0$, $|\xi_t - \sqrt{2\mu\Delta_\gamma}| \leq \epsilon$, for $t \geq \Omega(\frac{\log(1/\epsilon)}{\log(1/\lambda)})$ (eventually achieves **full acceleration**).*

To establish Theorem 1.1, we revisit and develop an alternative analysis to Nesterov’s analysis for the Euclidean case (§2). We use the technique of *potential functions* (also referred to as *Lyapunov functions*) (Lyapunov, 1992) for our analysis—see §5 for details. Interestingly, the parameters of the algorithm determined—from first principles—by our potential function analysis *exactly* satisfy the complicated recursive relations derived by Nesterov, thereby providing a simple alternative to his famous estimate sequence technique as a byproduct (§2.3). Further, we also develop a fixed-point based analysis of how accelerated convergence rates are derived from such complicated relations (§2.4), again providing a simple alternate to Nesterov’s original analysis based on clever algebras.

Our new analysis is then applied to the Riemannian case (§3 and §4). By carefully choosing potential function with the notion of projected distances, we nail down the main difficulty in the non-Euclidean case through the distortion in such distances (§3.2). We then demonstrate that our analysis for a simplified setting already recovers the local acceleration results by (Zhang and Sra, 2018). To tackle global acceleration, we establish a novel metric distortion inequality based on comparison theorems in Riemannian geometry (§4.1). Our metric distortion result engenders a distortion rate that is accessible to algorithm at each iteration. Finally, we show that such chosen tractable distortion rate decreases over iterations (§4.2), thereby obtaining Theorem 1.1.

1.2. Related work

Recently, a few efforts have been made to answer the main question of this paper. The first attempt is in (Liu et al., 2017), where the authors consider some nonlinear equations ((4) and (5) therein) and claim that the solutions to those equations yield accelerated schemes. However, it is *a priori* unclear whether such equations are even feasible and whether solving them is tractable.

As for different approach, Alimisis et al. (2020) establish a Riemannian analogue of the differential-equation approach to accelerated methods (Su et al., 2014). In particular, they propose and analyze several second-order ODEs on Riemannian manifolds that achieve accelerated convergence rates. Employing discretization results from the Euclidean case (Betancourt et al., 2018; Shi et al., 2019), they also derive some first-order methods from the ODEs. However, it is not clear whether the resulting methods achieve acceleration because such discretization techniques do *not* directly yield Nesterov’s accelerated method even in the Euclidean case.

The most concrete progress is the work (Zhang and Sra, 2018) that proves accelerated convergence of their algorithm, albeit only *locally*: acceleration is shown to hold within a neighborhood whose radius vanishes as the condition number L/μ and the curvature bound κ grow. Indeed, their work does not characterize how the algorithm behaves outside such a local neighborhood. This is in stark contrast with our *global* acceleration result. See §3.2 for a detailed comparison.

On a less related note, Siegel (2019) develop accelerated gradient methods for functions on *Stiefel manifolds*, by designing (i) a non-convex analogue of the adaptive restart methods due to O’donoghue and Candes (2015), and (ii) an analogue of momentum steps specifically for Stiefel manifolds. However, the accelerated performance is only verified *numerically*.

2. Warm up in the Euclidean case: alternative analysis of Nesterov’s optimal method

Before we consider the Riemannian setting, we first illustrate our analysis in the Euclidean case. In particular, we consider Nesterov’s optimal method which is derived based on an ingenious construction called an *estimate sequence* (2018, Ch. 2.2.1): For $t \geq 0$, the iterates are updated as

$$x_{t+1} \leftarrow y_t + \alpha_{t+1}(z_t - y_t) \quad (2.1a)$$

$$y_{t+1} \leftarrow x_{t+1} - \gamma_{t+1} \nabla f(x_{t+1}) \quad (2.1b)$$

$$z_{t+1} \leftarrow x_{t+1} + \beta_{t+1}(z_t - x_{t+1}) - \eta_{t+1} \nabla f(x_{t+1}). \quad (2.1c)$$

for given initial iterates $y_0 = z_0 \in \mathbb{R}^n$. This construction yields optimal first-order methods that achieve the lower bounds under the black-box complexity model (Nemirovsky and Yudin, 1983). Note that without resorting to estimate sequences, the updates (2.1) can be also derived via the *linear coupling* framework of Allen-Zhu and Orecchia (2014). See Appendix A for details.

Despite its fundamental nature, there is a well-known puzzling aspect of Nesterov’s construction. In order to guarantee the standing assumption of the estimate sequence technique (2018, (2.2.3)), Nesterov’s original analysis (2018, page 87) finds complicated recursive relations between parameters $\alpha, \beta, \gamma, \eta$ via some non-trivial algebraic “tricks.” These tricks are carried out in a fortuitous manner, obscuring the driving principle and the scope of the underlying technique. Notably, the work (Zhang and Sra, 2018) favors estimate sequences over other approaches, but still achieves only local acceleration. Therefore, toward our goal of obtaining global acceleration, we revisit Euclidean analysis of acceleration from first-principles.

In particular, we provide an alternative analysis of iteration (2.1) that sheds new light on understanding the scope of Nesterov’s original analysis. Our analysis employs a *potential function*,¹ a classical tool for studying the stability of dynamical systems (Lyapunov, 1992), which has received a resurgence of interest due to its success in analyzing first-order optimization methods (see §5 for details). Roughly, the convergence analysis based on potential-functions proceeds as follows:

1. **Choose a potential function:** First, choose a performance measure \mathcal{E}_t defined for each step that “measures” how close the iterates at step t are to the optimal solution. Having chosen \mathcal{E}_t , define the potential function as $\Phi_t := A_t \mathcal{E}_t$ for some quantity A_t to be determined.
2. **Ensure potential decrease:** Choose an increasing sequence A_t for which Φ_t is decreasing.

Once Φ_t is chosen as above, it immediately implies that $\mathcal{E}_t \leq \mathcal{E}_0/A_t$, yielding a convergence rate of $O(1/A_t)$ under the chosen performance measure. See §5 for a discussion positioning our proposed potential function to related as well as different techniques for analyzing acceleration.

2.1. Choosing the potential function

The key to potential function based analysis is to choose the “correct” performance measure. For an iterate u_t at step t , two prototypical choices might be (i) the suboptimality $\mathcal{E}_t = f(u_t) - f(x_*)$;

¹ Also known as *Lyapunov function* in control theory or *invariant* in theoretical computer science and mathematics

and (ii) the distance to an optimal point $\|u_t - x_*\|$. Indeed, many existing analyses correspond to choosing either one for performance measure, as explicitly observed in (Bansal and Gupta, 2019).

For iteration (2.1), it turns out that a weighted sum of the suboptimality $f(y_t) - f(x_*)$ and the distance $\|z_t - x_*\|^2$ is the “correct” performance measure, i.e., we choose the potential function as

$$\Phi_t := A_t \cdot (f(y_t) - f(x_*)) + B_t \cdot \|z_t - x_*\|^2 \quad (2.2)$$

for some $A_t > 0$ and $B_t \geq 0$. By taking a weighted sum of the two measures, this performance measure does not require either one to be monotonically decreasing over iterations. This property, also known as *non-relaxational property*, was indeed the main innovation in Nesterov’s illustrious paper (1983). The reason why we choose y_t for the cost and z_t for the distance will be clearer when we carry out the analysis. See Remark 2.3.

The current form of the potential function (2.2) is not new; indeed it also appears in prior works (Wilson et al., 2016; Diakonikolas and Orecchia, 2019; Bansal and Gupta, 2019), although with different motivations; see §5 for precise details. Moreover, this potential also has an interpretation under the linear coupling framework of (Allen-Zhu and Orecchia, 2014); see Appendix A.

2.2. Potential difference calculations

Having chosen the potential function (2.2), the main goal now is to choose the parameters A_{t+1} , B_{t+1} , α_{t+1} , β_{t+1} , γ_{t+1} , η_{t+1} so that the potential decreases, i.e., $\Phi_{t+1} - \Phi_t \leq 0$. To that end, we first express the potential difference $\Phi_{t+1} - \Phi_t$ more simply and derive a manageable upper bound using *first principles*. Using definition (2.2), the difference $\Phi_{t+1} - \Phi_t$ can be split into two parts:

$$A_{t+1} \cdot (f(y_{t+1}) - f(x_*)) - A_t \cdot (f(y_t) - f(x_*)) \quad (2.3)$$

$$+ B_{t+1} \cdot \|z_{t+1} - x_*\|^2 - B_t \cdot \|z_t - x_*\|^2. \quad (2.4)$$

Since $\alpha, \beta, \gamma, \eta$ will only appear with index $t + 1$, we drop their subscripts for simplicity. We first relate the terms for step $t + 1$ with those for step t . To do that, we reinterpret (2.1). Using the notation $\text{Grad}_{s \cdot \nabla}(x) := x - s \cdot \nabla$, the updates (2.1b) and (2.1c) can be rewritten as

$$y_{t+1} = \text{Grad}_{\gamma \cdot \nabla f(x_{t+1})}(x_{t+1}) \quad (2.1b')$$

$$z_{t+1} = \text{Grad}_{\eta \cdot \nabla f(x_{t+1})}(x_{t+1} + \beta(z_t - x_{t+1})), \quad (2.1c')$$

respectively. Now the difference between (2.1b') and (2.1c') is clear: the first one is an *exact* gradient step in the sense that $\nabla = \nabla f(x)$, while the second one is *inexact*. Hence, in relating the terms for step $t + 1$ with those for step t , we need to invoke different analyses for two different gradient steps.

Let us now review folklore analyses for gradient steps; the proofs are provided in Appendix C.1. First, when the gradient step is exact, the following result is well-known for an L -smooth cost:

Proposition 2.1 *Assume $\nabla = \nabla f(x)$, and let $y = \text{Grad}_{s \cdot \nabla}(x)$. If f is L -smooth, then the gradient step decreases cost: $f(y) - f(x) \leq -s(1 - Ls/2) \|\nabla\|^2$.*

When the gradient step is inexact, one can only guarantee the following weaker result:

Proposition 2.2 ² *Let $z = \text{Grad}_{s \cdot \nabla}(x)$. Then, for any x_* , $\|z - x_*\|^2 - \|x - x_*\|^2 = s^2 \|\nabla\|^2 + 2s \langle \nabla, x_* - x \rangle$, i.e., (inexact) gradient step decreases the distance to x_* as long as direction $-\nabla$ is well aligned with the vector $x_* - x$ and has sufficiently small norm.*

²Actually, the fact that this type of analyses (perhaps, better known as mirror descent analyses) can be applied to inexact gradient steps has brought about many successful applications in *online optimization* (see e.g. (Bubeck, 2011)).

Remark 2.3 These observations reveal why we use y_t for the cost term and z_t for the distance term in (2.2): Proposition 2.1 deals with the cost, while Proposition 2.2 deals with the distance.

Now we apply Proposition 2.1 to (2.1b') and Proposition 2.2 to (2.1c'). For clarity, we denote:

$$\Delta_\gamma := \gamma(1 - L\gamma/2), \quad \nabla := \nabla f(x_{t+1}), \quad X := -(x_* - x_{t+1}), \quad \text{and } W := z_t - x_{t+1}. \quad (2.5)$$

With this notation, Propositions 2.1 and 2.2 imply: $f(y_{t+1}) \leq f(x_{t+1}) - \Delta_\gamma \|\nabla\|^2$ and $\|z_{t+1} - x_*\|^2 = \|X + \beta W\|^2 + \eta^2 \|\nabla\|^2 - 2\eta \langle \nabla, X + \beta W \rangle$. Plugging these two back into to (2.3) and (2.4), one can derive the following upper bound on $\Phi_{t+1} - \Phi_t$ in terms of the vectors ∇, X, W from *first principles* (i.e., using only smoothness and (strong) convexity; see Appendix C.2):

$$\begin{aligned} \Phi_{t+1} - \Phi_t &\leq C_1 \cdot \|W\|^2 + C_2 \cdot \|X\|^2 + C_3 \|\nabla\|^2 + C_4 \cdot \langle W, X \rangle + C_5 \cdot \langle W, \nabla \rangle + C_6 \cdot \langle X, \nabla \rangle, \quad (2.6) \\ \text{where } \begin{cases} C_1 := \beta^2 B_{t+1} - B_t - \frac{\mu}{2} \frac{\alpha^2}{(1-\alpha)^2} A_t, & C_2 := B_{t+1} - B_t - \frac{\mu}{2} (A_{t+1} - A_t), \\ C_3 := \eta^2 B_{t+1} - \Delta_\gamma \cdot A_{t+1}, & C_4 := 2 \cdot (\beta B_{t+1} - B_t), \\ C_5 := \frac{\alpha}{1-\alpha} A_t - 2\beta\eta B_{t+1}, & \text{and } C_6 := (A_{t+1} - A_t) - 2\eta B_{t+1}. \end{cases} \end{aligned}$$

The reader may have noticed that in (2.5) the vectors ∇, X, W are rooted at the same point x_{t+1} . This choice is deliberate, and will be crucial in the Riemannian setting, because there the vectors have to lie in the same *tangent space* for their compatibility. See Appendix C.5.

2.3. Ensuring potential decrease

Having established the bound (2.6), our goal is to now choose $A_{t+1}, B_{t+1}, \alpha, \beta, \gamma, \eta$ given A_t, B_t so that (2.6) is non-positive (recall that we have dropped indices of $\alpha_{t+1}, \beta_{t+1}, \gamma_{t+1}, \eta_{t+1}$). In general, it is difficult to ensure non-positivity of a symbolic expression; but since (2.6) is a quadratic form, one avenue might be to turn it into a *negative sum of squares* (“–SoS”). The simplest strategy to make it “–SoS” would be to choose parameters so that the coefficients C_4, C_5, C_6 of the cross terms are set to 0, while $C_1, C_2, C_3 \leq 0$. It turns out this strategy *fully* determines the parameters.

- *Coefficients of cross terms characterize α, β, η in terms of A_{t+1}, B_{t+1} :* From $C_6 = 0$, we get $\eta = (A_{t+1} - A_t)/(2B_{t+1})$, and from $C_4 = 0$, we get $\beta = B_t/B_{t+1}$. Plugging these into $C_5 = 0$, we obtain the equation $\frac{\alpha}{1-\alpha} A_t = (A_{t+1} - A_t) B_t / B_{t+1}$. To summarize:

$$\eta = \frac{A_{t+1} - A_t}{2B_{t+1}}, \quad \beta = \frac{B_t}{B_{t+1}} \quad \text{and} \quad \frac{\alpha}{1-\alpha} = \frac{(A_{t+1} - A_t) B_t}{A_t B_{t+1}}. \quad (2.7)$$

- *For a fixed γ , coefficients of squared terms determines A_{t+1}, B_{t+1} based on given A_t, B_t :* Beginning with $C_3 \leq 0$, we replace η with the one from (2.7) to obtain:

$$(A_{t+1} - A_t)^2 / (4\Delta_\gamma \cdot A_{t+1}) \leq B_{t+1}. \quad (2.8)$$

Plugging (2.8) into $C_2 \leq 0$, we get an inequality only in terms of A_{t+1} (assuming γ is fixed):

$$(A_{t+1} - A_t)^2 / (4\Delta_\gamma \cdot A_{t+1}) - (A_{t+1} - A_t) \frac{\mu}{2} \leq B_t. \quad (2.9)$$

Recall from the potential function analysis that we need to choose A_{t+1} as large as possible. It turns out that the largest possible A_{t+1} satisfies (2.9) as an equality (hence (2.8) as well). To see this, let us follow Nesterov's notation and consider the *suboptimality shrinking ratio*

$1 - \xi := A_t/A_{t+1}$.³ With this ratio, inequality (2.9) can be rewritten as:

$$\xi(\xi - 2\mu\Delta_\gamma)/(1 - \xi) \leq 4\Delta_\gamma \cdot B_t/A_t. \quad (2.10)$$

In (2.10) note that the RHS is a nonnegative constant (assuming $\Delta_\gamma > 0$ is already chosen) and the LHS is an increasing function on $[2\mu\Delta_\gamma, 1)$ whose value is 0 at $2\mu\Delta_\gamma$ and approaches $+\infty$ as $\xi \rightarrow 1$. Hence, the largest ξ (equivalently, the largest A_{t+1}) satisfies (2.10) (or equivalently, (2.9)) with equality. Consequently, this choice of ξ also satisfies (2.8) with equality. One can then verify that this choice satisfies $\beta^2 B_{t+1} \leq B_t$ and hence implies $C_1 \leq 0$ (see Appendix C.3).

- *Lastly, choose γ from $(0, 2/L)$:* Now the last variable to determine is γ . The above calculations are valid as long as $\Delta_\gamma > 0$, so we can arbitrarily choose γ in $(0, 2/L)$. Note that most accelerated methods in the literature choose $\gamma = 1/L$ since it is the maximizer of Δ_γ .

Furthermore, note that one can express the other variables $B_{t+1}, \alpha, \beta, \eta$ via equation (2.7) and the equality version of (2.8) (see Appendix C.3). Therefore, our findings can be summarized in the main result of this section as follows (after recovering the indices of $\alpha, \beta, \gamma, \eta$):

Theorem 2.1 (Parameter choice for potential decrease) *For $\gamma_{t+1} \in (0, 2/L)$, $\Delta_\gamma := \gamma_{t+1}(1 - L\gamma_{t+1}/2)$, and given y_t, z_t and $A_t, B_t > 0$,*

1. *Compute $\xi_{t+1} \in [2\mu\Delta_\gamma, 1)$ satisfying $\frac{\xi_{t+1}(\xi_{t+1} - 2\mu\Delta_\gamma)}{1 - \xi_{t+1}} = 4\Delta_\gamma \cdot \frac{B_t}{A_t}$.* (2.11)
2. *Determine parameters based on ξ : $A_{t+1} = \frac{A_t}{1 - \xi_{t+1}}$, $B_{t+1} = \frac{\xi_{t+1}^2}{1 - \xi_{t+1}} \cdot \frac{A_t}{4\Delta_\gamma}$, $\alpha_{t+1} = \frac{\xi_{t+1} - 2\mu\Delta_\gamma}{1 - 2\mu\Delta_\gamma}$, $\beta_{t+1} = 1 - 2\mu\Delta_\gamma\xi_{t+1}^{-1}$, and $\eta_{t+1} = 2\Delta_\gamma\xi_{t+1}^{-1}$.*

Then, y_{t+1}, z_{t+1} defined as per (2.1) satisfy $\Phi_{t+1} \leq \Phi_t$ (see (2.2)). In other words,

$$f(y_{t+1}) - f(x_*) + \frac{\xi_{t+1}^2}{4\Delta_\gamma} \cdot \|z_{t+1} - x_*\|^2 \leq (1 - \xi_{t+1}) \cdot \left[f(y_t) - f(x_*) + \frac{B_t}{A_t} \cdot \|z_t - x_*\|^2 \right].$$

Proof An immediate consequence of the above calculations; See Appendix C.3 for details. ■

A noteworthy outcome of the above analysis is that the parameter choices in Theorem 2.1 *exactly match* those of Nesterov's "General Scheme for Optimal Method" (2018, (2.2.1)). Hence, our approach actually recovers Nesterov's optimal method that encompasses both strongly and non-strongly convex costs, without resorting to the estimate sequence technique. Another important byproduct of our analysis is the convergence of z_t to x_* for $\mu > 0$ (in which case, $\xi > 0$), a property proved via additional analysis in the literature (see e.g., (Gasnikov and Nesterov, 2018, Corollary 1)). More importantly, this convergence plays a crucial role in the Riemannian setting. See §4.2.

Observe that upon applying Theorem 2.1 recursively, we can deduce that

$$f(y_t) - f(x_*) = O((1 - \xi_1)(1 - \xi_2) \cdots (1 - \xi_t)). \quad (2.12)$$

Thus, to identify the convergence rate of iteration (2.1) with parameters chosen via Theorem 2.1, we only need to study how the sequence $\{\xi_t\}$ evolves. This evolution is the focus of the next subsection.

³More precisely, Nesterov's estimate sequence analysis finds $\alpha_i \in (0, 1)$ such that $f(y_t) - f(x_*) \leq \prod_{i=1}^t (1 - \alpha_i) \cdot [f(x_0) - f(x_*) + C \|x_0 - x_*\|^2]$ for some constant $C > 0$, where x_0 is an initial iterate (see (2018, Theorem 2.2.1)). Note that these α_i 's exactly correspond to our suboptimality shrinking ratios.

2.4. Identifying the convergence rate of (2.1): a simple analysis based on fixed-point iteration

For studying the evolution of ξ_t , we consider the strongly convex case ($\mu > 0$) and assume that γ_t is fixed to be a constant $\gamma \in (0, 2/L)$. This assumption is not stringent as most accelerated methods in the literature choose $\gamma_t \equiv 1/L$.

Our approach below offers an alternative to its counterpart in Nesterov's book (2018, Lemma 2.2.4): In contrast to Nesterov's analysis based on clever algebraic manipulations, our approach *directly* analyzes the evolution of the sequence by drawing a connection to *fixed point iterations*. Moreover, this approach will generalize better to the more complicated Riemannian setting. As a byproduct of our simple approach, we were able to remove the technical condition on ξ_0 required by Nesterov's analysis. See Remark 2.4.

First, let us pinpoint the recursive relation satisfied by ξ_t . Applying Theorem 2.1 for $t \leftarrow t - 1$, we have $\frac{B_t}{A_t} = \frac{\xi_t^2}{4\Delta_\gamma}$. Hence, (2.11) implies the following *nonlinear* recursive relation on ξ_t 's:

$$\xi_{t+1}(\xi_{t+1} - 2\mu\Delta_\gamma)/(1 - \xi_{t+1}) = \xi_t^2. \quad (2.13)$$

Now, our objective is to characterize the evolution of ξ_t from (2.13). Intuitively, (2.13) can be construed as the recursive relation for computing the root of $\phi(v) = \psi(v)$, where $\phi(v) := \frac{v(v-2\mu\Delta_\gamma)}{(1-v)}$ and $\psi(v) := v^2$. Since the root is equal to $v = \sqrt{2\mu\Delta_\gamma}$, one can guess that $\xi_t \rightarrow \sqrt{2\mu\Delta_\gamma}$. See Figure 1 for illustration. The following lemma confirms this guess.




Figure 1: An illustration of the evolution of (2.13) for $2\mu\Delta_\gamma = 0.25$. We plot $\phi = \frac{v(v-2\mu\Delta_\gamma)}{(1-v)}$ in blue and $\psi(v) = v^2$ in red.

Lemma 2.1 (Evolution of (2.13)) *For an arbitrary initial value $\xi_0 \geq 0$, let ξ_t ($t \geq 1$) be the sequence of numbers defined as per (2.13). Then, $\xi_t \in [2\mu\Delta_\gamma, 1)$ for all $t \geq 1$. Furthermore, if*

$$\begin{cases} \xi_0 > \sqrt{2\mu\Delta_\gamma}, \\ \xi_0 = \sqrt{2\mu\Delta_\gamma}, \\ \xi_0 < \sqrt{2\mu\Delta_\gamma}, \end{cases} \quad \text{then} \quad \begin{cases} \xi_t \searrow \sqrt{2\mu\Delta_\gamma} \text{ as } t \rightarrow \infty. \\ \xi_t \equiv \sqrt{2\mu\Delta_\gamma}. \\ \xi_t \nearrow \sqrt{2\mu\Delta_\gamma} \text{ as } t \rightarrow \infty. \end{cases} \quad \text{In particular, the convergences are geometric.}$$

Proof The proof and the formal statement (Lemma C.1) are provided in Appendix C.4. ■

Lemma 2.1 delivers the desired accelerated convergence rate:

Corollary 2.1 *If $\xi_0 \geq \sqrt{2\mu\Delta_\gamma}$, then $f(y_t) - f(x_*) = O(\prod_{i=1}^t (1 - \sqrt{2\mu\Delta_\gamma})) = O(\exp(-t\sqrt{2\mu\Delta_\gamma}))$. In particular, setting $\gamma = 1/L$, $f(y_t) - f(x_*) = O(\exp(-t\sqrt{\mu/L}))$.*

Remark 2.4 (Removing technical conditions in Nesterov's analysis) *Nesterov's original analysis requires a technical condition on the initial value ξ_0 : $\sqrt{\mu/L} \leq \xi_0 \leq (2(3+\mu/L))/(3+\sqrt{21+4\mu/L})$ (Nesterov, 2018, (2.2.21)). In contrast, our analysis reveals that the upper bound on ξ_0 is not needed; the lower bound is also not needed in the sense that ξ_t converges to $\sqrt{\mu/L}$, the accelerated rate.*

3. Generalization to the non-Euclidean case: Riemannian potential function analysis

This section develops the first key ingredient towards obtaining our main theorem (Theorem 4.1), namely, Theorem 3.1 that is a Riemannian analogue of Theorem 2.1. We begin by introducing some preliminaries and stating the Riemannian analogue of iteration (2.1).

3.1. Riemannian geometry and Riemannian analogue of Nesterov method

We recall below some basic concepts from Riemannian geometry, and defer to textbooks (e.g., (Jost, 2008; Burago et al., 2001)) for an in depth introduction. A *Riemannian manifold* is a smooth manifold M equipped with a smoothly varying inner product $\langle \cdot, \cdot \rangle_x$ (called the *Riemannian metric*) defined for each $x \in M$ on the tangent space $T_x M$. One can define geometric concepts such as angles, length of curves, and surface areas on a Riemannian manifold M . With the concept of length of curves, one can introduce a distances function d on M and consequently view (M, d) as a metric space. Length also allows us to define analogues of straight lines, namely *geodesics*: A curve is a *geodesic* if it is locally distance minimizing. The notions of curvature can be also defined in M . We focus on *sectional curvature*, which characterizes the notion of curvature by measuring Gaussian curvatures of 2-dimensional submanifolds of M . We make the following key assumption on curvature:

Assumption 1 *We assume that the sectional curvature is lower bounded by $-\kappa$ for some nonnegative constant κ . This is a widely used standard assumption in Riemannian geometry; for a textbook treatment dedicated to this assumption see e.g., (Burago et al., 2001, Chapter 10).*

Operations on Riemannian manifolds. One can define analogues of vector addition and subtraction in Riemannian manifolds via the concept of *exponential map*. An exponential map $\text{Exp}_x : T_x M \rightarrow M$ maps $v \in T_x M$ to $g(1) \in M$ for a geodesic g with $g(0) = x$ and $g'(0) = v$. We remark that it is always well-defined locally. Notice that $\text{Exp}_x(v) \in M$ is an analogue of vector addition “ $x + v$.” Similarly, the inverse map $\text{Exp}_x^{-1}(y) \in T_x M$ is a non-Euclidean analogue of vector subtraction “ $y - x$.” For Exp_x^{-1} to be well-defined for each x , we assume that any two points on M are connected by a unique geodesic. This property is called *uniquely geodesic*, and it is valid locally for general Riemannian manifolds and globally for Hadamard manifolds (Riemannian manifolds with global non-positive sectional curvatures). We assume further that $\text{Exp}_x, \text{Exp}_x^{-1}$ can be computed at each x , which is in fact the case for many widely used matrix manifolds (Absil et al., 2009).

Convexity on Riemannian manifolds. The notion of convexity can be extended to Riemannian manifolds using geodesics where a convex combination of two points is defined along the geodesic connecting them. This generalized notion of convexity is called *geodesic convexity* (*g-convexity* for short) (Gromov, 1978). In particular, one can define geodesic-smoothness and (strong) *g-convexity* for a function $f : M \rightarrow \mathbb{R}$ akin to their Euclidean counterparts.

Assumption 2 *We assume that the cost function f is geodesically L -smooth and μ -strongly convex (formal definitions in Appendix C.5; see also (Zhang and Sra, 2016, Section 2.3)).*

Now we are ready to generalize Nesterov's method (2.1) to Riemannian manifolds. Using the above noted Riemannian analogues of vector addition and subtractions, (2.1) becomes

$$x_{t+1} \leftarrow \text{Exp}_{y_t}(\alpha_{t+1} \text{Exp}_{y_t}^{-1}(z_t)) \quad (3.1a)$$

$$y_{t+1} \leftarrow \text{Exp}_{x_{t+1}}(-\gamma_{t+1} \nabla f(x_{t+1})) \quad (3.1b)$$

$$z_{t+1} \leftarrow \text{Exp}_{x_{t+1}}(\beta_{t+1} \text{Exp}_{x_{t+1}}^{-1}(z_t) - \eta_{t+1} \nabla f(x_{t+1})) . \quad (3.1c)$$

Note that updates (3.1b) and (3.1c) are well-defined since $\nabla f(x_{t+1})$ lies in the tangent space $T_x M$. Having formalized the Riemannian version of (2.1), we study its potential function analysis.

3.2. Riemannian potential function analysis and metric distortion

Since iteration (3.1) is a direct analog of its Euclidean counterpart (2.1), one may be tempted to study the potential function $\Psi_t := A_t \cdot (f(y) - f(x_*)) + B_t \cdot d(z_t, x_*)^2$ that is a direct analog of (2.2). However, it turns out that the following less direct potential function is much more advantageous:

$$\Psi_t := A_t \cdot (f(y_t) - f(x_*)) + B_t \cdot \|\text{Exp}_{x_t}^{-1}(z_t) - \text{Exp}_{x_t}^{-1}(x_*)\|^2. \quad (3.2)$$

The distance term in (3.2) is preferable to $d(z_t, x_*)^2$ because it lets us use Euclidean geometry (since it is defined on the tangent space $T_{x_t}M \cong \mathbb{R}^n$) to control it. To simplify notation, we define:

Definition 3.1 (Projected distance) *For any three points $u, v, w \in M$, the projected distance between v and w relative to u is defined as $d_u(v, w) := \|\text{Exp}_u^{-1}(v) - \text{Exp}_u^{-1}(w)\|$.*

However, there is one fundamental hurdle *inherent* to comparing distances in the Riemannian setting: we need to handle the *incompatibility* of metrics between two different points. A key advantage of the potential function analysis is that one only needs to focus on comparing the distances appearing in *adjacent* terms, namely Ψ_t and Ψ_{t+1} , which simplifies the argument considerably. Motivated by (3.2), we define the following quantity for comparing distances:

Definition 3.2 (Valid distortion rate) *We say δ_t is a valid distortion rate at iteration $t \geq 1$ if the following inequality holds: $d_{x_t}(z_{t-1}, x_*)^2 \leq \delta_t \cdot d_{x_{t-1}}(z_{t-1}, x_*)^2$.*

Assuming the existence of valid distortion rates at *each* iteration, we can analyze iteration (3.1) analogously to the analysis in §2.2 and §2.3, to obtain the main result of this section:

Theorem 3.1 (Riemannian analogue of Theorem 2.1) *Assume that δ_t is a valid distortion rate at iteration t . For $\gamma_{t+1} \in (0, 2/L)$, $\Delta_\gamma := \gamma_{t+1}(1 - L\gamma_{t+1}/2)$, and given y_t, z_t and $A_t, B_t > 0$,*

1. *Compute $\xi_{t+1} \in [2\mu\Delta_\gamma, 1)$ satisfying $\frac{\xi_{t+1}(\xi_{t+1} - 2\mu\Delta_\gamma)}{1 - \xi_{t+1}} = \frac{4\Delta_\gamma}{\delta_{t+1}} \cdot \frac{B_t}{A_t}$.* (3.3)
2. *Compute $A_{t+1}, B_{t+1}, \alpha_{t+1}, \beta_{t+1}, \eta_{t+1}$ as in Theorem 2.1.*

Then, y_{t+1}, z_{t+1} generated via iteration (3.1) satisfy $\Psi_{t+1} \leq \Psi_t$ (see (3.2)). In other words,

$$f(y_{t+1}) - f(x_*) + \frac{\xi_{t+1}^2}{4\Delta_\gamma} \cdot d_{x_{t+1}}(z_{t+1}, x_*)^2 \leq (1 - \xi_{t+1}) \cdot [f(y_t) - f(x_*) + \frac{B_t}{A_t} \cdot d_{x_t}(z_t, x_*)^2].$$

Proof The proof resembles the arguments in §2.2 and §2.3, modulo the appearance of valid distortion rates in (3.3). See Appendix C.5 for details. ■

As before, we can deduce from Theorem 3.1 the suboptimality gap bound (2.12). Hence, to identify the convergence rate we only need to determine the evolution of $\{\xi_t\}$. We provide an illustrative example below, before moving onto the full accelerated algorithm in §4.

Illustrative example: constant distortion rate. Consider the simplified case where $\delta_t \equiv \delta \geq 1$ for all $t \geq 0$. Under this constant distortion condition, similarly to recursion (2.13), we can obtain a recursive relation on $\{\xi_t\}$ by choosing $\gamma_t \equiv \gamma \in (0, 2/L)$:

$$\xi_{t+1}(\xi_{t+1} - 2\mu\Delta_\gamma)/(1 - \xi_{t+1}) = \xi_t^2/\delta. \quad (3.4)$$

Here the only difference relative to (2.13) is now the RHS is divided by δ . Analogously to Lemma 2.1, one can establish geometric convergence of ξ_t to the fixed point $\xi(\delta)$ of equation (3.4) (see Lemma C.1). Solving for $\xi(\delta)$ explicitly, we obtain the following analogue of Corollary 2.1:

Corollary 3.1 *If $\xi_0 \geq \xi(\delta) := \sqrt{(\delta-1)^2 + 8\delta\mu\Delta_\gamma}/2 - (\delta-1)/2$, then the following convergence rate holds: $f(y_t) - f(x_*) = O(\prod_{i=1}^t (1 - \xi(\delta))) = O(\exp(-t \cdot \xi(\delta)))$. In particular, setting $\gamma = 1/L$, $f(y_t) - f(x_*) = O((\exp(-\frac{t}{2}\{\sqrt{(\delta-1)^2 + 4\delta\mu/L} - \frac{t}{2}(\delta-1)\}))$.*

A notable aspect of Corollary 3.1 is that it characterizes a trade-off between the metric distortion and the convergence rate of the resulting algorithm. This point is elaborated by the following remark:

Remark 3.3 (Properties of $\xi(\delta)$) *When there is no distortion, i.e., $\delta = 1$, then $\xi(1) = \sqrt{2\mu\Delta_\gamma}$ since (3.4) becomes (2.13). Moreover, one can verify that $\xi(\delta)$ is (strictly) decreasing in δ , implying that the algorithm's performance gets worse as the distortion gets severer (see Appendix C.8 for verification). Hence, $\xi(\delta) > \lim_{\delta \rightarrow \infty} \xi(\delta) = 2\mu\Delta_\gamma$ for all $\delta > 1$, implying that the convergence rate is always **strictly** better than gradient descent no matter how severe the distortion is.*

Interestingly, our illustrative analysis already recovers the local acceleration result of Zhang and Sra (2018). More specifically, they showed that if the distance between initial iterates and the global optimum is bounded by $1/20 \cdot \kappa^{1/2}(L/\mu)^{-3/4}$ (see Appendix F therein), then the distortion is bounded by $\delta = 1 + 1/5 \cdot (L/\mu)^{-1/2}$. Simplifying the value $\xi(\delta)$ for this choice of δ , we obtain the following strengthening of their main result (Zhang and Sra, 2018, Theorem 3):

Corollary 3.2 (Local acceleration) *Let $\delta = 1 + \frac{1}{5} \cdot (\mu/L)^{1/2}$, $\gamma = 1/L$ and $\xi_0 \geq \xi(\delta)$. Then, assuming $d(x_0, x_*) \leq \frac{1}{20} \cdot \kappa^{1/2}(\mu/L)^{3/4}$, we have $f(y_t) - f(x_*) = O(\exp(-\frac{9}{10}t\sqrt{\mu/L}))$. In particular, $\xi_t = \xi(\delta)$ for all $t \geq 0$, recovers (Zhang and Sra, 2018, Algorithm 2).*

4. Riemannian Accelerated Gradient Method

Thus far, the analysis assumed existence of valid distortion rates. Now the question is: *are valid distortion rates available to the method?* We provide a positive answer below and therewith propose a new Riemannian accelerated gradient method. For clarity, we will focus on Hadamard manifolds; the development for non-Hadamard manifolds is analogous and is provided in Appendix D.

4.1. Valid distortion rates and Riemannian accelerated gradient method

We estimate metric distortion by first invoking a classical comparison theorem of Rauch (1951).

Proposition 4.1 *Let $x, y, z \in M$, a Riemannian manifold with curvature lower bounded by $-\kappa < 0$. Let $S_\kappa(r) := (\frac{\sinh(\sqrt{\kappa}r)}{\sqrt{\kappa}r})^2$; then, we have $d(y, z)^2 \leq S_\kappa(\max\{d(x, y), d(x, z)\}) \cdot d_x(y, z)^2$.*

Proof A direct consequence of the Rauch comparison theorem; see Appendix B for completeness. ■

Proposition 4.1 applied to x_t, z_t, x_* yields $d(z_t, x_*)^2 \leq S_\kappa(\max\{d(x_t, z_t), d(x_t, x_*)\}) \cdot d_{x_t}(z_t, x_*)^2$. Next, from Topogonov's comparison theorem (see e.g. (Burago et al., 2001, Section 6.5)) we know that $d_{x_{t+1}}(z_t, x_*) \leq d(z_t, x_*)$ since M is a Hadamard manifold. Combining these two inequalities, we see that $\delta_t = S_\kappa(\max\{d(x_t, z_t), d(x_t, x_*)\})$ is a valid distortion rate. However, the issue with this argument is that the distortion rate depends on $d(x_t, x_*)$, which is in general unavailable to the algorithm. We overcome this fundamental difficulty by developing a new distortion inequality.

Lemma 4.1 (Improved metric distortion inequality) *Let x, y, z be points on Riemannian manifold M with sectional curvatures lower bounded by $-\kappa < 0$. Then for $T_\kappa : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 1}$ defined as*

$$T_\kappa(r) := \begin{cases} \max\left\{1 + 4\left(\frac{\sqrt{\kappa}r}{\tanh(\sqrt{\kappa}r)} - 1\right), \left(\frac{\sinh(2\sqrt{\kappa}r)}{2\sqrt{\kappa}r}\right)^2\right\}, & \text{if } r > 0, \\ 1, & \text{if } r = 0, \end{cases} \quad (4.1)$$

the following inequality holds: $d(y, z)^2 \leq T_\kappa(d(x, y)) \cdot d_x(y, z)^2$.

Proof The proof uses Proposition 4.1 and a Riemannian trigonometric inequality due to (Zhang and Sra, 2016, Lemma 6). See Appendix B for a formal statement and the proof. ■

Note that T_κ behaves similarly to S_κ . Most importantly, $\lim_{r \rightarrow 0+} T_\kappa(r) = 1$, implying that the effect of distortion diminishes as the distance decreases. Hence, one can essentially regard Lemma 4.1 as a version of Proposition 4.1 in which the term $\max\{d(x, y), d(x, z)\}$ is replaced with $d(x, y)$. Thanks to Lemma 4.1, now we have $T_\kappa(d(x_t, z_t))$ as a valid distortion rate, which is *accessible* to the algorithm at iteration t . Therefore, we propose the following algorithm:

Algorithm 1 (Riemannian accelerated gradient method)

Input: $x_0 = y_0 = z_0 \in M$; constant $\xi_0 > 0$; $\gamma \in (0, 2/L)$; $\Delta_\gamma := \gamma(1 - L\gamma/2)$; integer T .
for $t = 0, 1, 2, \dots, T$:
 Compute the distortion rate $\delta_{t+1} := T_\kappa(d(x_t, z_t))$ as per (4.1).
 Find $\xi_{t+1} \in [2\mu\Delta_\gamma, 1)$ such that $\frac{\xi_{t+1}(\xi_{t+1} - 2\mu\Delta_\gamma)}{1 - \xi_{t+1}} = \frac{1}{\delta_{t+1}} \cdot \xi_t^2$.
 Compute $\alpha_{t+1} := \frac{\xi_{t+1} - 2\mu\Delta_\gamma}{1 - 2\mu\Delta_\gamma}$, $\beta_{t+1} := 1 - 2\mu\Delta_\gamma \xi_{t+1}^{-1}$, and $\eta_{t+1} := 2\Delta_\gamma \xi_{t+1}^{-1}$.
 Update the next step iterates as per (3.1) with $\gamma_{t+1} := \gamma$.
end for

4.2. Convergence rate analysis of the proposed method

Having proposed the algorithm, our final task is to analyze its convergence rate. From Remark 3.3, we know the algorithm achieves a *full* acceleration when δ_t is close to 1. Due to the property $\lim_{r \rightarrow 0+} T_\kappa(r) = 1$, one therefore needs to show that $d(x_t, z_t)$ is close to 0. Although $d(x_0, z_0) = 0$, one can quickly notice that this is obviously not true for $t \geq 1$.

Now one natural follow-up question is whether $d(x_t, z_t)$ shrinks over iterations. As we have seen in §2.3, the convergence of the iterates to the optimal point is a direct consequence of our potential function analysis. Similarly, one can immediately see that $d_{x_t}(z_t, x_*) \rightarrow 0$. It turns out that from this shrinking projected distance, one can also deduce $d(x_t, z_t) \rightarrow 0$ under mild conditions:

Lemma 4.2 (Shrinking $d(x_t, z_t)$) Assume $0 < \mu$; let $D_0 := f(x_0) - f(x_*) + \frac{1}{4\Delta_\gamma} \xi_0^2 \cdot d(x_0, x_*)^2$. If $1 < \gamma L < 2 - \xi_t$ and $\xi_t > 2\mu\Delta_\gamma$ hold at iteration $t \geq 1$, then Algorithm 1 satisfies: $d(x_t, z_t) \leq \mathcal{C}_{\mu, L, \gamma} [D_0 \prod_{j=1}^{t-1} (1 - \xi_j)]^{1/2}$, where $\mathcal{C}_{\mu, L, \gamma} > 0$ is a constant depending only on μ, L, γ .

Proof The proof relies on elementary geometric inequalities (see Appendix C.6). ■

Note that the assumption $\gamma L \in (1, 2 - \xi_t]$ can be roughly read as “ $\gamma L \in (1, 2 - \sqrt{\mu/L}]$ ” because Remark 3.3 ensures that $\xi(\delta) \leq \sqrt{2\mu\Delta_\gamma} < \sqrt{\mu/L}$ for all $\delta \geq 1$. More precisely, since ξ_t quickly converges to the fixed point, one can easily ensure $\xi_t \leq \sqrt{\mu/L}$ after few iterations. Formalizing this argument, we finally obtain our main theorem (which formalizes Theorem 1.1):

Theorem 4.1 (Eventual full acceleration of Algorithm 1) Assume that $0 < \mu < L$. For any $\xi_0 > 0$, $\gamma L \in (1, 2 - \sqrt{\mu/L}]$, let $\Delta_\gamma := \gamma(1 - L\gamma/2)$ and $\lambda := 1 - \frac{8\mu\Delta_\gamma}{5 + \sqrt{5}} \in (0, 1)$. Then, Algorithm 1 satisfies the following accelerated convergence:

$$f(y_t) - f(x_*) = O((1 - \xi_1)(1 - \xi_2) \cdots (1 - \xi_t)), \quad (4.2)$$

where $\{\xi_t\}$ is a sequence such that (i) $\xi_t > 2\mu\Delta_\gamma \forall t \geq 0$ and (ii) for all $\epsilon > 0$, $|\xi_t - \sqrt{2\mu\Delta_\gamma}| \leq \epsilon$ whenever $t = \Omega\left(\frac{\log(1/\epsilon)}{\log(1/\lambda)}\right)$, where the constant involved in $\Omega(\cdot)$ depends only on μ, L, γ, κ .

Proof (4.2) is immediate from Theorem 3.1. For the convergence of $\{\xi_t\}$, see Appendix C.7. ■

Since $\Delta_\gamma \rightarrow 1/(2L)$ as $\gamma \rightarrow 1/L$, one can achieve the convergence rate *arbitrarily* close to the full acceleration rate by choosing γ bigger but sufficiently close to $1/L$. This concludes our main results.

5. Comparison with other potential function analyses

In this section, we compare existing potential function analyses with our approach. For a survey, see e.g. (Taylor and Bach, 2019, Appendix B).

Other approaches to (2.2). The potential function (2.2) has appeared in prior work on accelerated methods, corroborating its *suitability*. Compared with our analysis, the main difference is that the existing analyses either work for (i) the case $\mu = 0$, or (ii) just the fixed-step case $\xi_t = \xi(\delta)$. We highlight that our analysis is the first to recover—from first principles—Nesterov’s general scheme that smoothly interpolates the cases $\mu = 0$ and $\mu > 0$. Moreover, our analysis allows ξ_t to vary, which is crucial in the Riemannian case where the recursive relation changes over iterations.

Function (2.2) appears in (Wilson et al., 2016, Proposition 4) within the context of a continuous dynamics approach to acceleration. That work studies methods for discretizing *accelerated* ODEs derived in (Su et al., 2014; Wibisono et al., 2016) to transform the continuous dynamics into discrete methods. In that context, they show that (2.2) is a discretization of a *canonical* Lyapunov function. Another appearance is in (Diakonikolas and Orecchia, 2019), where they extend the continuous dynamics view via an *approximate duality gap technique*. Roughly, to analyze a first-order method, they consider an upper bound U_t and a lower bound L_t on the optimal value $f(x_*)$. Their analysis then proceeds by showing the gap $G_t := U_t - L_t$ diminishes with the rate α_t , i.e., $\alpha_t G_t$ is decreasing, which corresponds to showing $A_t \mathcal{E}_t$ is decreasing in our language (§2). Although motivated mostly for continuous dynamics, their techniques cover discrete methods with some modifications. In particular, their choice of G_t for accelerated method corresponds to (2.2) (see §4.2 therein).

Yet another appearance of (2.2) is (Bansal and Gupta, 2019, (5.50)), wherein the motivation was to modify the potential function analyses for gradient descent to cover accelerated methods. They propose the idea of running *two different* gradient steps and linearly combining them to achieve desired accelerated convergence. Following their argument, it turns out (2.2) is the right choice. Indeed, their approach bears resemblance to the *linear coupling* framework (Allen-Zhu and Orecchia, 2014), in which (2.2) has even more *canonical* interpretations; see Appendix A.

SDP-based approaches. The primary distinction between our approach and most SDP-based approaches (Drori and Teboulle, 2014; Lessard et al., 2016; Taylor et al., 2018; Taylor and Bach, 2019) is that our analysis is *analytical*, whereas the analyses therein are *numerical*. More specifically, the existing works require *numeric values* of parameters (e.g., α, β, L, μ) because they find suitable potential functions via solving SDPs. Note that one *cannot* solve SDPs unless the numeric coefficients are given. Abstractly, our choice of parameters in Theorem 2.1 can be interpreted as an *analytical* solution to the *symbolic* versions of SDPs formulated in the prior works.

Notable exceptions are (Kim and Fessler, 2016; Hu and Lessard, 2017; Safavi et al., 2018; Cyrus et al., 2018; Aybat et al., 2018), in which small SDPs are solved *analytically*. Specifically, some optimized step sizes for Nesterov’s method are derived via solving small SDPs explicitly in (Kim and Fessler, 2016; Safavi et al., 2018); robust versions of gradient methods are derived analytically via classical control-theoretic arguments in (Cyrus et al., 2018; Aybat et al., 2018), and Nesterov’s method is reinterpreted using *dissipativity theory* in (Hu and Lessard, 2017). Indeed borrowing the

dissipativity interpretation from (Hu and Lessard, 2017), one can interpret our calculations in §2.3 as finding a *analytic* solution to the dissipation inequality (Theorem 2 therein).

6. Conclusion

In this paper, we established the first *global* accelerated methods for non-Euclidean setting. Our analysis demonstrated that our proposed scheme is always faster than gradient descent, and achieves the accelerated convergence after few iterations. This establishment makes a solid progress towards understanding acceleration under non-Euclidean settings. Our analysis builds on the potential function analysis, a recent paradigm of analyzing first-order methods. Remarkably, our analysis for the Euclidean case *exactly* recovers Nesterov’s optimal method based on estimate sequence, shedding a new light on the scope of the technique which has puzzled researchers for many years.

References

- P-A Absil, Robert Mahony, and Rodolphe Sepulchre. *Optimization algorithms on matrix manifolds*. Princeton University Press, 2009.
- Naman Agarwal, Nicolas Boumal, Brian Bullins, and Coralia Cartis. Adaptive regularization with cubics on manifolds. *arXiv:1806.00065*, 2018.
- Foivos Alimisis, Antonio Orvieto, Gary Bécigneul, and Aurelien Lucchi. A continuous-time perspective for modeling acceleration in riemannian optimization. *To appear at AISTATS*, 2020.
- Zeyuan Allen-Zhu and Lorenzo Orecchia. Linear coupling: An ultimate unification of gradient and mirror descent. *arXiv preprint arXiv:1407.1537*, 2014.
- Necdet Serhat Aybat, Alireza Fallah, Mert Gurbuzbalaban, and Asuman Ozdaglar. Robust accelerated gradient methods for smooth strongly convex functions. *arXiv preprint arXiv:1805.10579*, 2018.
- Miroslav Bacák. *Convex analysis and optimization in Hadamard spaces*, volume 22. Walter de Gruyter GmbH & Co KG, 2014.
- Nikhil Bansal and Anupam Gupta. Potential-function proofs for gradient methods. *Theory of Computing*, 15(4):1–32, 2019. doi: 10.4086/toc.2019.v015a004.
- Glaydston C Bento, Orizon P Ferreira, and Jefferson G Melo. Iteration-complexity of gradient, subgradient and proximal point methods on riemannian manifolds. *Journal of Optimization Theory and Applications*, 173(2):548–562, 2017.
- Michael Betancourt, Michael I Jordan, and Ashia C Wilson. On symplectic optimization. *arXiv preprint arXiv:1802.03653*, 2018.
- Martin R Bridson and André Haeffliger. *Metric spaces of non-positive curvature*, volume 319. Springer Science & Business Media, 2013.
- Sébastien Bubeck. Introduction to online optimization. *Lecture Notes*, 2, 2011.
- Dmitri Burago, IU D Burago, Yuri Burago, Sergei A Ivanov, and Sergei Ivanov. *A course in metric geometry*, volume 33. American Mathematical Soc., 2001.
- Peter Bürgisser, Cole Franks, Ankit Garg, Rafael Oliveira, Michael Walter, and Avi Wigderson. Towards a theory of non-commutative optimization: geodesic 1st and 2nd order methods for

- moment maps and polytopes. In *2019 IEEE 60th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 845–861. IEEE, 2019.
- Dario Cordero-Erausquin, Robert J McCann, and Michael Schmuckenschläger. A riemannian interpolation inequality à la borell, brascamp and lieb. *Inventiones mathematicae*, 146(2):219–257, 2001.
- Saman Cyrus, Bin Hu, Bryan Van Scoy, and Laurent Lessard. A robust accelerated optimization algorithm for strongly convex functions. In *2018 Annual American Control Conference (ACC)*, pages 1376–1381. IEEE, 2018.
- Jelena Diakonikolas and Lorenzo Orecchia. The approximate duality gap technique: A unified theory of first-order methods. *SIAM Journal on Optimization*, 29(1):660–689, 2019.
- Yoel Drori and Marc Teboulle. Performance of first-order methods for smooth convex minimization: a novel approach. *Mathematical Programming*, 145(1-2):451–482, 2014.
- Ramsay Dyer, Gert Vegter, and Mathijs Wintraecken. Riemannian simplices and triangulations. *Geometriae Dedicata*, 179(1):91–138, 2015.
- Alexander Vladimirovich Gasnikov and Yu E Nesterov. Universal method for stochastic composite optimization problems. *Computational Mathematics and Mathematical Physics*, 58(1):48–64, 2018.
- Navin Goyal and Abhishek Shetty. Sampling and optimization on convex sets in riemannian manifolds of non-negative curvature. In *Conference on Learning Theory*, pages 1519–1561, 2019.
- Mikhail Gromov. Manifolds of negative curvature. *Journals of Differential Geometry*, 13(2):223–230, 1978.
- Bin Hu and Laurent Lessard. Dissipativity theory for nesterov's accelerated method. In *Proceedings of the 34th International Conference on Machine Learning-Volume 70*, pages 1549–1557. JMLR. org, 2017.
- Jürgen Jost. *Riemannian geometry and geometric analysis*, volume 42005. Springer, 2008.
- Hiroyuki Kasai, Hiroyuki Sato, and Bamdev Mishra. Riemannian stochastic variance reduced gradient on grassmann manifold. *arXiv preprint arXiv:1605.07367*, 2016.
- Donghwan Kim and Jeffrey A Fessler. Optimized first-order methods for smooth convex minimization. *Mathematical programming*, 159(1-2):81–107, 2016.
- Laurent Lessard, Benjamin Recht, and Andrew Packard. Analysis and design of optimization algorithms via integral quadratic constraints. *SIAM Journal on Optimization*, 26(1):57–95, 2016.
- Yuanyuan Liu, Fanhua Shang, James Cheng, Hong Cheng, and Licheng Jiao. Accelerated first-order methods for geodesically convex optimization on riemannian manifolds. In *Advances in Neural Information Processing Systems*, pages 4868–4877, 2017.
- Aleksandr Mikhailovich Lyapunov. The general problem of the stability of motion. *International journal of control*, 55(3):531–534, 1992.
- Arkadii Semenovich Nemirovsky and David Borisovich Yudin. Problem complexity and method efficiency in optimization. 1983.
- Yurii Nesterov. A method for unconstrained convex minimization problem with the rate of convergence $O(1/k^2)$. In *Doklady AN USSR*, volume 269, pages 543–547, 1983.

- Yurii Nesterov. *Lectures on convex optimization*, volume 137. Springer, 2018.
- Brendan O’donoghue and Emmanuel Candes. Adaptive restart for accelerated gradient schemes. *Foundations of computational mathematics*, 15(3):715–732, 2015.
- Harry Ernest Rauch. A contribution to differential geometry in the large. *Annals of Mathematics*, pages 38–55, 1951.
- Sam Safavi, Bikash Joshi, Guilherme França, and José Bento. An explicit convergence rate for nesterov’s method from sdp. In *2018 IEEE International Symposium on Information Theory (ISIT)*, pages 1560–1564. IEEE, 2018.
- Bin Shi, Simon S Du, Weijie J Su, and Michael I Jordan. Acceleration via symplectic discretization of high-resolution differential equations. *arXiv preprint arXiv:1902.03694*, 2019.
- Jonathan W Siegel. Accelerated optimization with orthogonality constraints. *arXiv preprint arXiv:1903.05204*, 2019.
- Weijie Su, Stephen Boyd, and Emmanuel Candes. A differential equation for modeling nesterov’s accelerated gradient method: Theory and insights. In *Advances in Neural Information Processing Systems*, pages 2510–2518, 2014.
- Adrien Taylor and Francis Bach. Stochastic first-order methods: non-asymptotic and computer-aided analyses via potential functions. *arXiv preprint arXiv:1902.00947*, 2019.
- Adrien Taylor, Bryan Van Scoy, and Laurent Lessard. Lyapunov functions for first-order methods: Tight automated convergence guarantees. *arXiv preprint arXiv:1803.06073*, 2018.
- Constantin Udriste. *Convex functions and optimization methods on Riemannian manifolds*, volume 297. Springer Science & Business Media, 1994.
- Melanie Weber and Suvrit Sra. Nonconvex stochastic optimization on manifolds via riemannian frank-wolfe methods. *arXiv preprint arXiv:1910.04194*, 2019.
- Andre Wibisono, Ashia C Wilson, and Michael I Jordan. A variational perspective on accelerated methods in optimization. *proceedings of the National Academy of Sciences*, 113(47):E7351–E7358, 2016.
- Ashia C Wilson, Benjamin Recht, and Michael I Jordan. A lyapunov analysis of momentum methods in optimization. *arXiv preprint arXiv:1611.02635*, 2016.
- Hongyi Zhang and Suvrit Sra. First-order methods for geodesically convex optimization. In *Conference on Learning Theory*, pages 1617–1638, 2016.
- Hongyi Zhang and Suvrit Sra. An estimate sequence for geodesically convex optimization. In *Conference On Learning Theory*, pages 1703–1723, 2018.
- Hongyi Zhang, Sashank J Reddi, and Suvrit Sra. Riemannian svrg: Fast stochastic optimization on riemannian manifolds. In *Advances in Neural Information Processing Systems*, pages 4592–4600, 2016.
- Jingzhao Zhang, Hongyi Zhang, and Suvrit Sra. R-spider: A fast riemannian stochastic optimization algorithm with curvature independent rate. *arXiv preprint arXiv:1811.04194*, 2018.
- Pan Zhou, Xiaotong Yuan, Shuicheng Yan, and Jiashi Feng. Faster first-order methods for stochastic non-convex optimization on riemannian manifolds. *IEEE transactions on pattern analysis and machine intelligence*, 2019.

Appendix A. Interpretations via linear coupling

Recently, [Allen-Zhu and Orecchia \(2014\)](#) established a framework of designing fast first-order methods called *linear coupling*. The principle observation therein is that the two most fundamental first-order methods, namely *gradient* and *mirror* descent, have *complementary* performances, and one can therefore design faster first-order methods by *linearly coupling* the two methods. In this section, we will discuss how one can derive from linear coupling (i) the Nesterov's optimal method iterations (2.1) and (ii) our choice of potential function (2.2).

A.1. Nesterov's iteration from linear coupling.

One can actually derive the iteration updated rules (2.1) from linear coupling. To illustrate, let us denote by $\text{Grad}_{s,\nabla}(x)$ and $\text{Mirr}_{s,\nabla}(x)$ a single step of the gradient and mirror step, respectively. If we choose the Bregman divergence associated with the mirror descent to be $D(u, v) = \frac{1}{2} \|u - v\|_2^2$, then (2.1) can be rewritten as follows:

$$\begin{aligned} x_{t+1} &\leftarrow \alpha_{t+1} z_t + (1 - \alpha_{t+1}) y_t \\ w_{t+1} &\leftarrow \tilde{\beta}_{t+1} z_t + (1 - \tilde{\beta}_{t+1}) y_t \\ \nabla_{t+1} &\leftarrow \nabla f(x_{t+1}) \\ y_{t+1} &\leftarrow \text{Grad}_{\gamma_{t+1} \nabla_{t+1}}(x_{t+1}) \\ z_{t+1} &\leftarrow \text{Mirr}_{\eta_{t+1} \nabla_{t+1}}(w_{t+1}), \end{aligned}$$

where $\tilde{\beta}_{t+1} = \alpha_{t+1} + (1 - \alpha_{t+1})\beta_{t+1}$. Note that these steps clearly respect linear coupling: for each step, we compute two different linear combinations of z_t and y_t and run gradient and mirror step from each combination to obtain the next iterates y_{t+1} and z_{t+1} , respectively. Indeed, the original algorithm considered in the paper ([Allen-Zhu and Orecchia, 2014](#)) chooses $\beta' \equiv 1$ and is hence a special case of the above steps. One concrete advantage of viewing (2.1) as above is that then (2.1) can be naturally generalized to other settings where the smoothness of f is defined with respect to a norm different from ℓ_2 .

A.2. Choosing potential function from linear coupling.

Another advantage of linear coupling is that one can naturally derive our choice of potential function (2.2). To see this, we first note that the folklore analysis of gradient descent deals with the cost value $f(y)$, while that of mirror descent deals with the distance to the optimal, or more generally, the Bregman divergence $D(z, x_*)$. (See e.g. ([Allen-Zhu and Orecchia, 2014](#), Section 2) for details.) Since the algorithm is a *linear combination* of the two methods, it is then natural to consider a *linear combination* of the two performance measures, arriving at (2.2) since our case corresponds to the case where the Bregman divergence is chosen as $D(z, x_*) = \frac{1}{2} \|z - x_*\|_2^2$.

Appendix B. Some inequalities from Riemannian geometry (proof of Lemma 4.1)

The main technical difficulty of analyzing optimization methods over Riemannian manifolds lies in handling its non-Euclidean metric. One machinery to overcome this difficulty is a classical comparison theorem due to [Rauch \(1951\)](#). At a high level, the theorem compares the exponential map on the manifold of interest to that on the manifold of constant sectional curvatures.

Proposition B.1 (Rauch comparison theorem) *Let M be a Riemannian manifold with sectional curvatures lower bounded by $-\kappa < 0$. Then, for any $x \in M$ and $u \in T_x M$, the following upper bound on the operator norm of the differential of the exponential map holds:*

$$\|d(\text{Exp}_x)_u\|_{\text{op}} \leq \frac{\sinh(\sqrt{\kappa} \|u\|)}{\sqrt{\kappa} \|u\|}.$$

Proof Let $u_0 := u / \|u\|$. First, it follows from the definition that the exponential map is radially isometric, i.e., $\|d(\text{Exp}_x)_u(u_0)\| = \|u_0\|$. Next, due to Rauch Comparison Theorem, for any v orthogonal to u , we have $\|v\| \leq \|d(\text{Exp}_x)_u(v)\| \leq \frac{\sinh(\sqrt{\kappa} \|u\|)}{\sqrt{\kappa} \|u\|} \|v\|$. Since any vector in $T_u(T_x M)$ can be represented as a linear combination of u_0 and vectors orthogonal to u_0 , the proof follows. ■

The above consequence of Rauch comparison theorem then implies the following metric distortion inequality, which can be seen as a global version of the ones employed in the prior arts (Dyer et al., 2015, Lemma 9) and (Zhang and Sra, 2018, Theorem 2).

Proposition B.2 (Restatement of Proposition 4.1) *Let x, y, z be points on Riemannian manifold M with sectional curvatures lower bounded by $-\kappa < 0$. Then, for the following inequality holds:*

$$d(y, z) \leq \frac{\sinh(\sqrt{\kappa} \max\{d(x, y), d(x, z)\})}{\sqrt{\kappa} \max\{d(x, y), d(x, z)\}} \cdot d_x(y, z).$$

Proof To upper bound $d(y, z)$ in terms of $d_x(y, z)$, consider a path $p : [0, 1] \rightarrow T_x M$ defined as $p(t) = (1 - t) \cdot \text{Exp}_x^{-1}(y) + t \cdot \text{Exp}_x^{-1}(z)$. Then, its image $\text{Exp}_x(p)$ is a path on M connecting y from z . By definition of the distance on the manifold, $d(y, z)$ is clearly upper bounded by the length of $\text{Exp}_x(p)$. On the other hand, using Proposition B.1, the length of $\text{Exp}_x(p)$ can be upper bounded as follows (since $\|p'(t)\| = \|\text{Exp}_x^{-1}(y) - \text{Exp}_x^{-1}(z)\| = d_x(y, z)$):

$$\begin{aligned} \int_0^1 \left\| \frac{d}{dt} \text{Exp}_x(p(t)) \right\| dt &\leq \int_0^1 \|d(\text{Exp}_x)_{p(t)}\|_{\text{op}} \cdot \|p'(t)\| dt \\ &\leq S_\kappa(\max\{d(x, y), d(x, z)\}) \cdot d_x(y, z), \end{aligned}$$

where the last inequality follows from the fact that $\|p(t)\|$ is upper bounded by $\max\{\|p(0)\|, \|p(1)\|\} = \max\{d(x, y), d(x, z)\}$. ■

Next machinery is a user-friendly global trigonometric inequality from (Zhang and Sra, 2016, Lemma 6) and (Cordero-Erausquin et al., 2001, Lemma 3.12):

Proposition B.3 (Global trigonometric inequality) *Let M be a Riemannian manifold with sectional curvatures lower bounded by $-\kappa < 0$. Let x, y, z be the vertices of a geodesic triangle with the lengths of the opposite side being a, b, c , respectively, and A be the angle of the triangle at the vertex x , then we have the following inequality:*

$$a^2 \leq \frac{\sqrt{\kappa} c}{\tanh(\sqrt{\kappa} c)} \cdot b^2 + c^2 - 2bc \cos A.$$

Proof See (Zhang and Sra, 2016, Section 3.1). ■

Now based on these two machineries, we prove the improved metric distortion inequality (Lemma 4.1):

Lemma B.1 (Formal statement of Lemma 4.1) *Let x, y, z be points on Riemannian manifold M with sectional curvatures lower bounded by $-\kappa < 0$. For function $\widehat{T}_\kappa : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 1}$ defined as*

$$\widehat{T}_\kappa(r) := \begin{cases} \min_{\epsilon > 0} \max \left\{ 1 + (1 + \epsilon^{-1})^2 \left(\frac{\sqrt{\kappa}r}{\tanh(\sqrt{\kappa}r)} - 1 \right), \left(\frac{\sinh((1+\epsilon)\sqrt{\kappa} \cdot r)}{(1+\epsilon)\sqrt{\kappa} \cdot r} \right)^2 \right\} & \text{if } r > 0, \\ 1, & \text{if } r = 0, \end{cases}$$

the following inequality holds: $d(y, z)^2 \leq \widehat{T}_\kappa(d(x, y)) \cdot d_x(y, z)^2$.

Remark B.4 (Properties of \widehat{T}_κ) *Note that $\widehat{T}_\kappa \leq T_\kappa$ since T_κ corresponds to choosing $\epsilon = 1$ for all r in \widehat{T}_κ . Hence Lemma B.1 implies Lemma 4.1. In particular, the followings are true about $\widehat{T}_\kappa(r)$:*

1. $\lim_{r \rightarrow 0+} \widehat{T}_\kappa(r) = 1$.
2. *For any $\epsilon > 0$, there exists $R_\epsilon > 0$ such that $\widehat{T}_\kappa(r) \leq \left(\frac{\sinh((1+\epsilon)\sqrt{\kappa} \cdot r)}{(1+\epsilon)\sqrt{\kappa} \cdot r} \right)^2$ for $r > R_\epsilon$.*

Hence, one can essentially regard \widehat{T}_κ as a proxy for S_κ .

Proof [Proof of Lemma B.1] Let us fix an arbitrary constant $\epsilon > 0$. We will separately handle two cases: (i) $(1 + \epsilon) \cdot d(x, y) < d(x, z)$ and (ii) $(1 + \epsilon) \cdot d(x, y) \geq d(x, z)$. Let us begin with the first case. Applying the trigonometric lemma to $\triangle xyz$, and letting $\zeta := \frac{\sqrt{\kappa}d(x, y)}{\tanh(\sqrt{\kappa}d(x, y))}$, we obtain:

$$\begin{aligned} d(y, z)^2 &\leq d(x, y)^2 + \zeta \cdot d(x, z)^2 - 2 \langle \text{Exp}_x^{-1}(y), \text{Exp}_x^{-1}(z) \rangle \\ &= (\zeta - 1) \cdot d(x, z)^2 + d(x, y)^2 + d(x, z)^2 - 2 \langle \text{Exp}_x^{-1}(y), \text{Exp}_x^{-1}(z) \rangle \\ &= (\zeta - 1) \cdot d(x, z)^2 + d_x(y, z)^2. \end{aligned}$$

where the last line follows from the Euclidean law of cosine. On the other hand, from the Euclidean triangle inequality (consider the triangle $\triangle xyz$ in the tangent space $T_x M$), $d_x(y, z) \geq (d(x, z) - d(x, y)) > \frac{\epsilon}{1+\epsilon} \cdot d(x, z)$. Hence, combining these two, we get

$$\begin{aligned} d(y, z)^2 &\leq (\zeta - 1) \cdot d(x, z)^2 + d_x(y, z)^2 \\ &\leq (1 + \epsilon^{-1})^2 \cdot (\zeta - 1) \cdot d_x(y, z)^2 + d_x(y, z)^2 \\ &= \left[1 + (1 + \epsilon^{-1})^2 \cdot (\zeta - 1) \right] \cdot d_x(y, z)^2. \end{aligned} \tag{B.1}$$

Next, for the case $(1 + \epsilon) \cdot d(x, y) \geq d(x, z)$, Proposition B.2 implies:

$$d(y, z)^2 \leq \left(\frac{\sinh((1 + \epsilon)\sqrt{\kappa} \cdot d(x, y))}{(1 + \epsilon)\sqrt{\kappa} \cdot d(x, y)} \right)^2 \cdot d_x(y, z)^2. \tag{B.2}$$

Therefore, combining (B.1) and (B.2), the proof is completed. ■

Appendix C. Proofs/calculations missing from the main text

C.1. Proofs of folklore analyses of gradient steps (Propositions 2.1 and 2.2)

Proof of Proposition 2.1: By the L -smoothness of f , we have $f(y) \leq f(x) + \langle \nabla f(x), y - x \rangle + \frac{L}{2} \|x - y\|^2 = f(x) + \langle \nabla f(x), -s \nabla f(x) \rangle + \frac{L}{2} \|-s \nabla f(x)\|^2 = f(x) - s \left(1 - \frac{Ls}{2}\right) \|\nabla f(x)\|^2$. ■

Proof of Proposition 2.2: The proof is an elementary calculation: $\|z - x_*\|^2 = \|z - x + x - x_*\|^2 = \|z - x\|^2 + \|x - x_*\|^2 + 2 \langle z - x, x - x_* \rangle = s^2 \|\nabla f(x)\|^2 + \|x - x_*\|^2 + 2s \langle \nabla f(x), x_* - x \rangle$. ■

C.2. Derivation of the upper bound on the potential difference (2.6)

First, one can easily express (2.4) in terms of ∇, X, W using Proposition 2.2:

$$\begin{aligned}
 (2.4) &= B_{t+1} \cdot \|z_{t+1} - x_*\|^2 - B_t \cdot \|z_t - x_*\|^2 \\
 &= B_{t+1} \cdot \|X + \beta W\|^2 + B_{t+1} \eta^2 \cdot \|\nabla\|^2 - 2B_{t+1} \eta \cdot \langle \nabla, X + \beta W \rangle - B_t \cdot \|W + X\|^2 \\
 &= (B_{t+1} - B_t) \cdot \|X\|^2 + (\beta^2 B_{t+1} - B_t) \cdot \|W\|^2 + \eta^2 B_{t+1} \cdot \|\nabla\|^2 \\
 &\quad + 2(\beta B_{t+1} - B_t) \cdot \langle X, W \rangle - 2\beta \eta B_{t+1} \langle W, \nabla \rangle - 2\eta B_{t+1} \cdot \langle X, \nabla \rangle.
 \end{aligned} \tag{C.1}$$

For (2.3), we apply Proposition 2.1 and rearrange terms to obtain:

$$\begin{aligned}
 (2.3) &\leq A_{t+1} \cdot (f(x_{t+1}) - f(x_*)) - \Delta_\gamma A_{t+1} \cdot \|\nabla\|^2 - A_t \cdot (f(y_t) - f(x_*)) \\
 &= A_t \cdot (f(x_{t+1}) - f(y_t)) + (A_{t+1} - A_t) \cdot (f(x_{t+1}) - f(x_*)) - \Delta_\gamma A_{t+1} \cdot \|\nabla\|^2.
 \end{aligned}$$

Now using the inequality $f(u) - f(v) \leq \langle \nabla f(u), u - v \rangle - \frac{\mu}{2} \|u - v\|^2$ for all u, v ($\because \mu$ -strong convexity of f) together with the identity $x_{t+1} - y_t = \frac{\alpha}{1-\alpha}(z_t - x_{t+1}) = \frac{\alpha}{1-\alpha}W$, one can also derive an upper bound on (2.3) in terms of ∇, W, X :

$$\begin{aligned}
 (2.3) &\leq \frac{\alpha}{1-\alpha} A_t \cdot \langle \nabla, W \rangle - \frac{\mu}{2} \left(\frac{\alpha}{1-\alpha} \right)^2 A_t \cdot \|W\|^2 \\
 &\quad + (A_{t+1} - A_t) \cdot \langle \nabla, X \rangle - \frac{\mu}{2} (A_{t+1} - A_t) \cdot \|X\|^2 - \Delta_\gamma A_{t+1} \cdot \|\nabla\|^2 \\
 &= -\frac{\mu}{2} \left(\frac{\alpha}{1-\alpha} \right)^2 A_t \cdot \|W\|^2 - \frac{\mu}{2} (A_{t+1} - A_t) \cdot \|X\|^2 - \Delta_\gamma A_{t+1} \cdot \|\nabla\|^2 \\
 &\quad + \frac{\alpha}{1-\alpha} A_t \cdot \langle W, \nabla \rangle + (A_{t+1} - A_t) \cdot \langle X, \nabla \rangle.
 \end{aligned} \tag{C.2}$$

Putting (C.1) and (C.2) together, we obtain (2.6).

C.3. Proof of Theorem 2.1

From the equality version of (2.8), i.e., $B_{t+1} = (A_{t+1} - A_t)^2 / (4\Delta_\gamma \cdot A_{t+1})$, one can easily express B_{t+1} in terms of ξ_{t+1} . More specifically, using the relation $1 - \xi_{t+1} := A_t / A_{t+1}$, we have $(\frac{A_{t+1} - A_t}{A_{t+1}})^2 = \xi_{t+1}^2$, and hence:

$$B_{t+1} = \left(\frac{A_{t+1} - A_t}{A_{t+1}} \right)^2 \cdot \frac{A_{t+1}}{4\Delta_\gamma} = \frac{\xi_{t+1}^2}{1 - \xi_{t+1}} \cdot \frac{A_t}{4\Delta_\gamma}. \tag{C.3}$$

From this, one can also conclude that

$$\frac{A_{t+1}}{B_{t+1}} = \frac{A_t}{(1 - \xi_{t+1})} \cdot \frac{1 - \xi_{t+1}}{\xi_{t+1}^2} \cdot \frac{4\Delta_\gamma}{A_t} = \frac{4\Delta_\gamma}{\xi_{t+1}^2}. \quad (\text{C.4})$$

Now from (2.7), (2.11), (C.3) and (C.4), one can express $\alpha_{t+1}, \beta_{t+1}, \eta_{t+1}$ in terms of ξ_{t+1} :

$$\begin{aligned} \eta_{t+1} &\stackrel{(2.7)}{=} \frac{A_{t+1} - A_t}{2B_{t+1}} = \frac{A_{t+1} - A_t}{A_{t+1}} \cdot \frac{A_{t+1}}{2B_{t+1}} \stackrel{(2.7) \& (C.4)}{=} \xi_{t+1} \cdot \frac{2\Delta_\gamma}{\xi_{t+1}^2} = 2\Delta_\gamma \xi_{t+1}^{-1}, \\ \beta_{t+1} &\stackrel{(2.7)}{=} \frac{B_t}{B_{t+1}} \stackrel{(C.3)}{=} \frac{1 - \xi_{t+1}}{\xi_{t+1}^2} \cdot 4\Delta_\gamma \cdot \frac{B_t}{A_t} \stackrel{(2.11)}{=} \frac{\xi_{t+1} - 2\mu\Delta_\gamma}{\xi_{t+1}} = 1 - 2\mu\Delta_\gamma \xi_{t+1}^{-1}, \quad \text{and} \\ \frac{\alpha_{t+1}}{1 - \alpha_{t+1}} &\stackrel{(2.7)}{=} \frac{(A_{t+1} - A_t)B_t}{A_t B_{t+1}} = \frac{A_{t+1} - A_t}{A_{t+1}} \cdot \frac{B_t}{A_t} \cdot \frac{A_{t+1}}{B_{t+1}} \\ &\stackrel{(2.11) \& (C.4)}{=} \xi_{t+1} \cdot \frac{\xi_{t+1}(\xi_{t+1} - 2\mu\Delta_\gamma)}{4\Delta_\gamma(1 - \xi_{t+1})} \cdot \frac{4\Delta_\gamma}{\xi_{t+1}^2} = \frac{\xi_{t+1} - 2\mu\Delta_\gamma}{1 - \xi_{t+1}}. \end{aligned}$$

With the above choices of parameters, one can easily check that $\alpha_{t+1}, \beta_{t+1}$ both lie in $[0, 1]$ since $\xi_{t+1} \in [2\mu\Delta_\gamma, 1)$. In particular, $\beta_{t+1}^2 B_{t+1} \leq \beta_{t+1} B_{t+1} = B_t$, implying $C_1 \leq 0$. Therefore, the above choices of parameters satisfy $C_1, C_2, C_3 \leq 0$ and $C_4, C_5, C_6 = 0$, and consequently, $\Phi_{t+1} \leq \Phi_t$ since $\Phi_{t+1} - \Phi_t \leq (2.6)$. This completes the proof of Theorem 2.1.

C.4. Analysis of recursive relations ((2.13) and (3.4))

For simplicity, we replace $2\mu\Delta_\gamma$ with a constant $a \in (0, 1)$ and consider:

$$\frac{\xi_{t+1}(\xi_{t+1} - a)}{1 - \xi_{t+1}} = \frac{1}{\delta} \cdot \xi_t^2. \quad (\text{C.5})$$

In particular, $\delta = 1$ and $a = 2\mu\Delta_\gamma$ recovers (2.13). Below, we state and prove a formal statement of Lemma 2.1 which demonstrates the geometric convergence of the ξ_t to the fixed point.

Lemma C.1 *For any constants $\delta \geq 1$, $a \in (0, 1)$, and an initial value $\xi_0 \geq 0$, Then the followings properties are true about (C.5):*

1. $\xi(\delta) := (\sqrt{(\delta - 1)^2 + 4\delta a} - (\delta - 1))/2$ is the unique fixed point of (C.5).
2. $\lim_{t \rightarrow \infty} \xi_t \searrow \xi(\delta)$ if $\xi_0 > \xi(\delta)$; $\xi_t \equiv \xi(\delta)$ if $\xi_0 = \xi(\delta)$; and $\lim_{t \rightarrow \infty} \xi_t \nearrow \sqrt{2\mu\Delta_\gamma}$ if $0 \leq \xi_0 < \xi(\delta)$.
3. $|\xi_t - \xi(\delta)| \leq \left(\frac{1}{\sqrt{\delta}} \left(1 - \frac{4}{5 + \sqrt{5}} \cdot \frac{a}{\sqrt{\delta}} \right) \right)^t |\xi_0 - \xi(\delta)|$, i.e., the convergences are geometrical.

Proof Define $\phi(v) := \frac{v(v-a)}{1-v}$ and $\psi(v) := \frac{1}{\delta} v^2$. Then, using these notations, (C.5) becomes

$$\phi(\xi_{t+1}) = \psi(\xi_t). \quad (\text{C.6})$$

Now, in order to understand (C.6), let us study the properties of the two functions. First, it is straightforward to check that ψ is increasing on $\mathbb{R}_{\geq 0}$ and ϕ is increasing on $[a, 1)$ with $\phi(a) = 0$ and $\lim_{v \rightarrow 1-} \phi(v) = \infty$. Indeed, ϕ is increasing since $\frac{d}{dv} \phi(v) = \frac{1-a}{(1-v)^2} - 1 \geq \frac{1}{1-a} - 1 > 0$.

Hence, one can consider the inverse of the restriction $\phi|_{[a,1]}$. We will write the inverse as ϕ^{-1} . Letting $\tau := \phi^{-1} \circ \psi$, (C.6) can be written as the standard recursive relation form

$$\xi_{t+1} = \tau(\xi_t). \quad (\text{C.7})$$

Note that $\tau : \mathbb{R}_{\geq 0} \rightarrow [a, 1]$, and hence, $\xi_t \in (a, 1]$ for $t \geq 1$. Solving $\phi(v) = \psi(v)$ on $v \in [a, 1]$ yields $v = \xi(\delta)$. Hence, $\xi(\delta)$ is the unique fixed point of (C.7). From this, together with the fact that both ϕ and ψ are increasing, we have $\phi < \psi$ for $x \in [a, \xi(\delta))$, and $\phi > \psi$ for $x \in (\xi(\delta), 1)$. Hence, $\{\xi_t\}$ is increasing if $\xi_0 \in [0, \xi(\delta))$ and decreasing if $\xi_0 > \xi(\delta)$.

Now we prove the geometric convergence of (C.7) to $\xi(\delta)$. To that end, let us first compute τ :

$$\tau(v) = \phi^{-1}(\psi(v)) = \phi^{-1}(v^2/\delta) = \frac{1}{2} \left(\sqrt{(v^2/\delta - a)^2 + 4v^2/\delta} - (v^2/\delta - a) \right).$$

By mean value theorem, the key to showing geometric convergences of (C.7) is to bound the derivative of τ . In particular, if we can establish that $|\tau'(v)| \leq K < 1$ for $v \in [a, 1]$, then we have

$$|\xi_{t+1} - \xi(\delta)| = |\tau(\xi_t) - \tau(\xi(\delta))| \leq K \cdot |\xi_t - \xi(\delta)|. \quad (\text{C.8})$$

Defining $\theta(v) = \frac{v(v^2-a)+2v}{\sqrt{(v^2-a)^2+4v^2}} - v$, one can express τ in terms of θ :

$$\begin{aligned} \tau'(v) &= \frac{\frac{v}{\delta}(v^2/\delta - a) + 2\frac{v}{\delta}}{\sqrt{(v^2/\delta - a)^2 + 4v^2/\delta}} - \frac{v}{\delta} = \frac{1}{\sqrt{\delta}} \cdot \left(\frac{\frac{v}{\sqrt{\delta}}(v^2/\delta - a) + 2\frac{v}{\sqrt{\delta}}}{\sqrt{(v^2/\delta - a)^2 + 4v^2/\delta}} - \frac{v}{\sqrt{\delta}} \right) \\ &= \frac{1}{\sqrt{\delta}} \cdot \theta(v/\sqrt{\delta}) \end{aligned}$$

Hence, it suffices to show that $\theta(v) < 1$ for $v \in (0, 1)$.

Proposition C.1 *The following inequality holds for $v \in (0, 1)$: $0 \leq \theta(v) < 1 - \frac{4}{5+\sqrt{5}} \cdot v$.*

Proof $\theta(v) \geq 0$ trivially holds since, τ is a composition of increasing functions and hence increasing. Now let us prove the upper bound. We first consider the case $a < v \leq \sqrt{a}$. Since $v^2 \leq a$, we have

$$\theta(v) = \frac{-v(a - v^2) + 2v}{\sqrt{(v^2 - a)^2 + 4v^2}} - v \leq \frac{2v}{\sqrt{(v^2 - a)^2 + 4v^2}} - v \leq 1 - v.$$

Next, consider the case $v > \sqrt{a}$. Then, $v^2 > a$, and hence

$$\begin{aligned} \theta(v) &= \frac{v(v^2 - a) + 2v}{\sqrt{(v^2 - a)^2 + 4v^2}} - v = \frac{2v}{\sqrt{(v^2 - a)^2 + 4v^2}} - v \cdot \frac{\sqrt{(v^2 - a)^2 + 4v^2} - (v^2 - a)}{\sqrt{(v^2 - a)^2 + 4v^2}} \\ &= \frac{2v}{\sqrt{(v^2 - a)^2 + 4v^2}} - v \cdot \frac{4v^2}{(v^2 - a)^2 + 4v^2 + (v^2 - a)\sqrt{(v^2 - a)^2 + 4v^2}} \\ &\stackrel{(\clubsuit)}{\leq} 1 - v \cdot \frac{4v^2}{v^2 + 4v^2 + v\sqrt{v^2 + 4v^2}} = 1 - \frac{4}{5 + \sqrt{5}} \cdot v. \end{aligned}$$

where (\clubsuit) follows since $v \in (\sqrt{a}, 1)$; in particular, it follows that $0 \leq v^2 - a \leq v^2 \leq v$. Combining the two cases, we complete the proof. \blacksquare

From Proposition C.1 and (C.8), the geometric convergence follows. \blacksquare

C.5. Proof of Theorem 3.1

We first introduce the definitions of geodesical smoothness and (strong) convexity for completeness. For simplicity, we assume that the function $f : M \rightarrow \mathbb{R}$ is differentiable throughout the definitions, and following the notation for the Euclidean case, we will denote by $\nabla f(x)$ the gradient of f at x .

Definition C.2 (Geodesical (strong) convexity) f is said to be geodesically μ -strongly convex if

$$f(y) \geq f(x) + \langle \nabla f(x), \text{Exp}_x^{-1}(y) \rangle_x + \frac{\mu}{2} d(x, y)^2 \quad \text{for any } x, y \in M,$$

where $\langle \cdot, \cdot \rangle_x$ denotes the inner product in the tangent space of x induced by the Riemannian metric.

Definition C.3 (Geodesical smoothness) $f : M \rightarrow \mathbb{R}$ is said to be geodesically L -smooth if

$$f(y) \leq f(x) + \langle \nabla f(x), \text{Exp}_x^{-1}(y) \rangle_x + \frac{L}{2} d(x, y)^2 \quad \text{for any } x, y \in M.$$

An Equivalent definition is

$$\|\nabla f(x) - \Gamma_y^x \nabla f(y)\|_x \leq L \cdot d(x, y) \quad \text{for any } x, y \in M,$$

where Γ_y^x is the parallel transport from y to x .

Now with these definitions, one can establish analogues of Propositions 2.1 and 2.2:

Proposition C.4 Let $y = \text{Exp}_x(-s \cdot \nabla f(x))$. If f is geodesically L -smooth, then $f(y) - f(x) \leq -s(1 - Ls/2) \|\nabla f(x)\|_x^2$.

Proof Identical to that of Proposition 2.1. ■

Proposition C.5 Let $z = \text{Exp}_u(v - s \cdot \nabla f(u))$ for some vector $v \in T_x M$. Then, for any x_* , $d_u(z, x_*)^2 - d_u(\text{Exp}_u(v), x_*)^2 = s^2 \|\nabla f(u)\|_u^2 + 2s \langle \nabla f(u), \text{Exp}_u^{-1}(x_*) - v \rangle_u$.

Proof The proof follows immediately from the definition of the projected distances (Definition 3.1):

$$\begin{aligned} d_u(z, x_*)^2 &= \|\text{Exp}_u^{-1}(z) - \text{Exp}_u^{-1}(x_*)\|_u^2 \\ &= \|v - s \cdot \nabla f(u) - \text{Exp}_u^{-1}(x_*)\|_u^2 \\ &= \|v - \text{Exp}_u^{-1}(x_*)\|_u^2 + \|-s \cdot \nabla f(u)\|_u^2 + 2 \langle -s \cdot \nabla f(u), v - \text{Exp}_u^{-1}(x_*) \rangle_u \\ &= d_u(\text{Exp}_u(v), x_*)^2 + s^2 \|\nabla f(u)\|_u^2 + 2 \langle \nabla f(u), \text{Exp}_u^{-1}(x_*) - v \rangle_u. \end{aligned} \quad \blacksquare$$

Now we prove Theorem 3.1. It turns out one can establish an upper bound of the potential difference $\Psi_{t+1} - \Psi_t$ similar to (2.6). Here the difference is that instead of W, X, ∇ , we now have the following three vectors in the same tangent space $T_x M$:

$$\widetilde{W} := \text{Exp}_{x_{t+1}}^{-1}(z_t), \quad \widetilde{X} := -\text{Exp}_{x_{t+1}}^{-1}(x_*), \quad \text{and} \quad \widetilde{\nabla} := \text{Grad} f(x_{t+1}). \quad (\text{C.9})$$

As pointed out in §2.2, the fact that the three vectors are in the same tangent space (since they are all rooted at x_{t+1}) is crucial for the analysis to follow. With these vectors and Propositions C.4 and C.5, one can derive the following upper bound similarly to Appendix C.2 (here $\|\cdot\|$ denotes $\|\cdot\|_{x_{t+1}}$):

$$\begin{aligned} & \tilde{C}_1 \cdot \|\tilde{W}\|^2 + \tilde{C}_2 \cdot \|\tilde{X}\|^2 + \tilde{C}_3 \|\tilde{\nabla}\|^2 + \tilde{C}_4 \cdot \langle \tilde{W}, \tilde{X} \rangle + \tilde{C}_5 \cdot \langle \tilde{W}, \tilde{\nabla} \rangle + \tilde{C}_6 \cdot \langle \tilde{X}, \tilde{\nabla} \rangle, \quad (\text{C.10}) \\ \text{where } & \begin{cases} \tilde{C}_1 := \beta_{t+1}^2 B_{t+1} - \frac{B_t}{\delta_{t+1}} - \frac{\mu}{2} \frac{\alpha_{t+1}^2}{(1-\alpha_{t+1})^2} A_t, & \tilde{C}_2 := B_{t+1} - \frac{B_t}{\delta_{t+1}} - \frac{\mu}{2} (A_{t+1} - A_t), \\ \tilde{C}_3 := \eta_{t+1}^2 B_{t+1} - \Delta_\gamma \cdot A_{t+1}, & \tilde{C}_4 := 2 \cdot \left(\beta_{t+1} B_{t+1} - \frac{B_t}{\delta_{t+1}} \right), \\ \tilde{C}_5 := \frac{1-\alpha}{\alpha} A_t - 2\beta_{t+1} \eta_{t+1} B_{t+1}, \quad \text{and} & \tilde{C}_6 := (A_{t+1} - A_t) - 2\eta_{t+1} B_{t+1}. \end{cases} \end{aligned}$$

Notice the similarity between (C.10) and (2.6): the only difference is that B_t 's in (2.6) are replaced with B_t/δ_{t+1} 's. Indeed, this difference is precisely attributed to the definition of valid distortion rate (Definition 3.2). More specifically, in the derivation of (C.10), we use $-B_t \cdot d_{x_t}(z_t, x_*)^2 \leq -\frac{B_t}{\delta_t} \cdot d_{x_{t+1}}(z_t, x_*)^2$, which precisely accounts for the appearance of B_t/δ_{t+1} instead of B_t :

We now follow Section 2.3 to make (C.10) “–SoS.” First, from $\tilde{C}_4 = \tilde{C}_5 = \tilde{C}_6 = 0$, we get:

$$\eta_{t+1} = \frac{A_{t+1} - A_t}{2B_{t+1}}, \quad \beta_{t+1} = \frac{B_t}{\delta_{t+1} B_{t+1}}, \quad \text{and} \quad \frac{\alpha_{t+1}}{1 - \alpha_{t+1}} = \frac{(A_{t+1} - A_t)B_t}{\delta_{t+1} A_t B_{t+1}}. \quad (\text{C.11})$$

Moving on, one can similarly obtain the following from $\tilde{C}_3 \leq 0$ and $\tilde{C}_2 \leq 0$:

$$\frac{(A_{t+1} - A_t)^2}{4\Delta_\gamma \cdot A_{t+1}} \leq B_{t+1} \quad \text{and} \quad (\text{C.12})$$

$$\frac{(A_{t+1} - A_t)^2}{4\Delta_\gamma \cdot A_{t+1}} - (A_{t+1} - A_t) \frac{\mu}{2} \leq \frac{B_t}{\delta_{t+1}}. \quad (\text{C.13})$$

Again, using the suboptimality shrinking ratio $1 - \xi_{t+1} := A_t/A_{t+1}$, (C.13) becomes

$$\frac{\xi_{t+1}(\xi_{t+1} - 2\mu\Delta_\gamma)}{1 - \xi_{t+1}} \leq \frac{4\Delta_\gamma}{\delta_{t+1}} \cdot \frac{B_t}{A_t}. \quad (\text{C.14})$$

Again, due to the LHS being an increasing function on $\xi_{t+1} \in [2\mu\Delta_\gamma, 1)$, the largest ξ_{t+1} (or equivalently, the largest A_{t+1}) satisfies (C.14) (or equivalently, (C.13)) with equality:

$$\frac{\xi_{t+1}(\xi_{t+1} - 2\mu\Delta_\gamma)}{1 - \xi_{t+1}} = \frac{4\Delta_\gamma}{\delta_{t+1}} \cdot \frac{B_t}{A_t}. \quad (\text{C.15})$$

Consequently, such choice of ξ_{t+1} (or corresponding A_{t+1}) also satisfies (C.12) with equality.

Now one can follow the calculations in Appendix C.3 to express parameters in terms of ξ_{t+1} . From the equality version of (C.12), one can easily deduce the identical relations as (C.3) and (C.4). Now using (C.3), (C.4), (C.11), and (C.15), one can express $\alpha_{t+1}, \beta_{t+1}, \eta_{t+1}$ in terms of ξ_{t+1} . It turns out that the final expressions do not depend on δ_{t+1} and are *identical* to Appendix C.3:

$$\begin{aligned} \eta_{t+1} & \stackrel{(\text{C.11})}{=} \frac{A_{t+1} - A_t}{2B_{t+1}} = \frac{A_{t+1} - A_t}{A_{t+1}} \cdot \frac{A_{t+1}}{2B_{t+1}} \stackrel{(\text{C.11})\&(\text{C.4})}{=} \xi_{t+1} \cdot \frac{2\Delta_\gamma}{\xi_{t+1}^2} = 2\Delta_\gamma \xi_{t+1}^{-1}, \\ \beta_{t+1} & \stackrel{(\text{C.11})}{=} \frac{B_t}{\delta_{t+1} B_{t+1}} \stackrel{(\text{C.3})}{=} \frac{1 - \xi_{t+1}}{\xi_{t+1}^2} \cdot \frac{4\Delta_\gamma}{\delta_{t+1}} \cdot \frac{B_t}{A_t} \stackrel{(\text{C.15})}{=} \frac{\xi_{t+1} - 2\mu\Delta_\gamma}{\xi_{t+1}} = 1 - 2\mu\Delta_\gamma \xi_{t+1}^{-1}, \quad \text{and} \\ \frac{\alpha_{t+1}}{1 - \alpha_{t+1}} & \stackrel{(\text{C.11})}{=} \frac{(A_{t+1} - A_t)B_t}{\delta_{t+1} A_t B_{t+1}} = \frac{A_{t+1} - A_t}{A_{t+1}} \cdot \frac{B_t}{\delta_{t+1} A_t} \cdot \frac{A_{t+1}}{B_{t+1}} \\ & \stackrel{(\text{C.15})\&(\text{C.4})}{=} \xi_{t+1} \cdot \frac{\xi_{t+1}(\xi_{t+1} - 2\mu\Delta_\gamma)}{4\Delta_\gamma(1 - \xi_{t+1})} \cdot \frac{4\Delta_\gamma}{\xi_{t+1}^2} = \frac{\xi_{t+1} - 2\mu\Delta_\gamma}{1 - \xi_{t+1}}. \end{aligned}$$

With the above choices of parameters, one can again check that $\alpha_{t+1}, \beta_{t+1}$ both lie in $[0, 1]$ since $\xi_{t+1} \in [2\mu\Delta_\gamma, 1)$. In particular, $\beta_{t+1}^2 B_{t+1} \leq \beta_{t+1} B_{t+1} = B_t / \delta_{t+1}$, implying $\tilde{C}_1 \leq 0$. Therefore, the above choices of parameters satisfy $\tilde{C}_1, \tilde{C}_2, \tilde{C}_3 \leq 0$ and $\tilde{C}_4, \tilde{C}_5, \tilde{C}_6 = 0$, and consequently, $\Psi_{t+1} \leq \Psi_t$ since $\Psi_{t+1} - \Psi_t \leq (\text{C.10})$. This completes the proof of Theorem 3.1.

C.6. Proofs of distance shrinking results (Lemma 4.2)

We first show the following result which is a direct consequence of Theorem 3.1:

Proposition C.6 *Assume that $\mu > 0$ and $\gamma \in (0, 2/L)$. Let $D_0 := f(x_0) - f(x_*) + \frac{1}{4\Delta_\gamma} \xi_0^2 \cdot d(x_0, x_*)^2$. Then, x_t, y_t, z_t generated from Algorithm 1 satisfy the following distance bounds:*

1. $d_{x_t}(z_t, x_*) \leq \sqrt{D_0 \prod_{j=1}^t (1 - \xi_j)} \cdot \sqrt{\frac{1}{\mu^2 \Delta_\gamma}}.$
2. $d(y_t, x_*) \leq \sqrt{D_0 \prod_{j=1}^t (1 - \xi_j)} \cdot \sqrt{\frac{2}{\mu}}.$
3. $d_{x_t}(y_t, z_t) \leq \sqrt{D_0 \prod_{j=1}^t (1 - \xi_j)} \cdot \left(\sqrt{\frac{2}{\mu}} + \sqrt{\frac{1}{\mu^2 \Delta_\gamma}} \right).$

Proof By recursively applying Theorem 3.1, we have the following for any $t \geq 1$:

$$\begin{aligned} f(y_t) - f(x_*) + \frac{1}{4\Delta_\gamma} \xi_t^2 \cdot d_{x_t}(z_t, x_*)^2 &\leq \prod_{j=1}^t (1 - \xi_j) \cdot \left[f(y_0) - f(x_*) + \frac{1}{4\Delta_\gamma} \xi_0^2 \cdot d_{x_0}(z_0, x_*)^2 \right] \\ &= \prod_{j=1}^t (1 - \xi_j) \cdot D_0, \end{aligned}$$

where the equality follows since $x_0 = y_0 = z_0$ (which implies $d_{x_0}(z_0, x_*) = d(x_0, x_*)$).

Hence, the bound on $d_{x_t}(z_t, x_*)$ follows immediately due to $\xi_t \in [2\mu\Delta_\gamma, 1)$. As for the bound on $d(y_t, x_*)$, it follows from the μ -strong g-convexity of f (Definition C.2), which implies $\frac{\mu}{2} \cdot d(y_t, x_*)^2 \leq f(y_t) - f(x_*)$. Lastly, the bound on $d_{x_t}(y_t, z_t)$ follows from

$$d_{x_t}(y_t, z_t) \leq d_{x_t}(y_t, x_*) + d_{x_t}(z_t, x_*) \leq d(y_t, x_*) + d_{x_t}(z_t, x_*)$$

which is a consequence of the (Euclidean) triangle inequality together with the fact that the projected distances are shorter than the actual distances (a property of Hadamard manifolds; see e.g. (Burago et al., 2001, Section 6.5)). \blacksquare

From Proposition C.6, we established that the projected distance $d_{x_t}(y_t, z_t)$ is shrinking over iterations. Now one can also demonstrate that $d(y_t, z_t)$ is shrinking under mild conditions, and that is what the next proposition is about.

Proposition C.7 *Let $D_0 := f(x_0) - f(x_*) + \frac{1}{4\Delta_\gamma} \xi_0^2 \cdot d(x_0, x_*)^2$. If $\gamma L > 1$, $\gamma L \leq 2 - \xi_{t+1}$ and $\xi_{t+1} > 2\mu\Delta_\gamma$ hold for $t \geq 0$, then Algorithm 1 satisfies:*

$$d(y_t, z_t) \leq \frac{\sqrt{D_0 \prod_{j=1}^t (1 - \xi_j)}}{1 - 2\mu\Delta_\gamma \xi_{t+1}^{-1}} \cdot \frac{\left(\sqrt{\frac{2}{\mu}} + \sqrt{\frac{1}{\mu^2 \Delta_\gamma}} + \frac{L}{\mu} \sqrt{\frac{2}{\mu}} \right) (1 - 2\mu\Delta_\gamma)}{(\gamma L - 1)(\gamma L - 1 + 2\mu\Delta_\gamma)}.$$

Remark C.8 A careful reader might realize that the appearance of the term $1 - 2\mu\Delta_\gamma\xi_{t+1}^{-1}$ in the denominator of the bound could be potentially problematic as this could be arbitrarily small in general when ξ_{t+1} is very close to $2\mu\Delta_\gamma$. However, the appearance of this term will not be problematic for our purpose since the term will be canceled out with the algorithm parameter $\beta_{t+1} = 1 - 2\mu\Delta_\gamma\xi_{t+1}^{-1}$ (see Algorithm 1) when we use Proposition C.7 to bound $d(x_t, z_t)$.

Proof First, from (3.1b) and (3.1c) together with the triangle inequality,

$$\begin{aligned} d_{x_{t+1}}(y_{t+1}, z_{t+1}) &= \left\| -\gamma \nabla f(x_{t+1}) - \beta_{t+1} \text{Exp}_{x_{t+1}}^{-1}(z_t) + \eta_{t+1} \nabla f(x_{t+1}) \right\|_{x_{t+1}} \\ &\geq \beta_{t+1} \left\| \text{Exp}_{x_{t+1}}^{-1}(z_t) \right\|_{x_{t+1}} - |\eta_{t+1} - \gamma| \cdot \|\nabla f(x_{t+1})\|_{x_{t+1}} \\ &= \beta_{t+1} \cdot d(x_{t+1}, z_t) - |\eta_{t+1} - \gamma| \cdot \|\nabla f(x_{t+1})\|. \end{aligned}$$

From (3.1a), we have $d(x_{t+1}, z_t) = (1 - \alpha_{t+1}) \cdot d(y_t, z_t)$. Hence, the above inequality becomes

$$\begin{aligned} \beta_{t+1}(1 - \alpha_{t+1}) \cdot d(y_t, z_t) &\leq d_{x_{t+1}}(y_{t+1}, z_{t+1}) + |\eta_{t+1} - \gamma| \cdot \|\nabla f(x_{t+1})\| \\ &\leq d_{x_{t+1}}(y_{t+1}, z_{t+1}) + L|\eta_{t+1} - \gamma| \cdot d(x_{t+1}, x_*), \end{aligned}$$

where the last inequality follows from the geodesically L -smoothness of f , which implies $\|\nabla f(x_{t+1})\| \leq L \cdot d(x_{t+1}, x_*)$. Due to the Riemannian triangle inequality, we have $d(x_{t+1}, x_*) \leq d(x_{t+1}, y_t) + d(y_t, x_*)$, and hence the upper bound becomes

$$\begin{aligned} &d_{x_{t+1}}(y_{t+1}, z_{t+1}) + L|\eta_{t+1} - \gamma| \cdot d(x_{t+1}, y_t) + L|\eta_{t+1} - \gamma| \cdot d(y_t, x_*), \\ &= d_{x_{t+1}}(y_{t+1}, z_{t+1}) + L\alpha_{t+1}|\eta_{t+1} - \gamma| \cdot d(y_t, z_t) + L|\eta_{t+1} - \gamma| \cdot d(y_t, x_*), \end{aligned}$$

where the second line follows from the identity $d(x_{t+1}, y_t) = \alpha_{t+1} \cdot d(y_t, z_t)$ (\because (3.1a)). Moving $L\alpha_{t+1}|\eta_{t+1} - \gamma| \cdot d(y_t, z_t)$ term to the LHS, we then obtain:

$$\mathcal{K} \cdot d(y_t, z_t) \leq d_{x_{t+1}}(y_{t+1}, z_{t+1}) + L|\eta_{t+1} - \gamma| \cdot d(y_t, x_*), \quad (\text{C.16})$$

where $\mathcal{K} := \beta_{t+1}(1 - \alpha_{t+1}) - L\alpha_{t+1}|\eta_{t+1} - \gamma|$.

Having established (C.16), it is straightforward to see that Proposition C.7 is a direct consequence the following two statements:

1. The RHS is upper bounded by $\sqrt{D_0 \prod_{j=1}^t (1 - \xi_j)} \cdot \left(\sqrt{\frac{2}{\mu}} + \sqrt{\frac{1}{\mu^2 \Delta_\gamma}} + \frac{L}{\mu} \sqrt{\frac{2}{\mu}} \right)$.
2. $\mathcal{K} \geq \frac{1 - 2\mu\Delta_\gamma\xi_{t+1}^{-1}}{1 - 2\mu\Delta_\gamma} \cdot (\gamma L - 1)(\gamma L - 1 + 2\mu\Delta_\gamma)$, where the lower bound is always positive since $\gamma L > 1$ together with the fact that $1 - 2\mu\Delta_\gamma\xi_{t+1}^{-1} > 1 - 2\mu\Delta_\gamma \cdot (2\mu\Delta_\gamma)^{-1} = 0$.

Hence, the proof is completed as soon as we prove the above two statements.

We first prove the first statement. From Proposition C.6, we have

- $d_{x_{t+1}}(y_{t+1}, z_{t+1}) \leq \sqrt{D_0 \prod_{j=1}^{t+1} (1 - \xi_j)} \cdot \left(\sqrt{\frac{2}{\mu}} + \sqrt{\frac{1}{\mu^2 \Delta_\gamma}} \right)$
- $L|\eta_{t+1} - \gamma| \cdot d(y_t, x_*) \leq L|\eta_{t+1} - \gamma| \cdot \sqrt{D_0 \prod_{j=1}^t (1 - \xi_j)} \cdot \sqrt{\frac{2}{\mu}}.$

Moreover, since $\eta_{t+1} = 2\Delta_\gamma \xi_{t+1}^{-1} = \gamma(2 - \gamma L)\xi_{t+1}^{-1}$, it is straightforward from the assumption $\gamma L \leq 2 - \xi_{t+1}$ that $\eta_{t+1} \geq \gamma$. Hence, using inequalities (i) $1 - \xi_{t+1} \leq 1$ and (ii) $L|\eta_{t+1} - \gamma| \leq L\eta_{t+1} = 2L\Delta_\gamma \xi_{t+1}^{-1} \leq \frac{L}{\mu}$, we obtain the first statement.

Now, let us prove the second statement. From the parameter choices in Algorithm 1 together with the fact $\eta_{t+1} \geq \gamma$ we have established above, one can simplify and lower bound \mathcal{K} as follows:

$$\begin{aligned} \mathcal{K} &= (1 - 2\mu\Delta_\gamma \xi_{t+1}^{-1}) \frac{1 - \xi_{t+1}}{1 - 2\mu\Delta_\gamma} - L \frac{\xi_{t+1} - 2\mu\Delta_\gamma}{1 - 2\mu\Delta_\gamma} (2\Delta_\gamma \xi_{t+1}^{-1} - \gamma) \\ &= \frac{1 - 2\mu\Delta_\gamma \xi_{t+1}^{-1}}{1 - 2\mu\Delta_\gamma} \cdot [1 - \xi_{t+1} - 2L\Delta_\gamma + \gamma L \xi_{t+1}] \\ &= \frac{1 - 2\mu\Delta_\gamma \xi_{t+1}^{-1}}{1 - 2\mu\Delta_\gamma} \cdot [(\gamma L - 1)^2 + (\gamma L - 1)\xi_{t+1}] \\ &> \frac{1 - 2\mu\Delta_\gamma \xi_{t+1}^{-1}}{1 - 2\mu\Delta_\gamma} \cdot [(\gamma L - 1)^2 + (\gamma L - 1) \cdot 2\mu\Delta_\gamma] . \end{aligned}$$

where the last line follows from the fact $\xi_{t+1} > 2\mu\Delta_\gamma$ and $\gamma L - 1 > 0$. ■

Now we are ready to provide the formal statement and the proof of Lemma 4.2:

Lemma C.2 (Formal statement of Lemma 4.2) *Assume that $\mu > 0$. Let $D_0 := f(x_0) - f(x_*) + \frac{1}{4\Delta_\gamma} \xi_0^2 \cdot d(x_0, x_*)^2$. If $\gamma L > 1$, $\gamma L \leq 2 - \xi_{t+1}$ and $\xi_{t+1} > 2\mu\Delta_\gamma$ hold, then Algorithm 1 satisfies:*

$$d(x_{t+1}, z_{t+1}) \leq \mathcal{C}_{\mu, L, \gamma} \cdot \sqrt{D_0 \prod_{j=1}^t (1 - \xi_j)},$$

where $\mathcal{C}_{\mu, L, \gamma} = \frac{\left(\sqrt{\frac{2}{\mu}} + \sqrt{\frac{1}{\mu^2 \Delta_\gamma}} + \frac{L}{\mu} \sqrt{\frac{2}{\mu}}\right)(2L\Delta_\gamma + 1 - 2\mu\Delta_\gamma)}{(\gamma L - 1)(\gamma L - 1 + 2\mu\Delta_\gamma)} + \frac{L}{\mu} \sqrt{\frac{2}{\mu}}.$

Proof From (3.1c), one can use the Euclidean triangle inequality on $T_{x_{t+1}}M$ to obtain:

$$\begin{aligned} d(x_{t+1}, z_{t+1}) &\leq \beta_{t+1} \cdot d(x_{t+1}, z_t) + \eta_{t+1} \cdot \|\nabla f(x_{t+1})\|_{x_{t+1}} \\ &\stackrel{(\clubsuit)}{\leq} \beta_{t+1} \cdot d(x_{t+1}, z_t) + L\eta_{t+1} \cdot d(x_{t+1}, x_*) \\ &\stackrel{(\spadesuit)}{\leq} \beta_{t+1} \cdot d(x_{t+1}, z_t) + L\eta_{t+1} \cdot d(x_{t+1}, y_t) + L\eta_{t+1} \cdot d(y_t, x_*) \\ &\stackrel{(\heartsuit)}{=} (\beta_{t+1}(1 - \alpha_{t+1}) + L\eta_{t+1}\alpha_{t+1}) \cdot d(y_t, z_t) + L\eta_{t+1} \cdot d(y_t, x_*) \\ &\stackrel{(\diamondsuit)}{=} \frac{2L\Delta_\gamma + 1 - \xi_{t+1}}{1 - 2\mu\Delta_\gamma} (1 - 2\mu\Delta_\gamma \xi_{t+1}^{-1}) \cdot d(y_t, z_t) + 2L\Delta_\gamma \xi_{t+1}^{-1} \cdot d(y_t, x_*) , \end{aligned}$$

where (\clubsuit) is due to the geodesically L -smoothness of f , which implies $\|\nabla f(x_{t+1})\| \leq L \cdot d(x_{t+1}, x_*)$; (\spadesuit) is due to Riemannian triangle inequality; (\heartsuit) is due to (3.1a); and (\diamondsuit) follows from the choice of parameters in Algorithm 1.

Now after we apply Propositions C.6 and C.7 to the last upper bound, and use the fact $\xi_{t+1} \in [2\mu\Delta_\gamma, 1)$ to upper bound ξ_{t+1} 's in the resulting upper bound, Lemma C.2 readily follows. ■

C.7. Proof of Theorem 4.1

We first demonstrate that regardless of what initial value $\xi_0 \geq 0$ we choose, ξ_t becomes less than $\sqrt{\mu/L}$ after few iterations. Before the demonstration, we denote by $\xi_{t+1} = \tau_{t+1}(\xi_t)$ the relation with which $\{\xi_t\}$ is produced in Algorithm 1. In other words, $\xi_{t+1} = \tau_{t+1}(\xi_t)$ is equivalent to (3.4) with $\delta = \delta_{t+1}$.

Proposition C.9 *If $\xi_0 > \sqrt{\mu/L}$, then $\xi_t \leq \sqrt{\mu/L}$ for all t greater than or equal to*

$$\frac{\log \left((\xi_0 - \sqrt{2\mu\Delta_\gamma}) / (\sqrt{\mu/L} - \sqrt{2\mu\Delta_\gamma}) \right)}{\log \left(1 / \left(1 - \frac{8\mu\Delta_\gamma}{5+\sqrt{5}} \right) \right)}. \quad (\text{C.17})$$

If $\xi_0 < \sqrt{\mu/L}$, then $\xi_t \leq \sqrt{\mu/L}$ for all $t \geq 0$.

Proof At some iteration t , we consider the two cases depending on whether $\xi_t \leq \sqrt{2\mu\Delta_\gamma}$ or not:

1. First, if $\xi_t \leq \sqrt{2\mu\Delta_\gamma}$, then we have $\xi_{t'} \leq \sqrt{2\mu\Delta_\gamma}$ for all $t' \geq t$ because $\xi(\delta_{t'})$ is less than $\sqrt{2\mu\Delta_\gamma}$ for all t' and the relation at each step, $\xi_{t'+1} = \tau_{t'+1}(\xi_{t'})$, will only bring $\xi_{t'}$ closer to $\xi(\delta_{t'})$ due to Lemma C.1.
2. Now consider the case $\xi_t > \sqrt{2\mu\Delta_\gamma}$. We may assume that $\xi_{t+1} > \sqrt{2\mu\Delta_\gamma}$ (otherwise, $\xi_{t'} \leq \sqrt{2\mu\Delta_\gamma}$ for $t' \geq t+1$ due to the first case). Then, the mean value theorem implies:

$$\begin{aligned} \xi_{t+1} - \sqrt{2\mu\Delta_\gamma} &= \tau_{t+1}(\xi_t) - \tau_{t+1}(\tau_{t+1}^{-1}(\sqrt{2\mu\Delta_\gamma})) \\ &\stackrel{(\clubsuit)}{\leq} \frac{1}{\sqrt{\delta_{t+1}}} \left(1 - \frac{4}{5+\sqrt{5}} \cdot \frac{2\mu\Delta_\gamma}{\sqrt{\delta_{t+1}}} \right) \cdot \left(\xi_t - \tau_{t+1}^{-1}(\sqrt{2\mu\Delta_\gamma}) \right) \\ &\stackrel{(\spadesuit)}{<} \left(1 - \frac{4}{5+\sqrt{5}} \cdot 2\mu\Delta_\gamma \right) \cdot \left(\xi_t - \sqrt{2\mu\Delta_\gamma} \right), \end{aligned}$$

where (\clubsuit) is due to Proposition C.1 together with $\xi_{t+1} > \sqrt{2\mu\Delta_\gamma} \Rightarrow \xi_t > \tau_{t+1}^{-1}(\sqrt{2\mu\Delta_\gamma})$; (\spadesuit) follows since the maximum of $\frac{1}{\sqrt{\delta}}(1 - \frac{4}{5+\sqrt{5}} \frac{2\mu\Delta_\gamma}{\sqrt{\delta}})$ is achieved by $\delta = 1$ and $\sqrt{2\mu\Delta_\gamma} < \tau_{t+1}^{-1}(\sqrt{2\mu\Delta_\gamma})$ due to $\sqrt{2\mu\Delta_\gamma} \geq \xi(\delta_{t+1})$ and Lemma C.1. Hence, the distance between ξ_t and $\sqrt{2\mu\Delta_\gamma}$ shrinks geometrically.

Combining the two cases, we conclude the proof. ■

We now study the rate of convergence of $\{\xi_t\}$. To that end, we first study the convergence of $\{\xi(\delta_t)\}$. For simplicity, we assume that $\xi_0 \leq \sqrt{\mu/L}$. By Proposition C.9, the arguments below remain true for $\xi_0 > \sqrt{\mu/L}$ after we substitute $t \leftarrow t + (\text{C.17})$. We first characterize the behaviour of $\xi(\delta)$ near $\delta = 1$:

Proposition C.10 *Let $\xi(\delta) := \frac{1}{2} \left(\sqrt{(\delta-1)^2 + 8\delta\mu\Delta_\gamma} - (\delta-1) \right)$ for $\delta \geq 1$. Then, $0 \leq \sqrt{2\mu\Delta_\gamma} - \xi(\delta) \leq \frac{1}{2}(\delta-1)$ for $1 \leq \delta \leq 1 + 3/(1 + (4\mu\Delta_\gamma)^{-1})$.*

Proof For simplicity, let us write $\delta = 1 + d$. Then, $\xi(1 + d) = \frac{1}{2} \left(\sqrt{d^2 + 8\mu\Delta_\gamma(1 + d)} - d \right)$. Using the inequality $\sqrt{1 + r} \geq 1 + \frac{1}{3}r$ for $0 \leq r \leq 3$, we get the following as long as $d + \frac{1}{8\mu\Delta_\gamma}d^2 \leq 3$:

$$\begin{aligned} \xi(1 + d) &\geq \sqrt{2\mu\Delta_\gamma} \cdot \left(1 + \frac{1}{3}d + \frac{1}{24\mu\Delta_\gamma}d^2 \right) - \frac{1}{2}d \\ &\geq \sqrt{2\mu\Delta_\gamma} - \left(\frac{1}{2} - \frac{\sqrt{2\mu\Delta_\gamma}}{3} \right) d. \end{aligned}$$

Now all we need to check is that $d \leq 3/(1 + \frac{1}{4\mu\Delta_\gamma})$ implies $d + \frac{1}{8\mu\Delta_\gamma}d^2 \leq 3$. Indeed, if $d \leq 3/(1 + \frac{1}{4\mu\Delta_\gamma})$, then we have $d \leq 3/(1 + \frac{1}{4\mu\Delta_\gamma}) \leq 3/(3/2) = 2$, and hence $d + \frac{1}{8\mu\Delta_\gamma}d^2 = d \left(1 + \frac{d}{8\mu\Delta_\gamma} \right) \leq d \left(1 + \frac{1}{4\mu\Delta_\gamma} \right) \leq 3$. \blacksquare

Next, we characterize the behaviour of the function $T_\kappa(r)$ near $r = 1$.

Proposition C.11 $T_\kappa(r) \leq 1 + 2\kappa r^2$ for $0 \leq r \leq \frac{1}{2\sqrt{\kappa}}$.

Proof Using Taylor expansion, one easily verify for $0 \leq r \leq \frac{1}{2\sqrt{\kappa}}$ that

$$\frac{\sqrt{\kappa}r}{\tanh(\sqrt{\kappa}r)} \leq 1 + \frac{\kappa}{2}r^2 \quad \text{and} \quad \left(\frac{\sinh(2\sqrt{\kappa}r)}{2\sqrt{\kappa}r} \right)^2 \leq 1 + 2\kappa r^2.$$

Hence, from the definition of T_κ (see (4.1)), we obtain the desired bound on T_κ . \blacksquare

Combining Propositions C.10 and C.11, we obtain the following results:

Proposition C.12 $\sqrt{2\mu\Delta_\gamma} - \xi(T_\kappa(r)) \leq \kappa r^2$ for $0 \leq r \leq \sqrt{\frac{3}{1+(4\mu\Delta_\gamma)^{-1}}} \cdot \frac{1}{2\sqrt{\kappa}}$.

Proof Note that $\frac{3}{1+(4\mu\Delta_\gamma)^{-1}} \leq \frac{3}{1+2L/\mu} \leq 1$, and hence, $\sqrt{\frac{3}{1+(4\mu\Delta_\gamma)^{-1}}} \cdot \frac{1}{2\sqrt{\kappa}} \leq \frac{1}{2\sqrt{\kappa}}$. Thus, one can apply Proposition C.11 for $0 \leq r \leq \sqrt{\frac{3}{1+(4\mu\Delta_\gamma)^{-1}}} \cdot \frac{1}{2\sqrt{\kappa}}$, and obtain $T_\kappa(r) \leq 1 + 2\kappa r^2$. Hence, $T_\kappa(r) \leq 1 + \frac{1}{2} \cdot \frac{3}{1+(4\mu\Delta_\gamma)^{-1}}$ within the range. Hence, by Proposition C.10, one then obtains $\sqrt{2\mu\Delta_\gamma} - \xi(T_\kappa(r)) \leq \kappa r^2$ for $0 \leq r \leq \sqrt{\frac{3}{1+(4\mu\Delta_\gamma)^{-1}}} \cdot \frac{1}{2\sqrt{\kappa}}$. \blacksquare

Now let $\mathcal{D}_{\kappa,\mu,\gamma} := \sqrt{\frac{3}{1+(4\mu\Delta_\gamma)^{-1}}} \cdot \frac{1}{2\sqrt{\kappa}}$. Then by Lemma C.2, one can deduce that $d(x_{t+1}, z_{t+1}) \leq \mathcal{D}_{\kappa,\mu,\gamma}$ whenever $t \geq 2 \frac{\log(\mathcal{C}_{\mu,L,\gamma} \cdot \sqrt{D_0}/\mathcal{D}_{\kappa,\mu,\gamma})}{-\log(1-2\mu\Delta_\gamma)}$. Therefore, Proposition C.12 implies that for $t \geq 2 \frac{\log(\mathcal{C}_{\mu,L,\gamma} \cdot \sqrt{D_0}/\mathcal{D}_{\kappa,\mu,\gamma})}{-\log(1-2\mu\Delta_\gamma)}$, the following bound holds:

$$\sqrt{2\mu\Delta_\gamma} - \xi(T_\kappa(d(x_{t+1}, z_{t+1}))) \leq \kappa \mathcal{C}_{\mu,L,\gamma}^2 D_0 (1 - 2\mu\Delta_\gamma)^t.$$

From this, it follows that $\xi(T_\kappa(d(x_{t+1}, z_{t+1}))) \in [\sqrt{2\mu\Delta_\gamma} - \epsilon/2, \sqrt{2\mu\Delta_\gamma}]$ whenever

$$t \geq \max \left\{ 2 \frac{\log(\mathcal{C}_{\mu,L,\gamma} \cdot \sqrt{D_0}/\mathcal{D}_{\kappa,\mu,\gamma})}{-\log(1-2\mu\Delta_\gamma)}, \frac{\log(2\kappa \mathcal{C}_{\mu,L,\gamma}^2 D_0/\epsilon)}{-\log(1-2\mu\Delta_\gamma)} \right\}.$$

Now having established the convergence of $\{\xi(\delta_t)\}$, we translate it into the convergence of $\{\xi_t\}$. Similarly to the proof of Proposition C.9, one can prove that for any $T \geq 0$,

$$|\xi_{T+t} - \xi(\delta_T)| \leq \left(1 - \frac{8\mu\Delta_\gamma}{5 + \sqrt{5}}\right)^t |\xi_T - \xi(\delta_T)|.$$

From this, one can conclude that $\xi_{t+1} \in [\sqrt{2\mu\Delta_\gamma} - \epsilon, \sqrt{2\mu\Delta_\gamma}]$ whenever

$$t \geq \max \left\{ 2 \frac{\log(\mathcal{C}_{\mu,L,\gamma} \cdot \sqrt{D_0}/\mathcal{D}_{\kappa,\mu,\gamma})}{-\log(1 - 2\mu\Delta_\gamma)}, \frac{\log(2\kappa\mathcal{C}_{\mu,L,\gamma}^2 D_0/\epsilon)}{-\log(1 - 2\mu\Delta_\gamma)} \right\} + \frac{\log(2\sqrt{2\mu\Delta_\gamma}/\epsilon)}{-\log\left(1 - \frac{8\mu\Delta_\gamma}{5+\sqrt{5}}\right)},$$

concluding the proof of the convergence rate of $\{\xi_t\}$ in Theorem 4.1.

C.8. Justification of Remark 3.3

In this section, we verify that $\xi(\delta)$ is decreasing. Note that for $\delta \geq 1$ we have

$$\frac{d}{d\delta} \xi(\delta) = \frac{2(\delta - 1) + 8\mu\Delta_\gamma}{4\sqrt{(\delta - 1)^2 + 8\mu\Delta_\gamma\delta}} - \frac{1}{2} = \frac{2(\delta - 1) + 8\mu\Delta_\gamma - 2\sqrt{(\delta - 1)^2 + 8\mu\Delta_\gamma\delta}}{4\sqrt{(\delta - 1)^2 + 8\mu\Delta_\gamma\delta}} < 0,$$

where the last inequality is due to the fact that $((\delta - 1) + 4\mu\Delta_\gamma)^2 = (\delta - 1)^2 + 8\mu\Delta_\gamma(\delta - 1) + 16\mu^2\Delta_\gamma^2 < (\delta - 1)^2 + 8\mu\Delta_\gamma\delta - 8\mu\Delta_\gamma(1 - 2\mu\Delta_\gamma) < (\delta - 1)^2 + 8\mu\Delta_\gamma\delta$ since $2\mu\Delta_\gamma < 1$.

Appendix D. Extension to the non-Hadamard case

Let us now assume that the sectional curvatures of M is upper bounded by $\sigma \geq 0$. In particular, $\sigma = 0$ corresponds to the Hadamard case. We first pinpoint the main differences: Unlike the Hadamard case, M now may not be uniquely geodesic. Instead, one can only guarantee the property within a local neighborhood of M . Consequently, the notion of convexity can be only guaranteed within a local neighborhood of M . For instance, manifolds with positive sectional curvatures (e.g. spheres) are compact, and hence, they do not admit globally geodesically convex functions other than the constant function. Following the prior arts (Dyer et al., 2015; Zhang and Sra, 2018), we make the following assumptions to avoid any further complications:

Assumption 3 *The domain $N \subset M$ of f is uniquely geodesic with the diameter bounded by $\frac{\pi}{2\sqrt{\sigma}}$.*

Assumption 4 *All the iterates of (3.1) (whose parameters to be chosen later) remain in N .*

The analysis for the non-Hadamard case is identical to that for the Hadamard case, modulo one additional geometric inequality due to (Zhang and Sra, 2018):

Proposition D.1 ((Zhang and Sra, 2018, Lemma 7)) *Let x, y, z be points on Riemannian manifold M with sectional curvatures upper bounded by $\sigma \geq 0$. If $d(x, z) \leq \frac{\pi}{2\sigma}$, then*

$$d_x(y, z)^2 \leq (1 + 2 \cdot d(x, y)^2) \cdot d(y, z)^2.$$

Applying Proposition D.1 to Lemma B.1, we obtain the following metric distortion inequality:

Lemma D.1 (Modification of Lemma B.1) *Let x, x', y, z be points on Riemannian manifold M with sectional curvatures upper and lower bounded by σ and $-\kappa < 0$, respectively. For $\widehat{T}_\kappa : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 1}$ defined as in Lemma B.1, we have*

$$d_{x'}(y, z)^2 \leq \widehat{T}_\kappa(d(x, y)) \cdot (1 + 2 \cdot d(x', y)^2) \cdot d_x(y, z)^2.$$

From Lemma D.1, one can conclude that at iteration t ,

$$T_\kappa d(x_t, z_t) \cdot (1 + 2 \cdot d(y_t, z_t)^2) \tag{D.1}$$

is a valid distortion rate. Thus, one can use (D.1) in lieu of $T_\kappa(d(x_t, z_t))$ for the valid distortion rate in Algorithm 1. Then the rest proceeds in the exactly same manner; in particular, one can similarly establish the distance shrinking results as in Appendix C.6 to corroborate Theorem 4.1 for this case. We skip the details since they significantly overlap with the Hadamard case.