

Every finite abelian group is a subgroup of the additive group of a finite simple left brace*

F. Cedó

E. Jespers

J. Okniński

Abstract

Left braces, introduced by Rump, have turned out to provide an important tool in the study of set theoretic solutions of the quantum Yang-Baxter equation. In particular, they have allowed to construct several new families of solutions. A left brace $(B, +, \cdot)$ is a structure determined by two group structures on a set B : an abelian group $(B, +)$ and a group (B, \cdot) , satisfying certain compatibility conditions. The main result of this paper shows that every finite abelian group A is a subgroup of the additive group of a finite simple left brace B with metabelian multiplicative group with abelian Sylow subgroups. This result complements earlier unexpected results of the authors on an abundance of finite simple left braces.

1 Introduction

In order to investigate a question posed by Drinfeld [9], Rump, in [11], introduced a new algebraic structure, called a left brace, to study non-degenerate involutive set-theoretic solutions of the Yang-Baxter equation. Using an equivalent formulation, given in [7], a left brace is a set B equipped with two operations $+$ and \cdot such that $(B, +)$ is an abelian group, (B, \cdot) is a group and $a \cdot (b + c) + a = a \cdot b + a \cdot c$, for all $a, b, c \in B$. In [2] it was shown that all non-degenerate involutive solutions on a finite set can be explicitly determined from finite left braces. Hence the study of left braces becomes essential to classify all solutions. Moreover, intriguingly, braces have shown up in several areas of mathematics (see for example the surveys [6, 12]). One of the fundamental problems is to classify the building blocks of all finite simple left braces, that is describe all finite simple left braces. It is well-known that the additive Sylow p -subgroup B_p of a finite brace B is a left subbrace of B and, in [3], it has been shown that B is an iterated matched product of all B_p . Rump, in [11], has shown that if $B = B_p$, then B is a simple left brace precisely when B has order p ; more generally, the conclusion holds if (B, \cdot) is a finite nilpotent group. In particular, such a brace is trivial, i.e. the two operations $+$ and \cdot coincide. Recall that Etingof, Schedler and Soloviev [10] have shown that if B is finite, then (B, \cdot) is a solvable group. Also if $B_p \neq B$, there are some natural constraints on the order of a finite simple left brace B . For example, Smoktunowicz, in [13], showed that if $|B| = p^n q^m$ (with p and q different prime numbers and n, m positive integers), then $p|(q^t - 1)$ and $q|(p^s - 1)$ for some $0 < t \leq m$ and $0 < s \leq n$. However, these conditions are not sufficient for simplicity of B . For example, there is no simple left brace of order $p^n q$, where p and q are distinct primes and n is the multiplicative order of p in the unit group of $\mathbb{Z}/(q)$ (see [3, Remark 5.3]). Bachiller, in [1, Theorem 6.3 and Section 7], produced the first example of a non-trivial finite simple left brace. This initiated a program of constructing and describing finite simple left braces [3, 4]. One of the structural approaches is via the matched product of the Sylow subgroups; this allowed to construct new classes of examples and provided some necessary conditions for simplicity. In [8], it has been proven that there is an abundance of finite simple left braces, indeed, for any positive integer $n > 1$

*The first author was partially supported by the grants MINECO-FEDER MTM2017-83487-P and AGAUR 2017SGR1725 (Spain). The second author is supported in part by Onderzoeksraad of Vrije Universiteit Brussel and Fonds voor Wetenschappelijk Onderzoek (Belgium). The third author is supported by the National Science Centre grant. 2016/23/B/ST1/01045 (Poland). 2010 MSC: Primary 16T25, 20F16, 20E22. Keywords: Yang-Baxter equation, set-theoretic solution, brace, simple, A-group.

and distinct prime numbers p_1, p_2, \dots, p_n , there exist positive integers l_1, l_2, \dots, l_n , such that, for each n -tuple of integers $m_1 \geq l_1, m_2 \geq l_2, \dots, m_n \geq l_n$, there exists a simple left brace of order $p^{m_1} p^{m_2} \dots p^{m_n}$ that has a metabelian multiplicative group with abelian Sylow subgroups. The construction of these simple braces is via asymmetric products, as introduced by Catino, Colazzo and Stefanelli in [5]. This not only provided constructions of new classes of simple left braces, but also all previously known constructions have been interpreted as asymmetric products. Furthermore, in [4], a construction is given of finite simple left braces with a multiplicative group (B, \cdot) that is solvable of arbitrary derived length. In this paper, we focus on the additive group $(B, +)$ and we discover new examples of finite simple left braces. Our main result reads as follows:

For every finite abelian group A there exists a finite simple left brace B such that A is a subgroup of the additive group of B .

In fact, the simple left braces B that we construct have metabelian multiplicative groups (B, \cdot) and have abelian Sylow subgroups (so called A-groups [14]), and additive group isomorphic to

$$\prod_{i \in \mathbb{Z}/(m)} (\mathbb{Z}/(p_i^{n_i}))^{2s_i l_{i-1} + 1},$$

where $m > 1$, p_1, \dots, p_m are any distinct prime numbers, $n_1, \dots, n_m, s_1, \dots, s_m$ are any positive integers, and $l_i = p_i^{n_i} - p_i^{n_i-1}$. In particular, we thus also construct examples that have elements of additive order 2^n (for any $n > 1$). None of the known previous constructions included such elements.

2 Preliminaries

A left brace is a set B with two binary operations, $+$ and \cdot , such that $(B, +)$ is an abelian group, (B, \cdot) is a group, and for every $a, b, c \in B$,

$$a \cdot (b + c) + a = a \cdot b + a \cdot c.$$

In any left brace B there is an action $\lambda: (B, \cdot) \rightarrow \text{Aut}(B, +)$, called the lambda map of B , defined by $\lambda(a) = \lambda_a$ and $\lambda_a(b) = a \cdot b - a$, for $a, b \in B$. A trivial brace is a left brace B such that $ab = a + b$, for all $a, b \in B$, i.e. all $\lambda_a = \text{id}$.

A left ideal of a left brace B is a subgroup L of the additive group of B such that $\lambda_a(b) \in L$, for all $b \in L$ and all $a \in B$. An ideal of a left brace B is a normal subgroup I of the multiplicative group of B such that $\lambda_a(b) \in I$, for all $b \in I$ and all $a \in B$. Note that $ab^{-1} = a - \lambda_{ab^{-1}}(b)$ and $a - b = a\lambda_{a^{-1}b}(b^{-1})$, for all $a, b \in B$. Hence, every left ideal L of B also is a subgroup of the multiplicative group of B , and every ideal I of a left brace B also is a subgroup of the additive group of B , and then B/I is a left brace, the quotient brace B modulo I . A non-zero left brace B is simple if $\{0\}$ and B are the only ideals of B .

The following result is useful to prove simplicity in many cases.

Lemma 2.1 (*Lemma 2.5 in [3]*) *If I is an ideal of a left brace B , then $(\lambda_b - \text{id})(a) \in I$, for all $a \in B$ and $b \in I$.*

In [5] Catino, Colazzo and Stefanelli introduced the asymmetric product of two left braces. Let S and T be two (additive) abelian groups. Recall that a (normalized) symmetric 2-cocycle on T with values in S is a map $b: T \times T \rightarrow S$ such that

- (i) $b(0, 0) = 0$;
- (ii) $b(t_1, t_2) = b(t_2, t_1)$;
- (iii) $b(t_1 + t_2, t_3) + b(t_1, t_2) = b(t_1, t_2 + t_3) + b(t_2, t_3)$,

for all $t_1, t_2, t_3 \in T$.

Theorem 2.2 (Catino, Colazzo, Stefanelli [5]) Let T and S be two left braces. Let $b: T \times T \longrightarrow S$ be a symmetric 2-cocycle on $(T, +)$ with values in $(S, +)$, and let $\alpha: (S, \cdot) \longrightarrow \text{Aut}(T, +, \cdot)$ be a homomorphism of groups such that

$$s \cdot b(t_2, t_3) + b(t_1 \cdot \alpha_s(t_2 + t_3), t_1) = b(t_1 \cdot \alpha_s(t_2), t_1 \cdot \alpha_s(t_3)) + s,$$

where $\alpha_s = \alpha(s)$, for all $s \in S$ and $t_1, t_2, t_3 \in T$. Then the addition and multiplication on $T \times S$ given by

$$\begin{aligned} (t_1, s_1) + (t_2, s_2) &= (t_1 + t_2, s_1 + s_2 + b(t_1, t_2)), \\ (t_1, s_1) \cdot (t_2, s_2) &= (t_1 \cdot \alpha_{s_1}(t_2), s_1 \cdot s_2), \end{aligned}$$

define a structure of left brace on $T \times S$, called the asymmetric product of T by S (via b and α). It is denoted as $T \rtimes_\circ S$.

Note that, in $T \rtimes_\circ S$,

$$(t_1, s_1) - (t_2, s_2) = (t_1 - t_2, s_1 - s_2 - b(t_1 - t_2, t_2)),$$

for all $t_1, t_2 \in T$ and $s_1, s_2 \in S$. Thus the lambda map of $T \rtimes_\circ S$ is defined by

$$\lambda_{(t_1, s_1)}(t_2, s_2) = (\lambda_{t_1} \alpha_{s_1}(t_2), \lambda_{s_1}(s_2) - b(\lambda_{t_1} \alpha_{s_1}(t_2), t_1)). \quad (1)$$

Note that every symmetric bi-additive map $b: T \times T \longrightarrow S$ is a symmetric 2-cocycle on $(T, +)$ with values in $(S, +)$. In this case, the condition on b and α is equivalent to the following two conditions:

$$\begin{aligned} \lambda_s(b(t_2, t_3)) &= b(\alpha_s(t_2), \alpha_s(t_3)), \\ b(t_2, t_3) &= b(\lambda_{t_1}(t_2), \lambda_{t_1}(t_3)). \end{aligned}$$

In the construction below T and S will be trivial braces. Hence, these two conditions are reduced to

$$b(t_2, t_3) = b(\alpha_s(t_2), \alpha_s(t_3)). \quad (2)$$

Moreover, in this case

$$\lambda_{(t_1, s_1)}(t_2, s_2) = (\alpha_{s_1}(t_2), s_2 - b(\alpha_{s_1}(t_2), t_1)).$$

Let A, B be abelian additive groups. Let $b: A \times A \longrightarrow B$ be a symmetric bi-additive map. We define an addition on $A \times B$ by

$$(x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2 + b(x_1, x_2)),$$

for all $x_1, x_2 \in A$ and $y_1, y_2 \in B$. We denote by $G_b(A, B)$ the abelian group $A \times B$ with the above addition. Note that if b is the zero map 0, then $G_0(A, B)$ is just the direct product of the groups A and B .

Lemma 2.3 If B is finite of odd order, then $G_b(A, B) \cong G_0(A, B)$ for every symmetric bi-additive map $b: A \times A \longrightarrow B$.

Proof. Let $|B| = 2n + 1$. We define $\varphi_b: G_b(A, B) \longrightarrow G_0(A, B)$ by $\varphi_b(x, y) = (x, y + nb(x, x))$, for all $x \in A$ and $y \in B$. It is clear that φ is bijective. For $x_1, x_2 \in A$ and $y_1, y_2 \in B$, we have that

$$\begin{aligned} \varphi_b((x_1, y_1) + (x_2, y_2)) &= \varphi_b(x_1 + x_2, y_1 + y_2 + b(x_1, x_2)) \\ &= (x_1 + x_2, y_1 + y_2 + b(x_1, x_2) + nb(x_1 + x_2, x_1 + x_2)) \\ &= (x_1 + x_2, y_1 + y_2 + b(x_1, x_2) + nb(x_1, x_1) + nb(x_2, x_2) + 2nb(x_1, x_2)) \\ &= (x_1 + x_2, y_1 + y_2 + nb(x_1, x_1) + nb(x_2, x_2)) \\ &= \varphi_b(x_1, y_1) + \varphi_b(x_2, y_2). \end{aligned}$$

Thus the result follows. \blacksquare

The result is not true for B of even order. For example, let $A = B = \mathbb{Z}/(2)$ and let $b: A \times A \longrightarrow B$ be the map defined by $b(x, y) = xy$. Note that in $G_b(A, B)$ we have that $(1, 0) + (1, 0) = (0, 1)$. Then $G_b(A, B) \cong \mathbb{Z}/(4) \not\cong \mathbb{Z}/(2) \times \mathbb{Z}/(2) = G_0(A, B)$.

3 The construction

Let $m > 1$ be an integer. For $i \in \mathbb{Z}/(m)$, let p_i be a prime number and let n_i and s_i be positive integers. We assume that $p_i \neq p_j$ for $i \neq j$. Consider the rings $R_i = \mathbb{Z}/(p_i^{n_i})$ and the polynomials

$$q_i(x) = \sum_{k=0}^{p_i-1} x^{kp_i^{n_i-1}} \in R_{i+1}[x],$$

for $i \in \mathbb{Z}/(m)$. Note that $(x^{p_i^{n_i-1}} - 1)q_i(x) = x^{p_i^{n_i}} - 1$. Put $l_i = \deg(q_i(x))$. Let $C_i \in M_{l_{i-1}}(R_i)$ be the companion matrix of $q_{i-1}(x)$. Note that C_i is invertible in $M_{l_{i-1}}(R_i)$ and has order $p_i^{n_{i-1}}$. Let $T_i = R_i^{2l_{i-1}}$. We define the symmetric bilinear form $b_i: T_i \times T_i \rightarrow R_i$ by

$$b_i(u, v) = u \left(\begin{array}{c|c} 0 & I_{l_{i-1}} \\ \hline I_{l_{i-1}} & 0 \end{array} \right) v^t,$$

for all $u, v \in T_i$. We also define $f_i \in \text{Aut}(T_i)$ by

$$f_i(u) = u \left(\begin{array}{c|c} C_i^t & 0 \\ \hline 0 & C_i^{-1} \end{array} \right),$$

for all $u \in T_i$. One can check that $f_i \in O(T_i, b_i)$, the orthogonal group of b_i . This is a consequence of the fact that

$$\left(\begin{array}{c|c} C_i^t & 0 \\ \hline 0 & C_i^{-1} \end{array} \right) \left(\begin{array}{c|c} 0 & I_{l_{i-1}} \\ \hline I_{l_{i-1}} & 0 \end{array} \right) \left(\begin{array}{c|c} C_i^t & 0 \\ \hline 0 & C_i^{-1} \end{array} \right)^t = \left(\begin{array}{c|c} 0 & I_{l_{i-1}} \\ \hline I_{l_{i-1}} & 0 \end{array} \right).$$

For every ring R , by the trivial brace R will mean $(R, +, +)$.

Consider the trivial braces $T = T_1^{s_1} \times \cdots \times T_m^{s_m}$ and $S = R_1 \times \cdots \times R_m$. We define the symmetric bi-additive map $b: T \times T \rightarrow S$ by

$$\begin{aligned} & b((u_{1,1}, \dots, u_{1,s_1}, \dots, u_{m,1}, \dots, u_{m,s_m}), (v_{1,1}, \dots, v_{1,s_1}, \dots, v_{m,1}, \dots, v_{m,s_m})) \\ &= \left(\sum_{j=1}^{s_1} b_1(u_{1,j}, v_{1,j}), \dots, \sum_{j=1}^{s_m} b_m(u_{m,j}, v_{m,j}) \right), \end{aligned}$$

for all $u_{i,j}, v_{i,j} \in T_i$. We define $\alpha: S \rightarrow \text{Aut}(T)$ by $\alpha(a_1, \dots, a_m) = \alpha_{(a_1, \dots, a_m)}$ and

$$\begin{aligned} & \alpha_{(a_1, \dots, a_m)}(u_{1,1}, \dots, u_{1,s_1}, \dots, u_{m,1}, \dots, u_{m,s_m}) \\ &= (f_1^{a_m}(u_{1,1}), \dots, f_1^{a_m}(u_{1,s_1}), f_2^{a_1}(u_{2,1}), \dots, f_2^{a_1}(u_{2,s_2}), \dots, f_m^{a_{m-1}}(u_{m,1}), \dots, f_m^{a_{m-1}}(u_{m,s_m})), \end{aligned}$$

for all $a_i \in R_i$, and all $u_{i,j} \in T_i$. Note that, since f_i has order $p_i^{n_{i-1}}$ and $a_{i-1} \in \mathbb{Z}/(p_i^{n_{i-1}})$, the map α is well-defined. Clearly α is a group homomorphism from the group $(S, +)$ to $\text{Aut}(T, +, +)$. Since $f_i \in O(T_i, b_i)$, we have that

$$\begin{aligned} & b(\alpha_{(a_1, \dots, a_m)}(u_{1,1}, \dots, u_{1,s_1}, \dots, u_{m,1}, \dots, u_{m,s_m}), \alpha_{(a_1, \dots, a_m)}(v_{1,1}, \dots, v_{1,s_1}, \dots, v_{m,1}, \dots, v_{m,s_m})) \\ &= b((f_1^{a_m}(u_{1,1}), \dots, f_1^{a_m}(u_{1,s_1}), f_2^{a_1}(u_{2,1}), \dots, f_2^{a_1}(u_{2,s_2}), \dots, f_m^{a_{m-1}}(u_{m,1}), \dots, f_m^{a_{m-1}}(u_{m,s_m})), \\ & \quad (f_1^{a_m}(v_{1,1}), \dots, f_1^{a_m}(v_{1,s_1}), f_2^{a_1}(v_{2,1}), \dots, f_2^{a_1}(v_{2,s_2}), \dots, f_m^{a_{m-1}}(v_{m,1}), \dots, f_m^{a_{m-1}}(v_{m,s_m}))), \\ &= \left(\sum_{j=1}^{s_1} b_1(f_1^{a_m}(u_{1,j}), f_1^{a_m}(v_{1,j})), \dots, \sum_{j=1}^{s_m} b_m(f_m^{a_{m-1}}(u_{m,j}), f_m^{a_{m-1}}(v_{m,j})) \right), \\ &= \left(\sum_{j=1}^{s_1} b_1(u_{1,j}, v_{1,j}), \dots, \sum_{j=1}^{s_m} b_m(u_{m,j}, v_{m,j}) \right), \\ &= b((u_{1,1}, \dots, u_{1,s_1}, \dots, u_{m,1}, \dots, u_{m,s_m}), (v_{1,1}, \dots, v_{1,s_1}, \dots, v_{m,1}, \dots, v_{m,s_m})), \end{aligned}$$

for all $a_i \in R_i$ and all $u_{i,j}, v_{i,j} \in T_i$.

Hence, (2) is satisfied and we can construct the asymmetric product $T \rtimes_\alpha S$ of T by S via α and b .

Lemma 3.1 *The Sylow p_i -subgroup of the additive group of $T \rtimes_\circ S$ is*

$$A_i = \{0\} \times \cdots \times \{0\} \times T_i^{s_i} \times \{0\} \times \cdots \times \{0\} \times R_i \times \{0\} \times \cdots \times \{0\}.$$

Furthermore $A_i \cong (\mathbb{Z}/(p_i^{n_i}))^{2s_i l_{i-1} + 1}$.

Proof. Note that $|T \rtimes_\circ S| = \prod_{j \in \mathbb{Z}/(m)} p_j^{n_j(2s_j l_{j-1} + 1)}$ and $|A_i| = p_i^{n_i(2s_i l_{i-1} + 1)}$. Thus to prove that A_i is the Sylow p_i -subgroup of the additive subgroup of $T \rtimes_\circ S$, it is enough to show that A_i is an additive subgroup of $T \rtimes_\circ S$. But this is obvious from the definition of the addition in $T \rtimes_\circ S$. Therefore the first part of the result follows.

Let $u, v \in A_i$. Then

$$u = ((0, \dots, 0, u_{i,1}, \dots, u_{i,s_i}, 0, \dots, 0), (0, \dots, 0, a_i, 0, \dots, 0)) \in A_i$$

and

$$v = ((0, \dots, 0, v_{i,1}, \dots, v_{i,s_i}, 0, \dots, 0), (0, \dots, 0, c_i, 0, \dots, 0)) \in A_i,$$

for some $u_{i,k}, v_{i,k} \in T_i$ and $a_i, c_i \in R_i$. Since $T_i = R_i^{2l_{i-1}}$, there exist $a_{i,k,j}, c_{i,k,j} \in R_i$ such that

$$u_{i,k} = (a_{i,k,1}, \dots, a_{i,k,2l_{i-1}}) \quad \text{and} \quad v_{i,k} = (c_{i,k,1}, \dots, c_{i,k,2l_{i-1}}).$$

We define $\varphi: A_i \longrightarrow (\mathbb{Z}/(p_i^{n_i}))^{2s_i l_{i-1} + 1}$ by

$$\begin{aligned} \varphi(u) &= (a_{i,1,1}, \dots, a_{i,1,2l_{i-1}}, a_{i,2,1}, \dots, a_{i,2,2l_{i-1}}, \dots, a_{i,s_i,1}, \dots, a_{i,s_i,2l_{i-1}}, \\ &\quad a_i - \sum_{j=1}^{s_i} \sum_{k=1}^{l_{i-1}} a_{i,j,k} a_{i,j,k+l_{i-1}}). \end{aligned}$$

Note that

$$\begin{aligned} u + v &= ((0, \dots, 0, u_{i,1} + v_{i,1}, \dots, u_{i,s_i} + v_{i,s_i}, 0, \dots, 0), (0, \dots, 0, a_i + c_i + \sum_{j=1}^{s_i} b_i(u_{i,j}, v_{i,j}), 0, \dots, 0)) \\ &= ((0, \dots, 0, u_{i,1} + v_{i,1}, \dots, u_{i,s_i} + v_{i,s_i}, 0, \dots, 0), \\ &\quad (0, \dots, 0, a_i + c_i + \sum_{j=1}^{s_i} \sum_{k=1}^{l_{i-1}} (a_{i,j,k} c_{i,j,k+l_{i-1}} + a_{i,j,k+l_{i-1}} c_{i,j,k}), 0, \dots, 0)) \end{aligned}$$

and then

$$\begin{aligned} \varphi(u + v) &= (a_{i,1,1} + c_{i,1,1}, \dots, a_{i,1,2l_{i-1}} + c_{i,1,2l_{i-1}}, a_{i,2,1} + c_{i,2,1}, \dots, a_{i,2,2l_{i-1}} + c_{i,2,2l_{i-1}}, \\ &\quad \dots, a_{i,s_i,1} + c_{i,s_i,1}, \dots, a_{i,s_i,2l_{i-1}} + c_{i,s_i,2l_{i-1}}, \\ &\quad a_i + c_i + \sum_{j=1}^{s_i} \sum_{k=1}^{l_{i-1}} (a_{i,j,k} c_{i,j,k+l_{i-1}} + a_{i,j,k+l_{i-1}} c_{i,j,k}) \\ &\quad - \sum_{j=1}^{s_i} \sum_{k=1}^{l_{i-1}} (a_{i,j,k} + c_{i,j,k})(a_{i,j,k+l_{i-1}} + c_{i,j,k+l_{i-1}})) \\ &= (a_{i,1,1} + c_{i,1,1}, \dots, a_{i,1,2l_{i-1}} + c_{i,1,2l_{i-1}}, a_{i,2,1} + c_{i,2,1}, \dots, a_{i,2,2l_{i-1}} + c_{i,2,2l_{i-1}}, \\ &\quad \dots, a_{i,s_i,1} + c_{i,s_i,1}, \dots, a_{i,s_i,2l_{i-1}} + c_{i,s_i,2l_{i-1}}, \\ &\quad a_i + c_i - \sum_{j=1}^{s_i} \sum_{k=1}^{l_{i-1}} (a_{i,j,k} a_{i,j,k+l_{i-1}} + c_{i,j,k+l_{i-1}} c_{i,j,k}) \\ &= \varphi(u) + \varphi(v). \end{aligned}$$

Hence φ is a homomorphism of groups. Clearly φ is bijective. Therefore the result follows. \blacksquare

Remark 3.2 Note that the multiplicative group of the left brace $T \rtimes_\alpha S$ constructed above is the semidirect product $T \rtimes_\alpha S$ of T and S via α . Hence, it is metabelian. Since A_i is the additive Sylow p_i -subgroup of the left brace $T \rtimes_\alpha S$, it is clear that A_i is a left ideal. Thus the multiplicative group of A_i is a multiplicative Sylow p_i -subgroup of $T \rtimes_\alpha S$. By the definition of α , it is easy to see that the multiplicative group of A_i is isomorphic to the direct product $T_i^{s_i} \times R_i = (\mathbb{Z}/(p_i^{n_i}))^{2s_i l_{i-1} + 1}$. Hence the Sylow subgroups of the multiplicative group of the left brace $T \rtimes_\alpha S$ are abelian, i.e. $T \rtimes_\alpha S$ is an A-group.

Proposition 3.3 *With the above notation, the left brace $T \rtimes_\alpha S$ is simple.*

Proof. Let I be a non-zero ideal of $T \rtimes_\alpha S$. Let $((u_{1,1}, \dots, u_{1,s_1}, \dots, u_{m,1}, \dots, u_{m,s_m}), (a_1, \dots, a_m)) \in I$ be a non-zero element. By symmetry, we may assume that $(u_{1,1}, \dots, u_{1,s_1}, a_1) \neq (0, \dots, 0)$. Since p_1, \dots, p_m are different prime numbers, we also may assume that $(u_{i,1}, \dots, u_{i,s_i}, a_i) = (0, \dots, 0)$, for all $i \neq 1$. Thus $((u_{1,1}, \dots, u_{1,s_1}, 0, \dots, 0), (a_1, 0, \dots, 0)) \in I$.

We shall prove that $((0, \dots, 0), (p_1^{n_1-1}, 0, \dots, 0)) \in I$.

Note that if $(u_{1,1}, \dots, u_{1,s_1}) = (0, \dots, 0)$, then $a_1 \neq 0$ and, in this case, $((0, \dots, 0), (p_1^{n_1-1}, 0, \dots, 0)) \in I$, because every nontrivial additive subgroup of $\mathbb{Z}/(p_i^{n_i})$ contains $p_i^{n_i-1}$.

Suppose that $(u_{1,1}, \dots, u_{1,s_1}) \neq (0, \dots, 0)$. In this case, since b_1 is non-singular, there exists $(v_{1,1}, \dots, v_{1,s_1}) \in T_1^{s_1}$ such that $\sum_{j=1}^{s_1} b_1(u_{1,j}, v_{1,j}) \neq 0$. Now, since I is closed by the lambda maps, we get

$$\begin{aligned} & \lambda_{((v_{1,1}, \dots, v_{1,s_1}, 0, \dots, 0), (0, \dots, 0))}((u_{1,1}, \dots, u_{1,s_1}, 0, \dots, 0), (a_1, 0, \dots, 0)) \\ &= ((u_{1,1}, \dots, u_{1,s_1}, 0, \dots, 0), (a_1 - \sum_{j=1}^{s_1} b_1(u_{1,j}, v_{1,j}), 0, \dots, 0)) \in I. \end{aligned}$$

Hence

$$\begin{aligned} & ((u_{1,1}, \dots, u_{1,s_1}, 0, \dots, 0), (a_1, 0, \dots, 0)) \\ & - ((u_{1,1}, \dots, u_{1,s_1}, 0, \dots, 0), (a_1 - \sum_{j=1}^{s_1} b_1(u_{1,j}, v_{1,j}), 0, \dots, 0)) \\ &= ((0, \dots, 0), (\sum_{j=1}^{s_1} b_1(u_{1,j}, v_{1,j}), 0, \dots, 0)) \in I, \end{aligned}$$

and thus also $((0, \dots, 0), (p_1^{n_1-1}, 0, \dots, 0)) \in I$ in this case.

Now, by Lemma 2.1, we have that

$$\begin{aligned} & (\lambda_{((0, \dots, 0), (p_1^{n_1-1}, 0, \dots, 0))} - \text{id})((0, \dots, 0, v_{2,1}, \dots, v_{2,s_2}, 0, \dots, 0), (0, \dots, 0)) \\ &= ((0, \dots, 0, (f_2^{p_1^{n_1-1}} - \text{id})(v_{2,1}), \dots, (f_2^{p_1^{n_1-1}} - \text{id})(v_{2,s_2}), 0, \dots, 0), \\ & \quad (0, -\sum_{j=1}^{s_2} b_2(v_{2,j}, (f_2^{p_1^{n_1-1}} - \text{id})(v_{2,j})), 0, \dots, 0)) \in I, \end{aligned} \tag{3}$$

for every $(v_{2,1}, \dots, v_{2,s_2}) \in T_2^{s_2}$. Note that

$$q_1(x) = p_1 + (x^{p_1^{n_1-1}} - 1) \sum_{j=1}^{p_1-1} j x^{(p_1-j-1)p_1^{n_1-1}}.$$

Hence

$$0 = q_1(C_2) = p_1 I_{l_2} + (C_2^{p_1^{n_1-1}} - I_{l_2}) \sum_{j=1}^{p_1-1} j C_2^{(p_1-j-1)p_1^{n_1-1}}.$$

Since $p_1 \in R_2$ is invertible, this implies that $f_2^{p_1^{n_1-1}} - \text{id}$ is invertible. Hence, by the definition of b_2 , there exist $v_{2,j}, v'_{2,j} \in T_2$ such that

$$\sum_{j=1}^{s_2} b_2((f_2^{p_1^{n_1-1}} - \text{id})(v_{2,j}), v'_{2,j}) = 1. \quad (4)$$

Indeed, there exists $v_{2,1} \in T_2$ such that $(f_2^{p_1^{n_1-1}} - \text{id})(v_{2,1}) = (1, 0, \dots, 0)$. Let $v'_{2,1} = (1, \dots, 1)$ and $v'_{2,j} = (0, \dots, 0)$, for all $1 < j \leq s_2$. Then $b_2((1, 0, \dots, 0), (1, \dots, 1)) = 1$ and thus we get (4). By (3)

$$\begin{aligned} & ((0, \dots, 0, (f_2^{p_1^{n_1-1}} - \text{id})(v_{2,1}), \dots, (f_2^{p_1^{n_1-1}} - \text{id})(v_{2,s_2}), 0, \dots, 0), \\ & (0, -\sum_{j=1}^{s_2} b_2(v_{2,j}, (f_2^{p_1^{n_1-1}} - \text{id})(v_{2,j})), 0, \dots, 0)) \in I. \end{aligned} \quad (5)$$

Then, since I is closed by the lambda maps, applying $\lambda_{((0, \dots, 0, v'_{2,1}, \dots, v'_{2,s_2}, 0, \dots, 0), (0, \dots, 0))}$ to the element in (5), we have that

$$\begin{aligned} & \lambda_{((0, \dots, 0, v'_{2,1}, \dots, v'_{2,s_2}, 0, \dots, 0), (0, \dots, 0))}((0, \dots, 0, (f_2^{p_1^{n_1-1}} - \text{id})(v_{2,1}), \dots, (f_2^{p_1^{n_1-1}} - \text{id})(v_{2,s_2}), 0, \dots, 0), \\ & (0, -\sum_{j=1}^{s_2} b_2(v_{2,j}, (f_2^{p_1^{n_1-1}} - \text{id})(v_{2,j})), 0, \dots, 0)) \\ & = ((0, \dots, 0, (f_2^{p_1^{n_1-1}} - \text{id})(v_{2,1}), \dots, (f_2^{p_1^{n_1-1}} - \text{id})(v_{2,s_2}), 0, \dots, 0), \\ & (0, -\sum_{j=1}^{s_2} b_2(v_{2,j}, (f_2^{p_1^{n_1-1}} - \text{id})(v_{2,j})) - \sum_{j=1}^{s_2} b_2((f_2^{p_1^{n_1-1}} - \text{id})(v_{2,j}), v'_{2,j}), 0, \dots, 0)) \\ & = ((0, \dots, 0, (f_2^{p_1^{n_1-1}} - \text{id})(v_{2,1}), \dots, (f_2^{p_1^{n_1-1}} - \text{id})(v_{2,s_2}), 0, \dots, 0), \\ & (0, -\sum_{j=1}^{s_2} b_2(v_{2,j}, (f_2^{p_1^{n_1-1}} - \text{id})(v_{2,j})) - 1, 0, \dots, 0)) \in I, \end{aligned}$$

where the last equality follows by (4). Hence, this together with (5) yields

$$\begin{aligned} & ((0, \dots, 0, (f_2^{p_1^{n_1-1}} - \text{id})(v_{2,1}), \dots, (f_2^{p_1^{n_1-1}} - \text{id})(v_{2,s_2}), 0, \dots, 0), \\ & (0, -\sum_{j=1}^{s_2} b_2(v_{2,j}, (f_2^{p_1^{n_1-1}} - \text{id})(v_{2,j})), 0, \dots, 0)) \\ & - ((0, \dots, 0, (f_2^{p_1^{n_1-1}} - \text{id})(v_{2,1}), \dots, (f_2^{p_1^{n_1-1}} - \text{id})(v_{2,s_2}), 0, \dots, 0), \\ & (0, -\sum_{j=1}^{s_2} b_2(v_{2,j}, (f_2^{p_1^{n_1-1}} - \text{id})(v_{2,j})) - 1, 0, \dots, 0)) \\ & = ((0, \dots, 0), (0, 1, 0, \dots, 0)) \in I. \end{aligned} \quad (6)$$

Now we have that

$$((0, \dots, 0), (0, \sum_{j=1}^{s_2} b_2(w_{2,j}, (f_2^{p_1^{n_1-1}} - \text{id})(w_{2,j})), 0, \dots, 0)) \in I,$$

for all $(w_{2,1}, \dots, w_{2,s_2}) \in T_2^{s_2}$, and by (3), we also have that

$$\begin{aligned} & ((0, \dots, 0, (f_2^{p_1^{n_1-1}} - \text{id})(w_{2,1}), \dots, (f_2^{p_1^{n_1-1}} - \text{id})(w_{2,s_2}), 0, \dots, 0), \\ & (0, -\sum_{j=1}^{s_2} b_2(w_{2,j}, (f_2^{p_1^{n_1-1}} - \text{id})(w_{2,j})), 0, \dots, 0)) \\ & + ((0, \dots, 0), (0, \sum_{j=1}^{s_2} b_2(w_{2,j}, (f_2^{p_1^{n_1-1}} - \text{id})(w_{2,j})), 0, \dots, 0)) \in I, \end{aligned}$$

for all $(w_{2,1}, \dots, w_{2,s_2}) \in T_2^{s_2}$. Thus

$$\begin{aligned}
& ((0, \dots, 0, (f_2^{p_1^{n_1-1}} - \text{id})(w_{2,1}), \dots, (f_2^{p_1^{n_1-1}} - \text{id})(w_{2,s_2}), 0, \dots, 0), \\
& \quad (0, -\sum_{j=1}^{s_2} b_2(w_{2,j}, (f_2^{p_1^{n_1-1}} - \text{id})(w_{2,j})), 0, \dots, 0)) \\
& \quad + ((0, \dots, 0), (0, \sum_{j=1}^{s_2} b_2(w_{2,j}, (f_2^{p_1^{n_1-1}} - \text{id})(w_{2,j})), 0, \dots, 0)) \\
& = ((0, \dots, 0, (f_2^{p_1^{n_1-1}} - \text{id})(w_{2,1}), \dots, (f_2^{p_1^{n_1-1}} - \text{id})(w_{2,s_2}), 0, \dots, 0), (0, \dots, 0)) \in I,
\end{aligned}$$

for all $(w_{2,1}, \dots, w_{2,s_2}) \in T_2^{s_2}$. Since $f_2^{p_1^{n_1-1}} - \text{id}$ is invertible, this yields that

$$((0, \dots, 0, t_{2,1}, \dots, t_{2,s_2}, 0, \dots, 0), (0, \dots, 0)) \in I,$$

for all $(t_{2,1}, \dots, t_{2,s_2}) \in T_2^{s_2}$. Since, by (6), $((0, \dots, 0), (0, 1, 0, \dots, 0)) \in I$, we have that

$$((0, \dots, 0, t_{2,1}, \dots, t_{2,s_2}, 0, \dots, 0), (0, d_2, 0, \dots, 0)) \in I,$$

for all $(t_{2,1}, \dots, t_{2,s_2}, d_2) \in T_2^{s_2} \times R_2$. Now, by a similar argument, one can prove that

$$((0, \dots, 0, t_{i,1}, \dots, t_{i,s_i}, 0, \dots, 0), (0, \dots, 0, d_i, 0, \dots, 0)) \in I,$$

for all $(t_{i,1}, \dots, t_{i,s_i}, d_i) \in T_i^{s_i} \times R_i$, for all $i \in \mathbb{Z}/(m)$, and the result follows. ■

Hence, as an easy consequence of Proposition 3.3, Lemma 3.1 and Remark 3.2, we get the desired result.

Theorem 3.4 *Let A be a finite abelian group. Then A is a subgroup of the additive group of a finite simple left brace B with metabelian multiplicative group with abelian Sylow subgroups.*

References

- [1] D. Bachiller, Extensions, matched products and simple braces, J. Pure Appl. Algebra 222 (2018), 1670–1691.
- [2] D. Bachiller, F. Cedó and E. Jespers, Solutions of the Yang-Baxter equation associated with a left brace, J. Algebra 463 (2016), 80–102.
- [3] D. Bachiller, F. Cedó, E. Jespers and J. Okniński, Iterated matched products of finite braces and simplicity; new solutions of the Yang-Baxter equation, Trans. Amer. Math. Soc. 370 (2018), 4881–4907.
- [4] D. Bachiller, F. Cedó, E. Jespers and J. Okniński, Asymmetric product of left braces and simplicity; new solutions of the Yang-Baxter equation, Communications in Contemporary Mathematics 21 (2019), 1850042 (30 pages).
- [5] F. Catino, I. Colazzo and P. Stefanelli, Regular subgroups of the affine group and asymmetric product of braces, J. Algebra 455 (2016), 164–182.
- [6] F. Cedó, Left braces: solutions of the Yang-Baxter equation, Adv. Group Theory Appl. 5 (2018), 33–90.
- [7] F. Cedó, E. Jespers and J. Okniński, Braces and the Yang-Baxter equation, Commun. Math. Phys. 327 (2014), 101–116.
- [8] F. Cedó, E. Jespers and J. Okniński, An abundance of simple left braces with abelian multiplicative Sylow subgroups (to appear in Revista Matemática Iberoamericana).
- [9] V. G. Drinfeld, On unsolved problems in quantum group theory. Quantum Groups, Lecture Notes Math. 1510, Springer-Verlag, Berlin, 1992, 1–8.

- [10] P. Etingof, T. Schedler and A. Soloviev, Set-theoretical solutions to the quantum Yang-Baxter equation, *Duke Math. J.* 100 (1999), 169–209.
- [11] W. Rump, Braces, radical rings, and the quantum Yang-Baxter equation, *J. Algebra* 307 (2007), 153–170.
- [12] W. Rump, The brace of a classical group, *Note Mat.* 34 (2014), no. 1, 115–144.
- [13] A. Smoktunowicz, A note on set-theoretic solutions of the Yang–Baxter equation, *J. Algebra* 500 (2018), 3–18.
- [14] D.R. Taunt, On A-groups, *Proc. Cambridge Philos. Soc.* 45 (1949), 24–42.

F. Cedó
 Departament de Matemàtiques
 Universitat Autònoma de Barcelona
 08193 Bellaterra (Barcelona), Spain
 cedo@mat.uab.cat

E. Jespers
 Department of Mathematics
 Vrije Universiteit Brussel
 Pleinlaan 2, 1050 Brussel, Belgium
 Eric.Jespers@vub.be

J. Okniński
 Institute of Mathematics
 Warsaw University
 Banacha 2, 02-097 Warsaw, Poland
 okninski@mimuw.edu.pl