

Time and Observables in Covariant Quantum Theory

Natalia Gorobey¹ and Alexander Lukyanenko^{1,*}

¹*Peter the Great Saint Petersburg Polytechnic University,
Polytekhnicheskaya 29, 195251, St. Petersburg, Russia*

A modification of the covariant theory is proposed in which the self-energy of the system, corresponding to time-like degrees of freedom in the configuration space, preserves the classical law of change in quantum theory. As a result, proper time in covariant quantum theory takes on a dynamic meaning. As applications of the new formalism, a modification of the relativistic quantum mechanics of a scalar particle and a homogeneous model of the universe is considered.

I. INTRODUCTION

In theories based on the principle of covariance, including reparametrization of time, there is a problem with the determination of this time after quantization. Here we focus on the covariant approach to quantization in which the Batalin-Fradkin-Vilkovysky theorem (BFV) [1–3] and the concept of BRST invariance [4, 5] play the central role. However, for a simplest covariant system — a relativistic particle, the result of this formalism reduces to the well-known representation of the Feynman propagator in the form of an integral over the particle’s proper time [6]. This representation of the Green’s function for the Dirac and Klein-Gordon operators was first introduced by Fock [7] and Schwinger [8] and is now widely used for calculations [9]. The Feynman propagator serves to solve the scattering problem in relativistic quantum mechanics in the framework of perturbation theory [10]. It should be expected that in the general case the propagator contains a multiple integral over the parameters of the system’s proper time (multi-arrow time in the theory of gravity [11]), as well as over the remaining parameters of the full symmetry group. In this paper, a simplified version of the construction of a propagator of covariant theory is proposed: the amplitude of the transition is determined on the space of the symmetry group and then it is integrated over all group parameters. Thus, proper time is also excluded from the dynamics. To restore the dynamic meaning of proper time, based on this integral representation of the covariant propagator, a modification of the covariant quantum theory is proposed in this paper. The proposed modification can be considered as an approximation in which certain dynamic variables (in each case it is clear which ones) are limited by the classical laws of their motion. Its result is precisely the removal of integration in proper time. Formally, the approximation is reduced to the linearization of quadratic in momenta (Hamiltonian) constraints with respect to the corresponding canonical momenta and the appearance of δ -functions that remove integration over proper time. Together with the time parameter in a modified theory, an observable is associated with it that has

the physical meaning of the system’s own energy, corresponding to time-like degrees of freedom in the configuration space. As examples, the quantum mechanics of a scalar relativistic particle and a homogeneous model of the Friedman universe are considered.

In the section II, a simplified version of the covariant quantum theory is formulated, which then will be the basis for its modification. In the section III, we consider a modification of relativistic quantum mechanics, in which the x_0 -coordinate of a particle in Minkowski space serves as the evolution parameter, and the corresponding observable is the particle’s own energy. In the section IV, we consider a modified quantum theory of the homogeneous Friedmann universe, in which the radius of the universe serves as the evolution parameter, and the energy of space is the observable associated with it. In this case, the pair of variables (the radius of the universe and the energy of the space) can exchange their roles (time and the observable). This result is consistent with the proposal to consider space energy density as a parameter of time in cosmology [12].

II. PARAMETERS OF THE SYMMETRY GROUP AS DYNAMIC VARIABLES

We take the initial action of the covariant theory in the canonical form [13]

$$I = \int_0^1 d\tau \left[p_k \dot{q}_k - \lambda_a \varphi_a \right], \quad (1)$$

where the Hamiltonian $h(p, q)$ is set equal to zero. Moreover, in the set of constraints $\varphi_a(p, q)$ we assume the presence of quadratic in momenta (Hamiltonian) constraints, as is the case of a relativistic particle and cosmology that interest us. The constraints form a closed Lie algebra with respect to the Poisson brackets,

$$\{\varphi_a, \varphi_b\} = C_{abd} \varphi_d, \quad (2)$$

where the structural "constants" C_{abd} generally depend on the canonical variables. In the theory of gravity, they depend on the 3D metric [14]. Here, for simplicity, we consider them to be constants. They obey Jacobi’s identity:

$$C_{bcd} C_{adp} - C_{abd} C_{dcp} - C_{acd} C_{bdp} = 0. \quad (3)$$

* alex.lukyan@mail.ru

The action Eq. (1) is invariant with respect to gauge transformations ($\varepsilon_d(0) = \varepsilon_d(1) = 0$),

$$\delta q_k = \varepsilon_a \{q_k, \varphi_a\}, \delta p_k = \varepsilon_a \{p_k, \varphi_a\}, \quad (4)$$

provided that the Lagrange multipliers are converted as follows:

$$\delta \lambda_a = \dot{\varepsilon}_a - C_{bda} \lambda_b \varepsilon_d. \quad (5)$$

We introduce the explicit dependence of the Lagrange multipliers λ_a on the group parameters s_a for which $\varepsilon_a = \delta s_a$ are infinitesimal variations, and write the relation Eq. (5) in the form of a functional differential equation

$$\frac{\delta \lambda_a(\tau)}{\delta s_d(\tau')} = \delta_{ad} \frac{d}{d\tau} \delta(\tau - \tau') - C_{bda} \lambda_b \delta(\tau - \tau'), \quad (6)$$

which is supplemented by the initial condition $\lambda_a(0) = 0$. The solution of Eq. (6) is linear with respect to the velocities \dot{s}_a :

$$\lambda_b = \dot{s}_a \Lambda_{ab}, \quad (7)$$

where the Taylor series of Λ_{ab} in s up to the second order is

$$\Lambda_{ab} = \delta_{ab} - \frac{1}{2!} C_{adb} s_d + \frac{1}{3!} C_{ad'b'} C_{b'db} s_{d'} s_d + \dots \quad (8)$$

Thus, the original coordinate set q_k is supplemented with group parameters s_a that are now explicitly present in the action:

$$I = \int_0^1 d\tau \left[p_k \dot{q}_k - \dot{s}_b \Lambda_{ba}(s) \varphi_a \right]. \quad (9)$$

It remains to write down the action Eq. (9) in canonical form:

$$I = \int_0^1 d\tau \left[p_k \dot{q}_k + P_a \dot{s}_a - \eta_a \Phi_a \right], \quad (10)$$

where

$$\Phi_a = P_a + \Lambda_{ab} \varphi_b. \quad (11)$$

A set of constraints obeys the simple algebra of Poisson brackets

$$\{\Phi_a, \Phi_b\} = 0 \quad (12)$$

at the condition

$$\frac{\partial \Lambda_{ap}}{\partial s_q} - \frac{\partial \Lambda_{sp}}{\partial s_a} + \Lambda_{ab} \Lambda_{qr} C_{brp} = 0. \quad (13)$$

The relation (13) holds in our case with the indicated accuracy as a consequence of the Jacobi identity Eq. (3).

We quantize the theory based on the new canonical form of the action Eq. (10). For the wave function $\psi(s, q)$, we obtain the system of wave equations

$$i\hbar \frac{\partial \psi}{\partial s_a} = \Lambda_{ab}(s) \varphi_b(q, \hat{p}) \psi, \quad (14)$$

which have the form of an equation of evolution on the space of a symmetry group. In our formulation, the covariance condition is that, since the group parameters s_a are unobservable, the solution of the system Eq. (14) should be integrated over the entire range of their variation. So, in the presence of an abelian gauge subgroup, the corresponding part of the equations of the system Eq. (14) has the form

$$\frac{\partial \psi}{\partial s_a} = \frac{\partial \psi}{\partial q_a}, \quad (15)$$

where q_a are (longitudinal) coordinates, on which the wave function should not depend. According to Eq. (15), the wave function depends on the sum $q_a + s_a$ so that integration over s_a in the range from $-\infty$ to $+\infty$ excludes the dependence on q_a .

The group parameters corresponding to the Hamiltonian constraints have the meaning of the proper times of the system. One should integrate over them within the limits $[0, +\infty)$ [6]. It is these integrals that are the subject of concern in our subsequent constructions, and we will still refine this construction in the process of its application.

III. MODIFIED PROPAGATOR FOR RELATIVISTIC PARTICLE

We take the initial action of a relativistic particle in a parameterized form (speed of light $c = 1$) in a form

$$\tilde{I} = \int_0^1 d\tau \left[\frac{1}{4} \left(\frac{\dot{x}_0}{N} - \frac{\dot{x}_i}{N} \right)^2 + m^2 N + e\varphi(x_0, x_i) \dot{x}_0 \right], \quad (16)$$

where the dot denotes the derivative with respect to the parameter τ on its world line. Here we restrict ourselves to the interaction with the electric field of strength $E_i = -\partial_i \varphi$. The Hamiltonian constraint

$$H = (p_0 - e\varphi)^2 - p_i^2 - m^2 \quad (17)$$

is quadratic in momenta. Our task in the future is to remove the integration over the particle's proper time

$$s = \int_0^1 d\tau N(\tau) \quad (18)$$

in the Fock-Schwinger (FS) representation for the Feynman propagator in an external electric field. We modify the dynamics of the coordinate x_0 of Minkowski space, adding to the action (16) its variation generated by the infinitesimal time shift of x_0 ,

$$\delta x_0 = -\frac{\dot{x}_0}{N} \varepsilon. \quad (19)$$

A new action,

$$\int_0^1 d\tau \left[\frac{1}{4} \frac{\dot{x}_0^2}{N} \left(1 - \frac{\dot{\varepsilon}}{N} \right) - \frac{1}{4} \frac{\dot{x}_i^2}{N} + m^2 N + e\varphi \dot{x}_0 + e\partial_i \varphi \frac{\dot{x}_0 \dot{x}_i}{N} \varepsilon \right], \quad (20)$$

contains the time shift ε as an independent variable. To go to the canonical form of the modified theory, we find the canonical momenta,

$$p_0 = \frac{1}{2} \frac{\dot{x}_0}{N} \left(1 - \frac{\dot{\varepsilon}}{N} \right) + e\varphi + e\partial_i \varphi \frac{\dot{x}_i}{N} \varepsilon, \quad (21)$$

$$p_i = -\frac{1}{2} \frac{\dot{x}_i}{N} + e\partial_i \varphi \frac{\dot{x}_0}{N} \varepsilon, \quad (22)$$

$$-P_\varepsilon = -\frac{1}{4} \frac{\dot{x}_0^2}{N^2}, \quad (23)$$

and a modified Hamiltonian,

$$\tilde{h} = N \left[2\sqrt{P_\varepsilon} (p_0 - e\varphi) - \left(p_i + 2e\partial_i \varphi \varepsilon \sqrt{P_\varepsilon} \right)^2 - P_\varepsilon - m^2 \right]. \quad (24)$$

The Hamiltonian function is linear in the momentum p_0 . The pair of canonical variables $(\varepsilon, P_\varepsilon)$ appears and ε will be considered as the canonical momentum. The canonical coordinate P_ε has the physical meaning of the particle's own energy. After the replacement

$$\tilde{p}_i = p_i + 2e\partial_i \varphi \varepsilon \sqrt{P_\varepsilon}, \quad (25)$$

the canonical form of the modified action takes the form

$$\tilde{I} = \int_0^s ds \left[\tilde{p}_i \dot{x}_i + p_0 \dot{x}_0 - \varepsilon \left(\dot{P}_\varepsilon + 2e\partial_i \varphi \dot{x}_0 \sqrt{P_\varepsilon} \right) - \tilde{H} \right], \quad (26)$$

where,

$$\tilde{H} = 2\sqrt{P_\varepsilon} (p_0 - e\varphi) - \tilde{p}_i^2 - P_\varepsilon - m^2. \quad (27)$$

Here we proceed to the parametrization of the world line of the particle with its proper time Eq. (18). The following dynamic constraint is also added to the action Eq. (26):

$$\dot{P}_\varepsilon + 2e\partial_i \varphi \dot{x}_i \sqrt{P_\varepsilon} = 0, \quad (28)$$

which represents the classical law of particle energy change in an external electric field, and the momentum ε plays the role of the Lagrange multiplier.

In the formal representation of the modified propagator for a particle as a functional integral on the phase space,

$$\tilde{K} = \int_0^\infty ds \int \prod_{s,i} \frac{dp_0 dx_0}{2\pi\hbar} \frac{d\tilde{p}_i dx_i}{2\pi\hbar} \frac{dP_\varepsilon d\varepsilon}{2\pi\hbar} \exp \left\{ \frac{i}{\hbar} \tilde{I} \right\}, \quad (29)$$

after functional integration over momenta, we obtain the product of δ -functions in the integral on the augmented configuration space:

$$\begin{aligned} \tilde{K} &= \int_0^\infty ds \int \prod_{s,i} \frac{dx_i}{\sqrt{2\pi\hbar}} \frac{dx_0}{\sqrt{2\pi\hbar}} \frac{dP_\varepsilon}{\sqrt{2\pi\hbar}} \delta \left(\dot{x}_0 - 2\sqrt{P_\varepsilon} \right) \\ &\quad \times \delta \left(\dot{P}_\varepsilon + 2e\partial_i \varphi \dot{x}_i \sqrt{P_\varepsilon} \right) \\ &\quad \times \exp \left\{ \frac{i}{\hbar} \int_0^s ds \left[\dot{x}_i^2 + P_\varepsilon + m^2 + e\varphi \right] \right\}. \end{aligned} \quad (30)$$

The propagator Eq. (30) serves as the basis for a new dynamic interpretation of relativistic quantum mechanics. For definiteness, we consider the scattering problem. In the absence of an external electric field, the integrals over all internal values of the coordinates x_0 and P_ε are removed and the product of the δ -function remains

$$\delta \left(\frac{x'_0 - x_0}{s} - 2\sqrt{P'_\varepsilon} \right) \delta \left(\frac{P'_\varepsilon - P_\varepsilon}{s} \right). \quad (31)$$

Then, the integral over the proper time in Eq. (30) is removed by the first δ -function. Thus, the physical parameter of time, as expected, is x_0 . In this asymptotic region, according to the Hamiltonian constraint Eq. (27), the wave equation has the form of the Schrödinger equation

$$i\hbar 2\sqrt{P_\varepsilon} \frac{\partial \psi}{\partial x_0} = [-\hbar^2 \Delta + P_\varepsilon + m^2] \psi, \quad (32)$$

in which the parameter x_0 plays the role of the classical time parameter. However, if we put $P_\varepsilon = E$, it gives the relativistic dispersion law $E^2 - p_i^2 - m^2 = 0$ for a de Broglie plane wave. In the interaction region, the integrals over all internal values of the coordinates x_0 and P_ε are also removed, but now we obtain the proper time s in the form of the functional of the trajectory $x_i(s)$ of the particle. Therefore, the integral over the proper time in Eq. (30) should be located to the right of the functional integral along the particle trajectory. We note that the classical law of the change in the self-energy of the particle P_ε with time excludes the processes of annihilation and production of particle-antiparticle pairs in the approximation considered here. A consistent consideration of these processes is possible only after secondary quantization. In cosmology we want to avoid this.

IV. MODIFIED PROPAGATOR FOR FRIEDMAN'S HOMOGENEOUS UNIVERSE

The dynamics of the Friedman universe is determined by the action

$$I = \int_0^1 d\tau \left[-\frac{a}{2g} \left(\frac{\dot{a}}{N} - N \right) + 2\pi^2 a^3 \frac{1}{2} \left(\frac{\dot{\phi}_k}{N} - V(\phi) \right) \right], \quad (33)$$

where $g = 2G/3\pi$, G is the Newtonian gravitational constant and a is the radius of the universe. We consider the set of scalar fields ϕ_k as matter and N is the ADM lapse function [14]. As in the case of a relativistic particle, we have here one Hamiltonian constraint and an additional integration over the proper time in the path integral representing the wave function of the universe [15]. In the Euclidean form of quantum theory, this integral diverges due to the uncertainty of the sign of the action. This theory only makes sense in the semiclassical approximation. We will further search for the dynamic meaning of this quantum theory in its modification. An infinitesimal time shift,

$$\delta a = -\frac{\dot{a}}{N}\varepsilon, \quad (34)$$

leads to a modified action

$$\begin{aligned} \tilde{I} = & \int_0^1 d\tau \left[-\frac{1}{2g} \frac{a\dot{a}^2}{N} \left(1 - \frac{\dot{\varepsilon}}{N} \right) + \frac{a}{2g} \left(N + \dot{\varepsilon} \right) \right. \\ & \left. + \pi^2 \left(a^3 - 3a^2\dot{a}\varepsilon \right) \left(\frac{\dot{\phi}_k}{N} - V(\phi) N \right) \right]. \quad (35) \end{aligned}$$

Compared to a relativistic particle, the interaction of a time-like dynamic variable a with matter is more complex dynamics. For the energy of the space P_ε from Eq. (35) we obtain the equation of motion

$$\dot{P}_\varepsilon + 3\pi^2 a^2 \dot{a} \left(\frac{\dot{\phi}_k}{N^2} - V(\phi) \right) = 0. \quad (36)$$

We turn to the canonical form of the modified action. We find the canonical momenta:

$$p_a = -\frac{1}{g} \frac{a\dot{a}}{N} \left(1 - \frac{\dot{\varepsilon}}{N} \right) - 3\pi^2 a^2 \varepsilon \left(\frac{\dot{\phi}_k}{N^2} - V(\phi) \right), \quad (37)$$

$$p_{\phi_k} = 2\pi^2 \left(a^3 - 3a^2\dot{a}\varepsilon \right) \frac{\dot{\phi}_k}{N}, \quad (38)$$

$$P_\varepsilon = \frac{a}{2g} \left(\frac{\dot{a}}{N^2} + 1 \right). \quad (39)$$

The modified Hamiltonian constraint is now linear in momentum p_a :

$$\begin{aligned} \tilde{H} = & p_a \sqrt{\frac{2gP_\varepsilon}{a} - 1} + \frac{a}{2g} \left(\sqrt{\frac{2gP_\varepsilon}{a} - 1} - 1 \right) \\ & + \frac{1}{2} \left[\frac{p_{\phi_k}^2}{2\pi^2 (a^3 - 3a^2\zeta)} + 2\pi^2 (a^3 - 3a^2\zeta) V(\phi) \right], \quad (40) \end{aligned}$$

where

$$\zeta = \dot{a}\varepsilon = \varepsilon \sqrt{\frac{2gP_\varepsilon}{a} - 1}. \quad (41)$$

Here the difficulty arises: the infinitesimal time shift ε , which serves as the canonical momentum, falls into the denominator. It will become a problem after quantization. We write, however, a modified propagator in the form of a formal continual integral in the canonical form,

$$\begin{aligned} \tilde{K} = & \int_0^\infty ds \int \prod_s \frac{dp_a da}{2\pi\hbar} \frac{d\varepsilon dP_\varepsilon}{2\pi\hbar} \prod_{k=1}^\alpha \frac{dp_{\phi_k} d\phi_k}{2\pi\hbar} \frac{d\lambda d\zeta}{2\pi\hbar} J \\ & \times \exp \left(\frac{i}{\hbar} \tilde{I} \right), \quad (42) \end{aligned}$$

where

$$\tilde{I} = \int_0^s ds \left[p_{\phi_k} \dot{\phi}_k + p_a \dot{a} - \varepsilon \dot{P}_\varepsilon + \lambda (\zeta - \dot{a}\varepsilon) - \tilde{H} \right]. \quad (43)$$

Here, we took into account the relation Eq. (41), λ is the corresponding Lagrange multiplier, and J will be the ‘‘price’’ for the mentioned complication in the Hamiltonian constraint. Momentum integration gives:

$$\begin{aligned} \tilde{K} = & \int_0^\infty ds \int \prod_s \frac{dP_\varepsilon da}{2\pi\hbar} \frac{dp_a da}{2\pi\hbar} \\ & \times \prod_k \frac{d\phi_k}{(2\pi\hbar)^{\alpha/2} [2\pi^2 (a^3 - 3a^2\zeta)]^{\alpha/2}} \\ & \times \frac{d\lambda d\zeta}{2\pi\hbar} J \delta \left(\dot{a} - \sqrt{\frac{2gP_\varepsilon}{a} - 1} \right) \delta \left(\dot{P}_\varepsilon + \lambda \dot{a} \right) \\ & \times \exp \left(-\frac{i}{\hbar} \int_0^s ds \lambda \zeta \right) \\ & \times \exp \left\{ -\frac{i}{\hbar} \int_0^s ds \left[\frac{a}{2g} \left(\sqrt{\frac{2gP_\varepsilon}{a} - 1} - 1 \right) \right] \right\} \\ & \times \exp \left\{ -\frac{i}{\hbar} \int_0^s ds \left[\pi^2 (a^3 - 3a^2\zeta) \right. \right. \\ & \left. \left. \times \left(\frac{\dot{\phi}_k}{N} - V(\phi) \right) \right] \right\}. \quad (44) \end{aligned}$$

Thus, in order to get a reasonable expected result, you should put

$$J = [2\pi^2 (a^3 - 3a^2\zeta)]^{\alpha/2}. \quad (45)$$

The integral over ζ is now taken and gives the δ -function

$$\delta \left(\lambda - 3a^2 \left(\frac{\dot{\phi}_k}{N} - V(\phi) \right) \right). \quad (46)$$

Then the integral over λ is taken. Finally, we obtain a modified propagator containing two δ -functions:

$$\begin{aligned} \tilde{K} = & \int_0^\infty ds \int \prod_s \frac{dP_\varepsilon da}{2\pi\hbar} \prod_k \frac{d\phi_k}{(2\pi\hbar)^{\alpha/2}} \delta \left(a - \sqrt{\frac{2gP_\varepsilon}{a} - 1} \right) \\ & \times \delta \left(\dot{P}_\varepsilon + 3a^2 \dot{a} \left(\dot{\phi}_k^2 - V(\phi) \right) \right) \\ & \times \exp \left\{ -\frac{i}{\hbar} \int_0^s ds \left[\frac{a}{2g} \left(\sqrt{\frac{2gP_\varepsilon}{a} - 1} - 1 \right) \right] \right\} \\ & \times \exp \left\{ -\frac{i}{\hbar} \int_0^s ds \left[\pi^2 a^3 \left(\dot{\phi}_k^2 - V(\phi) \right) \right] \right\}. \end{aligned} \quad (47)$$

These δ -functions remove the integration over a , P_ε in the functional integral. The remaining two δ -functions, that depend on the boundary values of the variables a , P_ε in the propagator (any of them), take the integral in proper time and determine the physical time.

There are two possibilities for this. The most familiar is the choice of the radius a as a parameter of time with the beginning in the singularity $a = 0$. Now we need to set the initial state of the universe near the singularity, as well as the initial value of the energy of the space P_ε . The difficulty is that the Hamiltonian constraint (40) and the corresponding Schrödinger equation in quantum theory are connected by the parameter ζ to the dynamic constraint Eq. (39). To unlock this connection, we will make further approximations. Near the singularity, the constraint Eq. (40) is simplified:

$$\tilde{H} \cong \sqrt{\frac{2gP_\varepsilon}{a}} p_a - \frac{p_{\phi_k}^2}{12\pi^2 a^2 \zeta}. \quad (48)$$

The parameter ζ can be found from the condition for the extremum of the action Eq. (43),

$$\zeta = \frac{\sqrt{p_{\phi_k}^2}}{2\sqrt{3\pi a}\sqrt{\lambda}}, \quad (49)$$

and λ can be found from Eq. (46) as a function (functional) of $\phi_k(s)$. However, if we continue to use the quasi-classical approximation to calculate the path integral Eq. (47), the arbitrary trajectory of the scalar fields $\phi_k(s)$ in Eq. (46) should be replaced by the classical trajectory of their motion (together with a and P_ε). In the theory of inflation [16], it is customary to consider this dynamics, starting from the stage when the potential energy $V(\phi)$ is maximum. However, we will move the time reference closer to the singularity, assuming that inflation was preceded by a stage of rapid change of fields, so,

$$\dot{\phi}_k^2 \gg V(\phi). \quad (50)$$

In this case,

$$\zeta \cong \frac{\sqrt{p_{\phi_k}^2}}{2\pi\sqrt{\dot{\phi}_k}}, \quad (51)$$

and the Schrödinger equation, corresponding in the quantum communication theory Eq. (48), takes the form:

$$i\hbar \frac{\partial \psi}{\partial \sigma} = \frac{\hbar}{2\pi} \sqrt{-\Delta_{\phi_k}} \psi, \quad (52)$$

where under the root sign is the Laplace operator in the configuration space of scalar fields, and

$$d\sigma = \sqrt{d\phi_k^2} \quad (53)$$

is an element of length in this space. Thus, the initial state of the universe near a singularity is described by a de Broglie plane wave in the configuration space of matter fields with initial momenta p_{ϕ_k} and energy

$$W = \|p_\phi\| = \sqrt{p_{\phi_k}^2}. \quad (54)$$

To find the energy of the space P_ε near the singularity, we simplify the consideration further by restricting ourselves to a single scalar field ϕ . From the equation of motion of this field (taking into account Eq. (50)),

$$\ddot{\phi} + 3\frac{\dot{a}}{a}\dot{\phi} = 0, \quad (55)$$

we find

$$\dot{\phi} = \frac{\Phi}{a^3}. \quad (56)$$

After this, from Eq. (36) we obtain

$$P_\varepsilon = \frac{\Phi^2}{a^3}. \quad (57)$$

Thus, at the beginning $a = 0$, we have a singularity: the energy of space there is infinite. It with expansion decreases according to Eq. (57). The energy of space reaches its minimum value at the boundary of the inflation stage when $(\dot{\phi})^2 = V(\phi)$. What follows is an exponential increase in the energy of space at the inflation stage. Starting from the indicated minimum value, the energy of the space P_ε can be considered as a physical parameter of time [12].

V. CONCLUSIONS

The variant of covariant quantum theory proposed in this paper and its subsequent modification with additional conditions on the dynamics of time-like degrees of freedom allow us to solve the problem of time in it. On the one hand, proper time is clearly added to the dynamic variables of the theory at the classical level. On the other hand, in quantum theory, this time disappears as a result of FS integration (the condition for the propagator covariance). Its dynamic meaning is restored by the conditions of the classical dynamics of time-like degrees of freedom. These conditions lead to the appearance of an additional dynamic variable - self-energy, for

which the classical law of its change at the classical and quantum level is formulated. Thus, the modification is focused here on time-like degrees of freedom of the system (coordinate x_0 of Minkowski space and radius a in the mini-superspace of the Friedmann universe) and represents a variant of combining the principles of covariance

and quantization.

VI. ACKNOWLEDGEMENTS

The authors would like to thank A. V. Goltsev and V. A. Franke for useful discussions.

-
- [1] E. S. Fradkin and G. A. Vilkovisky, *Phys. Lett.* **55B**, 224 (1975).
 - [2] I. A. Batalin and G. A. Vilkovisky, *Phys. Lett.* **69B**, 309 (1977).
 - [3] M. Henneaux, *Physics Reports* **126**, 1 (1985).
 - [4] T. Kugo and I. Ojima, *Suppl. Progr. Theor. Phys.* **66**, 1 (1979).
 - [5] F. R. Ore and P. van Nieuwenhuisen, *Nucl. Phys. B* **204**, 317 (1982).
 - [6] Jan Govaerts, A note of the Fradkin-Vilkovisky theorem, CERN-TH 5010/88 (1988).
 - [7] V. A. Fock, *Izv. Acad. Nauk SSSR*, 551 (1937).
 - [8] J. Schwinger, On gauge invariance and vacuum polarization, *Phys. Rev.* **82**, No 5, 664–679 (1951).
 - [9] B.S. De Witt, in General Relativity, An Einstein sesentary survey edited by S.W. Hawking and W. Israel, Cambridge University Press, Cambridge, 463 (1983).
 - [10] J. D. Bjorken, Sidney D.Drell, Relativistic quantum fields, Mc Graw-Hill Book Company (1978).
 - [11] C. W. Misner, K. S. Thorne and J. A. Wheeler, Gravitation, W. H. Freeman and Company, San Francisco (1973).
 - [12] N. N. Gorobey and A. S. Lukyanenko, St. Petersburg Polytechnical State University Journal *Physics and Mathematics* **11**, No 1, 147-156 (2018).
 - [13] L. D. Faddeev and A. A. Slavnov, Gauge fields: An introduction to quantum theory, Westview Press, 2d edition, 236 (1993).
 - [14] R. Arnoivitt, S. Deser and C. Misner, Dinamical Structure and Definition of Energy in General Relativity, *Phys. Rev.* **116**, No 5, 1322-1330 (1959).
 - [15] J. B. Hartle, S. W. Hawking and T. Hertog, arXiv:0803.1663v1 [hep-th] 11 Mar 2008.
 - [16] A. Linde, Inflation, Quantum Cosmology and the Anthropic Principle, In “Science and Ultimate Reality: From Quantum to Cosmos”, honoring John Wheeler’s 90th birthday. J. D. Barrow, P. C. W. Davies, & C. L. Harper eds. Cambridge University Press (2003).