

ARTICLE TEMPLATE

Inexact Relative Smoothness and Strong Convexity for Optimization and Variational Inequalities by Inexact Model.

Fedor Stonyakin^{a,b} and Alexander Tyurin^{b,d} and Alexander Gasnikov^{b,c,d} and Pavel Dvurechensky^{e,c} and Artem Agafonov^b and Darina Dvinskikh^{b,e} and Mohammad Alkousa^{d,b} and Dmitry Pasechnyuk^b and Sergei Artamonov^d and Victorya Piskunova^a

^aV. Vernadsky Crimean Federal University, Simferopol, Republic of Crimea; ^bMoscow Institute of Physics and Technology, Dolgoprudny, Russia; ^cInstitute for Information Transmission Problems, Moscow, Russia; ^dNational Research University Higher School of Economics, Moscow, Russia; ^eWeierstrass Institute for Applied Analysis and Stochastics, Berlin, Germany

ARTICLE HISTORY

Compiled December 21, 2021

ABSTRACT

In this paper, we propose a general algorithmic framework for first-order methods in optimization in a broad sense, including minimization problems, saddle-point problems, and variational inequalities. This framework allows obtaining many known methods as a special case, the list including accelerated gradient method, composite optimization methods, level-set methods, Bregman proximal methods. The idea of the framework is based on constructing an inexact model of the main problem component, i.e. objective function in optimization or operator in variational inequalities. Besides reproducing known results, our framework allows constructing new methods, which we illustrate by constructing a universal conditional gradient method and a universal method for variational inequalities with a composite structure. This method works for smooth and non-smooth problems with optimal complexity without a priori knowledge of the problem's smoothness. As a particular case of our general framework, we introduce relative smoothness for operators and propose an algorithm for variational inequalities (VIs) with such operators. We also generalize our framework for relatively strongly convex objectives and strongly monotone variational inequalities.

KEYWORDS

Convex optimization; composite optimization; proximal method; level-set method; variational inequality; universal method; mirror prox; acceleration; relative smoothness; saddle-point problem.

The work in Sections 2, 3.1, Definitions 4.1 and 4.6, Algorithm 3 and Appendix C was supported by the Ministry of Science and Higher Education of the Russian Federation (Goszadaniye in MIPT, project 075-00337-20-03). The work in Sections 3.2 and 3.3 was funded by Russian Foundation for Basic Research grant, project number 19-31-90062. The work in Definition 4.9 and Theorems 4.8, 4.11 was supported by the grant of the President of the Russian Federation, code MC-15.2020.1. The work in Section 5 was supported by Russian Science Foundation, project number 18-71-10044. The work in Section 6 was supported by Russian Foundation for Basic Research, project number 18-29-03071 mk

CONTACT Fedor Stonyakin. Email: fedyor@mail.ru

1. Introduction

In this paper we consider the following optimization problem

$$\min_{x \in Q} f(x), \quad (1)$$

where Q is a convex subset of finite-dimensional vector space E , f is generally a non-convex function.

Most minimization methods for such problems are constructed using some model of the objective f at the current iterate x_k . This can be a quadratic model based on the L -smoothness of the objective

$$f(x_k) + \langle \nabla f(x_k), x - x_k \rangle + \frac{L}{2} \|x - x_k\|_2^2. \quad (2)$$

The step of gradient method is obtained by the minimization of this model [59] (see the orange, dashed line in Fig. 1). More general models are constructed based on regularized second-order Taylor expansion [63] or other Taylor-like models [18] as well as other objective surrogates [47]. Another example is the conditional gradient method [27], where a linear model of the objective is minimized on every iteration. Adaptive choice of the parameter of the model with provably small computational overhead was proposed in [63] and applied to first-order methods in [24, 55, 56]. Recently, the first-order optimization methods were generalized to the so-called relative smoothness framework [7, 46, 64], where $\frac{1}{2}\|x - x_k\|_2^2$ in the quadratic model (2) for the objective is replaced by general Bregman divergence.

The literature on the first-order methods [9, 16, 23] considers also gradient methods with inexact information, relaxing the model (2) to

$$f_\delta(x_k) + \langle \nabla f_\delta(x_k), x - x_k \rangle + \frac{L}{2} \|x - x_k\|_2^2 + \delta, \quad (3)$$

with $(f_\delta, \nabla f_\delta)$ called inexact oracle and this model being an upper bound for the objective. In particular, this relaxation allows to obtain universal gradient methods [56].

One of the goals of this paper is to describe and analyze the first-order optimization methods which use a very general *inexact model* of the objective function, the idea being to replace the linear part in (3) by a general function $\psi_\delta(x, x_k)$ and the squared norm by general Bregman divergence (see the red, dotted line in Fig. 1). The resulting model includes as a particular case inexact oracle model and relative smoothness framework and allows one to obtain many optimization methods as a particular case, including conditional gradient method [27], Bregman proximal gradient method [11] and its application to optimal transport [75] and Wasserstein barycenter [71] problems, general Catalyst acceleration technique [45], (accelerated) composite gradient methods [8, 55], (accelerated) level and bundle-type methods [43, 53, 61], gradient methods in the relative smoothness framework [7, 46]. First attempts to propose inexact model generalization were made in [30, 71] for non-accelerated methods and in [32] for accelerated methods, yet without a relative smoothness paradigm. In this paper, we propose the inexact model in a very general setting, including adaptivity of the algorithms to the parameter L , possible relative strong convexity, and relative smoothness. We also provide convergence rates for the gradient method and acceler-




Figure 1.: **Orange**, dashed line and x_{k+1} illustrate the standard upper-bound and a step of the classical gradient method. **Red**, dotted line and \tilde{x}_{k+1} illustrate the upper-bound based on our notion of inexact model and a step of the proposed method. We can see that with better upper-bound, the next iterate is closer to the minimum point.

ated gradient method using an inexact model of the objective. As an application of our general framework, we develop a universal conditional gradient method, providing a parameter-free generalization of the results in [57].

We believe that our model is flexible enough to be extended for problems with primal-dual structure² [52, 54, 57], e.g. for problems with linear constraints [2, 12, 35, 62]; for random block-coordinate descent [26]; for tensor methods [31, 58]; for distributed optimization setting [19, 20, 68, 74]; and adaptive stochastic optimization [37, 65].

Optimization problem (1) is tightly connected with variational inequality (VI)

$$\text{Find } x_* \in Q \text{ s.t. } \langle g(x_*), x_* - x \rangle \leq 0, \forall x \in Q, \quad (4)$$

where $g(x) = \nabla f(x)$. A special VI is also equivalent to finding a saddle-point of a convex-concave function

$$\min_{u \in Q_1} \max_{v \in Q_2} f(u, v) \quad (5)$$

for $x = (u, v)$ and $g(x) = (\nabla_u f(u, v), -\nabla_v f(u, v))$. This motivates the second part of this paper, which consists of the generalization of the inexact model of the objective function to an inexact model for an operator in variational inequality. In particular, we extend the relative smoothness paradigm to variational inequalities with monotone and strongly monotone operators and provide a generalization of the Mirror-Prox method [51], its adaptive version [29] (see also [3, 4, 48, 49] for other types of adaptive extra-gradient methods), and universal version [69] to variational inequalities with

²see recent results on this generalization in [73].

such general inexact model of the operator. As a particular case, our approach allows us to partially reproduce the results of [10]. We also apply the general framework for variational inequalities to saddle-point problems.

To sum up, we present a unified view on inexact models for optimization problems, variational inequalities, and saddle-point problems.

The structure of the paper is the following. In Section 2, we introduce an inexact model of the objective in optimization and provide several examples to illustrate the flexibility and generality of the proposed model. In particular, we demonstrate that relative smoothness and strong convexity are the special cases of our general framework.

In Section 3, we consider an adaptive gradient method (GM) and an adaptive fast gradient method (FGM). FGM has a better convergence rate yet is not adapted to the relative smoothness paradigm. In Subsection 3.3, we construct a universal conditional gradient (Frank–Wolfe) method using FGM with inexact projection. To the best of our knowledge, this is the first attempt to combine the Frank–Wolfe method [36, 38] and the universal method [56]. In Section 4, we generalize the inexact model to variational inequalities and saddle-point problems for the case of monotone and strongly monotone operators. In the former case, we construct an adaptive generalization of the Mirror-Prox algorithm for variational inequalities and saddle-point problems with such an inexact model. In the latter case, the proposed algorithm is accelerated by the restart technique to have the linear rate of convergence. We especially consider the case of m -strong convexity of the model. The natural motivations for such a formulation are composite saddle-point problems and mixed variational inequalities with an m -strongly convex composite.

The contribution of this paper is follows:

- (1) We introduce an inexact (δ, L, μ, m, V) -model for optimization problems and obtain convergence rates for adaptive GM for optimization problems with this model.
- (2) We introduce an inexact $(\delta, L, \mu, m, V, \|\cdot\|)$ -model for optimization problems and obtain convergence rates for adaptive FGM for optimization problems with such a model. Using FGM with the inexact model we construct a universal conditional gradient (Frank–Wolfe) method.
- (3) We propose generalizations of the above models, namely (δ, L, V) -model and (δ, L, μ, V) -model for variational inequalities and saddle-point problems. As a special case, we introduce relative smoothness for operators in variational inequalities, thus, generalizing [46] from optimization to variational inequalities. We obtain convergence rates for adaptive versions of the Mirror-Prox algorithm for problems with an inexact model. In Example 4.6, we show that our method is able to solve variational inequalities with singularities, for which the application of existing Mirror-Prox algorithms is not theoretically justified.³

To demonstrate the practical performance of the proposed algorithms, several numerical experiments were performed for different examples. First, in Appendix D, we consider a constrained convex optimization problem with non-smooth functional constraints motivated by three different geometrical problems: an analogue of the *smallest covering circle problem*, the *problem of the best approximation of the distance from the given set*, and the *Fermat–Torricelli–Steiner problem*. Further, in Appendix E, we test our algorithms on a minimization problem of a polynomial-like function, which

³As an exception we mention works [3, 4] that appeared after the first version of this paper appeared as a preprint [70].

is shown in [46] to be relatively smooth, under non-smooth functional constraints. This is followed by numerical experiments in Appendix F for the D-optimal design problem with the relatively smooth objective and simplex constraints, which has wide applications in statistics and computational geometry. We report the advantages of the proposed method and the possibility to control the convergence of the method by adjusting the line-search parameter. Finally, in Appendix G we describe the results of the experiments for the resource sharing problem considered as an example in [4].

2. Inexact Model in Minimization Problems. Definitions and Examples

In this section, we introduce a general notation which is used in the paper and our main definition of the inexact model for optimization problems. To motivate this definition and illustrate its generality, we give several examples of known situations which are covered by our inexact model.

As said, we start with the general notation. Let E be an n -dimensional real vector space and E^* be its dual. We denote the value of a linear function $g \in E^*$ at $x \in E$ by $\langle g, x \rangle$. Let $\|\cdot\|$ be some norm on E , $\|\cdot\|_*$ be its dual, defined by $\|g\|_* = \max_x \{\langle g, x \rangle, \|x\| \leq 1\}$. We use $\nabla f(x)$ to denote the gradient of a differentiable function f at a point $x \in \text{dom} f$ and, with a slight abuse of notation, a subgradient of a convex function f at a point $x \in \text{dom} f$. The whole subdifferential of a convex function f at a point $x \in \text{dom} f$ is denoted by $\partial f(x)$. Let d be a convex function on Q , which is continuously differentiable on the relative interior $\text{ri}Q$ of Q . Let $V[y](x) = d(x) - d(y) - \langle \nabla d(y), x - y \rangle$, $x \in Q, y \in \text{ri}Q$ be the corresponding Bregman divergence. Most typically it is assumed that d is 1-strongly convex on Q w.r.t. $\|\cdot\|$ -norm, which we refer to as (1-SC) assumption w.r.t. $\|\cdot\|$ -norm. More precisely, this assumption means that, for all $x \in Q, y \in \text{ri}Q$, $d(x) - d(y) - \langle \nabla d(y), x - y \rangle \geq \frac{1}{2}\|x - y\|^2$. We underline that, in general, we do not make this assumption, and, in what follows, we explicitly write if this assumption is made.

Definition 2.1 (Inexact model in optimization). Let $\delta, L, \mu, m \geq 0$. We say that $\psi_\delta(x, y)$ is a (δ, L, μ, m, V) -model of the function f at a given point y iff, for all $x \in Q$,

$$\mu V[y](x) \leq f(x) - (f_\delta(y) + \psi_\delta(x, y)) \leq LV[y](x) + \delta, \quad (6)$$

$\psi_\delta(x, y)$ is convex in x , satisfies $\psi_\delta(x, x) = 0$ for all $x \in Q$ and

$$\psi(x) \geq \psi(z) + \langle g, x - z \rangle + mV[z](x), \quad \forall x, z \in Q, \forall g \in \partial\psi(z), \quad (7)$$

where for fixed $y \in Q$ and any $x \in Q$ we denote $\psi(x) = \psi_\delta(x, y)$. In other words, $\psi(x)$ is relatively m -strongly convex relative to d .

Note that in Definition 2.1 we allow L to depend on δ . Definition 2.1 is a generalization of (δ, L) -model from [30, 32, 71], where $\mu = 0$ and $m = 0$. Below, we denote $(\delta, L, 0, 0, V)$ -model as (δ, L) -model. Thus, our framework allows obtaining (accelerated) gradient methods for composite optimization and their counterparts for inexact oracle models.

Let us illustrate the above definition by several examples.

Example 2.2. Relative smoothness and relative strong convexity, [7, 46]. Assume that $d(x)$ is differentiable, and in (1), the objective f is differentiable and

relatively smooth [7, 46] relative to d , i.e.

$$f(x) - f(y) - \langle \nabla f(y), x - y \rangle \leq (d(x) - d(y) - \langle \nabla d(y), x - y \rangle) = LV[y](x), \quad \forall x, y \in Q$$

and relatively strongly convex [46] relative to d , i.e.

$$\mu V[y](x) = \mu (d(x) - d(y) - \langle \nabla d(y), x - y \rangle) \leq f(x) - f(y) - \langle \nabla f(y), x - y \rangle, \quad \forall x, y \in Q.$$

Then, clearly, Definition 2.1 holds with $m = 0$, $\delta = 0$, $\psi_\delta(x, y) = \langle \nabla f(y), x - y \rangle$. Importantly, the function d is not necessarily strongly convex.

Note that if $V[y](x) \leq C_n \|x - y\|^2$ for some constant $C_n = O(\log n)$, the condition of (μC_n) -strong convexity w.r.t. norm $\|\cdot\|$, namely $\mu C_n \|x - y\|^2 + f_\delta(y) + \psi_\delta(x, y) \leq f(x)$ implies the left inequality in (6).

One of the main applications of general relative smoothness and strong convexity is the step of tensor methods which use the derivatives of the objective of the order higher than 2 [31, 58]. Thus, our framework allows obtaining gradient method for optimization with relative smoothness and strong convexity and extending them to the case of inexact oracle setting.

Example 2.3. Composite optimization, [8, 55]. Assume that in (1), $f(x) = g(x) + h(x)$ with L -smooth w.r.t. norm $\|\cdot\|$ term g and simple convex term h , which is usually called composite. In this case we assume that $V[y](x)$ satisfies (1-SC) condition w.r.t $\|\cdot\|$, and define $f_\delta(x) = g(x) + h(x)$ and $\psi_\delta(x, y) = \langle \nabla g(y), x - y \rangle + h(x) - h(y)$. We have

$$\begin{aligned} f(x) - (f_\delta(y) + \psi_\delta(x, y)) &= g(x) + h(x) - (g(y) + h(y) + \langle \nabla g(y), x - y \rangle + h(x) - h(y)) \\ &= g(x) - (g(y) + \langle \nabla g(y), x - y \rangle). \end{aligned}$$

By convexity of g , we have $0 \leq g(x) - (g(y) + \langle \nabla g(y), x - y \rangle)$. At the same time, by the L -smoothness of g and (1-SC) condition for $V[y](x)$,

$$g(x) - (g(y) + \langle \nabla g(y), x - y \rangle) \leq \frac{L}{2} \|x - y\|^2 \leq LV[y](x).$$

From the combination of the above three relations, it is clear that (6) holds with $\delta = 0$ and $\mu = 0$ and we are in the situation of Definition 2.1 with $m = 0$ since $\psi_\delta(x, y)$ is convex in x . If h turns out to be differentiable and relatively m -strongly convex relative to d [46], i.e. $h(x) - h(y) - \langle \nabla h(y), x - y \rangle \geq mV[y](x)$, then (7) holds, but in (6) we have $\mu = 0$. On the other hand, if g turns out to be relatively μ -strongly convex relative to d [46], i.e. $g(x) - g(y) - \langle \nabla g(y), x - y \rangle \geq \mu V[y](x)$, then (6) holds with $\delta = 0$, but in (7) we have $m = 0$.

A particular example is the following minimization problem [1] motivated by traffic demands matrix estimation from link loads:

$$f(x) = \frac{1}{2} \|Ax - b\|_2^2 + m \sum_{k=1}^n x_k \ln x_k \rightarrow \min_{x \in S_n(1)},$$

where $S_n(1) = \{x \in \mathbb{R}_+^n \mid \sum_{i=1}^n x_i = 1\}$ is the standard unit simplex in \mathbb{R}^n . In this case $g(x) = \frac{1}{2} \|Ax - b\|_2^2$ and $h(x) = m \sum_{k=1}^n x_k \ln x_k$. If we choose $\|\cdot\| = \|\cdot\|_1$ and

$d(x) = \sum_{k=1}^n x_k \ln x_k$, then $d(x)$ satisfies (1-SC) condition, $V[y](x) = \sum_{k=1}^n x_k \ln(x_k/y_k)$, and $h(x)$ is relatively m -strongly convex relative to d . Since g is a quadratic function, by using the fact that Lipschitz gradient is equivalent to the bounded Hessian, we obtain that g has a Lipschitz-continuous gradient w.r.t. $\|\cdot\|_1$ with the constant $L = \max_{\|h\|_1 \leq 1} \langle h, A^T A h \rangle = \max_{k=1, \dots, n} \|A_k\|_2^2$, where A_k is the k -th column of A . Thus, we obtain that $\psi_\delta(x, y) = \langle \nabla g(y), x - y \rangle + h(x) - h(y)$ is a $(0, L, 0, m, V)$ -model. At the same time, the part g is not necessarily strongly convex.

Example 2.4. Superposition of functions, [53]. Assume that in (1) [43, 53] $f(x) := g(g_1(x), \dots, g_m(x))$, where each function $g_k(x)$ is a smooth convex function with L_k -Lipschitz gradient w.r.t. $\|\cdot\|$ -norm for all k . The function $g(x)$ is an M -Lipschitz convex function w.r.t 1-norm, non-decreasing in each of its arguments. The chosen Bregman divergence $V[y](x)$ is assumed to satisfy (1-SC). We define $f_\delta(y) := f(y)$ and

$$\psi_\delta(x, y) = g(g_1(y) + \langle \nabla g_1(y), x - y \rangle, \dots, g_m(y) + \langle \nabla g_m(y), x - y \rangle) - f(y).$$

By convexity and L_k -smoothness of each $g_k(x)$, we have, for all $k = 1, \dots, m$

$$g_k(y) + \langle \nabla g_k(y), x - y \rangle \leq g_k(x) \leq g_k(y) + \langle \nabla g_k(y), x - y \rangle + \frac{L_k}{2} \|y - x\|^2.$$

By monotonicity and Lipschitz-continuity of g w.r.t. 1-norm, we have

$$\begin{aligned} & g(g_1(y) + \langle \nabla g_1(y), x - y \rangle, \dots, g_m(y) + \langle \nabla g_m(y), x - y \rangle) \leq g(g_1(x), \dots, g_m(x)) \\ & \leq g \left(g_1(y) + \langle \nabla g_1(y), x - y \rangle + \frac{L_1}{2} \|y - x\|^2, \dots, g_m(y) + \langle \nabla g_m(y), x - y \rangle + \frac{L_m}{2} \|y - x\|^2 \right) \\ & \leq g(g_1(y) + \langle \nabla g_1(y), x - y \rangle, \dots, g_m(y) + \langle \nabla g_m(y), x - y \rangle) + M \frac{\sum_{i=1}^m L_i}{2} \|x - y\|^2. \end{aligned}$$

From this, by the (1-SC) condition for $V[y](x)$, we obtain

$$\begin{aligned} 0 & \leq f(x) - (f(y) + g(g_1(y) + \langle \nabla g_1(y), x - y \rangle, \dots, g_m(y) + \langle \nabla g_m(y), x - y \rangle) - f(y)) \\ & \leq M \frac{\sum_{i=1}^m L_i}{2} \|x - y\|^2 \leq MV[y](x) \sum_{i=1}^m L_i, \quad \forall x, y \in Q, \end{aligned}$$

which is (6) with $\mu = 0$ and $\delta = 0$ given in the above definition of $f_\delta(y)$ and $\psi_\delta(x, y)$. Clearly, $\psi_\delta(x, y)$ is convex in x as a composition of non-decreasing convex function g with linear functions. Thus, (7) holds with $m = 0$, and we obtain a $(0, M \cdot (\sum_{i=1}^m L_i))$ -model of f . As a result, our framework allows to obtain (accelerated) level gradient methods considered in [43, 53] as a special case. Moreover, we generalize these methods for the case of inexact oracle information.

Example 2.5. Proximal method, [11]. Let us consider the optimization problem (1), where f is an arbitrary convex function (not necessarily smooth). Then, for arbitrary $L \geq 0$, $\psi_\delta(x, y) = f(x) - f(y)$ is $(0, L)$ -model of f with $f_\delta(y) = f(y)$ at a given point y . Thus, our framework allows obtaining (Bregman) proximal gradient methods [11, 67] as a special case and extends them to the case of inexact oracle setting. In

particular, based on this model (with Bregman divergence to be Kullback–Leibler divergence), we propose in [71] proximal Sinkhorn’s algorithm for Wasserstein distance calculation problem and in [40] proximal IBP for Wasserstein barycenter problem.

Example 2.6. Min-min problem. Assume that in (1) $f(x) := \min_{z \in Z} F(z, x)$, where the set Z is convex and bounded, the function F is smooth and convex w.r.t. all variables. Moreover, assume that

$$\|\nabla F(z', x') - \nabla F(z, x)\|_2 \leq L \|(z', x') - (z, x)\|_2, \quad \forall z, z' \in Z, x, x' \in \mathbb{R}^n.$$

Let $V[y](x) = \frac{1}{2}\|x - y\|_2^2$. Assume that for any $y \in Q$ it is possible to find an approximate solution of $\min_{z \in Z} F(z, x)$, i.e. a point $\tilde{z}_\delta(y) \in Z$ such that

$$\langle \nabla_z F(\tilde{z}_\delta(y), y), z - \tilde{z}_\delta(y) \rangle \geq -\delta, \quad \forall z \in Z.$$

Then $F(\tilde{z}_\delta(y), y) - f(y) \leq \delta$ and $\psi_\delta(x, y) = \langle \nabla_z F(\tilde{z}_\delta(y), y), x - y \rangle$ is $(6\delta, 2L, 0, 0, V)$ -model of f with $f_\delta(y) = F(\tilde{z}_\delta(y), y) - 2\delta$ at a given point y .

Example 2.7. Inexact oracle, [16]. We say that f is equipped with (δ, L) -oracle, if for any $y \in Q$ we can find a pair $(f_\delta(y), \nabla f_\delta(y))$ such that

$$0 \leq f(x) - (f_\delta(y) + \langle \nabla f_\delta(y), x - y \rangle) \leq \frac{L}{2} \|x - y\|^2 + \delta,$$

for all $x \in Q$ (see [16]). Clearly, under the assumption (1-SC), the function $\langle \nabla f_\delta(y), x - y \rangle$ is a (δ, L) -model. Examples of inexact models in 2.8, 2.9, 2.10 are also described by inexact (δ, L) -oracle. The detailed proofs can be found in [16].

Example 2.8. Saddle-point problem, [16]. Assume that in (1) $Q = \mathbb{R}^n$, $f(x) = \max_{z \in Z} [\langle x, b - Az \rangle - \phi(z)]$, where $\phi(z)$ is a μ -strongly convex function w.r.t. p -norm ($1 \leq p \leq 2$). Then f is smooth and convex and its gradient is Lipschitz continuous with constant $L = \frac{1}{\mu} \max_{\|z\|_p \leq 1} \|Az\|_2^2$. If $z_\delta(y) \in Z$ is an approximate solution to the auxiliary max-problem, i.e.

$$\max_{z \in Z} [\langle y, b - Az \rangle - \phi(z)] - [\langle y, b - Az_\delta(y) \rangle - \phi(z_\delta(y))] \leq \delta,$$

then $\psi_\delta(x, y) = \langle b - Az_\delta(y), x - y \rangle$ is $(\delta, 2L, 0, 0, V)$ -model of f with $f_\delta(y) = \langle y, b - Az_\delta(y) \rangle - \phi(z_\delta(y))$ at the point y if we define $V[y](x) = \frac{1}{2}\|x - y\|_2^2$.

Example 2.9. Augmented Lagrangians, [16]. Let $\phi(z)$ be a smooth function. Consider the problem

$$\min_{Az=b, z \in Q} \phi(z) + \frac{\mu}{2} \|Az - b\|_2^2$$

and the corresponding dual problem

$$\min_{x \in \mathbb{R}^n} \left\{ f(x) = \max_{z \in Q} \underbrace{\left(\langle x, b - Az \rangle - \phi(z) - \frac{\mu}{2} \|Az - b\|_2^2 \right)}_{\Lambda(x, z)} \right\}.$$

If $z_\delta(y)$ is an approximate solution of auxiliary max-problem, i.e.

$$\max_{z \in Q} \langle \nabla_z \Lambda(y, z_\delta(y)), z - z_\delta(y) \rangle \leq \delta,$$

then $\psi_\delta(x, y) = \langle b - Az_\delta(y), x - y \rangle$ is $(\delta, \mu^{-1}, 0, 0, V)$ -model of f with

$$f_\delta(y) = \langle y, b - Az_\delta(y) \rangle - \phi(z_\delta(y)) - \frac{\mu}{2} \|Az_\delta(y) - b\|_2^2$$

at the point y if we take $V[y](x) = \frac{1}{2} \|x - y\|_2^2$.

Example 2.10. Moreau envelope of the objective function, [16]. Let us consider the following optimization problem:

$$\min_{x \in \mathbb{R}^n} \left\{ f_L(x) := \min_{z \in Q} \underbrace{\left\{ f(z) + \frac{L}{2} \|z - x\|_2^2 \right\}}_{\Lambda(x, z)} \right\}. \quad (8)$$

Assume that f is convex and, for some $z_L(y)$,

$$\max_{z \in Q} \left\{ \Lambda(y, z_L(y)) - \Lambda(y, z) + \frac{L}{2} \|y - z_L(y)\|_2^2 \right\} \leq \delta.$$

Then $\psi_\delta(x, y) = \langle L(y - z_L(y)), x - y \rangle$ is $(\delta, L, 0, 0, V)$ -model of f with

$$f_\delta(y) = f(z_L(y)) + \frac{L}{2} \|z_L(y) - y\|_2^2 - \delta$$

at the point y if we take $V[y](x) = \frac{1}{2} \|x - y\|_2^2$.

Example 2.11. Clustering by Electoral Model, [71]. Another example of an optimization problem that allows for $(\delta, L, 0, m, V)$ -model with strong convexity of the function $\psi_\delta(x, y)$ is proposed in [71] to address a *non-convex* optimization problem which arises in an electoral model for clustering introduced in [60]. In this model, voters (data points) select a party (cluster) iteratively by minimizing the following function

$$\min_{z \in S_n(1), p \in \mathbb{R}_+^m} \left\{ f_{\mu_1, \mu_2}(x = (z, p)) = g(x) + \mu_1 \sum_{k=1}^n z_k \ln z_k + \frac{\mu_2}{2} \|p\|_2^2 \right\}$$

where $S_n(1)$ is the standard unit simplex in \mathbb{R}^n . Let us choose $\|x\|^2 = \|z\|_1^2 + \|p\|_2^2$ and

assume that, in general non-convex, $g(x)$ has L_g -Lipschitz continuous gradient

$$\|\nabla g(x) - \nabla g(y)\|_* \leq L_g \|x - y\| \quad \forall x, y \in S_n(1) \times \mathbb{R}_+^m$$

and $L_g \leq \mu_1$, and $L_g \leq \mu_2$. It can be shown (see [71]) that

$$\begin{aligned} \psi_\delta(x, y) &= \langle \nabla g(y), x - y \rangle - L_g \cdot \text{KL}(z_x|z_y) - \frac{L_g}{2} \|p_x - p_y\|_2^2 \\ &\quad + \mu_1(\text{KL}(z_x|\mathbf{1}) - \text{KL}(z_y|\mathbf{1})) + \frac{\mu_2}{2} (\|p_x\|_2^2 - \|p_y\|_2^2) \end{aligned}$$

is a $(0, 2L_g, 0, \min\{\mu_1, \mu_2\} - L_g, V)$ -model of $f_{\mu_1, \mu_2}(x)$. Here $\text{KL}(z_x|z_y) = \sum_{i=1}^m [z_x]_i \ln([z_x]_i/[z_y]_i)$ and

$$V[y](x) = \text{KL}(z_x|z_y) + \frac{1}{2} \|p_x - p_y\|_2^2.$$

We finish this section by defining an approximate solution to an optimization problem. This definition will be used to allow inexact solutions of auxiliary minimization problems in each iteration of our algorithms.

Definition 2.12. For a convex optimization problem $\min_{x \in Q} \Psi(x)$ and $\tilde{\delta} \geq 0$, we denote by $\text{Arg min}_{x \in Q}^{\tilde{\delta}} \Psi(x)$ a set of points \tilde{x} such that

$$\exists h \in \partial \Psi(\tilde{x}): \forall x \in Q \rightarrow \langle h, x - \tilde{x} \rangle \geq -\tilde{\delta}. \quad (9)$$

We denote by $\text{argmin}_{x \in Q}^{\tilde{\delta}} \Psi(x)$ some element of $\text{Arg min}_{x \in Q}^{\tilde{\delta}} \Psi(x)$.

3. Gradient Method with Inexact Model.

In this section, we consider adaptive gradient-type methods for problems with the (δ, L, μ, m, V) -model of the objective. First, we consider the non-accelerated gradient method and then its accelerated version. For each algorithm, we obtain convergence rates that have two terms. The first term decreases as iterations go and corresponds to the case of the exact model with $\delta = 0$ and the exact solution of the auxiliary problems with $\tilde{\delta} = 0$. The second term is non-decreasing as iterations go and corresponds to errors in the model and the solutions of the auxiliary problems. Non-accelerated Algorithm 1 has a slower decaying part, but the constant term depending on errors $\delta, \tilde{\delta}$. Accelerated Algorithm 2 has a faster decaying term, but increasing term depending on errors $\delta, \tilde{\delta}$. Moreover, to obtain acceleration, we have to consider Algorithm 2 for a narrower class of problems with $(\delta, L, \mu, m, V, \|\cdot\|)$ -models (for a precise definition, see Definition 3.3).

3.1. Adaptive Gradient Method with (δ, L, μ, m, V) -Model

In this section, we consider the adaptive gradient method for problem (1), which uses a (δ, L, μ, m, V) -model of the objective. For the case when $\mu + m > 0$, our method has a linear convergence, and for a more general case $\mu = 0$ and $m = 0$, we prove a sublinear convergence rate.

We assume that in each iteration k , the method has access to $(\delta, \bar{L}_{k+1}, \mu, m, V)$ -model of f (see Definition 2.1). In general, constant \bar{L}_{k+1} may vary from iteration to iteration and we only assume that the $(\delta, \bar{L}_{k+1}, \mu, m, V)$ -model exists. We do not use \bar{L}_{k+1} in Algorithm 1 explicitly and, moreover, our method is adaptive to this constant. Constant $\tilde{\delta}$ is an error in terms of Definition 2.12 and there is no need to know it explicitly in Algorithm 1, $\tilde{\delta}$ only appears in Theorem 3.1.

Algorithm 1 Adaptive gradient method with (δ, L, μ, m, V) -model

- 1: **Input:** x_0 is the starting point, $\delta > 0$ is the oracle error and $L_0 > 0$.
- 2: Set $S_0 := 0$
- 3: **for** $k \geq 0$ **do**
- 4: Find the smallest integer $i_k \geq 0$ such that

$$f_\delta(x_{k+1}) \leq f_\delta(x_k) + \psi_\delta(x_{k+1}, x_k) + L_{k+1}V[x_k](x_{k+1}) + \delta, \quad (10)$$

where $L_{k+1} = 2^{i_k-1}L_k$, $\alpha_{k+1} := \frac{1}{L_{k+1}}$, $S_{k+1} := S_k + \alpha_{k+1}$, and, for some $\tilde{\delta} \geq 0$

$$\phi_{k+1}(x) := \psi_\delta(x, x_k) + L_{k+1}V[x_k](x), \quad x_{k+1} := \arg \min_{x \in Q}^{\tilde{\delta}} \phi_{k+1}(x). \quad (11)$$

Here the inexact minimization is in accordance with Definition 2.12.

- 5: **end for**
-

For $L_k > \mu$ and all $k \geq 0$, we denote

$$q_k \stackrel{\text{def}}{=} \frac{L_k - \mu}{L_k + m} = 1 - \frac{m + \mu}{L_k + m} \leq 1,$$

and $q_k = 0$ for $L_k \leq \mu$. We assume that $Q_j^k \stackrel{\text{def}}{=} \prod_{i=j}^k q_i$ and $Q_j^k = 1$ for $j > k$.

Theorem 3.1. *Assume that $\psi_\delta(x, y)$ is a (δ, L, μ, m, V) -model according to Definition 2.1. Denote by $y_N = x_{j(N)}$, where $j(N) = \operatorname{argmin}_{k=1, \dots, N} f_\delta(x_k)$. Then, after N iterations of Algorithm 1, we have*

$$f(y_N) - f(x_*) \leq \min \left\{ (L_N + m)Q_1^N, \frac{1}{\sum_{i=1}^N \frac{1}{L_i + m}} \right\} V[x_0](x_*) + \tilde{\delta} + 3\delta, \quad (12)$$

$$V[x_N](x_*) \leq Q_1^N V[x_0](x_*) + (\tilde{\delta} + 2\delta) \sum_{i=1}^N \frac{Q_{i+1}^N}{L_i + m}. \quad (13)$$

To prove Theorem 3.1 we need the following lemma.

Lemma 3.2. *Let $\psi(x)$ be a relatively m -strongly convex function relative to d with $m \geq 0$, i.e. (7) holds, and*

$$y = \operatorname{argmin}_{x \in Q}^{\tilde{\delta}} \{\psi(x) + \beta V[z](x)\}, \quad (14)$$

where $\beta \geq 0$. Then

$$\psi(x) + \beta V[z](x) \geq \psi(y) + \beta V[z](y) + (\beta + m)V[y](x) - \tilde{\delta}, \quad \forall x \in Q.$$

Proof. By optimality conditions in (14) and Definition 2.12,

$$\exists g \in \partial\psi(y), \quad \langle g + \beta \nabla_y V[z](y), x - y \rangle \geq -\tilde{\delta}, \quad \forall x \in Q.$$

From this we obtain

$$\psi(x) - \psi(y) \stackrel{(7)}{\geq} \langle g, x - y \rangle + mV[y](x) \geq \langle \beta \nabla_y V[z](y), y - x \rangle - \tilde{\delta} + mV[y](x),$$

which together with equality

$$\begin{aligned} \langle \nabla_y V[z](y), y - x \rangle &= \langle \nabla d(y) - \nabla d(z), y - x \rangle = d(y) - d(z) - \langle \nabla d(z), y - z \rangle + \\ &\quad + d(x) - d(y) - \langle \nabla d(y), x - y \rangle - d(x) + d(z) + \langle \nabla d(z), x - z \rangle = \\ &= V[z](y) + V[y](x) - V[z](x) \end{aligned}$$

complete the proof. The latter chain of equalities follows from the definition of Bregman divergence, see the beginning of Section 2. \square

Proof of Theorem 3.1. Since by Definition 2.1 with $x = y$, we have

$$f(x) - \delta \leq f_\delta(x) \leq f(x), \quad (15)$$

and using (10), we have the following series of inequalities

$$f(x_N) \leq f_\delta(x_N) + \delta \stackrel{(10)}{\leq} f_\delta(x_{N-1}) + \psi_\delta(x_N, x_{N-1}) + L_N V[x_{N-1}](x_N) + 2\delta.$$

Using Lemma 3.2 for (11) with $y = x_N$, $\beta = L_N$, $z = x_{N-1}$, $\psi(x) = \psi_\delta(x, x_{N-1})$, we have

$$f(x_N) \leq f_\delta(x_{N-1}) + \psi_\delta(x, x_{N-1}) + L_N V[x_{N-1}](x) - (L_N + m) V[x_N](x) + \tilde{\delta} + 2\delta.$$

Further, applying the left inequality (6), we have

$$f(x_N) \leq f(x) + (L_N - \mu) V[x_{N-1}](x) - (L_N + m) V[x_N](x) + \tilde{\delta} + 2\delta. \quad (16)$$

Let us first consider the case $L_N > \mu$. Taking $x = x_*$ in the above inequality and using inequality $f(x_*) \leq f(x_N)$, we obtain

$$(L_N + m) V[x_N](x_*) \leq (L_N - \mu) V[x_{N-1}](x_*) + \tilde{\delta} + 2\delta. \quad (17)$$

Thus, recursively we have that

$$V[x_N](x_*) \leq q_N V[x_{N-1}](x_*) + \frac{\tilde{\delta} + 2\delta}{L_N + m} \leq Q_1^N V[x_0](x_*) + (\tilde{\delta} + 2\delta) \sum_{i=1}^N \frac{Q_{i+1}^N}{L_i + m}.$$

The last inequality proves (13). Now we rewrite (16) for $x = x_*$ as

$$V[x_N](x_*) \leq \frac{1}{L_N + m}(f(x_*) - f(x_N) + \tilde{\delta} + 2\delta) + q_N V[x_{N-1}](x_*).$$

Recursively, we have

$$V[x_N](x_*) \leq \sum_{i=1}^N \left(\frac{Q_{i+1}^N}{L_i + m}(f(x_*) - f(x_i) + \tilde{\delta} + 2\delta) \right) + Q_1^N V[x_0](x_*).$$

From the definition of y_N and (15), we obtain

$$f(y_N) - \delta \leq f_\delta(y_N) \leq f_\delta(x_k) \leq f(x_k), \quad \forall k \in [1, N]. \quad (18)$$

Using that $V[x_N](x_*) \geq 0$ and (18), we get

$$\begin{aligned} Q_1^N V[x_0](x_*) &\geq \sum_{i=1}^N \left(\frac{Q_{i+1}^N}{L_i + m}(f(x_i) - f(x_*) - \tilde{\delta} - 2\delta) \right) \\ &\geq (f(y_N) - f(x_*)) \sum_{i=1}^N \frac{Q_{i+1}^N}{L_i + m} - (\tilde{\delta} + 3\delta) \sum_{i=1}^N \frac{Q_{i+1}^N}{L_i + m}. \end{aligned}$$

Dividing by $\sum_{i=1}^N \frac{Q_{i+1}^N}{L_i + m}$, we obtain

$$f(y_N) - f(x_*) \leq \frac{Q_1^N}{\sum_{i=1}^N \frac{Q_{i+1}^N}{L_i + m}} V[x_0](x_*) + \tilde{\delta} + 3\delta.$$

Since $\sum_{i=1}^N \frac{Q_{i+1}^N}{L_i + m} \geq \frac{1}{L_N + m}$ and $Q_i^N \geq Q_1^N$ for all $i \geq 1$, we get

$$f(y_N) - f(x_*) \leq \min \left\{ (L_N + m)Q_1^N, \frac{1}{\sum_{i=1}^N \frac{1}{L_i + m}} \right\} V[x_0](x_*) + \tilde{\delta} + 3\delta.$$

This proves (12).

In the opposite case $L_N \leq \mu$, we obtain from (18), (17) and (16) with $x = x_*$ that

$$\begin{aligned} V[x_N](x_*) &\leq \frac{\tilde{\delta} + 2\delta}{L_N + m} \\ f(y_N) &\leq f(x_N) + \delta \leq f(x_*) + \tilde{\delta} + 3\delta. \end{aligned}$$

The r.h.s. of these two inequalities can be bounded from above by respectively the r.h.s. of (13) and (12), which proves the latter inequalities. \square

Remark 1. Let us assume that $L_0 \leq L$, and we know that $\bar{L}_{k+1} \leq L$ for all $k \geq 0$ (or in other words, (δ, L, μ, m, V) -model exists for all $k \geq 0$). This means that if L_k in Algorithm 1 satisfies $L_k \geq L$, then inequality (10) holds. Thus, $L_k \leq 2L$ for all $k \geq 0$.

From this fact we can obtain that $\sum_{i=1}^N \frac{1}{L_i+m} \geq \frac{N}{2L+m}$ and $q_k \leq q \stackrel{\text{def}}{=} \frac{2L-\mu}{2L+m} = 1 - \frac{m+\mu}{2L+m}$. In view of (12) and (13), we have

$$f(y_N) - f(x_*) \leq \min \left\{ \frac{2L+m}{N}, (2L+m)q^N \right\} V[x_0](x_*) + \tilde{\delta} + 3\delta,$$

$$V[x_N](x_*) \leq q^N V[x_0](x_*) + (\tilde{\delta} + 2\delta) \sum_{i=1}^N \frac{Q_{i+1}^N}{L_i + m}.$$

The first term in the r.h.s. of the first equation above corresponds to exact gradient methods in the relatively smooth setting [7, 46]. According to [17], this part of our bound can be interpreted as optimal since [17] show that it can not be improved without additional assumptions on the Bregman divergence V .

Remark 2. The advantage of Algorithm 1 is that there is no need to know the true values of the parameters L and m . Using the standard argument [22] one can show that the number of oracle calls is less than $2N + \log_2 \frac{2L}{L_0}$, where N is the number of iterations of Algorithm 1.

Remark 3. In view of Theorem 3.1, if we would like to obtain the residual in the objective (12) to be smaller than ε , then it is sufficient to ensure $\tilde{\delta} = O(\varepsilon)$ and $\delta = O(\varepsilon)$.

3.2. Adaptive Fast Gradient Method with $(\delta, L, \mu, m, V, \|\cdot\|)$ -model

In this section, we slightly change the definition of inexact model and introduce $(\delta, L, \mu, m, V, \|\cdot\|)$ -model of the objective (see Definition 3.3). This change allows us to obtain an accelerated method for problems with this type of the model. The method is close to the accelerated mirror-descent type of methods (see [24, 42, 72]). In the new definition, we use the square of the norm in the r.h.s. of (6) instead of the function V , which gives the following modification of Definition 2.1.

Definition 3.3. Let $\delta, L, \mu, m \geq 0$. We say that $\psi_\delta(x, y)$ is a $(\delta, L, \mu, m, V, \|\cdot\|)$ -model of the function f at a given point y iff, for all $x \in Q$

$$\mu V[y](x) \leq f(x) - (f_\delta(y) + \psi_\delta(x, y)) \leq \frac{L}{2} \|x - y\|^2 + \delta \quad (19)$$

and $\psi_\delta(x, y)$ is convex in x , satisfies $\psi_\delta(x, x) = 0$ for all $x \in Q$ and

$$\psi(x) \geq \psi(z) + \langle g, x - z \rangle + mV[z](x), \quad \forall x, z \in Q \quad \forall g \in \partial\psi(z), \quad (20)$$

where for fixed $y \in Q$ and any $x \in Q$ we denote $\psi(x) = \psi_\delta(x, y)$. In other words, $\psi(x)$ is relatively m -strongly convex relative to d .

As in the previous subsection, we assume that there exists some constant \bar{L}_{k+1} such that $(\delta_k, \bar{L}_{k+1}, \mu, m, V, \|\cdot\|)$ -model of f exists at k -th step ($k = 0, \dots, N-1$) of Algorithm 2. Unlike Algorithm 1, we assume that the errors $\tilde{\delta}, \delta$ can depend on the iteration counter k , which is indicated by input sequences $\{\tilde{\delta}_k\}_{k \geq 0}$ and $\{\delta_k\}_{k \geq 0}$. For instance, this allows obtaining the Universal Fast Gradient Method in which different

values of $\{\delta_k\}_{k \geq 0}$ are required (see [5, 56]) in each iteration. Moreover, the sequence $\{\tilde{\delta}_k\}_{k \geq 0}$ is not used explicitly and only appears in convergence rates of Theorem 3.4.

Theorem 3.4. *Assume that $\psi_\delta(x, y)$ is a $(\delta, L, \mu, m, V, \|\cdot\|)$ -model according to Definition 3.3. Also assume that $V[y](x)$ satisfies (1-SC) condition w.r.t. $\|\cdot\|$ -norm. Then, after N iterations of Algorithm 2, we have*

$$f(x_N) - f(x_*) \leq \frac{V[u_0](x_*)}{A_N} + \frac{2 \sum_{k=0}^{N-1} A_{k+1} \delta_k}{A_N} + \frac{\sum_{k=0}^{N-1} \tilde{\delta}_k}{A_N}, \quad (21)$$

$$V[u_N](x_*) \leq \frac{V[u_0](x_*)}{(1 + A_N \mu + A_N m)} + \frac{2 \sum_{k=0}^{N-1} A_{k+1} \delta_k}{(1 + A_N \mu + A_N m)} + \frac{\sum_{k=0}^{N-1} \tilde{\delta}_k}{(1 + A_N \mu + A_N m)}. \quad (22)$$

Remark 4. Despite the adaptive structure of Algorithm 2, as in [56], it can be shown that on average the algorithm up to logarithmic terms requires four inexact computations of the objective function and two computations of $(\delta, L, \mu, m, V, \|\cdot\|)$ -model per iteration.

Algorithm 2 Fast adaptive gradient method with $(\delta, L, \mu, m, V, \|\cdot\|)$ -model

- 1: **Input:** x_0 is the starting point, $\mu \geq 0$, $m \geq 0$, $\{\delta_k\}_{k \geq 0}$ and $L_0 > 0$.
- 2: Set $y_0 := x_0$, $u_0 := x_0$, $\alpha_0 := 0$, $A_0 := \alpha_0$
- 3: **for** $k \geq 0$ **do**
- 4: Find the smallest integer $i_k \geq 0$ such that

$$f_{\delta_k}(x_{k+1}) \leq f_{\delta_k}(y_{k+1}) + \psi_{\delta_k}(x_{k+1}, y_{k+1}) + \frac{L_{k+1}}{2} \|x_{k+1} - y_{k+1}\|^2 + \delta_k, \quad (23)$$

where $L_{k+1} = 2^{i_k-1} L_k$, α_{k+1} is the largest root of the equation

$$A_{k+1}(1 + A_k \mu + A_k m) = L_{k+1} \alpha_{k+1}^2, \quad A_{k+1} := A_k + \alpha_{k+1}, \text{ and} \quad (24)$$

$$y_{k+1} := \frac{\alpha_{k+1} u_k + A_k x_k}{A_{k+1}}, \quad (25)$$

$$\phi_{k+1}(x) = \alpha_{k+1} \psi_{\delta_k}(x, y_{k+1}) + (1 + A_k \mu + A_k m) V[u_k](x) + \alpha_{k+1} \mu V[y_{k+1}](x),$$

$$u_{k+1} := \operatorname{argmin}_{x \in Q}^{\tilde{\delta}_k} \phi_{k+1}(x), \text{ for some } \tilde{\delta}_k \geq 0 \quad (26)$$

$$x_{k+1} := \frac{\alpha_{k+1} u_{k+1} + A_k x_k}{A_{k+1}}. \quad (27)$$

5: **end for**

In order to prove Theorem 3.4 we need the following lemma.

Lemma 3.5. *Let $\psi(x)$ be a relatively m -strongly convex function relative to d with $m \geq 0$, i.e. (7) holds, and*

$$y = \operatorname{argmin}_{x \in Q}^{\tilde{\delta}} \{\psi(x) + \beta V[z](x) + \gamma V[u](x)\},$$

where $\beta \geq 0$ and $\gamma \geq 0$. Then

$$\psi(x) + \beta V[z](x) + \gamma V[u](x) \geq \psi(y) + \beta V[z](y) + \gamma V[u](y) + (\beta + \gamma + m)V[y](x) - \tilde{\delta}, \quad \forall x \in Q.$$

We omit the proof of Lemma 3.5 since it is similar to the proof of Lemma 3.2.

Lemma 3.6. For all $x \in Q$, we have

$$\begin{aligned} & A_{k+1}f(x_{k+1}) - A_k f(x_k) + (1 + A_{k+1}\mu + A_{k+1}m)V[u_{k+1}](x) - (1 + A_k\mu + A_k m)V[u_k](x) \\ & \leq \alpha_{k+1}f(x) + 2\delta_k A_{k+1} + \tilde{\delta}_k. \end{aligned}$$

Proof. Since by Definition 3.3 with $x = y$, one has $f(x) - \delta \leq f_\delta(x) \leq f(x)$, and using (23), we have

$$f(x_{k+1}) \stackrel{(19)}{\leq} f_{\delta_k}(y_{k+1}) + \delta_k \stackrel{(23)}{\leq} f_{\delta_k}(y_{k+1}) + \psi_{\delta_k}(x_{k+1}, y_{k+1}) + \frac{L_{k+1}}{2} \|x_{k+1} - y_{k+1}\|^2 + 2\delta_k.$$

Substituting in this expression definitions (27) and (25) of the points x_{k+1} , y_{k+1} , we obtain that

$$\begin{aligned} f(x_{k+1}) & \leq f_{\delta_k}(y_{k+1}) + \psi_{\delta_k}\left(\frac{\alpha_{k+1}u_{k+1} + A_k x_k}{A_{k+1}}, y_{k+1}\right) \\ & \quad + \frac{L_{k+1}}{2} \left\| \frac{\alpha_{k+1}u_{k+1} + A_k x_k}{A_{k+1}} - \frac{\alpha_{k+1}u_k + A_k x_k}{A_{k+1}} \right\|^2 + 2\delta_k \\ & = f_{\delta_k}(y_{k+1}) + \psi_{\delta_k}\left(\frac{\alpha_{k+1}u_{k+1} + A_k x_k}{A_{k+1}}, y_{k+1}\right) + \frac{L_{k+1}\alpha_{k+1}^2}{2A_{k+1}^2} \|u_{k+1} - u_k\|^2 + 2\delta_k. \end{aligned}$$

Since $A_{k+1} = A_k + \alpha_{k+1}$ and, by Definition 3.3, $\psi_{\delta_k}(\cdot, y)$ is convex, we have

$$\begin{aligned} f(x_{k+1}) & \leq \frac{A_k}{A_{k+1}} (f_{\delta_k}(y_{k+1}) + \psi_{\delta_k}(x_k, y_{k+1})) + \frac{\alpha_{k+1}}{A_{k+1}} (f_{\delta_k}(y_{k+1}) + \psi_{\delta_k}(u_{k+1}, y_{k+1})) \\ & \quad + \frac{L_{k+1}\alpha_{k+1}^2}{2A_{k+1}^2} \|u_{k+1} - u_k\|^2 + 2\delta_k. \end{aligned}$$

In view of the definition (24) for the sequence α_{k+1} , we obtain

$$\begin{aligned} f(x_{k+1}) & \leq \frac{A_k}{A_{k+1}} (f_{\delta_k}(y_{k+1}) + \psi_{\delta_k}(x_k, y_{k+1})) + \frac{\alpha_{k+1}}{A_{k+1}} \left(f_{\delta_k}(y_{k+1}) + \psi_{\delta_k}(u_{k+1}, y_{k+1}) \right. \\ & \quad \left. + \frac{1 + A_k\mu + A_k m}{2\alpha_{k+1}} \|u_{k+1} - u_k\|^2 \right) + 2\delta_k. \end{aligned}$$

Using (1-SC) condition w.r.t. norm for V and the left inequality in (19), we get

$$\begin{aligned} f(x_{k+1}) & \leq \frac{A_k}{A_{k+1}} f_{\delta_k}(x_k) + \frac{\alpha_{k+1}}{A_{k+1}} \left(f_{\delta_k}(y_{k+1}) + \psi_{\delta_k}(u_{k+1}, y_{k+1}) \right. \\ & \quad \left. + \frac{1 + A_k\mu + A_k m}{\alpha_{k+1}} V[u_k](u_{k+1}) \right) + 2\delta_k. \end{aligned} \tag{28}$$

By Lemma 3.5 for the optimization problem in (26) with $\psi(x) = \alpha_{k+1}\psi_{\delta_k}(x, y_{k+1})$, $\beta = 1 + A_k\mu + A_k m$, $z = u_k$, $\gamma = \alpha_{k+1}\mu$, and $u = y_{k+1}$, it holds that

$$\begin{aligned} & \alpha_{k+1}\psi_{\delta_k}(u_{k+1}, y_{k+1}) + (1 + A_k\mu + A_k m)V[u_k](u_{k+1}) + \alpha_{k+1}\mu V[y_{k+1}](u_{k+1}) \\ & \quad + (1 + A_{k+1}\mu + A_{k+1}m)V[u_{k+1}](x) - \tilde{\delta}_k \\ & \leq \alpha_{k+1}\psi_{\delta_k}(x, y_{k+1}) + (1 + A_k\mu + A_k m)V[u_k](x) + \alpha_{k+1}\mu V[y_{k+1}](x). \end{aligned}$$

From the fact that $V[y_{k+1}](u_{k+1}) \geq 0$, we have

$$\begin{aligned} & \alpha_{k+1}\psi_{\delta_k}(u_{k+1}, y_{k+1}) + (1 + A_k\mu + A_k m)V[u_k](u_{k+1}) \\ & \leq \alpha_{k+1}\psi_{\delta_k}(x, y_{k+1}) + (1 + A_k\mu + A_k m)V[u_k](x) \\ & \quad - (1 + A_{k+1}\mu + A_{k+1}m)V[u_{k+1}](x) + \alpha_{k+1}\mu V[y_{k+1}](x) + \tilde{\delta}_k. \end{aligned} \tag{29}$$

Combining (28) and (29), we obtain

$$\begin{aligned} f(x_{k+1}) & \leq \frac{A_k}{A_{k+1}}f(x_k) + \frac{\alpha_{k+1}}{A_{k+1}} \left(f_{\delta_k}(y_{k+1}) + \psi_{\delta_k}(x, y_{k+1}) + \mu V[y_{k+1}](x) \right. \\ & \quad \left. + \frac{1 + A_k\mu + A_k m}{\alpha_{k+1}}V[u_k](x) - \frac{1 + A_{k+1}\mu + A_{k+1}m}{\alpha_{k+1}}V[u_{k+1}](x) + \frac{\tilde{\delta}_k}{\alpha_{k+1}} \right) + 2\delta_k. \end{aligned}$$

We finish the proof of Lemma 3.6 applying the left inequality in (19)

$$\begin{aligned} f(x_{k+1}) & \leq \frac{A_k}{A_{k+1}}f(x_k) + \frac{\alpha_{k+1}}{A_{k+1}}f(x) \\ & \quad + \frac{1 + A_k\mu + A_k m}{A_{k+1}}V[u_k](x) - \frac{1 + A_{k+1}\mu + A_{k+1}m}{A_{k+1}}V[u_{k+1}](x) + 2\delta_k + \frac{\tilde{\delta}_k}{A_{k+1}}. \end{aligned}$$

□

Proof of Theorem 3.4. We telescope the inequality in Lemma 3.6 for k from 0 to $N - 1$ and take $x = x_*$:

$$A_N f(x_N) \leq A_N f(x_*) + V[u_0](x_*) - (1 + A_N(\mu + m))V[u_N](x_*) + 2 \sum_{k=0}^{N-1} A_{k+1} \delta_k + \sum_{k=0}^{N-1} \tilde{\delta}_k. \tag{30}$$

Since $V[u_{k+1}](x_*) \geq 0$ for all $k \geq 0$, we have

$$A_N f(x_N) - A_N f(x_*) \leq V[u_0](x_*) + 2 \sum_{k=0}^{N-1} A_{k+1} \delta_k + \sum_{k=0}^{N-1} \tilde{\delta}_k.$$

The last inequality proves (21). Inequality (22) is a straightforward from (30) since $f(x) \geq f(x_*)$ for all $x \in Q$. □

The next lemma is proved in Appendix B and gives a lower bound for the growth rate of the sequence A_N , see [15, 33, 55].

Lemma 3.7. For all $N \geq 0$,

$$A_N \geq \max \left\{ \frac{1}{4} \left(\sum_{k=0}^{N-1} \frac{1}{\sqrt{L_{k+1}}} \right)^2, \frac{1}{L_1} \prod_{k=1}^{N-1} \left(1 + \sqrt{\frac{\mu+m}{4L_{k+1}}} \right)^2 \right\}.$$

Remark 5. Let us assume that the function f has L -Lipschitz continuous gradient. This means that for $L_k \geq L$, the inequality (23) always holds, whence, $L_k \leq 2L$, assuming that $L_0 \leq L$. Substituting this inequality in the bound of Lemma 3.7, we have $A_N \geq \frac{N^2}{4L}$ and

$$A_N \geq \frac{1}{2L} \left(1 + \frac{1}{4} \sqrt{\frac{\mu+m}{L}} \right)^{2(N-1)} \geq \frac{1}{2L} \exp \left(\frac{N-1}{4} \sqrt{\frac{\mu+m}{L}} \right).$$

In the last inequality we used inequality $\log(1+2x) \geq x$ for all $x \in [0, \frac{1}{4}]$. Combining Theorem 3.4 and Lemma 3.7 we have

$$\begin{aligned} f(x_N) - f(x_*) &\leq \min \left\{ \frac{8L}{N^2}, 2L \exp \left(-\frac{N-1}{4} \sqrt{\frac{\mu+m}{L}} \right) \right\} V[u_0](x_*) \\ &\quad + \frac{2 \sum_{k=0}^{N-1} A_{k+1} \delta_k}{A_N} + \frac{\sum_{k=0}^{N-1} \tilde{\delta}_k}{A_N}. \end{aligned} \quad (31)$$

The first term in the right hand side of inequality (31) is up to a constant factor optimal for μ -strongly convex functions with L -Lipschitz continuous gradient.

Note that in [15] for a non-adaptive fast gradient method with (δ, L, μ) -oracle, which is a particular case of our inexact model, for the case when δ_k is a constant it is shown that

$$\frac{\sum_{k=0}^{N-1} A_{k+1} \delta}{A_N} \leq \min \left\{ \left(\frac{1}{3} N + 2.4 \right), \left(1 + \sqrt{\frac{L}{\mu}} \right) \right\} \delta. \quad (32)$$

This means that for $\mu > 0$, the error δ does not accumulate.

Let us briefly discuss what happens if we can control the errors $\delta_k, \tilde{\delta}_k$ and choose them such that the residual in the objective is smaller than ε . For this, we need to ensure $\frac{\sum_{k=0}^{N-1} \tilde{\delta}_k}{A_N} = O(\varepsilon)$ for all $k \geq 0$. One way to guarantee it is to take $\tilde{\delta}_k = O(\varepsilon \alpha_{k+1})$ for all $k \geq 0$. From Lemma 3.7, $L_k \leq 2L$ for all $k \geq 0$, and (24), we have $\alpha_k \geq \frac{\sqrt{2k}}{4L}$. It means that the sequence $\{\tilde{\delta}_k\}_{k \geq 0}$ can be an increasing sequence.

Additionally, taking $\delta_k = O(\frac{\alpha_{k+1}}{A_{k+1}} \varepsilon)$, we can guarantee $\frac{\sum_{k=0}^{N-1} A_{k+1} \delta_k}{A_N} = O(\varepsilon)$. Therefore, the sequence $\{\delta_k\}_{k \geq 0}$, in general, should decrease with the rate of the order $1/k$ in order to obtain an ε -solution for problem (1).

Remark 6. In view of assumptions from Remark 5. For the case when $\mu = m = 0$ Algorithm 2 can guarantee the following convergence rate:

$$f(x_N) - f_* \leq \frac{8LV[x_0](x_*)}{N^2} + 2N\delta + \frac{4L\tilde{\delta}}{N}. \quad (33)$$

A similar result was shown in [32].

Remark 7. Let us analyze the convergence rate of $V[u_N](x_*)$ which is guaranteed by (22) in Theorem 3.4. Since we make a (1-SC) assumption on the function d , we have that $\frac{1}{2}\|u_N - x_*\|^2 \leq V[u_N](x_*)$ and the convergence rate for $V[u_N](x_*)$ gives also convergence rate for $\|u_N - x_*\|^2$. There are two different scenarios:

- (1) $\mu = m = 0$. In this case we have:

$$V[u_N](x_*) \leq V[u_0](x_*) + 2 \sum_{k=0}^{N-1} A_{k+1} \delta_k + \sum_{k=0}^{N-1} \tilde{\delta}_k.$$

We can only bound $V[u_N](x_*)$ by $V[u_0](x_*)$ up to additive noise.

- (2) $\mu + m > 0$. Using Lemma 3.7 we can see that Theorem 3.4 guarantees the linear convergence in argument up to additive noise.

Note that convergence rates for the objective and the argument are obtained for different sequences x_N and u_N , respectively.

3.3. Universal conditional gradient (Frank–Wolfe) method

Let us show an example of $(\delta, L, \mu, m, V, \|\cdot\|)$ -model's application. We assume that $\mu = 0, m = 0$ and use Algorithm 2 to propose a universal Frank–Wolfe method. In order to construct the universal Frank–Wolfe method let us introduce the following constraints to the optimization problem (1):

- (1) The set Q is bounded w.r.t $V[y](x)$: $\exists R_Q \in \mathbb{R} : V[y](x) \leq R_Q^2 \quad \forall x, y \in Q$.
- (2) The function $f(x)$ has Hölder-continuous subgradients:

$$\|\nabla f(x) - \nabla f(y)\|_* \leq L_\nu \|x - y\|^\nu \quad \forall x, y \in Q.$$

From this we can get, for any $\delta > 0$, an inequality (see [56])

$$0 \leq f(x) - f(y) - \langle \nabla f(y), x - y \rangle \leq \frac{L(\delta)}{2} \|x - y\|^2 + \delta \quad \forall x, y \in Q, \quad (34)$$

where

$$L(\delta) = L_\nu \left[\frac{L_\nu}{2\delta} \frac{1 - \nu}{1 + \nu} \right]^{\frac{1-\nu}{1+\nu}}.$$

Using the same arguments as in [32], we can show that the point $u_{k+1} = \arg \min_{x \in Q} \alpha_{k+1} \psi_{\delta_k}(x, y_{k+1})$ satisfies $u_{k+1} = \arg \min_{x \in Q} \tilde{\delta}_k \phi_{k+1}(x)$ with $\tilde{\delta}_k = 2R_Q^2$. It means that iteration u_{k+1} defined in this way satisfies (26). We also set $\delta_k = \varepsilon \frac{\alpha_{k+1}}{4A_{k+1}}$ as it was proposed in [56] for Universal Fast Gradient Method. This choice of δ_k and the fact that the objective function has Hölder continuous subgradient, using the same arguments as in Theorem 3 of [56], guarantees that in (21) $\frac{2 \sum_{k=0}^{N-1} A_{k+1} \delta_k}{A_N} \leq \frac{\varepsilon}{2}$. Combining it again with (21), $V[y](x) \leq R_Q^2$ and $\tilde{\delta}_k = 2R_Q^2$, we obtain from Theorem 3.4

that

$$f(x_N) - f(x_*) \leq \frac{R_Q^2}{A_N} + \frac{\varepsilon}{2} + \frac{2R_Q^2 N}{A_N} \leq \frac{3R_Q^2 N}{A_N} + \frac{\varepsilon}{2}.$$

Using the same arguments as in Theorem 3 of [56], we obtain that

$$A_N \geq \frac{N^{\frac{1+3\nu}{1+\nu}} \epsilon^{\frac{1-\nu}{1+\nu}}}{2^{\frac{3+5\nu}{1+\nu}} L_\nu^{\frac{2}{1+\nu}}}. \quad (35)$$

Using this inequality, we obtain the following upper bound for the number of steps in order to guarantee $f(x_N) - f(x_*) \leq \varepsilon$

$$N \leq \inf_{\nu \in (0,1]} \left[2^{\frac{3+4\nu}{\nu}} \left(\frac{L_\nu R_Q^{1+\nu}}{\varepsilon} \right)^{\frac{1}{\nu}} \right],$$

where the infimum can be taken since neither ν nor L_ν is not used in the algorithm. This bound justifies the word "universal" in the name of the algorithm since the algorithm is the same for any values of $\nu \in [0, 1]$ and works with the best possible convergence guarantee. We call the proposed algorithm conditional gradient (Frank–Wolfe) method since in a simple situation $\psi_{\delta_k}(x, y_{k+1}) = \langle \nabla f(y_{k+1}), x - y_{k+1} \rangle$, finding $u_{k+1} = \arg \min_{x \in Q} \alpha_{k+1} \psi_{\delta_k}(x, y_{k+1})$ is reduced to solving a linear minimization problem over the set Q . This method for composite optimization with Hölder-continuous gradient was analyzed in [57] and for a non-composite case was very recently analyzed in [76]. Our method is more general, since it is universal and does not require the knowledge of the parameters ν, L_ν , and also can utilize general inexact models described by Definition 3.3, including inexact gradients, composite optimization, inexact linear minimization oracle.

4. Inexact Model for Variational Inequalities

In this section, we go beyond minimization problems and propose an abstract inexact model counterpart for variational inequalities. As a special case in Example 4.3, we introduce relative smoothness for operators in the spirit of [46], where it was introduced for optimization problems. Further, we propose a generalization of the Mirror-Prox algorithm for this general case of the inexact model of the operator and abstract variational inequalities. One of the main features of our algorithm is its adaptation to generalized inexact parameter of smoothness. As a special case, we propose a universal method for variational inequalities with complexity $O\left(\left(\frac{1}{\varepsilon}\right)^{\frac{2}{1+\nu}}\right)$, where ε is the desired accuracy of the solution and ν is the Hölder exponent of the operator. According to the lower bounds in [66], this algorithm is optimal for $\nu = 0$ (bounded variation of the operator) and $\nu = 1$ (Lipschitz continuity of the operator). Based on the model for VI and functions, we introduce an inexact model for saddle-point problems (see Definition 4.9). We are also motivated by mixed variational inequalities [6, 39] and composite saddle-point problems [10].

Formally speaking, we consider the problem of finding the solution $x_* \in Q$ for VI

in the following abstract form

$$\psi(x, x_*) \geq 0 \quad \forall x \in Q \quad (36)$$

for some convex compact set $Q \subset \mathbb{R}^n$ and some function $\psi : Q \times Q \rightarrow \mathbb{R}$. Assuming the abstract monotonicity of the function ψ

$$\psi(x, y) + \psi(y, x) \leq 0 \quad \forall x, y \in Q, \quad (37)$$

any solution to (36) is a solution of the following inequality

$$\max_{x \in Q} \psi(x_*, x) \leq 0 \quad (38)$$

In the general case, we make an assumption about the existence of a solution x_* of the problem (36). As a particular case, if for some operator $g : Q \rightarrow \mathbb{R}^n$ we set $\psi(x, y) = \langle g(y), x - y \rangle \quad \forall x, y \in Q$, then (36) and (38) are equivalent, respectively, to a standard strong and weak variational inequality with the operator g .

We start with a concept of (δ, L, V) -model for problems (36) and (38).

Definition 4.1. We say that a function ψ has (δ, L, V) -model $\psi_\delta(x, y)$ for some fixed values $\delta > 0$ and $L = L(\delta) > 0$ if the following properties hold for each $x, y, z \in Q$:

- (i) $\psi(x, y) \leq \psi_\delta(x, y) + \delta$;
- (ii) $\psi_\delta(x, y)$ convex in the first variable;
- (iii) $\psi_\delta(x, x) = 0$;
- (iv) (*abstract δ -monotonicity*)

$$\psi_\delta(x, y) + \psi_\delta(y, x) \leq \delta; \quad (39)$$

- (v) (*generalized relative smoothness*)

$$\psi_\delta(x, y) \leq \psi_\delta(x, z) + \psi_\delta(z, y) + LV[z](x) + LV[y](z) + \delta. \quad (40)$$

Example 4.2. For some operator $g : Q \rightarrow \mathbb{R}^n$ and a convex function $h : Q \rightarrow \mathbb{R}^n$, the choice

$$\psi(x, y) = \langle g(y), x - y \rangle + h(x) - h(y) \quad (41)$$

leads to a *mixed variational inequality* [6, 39]

$$\langle g(y), y - x \rangle + h(y) - h(x) \leq 0, \quad (42)$$

which in the case of the monotonicity of the operator g implies

$$\langle g(x), y - x \rangle + h(y) - h(x) \leq 0. \quad (43)$$

Remark 8. Similarly to Definition 2.1 above, in general case, we do not need the (1-SC) assumption for $V[y](x)$ in Definition 4.1. In some situations we make (1-SC) assumption for $V[y](x)$ (see Example 4.7 and Section 5).

Note that for $\delta = 0$ the following analogue of (40) for some fixed $a, b > 0$

$$\psi(x, y) \leq \psi(x, z) + \psi(z, y) + a\|z - y\|^2 + b\|x - z\|^2 \quad \forall x, y, z \in Q \quad (44)$$

was introduced in [50]. Condition (44) is used in many works on equilibrium programming. Our approach allows us to work with non-Euclidean set-up without (1-SC) assumption and inexactness δ , which is important for the framework of universal methods [56] (see Example 4.7 below).

One can directly verify that if $\psi_\delta(x, y)$ is $(\delta/3, L, 0, 0, V)$ -model of the function f at a given point y then $\psi_\delta(x, y)$ is (δ, L, V) -model in the sense of Definition 4.1.

Let us consider some examples.

Example 4.3. Relative smoothness for optimization and VI. Let us consider a minimization problem (1) with the function f being smooth, convex and relatively L -smooth relative to d [46], i.e., for all $x, y \in Q$,

$$f(x) - f(y) - \langle \nabla f(y), x - y \rangle \leq LV[y](x).$$

In this case, (1) is equivalent to abstract VI (38) with $\psi_\delta(x, y) := \langle \nabla f(y), x - y \rangle$. Properties (i)-(iv) in Definition 4.1 obviously hold with $\delta = 0$. Let us check that (v) also holds. Indeed,

$$\begin{aligned} \psi_\delta(x, y) - \psi_\delta(x, z) - \psi_\delta(z, y) &= \langle \nabla f(y), x - y \rangle - \langle \nabla f(z), x - z \rangle - \langle \nabla f(y), z - y \rangle \\ &= (f(x) - f(z) - \langle \nabla f(z), x - z \rangle) + (f(z) - f(y) - \langle \nabla f(y), z - y \rangle) \\ &\quad - (f(x) - f(y) - \langle \nabla f(y), x - y \rangle) \leq LV[z](x) + LV[y](z), \end{aligned}$$

where we used relative L -smoothness and convexity of f . This example shows that our inexact model for VI as a particular case contains the concept of relative smoothness introduced in optimization. In this particular case, we say that an operator g is relatively L -smooth if

$$\langle g(y) - g(z), x - z \rangle \leq LV[z](x) + LV[y](z) \quad \forall x, y, z \in Q.$$

Let us make a precise definition

Definition 4.4. We say that an operator $g(x)$ is relatively smooth relative to function $d(x)$ if for the corresponding Bregman divergence $V[y](x)$ and a constant $L \geq 0$ it holds

$$\langle g(y) - g(z), x - z \rangle \leq LV[z](x) + LV[y](z) \quad \forall x, y, z \in Q. \quad (45)$$

This definition (and its inexact version when there is some error $\delta > 0$ added to the r.h.s.) contains as particular case several definitions which were introduced in the literature after the initial draft of this paper appeared as a preprint [70]. The definition of relative Lipschitzness, introduced in [13] and used to provide a simple explanation for accelerated methods, coincides with the above definition of relatively smooth operator. Two other examples are explained next.

Example 4.5 (Bregman continuity and Metric regularity). Let us analyze the connection of our notion of relative smoothness and the notion of Bregman continuity

introduced in [4] and Metric regularity introduced in [3]. Both papers consider local norm $\|\cdot\|_x$ which is continuous for $x \in Q$, its dual $\|\cdot\|_{x,*}$ defined in a standard way, Bregman function d and its Bregman divergence $V[z](x) = d(x) - d(z) - \langle \nabla d(z), x - z \rangle$ satisfying for some $K > 0$

$$V[z](x) \geq \frac{K}{2} \|x - z\|_z^2. \quad (46)$$

The paper [4] introduces a notion of β -Bregman continuous operators which satisfy

$$\|g(y) - g(z)\|_{z,*} \leq \beta \sqrt{2V[y](z)}. \quad (47)$$

Let us show that such operators are relatively smooth relative to d . Indeed, for any $x, y, z \in Q$,

$$\begin{aligned} \langle g(y) - g(z), x - z \rangle &\leq \|g(y) - g(z)\|_{z,*} \|x - z\|_z \stackrel{(47)}{\leq} \beta \sqrt{2V[y](z)} \|x - z\|_z \\ &\stackrel{(46)}{\leq} \beta \sqrt{2V[y](z)} \cdot \sqrt{\frac{2}{K} V[z](x)} \leq \frac{\beta}{\sqrt{K}} (V[z](x) + V[y](z)), \end{aligned}$$

where in the last inequality we used that $ab \leq \frac{1}{2}(a^2 + b^2)$ for any $a, b \geq 0$. Thus, g satisfies (45) with $L = \beta/\sqrt{K}$ and is relatively smooth relative to d .

The paper [3] uses additional assumption that the local metric is regular, i.e., for $\beta > 0$, satisfies

$$\|v\|_{z,*} \leq \|v\|_{y,*} (1 + \beta \|z - y\|_y). \quad (48)$$

The following two notions are also introduced by saying that the operator g is

- Metrically bounded if there exists some $M_0 > 0$ such that

$$\|g(z)\|_{z,*} \leq M_0, \quad \forall z \in Q.$$

- Metrically smooth if there exists some $M_1 > 0$ such that

$$\|g(y) - g(z)\|_{z,*} \leq M_1 \|y - z\|_y, \quad \forall y, z \in Q.$$

Let us show that in these two cases g is inexactly relatively smooth relative to d . Let g be metrically bounded. Then

$$\begin{aligned} \|g(y) - g(z)\|_{z,*} &\leq \|g(y)\|_{z,*} + \|g(z)\|_{z,*} \\ &\stackrel{(48)}{\leq} \|g(y)\|_{y,*} (1 + \beta \|z - y\|_y) + \|g(z)\|_{z,*} \leq 2M_0 + \beta M_0 \|z - y\|_y. \end{aligned} \quad (49)$$

Further, for any $\delta > 0$,

$$\begin{aligned}
\langle g(y) - g(z), x - z \rangle &\leq \|g(y) - g(z)\|_{z,*} \|x - z\|_z \stackrel{(49)}{\leq} (2M_0 + \beta M_0 \|z - y\|_y) \|x - z\|_z \\
&= 2M_0 \|z - y\|_y^0 \|x - z\|_z + \beta M_0 \|z - y\|_y \|x - z\|_z \\
&\stackrel{*}{\leq} \frac{1}{\delta} \frac{M_0^2}{2} (\|z - y\|_y^2 + \|x - z\|_z^2) + \frac{\delta}{2} + \frac{\beta M_0}{2} (\|z - y\|_y^2 + \|x - z\|_z^2) \\
&\stackrel{(46)}{\leq} \left(\frac{M_0^2}{2\delta} + \frac{\beta M_0}{2} \right) \frac{2}{K} (V[z](x) + V[y](z)) + \frac{\delta}{2},
\end{aligned}$$

where $*$ uses that, for any $a, b, c \geq 0$, any $\delta > 0$, any $\nu \in [0, 1]$ (and in particular for $\nu = 0$), it holds that $ab^\nu c \leq \left(\frac{1}{\delta}\right)^{\frac{1-\nu}{1+\nu}} \frac{a^{\frac{2}{1+\nu}}}{2} (b^2 + c^2) + \frac{\delta}{2}$, see [69][Lemma 1]. Thus, we obtain that, for any $\delta > 0$,

$$\langle g(y) - g(z), x - z \rangle \leq L(\delta)(V[z](x) + V[y](z)) + \frac{\delta}{2}$$

with

$$L(\delta) = \left(\frac{M_0}{\delta} + \beta \right) \frac{M_0}{K}, \quad (50)$$

i.e. (45) holds with any error $\delta > 0$ and g is inexactly relatively smooth relative to d . Note that this is also covered by our concept of inexact model.

Let g be metrically smooth. Then, from (46) it is straightforward that

$$\|g(y) - g(z)\|_{z,*} \leq M_1 \|y - z\|_y \leq M_1 \sqrt{\frac{2}{K}} V[y](z)$$

and, thus, g is M_1/\sqrt{K} -Bregman continuous, whence, as it was shown above, g is also relatively smooth relative to d with $L = M_1/K$.

Thus, we conclude that our algorithms for variational inequalities are also applicable for solving variational inequalities with Bregman continuous, metrically bounded and metrically smooth operators. After presenting our method and its convergence rate we discuss their convergence rate for these types of operators. Moreover, the methods which we develop in this paper for variational inequalities and saddle-point problems are applicable to such problems in the presence of inexact information, which is a more general setting than in [3, 4].

Example 4.6 (Resource sharing problem). An interesting example of a problem in which a variational inequality with a relatively smooth and monotone operator naturally arises is a resource sharing problem considered in [4] (Example 2.3). In this example, $g(x) = (l_1(x_1), \dots, l_n(x_n))$ with a loss function $l_i(x) = \frac{1}{\alpha_i - x_i}$, $i = 1, \dots, n$ where x belongs to the set of feasible resource allocations $Q = \{x = (x_1, \dots, x_n) : 0 \leq x_i < \alpha_i, x_1 + \dots + x_n = R\}$ for $R > 0$, and α_i denotes the capacity of the resource i . This operator has singularities corresponding to α_i . Let $d(x) = \sum_{i=1}^n \frac{1}{1-x_i}$ be the prox-function defined on the feasible set Q . Then the associated Bregman divergence

is

$$V[y](x) = \sum_{i=1}^n \frac{(x_i - y_i)^2}{(1 - x_i)(1 - y_i)^2}.$$

In [4] the authors show that the operator g is Bregman continuous. Since in the previous example we showed that such operators are relatively smooth, we conclude that our algorithms are also suitable for the resource sharing problem. Experiments on this problem can be found in Appendix G.

Example 4.7. Variational Inequalities with monotone Hölder continuous operator. Assume that V satisfies (1-SC) condition w.r.t. some norm $\|\cdot\|$ and for a monotone operator g there exists $\nu \in [0, 1]$ such that

$$\|g(x) - g(y)\|_* \leq L_\nu \|x - y\|^\nu \quad \forall x, y \in Q. \quad (51)$$

Then, we have $\langle g(z) - g(y), z - x \rangle \leq \|g(z) - g(y)\|_* \|z - x\| \leq L_\nu \|z - y\|^\nu \|z - x\|$

$$\leq \frac{L(\delta)}{2} \|z - x\|^2 + \frac{L(\delta)}{2} \|z - y\|^2 + \delta \leq L(\delta)V[z](x) + L(\delta)V[y](z) + \delta \quad (52)$$

with

$$L(\delta) = \left(\frac{1}{2\delta} \right)^{\frac{1-\nu}{1+\nu}} L_\nu^{\frac{2}{1+\nu}} \quad (53)$$

with arbitrary $\delta > 0$. In this case $\psi_\delta(x, y) := \langle g(y), x - y \rangle$ is a (δ, L, V) -model.

Note that for the previous two examples in Algorithm 3 and Theorem 4.8 we need $V[z](x)$ to satisfy (1-SC) condition.

Next, we introduce our adaptive method (Algorithm 3) for abstract variational inequalities with inexact (δ, L, V) -model. This method adapts to the local values of L and allows us to construct a universal method for variational inequalities by applying it to VI with Hölder condition (51) for $\delta = \frac{\varepsilon}{2}$ and $L = L(\frac{\varepsilon}{2})$.

Next, we state the convergence rate result for the proposed method.

Theorem 4.8. *For Algorithm 3 the following inequality holds*

$$-\frac{1}{S_N} \sum_{k=0}^{N-1} \frac{\psi_\delta(x, w_k)}{L_{k+1}} \leq \frac{V[z_0](x)}{S_N} + \delta + 2\tilde{\delta} \quad \forall x \in Q. \quad (58)$$

Moreover,

$$\max_{u \in Q} \psi(\hat{w}_N, u) \leq \frac{2L \max_{u \in Q} V[z_0](u)}{N} + 3\delta + 2\tilde{\delta},$$

and Algorithm 3 stops in no more than

$$\left\lceil \frac{2L \max_{u \in Q} V[z_0](u)}{\varepsilon} \right\rceil \quad (59)$$

Algorithm 3 Generalized Mirror Prox for VI

Require: accuracy $\varepsilon > 0$, oracle error $\delta > 0$, initial guess $L_0 > 0$, prox set-up: $d(x), V[z](x)$.

1: Set $k = 0, z_0 = \arg \min_{u \in Q} d(u)$.

2: **repeat**

3: Find the smallest integer $i_k \geq 0$ such that

$$\psi_\delta(z_{k+1}, z_k) \leq \psi_\delta(z_{k+1}, w_k) + \psi_\delta(w_k, z_k) + L_{k+1}(V[z_k](w_k) + V[w_k](z_{k+1})) + \delta, \quad (54)$$

where $L_{k+1} = 2^{i_k-1} L_k$ and

$$w_k = \underset{x \in Q}{\operatorname{argmin}}^{\tilde{\delta}} \{ \psi_\delta(x, z_k) + L_{k+1} V[z_k](x) \}. \quad (55)$$

$$z_{k+1} = \underset{x \in Q}{\operatorname{argmin}}^{\tilde{\delta}} \{ \psi_\delta(x, w_k) + L_{k+1} V[z_k](x) \}. \quad (56)$$

4: **until**

$$S_N := \sum_{k=0}^{N-1} \frac{1}{L_{k+1}} \geq \frac{\max_{x \in Q} V[x^0](x)}{\varepsilon}. \quad (57)$$

Ensure: $\hat{w}_N = \frac{1}{\sum_{k=0}^{N-1} \frac{1}{L_{k+1}}} \sum_{k=0}^{N-1} \frac{1}{L_{k+1}} w_k$.

iterations.

Before we prove the theorem, we note that similarly to [3], we can consider any nonempty bounded convex subset C of the set Q and have similar guarantees for $\max_{u \in C} \psi(\hat{w}_N, u)$ that plays a role of generalized restricted gap (or merit) function. Thus, there is no need for the set Q to be bounded.

Proof. After $(k+1)$ -th iteration ($k = 0, 1, 2, \dots$) from (55) and (56) we have, for each $u \in Q$,

$$\psi_\delta(w_k, z_k) \leq \psi_\delta(u, z_k) + L_{k+1} V[z_k](u) - L_{k+1} V[w_k](u) - L_{k+1} V[z_k](w_k) + \tilde{\delta}$$

and

$$\psi_\delta(z_{k+1}, w_k) \leq \psi_\delta(u, w_k) + L_{k+1} V[z_k](u) - L_{k+1} V[z_{k+1}](u) - L_{k+1} V[z_k](z_{k+1}) + \tilde{\delta}.$$

The first inequality means that

$$\psi_\delta(w_k, z_k) \leq \psi_\delta(z_{k+1}, z_k) + L_{k+1} V[z_k](z_{k+1}) - L_{k+1} V[w_k](z_{k+1}) - L_{k+1} V[z_k](w_k) + \tilde{\delta}.$$

Taking into account (54), we obtain, for all $u \in Q$,

$$-\psi_\delta(u, w_k) \leq L_{k+1} V[z_k](u) - L_{k+1} V[z_{k+1}](u) + \delta + 2\tilde{\delta}.$$

So, the following inequality holds:

$$-\sum_{k=0}^{N-1} \frac{\psi_\delta(u, w_k)}{L_{k+1}} \leq V[z_0](u) - V[z_N](u) + S_N(\delta + 2\tilde{\delta}).$$

By virtue of (40) and the choice of $L_0 \leq 2L$, it is guaranteed that $L_{k+1} \leq 2L \ \forall k = 0, N-1$ and we have from Definition 4.1

$$\begin{aligned} \max_{u \in Q} \psi(\hat{w}_N, u) &\leq \max_{u \in Q} \psi_\delta(\hat{w}_N, u) + \delta \\ &\leq -\frac{1}{S_N} \sum_{k=0}^{N-1} \frac{\psi_\delta(u, w_k)}{L_{k+1}} + 2\delta \leq \frac{2L \max_{u \in Q} V[z_0](u)}{N} + 3\delta + 2\tilde{\delta}. \end{aligned} \tag{60}$$

□

Remark 9 (Bregman continuity and Metric regularity). Let us consider the setting of Example 4.5. In all the cases we define $\psi_\delta(w, u) = \langle g(u), w - u \rangle$. First, if the operator g is β -Bregman continuous, then, as we show in Example 4.5, it is relatively smooth relative to d with constant $L = \beta/\sqrt{K}$. Applying Theorem 4.8 with $\delta = \tilde{\delta} = 0$, we obtain that

$$\max_{u \in Q} \langle g(u), \hat{w}_N - u \rangle \leq \frac{2\beta \max_{u \in Q} V[z_0](u)}{N\sqrt{K}},$$

which has the same dependence on N as the rate in [4].

Let us now consider M_0 -metrically bounded operator g . As it was shown in Example 4.5, this operator is inexactly relatively smooth, which means that $\psi_\delta(w, u) = \langle g(u), w - u \rangle$ is an inexact model with $L(\delta) = (\frac{M_0}{\delta} + \beta) \frac{M_0}{K}$ defined in (50). In this setting, we have $\tilde{\delta} = 0$ and set $\delta = \varepsilon/6$. Applying Theorem 4.8, we obtain

$$\max_{u \in Q} \langle g(u), \hat{w}_N - u \rangle \leq \frac{2 \max_{u \in Q} V[z_0](u)}{N} \left(\frac{6M_0}{\varepsilon} + \beta \right) \frac{M_0}{K} + \frac{\varepsilon}{2}.$$

Thus, to obtain an ε -solution, i.e. guarantee $\max_{u \in Q} \langle g(u), \hat{w}_N - u \rangle \leq \varepsilon$, it is sufficient to take $N \geq \Omega \left(\frac{M_0^2 \max_{u \in Q} V[z_0](u)}{K\varepsilon^2} \right)$, which has the same dependence on ε as the bound in [3].

Finally, we consider M_1 -metrically smooth operator g . According to Example 4.5, this operator is also relatively smooth with $L = M_1/K$. Applying Theorem 4.8, we obtain

$$\max_{u \in Q} \langle g(u), \hat{w}_N - u \rangle \leq \frac{2M_1 \max_{u \in Q} V[z_0](u)}{NK}.$$

A very close convergence rate is obtained in [3], yet without explicit dependence on M_1 and K .

Note that in all three examples the same rate holds for $\max_{u \in C} \langle g(u), \hat{w}_N - u \rangle$ where C is any non-empty bounded convex subset of Q . Thus, there is no need for the set Q to be bounded.

Remark 10. In the setting of Example 4.7, we set, for any desired accuracy $\varepsilon > 0$, $\delta = \frac{\varepsilon}{2}$ and $L = L\left(\frac{\varepsilon}{2}\right)$ according to (53). Then from (59), we obtain that Algorithm 3 has the following complexity to guarantee

$$\left\lceil 2 \inf_{\nu \in [0,1]} \left(\frac{2L_\nu}{\varepsilon} \right)^{\frac{2}{1+\nu}} \cdot \max_{u \in Q} V[z_0](u) \right\rceil. \quad (61)$$

Note that estimate (61) is optimal for variational inequalities and saddle-point problems in the cases $\nu = 0$ and $\nu = 1$.

Thus, the introduced concept of the (δ, L, V) -model for variational inequalities allows us to extend the previously proposed universal method for VI to a wider class of problems, including *mixed variational inequalities* [6, 39] and *composite saddle-point problems* [10].

Remark 11. The authors of [4] propose an adaptive Mirror-Prox algorithm for variational inequalities with Bregman continuous operator, i.e. satisfying inequality similar to (47). As we show above in Example 4.6, this setting is covered by our concept of relative smoothness for operators and, thus, also by our notion of inexact model for variational inequalities. At the same time their algorithm has an important difference with our algorithm in terms of the choice of the stepsize. In our Algorithm 3, the role of the stepsize is played by $1/L_{k+1}$, and it may happen, since $L_{k+1} = 2^{i_k-1}L_k$ with $i_k \geq 0$, that the stepsize is increasing between iterations leading to faster convergence in practice. On the contrary, the stepsize in [4] is non-increasing. We illustrate the importance of the possibility to increase the stepsize in computational experiments in Appendix D for non-smooth saddle-point problems by comparing our method with $i_k \geq 0$ and with $i_k \geq 1$. The same remark holds also for a very recent preprint [3], where an adaptive Mirror-Prox algorithm for metrically regular problems was proposed with optimal bounds for metrically bounded and metrically smooth operators. This method also uses decreasing stepsizes. Finally, both papers [3, 4] do not consider any inexactness for their adaptive methods. Only for the non-adaptive method in [4] they prove convergence rate in the setting of stochastic errors in the operator values, the setting we do not consider here.

Now we introduce an inexact model for saddle-point problems. The solution of variational inequalities reduces the so-called saddle-point problems, in which for a convex in u and concave in v functional $f(u, v) : \mathbb{R}^{n_1+n_2} \rightarrow \mathbb{R}$ ($u \in Q_1 \subset \mathbb{R}^{n_1}$ and $v \in Q_2 \subset \mathbb{R}^{n_2}$) needs to be found the point (u_*, v_*) such that:

$$f(u_*, v) \leq f(u_*, v_*) \leq f(u, v_*) \quad (62)$$

for arbitrary $u \in Q_1$ and $v \in Q_2$. Let $Q = Q_1 \times Q_2 \subset \mathbb{R}^{n_1+n_2}$. For $x = (u, v) \in Q$, we assume that $\|x\| = \sqrt{\|u\|_1^2 + \|v\|_2^2}$ ($\|\cdot\|_1$ and $\|\cdot\|_2$ are the norms in the spaces \mathbb{R}^{n_1} and \mathbb{R}^{n_2} , respectively). We agree to denote $x = (u_x, v_x)$, $y = (u_y, v_y) \in Q$.

It is well known that for a sufficiently smooth function f with respect to u and v the problem (62) reduces to VI with an operator $g(x) = (f'_u(u_x, v_x), -f'_v(u_x, v_x))$.

For saddle-point problems we propose some adaptation of the concept of the (δ, L, V) -model for abstract variational inequality.

Definition 4.9. We say that the function $\psi_\delta(x, y)$ ($\psi_\delta : \mathbb{R}^{n_1+n_2} \times \mathbb{R}^{n_1+n_2} \rightarrow \mathbb{R}$)

is a (δ, L, V) -model for the saddle-point problem (62) if the conditions (ii) – (v) of Definition 4.1 hold and in addition

$$f(u_y, v_x) - f(u_x, v_y) \leq -\psi_\delta(x, y) + \delta \quad \forall x, y \in Q. \quad (63)$$

Example 4.10. The proposed concept of the (δ, L, V) -model for saddle-point problems is quite applicable, for example, for composite saddle-point problems of the form considered in the popular paper [10]:

$$f(u, v) = \tilde{f}(u, v) + h(u) - \varphi(v) \quad (64)$$

for some convex in u and concave in v subdifferentiable functions \tilde{f} , as well as convex functions h and φ . In this case, we can put

$$\psi_\delta(x, y) = \langle \tilde{g}(y), x - y \rangle + h(u_x) + \varphi(v_x) - h(u_y) - \varphi(v_y), \quad (65)$$

where

$$\tilde{g}(y) = \begin{pmatrix} \tilde{g}_u(u_y, v_y) \\ -\tilde{g}_v(u_y, v_y) \end{pmatrix}.$$

for some $\tilde{g}_u \in \partial_u f(u_y, v_y)$ and $\tilde{g}_v \in -\partial_v f(u_y, v_y)$.

Theorem 4.8 implies

Theorem 4.11. *If for the saddle-point problem (62) there is a (δ, L, V) -model $\psi_\delta(x, y)$, then after stopping the algorithm we get a point*

$$\hat{y}_N = (u_{\hat{y}_N}, v_{\hat{y}_N}) := (\hat{u}_N, \hat{v}_N) := \frac{1}{S_N} \sum_{k=0}^{N-1} \frac{y_k}{L_{k+1}}, \quad (66)$$

for which the following inequality is true:

$$\max_{v \in Q_2} f(\hat{u}_N, v) - \min_{u \in Q_1} f(u, \hat{v}_N) \leq \frac{2L \max_{(u,v) \in Q} V[u_0, v_0](u, v)}{N} + 2\tilde{\delta} + 2\delta. \quad (67)$$

5. Inexact Model for Strongly Monotone VI

In this section similarly with the concept of (δ, L, μ, m, V) -model in optimization we consider an inexact model for VI with a stronger version of the monotonicity condition (39).

Definition 5.1. We say that functional ψ has (δ, L, μ, V) -model $\psi_\delta(x, y)$ at a given point y if the following properties hold for each $x, y, z \in Q$:

- (i) $\psi(x, y) \leq \psi_\delta(x, y) + \delta$;
- (ii) $\psi_\delta(x, y)$ convex in the first variable;
- (iii) $\psi_\delta(x, y)$ continuous in x and y ;
- (iii) $\psi_\delta(x, x) = 0$;

(iv) (μ -strong δ -monotonicity)

$$\psi_\delta(x, y) + \psi_\delta(y, x) + \mu\|x - y\|^2 \leq \delta; \quad (68)$$

(v) (*generalized relative smoothness*)

$$\psi_\delta(x, y) \leq \psi_\delta(x, z) + \psi_\delta(z, y) + LV[z](x) + LV[y](z) + \delta \quad (69)$$

for some fixed values $L > 0$, $\delta > 0$.

Remark 12. We note that we can not replace $\|x - y\|^2$ by $V[y](x)$ in (68) since it is essentially used in the proof of Theorem 5.2.

Now we propose a method with a linear rate of convergence for VI with (δ, L, μ, V) -model. We slightly modify the assumptions on prox-function $d(x)$. Namely, we assume that $\operatorname{argmin}_{x \in Q} d(x) = 0$ and that d is bounded on the unit ball in the chosen norm $\|\cdot\|$, that is

$$d(x) \leq \frac{\Omega}{2}, \quad \forall x \in Q : \|x\| \leq 1, \quad (70)$$

where Ω is some known constant. Note that for standard proximal setups $\Omega = O(\ln \dim E)$. Finally, we assume that we are given a starting point $x_0 \in Q$ and a number $R_0 > 0$ such that $\|x_0 - x_*\|^2 \leq R_0^2$, where x_* is the solution to abstract VI. The procedure of restating of Algorithm 3 is applicable for abstract strongly monotone variational inequalities.

Algorithm 4 Restarted Generalized Mirror Prox

Require: accuracy $\varepsilon > 0$, $\mu > 0$, Ω s.t. $d(x) \leq \frac{\Omega}{2} \forall x \in Q : \|x\| \leq 1$; x_0, R_0 s.t. $\|x_0 - x_*\|^2 \leq R_0^2$.

- 1: Set $p = 0$, $d_0(x) = R_0^2 d\left(\frac{x - x_0}{R_0}\right)$.
- 2: **repeat**
- 3: Set x_{p+1} as the output of Algorithm 3 after N_p iterations of Algorithm 3 with prox-function $d_p(\cdot)$ and stopping criterion $\sum_{k=0}^{N_p-1} \frac{1}{L_{k+1}} \geq \frac{\Omega}{\mu}$.
- 4: Set $R_{p+1}^2 = R_0^2 \cdot 2^{-(p+1)} + 2(1 - 2^{-(p+1)}) \frac{\delta + 2\tilde{\delta}}{\mu}$.
- 5: Set $d_{p+1}(x) \leftarrow R_{p+1}^2 d\left(\frac{x - x_{p+1}}{R_{p+1}}\right)$.
- 6: Set $p = p + 1$.
- 7: **until** $p > \log_2 \frac{R_0^2}{\varepsilon}$

Ensure: x_{p+1} .

Theorem 5.2. Assume that ψ_δ is a (δ, L, μ, V) -model for ψ . Also assume that the prox function $d(x)$ satisfies (70) and the starting point $x_0 \in Q$ and a number $R_0 > 0$ are such that $\|x_0 - x_*\|^2 \leq R_0^2$, where x_* is the solution to (38). Then, for each $p \geq 0$

$$\|x_p - x_*\|^2 \leq R_0^2 \cdot 2^{-p} + \frac{2\delta}{\mu} + \frac{4\tilde{\delta}}{\mu} \leq \varepsilon + \frac{2\delta}{\mu} + \frac{4\tilde{\delta}}{\mu}.$$

The total number of iterations of the inner Algorithm 3 does not exceed

$$\left\lceil \frac{2L\Omega}{\mu} \cdot \log_2 \frac{R_0^2}{\varepsilon} \right\rceil, \quad (71)$$

where Ω satisfies (70).

Proof. We show by induction that for $p \geq 0$

$$\|x_p - x_*\|^2 \leq R_0^2 \cdot 2^{-p} + 2(1 - 2^{-p}) \left(\frac{\delta}{\mu} + \frac{2\tilde{\delta}}{\mu} \right), \quad (72)$$

which leads to the statement of the Theorem. For $p = 0$ this inequality holds by the theorem's assumption. Assuming that (72) holds for some $p \geq 0$, our goal is to prove (72) for $p + 1$ considering the outer iteration $p + 1$. Observe that the function $d_p(x)$ defined in Algorithm 4 is 1-strongly convex w.r.t. the norm $\|\cdot\|/R_p$.

Using the definition of $d_p(\cdot)$ and (70), we have, since $x_p = \operatorname{argmin}_{x \in Q} d_p(x)$

$$V_p[x_p](x_*) = d_p(x_*) - d_p(x_p) - \langle \nabla d_p(x_p), x_* - x_p \rangle \leq d_p(x_*) \leq \frac{R_p^2 \Omega}{2}.$$

Denote by

$$S_{N_p} := \sum_{k=0}^{N_p-1} \frac{1}{L_{k+1}}.$$

Thus, by Theorem 4.8, taking $u = x_*$, we obtain

$$-\frac{1}{S_{N_p}} \sum_{k=0}^{N_p-1} \frac{\psi_\delta(x_*, w_k)}{L_{k+1}} \leq \frac{R_p^2 V_p[x_p](x_*)}{S_{N_p}} + \delta + 2\tilde{\delta} \leq \frac{\Omega R_p^2}{2S_{N_p}} + \delta + 2\tilde{\delta}.$$

Since the operator ψ is continuous and abstract monotone, we can assume that the solution to weak VI (36) is also a strong solution and $-\psi(w_k, x_*) \leq 0$, $k = 0, \dots, N_p - 1$ and, by Definition 5.1 (i), $-\psi_\delta(\omega_k, x_*) \leq \delta$ ($k = 0, \dots, N_p - 1$). This and (68) give, that for each $k = 0, \dots, N_p - 1$,

$$\begin{aligned} -\psi_\delta(x_*, w_k) &\geq -\delta - \psi_\delta(x_*, w_k) - \psi_\delta(w_k, x_*) \geq -\delta + \mu \|w_k - x_*\|^2, \\ -\psi_\delta(x_*, \omega_k) &\geq -\delta - \psi_\delta(x_*, \omega_k) - \psi_\delta(\omega_k, x_*) \geq -\delta + \mu \|\omega_k - x_*\|^2. \end{aligned}$$

Thus, by the convexity of the squared norm, we obtain

$$\begin{aligned} -2\delta + \mu \|x_{p+1} - x_*\|^2 &= -2\delta + \mu \left\| \frac{1}{S_{N_p}} \sum_{k=0}^{N_p-1} \frac{w_k}{L_{k+1}} - x_* \right\|^2 \leq -2\delta + \frac{\mu}{S_{N_p}} \sum_{k=0}^{N_p-1} \frac{\|w_k - x_*\|^2}{L_{k+1}} \\ &\leq -2\delta - \frac{1}{S_{N_p}} \sum_{k=0}^{N_p-1} \frac{\psi_\delta(x_*, w_k)}{L_{k+1}} \leq \frac{\Omega R_p^2}{2S_{N_p}} - \delta + 2\tilde{\delta}. \end{aligned}$$

Using the stopping criterion $S_{N_p} \geq \frac{\Omega}{\mu}$ we can put $N_p = \left\lceil \frac{2L\Omega}{\mu} \right\rceil$ and

$$\begin{aligned} \|x_{p+1} - x_*\|^2 &\leq \frac{R_p^2}{2} + \frac{\delta + 2\tilde{\delta}}{\mu} = \frac{1}{2} \left(R_0^2 \cdot 2^{-p} + 2(1 - 2^{-p}) \frac{\delta + 2\tilde{\delta}}{\mu} \right) + \frac{\delta + 2\tilde{\delta}}{\mu} \\ &= R_0^2 \cdot 2^{-(p+1)} + 2(1 - 2^{-(p+1)}) \left(\frac{\delta}{\mu} + \frac{2\tilde{\delta}}{\mu} \right), \end{aligned}$$

which finishes the proof by induction. \square

Remark 13. If for some $m > 0$ $\psi_\delta(x, y)$ is an m -strongly convex function in x , then for Algorithm 4 we can prove the following estimate

$$\|x_p - x_*\|^2 \leq R_0^2 \cdot 2^{-p} + \frac{2\delta}{m + \mu} + \frac{4\tilde{\delta}}{m + \mu} \leq \varepsilon + \frac{2\delta}{m + \mu} + \frac{4\tilde{\delta}}{m + \mu}$$

for each $p \geq 0$ and instead of (71) we obtain

$$\left\lceil \frac{2L\Omega}{m + \mu} \cdot \log_2 \frac{R_0^2}{\varepsilon} \right\rceil. \quad (73)$$

6. Conclusion

In this paper, we consider convex optimization problem (1). It is well known (see [16, 23, 34]) that if there is an inexact gradient $\nabla_\delta f(y)$ of f , s.t., for all $x, y \in Q$,

$$f(y) + \langle \nabla_\delta f(y), x - y \rangle - \delta_1 \leq f(x) \leq f(y) + \langle \nabla_\delta f(y), x - y \rangle + \frac{L}{2} \|x - y\|_2^2 + \delta_2, \quad (74)$$

then the corresponding versions of Gradient Method (GM) and Fast Gradient Method (FGM) have the convergence rate

$$f(x_N) - f(x_*) = O \left(\frac{LR^2}{N^p} + \delta_1 + N^{p-1} \delta_2 \right), \quad (75)$$

where $p = 1$ corresponds to GM and $p = 2$ corresponds to FGM, x_* is a solution of (1), R is an upper bound for $\|x_0 - x_*\|_2$. We show⁴ that under an appropriate generalization of (74) to

$$f(y) + \psi_\delta(x, y) - \delta_1 \leq f(x) \leq f(y) + \psi_\delta(x, y) + \frac{L}{2} \|x - y\|_2^2 + \delta_2$$

as well as appropriate generalizations of GM and FGM, the sequence generated by these methods satisfies (75). It should be noted that, despite there are many variants of FGM, we are aware of only one which can be generalized for problems with inexact model, namely accelerated mirror descent type of FGM [25, 42, 72]. An important

⁴For simplicity in the paper we consider the case $\delta_1 = \delta_2 = \delta$, but one can easily rewrite all the results of this paper to obtain (75). See [34] for details.

feature of this method is that it requires only one projection step on each iteration. A primal-dual extension of the proposed framework is made in [73].

We also show that in the case of μ -strongly convex objective (model) the estimate (75) can be improved to

$$f(x_N) - f(x_*) = O \left(\Delta f \exp \left(-O(1) \left(\frac{\mu}{L} \right)^{\frac{1}{p}} N \right) + \delta_1 + \left(\frac{L}{\mu} \right)^{\frac{p-1}{2}} \delta_2 \right),$$

where $\Delta f = f(x^0) - f(x_*)$, $p = 1$ for GM and $p = 2$ for restarted FGM.

In this paper we also propose a generalization of this inexact model framework for saddle-point problems and variational inequalities. We consider universal (adaptive) generalizations in the spirit of [56] and relative smoothness generalizations, generalizing the framework [7, 46] from optimization problems to saddle-point problems and VI. We also investigate the sensitivity of the convergence results to the accuracy of auxiliary minimization on each iteration.

Due to the lack of the space we only briefly mention here an extension of our framework for block-coordinate descent using the randomized version of FGM in [26] and stochastic optimization problems using the ideas from [30]. For the latter case we indicate that if we additionally assume that δ_1, δ_2 are independently chosen at each iteration random variables such that $\mathbb{E}\delta_1 = 0$ and $\delta_1, \sqrt{\delta_2}$ have correspondingly $(\delta'_1)^2$ -subgaussian variance and δ'_2 -subgaussian second moment [34] then with a high probability (75) changes to

$$f(x_N) - f(x_*) = \tilde{O} \left(\frac{LR^2}{N^p} + \frac{\delta'_1}{\sqrt{N}} + N^{p-1}\delta'_2 \right).$$

From this result and mini-batch trick [30] one can obtain the main estimates for convex and strongly convex stochastic optimization problems [21, 28, 34, 41].

As further generalizations we point a generalization for tensor methods [31, 58] and for incremental and variance reduction methods for finite-sum minimization [14, 44].

References

- [1] A. Anikin, P. Dvurechensky, A. Gasnikov, A. Golov, A. Gornov, Y. Maximov, M. Mendel, and V. Spokoiny, *Modern efficient numerical approaches to regularized regression problems in application to traffic demands matrix calculation from link loads*, in *Proceedings of International conference ITAS-2015. Russia, Sochi*. 2015. arXiv:1508.00858.
- [2] A.S. Anikin, A.V. Gasnikov, P.E. Dvurechensky, A.I. Tyurin, and A.V. Chernov, *Dual approaches to the minimization of strongly convex functionals with a simple structure under affine constraints*, *Computational Mathematics and Mathematical Physics* 57 (2017), pp. 1262–1276.
- [3] K. Antonakopoulos, E.V. Belmega, and P. Mertikopoulos, *Adaptive extra-gradient methods for min-max optimization and games*, arXiv:2010.12100 (2020).
- [4] K. Antonakopoulos, V. Belmega, and P. Mertikopoulos, *An adaptive mirror-prox method for variational inequalities with singular operators*, in *Advances in Neural Information Processing Systems 32*, H. Wallach, H. Larochelle, A. Beygelzimer, F. dAlché Buc, E. Fox, and R. Garnett, eds., Curran Associates, Inc., 2019, pp. 8455–8465.
- [5] D. Baimurzina, A. Gasnikov, E. Gasnikova, P. Dvurechensky, E. Ershov, M. Kubentaeva, and A. Lagunovskaya, *Universal similar triangulars method for searching equilibriums in*

- traffic flow distribution models*, Journal of Computational Mathematics and Mathematical Physics 59 (2019), pp. 21–36.
- [6] T.Q. Bao and P.Q. Khanh, *Some algorithms for solving mixed variational inequalities*, Acta Mathematica Vietnamica 31 (2006), pp. 77–98.
 - [7] H.H. Bauschke, J. Bolte, and M. Teboulle, *A descent lemma beyond lipschitz gradient continuity: first-order methods revisited and applications*, Mathematics of Operations Research 42 (2016), pp. 330–348.
 - [8] A. Beck and M. Teboulle, *A fast iterative shrinkage-thresholding algorithm for linear inverse problems*, SIAM Journal on Imaging Sciences 2 (2009), pp. 183–202. Available at <https://doi.org/10.1137/080716542>.
 - [9] L. Bogolubsky, P. Dvurechensky, A. Gasnikov, G. Gusev, Y. Nesterov, A.M. Raigorodskii, A. Tikhonov, and M. Zhukovskii, *Learning supervised pagerank with gradient-based and gradient-free optimization methods*, in *Advances in Neural Information Processing Systems 29*, D.D. Lee, M. Sugiyama, U.V. Luxburg, I. Guyon, and R. Garnett, eds., Curran Associates, Inc., 2016, pp. 4914–4922. arXiv:1603.00717.
 - [10] A. Chambolle and T. Pock, *A first-order primal-dual algorithm for convex problems with applications to imaging*, Journal of Mathematical Imaging and Vision 40 (2011), pp. 120–145.
 - [11] G. Chen and M. Teboulle, *Convergence analysis of a proximal-like minimization algorithm using bregman functions*, SIAM Journal on Optimization 3 (1993), pp. 538–543.
 - [12] A. Chernov, P. Dvurechensky, and A. Gasnikov, *Fast Primal-Dual Gradient Method for Strongly Convex Minimization Problems with Linear Constraints*, in *Discrete Optimization and Operations Research: 9th International Conference, DOOR 2016, Vladivostok, Russia, September 19-23, 2016, Proceedings*, Y. Kochetov, M. Khachay, V. Beresnev, E. Nurminski, and P. Pardalos, eds. Springer International Publishing, 2016, pp. 391–403.
 - [13] M.B. Cohen, A. Sidford, and K. Tian, *Relative Lipschitzness in Extragradient Methods and a Direct Recipe for Acceleration*, in *12th Innovations in Theoretical Computer Science Conference, ITCS 2021, January 6-8, 2021, Virtual Conference*, J.R. Lee, ed., LIPIcs Vol. 185. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021, pp. 62:1–62:18. Available at <https://doi.org/10.4230/LIPIcs.ITCS.2021.62>.
 - [14] A. Defazio, *A simple practical accelerated method for finite sums*, in *Advances in neural information processing systems*. 2016, pp. 676–684.
 - [15] O. Devolder, F. Glineur, and Y. Nesterov, *First-order methods with inexact oracle: the strongly convex case*, CORE Discussion Papers 2013/16 (2013).
 - [16] O. Devolder, F. Glineur, and Y. Nesterov, *First-order methods of smooth convex optimization with inexact oracle*, Mathematical Programming 146 (2014), pp. 37–75. Available at <http://dx.doi.org/10.1007/s10107-013-0677-5>.
 - [17] R.A. Dragomir, A. Taylor, A. d’Aspremont, and J. Bolte, *Optimal complexity and certification of bregman first-order methods*, arXiv:1911.08510 (2019).
 - [18] D. Drusvyatskiy, A.D. Ioffe, and A.S. Lewis, *Nonsmooth optimization using Taylor-like models: error bounds, convergence, and termination criteria*, Mathematical Programming (2019). arXiv:1610.03446.
 - [19] D. Dvinskikh, E. Gorbunov, A. Gasnikov, P. Dvurechensky, and C.A. Uribe, *On Primal-Dual Approach for Distributed Stochastic Convex Optimization over Networks*, in *2019 IEEE Conference on Decision and Control (CDC)*. 2019. (accepted), arXiv:1903.09844.
 - [20] P. Dvurechensky, D. Dvinskikh, A. Gasnikov, C.A. Uribe, and A. Nedić, *Decentralize and Randomize: Faster Algorithm for Wasserstein Barycenters*, in *Advances in Neural Information Processing Systems 31*, S. Bengio, H. Wallach, H. Larochelle, K. Grauman, N. Cesa-Bianchi, and R. Garnett, eds. Curran Associates, Inc., NIPS’18, 2018, pp. 10783–10793. Available at <http://papers.nips.cc/paper/8274-decentralize-and-randomize-faster-algorithm-for-wasserstein-barycenters.pdf>, arXiv:1802.04367.
 - [21] P. Dvurechensky and A. Gasnikov, *Stochastic intermediate gradient method for convex problems with stochastic inexact oracle*, Journal of Optimization Theory and

- Applications 171 (2016), pp. 121–145. Available at <http://dx.doi.org/10.1007/s10957-016-0999-6>.
- [22] P. Dvurechensky, A. Gasnikov, E. Gasnikova, S. Matsievsky, A. Rodomanov, and I. Usik, *Primal-Dual Method for Searching Equilibrium in Hierarchical Congestion Population Games*, in *Supplementary Proceedings of the 9th International Conference on Discrete Optimization and Operations Research and Scientific School (DOOR 2016) Vladivostok, Russia, September 19 - 23, 2016*. 2016, pp. 584–595. arXiv:1606.08988.
 - [23] P. Dvurechensky, A. Gasnikov, and D. Kamzolov, *Universal intermediate gradient method for convex problems with inexact oracle*, arXiv:1712.06036, Opt. Meth. & Software (accepted) (2019). Available at <http://dx.doi.org/10.1080/10556788.2019.1711079>.
 - [24] P. Dvurechensky, A. Gasnikov, and A. Kroshnin, *Computational Optimal Transport: Complexity by Accelerated Gradient Descent Is Better Than by Sinkhorn’s Algorithm*, in *Proceedings of the 35th International Conference on Machine Learning*, J. Dy and A. Krause, eds., Proceedings of Machine Learning Research Vol. 80. 2018, pp. 1367–1376. arXiv:1802.04367.
 - [25] P. Dvurechensky, A. Gasnikov, S. Omelchenko, and A. Tiurin, *Adaptive similar triangles method: a stable alternative to sinkhorn’s algorithm for regularized optimal transport*, arXiv:1706.07622 (2017).
 - [26] P. Dvurechensky, A. Gasnikov, and A. Tiurin, *Randomized similar triangles method: A unifying framework for accelerated randomized optimization methods (coordinate descent, directional search, derivative-free method)*, arXiv:1707.08486 (2017).
 - [27] M. Frank and P. Wolfe, *An algorithm for quadratic programming*, Naval Research Logistics Quarterly 3 (1956), pp. 95–110.
 - [28] A.V. Gasnikov and P.E. Dvurechensky, *Stochastic intermediate gradient method for convex optimization problems*, Doklady Mathematics 93 (2016), pp. 148–151.
 - [29] A.V. Gasnikov, P.E. Dvurechensky, F.S. Stonyakin, and A.A. Titov, *An adaptive proximal method for variational inequalities*, Computational Mathematics and Mathematical Physics 59 (2019), pp. 836–841. Available at <https://doi.org/10.1134/S0965542519050075>.
 - [30] A. Gasnikov, *Universal gradient descent*, arXiv preprint arXiv:1711.00394 (2017).
 - [31] A. Gasnikov, P. Dvurechensky, E. Gorbunov, E. Vorontsova, D. Selikhanovych, C.A. Uribe, B. Jiang, H. Wang, S. Zhang, S. Bubeck, Q. Jiang, Y.T. Lee, Y. Li, and A. Sidford, *Near Optimal Methods for Minimizing Convex Functions with Lipschitz p -th Derivatives*, in *Proceedings of the Thirty-Second Conference on Learning Theory*, A. Beygelzimer and D. Hsu, eds., Proceedings of Machine Learning Research Vol. 99, 25–28 Jun, Phoenix, USA. PMLR, 2019, pp. 1392–1393. Available at <http://proceedings.mlr.press/v99/gasnikov19b.html>, arXiv:1809.00382.
 - [32] A. Gasnikov and A. Tyurin, *Fast gradient descent for convex minimization problems with an oracle producing a (δ, l) -model of function at the requested point*, Computational Mathematics and Mathematical Physics 59 (2019), pp. 1085–1097.
 - [33] A.V. Gasnikov and Y.E. Nesterov, *Universal method for stochastic composite optimization problems*, Computational Mathematics and Mathematical Physics 58 (2018), pp. 48–64. First appeared in arXiv:1604.05275.
 - [34] E. Gorbunov, D. Dvinskikh, and A. Gasnikov, *Optimal decentralized distributed algorithms for stochastic convex optimization*, arXiv preprint arXiv:1911.07363 (2019).
 - [35] S.V. Guminov, Y.E. Nesterov, P.E. Dvurechensky, and A.V. Gasnikov, *Accelerated primal-dual gradient descent with linesearch for convex, nonconvex, and nonsmooth optimization problems*, Doklady Mathematics 99 (2019), pp. 125–128.
 - [36] Z. Harchaoui, A. Juditsky, and A. Nemirovski, *Conditional gradient algorithms for norm-regularized smooth convex optimization*, // Mathematical Programming 152 (2015), pp. 75–112.
 - [37] A.N. Iusem, A. Jofré, R.I. Oliveira, and P. Thompson, *Variance-based extragradient methods with line search for stochastic variational inequalities*, SIAM Journal on Optimization 29 (2019), pp. 175–206. arXiv:1703.00262.

- [38] M. Jaggi, *Revisiting Frank-Wolfe: Projection-Free Sparse Convex Optimization.*, in *ICML (1)*. 2013, pp. 427–435.
- [39] I. Konnov and R. Salahutdin, *Two-level iterative method for non-stationary mixed variational inequalities*, *Izvestija vysshih uchebnyh zavedenij. Matematika* 61 (2017), pp. 50–61.
- [40] A. Kroshnin, N. Tupitsa, D. Dvinskikh, P. Dvurechensky, A. Gasnikov, and C. Uribe, *On the Complexity of Approximating Wasserstein Barycenters*, in *Proceedings of the 36th International Conference on Machine Learning*, K. Chaudhuri and R. Salakhutdinov, eds., *Proceedings of Machine Learning Research* Vol. 97, 09–15 Jun, Long Beach, California, USA. PMLR, 2019, pp. 3530–3540. arXiv:1901.08686.
- [41] A. Kulunchakov and J. Mairal, *Estimate sequences for stochastic composite optimization: Variance reduction, acceleration, and robustness to noise*, arXiv preprint arXiv:1901.08788 (2019).
- [42] G. Lan, *An optimal method for stochastic composite optimization*, *Mathematical Programming* 133 (2012), pp. 365–397. Available at <https://doi.org/10.1007/s10107-010-0434-y>, First appeared in June 2008.
- [43] G. Lan, *Bundle-level type methods uniformly optimal for smooth and nonsmooth convex optimization*, *Mathematical Programming* 149 (2015), pp. 1–45.
- [44] G. Lan and Y. Zhou, *Random gradient extrapolation for distributed and stochastic optimization*, *SIAM Journal on Optimization* 28 (2018), pp. 2753–2782.
- [45] H. Lin, J. Mairal, and Z. Harchaoui, *A Universal Catalyst for First-order Optimization*, in *Proceedings of the 28th International Conference on Neural Information Processing Systems*, Cambridge, MA, USA. MIT Press, NIPS’15, 2015, pp. 3384–3392. Available at <http://dl.acm.org/citation.cfm?id=2969442.2969617>.
- [46] H. Lu, R.M. Freund, and Y. Nesterov, *Relatively smooth convex optimization by first-order methods, and applications*, *SIAM Journal on Optimization* 28 (2018), pp. 333–354.
- [47] J. Mairal, *Optimization with first-order surrogate functions*, in *International Conference on Machine Learning*. 2013, pp. 783–791.
- [48] Y. Malitsky, *Proximal extrapolated gradient methods for variational inequalities*, *Optimization Methods and Software* 33 (2018), pp. 140–164. Available at <https://doi.org/10.1080/10556788.2017.1300899>, PMID: 29348705.
- [49] Y. Malitsky, *Golden ratio algorithms for variational inequalities*, *Mathematical Programming* 184 (2020), pp. 383–410. Available at <https://doi.org/10.1007/s10107-019-01416-w>.
- [50] G. Mastroeni, *On auxiliary principle for equilibrium problems*, *Publicatione del Dipartimento di Matematica Dell’Universita di Pisa* 3 (2000), pp. 1244–1258.
- [51] A. Nemirovski, *Prox-method with rate of convergence $O(1/t)$ for variational inequalities with Lipschitz continuous monotone operators and smooth convex-concave saddle point problems*, *SIAM Journal on Optimization* 15 (2004), pp. 229–251.
- [52] A. Nemirovski, S. Onn, and U.G. Rothblum, *Accuracy certificates for computational problems with convex structure*, *Mathematics of Operations Research* 35 (2010), pp. 52–78.
- [53] A. Nemirovskii and Y. Nesterov, *Optimal methods of smooth convex minimization*, *USSR Computational Mathematics and Mathematical Physics* 25 (1985), pp. 21 – 30. Available at <http://www.sciencedirect.com/science/article/pii/0041555385901004>.
- [54] Y. Nesterov, *Primal-dual subgradient methods for convex problems*, *Mathematical Programming* 120 (2009), pp. 221–259. Available at <https://doi.org/10.1007/s10107-007-0149-x>, First appeared in 2005 as CORE discussion paper 2005/67.
- [55] Y. Nesterov, *Gradient methods for minimizing composite functions*, *Mathematical Programming* 140 (2013), pp. 125–161. First appeared in 2007 as CORE discussion paper 2007/76.
- [56] Y. Nesterov, *Universal gradient methods for convex optimization problems*, *Mathematical Programming* 152 (2015), pp. 381–404. Available at <http://dx.doi.org/10.1007/s10107-014-0790-0>.
- [57] Y. Nesterov, *Complexity bounds for primal-dual methods minimizing the model of objective function*, *Math. Program.* 171 (2018), pp. 311–330. Available at <https://doi.org/10.1007/s10107-018-0000-0>.

- 1007/s10107-017-1188-6.
- [58] Y. Nesterov, *Implementable tensor methods in unconstrained convex optimization*, Tech. Rep., CORE UCL, 2018. Available at https://alfresco.uclouvain.be/alfresco/service/guest/streamDownload/workspace/SpacesStore/aabc2323-0bc1-40d4-9653-1c29971e7bd8/coredp2018_05web.pdf, CORE Discussion Paper 2018/05.
 - [59] Y. Nesterov, *Lectures on convex optimization*, Vol. 137, Springer International Publishing, 2018.
 - [60] Y. Nesterov, *Soft clustering by convex electoral model*, CORE Discussion Papers 2018001, Université catholique de Louvain, Center for Operations Research and Econometrics (CORE), 2018. Available at <https://ideas.repec.org/p/cor/louvco/2018001.html>.
 - [61] Y. Nesterov and M.I. Florea, *Gradient methods with memory*, Optimization Methods and Software 0 (2021), pp. 1–18. Available at <https://doi.org/10.1080/10556788.2020.1858831>.
 - [62] Y. Nesterov, A. Gasnikov, S. Guminov, and P. Dvurechensky, *Primal-dual accelerated gradient methods with small-dimensional relaxation oracle*, arXiv:1809.05895 (2018). Submitted to Optimization Methods & Software.
 - [63] Y. Nesterov and B. Polyak, *Cubic regularization of newton method and its global performance*, Mathematical Programming 108 (2006), pp. 177–205. Available at <http://dx.doi.org/10.1007/s10107-006-0706-8>.
 - [64] P. Ochs, J. Fadili, and T. Brox, *Non-smooth non-convex bregman minimization: Unification and new algorithms*, arXiv preprint arXiv:1707.02278 (2017).
 - [65] A. Ogaltsov, D. Dvinskikh, P. Dvurechensky, A. Gasnikov, and V. Spokoiny, *Adaptive gradient descent for convex and non-convex stochastic optimization*, arXiv:1911.08380 (2019). Submitted to IFAC 2020 Congress.
 - [66] Y. Ouyang and Y. Xu, *Lower complexity bounds of first-order methods for convex-concave bilinear saddle-point problems*, arXiv preprint arXiv: 1808.02901 (2018).
 - [67] N. Parikh and S. Boyd, *Proximal algorithms*, Foundations and Trends® in Optimization 1 (2014), pp. 127–239. Available at <http://dx.doi.org/10.1561/2400000003>.
 - [68] K. Scaman, F. Bach, S. Bubeck, Y.T. Lee, and L. Massoulié, *Optimal Algorithms for Smooth and Strongly Convex Distributed Optimization in Networks*, in *Proceedings of the 34th International Conference on Machine Learning*, D. Precup and Y.W. Teh, eds., Proceedings of Machine Learning Research Vol. 70, 06–11 Aug, International Convention Centre, Sydney, Australia. PMLR, 2017, pp. 3027–3036. Available at <http://proceedings.mlr.press/v70/scaman17a.html>.
 - [69] F. Stonyakin, A. Gasnikov, P. Dvurechensky, M. Alkousa, and A. Titov, *Generalized Mirror Prox for monotone variational inequalities: Universality and inexact oracle*, arXiv:1806.05140 (2018).
 - [70] F. Stonyakin, A. Gasnikov, A. Tyurin, D. Pasechnyuk, A. Agafonov, P. Dvurechensky, D. Dvinskikh, A. Kroshnin, and V. Piskunova, *Inexact model: A framework for optimization and variational inequalities*, arXiv:1902.00990 (2019). WIAS Preprint No. 2679.
 - [71] F.S. Stonyakin, D. Dvinskikh, P. Dvurechensky, A. Kroshnin, O. Kuznetsova, A. Agafonov, A. Gasnikov, A. Tyurin, C.A. Uribe, D. Pasechnyuk, and S. Artamonov, *Gradient Methods for Problems with Inexact Model of the Objective*, in *Mathematical Optimization Theory and Operations Research*, M. Khachay, Y. Kochetov, and P. Pardalos, eds., Cham. Springer International Publishing, 2019, pp. 97–114. arXiv:1902.09001.
 - [72] P. Tseng, *On accelerated proximal gradient methods for convex-concave optimization*, Tech. Rep., MIT, 2008. Available at <http://www.mit.edu/~dimitrib/PTseng/papers/apgm.pdf>.
 - [73] A. Tyurin, *Primal-dual fast gradient method with a model*, arXiv preprint arXiv:1906.10107 (2019).
 - [74] C.A. Uribe, D. Dvinskikh, P. Dvurechensky, A. Gasnikov, and A. Nedić, *Distributed Computation of Wasserstein Barycenters over Networks*, in *2018 IEEE 57th Annual Conference on Decision and Control (CDC)*. 2018. Accepted, arXiv:1803.02933.

- [75] Y. Xie, X. Wang, R. Wang, and H. Zha, *A fast proximal point method for wasserstein distance*, arXiv preprint arXiv:1802.04307 (2018).
- [76] R. Zhao and R.M. Freund, *Analysis of the frank-wolfe method for logarithmically-homogeneous barriers, with an extension*, arXiv:2010.08999 (2020).

Appendix A. Auxiliary facts

Let us comment the inexact solution of the auxiliary problem.

Remark 14. We can show that if $\tilde{x} \in \text{Argmin}_{x \in Q}^{\tilde{\delta}} \Psi(x)$, then $\Psi(\tilde{x}) - \Psi(x_*) \leq \delta$. Indeed, we have $\Psi(x_*) \geq \Psi(\tilde{x}) + \langle h, x_* - \tilde{x} \rangle \geq \Psi(\tilde{x}) - \tilde{\delta}$. The converse statement is not always true. However, for some general cases we can resolve the problem (see [32] and Example A.1).

Example A.1. Let us show an example, how we can resolve the problem in Remark 14. Note, that if $\Psi(x)$ is μ -strongly convex; has L -Lipschitz continuous gradient in $\|\cdot\|$ norm. To say more precisely

$$L = \max_{\|h\| \leq 1, x \in [\tilde{x}, x_*]} \langle h, \nabla^2 \Psi(x) h \rangle.$$

and $R = \max_{x, y \in Q} \|x - y\|$, then $\Psi(\tilde{x}) - \Psi(x_*) \leq \tilde{\epsilon}$ entails that [71]

$$\tilde{\delta} \leq (LR + \|\nabla \Psi(x_*)\|_*) \sqrt{2\tilde{\epsilon}/\mu}, \quad (\text{A1})$$

where $x_* = \text{argmin}_{x \in Q} \Psi(x)$. If one can guarantee that $\nabla \Psi(x_*) = 0$, then (A1) can be improved $\tilde{\delta} \leq R\sqrt{2L\tilde{\epsilon}}$.

Appendix B. Proof for Lemma 3.7

Proof. In view of the definition (24) of sequence α_{k+1} , we have:

$$\begin{aligned} A_N &\leq A_N(1 + \mu A_{N-1} + m A_{N-1}) = L_N(A_N - A_{N-1})^2 \\ &\leq L_N(A_N^{1/2} - A_{N-1}^{1/2})^2(A_N^{1/2} + A_{N-1}^{1/2})^2 \leq 4L_N A_N (A_N^{1/2} - A_{N-1}^{1/2})^2. \end{aligned}$$

We can see that

$$A_N^{1/2} \geq A_{N-1}^{1/2} + \frac{1}{2\sqrt{L_N}}$$

and

$$A_N \geq \frac{1}{4} \left(\sum_{k=0}^{N-1} \frac{1}{\sqrt{L_{k+1}}} \right)^2.$$

For the case when $\mu + m > 0$ we obtain:

$$(\mu + m)A_{N-1}A_N \leq A_N(1 + \mu A_{N-1} + mA_{N-1}) \leq 4L_N A_N (A_N^{1/2} - A_{N-1}^{1/2})^2.$$

From the fact that $A_1 = 1/L_1$ and the last inequality we can show that

$$A_N^{1/2} \geq \left(1 + \sqrt{\frac{\mu + m}{4L_N}}\right) A_{N-1}^{1/2} \geq \frac{1}{\sqrt{L_1}} \prod_{k=1}^{N-1} \left(1 + \sqrt{\frac{\mu + m}{4L_{k+1}}}\right).$$

□

Appendix C. Fast gradient method with $(\delta, L, \mu, m, V, \|\cdot\|)$ -model. Restart technique.

Let us consider the case of a smooth strongly convex function f and show how to accelerate the work of Algorithm 2 using the restart technique. Let us assume that

$$\psi_\delta(x, x_*) \geq 0 \quad \forall x \in Q. \quad (\text{C1})$$

Note that this assumption is natural, e.g. $\psi_\delta(x, y) := \langle \nabla f(y), x - y \rangle \quad \forall x, y \in Q$. We also modify the concept of μ -strong convexity in the following way.

Definition C.1. We say that the function f is a left relative μ -strongly convex if the following inequality

$$\mu V[x](y) \leq f(x) - f(y) - \psi_\delta(x, y) \quad \forall x, y \in Q$$

holds.

Remark 15. Let us remind that if $d(x - y) \leq C_n \|x - y\|^2$ for $C_n = O(\log n)$, (where n is the dimension of vectors from Q) then $V[y](x) \leq C_n \|x - y\|^2$. This assumption is true for many standard proximal setups. In this case the condition of (μC_n) -strong convexity

$$\mu C_n \|x - y\|^2 + f_\delta(y) + \psi_\delta(x, y) \leq f(x)$$

entails right relative strong convexity:

$$\mu V[y](x) + f_\delta(y) + \psi_\delta(x, y) \leq f(x).$$

Note that the concepts of right and left relative strongly convexity from Definitions 2.1 and C.1 are equivalent in the case of an assumption from Remark 15 ($V[x](y) \leq C_n \|x - y\|^2$ for each $x, y \in Q$).

We show that using the restart technique can also accelerate the work of non-adaptive version of Algorithm 1 ($L_{k+1} = L$) for $(\delta, L, 0, 0, V, \|\cdot\|)$ -model and relative μ -strongly convex function f in the sense of Definition C.1:

$$\mu V[x](y) + f(y) + \psi_\delta(x, y) - \delta \leq f(x) \leq f(y) + \psi_\delta(x, y) + \frac{L}{2} \|x - y\|^2 + \delta.$$

for each $x, y \in Q$. By Theorem 3.4 and Remark 6, we have:

$$f(x_N) - f(x_*) \leq \frac{4LV[x_0](x_*)}{N^2} + \frac{4L\tilde{\delta}}{N} + 2N\delta. \quad (\text{C2})$$

Consider the case of relatively μ -strongly convex function f . We will use the restart technique to obtain the method for strongly convex functions.

Theorem C.2. *Let f be a left relative μ -strongly convex function and $\psi_\delta(x, y)$ is a $(\delta, L, 0, 0, V, \|\cdot\|)$ -model. Let δ and $\tilde{\delta}$ satisfy $\frac{4\mu\sqrt{10}}{L} \left(5\delta \left\lceil \sqrt{\frac{L}{\mu}} \right\rceil^3 + \tilde{\delta}L \left\lceil \sqrt{\frac{L}{\mu}} \right\rceil \right) \leq \varepsilon$. Then, using the restarts of Algorithm 1, we need*

$$N = \left\lceil \log_2 \frac{\mu R^2}{\varepsilon} \right\rceil \cdot \left\lceil \sqrt{\frac{10L}{\mu}} \right\rceil.$$

iterations to achieve an ε accuracy by function: $f(x_N) - f(x_*) \leq \varepsilon$.

Proof. By (C2) and Definition C.1, we have:

$$\mu V[x_{N_1}](x_*) \leq f(x_{N_1}) - f(x_*) \leq \frac{4LV[x_0](x_*)}{N^2} + \frac{4L\tilde{\delta}}{N} + 2N\delta. \quad (\text{C3})$$

Let's choose N_1 so that the following inequality holds:

$$\frac{4L\tilde{\delta}}{N_1} + 2N_1\delta \leq \frac{LV[x_0](x_*)}{N_1^2}. \quad (\text{C4})$$

We restart method as $V[x_{N_1}](x_*) \leq \frac{V[x_0](x_*)}{2}$. Using (C3), we obtain an estimation for the number of iterations on the first restart: $\frac{5L}{\mu N_1^2} \leq \frac{1}{2}$. Therefore, let's choose

$$N_1 = \left\lceil \sqrt{\frac{10L}{\mu}} \right\rceil. \quad (\text{C5})$$

Then after N_1 iterations we restart method. Similarly, we restart after N_2 iterations, such that $V[x_{N_2}](x_*) \leq \frac{V[x_{N_1}](x_*)}{2}$. We obtain $N_2 = \left\lceil \sqrt{\frac{10L}{\mu}} \right\rceil$. So, after p -th restart the total number of iterations is $M = p \cdot \left\lceil \sqrt{\frac{10L}{\mu}} \right\rceil$.

Now let's consider how many iterations are needed to achieve the accuracy $\varepsilon = f(x_{N_p}) - f(x_*)$. From (C2) and (C5) we take $p = \left\lceil \log_2 \frac{\mu R^2}{\varepsilon} \right\rceil$ and the total number of iterations is $M = \left\lceil \log_2 \frac{\mu R^2}{\varepsilon} \right\rceil \cdot \left\lceil \sqrt{\frac{10L}{\mu}} \right\rceil$.

We have chosen our errors as δ and $\tilde{\delta}$ to satisfy (C4). Indeed, from (C4) using

$N_k = \left\lceil \sqrt{\frac{10L}{\mu}} \right\rceil$ we can deduce the following inequality:

$$\varepsilon \geq \frac{4\sqrt{10}\mu}{L} \left(5\delta \left\lceil \sqrt{\frac{L}{\mu}} \right\rceil^3 + \tilde{\delta}L \left\lceil \sqrt{\frac{L}{\mu}} \right\rceil \right).$$

One can see that such a choice of δ and $\tilde{\delta}$ as above satisfies that inequality. \square

Remark 16. Partially, we can choose $\delta = O\left(\frac{\varepsilon L}{\mu \left\lceil \sqrt{\frac{10L}{\mu}} \right\rceil^3}\right)$ and $\tilde{\delta} = O\left(\frac{\varepsilon}{\mu \left\lceil \sqrt{\frac{10L}{\mu}} \right\rceil}\right)$.

Appendix D. Numerical results for some different geometrical problems

To show the advantages of the proposed Algorithm 3, a series of numerical experiments were performed for different examples.

We consider an example of a variational inequality with a non-smooth and non-strongly monotone operator, i.e. with $\nu = 0$, for which the proposed Algorithm 3, due to its adaptivity to the level of smoothness of the problem, in practice works with iteration complexity much smaller than the predicted by theory. We compare the performance of Algorithm 3, with two different approaches for step-size selection, the first one is the proposed procedure in this paper, i.e. with $L_{k+1} = 2^{i_k-1}L_k$, the second procedure with $L_{k+1} = 2^{i_k}L_k$, i.e. with a non-increasing stepsize similarly to [4], for three different geometrical problems with some non-smooth functional constraints.

Let Q be a convex compact set and $A_k \in \mathbb{R}^n, k = 1, \dots, N$ be a given set of N points.

The first problem is naturally connected with the analogue of the well-known *smallest covering circle problem*. For this problem we need to solve the following optimization problem

$$\min_{x \in Q} \left\{ f(x) := \max_{1 \leq k \leq N} \|x - A_k\|_2 \mid \varphi_p(x) := \sum_{i=1}^n \alpha_{pi}|x_i| - 1 \leq 0, p = 1, \dots, m \right\}. \quad (\text{D1})$$

The second problem is naturally connected with the analogue of the well-known *Fermat–Torricelli–Steiner problem*. For this problem we need to solve the following optimization problem

$$\min_{x \in Q} \left\{ f(x) := \sum_{k=1}^N \|x - A_k\|_2 \mid \varphi_p(x) := \sum_{i=1}^n \alpha_{pi}|x_i| - 1 \leq 0, p = 1, \dots, m \right\}. \quad (\text{D2})$$

The third problem is connected with *problem of the best approximation of the distance from the given set Q* . For this problem, let $A \notin Q$ be a given point, we need to solve the following optimization problem

$$\min_{x \in Q} \left\{ f(x) := \|x - A\|_2 \mid \varphi_p(x) := \sum_{i=1}^n \alpha_{pi}|x_i| - 1 \leq 0, p = 1, \dots, m \right\}. \quad (\text{D3})$$

The coefficients α_{pi} , in (D1), (D2) and (D3) are drawn from the standard normal distribution and then truncated to be positive.

The corresponding Lagrange saddle-point problem is defined as

$$\min_{x \in Q_1} \max_{\vec{\lambda} = (\lambda_1, \lambda_2, \dots, \lambda_m)^T \in Q_2} L(x, \lambda) := f(x) + \sum_{p=1}^m \lambda_p \varphi_p(x),$$

This problem is equivalent to the variational inequality with monotone bounded operator

$$g(x, \lambda) = \begin{pmatrix} \nabla f(x) + \sum_{p=1}^m \lambda_p \nabla \varphi_p(x), \\ (-\varphi_1(x), -\varphi_2(x), \dots, -\varphi_m(x))^T \end{pmatrix}.$$

For simplicity, we assume that there exist (potentially very large) bound for the optimal Lagrange multiplier $\vec{\lambda}^*$, which allows us to compactify the feasible set for the pair $(x, \vec{\lambda})$ to be an Euclidean ball of some radius.

We run Algorithm 3 for different values of n and m with standard Euclidean prox-structure and the starting point $(x^0, \vec{\lambda}^0) = \frac{1}{\sqrt{m+n}} \mathbf{1} \in \mathbb{R}^{n+m}$, where $\mathbf{1}$ is the vector of all ones. Points A_k , $k = 1, \dots, N$, in (D1), (D2) and the point A in (D3) are chosen randomly from standard normal distribution. The set Q_1 is chosen as the unit ball in \mathbb{R}^n , also Q_2 is chosen as the unit ball in \mathbb{R}_+^m .

All experiments were implemented in Python 3.4, on a computer fitted with Intel(R) Core(TM) i7-8550U CPU @ 1.80GHz, 1992 Mhz, 4 Core(s), 8 Logical Processor(s). RAM of the computer is 8 GB.

The results of the comparison, for problems (D1), (D2) and (D3), are represented in Tables D1, D2 and D3, respectively. These results demonstrate the number of the produced iterations to reach the ε -solution of the three considered problems, and the running time of the algorithm in seconds, with different values of $\varepsilon \in \{1/2^i, i = 1, 2, 3, 4, 5, 6\}$.

In general, from all conducted experiments, we can see that, in order to the number of iterations and the running time, the proposed procedure of the step-size selection in Algorithm 3 is the best, the efficiency of this procedure is represented by its very high execution speed, where by the proposed Algorithm 3 one needs a few seconds and part of seconds to achieve the solution and to reach its stopping criterion.

Table D1.: The results for the problem D1, with $n = 100, m = 100$ and $N = 50$.

$\frac{1}{\varepsilon}$	Algorithm 3, with $L_{k+1} = 2^{i_k-1} L_k$.		Algorithm 3, with $L_{k+1} = 2^{i_k} L_k$.	
	Iterations	Time (sec.)	Iterations	Time (sec.)
2	6	0.242	82	3.985
4	7	0.272	163	9.207
8	8	0.330	646	35.780
16	10	0.443	2587	142.047
32	11	0.479	10371	530.594
64	12	0.513	41558	2081.414

Table D2.: The results for the problem (D2), with $n = 100, m = 100$ and $N = 50$.

$\frac{1}{\varepsilon}$	Algorithm 3, with $L_{k+1} = 2^{i_k-1} L_k.$		Algorithm 3, with $L_{k+1} = 2^{i_k} L_k.$	
	Iterations	Time (sec.)	Iterations	Time (sec.)
2	8	2.997	95	27.527
4	9	3.459	189	52.421
8	11	4.346	378	105.997
16	12	4.124	756	218.519
32	13	4.615	1511	473.350
64	15	5.195	3021	881.917

Table D3.: The results for the problem D3, with $n = 500$ and $m = 100$.

$\frac{1}{\varepsilon}$	Algorithm 3, with $L_{k+1} = 2^{i_k-1} L_k.$		Algorithm 3, with $L_{k+1} = 2^{i_k} L_k.$	
	Iterations	Time (sec.)	Iterations	Time (sec.)
2	8	1.273	86	11.293
4	9	1.365	342	42.407
8	10	1.491	684	100.926
16	12	1.837	1367	171.827
32	13	1.982	2733	358.685
64	15	2.382	5465	732.999

Appendix E. Numerical results for relatively smooth objective function

Let

$$f(x) := \frac{1}{4} \|Ax - b\|_4^4 + \frac{1}{2} \|Cx - d\|_2^2.$$

where $A, C \in \mathbb{R}^{n \times n}$ and $b, d \in \mathbb{R}^n$ [46]. We consider the following constrained optimization problem

$$\min_{x \in Q} \left\{ f(x) \mid \varphi_p(x) := \sum_{i=1}^n \alpha_{pi} x_i - 1 \leq 0, p = 1, \dots, m \right\}, \quad (\text{E1})$$

where Q is a convex compact, α_{pi} are drawn from the standard normal distribution and then truncated to be positive. The corresponding Lagrange saddle-point problem and the monotone non-smooth operator associated with the variational inequality problem, defined similarly as in the Appendix D.

We run Algorithm 3, for different values of n and m with standard Euclidean prox-structure and the same inputs as in Appendix D. The results are represented in Table E1. These results demonstrate the number of the produced iterations, by Algorithm 3, to reach the ε -solution of (E1), and the running time of the algorithm in seconds, with different values of $\varepsilon \in \{1/2^i, i = 1, 2, 3, 4, 5, 6\}$.

Now let us consider the unconstrained case of this relatively smooth optimization

Table E1.: The results of Algorithm 3, for problem (E1), with different values of m and n .

$\frac{1}{\varepsilon}$	$n = 100, m = 10$		$n = 200, m = 10$	
	Iterations	Time (sec.)	Iterations	Time (sec.)
2	2486	12.603	5271	38.381
4	4792	23.380	10084	73.649
8	9380	43.940	19589	140.668
16	18503	86.753	38406	269.261
32	36686	186.048	75610	537.010
64	72990	369.457	149761	1066.677
	$n = 300, m = 10$		$n = 400, m = 10$	
	Iterations	Time (sec.)	Iterations	Time (sec.)
2	7921	79.872	11076	134.754
4	15581	145.664	22027	264.115
8	31082	318.656	43917	527.536
16	62085	612.985	87686	1041.121
32	124099	1258.532	175198	2098.559
64	248135	2424.470	350201	4189.408
	$n = 100, m = 50$		$n = 200, m = 50$	
	Iterations	Time (sec.)	Iterations	Time (sec.)
2	2491	38.386	5472	139.710
4	4813	72.719	10767	273.139
8	9456	142.670	21341	551.234
16	18803	282.562	42465	1077.094
32	37630	569.258	84703	2256.059
64	75438	1140.276	169159	4388.328

problem:

$$\min_x f(x) := \frac{1}{4}\|Ax - b\|_4^4 + \frac{1}{2}\|Cx - d\|_2^2, \quad (\text{E2})$$

where $A, C \in \mathbb{R}^{n \times n}$ and $b, d \in \mathbb{R}^n$. f is L -smooth relative to $h(x) := \frac{1}{4}\|x\|_2^4 + \frac{1}{2}\|x\|_2^2$ with constant

$$L = 3\|A\|^4 + 6\|A\|^3\|b\|_2 + 3\|A\|^2\|b\|_2^2 + \|C\|^2.$$

In Figure E1 we compare the performance of Gradient Method for relatively smooth problems [46] and Adaptive Gradient Method proposed in this paper. Components of A , C , b , and d are drawn from the uniform distribution $\mathcal{U}(0.95, 1.05)$. The horizontal axis is the number of solved auxiliary problems (11), and therefore the plot is not strictly decreasing. The graph shows that adaptive choice of local Lipschitz constant improves the convergence of the gradient method.

Appendix F. Numerical results for D-optimal design problem

Let us consider the D-optimal design problem:

$$\min_x f(x) := -\ln \det(HXH^\top) \quad \text{s.t. } \langle e, x \rangle = 1, \quad x \geq 0, \quad (\text{F1})$$




Figure E1.: The results of Adaptive GM (Algorithm 1) for problem (E2) with $n = 100$.

where $x \in \mathbb{R}^n$, $X = \text{Diag}(x)$, $H \in \mathbb{R}^{m \times n}$, $e = (1, 1, \dots, 1)^\top$. f is 1-smooth relative to $h(x) := -\sum_{j=1}^n \ln x_j$.

Note that simplicial constraints of the problem are satisfied for the solutions supplied by the considered methods (Algorithm 1) due to the use of a special procedure described in [46] for solving the auxiliary problem (11), which is based on the form of the function h and the structure of constraints of this particular problem.

In this experiment, we consider several cases for different L step sizes in the adaptive choice procedure in Algorithm 1. Instead of iteration over values of local Lipschitz constant with fixed step $\zeta = 2$:

$$L_{k+1} = 2^{i_k} L_k$$

we introduce $\zeta > 1$ as a hyperparameter of the algorithm:

$$L_{k+1} = \zeta^{i_k} L_k,$$

In Figure F1 we compare the performance of the Gradient Method for relatively smooth problems [46] and Adaptive Gradient Method proposed in this paper with different L stepsizes ζ . Part of diagonal components of H are drawn from the uniform distribution $\mathcal{U}(0, 1)$ and the other part from the $\mathcal{U}(0, 200)$, off-diagonal elements are sparsely drawn from $\mathcal{U}(0, 1)$ (proportion of nonzero elements is 0.05). The graph shows that the adaptive choice of the local Lipschitz constant improves the convergence of the gradient method, and the convergence rate can be controlled by the ζ hyperparameter tuning. The vertical axis of the graph measures $f(x^N) - f^*$ in logarithmic scale with f^* estimated as $f^* \geq f(x^N) - R$ where R is defined by (12).

Appendix G. Numerical results for resource sharing problem

In this section, we report the results of a series of numerical experiments for the *resource sharing problem* (RS-Problem) [4]. This problem can be briefly described as follows.

Consider a set of n resources $i \in I = \{1, 2, \dots, n\}$ serving a stream of demands that arrive at a rate of r per unit of time. If the load on the i -th resource is x_i , the expected




Figure F1.: The results of Adaptive GM (Algorithm 1) for problem (F1) with $m = 100$, $n = 200$.

service time in the standard Kleinrock model⁵ is given by the following loss function

$$l_i(x) = \frac{1}{\alpha_i - x_i},$$

where α_i denotes the capacity of the resource. The set of feasible resource allocations is $Q = \{(x_1, \dots, x_n) : 0 \leq x_i < \alpha_i, x_1 + \dots + x_n = R\}$. If we set the operator $g : Q \rightarrow \mathbb{R}^n$, such that $g(x) = (l_1(x_1), \dots, l_n(x_n))$. The point x^* is an equilibrium allocation if and only if it is a solution of the corresponding variational inequality problem for the operator g .

Let $d(x) = \sum_{i=1}^n \frac{1}{1-x_i}$, the prox set-up given on the feasible set Q , then the associated Bregman divergence is

$$V[y](x) = \sum_{i=1}^n \frac{(x_i - y_i)^2}{(1 - x_i)(1 - y_i)^2}$$

By this setting, the operator g is $\frac{1}{\sqrt{2}}$ -Bregman continuous [4] and relatively $\frac{1}{2}$ -smooth relative to d (see Examples 4.3 and 4.6).

For the experiments, in the operator g , we take $\alpha_i = \frac{\sqrt{3}}{2}$, for all $i = 1, \dots, n$ and $R = \frac{n}{2}$. We run Algorithm 3, for different values of n and ε . The starting point $(\frac{1}{2}, \dots, \frac{1}{2}) \in \mathbb{R}^n$, for this, in order to the stopping criterion (57), we have the following estimate $\max_{x \in Q} V[x^0](x) \leq 4n$.

We compare the performance of Algorithm 3, for $n = 100, 250, 500$ and $\varepsilon \in \{\frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \frac{1}{32}, \frac{1}{64}\}$, with three different approaches for step-size selection, the first one is a non-adaptive variant with fixed step-size $L = 0.5$, the second is the proposed procedure in this paper, i.e. with $L_{k+1} = 2^{i_k-1} L_k$, the third one is the procedure with $L_{k+1} = 2^{i_k} L_k$, i.e. non-increasing stepsize. The results of the comparison are presented in Tables G1.

As shown by the results of experiments, in Table G1, the required number of steps and required time to achieve the necessary stopping criterion increases if the stepsize

⁵see: L. Kleinrock: Queueing Systems, volume 1: Theory. John Wiley & Sons, New York, NY, 1975.

is not allowed to increase as in Algorithm 1 in [4]. Moreover, from the results listed in Table G1, we can see that the performance of the algorithm, in this case, is better than the non-adaptive version with the fixed stepsize.

Table G1.: The results for RS-Problem, with different values of n and ε .

$\frac{1}{\varepsilon}$	Non-Adaptive Algorithm 3, with $L = 0.5$.		Algorithm 3, with $L_{k+1} = 2^{i_k-1}L_k$.		Algorithm 3, with $L_{k+1} = 2^{i_k}L_k$.	
	Iterations	Time (sec.)	Iterations	Time (sec.)	Iterations	Time (sec.)
$n = 100$						
2	400	11.67684	5	0.14099	40	1.13174
4	800	20.13218	6	0.15329	80	2.31981
8	1600	40.18136	7	0.18168	160	4.55859
16	3200	89.0930	8	0.20135	320	11.27592
32	> 4000	> 100	9	0.23584	640	16.20664
64	> 4000	> 100	10	0.25457	1280	32.53708
$n = 250$						
2	1000	164.46193	6	1.0216	100	16.49735
4	2000	326.25757	7	1.3804	200	31.96703
8	> 3400	> 550	8	1.32271	400	63.91887
16	> 3400	> 550	9	1.52729	800	135.69973
32	> 3400	> 550	10	1.686	1600	264.14918
64	> 3400	> 550	11	1.82691	3200	525.27188
$n = 500$						
2	> 600	> 500	7	5.78455	200	167.49536
4	> 600	> 500	8	6.60655	400	331.99859
8	> 600	> 500	9	7.34555	> 554	> 500
16	> 600	> 500	10	8.38423	> 554	> 500
32	> 600	> 500	11	9.3096	> 554	> 500
64	> 600	> 500	12	10.15076	> 554	> 500

Since the running time of the non-adaptive variant with fixed step-size of Algorithm 3 and the variant with non-increasing stepsize is big (as we saw from the listed results in Table G1) and for more illustration of the results of the proposed Algorithm 3 (with $L_{k+1} = 2^{i_k-1}L_k$) and to show the slope of the lines, that illustrate the running time and the number of iterations as a function of ε , we run only proposed Algorithm 3 for $n = 100, 250, 500, 1000, 1500, 2000$ with $\varepsilon \in \{\frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \frac{1}{32}, \frac{1}{64}, \text{ and } 5 \times 10^{-i}, 10^{-i} : i = 3, 4, 5, 6\}$. (see Fig. G1 and Fig. G2).

As we see, the slope of the lines is much smaller than what is predicted by the theory, and the adaptivity of the method allows to accelerate the convergence in practice.




Figure G1.: The results of Algorithm 3. The running time as a function of ε , for RS-Problem, with different values of n and ε . The pictures on the left and the right show the results with $\varepsilon \in \{\frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \frac{1}{32}, \frac{1}{64}\}$. The picture on the middle shows the results with $\varepsilon \in \{5 \times 10^{-i}, 10^{-i} : i = 3, 4, 5, 6\}$.




Figure G2.: The results of Algorithm 3. The number of iterations as a function of ε , for RS-Problem, with different values of n and ε . The pictures on the left and the right show the results with $\varepsilon \in \{\frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \frac{1}{32}, \frac{1}{64}\}$. The picture on the middle shows the results with $\varepsilon \in \{5 \times 10^{-i}, 10^{-i} : i = 3, 4, 5, 6\}$.