

A note on the parameterized black hole quasinormal ringdown formalism

Masashi Kimura¹

¹*Department of Physics, Rikkyo University, Tokyo 171-8501, Japan*

(Dated: February 4, 2020)

Abstract

The parameterized black hole quasinormal ringdown formalism is useful to compute quasinormal mode (QNM) frequencies if a master equation for the gravitational perturbation around a black hole has a small deviation from the Regge-Wheeler or Zerilli equation. In this formalism, the deviation of QNM frequency from general relativity can be calculated by small deviation parameters and model independent coefficients. In this paper, we derive recursion relations for the model independent coefficients. Using these relations, the higher order coefficients are written only by the lower order coefficients. Thus, we only need the lower order coefficients when we numerically compute the model independent coefficients.

PACS numbers: 04.50.-h,04.70.Bw

I. INTRODUCTION

The master equations for the gravitational perturbation around the Schwarzschild black holes are known as the Regge-Wheeler or Zerilli equation [1, 2]. If we consider a gravity theory whose deviation from general relativity (GR) is very small, the master equations for the gravitational perturbation around spherically symmetric black holes sometimes have only small deviations from the Regge-Wheeler or Zerilli equation in the form:

$$f \frac{d}{dr} \left(f \frac{d}{dr} \Phi \right) + (\omega^2 - f(V^{(0)} + \delta V)) \Phi = 0, \quad (1)$$

with

$$f = 1 - \frac{r_H}{r}. \quad (2)$$

The back ground effective potential $V^{(0)}$ is

$$V^{(0)} = V_+ = \frac{9\lambda r_H^2 r + 3\lambda^2 r_H r^2 + \lambda^2(\lambda + 2)r^3 + 9r_H^3}{r^3(\lambda r + 3r_H)^2}, \quad (3)$$

where $\lambda = \ell^2 + \ell - 2$, for the even parity perturbation, and

$$V^{(0)} = V_- = \frac{\ell(\ell + 1)}{r^2} - \frac{3r_H}{r^3}, \quad (4)$$

for the odd parity perturbation. δV denotes the small deviation of the effective potential from the GR case. In [3], the form of δV is assumed to be

$$\delta V = \delta V_{\pm} = \frac{1}{r_H^2} \sum_{j=0}^{\infty} \alpha_j^{\pm} \left(\frac{r_H}{r} \right)^j, \quad (5)$$

where α_j^{\pm} are small parameters which depend on the details of gravity theories or physical situations. Note that many models are included in this parameterization, *e.g.*, [3–6]. The parameterized black hole quasinormal ringdown formalism [3] is useful to compute quasinormal mode (QNM) frequencies of the above systems Eqs. (1)-(5). At the first order of the small parameters α_j^{\pm} , the QNM frequency becomes

$$\begin{aligned} \omega_{\text{QNM}} &= \omega_0 + \delta\omega \\ &= \omega_0^{\pm} + \sum_{j=0}^{\infty} \alpha_j^{\pm} e_j^{\pm}, \end{aligned} \quad (6)$$

where $\omega_0 = \omega_0^\pm$ are the QNM frequencies for the GR case, *i.e.*, for the Schwarzschild black holes, and $\delta\omega = \sum_{j=0}^{\infty} \alpha_j^\pm e_j^\pm$ denote the deviation from the GR case. The coefficients e_j^\pm are constants which do not depend on α_j^\pm . In this sense, e_j^\pm can be considered as the model independent coefficients. In the previous work [3], the coefficients e_j^\pm were calculated numerically, but to obtain them in a high accuracy was technically not easy, especially for large j . In this paper, we derive recursion relations among e_j^\pm with different j . Using these relations, we can write e_j^\pm only from low j coefficients.

II. AMBIGUITY OF THE EFFECTIVE POTENTIAL AT THE FIRST ORDER OF SMALL PARAMETERS

Introducing a new master variable $\tilde{\Phi}$ as

$$\Phi = (1 + \epsilon X)\tilde{\Phi} + \epsilon Y f \frac{d\tilde{\Phi}}{dr} \quad (7)$$

where ϵ is a small parameter, which is assumed to be same order as α_j^\pm , and X, Y are functions of r . If we choose X as

$$X = c_1 - \frac{f}{2} \frac{dY}{dr}, \quad (8)$$

with a constant c_1 , Eq. (1) becomes

$$f \frac{d}{dr} \left(f \frac{d}{dr} \tilde{\Phi} \right) + (\omega^2 - f(V^{(0)} + \delta V + \delta W))\tilde{\Phi} = 0, \quad (9)$$

with

$$\delta W = \epsilon \left[-Y \frac{dV^{(0)}}{dr} + 2(\omega_0^2 - V^{(0)}) \frac{dY}{dr} + \frac{1}{2} \frac{d}{dr} \left(f \frac{d}{dr} \left(f \frac{dY}{dr} \right) \right) \right], \quad (10)$$

at the first order of ϵ . In the above derivation, we used the relation $\tilde{\Phi} = \Phi + \mathcal{O}(\epsilon)$ and that Φ satisfies Eq. (1). Unless the choice of X and Y do not change the QNM boundary conditions between Φ and $\tilde{\Phi}$, δW can be considered as an ambiguity of the effective potential. (see also Appendix. A for general discussion of the ambiguity of the effective potential for master equations with Schrödinger form.)

III. RECURSION RELATION FOR THE MODEL INDEPENDENT COEFFICIENTS

In this section, we derive recursion relations among e_j^\pm with different j . Let us consider the GR case, *i.e.*, $\delta V = 0$, where the master equation does not contain any correction term,

$$f \frac{d}{dr} \left(f \frac{d}{dr} \Phi \right) + (\omega^2 - fV^{(0)})\Phi = 0. \quad (11)$$

From the discussion in the previous section, this equation can be rewritten in the form of Eq. (9) with $\delta V = 0$.

A. Odd parity case

For the odd parity case, the back ground effective potential $V^{(0)}$ becomes V_- in Eq. (4). Setting the function Y as

$$Y = y_j \left(\frac{r_H}{r} \right)^j, \quad (12)$$

with an integer j and a constant y_j , δW in Eq. (10) becomes

$$\begin{aligned} \delta W = \epsilon y_j \left[-\frac{2j(\omega_0^-)^2}{r_H} \left(\frac{r_H}{r} \right)^{j+1} - \frac{(1+j)(-4\ell(\ell+1) + j(j+2))}{2r_H^3} \left(\frac{r_H}{r} \right)^{j+3} \right. \\ \left. + \frac{(2j+3)(-6 - 2\ell(\ell+1) + j(j+3))}{2r_H^3} \left(\frac{r_H}{r} \right)^{j+4} - \frac{(j-2)(j+2)(j+6)}{2r_H^3} \left(\frac{r_H}{r} \right)^{j+5} \right]. \end{aligned} \quad (13)$$

We can read α_j^- as

$$\alpha_{j+1}^- = -\frac{2j(\omega_0^-)^2}{r_H} \epsilon y_j, \quad (14)$$

$$\alpha_{j+3}^- = -\frac{(1+j)(-4\ell(\ell+1) + j(j+2))}{2r_H^3} \epsilon y_j, \quad (15)$$

$$\alpha_{j+4}^- = \frac{(2j+3)(-6 - 2\ell(\ell+1) + j(j+3))}{2r_H^3} \epsilon y_j, \quad (16)$$

$$\alpha_{j+5}^- = -\frac{(j-2)(j+2)(j+6)}{2r_H^3} \epsilon y_j, \quad (17)$$

and the QNM frequency at the first order becomes

$$\omega_{\text{QNM}} = \omega_0^- + \alpha_{j+1}^- e_{j+1}^- + \alpha_{j+3}^- e_{j+3}^- + \alpha_{j+4}^- e_{j+4}^- + \alpha_{j+5}^- e_{j+5}^-. \quad (18)$$

However, because now we are studying the system with Eq. (11), the QNM frequency Eq. (18) should be same as the GR case, *i.e.*, $\omega_{\text{QNM}} = \omega_0^-$. Thus, the recursion relation

$$\alpha_{j+1}^- e_{j+1}^- + \alpha_{j+1}^- e_{j+3}^- + \alpha_{j+1}^- e_{j+4}^- + \alpha_{j+1}^- e_{j+5}^- = 0 \quad (19)$$

should hold. Eq. (19) can be explicitly written as

$$\begin{aligned} & -4jr_H^2(\omega_0^-)^2 e_{j+1}^- - (j+1)(-4\ell(\ell+1) + j(j+2))e_{j+3}^- \\ & + (2j+3)(-6 - 2\ell(\ell+1) + j(j+3))e_{j+4}^- - (j-2)(j+2)(j+6)e_{j+5}^- = 0. \end{aligned} \quad (20)$$

We should note that while $Y \sim r$ at $r \rightarrow \infty$ for $j = -1$, the QNM boundary condition for $\tilde{\Phi}$ still corresponds to the QNM boundary condition for the original master variable Φ . Thus, we can put $j \geq -1$ to the above recursion relation, and we conclude that all e_j^- with $j \geq 3$ can be written by e_0^- , e_2^- and e_7^- .¹ For example, the relations

$$e_3^- = \frac{2(\ell-1)\ell(\ell+1)(\ell+2)(\ell^2+\ell-1)r_H^2(\omega_0^-)^2}{(\ell-1)^2\ell^2(\ell+1)^2(\ell+2)^2+36r_H^2(\omega_0^-)^2}e_0^- - \frac{12(\ell^2+\ell-2)r_H^2(\omega_0^-)^2}{(\ell-1)^2\ell^2(\ell+1)^2(\ell+2)^2+36r_H^2(\omega_0^-)^2}e_2^-, \quad (21)$$

$$e_4^- = -\frac{4}{15}r_H^2(\omega_0^-)^2e_0^- + \frac{2}{15}(4+\ell+\ell^2)e_3^-, \quad (22)$$

$$e_5^- = -\frac{1}{15}(3+\ell+\ell^2)r_H^2(\omega_0^-)^2e_0^- + \frac{1}{30}(12+\ell(\ell+1)(2+\ell+\ell^2))e_3^-, \quad (23)$$

$$e_6^- = -\frac{2}{35}(3+\ell+\ell^2)r_H^2(\omega_0^-)^2e_0^- + \frac{\ell(\ell+1)(\ell^2(\ell+1)^2+8)-4(6+5r_H^2(\omega_0^-)^2)}{35(\ell^2+\ell-2)}e_3^-. \quad (24)$$

...

hold.² We report that e_j^- with $j \geq 3$ evaluated from the numerical data of e_0^- , e_2^- , e_7^- in [3] agree with that in the previous work.

B. Even parity case

For the even parity case, the back ground effective potential $V^{(0)}$ becomes V_+ in Eq. (3). Setting the function Y as

$$Y = y_j \frac{(\lambda r + 3r_H)^3}{r_H^3} \left(\frac{r_H}{r}\right)^j, \quad (25)$$

¹ Note that e_7^- cannot be written by e_0^- , e_2^- because the coefficient of e_7^- in Eq. (20) with $j = 2$ vanishes.

² One may wonder how to express e_3^- by e_0^- and e_2^- . Eq. (20) with $j = -1, 0, 1, 2$ give four equations which contain e_0^- , e_2^- , e_3^- , e_4^- , e_5^- , e_6^- . Solving them with respect to e_3^- , e_4^- , e_5^- , e_6^- , we can write e_3^- by e_0^- and e_2^- .

with an integer j and a constant y_j , δW becomes

$$\begin{aligned}
\delta W = \frac{\epsilon y_j}{r_H^3} & \left[-2\lambda^3(j-3)r_H^2(\omega_0^+)^2 \left(\frac{r_H}{r}\right)^{j-2} - 18\lambda^2(j-2)r_H^2(\omega_0^+)^2 \left(\frac{r_H}{r}\right)^{j-1} \right. \\
& + \left(\frac{1}{2}\lambda^3(j-2)(4\lambda - (j-4)j+5) - 54\lambda(j-1)r_H^2(\omega_0^+)^2 \right) \left(\frac{r_H}{r}\right)^j \\
& + \left(\frac{1}{2}\lambda^2(6(\lambda-1)\lambda + (2\lambda-9)j^3 - 9(\lambda-3)j^2 + ((25-4\lambda)\lambda+6)j) - 54jr_H^2(\omega_0^+)^2 \right) \left(\frac{r_H}{r}\right)^{j+1} \\
& + \frac{\lambda(3(\lambda-4)\lambda - ((\lambda-18)\lambda+27)j^3 + 3(\lambda-9)\lambda j^2 + ((21-23\lambda)\lambda+27)j)}{2} \left(\frac{r_H}{r}\right)^{j+2} \\
& + \frac{9(\lambda^2(-(j^3+4j+2)) + \lambda(j+1)(6j^2+3j+4) - 3n(j+1)(j+2))}{2} \left(\frac{r_H}{r}\right)^{j+3} \\
& + \frac{9(3(j+1)(j+2)(2j+3) - \lambda(j(3j(j+3)+13)+9))}{2} \left(\frac{r_H}{r}\right)^{j+4} \\
& \left. - \frac{27(j+2)^3}{2} \left(\frac{r_H}{r}\right)^{j+5} \right]. \tag{26}
\end{aligned}$$

We can read $\alpha_{j-2}, \alpha_{j-1}, \dots, \alpha_{j+5}$ by comparing with $\delta W = r_H^{-2} \sum_{j=0}^{\infty} \alpha_j^{\pm} (r_H/r)^j$, and the QNM frequency becomes

$$\omega_{\text{QNM}} = \omega_0^+ + \sum_{k=j-2}^{j+5} \alpha_k^+ e_k^+. \tag{27}$$

Similar to the odd parity case, because the QNM frequency (27) should be same as the GR case, the recursion relation

$$\sum_{k=j-2}^{j+5} \alpha_k^+ e_k^+ = 0 \tag{28}$$

should hold. Using this, e_j^+ with $j \geq 7$ can be written by $e_0^+, e_1^+, \dots, e_6^+$.

C. Test scalar and vector fields

We can study the test scalar and vector fields similar to the above discussions. In that cases, the back ground effective potential becomes

$$V^{(0)} = V_s = \frac{\ell(\ell+1)}{r^2} - (1-s^2) \frac{r_H}{r^3}, \tag{29}$$

where $s = 0$ for the test massless scalar fields, $s = 1$ for the test massless vector fields. Note that $s = 2$ corresponds to the Regge-Wheeler potential V^- . We parameterize δV as

$$\delta V = \delta V_s = \frac{1}{r_H^2} \sum_{j=0}^{\infty} \beta_j^s \left(\frac{r_H}{r}\right)^j, \tag{30}$$

with the small parameters β_s , then the QNM frequency becomes

$$\omega_{\text{QNM}} = \omega_0^s + \sum_{j=0}^{\infty} \beta_j^s d_j^s, \quad (31)$$

where the coefficients d_j^s are constants which do not depend on β_j^s . Setting

$$Y = y_j \left(\frac{r_H}{r} \right)^j, \quad (32)$$

with an integer j and a constant y_j , after the same calculation as above discussions, we obtain a relation among d_j^s

$$\begin{aligned} & -4jr_H^2(\omega_0^s)^2 d_{j+1}^s - (j+1)(j-2\ell)(j+2\ell+2)d_{j+3}^s \\ & + (2j+3)(2+j(j+3) - 2\ell(\ell+1) - 2s^2)d_{j+4}^s - (j+2)(j+2-2s)(j+2+2s)d_{j+5}^s = 0. \end{aligned} \quad (33)$$

Using this relation, d_j^s with $j \geq 3$ can be written by d_0^s, d_2^s and d_7^s .

IV. REDUCTION OF THE EFFECTIVE POTENTIAL TO LOWER ORDER

We showed that e_j^\pm with high j can be written by that with low j . In a similar way, we can remove the higher order terms of r_H/r in δV by choosing the function Y appropriately. For simplicity, we only focus on the odd parity case in this section. Note that the similar discussion holds if we consider the cases of even parity, vector fields or scalar field.

Let us assume that the highest order term in δV is

$$\delta V^{\text{highest}} = \frac{\alpha_{n+5}^-}{r_H^2} \left(\frac{r_H}{r} \right)^{n+5}. \quad (34)$$

If we choose

$$Y = y_n \left(\frac{r_H}{r} \right)^n, \quad (35)$$

with

$$y_n = \frac{2r_H \alpha_{n+5}^-}{(n-2)(n+2)(n+6)}, \quad (36)$$

we can remove this term, and the new effective potential contains at most $O((r_H/r)^{n+4})$ terms. Repeating this process, we can write the effective potential δV in terms of $(r_H/r)^j$

with $j = 0, 1, 2, 7$.³ This implies that we only need to consider the case

$$\delta V = \frac{1}{r_H^2} \left[\alpha_0^- + \alpha_1^- \left(\frac{r_H}{r} \right) + \alpha_2^- \left(\frac{r_H}{r} \right)^2 + \alpha_7^- \left(\frac{r_H}{r} \right)^7 \right], \quad (37)$$

instead of Eq. (5). We should note that the coefficients $\alpha_0^-, \alpha_1^-, \alpha_2^-, \alpha_7^-$ become complicated functions of ω_0^- , in general.

V. SUMMARY AND DISCUSSION

In the parameterized black hole quasinormal ringdown formalism [3], the deviation of QNM frequency from GR is given by $\sum_j \alpha_j^\pm e_j^\pm$. While the parameters α_j^\pm , which depend on physical situations, can be read from the master equation, we need to numerically calculate the model independent coefficients e_j^\pm in advance [3]. Especially, it was technically not easy task to compute e_j^\pm with high j in a high accuracy.

In this paper, we derived recursion relations among e_j^\pm with different j . Using these relations, e_j^\pm with high j can be written by that with low j . This can efficiently reduce the calculation cost for computing e_j^\pm . The recursion relations can also be used to estimate the numerical calculation error for e_j^\pm . For example, if we numerically calculate e_0^-, e_2^-, e_3^- , we can estimate the error from $|1 - \text{LHS}/\text{RHS}|$, where LHS and RHS denote the left and right hand side of Eq. (21), respectively.

In a similar way, we also showed that δV can be rewritten in terms of $(r_H/r)^j$ with only low j using the ambiguity of the effective potential. It is also interesting to extend the present discussion to the coupled cases and the higher order cases [6], but we leave them for future work.

Acknowledgments

MK would like to thank Yasuyuki Hatsuda for useful discussions and comments on the paper.

³ Note that we cannot remove $(r_H/r)^7$ term because Eq. (36) diverges in that case. This is similar to why e_7^- cannot be written by e_0^- and e_2^- discussed in footnote. 1.

Appendix A: general discussion of the ambiguity of the effective potential

Let us consider the master equation with Schrödinger form

$$\frac{d^2}{dx^2}\Phi + (\omega^2 - V)\Phi = 0, \quad (\text{A1})$$

where V is the effective potential. For the case in Eq. (1), the relation between x and r is given by $d/dx = fd/dr$. In this section, we discuss the general differential transformations of the master variable which keep the master equation to be Schrödinger form. Defining a new master variable χ as⁴

$$\chi = A\Phi + B\frac{d\Phi}{dx}, \quad (\text{A2})$$

with

$$-B^2V' + B(2(\omega^2 - V)B' - A'') + A(2A' + B'') = 0, \quad (\text{A3})$$

χ satisfies the equation

$$\chi'' + (\omega^2 - (V + \delta W))\chi = 0, \quad (\text{A4})$$

with

$$\delta W = \frac{2A'}{B} + \frac{B''}{B}, \quad (\text{A5})$$

where A, B are functions of x and $'$ denotes derivative with respect to x . Note that Eq. (A3) is the condition such that the coefficient of χ' vanishes in the master equation. We can regard the term δW as an ambiguity of the effective potential. Eq. (A3) can be integrated as

$$(-\omega^2 + V)B^2 - A^2 + BA' - AB' = c_2, \quad (\text{A6})$$

with a constant c_2 . Defining

$$S := \frac{A}{B} + \frac{B'}{B}, \quad (\text{A7})$$

Eq. (A4) can be written as

$$\chi'' + \left(-\frac{c_2}{B^2} - S^2 - S'\right)\chi = 0. \quad (\text{A8})$$

⁴ Because Φ satisfies the second order differential equation Eq. (A1), the second or more derivatives of Φ can be written in terms of Φ and Φ' . For this reason, we only need to consider differential transformations which only contain at most first derivative of Φ .

1. $A = 1 - \epsilon X$ and $B = -\epsilon Y$ case

For $A = 1 - \epsilon X$ and $B = -\epsilon Y$ with the functions X, Y and a small parameter ϵ , because $\chi = \Phi$ at $\mathcal{O}(\epsilon^0)$, the relation

$$\Phi = (1 + \epsilon X)\chi + \epsilon Y\chi' + \mathcal{O}(\epsilon^2) \quad (\text{A9})$$

holds. From Eq. (A6), we obtain $X = c_3 - Y'/2 + \mathcal{O}(\epsilon)$ where $c_3 = (1 - \sqrt{-c_2})/\epsilon$. Then, δW becomes

$$\delta W = \frac{BV' - 2(\omega^2 - V)B' + A''}{A} = \epsilon \left[-YV' + 2(\omega^2 - V)Y' + \frac{1}{2}Y''' \right] + \mathcal{O}(\epsilon^2), \quad (\text{A10})$$

where we used Eq. (A3) in the first equality. This result is same as Eq. (10). Note that $d/dx = fd/dr$ and δW in this section corresponds to that in Sec. II multiplied by f .

2. $B = 1, c_2 = -\omega^2$ case

If we set $B = 1$ and $c_2 = -\omega^2$, Eq. (A6) becomes

$$V = S^2 - S' \quad (\text{A11})$$

and the master equation Eq. (A8) becomes

$$\chi'' + (\omega^2 - S^2 - S')\chi = 0. \quad (\text{A12})$$

This is nothing but the superpartner of the original system Eq. (A1).

-
- [1] T. Regge and J. A. Wheeler, Phys. Rev. **108**, 1063 (1957). doi:10.1103/PhysRev.108.1063
 - [2] F. J. Zerilli, Phys. Rev. Lett. **24**, 737 (1970). doi:10.1103/PhysRevLett.24.737
 - [3] V. Cardoso, M. Kimura, A. Maselli, E. Berti, C. F. B. Macedo and R. McManus, Phys. Rev. D **99**, no. 10, 104077 (2019) doi:10.1103/PhysRevD.99.104077 [arXiv:1901.01265 [gr-qc]].
 - [4] V. Cardoso, M. Kimura, A. Maselli and L. Senatore, Phys. Rev. Lett. **121**, no. 25, 251105 (2018) doi:10.1103/PhysRevLett.121.251105 [arXiv:1808.08962 [gr-qc]].
 - [5] O. J. Tattersall, arXiv:1911.07593 [gr-qc].
 - [6] R. McManus, E. Berti, C. F. B. Macedo, M. Kimura, A. Maselli and V. Cardoso, Phys. Rev. D **100**, no. 4, 044061 (2019) doi:10.1103/PhysRevD.100.044061 [arXiv:1906.05155 [gr-qc]].