arXiv:2001.09660v4 [cs.IT] 17 Feb 2020

The α-divergences associated with a pair of strictly comparable quasi-arithmetic means

Frank Nielsen*
Sony Computer Science Laboratories Inc.
Tokyo, Japan

Abstract

We generalize the family of α-divergences using a pair of strictly comparable weighted means. In particular, we obtain the 1-divergence in the limit case $\alpha \to 1$ (a generalization of the Kullback-Leibler divergence) and the 0-divergence in the limit case $\alpha \to 0$ (a generalization of the reverse Kullback-Leibler divergence). We state the condition for a pair of quasi-arithmetic means to be strictly comparable, and report the formula for the quasi-arithmetic α-divergences and its subfamily of bipower homogeneous α-divergences which belong to the Csisár's f-divergences. Finally, we show that these generalized quasi-arithmetic 1-divergences and 0-divergences can be decomposed as the sum of generalized cross-entropies minus entropies, and rewritten as conformal Bregman divergences using monotone embeddings.

Keywords: Kullback-Leibler divergence, α-divergences, comparable means, weighted quasi-arithmetic means, α-geometry, homogeneous divergences, conformal divergences, geometric divergence, monotone embeddings, conformal flattening.

1 Introduction

1.1 Statistical divergences

Consider a measurable space $(\mathcal{X}, \mathcal{F})$ (where $\mathcal{F}$ denotes the σ-algebra and $\mathcal{X}$ the sample space) equipped with a positive measure μ (e.g., usually the Lebesgue measure or the counting measure). The notion of *statistical dissimilarity* [4] $D[P : Q] = D_\mu[p_\mu : q_\mu]$ between two arbitrary probability measures with Radon-Nikodym (RN) densities $p_\mu = \frac{\mathrm{d}P}{\mathrm{d}\mu}$ and $q_\mu = \frac{\mathrm{d}Q}{\mathrm{d}\mu}$ with respect to μ is at the core of many algorithms in signal processing, information theory, information fusion, and machine learning among others. When those statistical dissimilarities are smooth, they are called *divergences* [2] in the literature. The most renown statistical divergence rooted in information theory [9] is the *Kullback-Leibler divergence* (KLD):

$$\mathrm{KL}_\mu[p_\mu : q_\mu] := \int_{\mathcal{X}} p_\mu(x) \log \frac{p_\mu(x)}{q_\mu(x)} \mathrm{d}\mu(x). \tag{1}$$

Since the KLD is independent of the reference measure μ, i.e., $\mathrm{KL}_\mu[p_\mu : q_\mu] = \mathrm{KL}_\nu[p_\nu : q_\nu]$ for $p_\mu = \frac{\mathrm{d}P}{\mathrm{d}\mu}$ and $q_\mu = \frac{\mathrm{d}Q}{\mathrm{d}\mu}$, and $p_\nu = \frac{\mathrm{d}P}{\mathrm{d}\nu}$ and $q_\nu = \frac{\mathrm{d}Q}{\mathrm{d}\nu}$ the RN derivatives with respect to another

*E-mail: Frank.Nielsen@acm.org

positive measure ν, we write concisely in the remainder:

$$\mathrm{KL}[p:q] = \int p \log \frac{p}{q} \mathrm{d}\mu, \tag{2}$$

instead of $\mathrm{KL}_\mu[p_\mu : q_\mu]$.

The KLD belongs to a parametric family of α-*divergences* [7] $I_\alpha[p:q]$ for $\alpha \in \mathbb{R}$:

$$I_\alpha[p:q] := \begin{cases} \frac{1}{\alpha(1-\alpha)} \left(1 - \int p^\alpha q^{1-\alpha} \mathrm{d}\mu\right), & \alpha \in \mathbb{R}\backslash\{0,1\} \\ I_1[p:q] = \mathrm{KL}[p:q], & \alpha = 1 \\ I_0[p:q] = \mathrm{KL}[q:p], & \alpha = 0 \end{cases} . \tag{3}$$

The α-divergences extended to positive densities (not necessarily normalized) play a central role in information geometry [2]:

$$I_\alpha[p:q] := \begin{cases} \frac{1}{\alpha(1-\alpha)} \int \left(\alpha p + (1-\alpha) q - p^\alpha q^{1-\alpha}\right) \mathrm{d}\mu, & \alpha \in \mathbb{R}\backslash\{0,1\} \\ I_1[p:q] = \mathrm{KL}_e[p:q], & \alpha = 1 \\ I_0[p:q] = \mathrm{KL}_e[q:p], & \alpha = 0 \end{cases} , \tag{4}$$

where KL_e denotes the *extended Kullback-Leibler divergence*:

$$\mathrm{KL}_e[p:q] := \int \left(p \log \frac{p}{q} + q - p\right) \mathrm{d}\mu. \tag{5}$$

The α-divergences are asymmetric for $\alpha \neq 0$ (i.e., $I_\alpha[p:q] \neq I_\alpha[q:p]$ for $\alpha \neq 0$) but exhibit the following *reference duality* [39]:

$$I_\alpha[q:p] = I_{1-\alpha}[p:q] = (I_\alpha)^*[p:q], \tag{6}$$

where we denoted by $D^*[p:q] := D[q:p]$, the *reverse divergence* for an arbitrary divergence D (e.g., $I_\alpha^*[p:q] = I_\alpha[q:p] = I_{1-\alpha}[p:q]$).

The α-divergences belong to the family of Csizár's f-divergences [10] which are defined for a convex function f satisfying by $f(1) = 0$ by:

$$I_f[p:q] := \int p f\left(\frac{q}{p}\right) \mathrm{d}\mu. \tag{7}$$

We have

$$I_\alpha[p:q] = I_{f_\alpha}[p:q], \tag{8}$$

with

$$f_\alpha(u) = \begin{cases} \frac{1}{\alpha(1-\alpha)}(\alpha + (1-\alpha)u - u^{1-\alpha}), & \alpha \in \alpha \in \mathbb{R}\backslash\{0,1\} \\ u - 1 - \log u, & \alpha = 1 \\ 1 - u + u \log u, & \alpha = -1 \end{cases} \tag{9}$$

In information geometry, α-divergences (and more generally f-divergences) are *invariant divergences* [2], and it is customary to rewrite the α-divergences using $\alpha_A = 1 - 2\alpha$ (i.e., $\alpha = \frac{1-\alpha_A}{2}$). Thus the *extended* α_A*-divergence* is defined by

$$\hat{I}_{\alpha_A}[p:q] = \begin{cases} \frac{4}{1-\alpha_A^2}\int\left(\frac{1-\alpha_A}{2}p + \frac{1+\alpha_A}{2}q - p^{\frac{1-\alpha_A}{2}}q^{\frac{1+\alpha_A}{2}}\right)\mathrm{d}\mu, & \alpha\in\mathbb{R}\backslash\{-1,1\} \\ \hat{I}_1[p:q] = \mathrm{KL}_e[p:q], & \alpha = 1 \\ \hat{I}_{-1}[p:q] = \mathrm{KL}_e[q:p], & \alpha = -1 \end{cases}, \quad (10)$$

and the reference duality is expressed by $\hat{I}_{\alpha_A}[q:p] = \hat{I}_{-\alpha_A}[p:q]$.

A statistical divergence $D[\cdot:\cdot]$ when evaluated on densities belonging to a given parametric family $\mathcal{P} = \{p_\theta \ : \ \theta\in\Theta\}$ of densities are equivalent to a corresponding *contrast function* [12]:

$$D_{\mathcal{P}}(\theta_1:\theta_2) := D[p_{\theta_1}:p_{\theta_2}]. \quad (11)$$

Although quite confusing, those contrast functions have also been called recently divergences in the literature [2]. Thus to disambiguate whether the divergence is a statistical divergence or a parameter divergence (i.e., contrast function), we choose to use the brackets for encapsulating arguments in statistical divergences and the parenthesis to encapsulate parameter arguments in divergences which are contrast functions.

A smooth divergence $D(\theta_1:\theta_2)$ induces a dualistic structure in information geometry [2]. For example, the KLD on the family Δ of probability mass functions on a finite alphabet $\mathcal{X}$ with equivalent contrast function a Bregman divergence induces a *dually flat space* [2]. More generally, the α_A-divergences on the probability simplex Δ induce the *α_A-geometry*.

The α-divergences are widely used in information sciences, see [3, 8, 38, 22, 17, 1] just to cite a few applications. The singly-parametric α-divergences have also been generalized to bi-parametric families of divergences like the (α,β)-divergences [2] or the $\alpha\beta$-divergences [37].

In this work, based on the observation that the term $\alpha p + (1-\alpha)q - p^\alpha q^{1-\alpha}$ (in the extended $I_\alpha[p:q]$ divergence for $\alpha\in(0,1)$ of Eq. 4) is a difference between a weighted arithmetic mean $A_{1-\alpha}(p,q) := \alpha p + (1-\alpha)q$ and a weighted geometric mean $G_{1-\alpha}(p,q) := p^\alpha q^{1-\alpha}$, we investigate a generalization of α-divergences with respect to a pair of abstract strictly comparable weighted means [25].

1.2 Divergences and decomposable divergences

A statistical divergence D shall satisfy the following two axioms:

D1. non-negativity. $D[p:q]\geq 0$ for all densities p and q,

D2. identity of indiscernibles. $D[p:q]=0$ if and only if $p=q$ μ-almost everywhere.

These axioms are a subset of the metric axioms since we do not consider the symmetry axiom nor the triangular inequality axiom of metric distances. See [14] for some common examples of *probability metrics* (e.g., total variation or Wasserstein metrics).

A divergence $D[p:q]$ is said *decomposable* [2] when it can be written as an integral of a scalar divergence $d(\cdot,\cdot)$:

$$D[p:q] = \int d(p(x):q(x))\mathrm{d}\mu(x), \quad (12)$$

or $D[p:q] = \int d(p:q)\mathrm{d}\mu$ for short.

The α-divergences are decomposable divergences since we have

$$I_\alpha[p:q] = \int i_\alpha(p(x):q(x))\mathrm{d}\mu \quad (13)$$

with the following *scalar α-divergence*:

$$i_\alpha(a:b) := \begin{cases} \frac{1}{\alpha(1-\alpha)}\left(\alpha a + (1-\alpha)b - a^\alpha b^{1-\alpha}\right), & \alpha \in \mathbb{R}\backslash\{0,1\} \\ i_1(a:b) = a\log\frac{a}{b} + b - a & \alpha = 1 \\ i_0(a:b) = i_1(b:a), & \alpha = 0 \end{cases} \tag{14}$$

1.3 Contributions and paper outline

The outline of the paper and the contributions are summarized as follows:

We define the generic α-divergences in §2 for two families of strictly comparable means (Definition 2). Then section 2.2 reports a closed-form formula (Theorem 3) for the quasi-arithmetic α-divergences induced by two strictly comparable quasi-arithmetic means with monotonically increasing generators f and g such that $f \circ g^{-1}$ is *strictly convex and differentiable*. In §2.3, we study the divergences I_0 and I_1 obtained in the limit cases when $\alpha \to 0$ and $\alpha \to 1$, respectively. We obtain generalized Kullback-Leibler divergences when $\alpha \to 1$ and generalized reverse Kullback-Leibler divergences when $\alpha \to 0$, which can be factorized as generalized cross-entropies minus entropies. In §2.4, we show how to express these generalized I_1-divergences and I_0-divergences as conformal Bregman representational divergences and briefly explain their induced conformally flat statistical manifolds. Section 3 explicits the subfamily of bipower homogeneous α-divergences which belong to the family of Csiszár f-divergences [10]. Finally, Section 4 summarizes the work and present several opportunities for future research directions.

2 The α-divergences based on a pair of means

2.1 The abstract (M, N) α-divergences

The point of departure for generalizing the α-divergences is to rewrite Eq. 4 for $\alpha \in \mathbb{R}\backslash\{0,1\}$ as

$$I_\alpha[p:q] = \frac{1}{\alpha(1-\alpha)} \int \left(A_{1-\alpha}(p:q) - G_{1-\alpha}(p:q)\right) \mathrm{d}\mu, \tag{15}$$

where A_λ and G_λ for $\lambda \in (0,1)$ stands for the weighted arithmetic and geometric means, respectively:

$$\begin{aligned} A_\lambda(x,y) &= (1-\lambda)x + \lambda y, \\ G_\lambda(x,y) &= x^{1-\lambda} y^\lambda. \end{aligned}$$

We choose the convention $A_0(x,y) = x$ and $A_1(x,y) = 1$ so that $\{A_\lambda(x,y)\}_{\lambda\in[0,1]}$ smoothly interpolates between x ($\lambda = 0$) and y ($\lambda = 1$).

In general, a *mean* $M(x,y)$ aggregates two values x and y of an interval I to produce an intermediate quantity which satisfies the *innerness property* [5]:

$$\min\{x,y\} \leq M(x,y) \leq \max\{x,y\}, \quad \forall x,y \in I. \tag{16}$$

A mean is said *strict* if the inequalities of Eq. 16 are strict whenever $x \neq y$. A mean M is said *reflexive* iff $M(x,x) = x$ for all $x \in I$. In the remainder, we consider $I = (0,\infty)$. By using the unique dyadic representation of any real $\lambda \in (0,1)$ (i.e., $\lambda = \sum_{i=1}^\infty \frac{d_i}{2^i}$ with $d_i \in \{0,1\}$, a binary digit) , one can build a *weighted mean* M_λ from any given mean M, see [25] for such a construction.

By analogy to the α-divergences, let us define the (decomposable) (M,N) *α-divergences* for a pair of weighted means $M_{1-\alpha}$ and $N_{1-\alpha}$ for $\alpha \in (0,1)$ as

$$I_\alpha^{M,N}[p:q] := \frac{1}{\alpha(1-\alpha)} \int (M_{1-\alpha}(p:q) - N_{1-\alpha}(p:q)) \,\mathrm{d}\mu. \tag{17}$$

The ordinary α-divergences for $\alpha \in (0,1)$ are recovered as the (A,G) α-divergences:

$$I_\alpha^{A,G}[p:q] = \frac{1}{\alpha(1-\alpha)} \int (A_{1-\alpha}(p:q) - G_{1-\alpha}(p:q)) \,\mathrm{d}\mu = I_{1-\alpha}[p:q] = I_\alpha[q:p] = I_\alpha^*[p:q]. \tag{18}$$

In order to define generalized α-divergences satisfying the axioms D1 and D2 of proper divergences, we need to characterize the class of acceptable means. We give a definition strengthening the notion of comparable means in [25]:

Definition 1 (Strictly comparable weighted means)**.** *A pair (M,N) of means are said* strictly comparable *whenever $M_\lambda(x,y) \leq N_\lambda(x,y)$ for all $x,y \in (0,\infty)$ with equality if and only if $x=y$, and for all $\lambda \in (0,1)$.*

For example, the inequality of the arithmetic and geometric means states that $A(x,y) \geq G(x,y)$: Means A and G are comparable, denoted by $A \geq G$. Furthermore, the arithmetic and geometric weighted means are distinct whenever $x \neq y$: Indeed, consider the equation $(1-\alpha)x+\alpha y = x^{1-\alpha}y^\alpha$ for $x,y>0$ and $x \neq y$. By taking the logarithm on both sides, we get

$$\log\left((1-\alpha)x+\alpha y\right) = (1-\alpha)\log x + \alpha \log y. \tag{19}$$

Since the logarithm is a strictly convex function, the only solution is $x=y$. Thus (A,G) is a pair of strictly comparable weighted means.

For a weighted mean M, define $\bar{M}_\alpha(x,y) := M_{1-\alpha}(x,y)$. We are ready to state the definition of generalized α-divergences:

Definition 2 ((M,N) α-divergences)**.** *The (M,N) α-divergences $I_\alpha^{M,N}[p:q]$ between two positive densities p and q for $\alpha \in (0,1)$ is defined for a pair of strictly comparable weighted means M_α and N_α with $M_\alpha \geq N_\alpha$ by:*

$$\begin{aligned} I_\alpha^{M,N}[p:q] &:= \frac{1}{\alpha(1-\alpha)} \int (M_{1-\alpha}(p:q) - N_{1-\alpha}(p:q)) \,\mathrm{d}\mu, & (20)\\ &= \frac{1}{\alpha(1-\alpha)} \int \left(\bar{M}_\alpha(p:q) - \bar{N}_\alpha(p:q)\right) \mathrm{d}\mu. & (21) \end{aligned}$$

Using $\alpha = \frac{1-\alpha_A}{2}$, we can rewrite this divergence as

$$\begin{aligned} \hat{I}_{\alpha_A}^{M,N}[p:q] &:= \frac{4}{1-\alpha_A^2} \int \left(M_{\frac{1+\alpha_A}{2}}(p:q) - N_{\frac{1+\alpha_A}{2}}(p:q)\right) \mathrm{d}\mu, & (22)\\ &= \frac{4}{1-\alpha_A^2} \int \left(\bar{M}_{\frac{1-\alpha_A}{2}}(p:q) - \bar{N}_{\frac{1-\alpha_A}{2}}(p:q)\right) \mathrm{d}\mu. & (23) \end{aligned}$$

A weighted mean M_α is said *symmetric* iff $M_\alpha(x,y) = M_{1-\alpha}(y,x)$. When both the weighted means M and N are symmetric, we have the following *reference duality* [39]:

$$I_\alpha^{M,N}[p:q] = I_{1-\alpha}^{M,N}[q:p]. \tag{24}$$

We consider symmetric means in the remainder.

In the limit cases of $\alpha \to 0$ or $\alpha \to 1$, we define the 0-*divergence* $I_0^{M,N}[p:q]$ and the 1-*divergence* $I_1^{M,N}[p:q]$, respectively, by

$$I_0^{M,N}[p:q] = \lim_{\alpha \to 0} I_\alpha^{M,N}[p:q], \tag{25}$$

$$I_1^{M,N}[p:q] = \lim_{\alpha \to 1} I_\alpha^{M,N}[p:q] = I_0^{M,N}[q:p], \tag{26}$$

provided that those limits exist.

2.2 The quasi-arithmetic α-divergences

A *quasi-arithmetic mean* (QAM) is defined for a continuous and strictly monotonic function $f : I \subset \mathbb{R} \to J \subset \mathbb{R}$ as:

$$M^f(x,y) := f^{-1}\left(\frac{f(x)+f(y)}{2}\right). \tag{27}$$

Function f is called the generator of the quasi-arithmetic mean. These strict and reflexive quasi-arithmetic means are also called Kolmogorov means [19], Nagumo means [23] or de Finetti means [11], or quasi-linear means [15] in the literature. These means are called quasi-arithmetic means because they can be interpreted as arithmetic means on the arguments $f(x)$ and $f(y)$:

$$f(M^f(x,y)) = \frac{f(x)+f(y)}{2} = A(f(x),f(y)). \tag{28}$$

QAMs are strict, reflexive and symmetric means.

Without loss of generality, we assume strictly increasing functions f instead of monotonic functions since $M^{-f} = M^f$. Indeed, $M^{-f}(x,y) = (-f)^{-1}(-f(M_f(x,y)))$ and $((-f)^{-1} \circ (-f))(u) = u$, the identity function. Notice that the composition $f_1 \circ f_2$ of two strictly monotonic increasing functions f_1 and f_2 is a strictly monotonic increasing function. Furthermore, we consider $I = J = (0,\infty)$ in the remainder since we apply these means on positive densities. Two quasi-arithmetic means M^f and M^g coincide if and only if $f(u) = ag(u) + b$ for some $a > 0$ and $b \in \mathbb{R}$ see [15]. The quasi-arithmetic means were considered in the axiomatization of the entropies by Rényi to define the α-entropies (see Eq. 2.11 of [36]).

By choosing $f_A(u) = u$, $f_G(u) = \log u$ or $f_H(u) = \frac{1}{u}$, we obtain the Pythagorean's arithmetic A, geometric G, and harmonic H means, respectively:

- the *arithmetic mean* (A): $A(x,y) = \frac{x+y}{2} = M^{f_A}(x,y)$,
- the *geometric mean* (G): $G(x,y) = \sqrt{xy} = M^{f_G}(x,y)$, and
- the *harmonic mean* (H): $H(x,y) = \frac{2}{\frac{1}{x}+\frac{1}{y}} = \frac{2xy}{x+y} = M^{f_H}(x,y)$.

More generally, choosing $f_{P_r}(u) = u^r$, we obtain the parametric family of *power means* (also called *Hölder means* [16]):

$$P_r(x,y) = \left(\frac{x^r+y^r}{2}\right)^{\frac{1}{r}} = M^{f_{P_r}}(x,y), \quad r \in \mathbb{R}\backslash\{0\}. \tag{29}$$

In order to get a *smooth family* of power means, we define the geometric mean in the limit case of $r \to 0$:

$$P_0(x,y) = \lim_{r\to 0} P_r(x,y) = G(x,y) = \sqrt{xy}. \tag{30}$$

It is known that the positively homogeneous quasi-arithmetic means, i.e. $M^f(\lambda a, \lambda b) = \lambda M^f(a,b)$ for $\lambda > 0$, coincide exactly with the family of power means. The weighted QAMs are given by

$$M^f_\alpha(p,q) = f^{-1}\left((1-\alpha)f(p) + \alpha f(q))\right) = f^{-1}\left(f(p) + \alpha(f(q) - f(p))\right) = M^f_{1-\alpha}(q,p). \tag{31}$$

The logarithmic mean $L(x,y)$ for $x > 0$ and $y > 0$:

$$L(x,y) = \frac{y-x}{\log y - \log x}$$

is an example of a homogeneous mean (i.e., $L(\lambda x, \lambda y) = \lambda L(x,y)$ for any $\lambda > 0$) that is *not* a QAM. Besides the family of QAMs, there exist many other families of means [5]: For example, let us mention the *Lagrangean means* [18] which intersect with the QAMs only for the arithmetic mean, or a generalization of the QAMs called the the *Bajraktarević means* [35].

Let us strengthen a recent theorem of [21] (Theorem 1, 2010):

Theorem 1 (Strictly comparable weighted QAMs)**.** *The pair (M^f, M^g) of quasi-arithmetic means obtained for two strictly increasing generators is strictly comparable provided that $f \circ g^{-1}$ is strictly convex.*

Proof. Since $f \circ g^{-1}$ is strictly convex, it is convex, and therefore it follows from Theorem 1 of [21] that $M^f_\alpha \geq M^g_\alpha$ for all $\alpha \in [0,1]$. Thus the very nice property of QAMs is that $M^f \geq M^g$ implies that $M^f_\alpha \geq M^g_\alpha$ for *any* $\alpha \in [0,1]$.

Now, let us consider the equation $M^f_\alpha(p,q) = M^g_\alpha(p,q)$ for $p \neq q$:

$$f^{-1}\left((1-\alpha)f(p) + \alpha f(q)\right) = g^{-1}\left((1-\alpha)g(p) + \alpha g(q)\right). \tag{32}$$

Since $f \circ g^{-1}$ is assumed strictly convex, and g is strictly increasing, we have $g(p) \neq g(q)$ for $p \neq q$, and we reach the following contradiction:

$$\begin{aligned}
(1-\alpha)f(p) + \alpha f(q) &= (f \circ g^{-1})\left((1-\alpha)g(p) + \alpha g(q)\right), &(33)\\
&< (1-\alpha)(f \circ g^{-1})(g(p)) + \alpha(f \circ g^{-1})(g(q)), &(34)\\
&< (1-\alpha)f(p) + \alpha f(q). &(35)
\end{aligned}$$

Thus $M^f_\alpha(p,q) \neq M^g_\alpha(p,q)$ for $p \neq q$, and $M^f_\alpha(p,q) = M^g_\alpha(p,q)$ for $p = q$. □

Note that the (A,G) α-divergences (i.e., the ordinary α-divergences) is a proper divergence satisfying both the properties D1 and D2 because $f_A(u) = u$ and $f_G(u) = \log u$, and hence $(f_A \circ f_G^{-1})(u) = \exp(u)$ is strictly convex on $(0,\infty)$.

Let us denote by $I^{f,g}_\alpha[p:q] := I^{M^f,M^g}_\alpha[p:q]$ the *quasi-arithmetic α-divergences*. Since the QAMs are symmetric means, we have $I^{f,g}_\alpha[p:q] = I^{f,g}_{1-\alpha}[q:p]$.

2.3 Limit cases of 1-divergences and 0-divergences

We seek a closed-form formula of the limit divergence $\lim_{\alpha\to 0} I_\alpha^{f,g}[p:q]$ when $\alpha \to 0$.

Lemma 1. *A first-order Taylor approximation of the quasi-arithmetic mean [30] M_α^f for a C_1 strictly increasing generator f when $\alpha \simeq 0$ yields*

$$M_\alpha^f(p:q) = p + \frac{\alpha(f(q)-f(p))}{f'(p)} + o(\alpha(f(q)-f(p))). \tag{36}$$

Proof. By taking the first-order Taylor expansion of $f^{-1}(x)$ at x_0 (i.e., Taylor polynomial of order 1), we get:

$$f^{-1}(x) = f^{-1}(x_0) + (x-x_0)(f^{-1})'(x_0) + o(x-x_0). \tag{37}$$

Using the property of the derivative of an inverse function:

$$(f^{-1})'(x) = \frac{1}{(f'(f^{-1})(x))}, \tag{38}$$

it follows that the first-order Taylor expansion of $f^{-1}(x)$ is:

$$f^{-1}(x) = f^{-1}(x_0) + (x-x_0)\frac{1}{(f'(f^{-1})(x_0))} + o(x-x_0). \tag{39}$$

Plugging $x_0 = f(p)$ and $x = f(p) + \alpha(f(q)-f(p))$, we get a *first-order approximation* of the weighted quasi-arithmetic mean M_α^f when $\alpha \to 0$:

$$M_\alpha^f(p,q) = p + \frac{\alpha(f(q)-f(p))}{f'(p)} + o(\alpha(f(q)-f(p))). \tag{40}$$

□

Let us introduce the following bivariate function:

$$E_f(p,q) := \frac{f(q)-f(p)}{f'(p)}. \tag{41}$$

Thus we obtain closed-form formula for the I_1-divergence and I_0-divergence:

Theorem 2 (Quasi-arithmetic I_1-divergence and I_0-divergence)**.** *The quasi-arithmetic I_1-divergence induced by two strictly increasing and differentiable functions f and g such that $f \circ g^{-1}$ is strictly convex is*

$$\begin{aligned} I_1^{f,g}[p:q] = \lim_{\alpha\to 1} I_\alpha^{f,g}[p:q]] &= \int (E_f(p,q) - E_g(p,q))\,\mathrm{d}\mu \geq 0, &(42)\\ &= \int \left(\frac{f(q)-f(p)}{f'(p)} - \frac{g(q)-g(p)}{g'(p)}\right)\mathrm{d}\mu. &(43)\end{aligned}$$

We have $I_0^{f,g}[p:q] = I_1^{f,g}[q:p]$.

Proof. Let us prove that $I_1^{f,g}$ is a proper divergence satisfying axioms D1 and D2. Note that a sufficient condition for $I_1^{f,g}[p:q] \geq 0$ is to check that

$$\begin{aligned} E_f(p,q) &\geq E_g(p,q), & (44)\\ \frac{f(q)-f(p)}{f'(p)} &\geq \frac{g(q)-g(p)}{g'(p)}. & (45) \end{aligned}$$

If $p = q$ μ-a.e. then clearly $I_1^{f,g}[p:q] = 0$. Consider $p \neq q$ (i.e., at some observation x: $p(x) \neq q(x)$).

We shall use the following property of a strictly convex and differentiable function h for $x < y$ (sometimes called the chordal slope lemma, see [25]):

$$h'(x) \leq \frac{h(y)-h(x)}{y-x} \leq h'(y). \quad (46)$$

We consider $h(x) = (f \circ g^{-1})(x)$ so that $h'(x) = \frac{f'(g^{-1}(x))}{g'(g^{-1}(x))}$. There are two cases to consider:

- $p < q$ and therefore $g(p) < g(q)$. Let $y = g(q)$ and $x = g(p)$ in Eq. 46. We have $h'(x) = \frac{f'(p)}{g'(p)}$ and $h'(y) = \frac{f'(q)}{g'(q)}$, and the double inequality of Eq. 46 becomes

  $$\frac{f'(p)}{g'(p)} \leq \frac{f(q)-f(p)}{g(q)-g(p)} \leq \frac{f'(q)}{g'(q)}.$$

  Since $g(q) - g(p) > 0$ and $g'(p) > 0$ and $f'(p) > 0$, we get

  $$\frac{g(q)-g(p)}{g'(p)} \leq \frac{f(q)-f(p)}{f'(p)}.$$

- $q < p$ and therefore $g(p) > g(q)$. Then the double inequality of Eq. 46 becomes

  $$\frac{f'(q)}{g'(q)} \leq \frac{f(q)-f(p)}{g(q)-g(p)} \leq \frac{f'(p)}{g'(p)}$$

  That is,

  $$\frac{f(q)-f(p)}{f'(p)} \geq \frac{g(q)-g(p)}{g'(p)},$$

  since $g(q) - g(p) < 0$.

Thus in both cases, we checked that $E_f(p(x), q(x)) \geq E_g(p(x), q(x))$. Therefore $I_1^{f,g}[p:q] \geq 0$ and since the QAMs are distinct $I_1^{f,g}[p:q] = 0$ iff $p(x) = q(x)$ μ-a.e. □

We can interpret the I_1 divergences as generalized KL divergences, and define generalized notions of cross-entropies and entropies. Since the KL divergence can be written as the cross-entropy minus the entropy, we can also decompose the I_1 divergences as follows:

$$\begin{aligned} I_1^{f,g}[p:q] &= \int \left(\frac{f(q)}{f'(p)} - \frac{g(q)}{g'(p)}\right) \mathrm{d}\mu - \int \left(\frac{f(p)}{f'(p)} - \frac{g(p)}{g'(p)}\right) \mathrm{d}\mu, & (47)\\ &= h_\times^{f,g}(p:q) - h^{f,g}(p), & (48) \end{aligned}$$

where $h_\times^{f,g}(p:q)$ denotes the (f,g)-*cross-entropy* (for a constant $c \in \mathbb{R}$):

$$h_\times^{f,g}(p:q) = \int \left(\frac{f(q)}{f'(p)} - \frac{g(q)}{g'(p)} \right) \mathrm{d}\mu + c, \tag{49}$$

and $h^{f,g}(p)$ stands for the (f,g)-*entropy* (self cross-entropy):

$$h^{f,g}(p) = h_\times^{f,g}(p:p) = \int \left(\frac{f(p)}{f'(p)} - \frac{g(p)}{g'(p)} \right) \mathrm{d}\mu + c. \tag{50}$$

We define the generalized (f,g)-*Kullback-Leibler divergence*:

$$\mathrm{KL}_{f,g}[p:q] := h_\times^{f,g}(p:q) - h^{f,g}(p). \tag{51}$$

When $f = f_A$ and $g = f_G$, we resolve the constant to $c = 0$, and recover the ordinary Shannon cross-entropy and entropy:

$$h_\times^{f_A,f_G}(p:q) = \int (q - p\log q)\mathrm{d}\mu = h_\times(p:q), \tag{52}$$

$$h^{f_A,f_G}(p:q) = h_\times^{f_A,f_G}(p:p) = \int (p - p\log p)\mathrm{d}\mu = h(p), \tag{53}$$

and we have the (f_A, f_G)-Kullback-Leibler divergence that is the extended Kullback-Leibler divergence:

$$\mathrm{KL}_{f_A,f_G}[p:q] = \mathrm{KL}_e[p:q] = h_\times(p:q) - h(p) = \int (p\log\frac{p}{q} + q - p)\mathrm{d}\mu. \tag{54}$$

Thus we have the (f,g)-cross-entropy and (f,g)-entropy expressed as

$$h_\times^{f,g}(p:q) = \int \left(\frac{f(q)}{f'(p)} - \frac{g(q)}{g'(p)} \right) \mathrm{d}\mu, \tag{55}$$

$$h^{f,g}(p) = \int \left(\frac{f(p)}{f'(p)} - \frac{g(p)}{g'(p)} \right) \mathrm{d}\mu. \tag{56}$$

In general, we can define the (f,g)-Jeffreys' divergence as:

$$J^{f,g}[p:q] = \mathrm{KL}^{f,g}[p:q] + \mathrm{KL}^{f,g}[q:p]. \tag{57}$$

Thus we define the quasi-arithmetic mean α-divergences as follows:

Theorem 3 (Quasi-arithmetic α-divergences)**.** *Let f and g be two strictly continuously increasing and differentiable functions on $(0,\infty)$ such that $f \circ g^{-1}$ is strictly convex. Then the quasi-arithmetic α-divergences induced by (f,g) for $\alpha \in [0,1]$ is*

$$I_\alpha^{f,g}[p:q] = \begin{cases} \frac{1}{\alpha(1-\alpha)} \int \left(M_{1-\alpha}^f(p:q) - M_{1-\alpha}^g(p:q) \right) \mathrm{d}\mu, & \alpha \in \mathbb{R}\backslash\{0,1\}. \\ I_1^{f,g}[p:q] = \int \left(\frac{f(q)-f(p)}{f'(p)} - \frac{g(q)-g(p)}{g'(p)} \right) \mathrm{d}\mu & \alpha = 1, \\ I_0^{f,g}[p:q] = I_1^{f,g}(q:p) = \int \left(\frac{f(p)-f(q)}{f'(q)} - \frac{g(p)-g(q)}{g'(q)} \right) \mathrm{d}\mu, & \alpha = 0. \end{cases} \tag{58}$$

When $f(u) = f_A(u) = u$ ($M^f = A$) and $g(u) = f_G(u) = \log u$ ($M^g = G$), we get:

$$I_1^{A,G}[p:q] = \int \left(q - p - p \log \frac{q}{p} \right) \mathrm{d}\mu = \mathrm{KL}_e[p:q] = I_1[p:q], \tag{59}$$

the Kullback-Leibler divergence (KLD) extended to positive densities, and $I_0 = \mathrm{KL}_e^*$ the reverse extended KLD.

We can rewrite the α-divergence $I_\alpha^{f,g}[p:q]$ for $\alpha \in (0,1)$ as

$$I_\alpha^{f,g}[p:q] = \frac{1}{\alpha(1-\alpha)} \left(S_{1-\alpha}^f(p:q) - S_{1-\alpha}^g(p:q) \right), \tag{60}$$

where

$$S_\lambda^h(p:q) := \int M_\lambda^h(p:q) \mathrm{d}\mu. \tag{61}$$

Zhang [39] (p. 188-189) considered the (A, M^ρ) α_A-divergences:

$$D_\alpha^\rho[p:q] = \frac{4}{1-\alpha^2} \int \left(\frac{1-\alpha}{2} p + \frac{1+\alpha}{2} q - \rho^{-1} \left(\frac{1-\alpha}{2} \rho(p) + \frac{1+\alpha}{2} \rho(q) \right) \right) \mathrm{d}\mu. \tag{62}$$

The formula he obtained for $D_{\pm 1}^\rho(p:q)$:

$$D_1^\rho[p:q] = \int \left(p - q - \left(\rho^{-1}\right)' (\rho(q))(\rho(p) - \rho(q)) \right) \mathrm{d}\mu = D_{-1}^\rho[q:p], \tag{63}$$

is in accordance with our generic formula of Eq. 42 since $(\rho^{-1}(x))' = \frac{1}{\rho'(\rho^{-1}(x))}$. Notice that $A_\alpha \geq P_\alpha^r$ for $r \leq 1$: The arithmetic weighted mean dominates the weighted power means P^r when $r \leq 1$.

Furthermore, by imposing the homogeneity condition $I_\alpha^{A,M^\rho}[\lambda p : \lambda q] = \lambda I_\alpha^{A,M^\rho}[p:q]$ for $\lambda > 0$, Zhang [39] obtained the class of the (α_A, β_A)-divergences for $(\alpha_A, \beta_A) \in [-1,1]^2$:

$$D_{\alpha_A,\beta_A}[p:q] = \frac{4}{1-\alpha_A^2} \frac{2}{1+\beta_A} \int \left(\frac{1-\alpha_A}{2} p + \frac{1+\alpha_A}{2} q - \left(\frac{1-\alpha_A}{2} p^{\frac{1-\beta_A}{2}} + \frac{1+\alpha_A}{2} q^{\frac{1-\beta_A}{2}} \right)^{\frac{2}{1-\beta_A}} \right) \mathrm{d}\mu. \tag{64}$$

2.4 Generalized KL divergences as conformal Bregman divergences on monotone embeddings

We can rewrite the generalized KLDs $I_1^{f,g}$ as a *conformal Bregman representational divergence* [32, 33, 34] as follows:

Theorem 4. *The generalized KLDs $I_1^{f,g}$ divergences are conformal Bregman representational divergences:*

$$I_1^{f,g}[p:q] = \int \frac{1}{f'(p)} B_F(g(q) : g(p)) \mathrm{d}\mu, \tag{65}$$

with $F = f \circ g^{-1}$ a strictly convex and differentiable Bregman convex generator.

Proof. For the Bregman strictly convex and differentiable generator $F = f \circ g^{-1}$, we expand the following conformal divergence:

$$\begin{aligned}
\frac{1}{f'(p)} B_F(g(q) : g(p)) &= \frac{1}{f'(p)} \left(F(g(q)) - F(g(p)) - (g(q) - g(p)) F'(g(p)) \right), && (66) \\
&= \frac{1}{f'(p)} \left((f(q) - f(p)) - (g(q) - g(p)) \frac{f'(p)}{g'(p)} \right), && (67)
\end{aligned}$$

since $(g^{-1} \circ g)(x) = x$ and $F'(g(x)) = \frac{f'(x)}{g'(x)}$. It follows that

$$\begin{aligned}
\frac{1}{f'(p)} B_F(g(q) : g(p)) &= \frac{f(q) - f(p)}{f'(p)} - \frac{g(q) - g(p)}{g'(p)}, && (68) \\
&= E_f(p, q) - E_g(p, q) = I_1^{f,g}[p : q]. && (69)
\end{aligned}$$

Hence, we easily check that $I_1^{f,g}[p : q] = \int \frac{1}{f'(p)} B_F(g(q) : g(p)) \mathrm{d}\mu \geq 0$ since $f'(p) > 0$ and $B_F \geq 0$. □

In general, for a functional generator f and a strictly monotonic representational function r, we can define the representational Bregman divergence [29] $B_{f \circ r^{-1}}(r(p) : r(q))$ provided that $F = f \circ r^{-1}$ is a Bregman generator (i.e., strictly convex and differentiable).

In [30], a generalization of the Bregman divergences was obtained using the *comparative convexity* induced by two abstract means M and N to define (M, N)-Bregman divergences as limit of scaled (M, N)-Jensen divergences. The skew (M, N)-Jensen divergences are defined for $\alpha \in (0, 1)$ by:

$$J_{F,\alpha}^{M,N}(p : q) = \frac{1}{\alpha(1 - \alpha)} \left(N_\alpha(F(p), F(q))) - F(M_\alpha(p, q))) \right), \quad (70)$$

where M_α and N_α are weighted means that should be *regular* [30] (i.e., homogeneous, symmetric, continuous and increasing in each variable). Then we can define the (M, N)-Bregman divergence as

$$\begin{aligned}
B_F^{M,N}[p : q] &= \lim_{\alpha \to 1^-} J_{F,\alpha}^{M,N}(p : q), \\
&= \lim_{\alpha \to 1^-} \frac{1}{\alpha(1 - \alpha)} \left(N_\alpha(F(p), F(q))) - F(M_\alpha(p, q))) \right).
\end{aligned}$$

The formula obtained in [30] for the quasi-arithmetic means M^f and M^g and a functional generator F that is (M^f, M^g)-convex is:

$$\begin{aligned}
B_F^{f,g}(p : q) &= \frac{g(F(p)) - g(F(q))}{g'(F(q))} - \frac{f(p) - f(q)}{f'(q)} F'(q), && (71) \\
&= \frac{1}{f'(F(q))} B_{g \circ F \circ f^{-1}}(f(p) : f(q)) \geq 0. && (72)
\end{aligned}$$

This is a conformal divergence [32] that can be written using the E_f terms as:

$$B_F^{f,g}(p : q) = E_g(F(q), F(p)) - E_f(q, p) F'(q). \quad (73)$$

A function F is (M^f, M^g)-convex iff $g \circ F \circ f^{-1}$ is (ordinary) convex [30].

The information geometry induced by a Bregman divergence (or equivalently by its convex generator) is a dually flat space [2, 27]. The dualistic structure induced by a conformal Bregman representational divergence is related to conformal flattening [33, 34].

Following the work of Ohara [33, 34], the *geometric divergence* $\rho(p,r)$ (a contrast function in affine differential geometry) induced by a pair (L,M) of strictly monotone smooth functions between two distributions p and r of the d-dimensional probability simplex Δ_d is defined by (Eq. 28 in [33]):

$$\rho(p:r) = \frac{1}{\Lambda(r)} \sum_{i=1}^{d+1} \frac{L(p_i) - L(r_i)}{L'(r_i)} = \frac{1}{\Lambda(r)} \sum_{i=1}^{d+1} E_L(r_i, p_i), \tag{74}$$

where $\Lambda(r) = \sum_{i=1}^{d+1} \frac{1}{L'(p_i)} p_i$. Affine immersions [20] can be interpreted as special embeddings.

Let ρ be a divergence (contrast function) and $({}^{\rho}g, {}^{\rho}\nabla, {}^{\rho}\nabla^*)$ be the induced statistical manifold structure with

$$\begin{aligned}
{}^{\rho}g_{ij}(p) &:= -(\partial_i)_p(\partial_j)_p\ \rho(p,q)|_{q=p}, & (75)\\
\Gamma_{ij,k}(p) &:= -(\partial_i)_p(\partial_j)_p(\partial_k)_q\ \rho(p,q)|_{q=p}, & (76)\\
\Gamma^*_{ij,k}(p) &:= -(\partial_i)_p(\partial_j)_q(\partial_k)_q\ \rho(p,q)|_{q=p}, & (77)
\end{aligned}$$

where $(\partial_i)_s$ denotes the tangent vector at s of a vector field ∂_i.

Consider a conformal divergence $\rho_\kappa(p:q) = \kappa(q)\rho(p:q)$ for a positive function $\kappa(q) > 0$, called the conformal factor. Then the induced statistical manifold [12, 2] $({}^{\rho_\kappa}g, {}^{\rho_\kappa}\nabla, {}^{\rho_\kappa}\nabla^*)$ is 1-conformally equivalent to $({}^{\rho}g, {}^{\rho}\nabla, {}^{\rho}\nabla^*)$ and we have

$$\begin{aligned}
{}^{\rho_\kappa}g &= \kappa\ {}^{\rho}g, & (78)\\
{}^{\rho}g({}^{\rho_\kappa}\nabla_X Y, Z) &= {}^{\rho}g({}^{\rho}\nabla_X Y, Z) - d(\log\kappa)(Z)\,{}^{\rho}g(X,Y). & (79)
\end{aligned}$$

The dual affine connections ${}^{\rho_\kappa}\nabla^*$ and ${}^{\rho}\nabla^*$ are projectively equivalent [20] (and ${}^{\rho}\nabla^*$ is said -1-conformally flat).

Conformal flattening [33, 34] consists in choosing the conformal factor κ such that $({}^{\rho_\kappa}g, {}^{\rho_\kappa}\nabla, {}^{\rho_\kappa}\nabla)$ becomes a dually flat space [2] equipped with a canonical Bregman divergence.

Therefore it follows that the statistical manifolds induced by the 1-divergence $I_1^{f,g}$ is a representational 1-conformally flat statistical manifold.

3 The subfamily of homogeneous (r,s)-power α-divergences

In particular, we can define the *(r,s)-power α-divergences* from two power means $P_r = M^{\mathrm{pow}_r}$ and $P_s = M^{\mathrm{pow}_s}$ with $r > s$ (and $P_r \geq P_s$) with the family of generators $\mathrm{pow}_l(u) = u^l$. Indeed, we check that $f_{rs}(u) := \mathrm{pow}_r \circ \mathrm{pow}_s^{-1}(u) = u^{\frac{r}{s}}$ is strictly convex on $(0,\infty)$ since $f''_{rs}(u) = \frac{r}{s}\left(\frac{r}{s} - 1\right) u^{\frac{r}{s}-2} > 0$ for $r > s$. Thus P_r and P_s are two QAMs which are both comparable and distinct. Table 1 lists the expressions of $E_r(p,q) := E_{\mathrm{pow}_r}(p,q)$ obtained from the power mean generators $\mathrm{pow}_r(u) = u^r$.

We conclude with the definition of the *(r,s)-power α-divergences*:

Corollary 1 (power α-divergences)**.** *Given $r > s$, the α-power divergences are defined for $r > s$ and $r, s \neq 0$ by*

$$I_\alpha^{r,s}[p:q] = \begin{cases} \frac{1}{\alpha(1-\alpha)} \int \left((\alpha p^r + (1-\alpha)q^r)^{\frac{1}{r}} - (\alpha p^s + (1-\alpha)q^s)^{\frac{1}{s}} \right) \mathrm{d}\mu, & \alpha \in \mathbb{R}\backslash\{0,1\}. \\ I_1^{r,s}[p:q] = \int \left(\frac{q^r - p^r}{r p^{r-1}} - \frac{q^s - p^s}{s p^{s-1}} \right) \mathrm{d}\mu & \alpha = 1, \\ I_0^{r,s}[p:q] = I_1^{r,s}(q:p) & \alpha = 0. \end{cases} \tag{80}$$

Power mean	$E_r(p,q)$
$P_r(r \in \mathbb{R}\backslash\{0\})$	$\frac{q^r-p^r}{rp^{r-1}}$
$Q(r=2)$	$\frac{q^2-p^2}{2p}$
$A(r=1)$	$q-p$
$G(r=0)$	$p\log\frac{q}{p}$
$H(r=-1)$	$-p^2\left(\frac{1}{q}-\frac{1}{p}\right)=p-\frac{p^2}{q}$

Table 1: Expressions of the terms E_r for the family of power means P_r, $r \in \mathbb{R}$.

When $r=0$, we get the following power α-divergences for $s<0$:

$$I_\alpha^{r,s}[p:q]=\begin{cases}\frac{1}{\alpha(1-\alpha)}\int\left(p^\alpha q^{1-\alpha}-(\alpha p^s+(1-\alpha)q^s)^{\frac{1}{s}}\right)\mathrm{d}\mu, & \alpha\in\mathbb{R}\backslash\{0,1\}.\\ I_1^{r,s}[p:q]=\int\left(p\log\frac{q}{p}-\frac{q^s-p^s}{sp^{s-1}}\right)\mathrm{d}\mu & \alpha=1,\\ I_0^{r,s}[p:q]=I_1^{r,s}[q:p] & \alpha=0.\end{cases} \tag{81}$$

When $s=0$, we get the following power α-divergences for $r>0$:

$$I_\alpha^{r,s}[p:q]=\begin{cases}\frac{1}{\alpha(1-\alpha)}\int\left((\alpha p^r+(1-\alpha)q^r)^{\frac{1}{r}}-p^\alpha q^{1-\alpha}\right)\mathrm{d}\mu, & \alpha\in\mathbb{R}\backslash\{0,1\}.\\ I_1^{r,s}[p:q]=\int\left(\frac{q^r-p^r}{rp^{r-1}}-p\log\frac{q}{p}\right)\mathrm{d}\mu & \alpha=1,\\ I_0^{r,s}[p:q]=I_1^{r,s}[q:p] & \alpha=0.\end{cases} \tag{82}$$

In particular, we get the following family of (A,H) α-*divergences*:

$$I_\alpha^{A,H}[p:q]=I_\alpha^{1,-1}[p:q]=\begin{cases}\frac{1}{\alpha(1-\alpha)}\int\left(\alpha p+(1-\alpha)q-\frac{pq}{\alpha q+(1-\alpha)p}\right)\mathrm{d}\mu, & \alpha\in\mathbb{R}\backslash\{0,1\}.\\ I_1^{1,-1}[p:q]=\int\left(q-2p+\frac{p^2}{q}\right)\mathrm{d}\mu & \alpha=1,\\ I_0^{1,-1}[p:q]=I_1^{1,-1}(q:p) & \alpha=0.\end{cases}, \tag{83}$$

and the family of (G,H) α-*divergences*:

$$I_\alpha^{G,H}[p:q]=I_\alpha^{0,-1}(p:q)=\begin{cases}\frac{1}{\alpha(1-\alpha)}\int\left(p^\alpha q^{1-\alpha}-\frac{pq}{\alpha q+(1-\alpha)p}\right)\mathrm{d}\mu, & \alpha\in\mathbb{R}\backslash\{0,1\}.\\ I_1^{0,-1}[p:q]=\int\left(p\log\frac{q}{p}-p+\frac{p^2}{q}\right)\mathrm{d}\mu & \alpha=1,\\ I_0^{0,-1}[p:q]=I_1^{0,-1}[q:p] & \alpha=0.\end{cases}. \tag{84}$$

The (r,s)-power α-divergences for $r,s\neq 0$ yield homogeneous divergences: $I_\alpha^{r,s}[\lambda p:\lambda q]=\lambda I_\alpha^{r,s}[p:q]$ for any $\lambda>0$ because the power means are homogeneous: $P_\alpha^r(\lambda x,\lambda y)=\lambda P_\alpha^r(x,y)=\lambda x P_\alpha^r\left(1,\frac{y}{x}\right)$. Thus the $I_\alpha^{r,s}$-divergences are Csiszár f-divergences [10]

$$I_\alpha^{r,s}[p:q]=\int p(x)f_{r,s}\left(\frac{q(x)}{p(x)}\right)\mathrm{d}\mu \tag{85}$$

for the generator

$$f_{r,s}(u)=\frac{1}{\alpha(1-\alpha)}(P_\alpha^r(1,u)-P^s(1,u)). \tag{86}$$

Thus the family of (r,s)-power α-divergences are *homogeneous divergences*:

$$I_\alpha^{r,s}[\lambda p:\lambda q]=\lambda I_\alpha^{r,s}[p:q],\quad \forall\lambda>0 \tag{87}$$

4 Conclusion, discussion and perspectives

For two comparable strict means $M \geq N$, one can define the (M,N)-*divergence* as

$$I^{M,N}[p:q] := 4\int (M(p,q) - N(p,q))\,\mathrm{d}\mu. \tag{88}$$

When the property of strict comparable means extend to their induced weighted means M_α and N_α (i.e., $M_\alpha \geq N_\alpha$), one can further define the (M,N) α-*divergences* for $\alpha \in (0,1)$:

$$I_\alpha^{M,N}[p:q] := \frac{1}{\alpha(1-\alpha)}\int (M_{1-\alpha}(p,q) - N_{1-\alpha}(p,q))\,\mathrm{d}\mu, \tag{89}$$

so that $I^{M,N}[p:q] = I_{\frac{1}{2}}^{M,N}[p:q]$. When the weighted means are symmetric, the reference duality holds (i.e., $I_\alpha^{M,N}[q:p] = I_{1-\alpha}^{M,N}[p:q]$), and we can define the (M,N)-equivalent of the Kullback-Leibler divergence, i.e., the (M,N) 1-divergence, as the limit case (when it exists): $I_1^{M,N}[p:q] = \lim_{\alpha\to 1} I_\alpha^{M,N}[p:q]$.

We proved that the quasi-arithmetic weighted means M_α^f and M_α^g are strictly comparable whenever $f\circ g^{-1}$ is *strictly* convex. In the limit cases of $\alpha \to 0$ and $\alpha \to 1$, we reported a closed-form formula for the equivalent of the Kullback-Leibler divergence and the reverse Kullback-Leibler divergence. We reported closed-form formula for the quasi-arithmetic α-divergences $I_\alpha^{f,g}(p:q)$ for $\alpha \in [0,1]$ (Theorem 3), and for the subfamily of homogeneous (r,s)-power α-divergences $I_\alpha^{r,s}(p:q)$ induced by power means (Corollary 1). The ordinary (A,G) α-divergences, the (A,H) α-divergences and the (G,H) α-divergences are examples of (r,s)-power α-divergences for $(r,s) = (1,0)$, $(r,s) = (1,-1)$ and $(r,s) = (0,-1)$, respectively.

Generalized α-divergences may prove useful in reporting closed-form formula between parametric densities: For example, consider the ordinary α-divergences between two scale Cauchy densities $p_1(x) = \frac{1}{\pi}\frac{s_1}{x^2+s_1^2}$ and $p_2(x) = \frac{1}{\pi}\frac{s_2}{x^2+s_2^2}$: There is no obvious closed-form for the ordinary α-divergences but we can report easily a closed-form for the (A,H) α-divergences following the calculus reported in [26]:

$$\begin{aligned} I_\alpha^{A,H}[p_1:p_2] &= \frac{1}{\alpha(1-\alpha)}\left(1 - \int H_{1-\alpha}(p_1(x),p_2(x))\mathrm{d}\mu(x)\right), &(90)\\ &= \frac{1}{\alpha(1-\alpha)}\left(1 - \frac{s_1 s_2}{(\alpha s_1 + (1-\alpha)s_2)s_{1-\alpha}}\right), &(91)\end{aligned}$$

with $s_\alpha = \sqrt{\frac{\alpha s_1 s_2^2 + (1-\alpha)s_2 s_1^2}{\alpha s_1 + (1-\alpha)s_2}}$. In general, instead of considering ordinary α-divergences in applications, one may consider the (r,s)-power α-divergences, and tune the three parameters (r,s,α) according to the various tasks (say, by cross-validation in supervised machine learning tasks). We note that the quasi-arithmetic means have been recently considered by Eguchi et al. [13] to define a novel non-parametric dualistic structure of information geometry via generalizations of the e-geodesics and the m-geodesics.

The elucidation the (f,g) α-*geometry* of these generalized α-divergences is left for future work. For the limit cases of $\alpha \to 0$ or of $\alpha \to 1$, we proved that the limit KL-type divergences amount to a conformal Bregman divergence on a monotone embedding, and briefly showed the connection with conformal divergences and conformal flattening. The geometry of conformal flattening [33, 34] and the relationships with the (ρ,τ)-monotone embeddings [24] shall be further studied.

Applications of (f, g) α-divergences to center-based clustering [31] and to generalized α-divergences in positive-definite matrix spaces [2] shall also be considered in future work. The quasi-arithmetic weighted means are convex if and only if the generators are differentiable with positive first derivatives with corresponding functions $-E_f$ of Eq. 41 convex (Theorem 4 of [6], i.e., convexity of the quasi-arithmetic weighted means does *not* depend on the weights). For example, when both quasiarithmetic means are convex means, the quasi-arithmetic α-divergence is the difference of two convex mean functions, and the k-means centroid computation amounts to solve a *Difference of Convex* (DC) program which can solved by the smooth DC Algorithm, DCA, called the Convex-ConCave Procedure [28]. Similarly, when $\alpha \in \{0, 1\}$, we get a DC program since $I_\alpha^{f,g}$ writes as a difference of convex functions.

References

[1] Vahideh Ahrari, Arezou Habibirad, and S Baratpour. Exponentiality test based on alpha-divergence and gamma-divergence. *Communications in Statistics-Simulation and Computation*, 48(4):1138–1152, 2019.

[2] S. Amari. *Information Geometry and Its Applications*. Applied Mathematical Sciences. Springer Japan, 2016.

[3] Shun-ichi Amari. Integration of stochastic models by minimizing α-divergence. *Neural computation*, 19(10):2780–2796, 2007.

[4] Michèle Basseville. Divergence measures for statistical data processing — an annotated bibliography. *Signal Processing*, 93(4):621–633, 2013.

[5] Peter S Bullen, Dragoslav S Mitrinovic, and M Vasic. *Means and their Inequalities*, volume 31. Springer Science & Business Media, 2013.

[6] Jacek Chudziak, Dorota Glazowska, Justyna Jarczyk, and Witold Jarczyk. On weighted quasi-arithmetic means which are convex. *Mathematical Inequalities & Applications*, 22(4):1123–1136, 2019.

[7] Andrzej Cichocki and Shun-ichi Amari. Families of alpha-beta-and gamma-divergences: Flexible and robust measures of similarities. *Entropy*, 12(6):1532–1568, 2010.

[8] Andrzej Cichocki, Hyekyoung Lee, Yong-Deok Kim, and Seungjin Choi. Non-negative matrix factorization with α-divergence. *Pattern Recognition Letters*, 29(9):1433–1440, 2008.

[9] Thomas M. Cover and Joy A. Thomas. *Elements of information theory*. John Wiley & Sons, 2012.

[10] Imre Csiszár. Information-type measures of difference of probability distributions and indirect observation. *Studia Scientiarum Mathematicarum Hungarica*, 2:229–318, 1967.

[11] Bruno De Finetti. Sul concetto di media. *Istituto italiano degli attuari*, 3:369–396, 1931.

[12] Shinto Eguchi. Geometry of minimum contrast. *Hiroshima Mathematical Journal*, 22(3):631–647, 1992.

[13] Shinto Eguchi, Osamu Komori, and Atsumi Ohara. Information geometry associated with generalized means. In *Information Geometry and its Applications IV*, pages 279–295. Springer, 2016.

[14] Alison L. Gibbs and Francis Edward Su. On choosing and bounding probability metrics. *International statistical review*, 70(3):419–435, 2002.

[15] G.H. Hardy, J.E. Littlewood, and G. Pólya. *Inequalities*. Cambridge Mathematical Library. Cambridge University Press, 1988.

[16] Otto Ludwig Holder. Über einen Mittelwertssatz. *Nachr. Akad. Wiss. Gottingen Math.-Phys. Kl.*, pages 38–47, 1889.

[17] Asif Iqbal and Abd-Krim Seghouane. An α-divergence-based approach for robust dictionary learning. *IEEE Transactions on Image Processing*, 28(11):5729–5739, 2019.

[18] Justyna Jarczyk. When Lagrangean and quasi-arithmetic means coincide. *J. Inequal. Pure Appl. Math.*, 8:71, 2007.

[19] Andrey Nikolaevich Kolmogorov. Sur la notion de moyenne. *Acad. Naz. Lincei Mem. Cl. Sci. His. Mat. Natur. Sez.*, 12:388–391, 1930.

[20] Takashi Kurose. On the divergences of 1-conformally flat statistical manifolds. *Tohoku Mathematical Journal, Second Series*, 46(3):427–433, 1994.

[21] Gyula Maksa and Zsolt Páles. Remarks on the comparison of weighted quasi-arithmetic means. *Colloquium Mathematicae*, 120(1):77–84, 2010.

[22] Yuzo Maruyama, Takeru Matsuda, and Toshio Ohnishi. Harmonic Bayesian prediction under α-divergence. *IEEE Transactions on Information Theory*, 65(9):5352–5366, 2019.

[23] Mitio Nagumo. Über eine klasse der mittelwerte. *Japanese journal of mathematics: transactions and abstracts*, 7(0):71–79, 1930.

[24] Jan Naudts and Jun Zhang. Rho–tau embedding and gauge freedom in information geometry. *Information Geometry*, 1(1):79–115, 2018.

[25] Constantin P. Niculescu and Lars-Erik Persson. *Convex functions and their applications: A contemporary approach*. Springer Science & Business Media, 2006. First edition.

[26] Frank Nielsen. Generalized Bhattacharyya and Chernoff upper bounds on Bayes error using quasi-arithmetic means. *Pattern Recognition Letters*, 42:25–34, 2014.

[27] Frank Nielsen. An elementary introduction to information geometry. *arXiv preprint arXiv:1808.08271*, 2018.

[28] Frank Nielsen and Sylvain Boltz. The Burbea-Rao and Bhattacharyya centroids. *IEEE Transactions on Information Theory*, 57(8):5455–5466, 2011.

[29] Frank Nielsen and Richard Nock. The dual Voronoi diagrams with respect to representational Bregman divergences. In *Sixth International Symposium on Voronoi Diagrams (ISVD)*, pages 71–78. IEEE, 2009.

[30] Frank Nielsen and Richard Nock. Generalizing skew Jensen divergences and Bregman divergences with comparative convexity. *IEEE Signal Processing Letters*, 24(8):1123–1127, 2017.

[31] Frank Nielsen, Richard Nock, and Shun-ichi Amari. On clustering histograms with k-means by using mixed α-divergences. *Entropy*, 16(6):3273–3301, 2014.

[32] Richard Nock, Frank Nielsen, and Shun-ichi Amari. On conformal divergences and their population minimizers. *IEEE Transactions on Information Theory*, 62(1):527–538, 2015.

[33] Atsumi Ohara. Conformal flattening for deformed information geometries on the probability simplex. *Entropy*, 20(3):186, 2018.

[34] Atsumi Ohara. Conformal flattening on the probability simplex and its applications to voronoi partitions and centroids. In *Geometric Structures of Information*, pages 51–68. Springer, 2019.

[35] Zsolt Páles and Amr Zakaria. On the equality of bajraktarević means to quasi-arithmetic means. *Results in Mathematics*, 75(1):19, 2020.

[36] Alfréd Rényi. On measures of entropy and information. In *Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability*. The Regents of the University of California, 1961. Volume 1: Contributions to the Theory of Statistics.

[37] Auxiliadora Sarmiento, Irene Fondón, Iván Durán-Díaz, and Sergio Cruces. Centroid-based clustering with $\alpha\beta$-divergences. *Entropy*, 21(2):196, 2019.

[38] Junichiro Wada and Yuta Kamahara. Studying malapportionment using α-divergence. *Mathematical Social Sciences*, 93:77–89, 2018.

[39] Jun Zhang. Divergence function, duality, and convex analysis. *Neural Computation*, 16(1):159–195, 2004.