

ON SHARP FRONTS AND ALMOST-SHARP FRONTS FOR SINGULAR SQG

CALVIN KHOR

*School of Mathematical Sciences, Beijing Normal University,
Beijing 100875, P. R. China*

JOSÉ L. RODRIGO

*Mathematics Research Centre, Zeeman Building,
University of Warwick, Coventry CV4 7AL,
United Kingdom*

ABSTRACT

In this paper we consider a family of active scalars with a velocity field given by $u = \Lambda^{-1+\alpha}\nabla^\perp\theta$, for $\alpha \in (0, 1)$. This family of equations is a more singular version of the two-dimensional Surface Quasi-Geostrophic (SQG) equation, which would correspond to $\alpha = 0$.

We consider the evolution of sharp fronts by studying families of almost-sharp fronts. These are smooth solutions with simple geometry in which a sharp transition in the solution occurs in a tubular neighbourhood (of size δ). We study their evolution and that of compatible curves, and introduce the notion of a spine for which we obtain improved evolution results, gaining a full power (of δ) compared to other compatible curves.

1. INTRODUCTION

In this paper, we develop and study the notions of sharp front, almost-sharp front, and spine curve for the following more singular

E-mail addresses: C.Khor@bnu.edu.cn, J.Rodrigo@warwick.ac.uk.
Date: 2024-03-14 15:11:36Z.

version of the two-dimensional SQG equation,

$$\begin{cases} \partial_t \theta + u \cdot \nabla \theta = 0, \\ u = \nabla^\perp \Lambda^{-1+\alpha} \theta. \end{cases} \quad (1.1)$$

We will focus in the range $\alpha \in (0, 1)$. Here $\theta = \theta(x, t) \in \mathbb{R}$, $x \in \mathbb{R}^2$, and $t \geq 0$ and the operator $\Lambda = |\nabla|$ is the half Laplacian on \mathbb{R}^2 . To simplify the presentation we will define $\Lambda^{-1+\alpha}$ as given by a convolution with the kernel $K(x) = |x|^{-1-\alpha} \in L^2_{\text{loc}}(\mathbb{R}^2)$,

$$\Lambda^{-1+\alpha} f(x) = K * f(x) = \int_{\mathbb{R}^2} \frac{f(y)}{|x - y|^{1+\alpha}} dy.$$

In \mathbb{R}^2 , usually one has $\Lambda^{-\beta} f = c_\beta |x|^{\beta-2} * f$ but we ignore c_β here. For $\alpha \in [-1, 0]$, the family interpolates between the 2D Euler equation ($\alpha = -1$) and the 2D SQG equation ($\alpha = 0$). The model with $\alpha \in (0, 1)$ is the natural extrapolation of this family and the focus of this paper.

Sharp fronts are weak solutions of (1.1) that have the special form

$$\theta(x, t) = \mathbf{1}_{x \in A(t)},$$

where A is a bounded simply connected domain with sufficiently regular boundary (say C^2), and $\mathbf{1}_U$ is the indicator function of the set U .

Periodic graph-type SQG sharp fronts have been considered in [13] and [6] by periodising one of the space variables. For SQG, sharp fronts have already been studied in the Sobolev setting in [9] and [2] (see Chae et. al. [1] as well).

Non-periodic graph-type sharp fronts of (1.1) were studied in the recent papers, and of Hunter, Shu, and Zhang. They gave two different approaches in [17] and [19] to deriving the correct contour dynamics equation in this setting, and proved [16] global existence under a certain smallness condition. They also study two-front dynamics [15] and approximate equations for sharp fronts [18]; these differ from the almost-sharp fronts considered in this paper, which are smooth functions solving (1.1) exactly, approximating a sharp front.

Finally, a second paper [20] by the authors has been prepared alongside this one that proves local existence of sharp fronts of (1.1) a bounded domain in the analytic setting. This is an analogous result to the existence theorem for SQG in [6].

Analogously to [7] and [8], we define an almost-sharp front (ASF) of size δ as a regular¹ approximation of a sharp front:

$$\theta(x, t) = \begin{cases} 1 & x \in A_{\text{in}}(t), \\ \text{regular} & x \in A_{\text{mid}}(t), \\ 0 & x \in \mathbb{R}^2 \setminus (A_{\text{in}}(t) \cup A_{\text{mid}}(t)), \end{cases}$$

where $\text{supp } \nabla \theta \subset A_{\text{mid}}$ and this ‘transition region’ has area $O(\delta)$, so that in the limit as $\delta \rightarrow 0$, one formally recovers a sharp front.

In the Appendix, we derive asymptotics for integrals appearing in the equation, which we use to write down an approximate equation for an almost-sharp front. In doing so, we discover a significant difference with [7], which corresponds to $\alpha = 0$: the appearance of a logarithm in the equation of [7] and the property $\log(ab) = \log a + \log b$ led to the development of an ‘approximate unwinding’ for their equation in [8]. However, in our case, the singularity is stronger and we have $|ab|^{-\alpha} \neq |a|^{-\alpha} + |b|^{-\alpha}$, so we have a nonlinear term that explodes as $\delta \rightarrow 0$. Regardless, the analogous function to the h function of [7] does satisfy a well behaved linear homogeneous equation in the limit as $\delta \rightarrow 0$.

The remainder of this paper is organised as follows. In Section 2, we set up some notation. Section 3 discusses sharp fronts, with almost-sharp fronts introduced in Section 4. The regularisation effect for Ω that leads to a limit equation for h is discussed in Section 5. Finally, Section 6 discusses the notion of compatible curves for almost-sharp fronts, and Section 7 introduces the notion of a spine, and establishes an improved evolution result.

2. NOTATION

Let $\mathbb{T} := \mathbb{R}/\mathbb{Z}$, with fundamental domain $[-1/2, 1/2)$. The 90° counter-clockwise rotation of $x = \begin{pmatrix} a \\ b \end{pmatrix} \in \mathbb{R}^2$ is $x^\perp := \begin{pmatrix} -b \\ a \end{pmatrix}$, and similarly $\nabla^\perp = \begin{pmatrix} -\partial_{x^2} \\ \partial_{x^1} \end{pmatrix}$. We will frequently employ the shorthand notation

$$\begin{aligned} \int_A F(g, f_*) \, ds_* &:= \int_A F(g(s), f(s_*)) \, ds_*, \\ \int_A \int_B F(g, f_*) \, ds_* \, d\xi_* &:= \int_A \int_B F(g(s, \xi), f(s_*, \xi_*)) \, ds_* \, d\xi_*. \end{aligned}$$

That is, evaluation at s, ξ is assumed unless a function is subscripted by $*$, and then we will assume it is evaluated at s_*, ξ_* .

¹We do not attempt to optimise the regularity conditions for the front or almost-sharp front, taking them to be C^2 throughout.

3. EVOLUTION EQUATION FOR A SHARP FRONT

In this section, we give the contour dynamics equation (CDE) for sharp fronts. Recall that $K(x) = |x|^{-1-\alpha}$. The equation we study (1.1) can be written equivalently as

$$\begin{cases} \partial_t \theta + u \cdot \nabla \theta = 0, \\ u = (\nabla^\perp K) * \theta. \end{cases} \quad (3.1)$$

Definition 3.1. We say that $\theta = \theta(x, t)$ for $x \in \mathbb{R}^2, t \geq 0$ is a weak solution to (3.1) if there exists $T > 0$ such that $\theta \in L^2(0, T; L^2(\mathbb{R}^2))$, $u = \nabla^\perp K * \theta \in L^2(0, T; L^2(\mathbb{R}^2; \mathbb{R}^2))$, and for any $\phi \in C_c^\infty((0, T) \times \mathbb{R}^2)$

$$\int_0^T \int_{\mathbb{R}^2} \theta \partial_t \phi + u \cdot \nabla \phi \, dx \, dt = 0.$$

Definition 3.2. (Sharp front) We say that a weak solution θ to (3.1) is a sharp-front solution to (3.1) if

- (1) for each $t \in [0, T]$, there exists a bounded simply connected closed set $A(t)$ with sufficiently regular boundary (C^2), parameterised (anti-clockwise) by $z = z(s, t)$, with $s \in \mathbb{T}, t \in [0, T]$,
- (2) θ is the indicator function of $A(t)$ for each $t \in (0, T)$,

$$\theta(x, t) = \mathbf{1}_{x \in A(t)} := \begin{cases} 1 & x \in A(t), \\ 0 & x \notin A(t). \end{cases}$$

It will be useful to have coordinates defined on a fixed domain \mathbb{T} . Therefore, we will not use arc-length, as the length of the curve may not be preserved in time. Instead, we use a uniform speed parameterisation (also see Section 4.1) $z : \mathbb{T} \rightarrow \mathbb{R}^2$ where the speed $|\frac{dz}{ds}| = L$ is constant in s , and $L(t)$ is the length of the curve at time t . If z is positively oriented, the 2D Frenet formulas are $\frac{dT}{ds} = L\kappa N$, $\frac{dN}{ds} = -L\kappa T$, with unit tangent $T = \frac{1}{L} \frac{dz}{ds}$ and unit normal $N := T^\perp$. The curvature is $\kappa = \frac{1}{L} \frac{dT}{ds} \cdot N = L^{-3} z_{ss} \cdot z_s^\perp$. Integration along z will be written as

$$\int_{z(\mathbb{T})} f(z) \, dl := L \int_{\mathbb{T}} f(z(s)) \, ds.$$

A similar argument to [13] yields a rigorous derivation of the evolution equation for z .

Proposition 3.3 (Evolution of a sharp front). *If $\theta = \mathbf{1}_A$ is a sharp front solution to (3.1), then the uniform speed counter-clockwise parameterisation $z : \mathbb{T} \rightarrow \mathbb{R}^2$ of ∂A with normal $N = \partial_s z^\perp$ satisfies the*

following CDE,

$$\partial_t z \cdot N = \left(- \int_{\mathbb{T}} K(z - z_*) (\partial_s z_* - \partial_s z) ds_* \right) \cdot N =: -\mathcal{I}(z) \cdot N. \quad (3.2)$$

4. ALMOST-SHARP FRONTS

In this chapter, we define an almost-sharp front as a regularisation (in a small strip) of a sharp front. Then we derive an asymptotic equation for an almost-sharp front, using an asymptotic lemma for a parameterised family of integrals. We also show that integrating across the transition region has a regularising effect, simplifying the asymptotic equation of an almost-sharp front.

Definition 4.1 (Almost-sharp front). An almost-sharp front (ASF) $\theta = \theta_\delta(x, t)$ to SQG is a family of solutions to (3.1), such that for each $\delta > 0$ sufficiently small, there exists a closed C^2 curve z bounding a simply connected region A , and a constant C^z with $0 < C^z < C$ for some fixed C independent of δ , such that with the following sets,

$$\begin{aligned} A_{\text{mid}}(t) &= \{x \in \mathbb{R}^2 : \text{dist}(x, \partial A(t)) \leq C^z \delta\}, \\ A_{\text{in}}(t) &= \{x \in A : \text{dist}(x, A(t)) > C^z \delta\}, \\ A_{\text{out}}(t) &= \mathbb{R}^2 \setminus (A_{\text{in}} \cup A_{\text{mid}}), \end{aligned}$$

θ is a C^2 function in \mathbb{R}^2 (as a function of x) such that

$$\theta(x, t) = \begin{cases} 1 & x \in A_{\text{in}}(t), \\ C^2 \text{ smooth} & x \in A_{\text{mid}}(t), \\ 0 & x \in A_{\text{out}}(t). \end{cases}$$

We also demand the following growth condition for θ_δ ,

$$\|\nabla \theta\|_{L^\infty} \lesssim \frac{1}{\delta}. \quad (4.1)$$

In particular θ is locally constant in $A_{\text{out}}(t) \cup A_{\text{in}}(t)$, and

$$\text{supp } \theta \subset A_{\text{mid}}(t) \cup A_{\text{in}}(t), \quad \text{supp } \nabla \theta(\cdot, t) \subset A_{\text{mid}}(t).$$

Note that the curve z in the above definition is not unique.

Definition 4.2 (Compatible curve). Any curve z satisfying the above definition for an ASF θ is called a compatible curve for θ .

4.1. Tubular neighbourhood coordinates. Given a curve $z(s, t)$ with uniform speed (denoted by $L(t)$), we define the tubular neighbourhood coordinates around z (for $\delta \ll 1$) by (we suppress the explicit dependence on t)

$$x(s, \xi) = z(s) + \delta \xi N(s), \quad s \in \mathbb{T}, \xi \in [-1, 1]. \quad (4.2)$$

We will denote by $L_1 := L(1 - \delta \kappa \xi)$, where as before κ is the curvature of z , and $L(t)$ is the length of the curve (which agrees with the uniform speed of the parameterisation). With s, ξ as above and setting $\tau = t$, then elementary calculations yield

$$\partial_{x^1} = \frac{T^1}{L_1} \partial_s - \frac{T^2}{\delta} \partial_\xi, \quad \partial_{x^2} = \frac{T^2}{L_1} \partial_s + \frac{T^1}{\delta} \partial_\xi, \quad \partial_t = A_0^1 \partial_s + A_0^2 \partial_\xi + \partial_\tau,$$

where $A_0 = \begin{pmatrix} A_0^1 \\ A_0^2 \end{pmatrix} = \begin{pmatrix} -x_\tau \cdot T / L_1 \\ -x_\tau \cdot N / \delta \end{pmatrix}$, and dx under the change of variables is $dx^1 dx^2 = \left| \det \frac{d(x^1, x^2, t)}{d(s, \xi, \tau)} \right| ds d\xi = L_1 \delta ds d\xi$.

4.2. Equation in tubular neighbourhood coordinates. Define Ω as θ expressed in the tubular neighbourhood coordinates, $\Omega(s, \xi, \tau) = \theta(x, t)$. Then the gradient of θ and its perpendicular can be written as

$$\nabla \theta = T \frac{\Omega_s}{L_1} + N \frac{\Omega_\xi}{\delta}, \quad \nabla^\perp \theta = N \frac{\Omega_s}{L_1} - T \frac{\Omega_\xi}{\delta}.$$

Since $u(x, t) = \int_{\mathbb{R}^2} \frac{\nabla^\perp \theta(x_*)}{|x - x_*|^{1+\alpha}} dx_*$ from (1.1) we obtain the following expression for u in terms of Ω (using $x = x(s, \xi, \tau)$ and $x_* = x(s_*, \xi_*, \tau)$),

$$u(x) = \iint_{\mathbb{T} \times [-1, 1]} \frac{\delta N_* \Omega_{s_*} - L_{1*} T_* \Omega_{\xi_*}}{|x - x_*|^{1+\alpha}} d\xi_* ds_*.$$

Then since

$$\begin{aligned} \nabla \theta(x) \cdot \nabla^\perp \theta(x_*) &= N_* \cdot T \frac{\Omega_{s_*} \Omega_s}{L_{1*} L_1} + N_* \cdot N \frac{\Omega_{s_*} \Omega_\xi}{\delta L_{1*}} \\ &\quad - T_* \cdot T \frac{\Omega_{\xi_*} \Omega_s}{\delta L_1} - T_* \cdot N \frac{\Omega_{\xi_*} \Omega_\xi}{\delta^2}, \end{aligned}$$

we can write the $u \cdot \nabla \theta$ term in the equation (3.1) as

$$\begin{aligned} u \cdot \nabla \theta(x) &= \iint_{\mathbb{T} \times [-1, 1]} \frac{1}{|x - x_*|^{1+\alpha}} \left(N_* \cdot T \frac{\delta \Omega_{s_*} \Omega_s}{L_1} + N_* \cdot N \Omega_{s_*} \Omega_\xi \right. \\ &\quad \left. - T_* \cdot T \frac{L_{1*} \Omega_{\xi_*} \Omega_s}{L_1} - T_* \cdot N \frac{L_{1*} \Omega_{\xi_*} \Omega_\xi}{\delta} \right) d\xi_* ds_*. \end{aligned}$$

Therefore, using $\frac{L_1}{L} = 1 - \frac{\delta(\xi_*\kappa_* - \xi\kappa)}{1 - \delta\kappa\xi}$, the equation (3.1) can be written in the new coordinates as (using $\nabla^\perp \Omega_* \cdot \nabla \Omega = \Omega_{s*} \Omega_\xi - \Omega_{\xi*} \Omega_s$)

$$\begin{aligned}
0 = & \Omega_\tau + \Omega_s \left(A_0^1 + \iint_{\mathbb{T} \times [-1,1]} \frac{\delta N_* \cdot T}{L_1 |x_* - x|^{1+\alpha}} \Omega_{s*} d\xi_* ds_* \right. \\
& + \iint_{\mathbb{T} \times [-1,1]} \frac{\delta(\xi_*\kappa_* - \xi\kappa) T_* \cdot T}{(1 - \delta\xi\kappa) |x_* - x|^{1+\alpha}} \Omega_{\xi*} d\xi_* ds_* \Big) \\
& + \left(\iint_{\mathbb{T} \times [-1,1]} \frac{T_* \cdot T}{|x_* - x|^{1+\alpha}} \nabla^\perp \Omega_* d\xi_* ds_* \right) \cdot \nabla \Omega \\
& + \frac{1}{\delta} \Omega_\xi \left(\iint_{\mathbb{T} \times [-1,1]} \frac{-L_1 T_* \cdot N}{|x_* - x|^{1+\alpha}} \Omega_{\xi*} d\xi_* ds_* - z_\tau \cdot N \right). \quad (4.2)
\end{aligned}$$

4.3. An approximate equation for an almost-sharp front. We rewrite (4.2) as (this defines the terms I_i)

$$0 = \Omega_\tau + \Omega_s (A_0^1 + \delta I_1 + \delta I_2) + I_3 \cdot \nabla \Omega + \frac{\Omega_\xi}{\delta} (I_4 - z_\tau \cdot N), \quad (4.3)$$

where $A_0^1 = \frac{-z_\tau \cdot T}{L} + O(\delta)$, and aside from explicit dependencies, I_i depends on δ through the coordinate function $x = z + \delta\xi N$ in (4.2).

We now want to find the leading order behaviour in $\delta \ll 1$. Asymptotic expansions for the integrals in (4.2) will be obtained using Lemma A.1 and Corollary A.2, which are deferred to Appendix A.

Theorem 4.3 (Approximate Equation for an ASF). *Given Ω , a δ -ASF solution of the generalised SQG equation (in the sense of Definition 4.1, tubular neighbourhood of the z) we have*

$$\begin{aligned}
o(1) = & \partial_\tau \Omega - \frac{z_\tau \cdot T}{L} \partial_s \Omega \\
& + (2 + 2\alpha) L \int_{-1}^1 \int_{\mathbb{T}} \frac{T_* \cdot N}{|z - z_*|^{3+\alpha}} (z - z_*) \cdot (\xi N - \xi_* N_*) \partial_\xi \Omega_* ds_* d\xi_* \partial_\xi \Omega \\
& + \int_{-1}^1 \int_{\mathbb{T}} \frac{L \kappa_* T_* \cdot N}{|z - z_*|^{1+\alpha}} \xi_* \partial_\xi \Omega_* ds_* d\xi_* \partial_\xi \Omega \\
& + \frac{C_{1,\alpha} \delta^{-\alpha}}{L} \int_{-1}^1 \frac{\nabla \Omega_*^\perp|_{s_*=s}}{|\xi - \xi_*|^\alpha} d\xi_* \cdot \nabla \Omega + \frac{C_{2,\alpha}}{L^{1+\alpha}} \int_{-1}^1 \nabla \Omega^\perp|_{s_*=s} d\xi_* \cdot \nabla \Omega \\
& + \int_{-1}^1 \int_{\mathbb{T}} \frac{T_* \cdot T \nabla^\perp \Omega_* \cdot \nabla \Omega}{|z - z_*|^{(1+\alpha)/2}} - \frac{\nabla^\perp \Omega_*|_{s_*=s} \cdot \nabla \Omega}{L^{1+\alpha} |\sin(s - s_*)|^{1+\alpha}} ds_* d\xi_*. \quad (4.4)
\end{aligned}$$

Remark 4.4. In (4.4), the integral term

$$(2 + 2\alpha)L \int_{-1}^1 \int_{\mathbb{T}} \frac{T_* \cdot N}{|z - z_*|^{3+\alpha}} (z - z_*) \cdot (\xi N - \xi_* N_*) \partial_\xi \Omega_* \, ds_* \, d\xi_* \, \partial_\xi \Omega$$

is well defined, despite the strength of the singularity in the denominator, since the dot product introduces a cancellation as long as say, $z \in C^3(\mathbb{T})$.

Proof of Theorem 4.3. To use the lemma, we will perform an approximation of the integral in s_* for fixed ξ, ξ_* , and s . For given s, ξ, ξ_* , when applying Lemma A.1, we will write $a_* = a(s_*) = a(s, s_*, \xi, \xi_*)$, and $a = a(s) = a(s, s, \xi, \xi_*)$.

We rewrite I_1, I_2, I_3, I_4 in (4.3) in a form suitable for Lemma A.1. We have $x = z + \xi \delta N$, and $x_* = z + \xi_* \delta N_*$ and so

$$\begin{aligned} x_{s_*} &= \partial_{s_*} x_* = L(1 - \xi_* \delta \kappa_*) T_*, \\ \tau^2 &= \delta^2 (\xi_* - \xi)^2, \\ g(s_*) &= |x_* - x|^2 - \tau^2, \\ g'(s_*) &= 2x_{s_*} \cdot x_* - 2x_{s_*} \cdot x = 2x_{s_*} \cdot (x_* - x), \\ g''(s_*) &= 2|x_{s_*}|^2 + 2x_{s_*} \cdot (x_* - x) \\ &= 2L^2(1 - \xi_* \delta \kappa_*)^2 + 2x_{s_*} \cdot (x_* - x). \end{aligned}$$

Since (as a function of s_* with s, ξ, ξ_* fixed) $g'(s) = 0$ and $g''(s) > 0$ for $\delta \ll 1$, g has a non-degenerate minimum at $s_* = s$, with $g(s) = \tau^2 - \tau^2 = 0$. As the curve z has no self-intersections, this is the unique minimum. Therefore we have (as in Lemma A.1)

$$G(\delta) = \sqrt{\frac{1}{2} g''(s)} = L(1 - \delta \kappa \xi_*).$$

(Note that $G \neq L_1$, defined earlier as $L(1 - \delta \kappa \xi)$.) For $a_* = a(s_*)$, Lemma A.1 gives the following expansion for $I = \int_{\mathbb{T}} \frac{a_*}{|g_* + \tau^2|^{(1+\alpha)/2}} \, ds_*$,

$$\begin{aligned} I &= \frac{a(s)}{G} C_{1,\alpha} \tau^{-\alpha} + \frac{a(s) C_{2,\alpha}}{G^{1+\alpha}} \\ &\quad + \int_{\mathbb{T}} \frac{a_*}{|g_*|^{(1+\alpha)/2}} - \frac{a(s)}{G^{1+\alpha} |\text{Sin}(s_* - s)|^{1+\alpha}} \, ds_* + O(\tau^{2-\alpha}), \quad \tau \rightarrow 0, \end{aligned}$$

where Sin is the rescaled $\text{Sin}(s) := \sin(\pi s)/\pi$, and $C_{1,\alpha}, C_{2,\alpha}$ are two known constants. Also, with $g_* = g(s_*, \delta)$, $G = G(\delta)$,

$$\begin{aligned} \partial_\delta g(s_*) &= 2\partial_\delta (x_* - x) \cdot (x_* - x) - 2\delta(\xi_* - \xi) \\ &= 2(\xi_* N_* - \xi N) \cdot (z_* - z) + 2\delta|\xi_* N_* - \xi N|^2 - 2\delta|\xi_* - \xi|^2, \end{aligned}$$

$$\partial_\delta G = -L\kappa\xi_*,$$

so that

$$\begin{aligned}\partial_\delta g(s_*, 0) &= 2(\xi_* N_* - \xi N) \cdot (z_* - z), \\ \partial_\delta(G^{-1-\alpha})(\delta) &= (-1 - \alpha)G^{-2-\alpha}\partial_\delta G = \frac{(1 + \alpha)}{L^{1+\alpha}(1 - \xi_* \delta \kappa)^{2+\alpha}} \kappa \xi_*.\end{aligned}$$

Corollary A.2 in this setting gives

$$\begin{aligned}I &= \frac{a(s)}{L} C_{1,\alpha} \delta^{-\alpha} |\xi_* - \xi|^{-\alpha} + \frac{a(s) C_{2,\alpha}}{L^{1+\alpha}} \\ &\quad + \int_{\mathbb{T}} \frac{a(s_*)}{|g(s_*, 0)|^{(1+\alpha)/2}} - \frac{a(s)}{L^{1+\alpha} |\sin(s_* - s)|^{1+\alpha}} ds_* \\ &\quad + \delta \int_{\mathbb{T}} \frac{(-2 - 2\alpha)a(s)}{|g(s_*, 0)|^{(3+\alpha)/2}} (\xi_* N_* - \xi N) \cdot (z_* - z) - \frac{\partial_\delta(G^{-1-\alpha})(0)a(s)}{|\sin s|^{1+\alpha}} ds_* \\ &\quad + O(\delta^{1-\alpha}).\end{aligned}$$

The terms I_1, I_2 are simpler. For I_1 , $a(s_*) = \frac{N_* \cdot T}{L(1 - \delta \kappa \xi)} \Omega_{s*}$, with $a(s) = 0$. Lemma A.1 gives

$$\delta I_1 = \delta \int_{-1}^1 \int_{\mathbb{T}} \frac{a_*}{|g_*|^{(1+\alpha)/2}} ds_* d\xi_* + O(\delta^{2-\alpha}) = O(\delta).$$

For I_2 , $a(s_*) = \frac{(\xi_* \kappa_* - \xi \kappa) T_* \cdot T}{1 - \delta \xi \kappa} \Omega_{\xi_*}$. Thus, Lemma A.1 gives

$$\begin{aligned}\delta I_2 &= \delta^{1-\alpha} \int_{-1}^1 |\xi - \xi_*|^{-\alpha} \frac{a}{L} C_{1,\alpha} d\xi_* + \delta \frac{a C_{2,\alpha}}{L^{1+\alpha}} \\ &\quad + \delta \int_{\mathbb{T}} \frac{a_*}{|g_*|^{(1+\alpha)/2}} - \frac{\pi^{1+\alpha} a}{L^{1+\alpha} |\sin(\pi(s_* - s))|^{1+\alpha}} ds_* + O(\delta^{1-\alpha}) \\ &= O(\delta^{1-\alpha}).\end{aligned}$$

For I_4 , since $L_{1*} = L(1 - \delta \kappa_* \xi_*)$, we write $I_4 = I_{4,1} + \delta I_{4,2}$, with

$$\begin{aligned}I_{4,1} &= \int_{-1}^1 \int_{\mathbb{T}} \frac{-LT_* \cdot N}{|x_* - x|^{1+\alpha}} \Omega_{\xi_*} d\xi_* ds_*, \\ I_{4,2} &= \int_{-1}^1 \int_{\mathbb{T}} \frac{L\kappa_* \xi_* T_* \cdot N}{|x_* - x|^{1+\alpha}} \Omega_{\xi_*} d\xi_* ds_*.\end{aligned}$$

For $I_{4,1}$, $a(s_*) = -LT_* \cdot N \Omega_{\xi_*}$, and $a(s) = 0$. Therefore, Lemma A.1 and Corollary A.2 gives (since $\partial_\delta g(s_*, 0) = L(x - x_*) \cdot (\xi N - \xi_* N_*)$)

$$I_{4,1} = -L \int_{\mathbb{T}} \frac{T_* \cdot N \int_{-1}^1 \Omega_{\xi_*} d\xi_*}{|z - z_*|^{1+\alpha}} ds_*$$

$$\begin{aligned}
& + (1 + \alpha)\delta \int_{-1}^1 \int_{\mathbb{T}} \frac{2LT_* \cdot N}{|z - z_*|^{3+\alpha}} (z - z_*) \cdot (\xi N - \xi_* N_*) \partial_\xi \Omega_* \, ds_* \, d\xi_* \\
& + O(\delta^{2-\alpha}).
\end{aligned}$$

Since we have chosen $\int_{-1}^1 \Omega_{\xi_*} \, d\xi_* = 1$, the first term is equal to the sharp front evolution term $-\mathcal{I}(z) \cdot N = z_\tau \cdot N$, and

$$\begin{aligned}
& \frac{1}{\delta} (I_{4,1} - z_\tau \cdot N) \\
& = (1 + \alpha) \int_{-1}^1 \int_{\mathbb{T}} \frac{2LT_* \cdot N}{|z - z_*|^{3+\alpha}} (z - z_*) \cdot (\xi N - \xi_* N_*) \partial_\xi \Omega_* \, ds_* \, d\xi_* + o(1).
\end{aligned}$$

For $I_{4,2}$, $a(s_*) = L\kappa_* \xi_* T_* \cdot N \Omega_{\xi_*}$. Lemma A.1 and Corollary A.2 give

$$\begin{aligned}
I_{4,2} & = \int_{-1}^1 \int_{\mathbb{T}} \frac{L\kappa_* \xi_* T_* \cdot N \Omega_{\xi_*}}{|x_* - x|^{1+\alpha}} \, d\xi_* \, ds_* \\
& = \int_{\mathbb{T}} \frac{L\kappa_* T_* \cdot N \int_{-1}^1 \xi_* \Omega_{\xi_*} \, d\xi_*}{|z_* - z|^{1+\alpha}} \, ds_* + O(\delta).
\end{aligned}$$

For I_3 , $a(s_*) = T_* \cdot T \nabla^\perp \Omega_*$. Applying Lemma A.1 gives

$$\begin{aligned}
I_3 & = \int_{-1}^1 \left(\frac{C_{1,\alpha}}{L(1 - \delta\kappa\xi_*)\delta^\alpha} \frac{\nabla^\perp \Omega_*|_{s_*=s}}{|\xi - \xi_*|^\alpha} + \frac{C_{2,\alpha}}{(L(1 - \delta\kappa\xi_*))^{1+\alpha}} \nabla^\perp \Omega_*|_{s_*=s} \right. \\
& \quad \left. + \int_{\mathbb{T}} \frac{T_* \cdot T \nabla^\perp \Omega_*}{|g_*|^{(1+\alpha)/2}} - \frac{\pi^{1+\alpha} \nabla^\perp \Omega_*|_{s_*=s}}{(L(1 - \delta\kappa\xi_*))^{1+\alpha} |\sin(\pi(s - s_*))|^{1+\alpha}} \, ds_* \right) d\xi_* \\
& + O(\delta^{1-\alpha}).
\end{aligned}$$

The remaining integral in s_* is dealt with by applying Corollary A.2,

$$\begin{aligned}
I_3 & = \frac{C_{1,\alpha}}{L\delta^\alpha} \int_{-1}^1 \frac{\nabla^\perp \Omega_*|_{s_*=s}}{|\xi - \xi_*|^\alpha} \, d\xi_* + \frac{C_{2,\alpha}}{L^{1+\alpha}} \int_{-1}^1 \nabla^\perp \Omega_*|_{s_*=s} \, d\xi_* \\
& + \int_{\mathbb{T}} \frac{T_* \cdot T \int_{-1}^1 \nabla^\perp \Omega_* \, d\xi_*}{|z - z_*|^{1+\alpha}} - \frac{\pi^{1+\alpha} \int_{-1}^1 \nabla^\perp \Omega_*|_{s_*=s} \, d\xi_*}{L^{1+\alpha} |\sin(\pi(s - s_*))|^{1+\alpha}} \, ds_* + O(\delta^{1-\alpha}),
\end{aligned}$$

completing the proof. \square

5. A REGULARISATION BY INTEGRATION ACROSS THE ALMOST-SHARP FRONT

As noticed in [7], the most singular terms in $\nabla \Omega^\perp \cdot \nabla \Omega_*|_{s_*=s}$ in the equation disappear upon integration in ξ over $[-1, 1]$, since for any even

integrable function F and two functions f, g , we have the cancellation

$$\iint_{\xi, \xi_* \in [-1, 1]} F(\xi - \xi_*) [f(\xi)g(\xi_*) - f(\xi_*)g(\xi)] \, d\xi \, d\xi_* = 0.$$

The integrals with $\nabla \Omega^\perp \cdot \nabla \Omega_*|_{s_*=s}$ are of this form, with $f = \partial_s \Omega$ and $g = \partial_\xi \Omega$, for fixed s . In particular, the term of order $\delta^{-\alpha}$ can be written in this form. This motivates shifting Ω by a constant so that

$$\Omega(s, \xi, t) = \begin{cases} -\frac{1}{2} & \xi \leq -1, \\ C^2 \text{ smooth} & \xi \in [-1, 1], \\ \frac{1}{2} & \xi \geq 1, \end{cases} \quad (5.1)$$

and we make the definition

$$h(s) := \int_{-1}^1 \Omega \, d\xi.$$

The identities $\int_{-1}^1 \xi \partial_\xi \Omega(s, \xi) \, d\xi = -h(s)$, and $\int_{-1}^1 \partial_\xi \Omega(s, \xi) \, d\xi = 1$ yield

$$\begin{aligned} o(1) &= \partial_\tau h - \frac{z_\tau \cdot T}{L} h' \\ &+ (2 + 2\alpha)L \int_{s_*} \frac{T_* \cdot N}{|z - z_*|^{3+\alpha}} (z - z_*) \cdot (hN - h_* N_*) \, ds_* \\ &+ \int_{\mathbb{T}} \frac{L\kappa_* T_* \cdot N}{|z - z_*|^{1+\alpha}} h_* \, ds_* + \int_{\mathbb{T}} \frac{T_* \cdot T(h'_* - h')}{|z - z_*|^{1+\alpha}} \, ds_*. \end{aligned}$$

Namely, in the limit $\delta \rightarrow 0$, h evolves via a linear homogenous integrodifferential equation that doesn't depend on Ω . Rewriting the ξ_* integrals in (4.4) using the two further identities

$$\int_{-1}^1 (\xi N - \xi_* N_*) \partial_\xi \Omega_* \, d\xi = (\xi N + h_* N_*), \quad \int_{-1}^1 \nabla \Omega^\perp|_{s_*=s} \, d\xi_* = \begin{pmatrix} -1 \\ h' \end{pmatrix},$$

we can treat h as an independently evolving function coupled with Ω via the following equation,

$$\begin{aligned}
o(1) = & \partial_\tau \Omega - \frac{z_\tau \cdot T}{L} \partial_s \Omega \\
& + (2 + 2\alpha)L \int_{\mathbb{T}} \frac{T_* \cdot N}{|z - z_*|^{3+\alpha}} (z - z_*) \cdot (\xi N + h_* N_*) ds_* \partial_\xi \Omega \\
& + \int_{\mathbb{T}} \frac{L \kappa_* T_* \cdot N}{|z - z_*|^{1+\alpha}} h_* ds_* \partial_\xi \Omega \\
& + \frac{C_{1,\alpha} \delta^{-\alpha}}{L} \int_{-1}^1 \frac{\nabla \Omega_*^\perp|_{s_*=s}}{|\xi - \xi_*|^\alpha} d\xi_* \cdot \nabla \Omega + \frac{C_{2,\alpha}}{L^{1+\alpha}} \begin{pmatrix} -1 \\ h' \end{pmatrix} \cdot \nabla \Omega \\
& + \int_{\mathbb{T}} \left(\frac{T_* \cdot T \begin{pmatrix} -1 \\ h'_* \end{pmatrix}}{|z - z_*|^{1+\alpha}} - \frac{\begin{pmatrix} -1 \\ h' \end{pmatrix}}{L^{1-\alpha} |\sin(s - s_*)|^{1+\alpha}} \right) ds_* \cdot \nabla \Omega.
\end{aligned}$$

Thus, using h reduces the understanding of the evolution of Ω to understanding a nonlinear system, where the main nonlinearity is in the single term $L^{-1} C_{1,\alpha} \delta^{-\alpha} \int_{-1}^1 \frac{\nabla \Omega_*^\perp|_{s_*=s}}{|\xi - \xi_*|^\alpha} d\xi_* \cdot \nabla \Omega$.

For SQG, this term can be expressed in terms of h and Ω , as a bilinear map, making the analysis completely different (see [8] for more details).

6. EVOLUTION OF COMPATIBLE CURVES

A sharp front solution to (1.1) is completely determined by the evolution of a curve. In this section, by considering how compatible curves are transported by (1.1) in the regime $\delta \ll 1$, we show that an almost-sharp front solution is approximately determined by the evolution of a compatible curve.

We will first present the main result of this section, Theorem 6.2 by relying on a fractional Leibniz rule (6.2). Lemma 6.3 replaces the more complicated (6.2) at the cost of a small loss in the error term. The proof is similar to Lemma 6.1 (see [4] - we present a short proof as the approach will be relevant for Lemma 6.3).

Lemma 6.1. *Let $0 < s < 1/2$ and suppose $A \subseteq \mathbb{R}^d$ is a bounded set with C^2 boundary in \mathbb{R}^d . Then $\mathbf{1}_A \in H^s(\mathbb{R}^d)$ with*

$$\|\Lambda^s \mathbf{1}_A\|_{L^2}^2 \lesssim_{d,s} |A|^{1-2s}.$$

Proof. We bound the Gagliardo seminorm $[\mathbf{1}_A]_{H^s}$ directly, which is known (see for instance [12]) to be equal to $\|\Lambda^s \mathbf{1}_A\|_{L^2}$ up to a constant depending on s alone. By definition,

$$[\mathbf{1}_A]_{H^s}^2 := \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{|\mathbf{1}_A(x) - \mathbf{1}_A(y)|^2}{|x - y|^{d+2s}} dy dx = 2 \int_{x \in A} \int_{y \in A^c} \frac{dy dx}{|x - y|^{d+2s}}.$$

Writing out a ‘layer cake’ decomposition (see for example [11, page 26]) with μ the $2d$ -dimensional Lebesgue measure,

$$[\mathbf{1}_A]_{H^s}^2 = \int_0^\infty \mu \left[x \in A, y \in A^c : |x - y| < \frac{1}{t^{1/(d+2s)}} \right] dt$$

From this, we have $[\mathbf{1}_A]_{H^s}^2 = \int_0^\infty m\left(\frac{1}{t^{1/(d+2s)}}\right) dt$, where $m(t)$ is

$$m(t) := \mu[x \in A, y \in A^c : |x - y| < t].$$

For any set $U \subseteq \mathbb{R}^d$, define $(U)_\varepsilon = \{x : d(x, U) < \varepsilon\}$. The change of variables $\tau = \frac{1}{t^{1/(d+2s)}}$, $dt = (d+2s)\tau^{-d-2s-1} d\tau$ yields

$$[\mathbf{1}_A]_{H^s}^2 = (2d+4s) \int_0^\infty \frac{m(\tau)}{\tau^{d+2s+1}} d\tau \lesssim_{d,s} \int_0^{\varepsilon_0} \frac{m(\tau)}{\tau^{d+2s+1}} d\tau + \int_{\varepsilon_0}^\infty \frac{m(\tau)}{\tau^{d+2s+1}} d\tau.$$

For some ε_0 to be chosen as follows. We can bound $m(\tau)$ using the following inclusion:

$$\{x \in A, y \in A^c : |x - y| < \tau\} \subset \{x \in A, y \in B_{\mathbb{R}^d}(x, \tau) : d(x, \partial A) < \tau\},$$

where $B_{\mathbb{R}^d}(x, \tau)$ is the ball around x of radius τ , which implies

$$m(\tau) \leq \mu[x \in A, y \in B_{\mathbb{R}^d}(x, \tau) : d(x, \partial A) < \tau] \lesssim |B_{\mathbb{R}^d}(0, 1)| \tau^{d+1},$$

since $(\partial A)_\tau$ is $O(\tau)$ by the C^2 regularity of the boundary. But for $\tau > |A|$, the following easier bound is better,

$$m(\tau) \leq \mu[x \in A : y \in B_{\mathbb{R}^d}(x, \tau)] \lesssim |B_{\mathbb{R}^d}(0, 1)| |A| \tau^d.$$

For the optimal bound, we choose $\varepsilon_0 = |A|$ to define I_1, I_2 , giving

$$\int_0^{\varepsilon_0} \frac{m(\tau)}{\tau^{d+2s+1}} d\tau \lesssim_d \int_0^{|A|} \frac{d\tau}{\tau^{2s}} = \frac{1}{1-2s} |A|^{1-2s} \lesssim_s |A|^{1-2s}$$

and

$$\int_{\varepsilon_0}^\infty \frac{m(\tau)}{\tau^{d+2s+1}} d\tau \lesssim_d |A| \int_{|A|}^\infty \frac{d\tau}{\tau^{2s+1}} = |A| \frac{|A|^{-2s}}{2s} \lesssim_s |A|^{1-2s}.$$

Combining these bounds gives the claimed bound on $[\mathbf{1}_A]_{H^s}^2$. \square

We now present the main result of this section.

Theorem 6.2. *Suppose that θ is an ASF solution, and z is a compatible curve as defined in (4.2). Then z satisfies the sharp front equation (in the weak sense) up to $O(\delta^{1-\alpha})$ errors,*

$$\partial_t z \cdot N = \left(- \int_{\mathbb{T}} K(z - z_*) (\partial_s z_* - \partial_s z) ds_* \right) \cdot N + O(\delta^{1-\alpha}).$$

Proof. The strategy of this proof is the same as [3]. For brevity of notation, we shall in this proof write

$$(\mathbf{1}_{\text{in}}, \mathbf{1}_{\text{mid}}, \mathbf{1}_{\text{out}}) := (\mathbf{1}_{A_{\text{in}}^z}, \mathbf{1}_{A_{\text{mid}}^z}, \mathbf{1}_{A_{\text{out}}^z}).$$

As in Proposition 3.3, the term $\iint_{\mathbb{R}^2, t} \partial_t \phi \theta$ brings out the time derivative of the C^2 boundary curve: since the set A_{mid} has measure $O(\delta)$, we see that if $z^\delta = z + \delta N$ parameterises the boundary of A_{in} ,

$$\begin{aligned} \iint_{\mathbb{R}^2 \times [0, \infty]} \partial_t \phi \theta dx dt &= \iint_{\{x \in \mathbb{R}^2, t \geq 0: x \in A_{\text{in}}(t)\}} \partial_t \phi dx dt + O(\delta) \\ &= \int_0^\infty \int_{\mathbb{T}} \partial_t(z^\delta) \nu^3 \phi(z^\delta, t) ds dt + O(\delta). \end{aligned}$$

Here, ν^3 is the third component of the outward normal ν 's as a vector in (x^1, x^2, t) -space, and

$$\begin{aligned} \partial_t(z^\delta) \nu^3 &= \partial_t(z^\delta) \cdot \partial_s((z^\delta)^\perp) = (L - \delta L \kappa) \partial_t(z + \delta N) \cdot \partial_s z^\perp \\ &= L \partial_t z \cdot N + O(\delta), \end{aligned}$$

where we used our definition of a compatible curve. Hence, as $\phi \in C^1$ and $dl = L ds$ for the uniform speed parameterised curve z , writing ∂A for the curve parameterised by z , we have

$$\iint_{\mathbb{R}^2 \times [0, \infty]} \partial_t \phi \theta dx dt = - \int_0^\infty \int_{\partial A} \phi(z) \partial_t z \cdot N dl dt + O(\delta).$$

We now treat the second term $\iint_{\mathbb{R}^2 \times [0, \infty]} u \cdot \nabla \phi \theta$: observe the following decomposition, where we have written $u = \nabla^\perp K * \theta = (\nabla^\perp K) * \theta$ as a convolution of θ with the kernel $\nabla^\perp K$, and used the bilinearity of $(f, g) \mapsto \int_{\mathbb{R}^2} \nabla^\perp K * f \cdot \nabla \phi g$, and $\theta = \theta \cdot (\mathbf{1}_{\text{in}} + \mathbf{1}_{\text{mid}} + \mathbf{1}_{\text{out}}) = \theta \mathbf{1}_{\text{mid}} + \mathbf{1}_{\text{in}}$:

$$\begin{aligned}
\int_{\mathbb{R}^2} u \cdot \nabla \phi \theta &= \int_{\mathbb{R}^2} \nabla^\perp K * \mathbf{1}_{\text{in}} \cdot \nabla \phi \mathbf{1}_{\text{in}} + \int_{\mathbb{R}^2} \nabla^\perp K * \mathbf{1}_{\text{in}} \cdot \nabla \phi \theta \mathbf{1}_{\text{mid}} \\
&\quad + \int_{\mathbb{R}^2} \nabla^\perp K * (\theta \mathbf{1}_{\text{mid}}) \cdot \nabla \phi \theta \mathbf{1}_{\text{mid}} + \int_{\mathbb{R}^2} \nabla^\perp K * (\theta \mathbf{1}_{\text{mid}}) \cdot \nabla \phi \mathbf{1}_{\text{in}} \\
&=: (\text{EVO}) + (\text{A}) + (\text{B}) + (\text{C}) \\
&= (\text{EVO}) + [(\text{A}) + (\text{B})] + [(\text{B}) + (\text{C})] - (\text{B}).
\end{aligned} \tag{6.1}$$

We will estimate separately each of the 4 terms in the last line of (6.1). Up to $O(\delta^{1-\alpha})$ errors, (EVO) will give us the evolution term, and the square-bracketed terms will use the C^ε regularity of $\theta = \theta \mathbf{1}_{\text{mid}} + \mathbf{1}_{\text{in}}$ that is not available when estimating (A) or (C) alone.

Control on $[(\text{A}) + (\text{B})]$. We proceed by splitting the kernel K ,

$$\begin{aligned}
[(\text{A}) + (\text{B})] &= \int_{\mathbb{R}^2} \nabla^\perp K * \theta \cdot \nabla \phi \mathbf{1}_{\text{mid}} \\
&= \int_{\mathbb{R}^2} ((\nabla^\perp K) \mathbf{1}_{|\cdot| > \delta}) * \theta \cdot \nabla \phi \mathbf{1}_{\text{mid}} + \int_{\mathbb{R}^2} ((\nabla^\perp K) \mathbf{1}_{|\cdot| < \delta}) * \theta \cdot \nabla \phi \mathbf{1}_{\text{mid}}
\end{aligned}$$

We note the bounds (recall that $\int_{|y| < \delta} K(y) dy = 0$)

$$\begin{aligned}
|((\nabla^\perp K) \mathbf{1}_{|\cdot| > \delta}) * \theta(x)| &\lesssim \|\theta\|_{L^\infty} \int_{r=\delta}^\infty \frac{r dr}{r^{2+\alpha}} \lesssim_{\theta, \alpha} \delta^{-\alpha}, \text{ and} \\
|((\nabla^\perp K) \mathbf{1}_{|\cdot| < \delta}) * \theta(x)| &= \left| \int_{|y| < \delta} K(y) \theta(x-y) dy \right| \\
&= \left| \int_{|y| < \delta} K(y) |y|^{\alpha'} \left(\frac{\theta(x-y) - \theta(x)}{|y|^{\alpha'}} \right) dy \right| \\
&\lesssim [\theta]_{C^{\alpha'}} \int_{r=0}^\delta \frac{r dr}{r^{2+\alpha-\alpha'}} \lesssim [\theta]_{C^{\alpha'}} \delta^{\alpha'-\alpha} = O(\delta^{-\alpha}),
\end{aligned}$$

where in the second inequality, we used the regularity assumption $[\theta]_{C^{\alpha'}} \lesssim \delta^{-\alpha'}$ for some $\alpha' > \alpha$. Hence both terms are integrals of $O(\delta^{-\alpha})$ functions that have support of size $O(\delta)$, due to the $\mathbf{1}_{\text{mid}}$ term. Therefore, $(\text{A}) + (\text{B}) = O(\delta^{1-\alpha})$.

Control on $[(\text{B}) + (\text{C})]$. Here, notice that

$$\begin{aligned}
\int_{\mathbb{R}^2} \nabla^\perp K * f \cdot g &= - \int_{\mathbb{R}^2} \partial_{x^2} K * f g^1 + \int_{\mathbb{R}^2} \partial_{x^1} K * f g^2 \\
&= - \int_{\mathbb{R}^2} f \partial_{x^2} K * g^1 + \int_{\mathbb{R}^2} f \partial_{x^1} K * g^2.
\end{aligned}$$

The important feature is that the two kernels $\partial_{x^1} K(-x), \partial_{x^2} K(-x)$ have the same $-2 - \alpha$ homogeneity as $\nabla^\perp K$, and have mean zero on the unit sphere. Hence, with $f = \theta \mathbf{1}_{\text{mid}}$ and $g = \nabla \phi \theta \in C^\epsilon$, we can repeat the proof as for [(A)+(B)], obtaining the same $O(\delta^{1-\alpha})$ estimate.

Control on (B). Writing $R = \nabla \Lambda^{-1}$ for the vector of Riesz transforms (see for instance [14]), we have $\nabla^\perp K * f = R^\perp \Lambda^\alpha f$, so that

$$\begin{aligned} |(B)| &= \left| \int_{\mathbb{R}^2} R^\perp \Lambda^{\alpha/2}(\theta \mathbf{1}_{\text{mid}}) \cdot \Lambda^{\alpha/2}(\nabla \phi \theta \mathbf{1}_{\text{mid}}) \right| \\ &\lesssim \|R^\perp \Lambda^{\alpha/2}(\theta \mathbf{1}_{\text{mid}})\|_{L^2} \|\Lambda^{\alpha/2}(\nabla \phi \theta \mathbf{1}_{\text{mid}})\|_{L^2} \\ &\lesssim \|\Lambda^{\alpha/2}(\theta \mathbf{1}_{\text{mid}})\|_{L^2} \|\Lambda^{\alpha/2}(\nabla \phi \theta \mathbf{1}_{\text{mid}})\|_{L^2}, \end{aligned}$$

where the last line uses the boundedness of $R^\perp : L^2 \rightarrow L^2$. To bound these terms, we will use the following fractional Leibniz rule of e.g. [10],

$$\|\Lambda^s(fg)\|_{L^2} \leq \|\Lambda^s f\|_{L^\infty} \|g\|_{L^2} + \|f\|_{L^\infty} \|\Lambda^s g\|_{L^2} \quad s \in (0, 1), \quad (6.2)$$

and the following easy estimate, valid for $s + \epsilon \leq 1$ that comes from bounding the following two terms separately (similarly to the earlier part of this proof) $\Lambda^s f(x) = \left(\int_{|x-y| \leq \delta} + \int_{|x-y| > \delta} \right) \frac{f(x) - f(y)}{|x-y|^{2+s}} dy$,

$$\|\Lambda^s f(x)\|_{L^\infty} \lesssim \delta^\epsilon [f]_{C^{s+\epsilon}} + \delta^{-s} \|f\|_{L^\infty}.$$

By interpolation, since $|\nabla \theta| \lesssim \frac{1}{\delta}$ (Definition 4.1), $[f]_{C^{s+\epsilon}} \lesssim \delta^{-s-\epsilon}$ for $s + \epsilon \leq 1$. Setting $s = \alpha/2$, $g = \mathbf{1}_{\text{mid}}$ and $f = \theta$ or $\nabla \phi \theta$, we obtain

$$\|\Lambda^{\alpha/2}(\theta \mathbf{1}_{\text{mid}})\|_{L^2} \lesssim \delta^{(1-\alpha)/2}, \quad \|\Lambda^{\alpha/2}(\nabla \phi \theta \mathbf{1}_{\text{mid}})\|_{L^2} \lesssim \delta^{(1-\alpha)/2}.$$

Together, these inequalities prove that $|(B)| \lesssim \delta^{1-\alpha}$.

Evolution term in (EVO). By following the proof of the analogous sharp front result (3.3) and using $z^\delta = z + \delta N$ again,

$$\begin{aligned} I_{\text{in}} &= \int_{\mathbb{T}} \phi(z^\delta) \left(\int_{\mathbb{T}} K(z^\delta - z_*^\delta) \partial_s z_*^\delta \cdot \partial_s (z_*^\delta)^\perp ds_* \right) ds \\ &= \int_{s \in I} \phi(z) \left(\int_{\mathbb{T}} K(z - z_*) (\partial_s z_* - \partial_s z) ds_* \right) \cdot N ds + O(\delta). \end{aligned}$$

This last line follows from a simple application of Mean Value Theorem, completing the proof. \square

In the above proof, we relied on a fractional Leibniz rule (6.2). The following lemma can serve as a weak replacement:

Lemma 6.3 (Hölder-Indicator Leibniz Rule). *Let $0 < s < 1/2$, $s < s'$. Suppose $A \subseteq \mathbb{R}^d$ is a bounded set with C^2 boundary and let $f \in C^{s'}(A)$. Then the extension of f by zero, $f\mathbf{1}_A$ belongs to $H^s(\mathbb{R}^d)$ with*

$$\|\Lambda^s(f\mathbf{1}_A)\|_{L^2(\mathbb{R}^d)}^2 \lesssim_{d,s} \|f\|_{L^\infty(A)}^2 |A|^{1-2s} + [f]_{C^{s'}(A)}^2 |A|^{1-\frac{2(s-s')}{d}}.$$

Proof. We again bound the Gagliardo seminorm $[f\mathbf{1}_A]_{H^s}^2$. We have

$$\begin{aligned} [f\mathbf{1}_A]_{H^s}^2 &= \left(2 \int_{A^c} \int_A + \int_A \int_A \right) \frac{|(f\mathbf{1}_A)(x) - (f\mathbf{1}_A)(y)|^2}{|x - y|^{d+2s}} dx dy \\ &= 2 \int_{A^c} \int_A \frac{|f(x)|^2}{|x - y|^{d+2s}} dx dy + \int_A \int_A \frac{|f(x) - f(y)|^2}{|x - y|^{d+2s}} dx dy. \end{aligned}$$

the proof of Lemma 6.1 (when only one of x, y is in A) yields

$$\int_{A^c} \int_A \frac{|f(x)|^2}{|x - y|^{d+2s}} dx dy \lesssim_{d,s} \|f\|_{L^\infty}^2 |A|^{1-2s},$$

but unlike Lemma 6.1, we do have contributions when $(x, y) \in A \times A$. That is, we need to obtain a bound on the integral

$$I_A := \int_A \int_A \frac{|f(x) - f(y)|^2}{|x - y|^{d+2s}} dx dy.$$

Using a layer cake decomposition again with

$$\tilde{m}(t) := \mu(x, y \in A, y \in A : |x - y| < t),$$

we can estimate I_A as follows:

$$\begin{aligned} I_A &= \int_0^\infty \mu \left[x \in A, y \in A : \frac{|f(x) - f(y)|^2}{|x - y|^{d+2s}} > t \right] dt \\ &\leq \int_0^\infty \mu \left[x \in A, y \in A : \frac{[f]_{C^{s'}}^2}{|x - y|^{d+2(s-s')}} > t \right] dt \\ &= \int_0^\infty \tilde{m} \left(([f]_{C^{s'}}^2 / t)^{\frac{1}{d+2(s-s')}} \right) dt \lesssim_{d,s,s'} [f]_{C^{s'}}^2 \int_0^\infty \frac{\tilde{m}(\tau)}{\tau^{d+2(s-s')+1}} d\tau, \end{aligned}$$

where in the last line we have changed variables $\tau = ([f]_{C^{s'}}^2 / t)^{\frac{1}{d+2(s-s')}}$ and ignored constants. Observe that by reasoning similarly to Lemma 6.1,

$$\tilde{m}(\tau) \lesssim_d \min(|A|^2, |A|\tau^d).$$

Hence, the optimal bound is obtained by splitting the integration region $\tau > 0$ into the sets $\tau \in [0, |A|^{1/d}]$ and $\tau \in [|A|^{1/d}, \infty]$, which yields

$$\begin{aligned} I_A &\lesssim [f]_{C^{s'}}^2 |A| \int_0^{|A|^{1/d}} \frac{d\tau}{\tau^{2(s-s')+1}} + [f]_{C^{s'}}^2 |A|^2 \int_{|A|^{1/d}}^\infty \frac{d\tau}{\tau^{d+2(s-s')+1}} \\ &= [f]_{C^{s'}}^2 (|A|^{1-\frac{2(s-s')}{d}} + |A|^{2-\frac{d+2(s-s')}{d}}) \lesssim [f]_{C^{s'}}^2 |A|^{1-\frac{2(s-s')}{d}}, \end{aligned}$$

so long as $s' > s$. As there is no contribution to $[f\mathbf{1}_A]_{H^s}^2$ when $(x, y) \in A^c \times A^c$, this concludes the proof. \square

Using Lemma 6.3 instead of the fractional Leibniz rule (6.2), we obtain the following weaker result.

Theorem 6.4. *Suppose that θ is an ASF solution, and z is a compatible curve as defined in (4.2). Then z satisfies the sharp front equation (in the weak sense) up to $O(\delta^{1-\alpha'})$ errors for any $\alpha' > \alpha$,*

$$\partial_t z \cdot N = \left(- \int_{\mathbb{T}} K(z - z_*) (\partial_s z_* - \partial_s z) ds_* \right) \cdot N + O(\delta^{1-\alpha'}).$$

7. THE SPINE OF AN ALMOST-SHARP FRONT

Here we introduce the concept of the spine, first considered by Felferman, Luli and Rodrigo in [5] for SQG in the periodic setting (our contours are however not graphs). To simplify the following calculations, assume without loss of generality Ω is given by (5.1).

Definition 7.1. Suppose an almost-sharp front has tubular neighbourhood coordinates (see Section 4.1) (s, ξ) for the transition region, induced by the compatible curve z .

We say that the curve \vec{S} is a spine for the almost-sharp front if \vec{S} is also a compatible curve, and there is a C^2 function of the uniform speed parameter $f = f(s)$ taking values in $[-C^z, C^z]$ such that

$$\int_{-C^z}^{C^z} (\xi_* - f(s_*)) \partial_\xi \Omega_* d\xi_* = 0,$$

or equivalently by the choice $\Omega|_{\xi=\pm C^z} = \pm 1/2$, $f(s_*) = - \int_{-C^z}^{C^z} \Omega_* d\xi_*$, and the corresponding spine is the curve \vec{S} given in (s, ξ) coordinates as $\xi = f(s)$, that is:

$$\vec{S}(s) = z(s) + \delta f(s) N(s).$$

The function f acts as a correction, so that for example, the base curve is also a spine if $f = 0$.

An immediate consequence of Definition 7.1 by integrating by parts is the following cancellation property for any constant $C \geq \|f\|_{L^\infty}$,

$$0 = \int_{\xi_* = f - C^z}^{f + C^z} \Omega_* d\xi_*, \quad (7.1)$$

where Ω is continuously extended to be constant on $|\xi| \geq C^z$ past the geometrically significant range that defines the tubular neighbourhood.

Indeed, for $D \gg 1$,

$$\int_{f-D}^{f+D} \Omega_* \, d\xi_* = \underbrace{\xi_* \Omega_* \Big|_{f-D}^{f+D}}_{=-f} - \underbrace{\int_{f-D}^{f+D} \xi_* \partial_\xi \Omega_* \, d\xi_*}_{=\int_{-C^z}^{C^z} \xi_* \partial_\xi \Omega_* \, d\xi_*} = \int_{-C^z}^{C^z} (\xi_* - f) \partial_\xi \Omega_* \, d\xi_* = 0.$$

We also have the following property.

Proposition 7.2 (Spine approximation property). *Let θ be a C^2 almost-sharp front, and let \vec{S} be the spine curve defined by the above (7.1). Then for any $\Gamma = \Gamma(x) \in C_c^2(\mathbb{R}^2)$, as $\delta \rightarrow 0$,*

$$\int_{x \in \mathbb{R}^2} \Gamma \nabla^\perp \theta \, dx = - \int_{\mathbb{T}} \Gamma(\vec{S}_*) \partial_s \vec{S}_* \, ds_* + O(\|\Gamma\|_{C^2} \delta^2). \quad (7.2)$$

We keep the $\|\Gamma\|_{C^2}$ dependence in (7.2) because we will require the use of test functions with $\|\Gamma\|_{C^2}$ that degenerate as $\delta \rightarrow 0$.

Proof. Write $\tilde{\Gamma} = \Gamma(x(s, \xi))$, with $x = z + \delta \xi N$. We are looking for the following expansion

$$\iint (\delta \partial_s \Omega_* N_* - L(1 - \delta \kappa_* \xi_*) \partial_\xi \Omega_* T_*) \tilde{\Gamma}_* \, ds_* \, d\xi_* + \int_{\mathbb{T}} \tilde{\Gamma}_* \partial_s \vec{S}_* \, ds_* \stackrel{?}{=} O(\delta^2). \quad (7.3)$$

We Taylor expand around \vec{S}_* in the direction of N_* to obtain

$$\begin{aligned} \tilde{\Gamma}_* &= \Gamma(x_*) = \Gamma(\vec{S}_*) + \nabla \Gamma(\vec{S}_*) \cdot (x_* - \vec{S}_*) + O(|x_* - \vec{S}_*|^2) \\ &= \Gamma(\vec{S}_*) + \delta \nabla \Gamma(\vec{S}_*) \cdot N_*(\xi_* - f_*) + O(\delta^2). \end{aligned}$$

Plugging into the left hand side (LHS) of (7.3) we have

LHS

$$\begin{aligned} &= \int_{\mathbb{T}} \left[\Gamma(\vec{S}_*) \left(\partial_s \vec{S}_* - T_* \int_{-1}^1 L(1 - \delta \kappa_* \xi_*) \partial_\xi \Omega_* \, d\xi_* + \delta N_* \int_{-1}^1 \partial_s \Omega_* \, d\xi_* \right) \right. \\ &\quad \left. - L \delta \nabla \Gamma(\vec{S}_*) \cdot N_* \int_{-1}^1 (\xi_* - f_*) \partial_\xi \Omega_* T_* \, d\xi_* \right] ds_* + O(\delta^2). \end{aligned} \quad (7.4)$$

The following identities (immediate from the definitions of \vec{S} and f),

$$\begin{aligned} 0 &= f'_* + \int_{-1}^1 \partial_s \Omega_* \, d\xi, \\ 0 &= \int_{-1}^1 (\xi_* - f_*) \partial_\xi \Omega_* \, d\xi, \text{ and} \end{aligned}$$

$$\begin{aligned}
\partial_s \vec{S}_* &= L(1 - \delta \kappa_* f_*) T_* + \delta f'_* N_* \\
&= L(1 - \delta \kappa_* f_*) T_* \int_{-1}^1 \partial_\xi \Omega_* d\xi + \delta f'_* N_*,
\end{aligned}$$

show that the right-hand side of (7.4) is of order δ^2 , as the terms in the square brackets vanish. \square

7.1. Evolution of a spine. Theorem 6.2 showed that any compatible curve evolves by the sharp front equation (3.2) up to an error of order $O(\delta^{1-\alpha})$. However, for the spine, we will improve this to $O(\delta^{2-\alpha})$.

Theorem 7.3. *Suppose that θ is an ASF solution. Then the spine curve \vec{S} in Definition 7.1 evolves (in the weak sense) according to the sharp front equation up to $O(\delta^{2-\alpha})$ error. That is,*

$$\partial_t \vec{S} \cdot N = \left(- \int_{s_* \in I} K(\vec{S} - \vec{S}_*)(\partial_s \vec{S}_* - \partial_s \vec{S}) ds_* \right) \cdot N + O(\delta^{2-\alpha}). \quad (7.5)$$

Proof. Without loss of generality, assume the constant for the base curve $C^z = 1$, and choose $\delta \ll 1$ so that we can extend the ξ coordinate to the range $[-3, 3]$, having a well defined neighbourhood of thickness 6δ . Recall that θ is a weak solution if for every $\Gamma \in C_c^\infty((0, \infty) \times \mathbb{R}^2)$,

$$0 = \iint_{t \geq 0, x \in \mathbb{R}^2} \theta \partial_t \Gamma + \theta(u \cdot \nabla \Gamma) dx dt. \quad (7.6)$$

Define for each time t the spine curve $\vec{S} = \vec{S}(s, t) \in A_{\text{in}}^{\vec{S}}(t)$, the inner region bounded by the closed curve $\vec{S} + 2\delta N$ with N the inward normal, the outer region $A_{\text{out}}^{\vec{S}}(t)$ bounded by $\vec{S} - 2\delta N$, and the tubular region $A_{\text{mid}}^{\vec{S}}(t)$ in the middle of radius 2δ . We give the names \vec{S}^+ and \vec{S}^- to the inner and outer boundary curves of $A_{\text{mid}}^{\vec{S}}$ respectively,

$$\vec{S}^+ := \vec{S} + 2\delta N = z + \delta(f + 2)N,$$

$$\vec{S}^- := \vec{S} - 2\delta N = z + \delta(f - 2)N.$$

We thus have for each t (up to null sets),

$$\mathbb{R}^2 = A_{\text{in}}^{\vec{S}}(t) \cup A_{\text{out}}^{\vec{S}}(t) \cup A_{\text{mid}}^{\vec{S}}(t),$$

$$A_{\text{in}}^{\vec{S}}(t) \subseteq \{x : \theta(x, t) = +1/2\},$$

$$A_{\text{out}}^{\vec{S}}(t) \subseteq \{x : \theta(x, t) = -1/2\}.$$

Also define the related partition of $\mathbb{R}^2 \times [0, \infty)$ by

$$\text{in} = \bigcup_{t \geq 0} A_{\text{in}}^{\vec{S}}(t) \times \{t\}, \quad \text{out} = \bigcup_{t \geq 0} A_{\text{out}}^{\vec{S}}(t) \times \{t\}, \quad \text{mid} = \bigcup_{t \geq 0} A_{\text{mid}}^{\vec{S}}(t) \times \{t\}.$$

We treat the two integrands $\theta \partial_t \Gamma$ and $\theta(u \cdot \nabla \Gamma)$ in (7.6) separately, with the three sets to integrate over. For the first integrand, we have

$$\left(\iint_{\text{in}} + \iint_{\text{out}} + \iint_{\text{mid}} \right) \theta \partial_t \Gamma \, dx \, dt =: I_{\text{in}} + I_{\text{out}} + I_{\text{mid}}.$$

In the tubular region, with $L_1(s, \xi) = L(1 - \delta \kappa(s) \xi) = L + O(\delta)$,

$$\begin{aligned} I_{\text{mid}} &= \iint_{t \geq 0, s \in \mathbb{T}} \int_{\xi=f-2}^{f+2} \Omega(s, \xi, t) \partial_t \Gamma(x(s, \xi), t) \delta L_1(s, \xi) \, d\xi \, ds \, dt \\ &= \iint_{t \geq 0, s \in \mathbb{T}} \int_{\xi=f-2}^{f+2} \Omega \partial_t \Gamma(x(s, \xi), t) L \delta \, d\xi \, ds \, dt + O(\delta^2). \end{aligned}$$

By the spine cancellation property (7.1) we have

$$\begin{aligned} I_{\text{mid}} &= \delta \iint_{t \geq 0, s \in \mathbb{T}} \int_{\xi=f-2}^{f+2} \Omega[\partial_t \Gamma(x, t) - \partial_t \Gamma(\vec{S}, t)] L \, d\xi \, ds \, dt + O(\delta^2) \\ &= O(\delta^2), \end{aligned}$$

since $|\partial_t \Gamma(x(s, \xi), t) - \partial_t \Gamma(\vec{S}(s), t)| \lesssim |x(s, \xi) - \vec{S}(s)| = O(\delta)$ uniformly in s and ξ . For I_{in} , we apply the Divergence theorem in 3D,

$$\begin{aligned} \iint_{(x,t) \in \text{in}} \theta \partial_t \Gamma \, dt \, dx &= \frac{1}{2} \iint_{(x,t) \in \text{in}} \nabla_{x^1, x^2, t} \cdot \begin{bmatrix} 0 \\ 0 \\ \Gamma \end{bmatrix} \, dt \, dx \\ &= \frac{1}{2} \iint_{t \geq 0, s \in \mathbb{T}} \Gamma(\vec{S}^+, t) \partial_t [\vec{S}^+] \cdot \partial_s [\vec{S}^+]^\perp \, ds \, dt. \end{aligned}$$

Similarly for I_{out} we obtain the term (note the minus sign from the opposite orientation)

$$\begin{aligned} \iint_{(x,t) \in \text{out}} \theta \partial_t \Gamma \, dt \, dx &= \frac{-1}{2} \iint_{t \geq 0, s \in \mathbb{T}} \Gamma(\vec{S}^-, t) (-\partial_t [\vec{S}^-]) \cdot \partial_s [\vec{S}^-]^\perp \, ds \, dt \\ &= \frac{1}{2} \iint_{t \geq 0, s \in \mathbb{T}} \Gamma(\vec{S}^-, t) \partial_t [\vec{S}^-] \cdot \partial_s [\vec{S}^-]^\perp \, ds \, dt. \end{aligned}$$

Since $\vec{S}^\pm = \vec{S} \pm O(\delta)$, we obtain by the approximation formula valid for C^2 functions, $f(a+b) + f(a-b) = 2f(a) + O(b^2)$, (with $O(b^2)$ constant depending on $\|f''\|_{L^\infty}$) that

$$\iint_{\text{in} \cup \text{out}} \theta \partial_t \Gamma \, dt \, dx = \iint_{t \geq 0, s \in \mathbb{T}} \Gamma(\vec{S}) \partial_t \vec{S} \cdot \partial_s \vec{S}^\perp \, ds \, dt + O(\delta^2),$$

with $O(\cdot)$ constant depending on $\|\Gamma\|_{C^2}$ and the geometry of the base curve z . We therefore obtain that $I_{\text{in}} + I_{\text{out}} + I_{\text{mid}} = \iint_{t \geq 0, x \in \mathbb{R}^2} \theta \partial_t \Gamma \, dt \, dx$

can be written as

$$I_{\text{in}} + I_{\text{out}} + I_{\text{mid}} = \iint_{t \geq 0, s \in \mathbb{T}} \Gamma(\vec{S}) \partial_t \vec{S} \cdot \partial_s \vec{S}^\perp \, ds \, dt + O(\delta^2).$$

For the second term, define $B(t)$ as the following integrand,

$$\int_{x \in \mathbb{R}^2, t \geq 0} \theta u \cdot \nabla \Gamma \, dx \, dt =: \int_{t \geq 0} B(t) \, dt.$$

We need to control $B(t)$. We symmetrise the integrand of $B(t)$ by reversing the roles of x and y , obtaining

$$2B(t) = \int_{x \in \mathbb{R}^2} \int_{y \in \mathbb{R}^2} \nabla_x^\perp |x - y|^{-1-\alpha} \cdot [\nabla \Gamma(x) - \nabla \Gamma(y)] \theta(x) \theta(y) \, dx \, dy.$$

We split $\mathbb{R}^2 \times \mathbb{R}^2 = C_0 \cup C_1 \cup C_2$, where the subscript in C_i depends on whether none of, one of, or both of x and y are in mid respectively,

$$\begin{aligned} C_0 &= (\text{mid})^c \times (\text{mid})^c, \\ C_1 &= [\text{mid} \times (\text{mid})^c] \cup [(\text{mid})^c \times \text{mid}], \\ C_2 &= \text{mid} \times \text{mid}. \end{aligned}$$

Then write $B_0(t) + B_1(t) + B_2(t) = B(t)$, where

$$B_i(t) = \frac{1}{2} \iint_{(x,y) \in C_i} \nabla_x^\perp |x - y|^{-1-\alpha} \cdot [\nabla \Gamma(x) - \nabla \Gamma(y)] \theta(x) \theta(y) \, dx \, dy,$$

B_1 is an error term. Indeed,

$$\begin{aligned} 2B_1(t) &= \iint_{(x,y) \in C_1} \nabla_x^\perp |x - y|^{-1-\alpha} \cdot [\nabla \Gamma(x) - \nabla \Gamma(y)] \theta(x) \theta(y) \, dx \, dy \\ &= \left(\iint_{x \in \text{mid}, y \in (\text{mid})^c} + \iint_{x \in (\text{mid})^c, y \in \text{mid}} \right) \nabla_x^\perp |x - y|^{-1-\alpha} \cdot [\nabla \Gamma(x) - \nabla \Gamma(y)] \theta(x) \theta(y) \, dx \, dy \\ &=: B_{11} + B_{12}. \end{aligned}$$

We show below that B_{12} is of order $O(\delta^{2-\alpha})$; B_{11} can be estimated similarly. Integrating by parts the derivative in $\nabla_x^\perp |x - y|^{-1-\alpha}$ and using the product rule $\nabla^\perp \cdot (fV) = \nabla^\perp f \cdot V + f \nabla^\perp \cdot V$, we are left with only the boundary terms because $\nabla^\perp \cdot \nabla \Gamma = 0$ and also $\nabla^\perp \theta|_{x \in (\text{mid})^c} = 0$. To get the sign on the integration by parts correct, note

$$\begin{aligned} \int_{\text{mid}^c} \nabla \cdot F(x) \, dx &= \int_{\partial \text{mid}^c} F(x) \cdot N_{\text{out}, \partial \text{mid}^c}(x) \, dx \\ &= \int_{\mathbb{T}} F(\vec{S}_*^-) \cdot (\partial_s \vec{S}_*^-)^\perp \, ds_* - \int_{\mathbb{T}} F(\vec{S}_*^+) \cdot (\partial_s \vec{S}_*^+)^\perp \, ds_* \end{aligned}$$

$$= - \sum_{\sigma=\pm 1} \int_{\mathbb{T}} F(\vec{S}_*^\sigma) \cdot (\partial_s \vec{S}_*^\sigma)^\perp \sigma \, ds_*$$

So as $\theta(\vec{S}^\sigma) = \frac{\sigma}{2}$, $\nabla^\perp \cdot F = \nabla \cdot (-F^\perp)$, and $\nabla \theta|_{\partial \text{mid}} = 0$, we have the following versions of Divergence Theorem:

$$\int_{\text{mid}^c} \nabla^\perp \cdot F(x) \theta(x) \, dx = \frac{1}{2} \sum_{\sigma=\pm 1} \int_{\mathbb{T}} F(\vec{S}_*^\sigma) \cdot (\partial_s \vec{S}_*^\sigma) \, ds_*, \quad (7.7)$$

$$\int_{\text{mid}^c} \nabla \cdot F(x) \theta(x) \, dx = -\frac{1}{2} \sum_{\sigma=\pm 1} \int_{\mathbb{T}} F(\vec{S}_*^\sigma) \cdot (\partial_s \vec{S}_*^\sigma)^\perp \, ds_*. \quad (7.8)$$

Therefore, we can write B_{12} as follows:

$$\begin{aligned} B_{12} &= \int_{\text{mid}} \theta(y) \int_{\partial \text{mid}^c} |x - y|^{-1-\alpha} [\nabla \Gamma(x) - \nabla \Gamma(y)] \cdot T_{\partial \text{mid}^c}(x) \theta(x) \, dl(x) \, dy \\ &= \int_{\text{mid}} \frac{\theta(y)}{2} \underbrace{\sum_{\sigma=\pm 1} \int_{\mathbb{T}} |\vec{S}_*^\sigma - y|^{-1-\alpha} [\nabla \Gamma(\vec{S}_*^\sigma) - \nabla \Gamma(y)] \cdot \partial_s \vec{S}_*^\sigma \, ds_*}_{=: G(y)} \, dy \end{aligned}$$

Writing the y -integral in tubular coordinates around \vec{S} , we can use the cancellation identity (7.1) to see that

$$2B_{12} = \int_{\mathbb{T}} \int_{\xi=f-2}^{f+2} \Omega(s, \xi) (G(y(s, \xi)) - G(\vec{S})) \delta L \, ds \, d\xi + O(\delta^2).$$

In what follows, we will write y for the parameterised point $y = y(s, \xi)$. If we can prove $|G(y) - G(\vec{S})| \lesssim \delta^{1-\alpha}$, this would imply that $B_{12} = O(\delta^{2-\alpha})$. We now use a smooth cut-off function $\rho_\delta(s) = \rho(s/\delta)$ with $\text{supp } \rho = [-1, 1]$, $\rho|_{[-1/2, 1/2]} = 1$ to split $G(y)$ into two parts,

$$\begin{aligned} G(y) &= \sum_{\sigma=\pm 1} \int_{\mathbb{T}} \underbrace{|\vec{S}_*^\sigma - y|^{-1-\alpha} [\nabla \Gamma(\vec{S}_*^\sigma) - \nabla \Gamma(y)] \cdot \partial_{s_*}(\vec{S}_*^\sigma)}_{\mathcal{G}(\vec{S}_*^\sigma, y)} \, ds_* \\ &= \sum_{\sigma=\pm 1} \int_{\mathbb{T}} \rho_\delta(s - s_*) \mathcal{G}(\vec{S}_*^\sigma, y) \cdot \partial_{s_*}(\vec{S}_*^\sigma) \, ds_* \\ &\quad + \sum_{\sigma=\pm 1} \int_{\mathbb{T}} (1 - \rho_\delta(s - s_*)) \mathcal{G}(\vec{S}_*^\sigma, y) \cdot \partial_{s_*}(\vec{S}_*^\sigma) \, ds_* \\ &=: G_1(y) + G_2(y). \end{aligned}$$

Note the function $\mathcal{G} = \mathcal{G}(x, y)$ defined by the above lines, with $x = \vec{S}_*^\sigma$. For G_1 , the support in s_* of $\rho_\delta(s - s_*)$ gives us the required control using

$|\mathcal{G}(x, y)| \lesssim_\Gamma |x - y|^{-\alpha}$, so that $|\mathcal{G}(\vec{S}_*^\sigma, y)| = O(|s_* - s|^{-\alpha})$. Therefore, we have the bound

$$|G_1| \leq \|\rho\|_{L^\infty} \|\partial_s \vec{S}\|_{L^\infty} \|\mathcal{G}(\vec{S}_*^\sigma, y)\|_{L^1_{s_*}[s-\delta, s+\delta]} = O_{\rho, \Gamma}(\delta^{1-\alpha}).$$

So it now suffices to study the derivative of G_2 , since

$$\begin{aligned} |G(y) - G(\vec{S})| &\lesssim |G_2(y) - G_2(\vec{S})| + O(\delta^{1-\alpha}) \\ &\lesssim \|\nabla G_2\|_{L^\infty} |y - \vec{S}| + O(\delta^{1-\alpha}) = \|\nabla G_2\|_{L^\infty} O(\delta) + O(\delta^{1-\alpha}). \end{aligned}$$

In the tubular coordinates $y = y(s, \xi)$, we have to control the two terms $\partial_s G_2(y(s, \xi))$ and $\partial_\xi G_2(y(s, \xi))/\delta$. The first term is

$$\begin{aligned} \partial_s G_2(y) &= \sum_{\sigma=\pm 1} \int_{\mathbb{T}} -\partial_{s_*} (1 - \rho_\delta(s - s_*)) \mathcal{G}(\vec{S}_*^\sigma, y) \cdot \partial_{s_*}(\vec{S}_*^\sigma) ds_* \\ &\quad + \sum_{\sigma=\pm 1} \int_{\mathbb{T}} (1 - \rho_\delta(s - s_*)) \partial_s \mathcal{G}(\vec{S}_*^\sigma, y) \cdot \partial_{s_*}(\vec{S}_*^\sigma) ds_* \\ &= \sum_{\sigma=\pm 1} \int_{\mathbb{T}} (1 - \rho_\delta(s - s_*)) (\partial_{s_*} \mathcal{G}(\vec{S}_*^\sigma, y) + \partial_s \mathcal{G}(\vec{S}_*^\sigma, y)) \cdot \partial_{s_*}(\vec{S}_*^\sigma) ds_* \\ &\quad + \sum_{\sigma=\pm 1} \int_{\mathbb{T}} (1 - \rho_\delta(s - s_*)) \mathcal{G}(\vec{S}_*^\sigma, y) \cdot \partial_{s_*}^2(\vec{S}_*^\sigma) ds_*. \end{aligned}$$

The worst term $\partial_{s_*} \mathcal{G}(\vec{S}_*^\sigma, y) + \partial_s \mathcal{G}(\vec{S}_*^\sigma, y)$ is $O(|s_* - s|^{-1-\alpha})$, and the cutoff function restricts the integration to the region $|s - s_*| \geq \delta/2$. Thus, $\partial_s G_2 = O(\delta^{-\alpha})$.

The other derivative $\frac{\partial_\xi}{\delta} G_2(y)$ is simpler since the cutoff ρ_δ does not depend on ξ ,

$$\frac{\partial_\xi}{\delta} G_2(y) = \sum_{\sigma=\pm 1} \int_{\mathbb{T}} (1 - \rho_\delta(s - s_*)) \underbrace{\frac{\partial_\xi}{\delta} \left(\frac{\nabla \Gamma(\vec{S}_*^\sigma) - \nabla \Gamma(y)}{|\vec{S}_*^\sigma - y|^{1+\alpha}} \right)}_{=: G_{2,1}(y)} \cdot \partial_{s_*} \vec{S}_*^\sigma ds_*$$

Note that $\frac{\partial_\xi}{\delta} G_{2,1}(y) = \nabla_y G_{2,1}(y) \cdot \frac{\partial_\xi}{\delta} y$, and $\frac{\partial_\xi}{\delta} y = N = O(1)$; so we have an $O(|s - s_*|^{-1-\alpha})$ integrand, integrated on the region $|s - s_*| > \delta/2$. Hence, $G_2(y) = O(\delta^{2-\alpha})$, so $B_{11}, B_{12} = O(\delta^{2-\alpha})$, and therefore

$$B_1 = O(\delta^{2-\alpha}).$$

$$2B_2(t)$$

$$= \iint_{(x,y) \in C_1} \nabla_x^\perp |x - y|^{-1-\alpha} \cdot [\nabla \Gamma(x) - \nabla \Gamma(y)] \theta(x) \theta(y) dx dy$$

$$= -\frac{1}{2} \sum_{\sigma=\pm 1} \int_{\mathbb{T}} \int_{\text{mid}} \theta(y) |\vec{S}_*^\sigma - y|^{-1-\alpha} [\nabla \Gamma(\vec{S}_*^\sigma) - \nabla \Gamma(y)] \cdot \partial_{s_*}(\vec{S}_*^\sigma) ds_* dy \quad (7.9)$$

$$- \int_{\text{mid}} \underbrace{\left(\int_{\text{mid}} |x - y|^{-1-\alpha} [\nabla \Gamma(x) - \nabla \Gamma(y)] \theta(y) dy \right)}_{=: Q(x)} \cdot \nabla^\perp \theta(x) dx \quad (7.10)$$

Above, the ∇_x^\perp never falls on $\nabla \Gamma$ due to $\nabla \cdot \nabla^\perp = 0$. For (7.9), the singularity is no worse than the one for B_1 and can be treated in exactly the same way. For (7.10), we aim to use (7.2) of Proposition 7.2, so we need to estimate $\|\nabla^2 Q\|_{C^2}$. In what follows, we concatenate vectors to denote a tensor e.g. $(UWV)_{ijk} = U_i W_j V_k$, and $(\nabla^2 F)_{ijk} = \partial_i \partial_j F_k$. Integrating by parts twice, and introducing the term $r = r(x)$,

$$\begin{aligned} \nabla^2 Q(x) &= \int_{\text{mid}} \nabla_x^2 (|x - y|^{-1-\alpha} [\nabla \Gamma(x) - \nabla \Gamma(y)]) \theta(y) dy \\ &= \int_{\text{mid}} \nabla_y^2 (|x - y|^{-1-\alpha} [\nabla \Gamma(x) - \nabla \Gamma(y)]) \theta(y) dy \end{aligned} \quad (7.11)$$

$$\begin{aligned} &+ (\nabla_x^2 - \nabla_y^2) (|x - y|^{-1-\alpha} [\nabla \Gamma(x) - \nabla \Gamma(y)]) \\ &= \int_{\partial \text{mid}} \nabla_y (|x - y|^{-1-\alpha} [\nabla \Gamma(x) - \nabla \Gamma(y)]) \theta(y) N_{\text{out}, \partial \text{mid}} dl(y) \quad (7.12) \\ &+ \int_{\text{mid}} \nabla_y (|x - y|^{-1-\alpha} [\nabla \Gamma(x) - \nabla \Gamma(y)]) \nabla \theta(y) dy + O(|x - y|^{-1-\alpha}). \end{aligned} \quad (7.13)$$

The boundary term (7.12) from integration by parts is

$$\begin{aligned} &\int_{\partial \text{mid}} \nabla_y (|x - y|^{-1-\alpha} [\nabla \Gamma(x) - \nabla \Gamma(y)]) \theta(y) N_{\text{out}, \partial \text{mid}} ds \\ &= \frac{1}{2} \sum_{\sigma=\pm 1} \int_{\mathbb{T}} \nabla_y (|x - y|^{-1-\alpha} [\nabla \Gamma(x) - \nabla \Gamma(y)]) \Big|_{y=\vec{S}_*^\sigma} (\partial_s \vec{S}_*^\sigma)^\perp ds. \end{aligned}$$

For the remaining term in (7.13), we have

$$\begin{aligned} &|\nabla_y (|x - y|^{-1-\alpha} [\nabla \Gamma(x) - \nabla \Gamma(y)])| \\ &= \left| \frac{(-1 - \alpha)(x - y)(\nabla \Gamma(x) - \nabla \Gamma(y)) + \nabla^2 \Gamma(y) |x - y|^2}{|x - y|^{3+\alpha}} \right| \\ &= \frac{(1 + \alpha) |\nabla^2 \Gamma(x)|}{|x - y|^{1+\alpha}} + O(|x - y|^{-\alpha}). \end{aligned}$$

All of these terms (7.11), (7.12), (7.13) are $O(\delta^{-\alpha})$ terms, which can be seen by using the asymptotic lemma A.1 to compute the s integral

to leading order. For instance, for (7.13), we write out the integral explicitly using the coordinates $y = z(s_*) + \delta \xi_* N(s_*)$, $x = z(s) + \delta \xi N(s)$, and the growth condition $|\nabla \theta| \lesssim \frac{1}{\delta}$ from (4.1),

$$\begin{aligned} & \left| \int_{\text{mid}} \nabla_y (|x - y|^{-1-\alpha} [\nabla \Gamma(x) - \nabla \Gamma(y)]) \nabla \theta(y) dy \right| \\ & \leq \frac{1}{\delta} (1 + \alpha) |\nabla^2 \Gamma(x)| \int_{\xi=f-2}^{f+2} \int_{\mathbb{T}} \frac{\delta L (1 - \delta \kappa_* \xi_*)}{|z - z_* + \delta(\xi N - \xi_* N_*)|^{1+\alpha}} ds_* d\xi_* \\ & = O(\delta^{-\alpha}), \end{aligned}$$

which follows by applying the asymptotic Lemma A.1. Thus, $\|\nabla^2 Q\|_{L^\infty} = O(\delta^{-\alpha})$, and Proposition 7.2 implies that

$$\begin{aligned} 2B_2(t) &= \int_{\mathbb{T}} \left(\int_{\text{mid}} |\vec{S} - y|^{-1-\alpha} [\nabla \Gamma(\vec{S}) - \nabla \Gamma(y)] \theta(y) dy \right) \cdot \partial_s \vec{S} ds \\ &\quad + O(\delta^{2-\alpha}). \end{aligned}$$

Reversing the order of integration and using the tubular coordinates we notice we can again use the spine condition (7.1) to bring out an extra cancellation. That is, defining H by

$$H(y_*) := \int_{\mathbb{T}} |\vec{S} - y_*|^{-1-\alpha} [\nabla \Gamma(\vec{S}) - \nabla \Gamma(y_*)] \cdot \partial_s \vec{S} ds,$$

we deduce that

$$\begin{aligned} 2B_2(t) &= \int_{f-2}^{f+2} \int_{\mathbb{T}} H(y_*) \Omega_* \delta L_{1*} ds_* d\xi_* + O(\delta^{2-\alpha}) \\ &= \delta \int_{f-2}^{f+2} \int_{\mathbb{T}} \Omega_* H(y_*) L ds_* d\xi_* + O(\delta^{2-\alpha}) \\ &= \delta \int_{f-2}^{f+2} \int_{\mathbb{T}} \Omega_* \left(H(y_*) - H(\vec{S}_*) \right) L ds_* d\xi_* + O(\delta^{2-\alpha}) \\ &= O(\delta^{2-\alpha}). \end{aligned}$$

The final term to estimate is

$$B_0(t) = \int_{(\text{mid})^c} \int_{(\text{mid})^c} \nabla_x^\perp |x - y|^{-1-\alpha} \cdot \nabla \Gamma(x) \theta(x) \theta(y) dx dy.$$

We use both gradients appearing in B_0 to integrate by parts (via the formulas (7.8), (7.7)), on which we obtain only boundary terms due to either of the two cancellations $\nabla^\perp \cdot \nabla = 0$ or $\nabla \theta|_{\partial \text{mid}} = 0$:

$$\begin{aligned} & B_0(t) \\ &= \int_{(\text{mid})^c} \int_{(\text{mid})^c} \nabla_x^\perp |x - y|^{-1-\alpha} \cdot \nabla \Gamma(x) \theta(x) \theta(y) dx dy \end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{2} \sum_{\sigma_1=\pm 1} \int_{(\text{mid})^c} \int_{\mathbb{T}} \nabla_x^\perp |x-y|^{-1-\alpha} \big|_{x=\vec{S}^{\sigma_1}} \cdot (\partial_s \vec{S}^{\sigma_1})^\perp \Gamma(\vec{S}^{\sigma_1}) \theta(y) \, ds \, dy \\
&= \frac{1}{2} \sum_{\sigma_1=\pm 1} \int_{(\text{mid})^c} \int_{\mathbb{T}} \nabla_y^\perp |\vec{S}^{\sigma_1} - y|^{-1-\alpha} \cdot (\partial_s \vec{S}^{\sigma_1})^\perp \Gamma(\vec{S}^{\sigma_1}) \theta(y) \, ds \, dy \\
&= \frac{1}{4} \sum_{\sigma_1=\pm 1} \sum_{\sigma_2=\pm 1} \int_{\mathbb{T}} \int_{\mathbb{T}} |\vec{S}^{\sigma_1} - \vec{S}_*^{\sigma_2}|^{-1-\alpha} \cdot (\partial_s \vec{S}^{\sigma_1})^\perp \cdot \partial_s \vec{S}_*^{\sigma_2} \Gamma(\vec{S}^{\sigma_1}) \, ds \, ds_* \\
&= -\frac{1}{4} \sum_{\sigma_1=\pm 1} \sum_{\sigma_2=\pm 1} \int_{\mathbb{T}} \int_{\mathbb{T}} |\vec{S}^{\sigma_1} - \vec{S}_*^{\sigma_2}|^{-1-\alpha} \partial_s \vec{S}^{\sigma_1} \cdot [\partial_s \vec{S}_*^{\sigma_2}]^\perp \Gamma(\vec{S}^{\sigma_1}) \, ds \, ds_* .
\end{aligned}$$

This sum of four terms will now be regrouped into two terms $B_0(t) = B_{00}(t) + B_{01}(t)$, one where $\sigma_1 = \sigma_2$ and one where $\sigma_1 = -\sigma_2$,

$$\begin{aligned}
B_{00}(t) &= -\frac{1}{4} \sum_{\sigma=\pm 1} \int_{\mathbb{T}} \int_{\mathbb{T}} |\vec{S}^\sigma - [\vec{S}^\sigma]_*|^{-1-\alpha} \partial_s \vec{S}^\sigma \cdot [\partial_s \vec{S}^\sigma]_*^\perp \Gamma(\vec{S}^\sigma) \, ds \, ds_* , \\
B_{01}(t) &= -\frac{1}{4} \sum_{\sigma=\pm 1} \int_{\mathbb{T}} \int_{\mathbb{T}} |\vec{S}^\sigma - [\vec{S}^{-\sigma}]_*|^{-1-\alpha} \partial_s \vec{S}^\sigma \cdot [\partial_s \vec{S}^{-\sigma}]_*^\perp \Gamma(\vec{S}^\sigma) \, ds \, ds_* .
\end{aligned}$$

For $B_{00}(t)$, we can use the formula $f(a+b) + f(a-b) = 2f(a) + O(b^2)$ to get that

$$B_{00}(t) = -\frac{1}{2} \int_{\mathbb{T}} \int_{\mathbb{T}} \frac{\partial_s \vec{S} \cdot \partial_s \vec{S}_*^\perp}{|\vec{S} - \vec{S}_*|^{1+\alpha}} \Gamma(\vec{S}) \, ds \, ds_* + O(\delta^2).$$

Treating $B_{01}(t) =: \mathcal{B}(\delta)$ as a function of δ , we need to prove that

$$\mathcal{B}(\delta) = \mathcal{B}(0) + O(\delta^{2-\alpha}),$$

since $\mathcal{B}(0) = -\frac{1}{2} \int_{\mathbb{T}} \int_{\mathbb{T}} \frac{\partial_s \vec{S} \cdot \partial_s \vec{S}_*^\perp}{|\vec{S} - \vec{S}_*|^{1+\alpha}} \Gamma(\vec{S}) \, ds \, ds_*$ with $B_{00}(t)$ gives the required evolution term:

$$B_{00}(t) + \mathcal{B}(0) = \int_{\mathbb{T}} \left(\int_{\mathbb{T}} \frac{\partial_s \vec{S}_* - \partial_s \vec{S}}{|\vec{S} - \vec{S}_*|^{1+\alpha}} \, ds_* \right) \cdot \partial_s \vec{S}^\perp \Gamma(\vec{S}) \, ds + O(\delta^2).$$

Symmetrizing as before, we obtain

$$\mathcal{B}(\delta) = -\frac{1}{8} \sum_{\sigma=\pm 1} \int_{\mathbb{T}} \int_{\mathbb{T}} \frac{\partial_s \vec{S}^\sigma \cdot [\partial_s \vec{S}^{-\sigma}]_*^\perp}{|\vec{S}^\sigma - [\vec{S}^{-\sigma}]_*|^{1+\alpha}} \left(\Gamma(\vec{S}^\sigma) - \Gamma([\vec{S}^{-\sigma}]_*) \right) \, ds \, ds_* .$$

Since (7.5) involves test functions supported on the curve \vec{S} , we may assume that Γ does not depend on ξ on a δ neighbourhood of \vec{S} , giving

$$\mathcal{B}(\delta) = -\frac{1}{8} \sum_{\sigma=\pm 1} \int_{\mathbb{T}} \int_{\mathbb{T}} \frac{\partial_s \vec{S}^\sigma \cdot [\partial_s \vec{S}^{-\sigma}]_*^\perp}{|\vec{S}^\sigma - [\vec{S}^{-\sigma}]_*|^{1+\alpha}} \left(\Gamma(\vec{S}) - \Gamma(\vec{S}_*) \right) \, ds \, ds_* .$$

Note that $\mathcal{B}(0)$ has the well-behaved $O(|s-s_*|^{1-\alpha})$ integrand. Recalling that $\mathcal{B}'(\tilde{\delta}) = \frac{\mathcal{B}(\tilde{\delta}) - \mathcal{B}(0)}{\tilde{\delta}}$ for some $\tilde{\delta} \in (0, \delta)$, it suffices to prove that

$$\mathcal{B}'(\delta) \stackrel{?}{=} O(\delta^{1-\alpha}),$$

since $s \mapsto s^{1-\alpha}$ is increasing for $0 < s$. On differentiation with respect to δ , a factor of σ appears, which means the sum over $\sigma = \pm 1$ becomes a symmetric difference. We expand the shorthand notation $\partial_s \vec{S}^\sigma \cdot [\partial_s \vec{S}^{-\sigma}]_*^\perp$ to find the derivative in δ ,

$$\partial_s \vec{S}^\sigma \cdot [\partial_s \vec{S}^{-\sigma}]_*^\perp = \partial_s \vec{S} \cdot \partial_s \vec{S}_*^\perp + \sigma \delta (\partial_s \vec{S} \cdot T_* + T \cdot \partial_s \vec{S}_*) + O(\delta^2).$$

Hence, its δ -derivative is some bounded function, say $E(s, s_*)$. When ∂_δ hits the kernel $|\vec{S}^\sigma - [\vec{S}^{-\sigma}]_*|^{-1-\alpha}$, we have

$$\begin{aligned} \partial_\delta |\vec{S}^\sigma - [\vec{S}^{-\sigma}]_*|^{-1-\alpha} &= (-1-\alpha) |\vec{S}^\sigma - [\vec{S}^{-\sigma}]_*|^{-3-\alpha} (\vec{S}^\sigma - [\vec{S}^{-\sigma}]_*) \cdot \partial_\delta (\vec{S}^\sigma - [\vec{S}^{-\sigma}]_*) \\ &= (-1-\alpha) |\vec{S}^\sigma - [\vec{S}^{-\sigma}]_*|^{-3-\alpha} [(\vec{S} - \vec{S}_*) \cdot \sigma(N + N_*) + O(\delta)]. \end{aligned}$$

With the cancellation from the symmetrisation in Γ , we see that we have

$$\begin{aligned} -8\mathcal{B}'(\delta) &= \sum_{\sigma=\pm 1} \sigma \int_{\mathbb{T}} \int_{\mathbb{T}} |\vec{S}^\sigma - [\vec{S}^{-\sigma}]_*|^{-1-\alpha} (\Gamma(\vec{S}) - \Gamma(\vec{S}_*)) \times \\ &\quad \left(E(s, s_*) + (-1-\alpha) \partial_s \vec{S} \cdot [\partial_s \vec{S}]_*^\perp \frac{\vec{S}^\sigma - [\vec{S}^{-\sigma}]_*}{|\vec{S}^\sigma - [\vec{S}^{-\sigma}]_*|^2} \right) ds ds_* + O(\delta). \end{aligned}$$

The factor of σ means that we can use the Mean Value Theorem in the form $f(x+\delta) - f(x-\delta) = O(\delta)$ to obtain that actually $\mathcal{B}'(\delta) = O(\delta)$, which finally implies the result. \square

8. ACKNOWLEDGEMENTS

Calvin Khor is supported by the studentship part of the ERC consolidator project n^o 616797. José L. Rodrigo is partially supported by the ERC consolidator project n^o 616797.

APPENDIX A. ASYMPTOTIC FOR A PARAMETERISED INTEGRAL

Lemma A.1. *Let $\alpha \in (0, 1)$ and $\mathbb{T} := \mathbb{R}/\mathbb{Z}$. For $\tau > 0$, let $I = I(\tau)$ denote the following family of integrals,*

$$I = \int_{s \in \mathbb{T}} \frac{a(s)}{|g(s) + \tau^2|^{(1+\alpha)/2}} ds,$$

where $a = a(s), g = g(s) \in C^\infty(\mathbb{T})$ and g has 0 as its unique global minimum at $s = 0$ that is non-degenerate, i.e. $g''(0) > 0$, $\operatorname{argmin} g = 0$, $g(0) = \min g = 0$. Then we have the asymptotic relation as $\tau \rightarrow 0$,

$$I = \frac{a(0)}{G} C_\alpha \tau^{-\alpha} - \frac{a(0)(\alpha^{-1} 2^{1+\alpha} + b_\alpha)}{G^{1+\alpha}} + \int_{s \in \mathbb{T}} \frac{a(s)}{|g(s)|^{(1+\alpha)/2}} - \frac{a(0)}{G^{1+\alpha} |\operatorname{Sin} s|^{1+\alpha}} \, ds + O(\tau^{2-\alpha}), \quad (\tau \rightarrow 0),$$

where:

- (1) $\operatorname{Sin} s := \sin(\pi s)/\pi$,
- (2) C_α is the constant $C_\alpha = \frac{\sqrt{\pi} \Gamma(\frac{\alpha}{2})}{\Gamma(\frac{\alpha+1}{2})} < \infty$ (which diverges as $\alpha \rightarrow 0$),
- (3) b_α is the constant $b_\alpha := \int_{-1/2}^{1/2} (\frac{1}{|s|^{1+\alpha}} - \frac{1}{|\operatorname{Sin} s|^{1+\alpha}}) \, ds < \infty$,
- (4) G is the constant $G := \sqrt{g''(0)/2}$, and
- (5) the $O(\tau^{2-\alpha})$ constant depends on $W^{3,\infty}$ norms of a and g .

Proof. The proof for the case $\alpha = 0$ can be found in [7]. We take $(-1/2, 1/2)$ as a fundamental domain for \mathbb{T} . We split $I = I_{\text{near}} + I_{\text{far}}$ into an integral I_{near} around the minimiser of g and I_{far} on the complement,

$$\begin{aligned} A_{\text{near}} &:= (s_-, s_+), & I_{\text{near}} &:= \int_{A_{\text{near}}} \frac{a(s)}{|g(s) + \tau^2|^{(1+\alpha)/2}} \, ds, \\ A_{\text{far}} &:= \mathbb{T} \setminus A_{\text{near}}, & I_{\text{far}} &:= \int_{A_{\text{far}}} \frac{a(s)}{|g(s) + \tau^2|^{(1+\alpha)/2}} \, ds, \end{aligned} \quad (\text{A.1})$$

where s_\pm are chosen (depending on g) sufficiently close to 0 so that we can choose new coordinates σ such that $g(s) = \sigma^2$, and that $g(s_-) = g(s_+) =: \sigma_0^2 \ll 1$. Also define $\bar{a}(\sigma)$ so that $\bar{a}(\sigma) d\sigma = a(s) ds$ in the integral. Thus $I_{\text{near}} = \int_{-\sigma_0}^{\sigma_0} \frac{\bar{a}(\sigma)}{|\sigma^2 + \tau^2|^{(1+\alpha)/2}} d\sigma = I_{\text{near},1} + I_{\text{near},2} + I_{\text{near},3}$, where

$$\begin{aligned} I_{\text{near},1} &= \bar{a}(0) \int_{-\sigma_0}^{\sigma_0} \frac{1}{|\sigma^2 + \tau^2|^{(1+\alpha)/2}} d\sigma, & I_{\text{near},2} &= \int_{-\sigma_0}^{\sigma_0} \frac{\bar{a}(\sigma) - \bar{a}(0)}{|\sigma|^{1+\alpha}} d\sigma, \\ I_{\text{near},3} &= \int_{-\sigma_0}^{\sigma_0} (\bar{a}(\sigma) - \bar{a}(0) - \bar{a}'(0)\sigma) \left(\frac{1}{|\sigma^2 + \tau^2|^{(1+\alpha)/2}} - \frac{1}{|\sigma|^{1+\alpha}} \right) d\sigma. \end{aligned}$$

By the regularity of \bar{a} , these integrals are well-defined. $I_{\text{near},1}$ is the only term that appears for a constant function $\bar{a} \equiv \bar{a}(0)$. $I_{\text{near},2}$ is bounded independent of τ . $I_{\text{near},3}$ can easily be seen to be $O(\tau^{2-\alpha})$ using

$$\left| \frac{1}{|\sigma^2 + \tau^2|^{(1+\alpha)/2}} - \frac{1}{\sigma^{1+\alpha}} \right| \leq \begin{cases} 2\sigma^{-1-\alpha} & |\sigma| \leq \tau, \\ \tau^2 \sigma^{-3-\alpha} & |\sigma| > \tau. \end{cases}$$

The first bound follows from the triangle inequality and $\frac{1}{|\sigma^2 + \tau^2|^{(1+\alpha)/2}} \leq \frac{1}{|\sigma|^{1+\alpha}}$; the second bound follows from $\frac{1+\alpha}{2} < 1$ and the Mean Value Theorem applied to $f(x) = x^{-(1+\alpha)/2}$, i.e. for some $\theta \in (0, 1)$,

$$f(x+h) - f(x) = f'(x+\theta h)h = -\left(\frac{1+\alpha}{2}\right) \frac{h}{|x+\theta h|^{(3+\alpha)/2}}, \quad (\text{A.2})$$

with $x = \sigma^2, h = \tau^2$, and $|x+\theta h|^{-(3+\alpha)/2} \leq |x|^{-(3+\alpha)/2}$. This implies

$$|I_{\text{near},3}| \lesssim_{\|a''\|_{L^\infty}} \int_{-\tau}^{\tau} 2\sigma^{2-1-\alpha} d\sigma + \int_{\tau \leq |\sigma| \leq \sigma_0} \tau^2 \sigma^{2-3-\alpha} d\sigma = O(\tau^{2-\alpha}).$$

We focus now on $I_{\text{near},1}$. Define

$$J := \int_{-\sigma_0}^{\sigma_0} \frac{1}{|\sigma^2 + \tau^2|^{(1+\alpha)/2}} d\sigma = 2\tau^{-\alpha} \int_0^{\sigma_0/\tau} \frac{1}{|\sigma^2 + 1|^{(1+\alpha)/2}} d\sigma.$$

In contrast with the $\alpha = 0$ case, the integrand is in $L^1(\mathbb{R})$, so we can easily write down the following expression with an error term,

$$\begin{aligned} J &= 2\tau^{-\alpha} \left(\int_0^\infty \frac{d\sigma}{|\sigma^2 + 1|^{(1+\alpha)/2}} - \int_{\sigma_0/\tau}^\infty \frac{d\sigma}{|\sigma^2 + 1|^{(1+\alpha)/2}} \right) \\ &= 2\tau^{-\alpha} \left(\int_0^\infty \frac{d\sigma}{|\sigma^2 + 1|^{(1+\alpha)/2}} - \int_{\sigma_0/\tau}^\infty \frac{d\sigma}{\sigma^{1+\alpha}} \right) \\ &\quad + \int_{\sigma_0/\tau}^\infty \frac{1}{\sigma^{1+\alpha}} - \frac{1}{|\sigma^2 + 1|^{(1+\alpha)/2}} d\sigma = C_\alpha \tau^{-\alpha} - \frac{2}{\alpha \sigma_0^\alpha} + \text{Rem}, \end{aligned}$$

where $C_\alpha = \int_{\mathbb{R}} \frac{d\sigma}{|\sigma^2 + 1|^{(1+\alpha)/2}} = \frac{\sqrt{\pi} \Gamma(\frac{\alpha}{2})}{\Gamma(\frac{\alpha+1}{2})} < \infty$ and the remainder term Rem satisfies (using (A.2) with $x = \sigma^2, h = 1$)

$$|\text{Rem}| \leq 2\tau^{-\alpha} \int_{\sigma_0/\tau}^\infty \frac{d\sigma}{\sigma^{3+\alpha}} = 2\tau^{-\alpha} \frac{\tau^{2+\alpha}}{(2+\alpha)\sigma_0^{2+\alpha}} = O(\tau^2), \quad \tau \rightarrow 0.$$

Hence we have that $J = C_\alpha \tau^{-\alpha} - \frac{2}{\alpha \sigma_0^\alpha} + O(\tau^2)$ as $\tau \rightarrow 0$.

Recall the transformation's defining equation $g(s) = \sigma^2$. We can write $g(s) = g''(0)s^2/2 + O(s^3)$ since $s = 0$ is a global nondegenerate minimum of g with $g(0) = 0$. Hence for $\sigma > 0$ (and therefore $s > 0$), $\sigma = s\sqrt{g''(0)/2 + O(s)} = s\sqrt{g''(0)/2} + O(s^2)$ as $s \rightarrow 0^+$, by the differentiability in h of $\sqrt{g''(0)/2 + h}$. The case $\sigma < 0$ is treated similarly, leading to

$$\sigma = \sqrt{g(s)} = \sqrt{\frac{g''(0)}{2}}s + O(s^2), \quad s \rightarrow 0.$$

and hence $\frac{d\sigma}{ds} \xrightarrow{s \rightarrow 0} \sqrt{\frac{g''(0)}{2}}$. Below, we use $G := \sqrt{\frac{g''(0)}{2}}$. Since $a(s) = \frac{d\sigma}{ds} \bar{a}(\sigma)$ we have $\bar{a}(0) = a(0)/G$, which allows us to rewrite $I_{\text{near},1}$,

$$I_{\text{near},1} = \frac{a(0)}{G} C_\alpha \tau^{-\alpha} - 2 \frac{a(0)}{\alpha G \sigma_0^\alpha} + O(\tau^2), \quad \tau \rightarrow 0.$$

Let us now treat $I_{\text{near},2}$. Let $0 < \sigma_1 \ll \sigma_0$, and let $s_{1-} < 0$, $s_{1+} > 0$ be the two unique numbers such that $g(s_{1\pm}) = \sigma_1^2$. Since $\bar{a}(\sigma) d\sigma = a(s) ds$, it is clear that $\frac{\bar{a}(\sigma) d\sigma}{|\sigma|^{1+\alpha}} = \frac{a(s) ds}{g(s)^{(1+\alpha)/2}}$. Hence, we only need to rewrite the other term of the difference $\frac{(\bar{a}(\sigma) - \bar{a}(0)) d\sigma}{|\sigma|^{1+\alpha}}$, which is $\bar{a}(0) \int_{\sigma_1}^{\sigma_0} \frac{d\sigma}{\sigma^{1+\alpha}}$. We would like to replace the integral in σ with an integral in s . Observe that as $0 < \sigma_1 < \sigma_0$ and $0 < s_{1+} < s_+$,

$$\int_{\sigma_1}^{\sigma_0} \frac{d\sigma}{\sigma^{1+\alpha}} = \frac{1}{\alpha} \left(\frac{1}{\sigma_1^\alpha} - \frac{1}{\sigma_0^\alpha} \right), \quad \text{and} \quad \int_{s_{1+}}^{s_+} \frac{ds}{s^{1+\alpha}} = \frac{1}{\alpha} \left(\frac{1}{s_{1+}^\alpha} - \frac{1}{s_+^\alpha} \right).$$

Thus, we have the following equality for any constant \tilde{C} ,

$$\int_{\sigma_1}^{\sigma_0} \frac{d\sigma}{\sigma^{1+\alpha}} = \tilde{C} \int_{s_{1+}}^{s_+} \frac{ds}{s^{1+\alpha}} + \frac{1}{\alpha} \underbrace{\left(\frac{1}{\sigma_1^\alpha} - \frac{\tilde{C}}{s_{1+}^\alpha} + \frac{\tilde{C}}{s_+^\alpha} - \frac{1}{\sigma_0^\alpha} \right)}_{\star}.$$

Treating s as a function $s = s(\sigma)$, the Inverse Function Theorem gives the asymptotic $s = G^{-1}\sigma + O(\sigma^2)$ for $\sigma \ll 1$. Setting $\tilde{C} = G^{-\alpha}$, as then the terms marked with a star \star become error terms for $\sigma_1 \ll 1$,

$$\star = \frac{1}{\sigma_1^\alpha} - \frac{1}{(Gs_{1+})^\alpha} = \frac{1}{\sigma_1^\alpha} - \frac{1}{(\sigma_1 + O(\sigma_1^2))^\alpha} \leq \frac{2O(\sigma_1^2)}{\alpha \sigma_1^{1+\alpha}} = O(\sigma_1^{1-\alpha}),$$

We can do a similar analysis for the integral $\int_{-\sigma_0}^{-\sigma_1} \frac{d\sigma}{|\sigma|^{1+\alpha}}$, yielding

$$\int_{-\sigma_0}^{-\sigma_1} \frac{d\sigma}{|\sigma|^{1+\alpha}} = \frac{1}{G^\alpha} \int_{s_-}^{s_{1-}} \frac{ds}{|s|^{1+\alpha}} + \frac{1}{\alpha} \underbrace{\left(\frac{1}{\sigma_1^\alpha} - \frac{1}{|Gs_{1-}|^\alpha} + \frac{1}{|Gs_-|^\alpha} - \frac{1}{\sigma_0^\alpha} \right)}_{=O(\sigma_1^{1-\alpha})}$$

which together yield (as $\int_{-\sigma_0}^{-\sigma_1} \frac{d\sigma}{|\sigma|^{1+\alpha}} = \int_{\sigma_1}^{\sigma_0} \frac{d\sigma}{|\sigma|^{1+\alpha}}$)

$$\begin{aligned} 2 \int_{\sigma_0}^{\sigma_1} \frac{d\sigma}{|\sigma|^{1+\alpha}} &= \int_{s_{1+}}^{s_+} \frac{ds}{G^\alpha |s|^{1+\alpha}} + \int_{s_-}^{s_{1-}} \frac{ds}{G^\alpha |s|^{1+\alpha}} \\ &\quad + \frac{1}{\alpha} \left(\frac{1}{|Gs_+|^\alpha} + \frac{1}{|Gs_-|^\alpha} - \frac{2}{|\sigma_0|^\alpha} \right) + O(\sigma_1^{1-\alpha}). \end{aligned}$$

Hence, we rewrite $I_{\text{near},2}$ as follows,

$$I_{\text{near},2} = \int_{-\sigma_0}^{\sigma_0} \frac{\bar{a}(\sigma) - \bar{a}(0)}{|\sigma|^{1+\alpha}} d\sigma$$

$$\begin{aligned}
&= \lim_{\sigma_1 \rightarrow 0} \left(\int_{\sigma_1}^{\sigma_0} \frac{\bar{a}(\sigma)}{|\sigma|^{1+\alpha}} d\sigma + \int_{-\sigma_0}^{-\sigma_1} \frac{\bar{a}(\sigma)}{|\sigma|^{1+\alpha}} d\sigma - 2 \int_{\sigma_1}^{\sigma_0} \frac{\bar{a}(0)}{|\sigma|^{1+\alpha}} d\sigma \right) \\
&= \lim_{\sigma_1 \rightarrow 0} \left(\int_{s_{1+}}^{s_+} \frac{a(s)}{|g(s)|^{1+\alpha}} - \frac{a(0)}{G^{1+\alpha}|s|^{1+\alpha}} ds \right. \\
&\quad \left. + \int_{s_-}^{s_{1-}} \frac{a(s)}{|g(s)|^{1+\alpha}} - \frac{a(0)}{G^{1+\alpha}|s|^{1+\alpha}} ds + O(\sigma_1^{1-\alpha}) \right) \\
&\quad - \frac{a(0)}{\alpha G^{1+\alpha} s_+^\alpha} - \frac{a(0)}{\alpha G^{1+\alpha} |s_-|^\alpha} + 2 \frac{a(0)}{\alpha G \sigma_0^\alpha} \\
&= \int_{s_-}^{s_+} \frac{a(s)}{|g(s)|^{1+\alpha}} - \frac{a(0)}{G^{1+\alpha}|s|^{1+\alpha}} ds \\
&\quad - \frac{a(0)}{\alpha G^{1+\alpha} s_+^\alpha} - \frac{a(0)}{\alpha G^{1+\alpha} |s_-|^\alpha} + 2 \frac{a(0)}{\alpha G \sigma_0^\alpha} + O(\sigma_1^{1-\alpha}). \tag{A.3}
\end{aligned}$$

The term $2 \frac{a(0)}{\alpha G \sigma_0^\alpha}$ here in (A.3) exactly cancels with the term with $-2 \frac{a(0)}{\alpha G \sigma_0^\alpha}$ in the equation (A.4) for $I_{\text{near},1}$. We therefore can write I_{near} as follows,

$$\begin{aligned}
I_{\text{near}} &= \frac{a(0)}{G} C_\alpha \tau^{-\alpha} + \int_{s_-}^{s_+} \frac{a(s)}{|g(s)|^{1+\alpha}} - \frac{a(0)}{G^{1+\alpha}|s|^{1+\alpha}} ds \\
&\quad - \frac{a(0)}{\alpha G^{1+\alpha} s_+^\alpha} - \frac{a(0)}{\alpha G^{1+\alpha} |s_-|^\alpha} + O(\tau^{2-\alpha}), \quad \tau \rightarrow 0. \tag{A.4}
\end{aligned}$$

To finish, we need to include I_{far} . Recall from (A.1) that $A_{\text{far}} = \mathbb{T} \setminus (s_-, s_+)$. Note that with s_\pm fixed, $g(s)^{-(1+\alpha)/2}$ is $L_s^\infty(A_{\text{far}})$, and the following error estimate holds, since $\frac{a(s)}{|g(s) + \tau^2|^{(1+\alpha)/2}}$ is smooth in $\tau \ll 1$:

$$\int_{A_{\text{far}}} \frac{a(s) ds}{|g(s) + \tau^2|^{(1+\alpha)/2}} = \int_{A_{\text{far}}} \frac{a(s) ds}{|g(s)|^{(1+\alpha)/2}} + O(\tau^2).$$

Since $\int_{A_{\text{far}}} \frac{ds}{|s|^{1+\alpha}} = \left(\int_{-1/2}^{s_-} + \int_{s_+}^{1/2} \right) \frac{ds}{|s|^{1+\alpha}} = \frac{-2^{1+\alpha}}{\alpha} + \frac{1}{\alpha s_+^\alpha} + \frac{1}{\alpha |s_-|^\alpha}$, We have

$$\begin{aligned}
I_{\text{far}} &= \int_{A_{\text{far}}} \frac{a(s)}{|g(s)|^{(1+\alpha)/2}} - \frac{a(0)}{G^{1+\alpha}|s|^{1+\alpha}} ds + \frac{a(0)}{G^{1+\alpha}} \int_{A_{\text{far}}} \frac{1}{|s|^{1+\alpha}} ds + O(\tau^2) \\
&= \int_{A_{\text{far}}} \frac{a(s)}{|g(s)|^{(1+\alpha)/2}} - \frac{a(0)}{G^{1+\alpha}|s|^{1+\alpha}} ds \\
&\quad - \frac{2^{1+\alpha} a(0)}{\alpha G^{1+\alpha}} + \frac{a(0)}{\alpha G^{1+\alpha} s_+^\alpha} + \frac{a(0)}{\alpha G^{1+\alpha} |s_-|^\alpha} + O(\tau^2). \tag{A.5}
\end{aligned}$$

The terms $\frac{a(0)}{\alpha G^{1+\alpha} s_+^\alpha} + \frac{a(0)}{\alpha G^{1+\alpha} |s_-|^\alpha}$ in (A.5) cancel $-\frac{a(0)}{\alpha G^{1+\alpha} s_+^\alpha} - \frac{a(0)}{\alpha G^{1+\alpha} |s_-|^\alpha}$ in (A.4), leaving an expression that does not depend on s_\pm . Therefore,

$$I = \frac{a(0)}{G} C_\alpha \tau^{-\alpha} - \frac{2^{1+\alpha} a(0)}{\alpha G^{1+\alpha}} + \int_{-1/2}^{1/2} \frac{a(s)}{|g(s)|^{(1+\alpha)/2}} - \frac{a(0)}{G^{1+\alpha} |s|^{1+\alpha}} ds + O(\tau^{2-\alpha}), \quad \tau \rightarrow 0.$$

Since a, g are 1-periodic functions, we rewrite this with the constant $b_\alpha := \int_{-1/2}^{1/2} \frac{1}{|s|^{1+\alpha}} - \frac{\pi^{1+\alpha}}{|\sin(\pi s)|^{1+\alpha}} ds \in \mathbb{R}$,

$$I = \frac{a(0)}{G} C_\alpha \tau^{-\alpha} - \frac{a(0)(\alpha^{-1} 2^{1+\alpha} + b_\alpha)}{G^{1+\alpha}} + \int_{s \in \mathbb{T}} \frac{a(s)}{|g(s)|^{(1+\alpha)/2}} - \frac{\pi^{1+\alpha} a(0)}{G^{1+\alpha} |\sin(\pi s)|^{1+\alpha}} ds + O(\tau^{2-\alpha}), \quad \tau \rightarrow 0,$$

as claimed. \square

Concerning the integral in Lemma A.1, we have the following result.

Corollary A.2. *Let $\mathbb{T} := \mathbb{R}/\mathbb{Z}$. For $\delta \in (-\delta_0, \delta_0)$ sufficiently small, let $H = H(\delta)$ denote the following family of integrals,*

$$H = \int_{s \in \mathbb{T}} \left(\frac{a(s)}{|g(s, \delta)|^{(1+\alpha)/2}} - \frac{a(0)}{G(\delta)^{1+\alpha} |\sin s|^{1+\alpha}} \right) ds,$$

where $a = a(s) \in C^\infty(\mathbb{T})$, $g = g(s, \delta) \in C^\infty(\mathbb{T} \times [0, \infty))$, and g has a unique global minimum that is non-degenerate with $\partial_s^2 g(0, \cdot) > c > 0$ for a constant c independent of δ , and

$$\operatorname{argmin} g(\cdot, \delta) = 0, \quad g(0, \delta) = \min g(\cdot, \delta) = 0$$

and $G(\delta) := \sqrt{\partial_s^2 g(0, \delta)/2}$. Then we have the first order Taylor expansion $H(\tau) = H(0) + H'(0)\delta + O(\delta^2)$ for $\tau \ll 1$, with

$$H'(0) = \int_{s \in \mathbb{T}} \left(\frac{a(s) \partial_\delta g(s, 0)(-1 - \alpha)}{|g(s, 0)|^{(3+\alpha)/2}} - \frac{\partial_\delta (G^{-1-\alpha})(0) a(0)}{|\sin s|^{1+\alpha}} \right) ds.$$

REFERENCES

- [1] D. Chae, P. Constantin, D. Córdoba, F. Gancedo, and J. Wu. Generalized Surface Quasi-Geostrophic equations with singular velocities. *Communications on Pure and Applied Mathematics*, 65(8):1037–1066, 2012.
- [2] A. Córdoba, D. Córdoba, and F. Gancedo. Uniqueness for SQG patch solutions. *ArXiv e-prints*, May 2016.
- [3] D. Córdoba, C. Fefferman, and J. Rodrigo. Almost sharp fronts for the Surface Quasi-Geostrophicequation. *Proceedings of the National Academy of Sciences of the United States of America*, 101(9):2687–2691, 2004.

- [4] D. Faraco and K. Rogers. The Sobolev norm of characteristic functions with applications to the Calderon inverse problem. *The Quarterly Journal of Mathematics*, 64(1):133–147, 2013.
- [5] C. Fefferman, G. Luli, and J. Rodrigo. The spine of an SQG almost-sharp front. *Nonlinearity*, 25(2):329–342, 2012.
- [6] C. Fefferman and J. Rodrigo. Analytic sharp fronts for the Surface Quasi-Geostrophic equation. *Comm. in Math. Physics*, 303(1):261–288, 2011.
- [7] C. Fefferman and J. Rodrigo. Almost sharp fronts for SQG: the limit equations. *Comm. in Math. Physics*, 313(1):131–153, 2012.
- [8] C. Fefferman and J. Rodrigo. Construction of Almost-Sharp Fronts for the Surface Quasi-Geostrophic equation. *Arch. for Rat. Mech. and Anal.*, 218(1):123 – 162, 2015.
- [9] F. Gancedo. Existence for the α -patch model and the QG sharp front in Sobolev spaces. *Advances in Mathematics*, 217(6):2569 – 2598, 2008.
- [10] C. Kenig, G. Ponce, and L. Vega. Well-posedness and scattering results for the generalized Korteweg-de Vries equation via the contraction principle. *Communications on Pure and Applied Mathematics*, 46(4):527–620, 1993.
- [11] E. Lieb and M. Loss. *Analysis*, volume 14 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2001.
- [12] E. Di Nezza, G. Palatucci, and E. Valdinoci. Hitchhiker’s guide to the fractional Sobolev spaces. *Bulletin des Sciences Mathematiques*, 136(5):521 – 573, 2012.
- [13] J. Rodrigo. The Vortex Patch problem for the Surface Quasi-Geostrophic equation. *Proc. of the National Academy of Sciences*, 101(9):2684–2686, 2004.
- [14] E. Stein. *Singular Integrals and Differentiability Properties of Functions (PMS-30)*. Princeton University Press, 1970.
- [15] J. K. Hunter, J. Shu, and Q. Zhang. Two-front solutions of the SQG equation and its generalizations. *arXiv preprint*, 2019.
- [16] J. K. Hunter, J. Shu, and Q. Zhang. Global Solutions of a Surface Quasi-Geostrophic Front Equation. *arXiv preprint*, 2018.
- [17] J. K. Hunter and J. Shu. Regularized and approximate equations for sharp fronts in the surface quasi-geostrophic equation and its generalizations. *Nonlinearity*, 31(6):2480–2517, 2018.
- [18] J. K. Hunter, J. Shu, and Q. Zhang. Local well-posedness of an approximate equation for SQG fronts, *J. Math. Fluid Mech.*, 20(4):1967–1984, 2018.
- [19] J. K. Hunter, J. Shu, and Q. Zhang. Contour dynamics for Surface Quasi-Geostrophic fronts. *arXiv preprint*, 2019.
- [20] C. Khor and J. L. Rodrigo. Local Existence of Analytic Sharp Fronts for Singular SQG. In preparation.