

Charmonium contribution to $B \rightarrow K\ell^+\ell^-$: testing the factorization approximation on the lattice

Katsumasa Nakayama^{*a,b}, Tsutomu Ishikawa^{b,c}, and Shoji Hashimoto^{b,c} (JLQCD collaboration)

^a NIC, DESY Zeuthen, Platanenallee 6, 15738 Zeuthen, Germany

^b KEK Theory Center, High Energy Accelerator Research Organization (KEK), Tsukuba 305-0801, Japan

^c School of High Energy Accelerator Science, The Graduate University for Advanced Studies (Sokendai), Tsukuba 305-0801, Japan

E-mail: katsumasa.nakayama@desy.de

We report the current status of a study of charmonium contribution to $B \rightarrow K\ell^+\ell^-$ on the lattice. Our lattice calculation tests the factorization approximation for this contribution. In order to control the problem of the artificial divergence, we focus on the low q^2 region with a small b-quark mass. We also take into account the renormalization constants of relevant four-quark operators calculated through the temporal moments. Results suggest a violation of the factorization approximation.

The 37th Annual International Symposium on Lattice Field Theory - LATTICE2019
16-22 June, 2019
Wuhan, China.

*Speaker.

1. Introduction

The rare decay $B \rightarrow K^{(*)}\ell^+\ell^-$ has received much attention as a clean probe of new physics since the Standard Model contribution is suppressed due to flavor-changing neutral-current. Sizeable difference from the Standard Model has been reported for the differential decay rate of $B \rightarrow K^{(*)}\ell^+\ell^-$ by LHCb [1, 2].

In order to confirm this tension, we have to control the uncertainty due to non-perturbative contributions. The experimental analysis of the $B \rightarrow K^{(*)}\ell^+\ell^-$ decays focused on the region where invariant mass squared of the final lepton pair q^2 is not close to the charmonium resonances. However, long-distance effects between the final state kaon and the virtual charmonium state could be significant even outside such resonance regions.

So far, theoretical estimates have been attempted by using the perturbative calculation and applying the factorization approximation, although the intermediate state can be more complex. In the factorization, we approximate the amplitude by a product of the $B \rightarrow K$ part and the charmonium resonance part. In other words, we ignore the interaction between the $B \rightarrow K$ form factor and the charmonium two-point function. The factorization approximation has been studied with experimental results and models, but reliable prediction for the $B \rightarrow K$ decay remains to be difficult [3, 4, 5, 6].

In this proceedings, we report the recent progress of the numerical lattice calculation to study the factorization approximation for the $B \rightarrow K\ell^+\ell^-$ amplitude. We calculate the $B \rightarrow K\ell^+\ell^-$ decay amplitude with and without the factorization. We take account of the renormalization constant and provide a test of the factorization approximation using an explicit lattice calculation.

2. $B \rightarrow K\ell^+\ell^-$ amplitude and the artificial divergence

In this section, we review the calculation of the decay amplitude with special emphasis on the artificial divergence. Avoiding such divergence is essential for the lattice calculation, and the problem is extensively studied for the calculation of $K \rightarrow \pi\ell^+\ell^-$ amplitude on the lattice [7, 8].

We consider the $B \rightarrow K$ amplitude with the charmonium contribution, which occurs through the weak effective Hamiltonian H_{eff} with the Fermi constant G_F , CKM matrix V_{cb}, V_{cs} , and Wilson coefficient C_i ,

$$H_{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{cs}^* V_{cb} (C_1 O_1^c + C_2 O_2^c). \quad (2.1)$$

The operators O_i^c , which include $c\bar{c}$ are

$$\begin{aligned} O_1^c &= (\bar{s}_i \gamma_\mu P_- c_j) (\bar{c}_j \gamma_\mu P_- b_i), \\ O_2^c &= (\bar{s}_i \gamma_\mu P_- c_i) (\bar{c}_j \gamma_\mu P_- b_j). \end{aligned} \quad (2.2)$$

where indices i and j represent the color index, and the chiral projection operator is defined as $P_- \equiv \frac{1-\gamma_5}{2}$.

We define the $B \rightarrow K\ell^+\ell^-$ decay amplitude for a four-momentum $q \equiv k - p$ as

$$A(q^2) = \int d^4x e^{iqx} \langle K(\mathbf{p}) | T [J_\mu(0) H_{\text{eff}}(x)] | B(\mathbf{k}) \rangle. \quad (2.3)$$




Figure 1: Setup of the lattice calculation of the $B \rightarrow K \ell^+ \ell^-$ amplitude through charmonium J/ψ resonances.

In order to calculate the amplitude, we integrate over the position of the weak effective Hamiltonian and define I_μ as

$$I_\mu = e^{-[E_K(\mathbf{p}) - E_B(\mathbf{k})]t_J} \int_{t_J - T_a}^{t_J + T_b} dt_H \int d^3 \mathbf{x} \int d^3 \mathbf{y} e^{-i\mathbf{q} \cdot \mathbf{x}} \langle K(t_K, \mathbf{p}) | T [J_\mu(t_J, \mathbf{x}) H_{\text{eff}}(t_H, \mathbf{y})] | B(0, \mathbf{k}) \rangle. \quad (2.4)$$

The setup of the lattice calculation is shown in Figure 1. We introduce t_H, t_J, t_K, T_a , and T_b to identify the time for each states and operators.

We can rewrite this quantity using the complete set of the intermediate states, which can be described by the spectral densities $\rho_1(E)$ for the states with strangeness, and $\rho_2(E)$ for those without strangeness. Namely,

$$I_\mu = - \int_0^\infty dE \frac{\rho_1(E)}{2E} \frac{\langle K(\mathbf{p}) | J_\mu(0) | E(\mathbf{k}) \rangle \langle E(\mathbf{k}) | H_{\text{eff}}(0) | B(\mathbf{k}) \rangle}{E_B(\mathbf{k}) - E} \left(1 - e^{(E_B(\mathbf{k}) - E)T_a} \right) \\ + \int_0^\infty dE \frac{\rho_2(E)}{2E} \frac{\langle K(\mathbf{p}) | H_{\text{eff}}(0) | E(\mathbf{p}) \rangle \langle E(\mathbf{p}) | J_\mu(0) | B(\mathbf{k}) \rangle}{E - E_K(\mathbf{p})} \left(1 - e^{-(E - E_K(\mathbf{p}))T_b} \right). \quad (2.5)$$

In this representation, the $T_{a,b} \rightarrow \infty$ limit of I_μ can be identified as the amplitude,

$$A(q^2) = -i \lim_{T_{a,b} \rightarrow \infty} I_\mu(T_a, T_b, \mathbf{k}, \mathbf{p}). \quad (2.6)$$

In order that the integral (2.5) stays finite, the energy of the intermediate state plays an essential role. Since $E - E_K(\mathbf{p}) > 0$ is always satisfied, $e^{-(E - E_K(\mathbf{p}))T_b}$ can be ignored in the $T_b \rightarrow \infty$ limit. On the other hand, $E_B(\mathbf{k}) - E < 0$ is not always satisfied, depending on the intermediate energy and the term $e^{(E_B(\mathbf{k}) - E)T_a}$ may diverge in the limit of large T_a . At the physical point of the quark masses, this artificial divergence can be hardly removed, since the number of such intermediate states is large. In this study, we set the b-quark mass smaller than that of the physical value in order to avoid this problem. Since the energy of the intermediate state E is bounded by the ground state energy of the K and J/ψ meson, we choose the b-quark mass to realize the condition, $E_B < E_{J/\psi} + E_K$.

With this unphysical b-quark mass, we can define the decay amplitude from the four-point correlators. In this work, however, we test the factorization approximation as the first step before proceeding to the extraction of the decay amplitude.




Figure 2: Factorization of the four-point correlator $B \rightarrow K \ell^+ \ell^-$ with charmonium J/ψ resonances.




Figure 3: A typical example of the non-factorizable contribution for $B \rightarrow K \ell^+ \ell^-$ with the charmonium. Gluon exchanging between B (K) and the charmonium can not be factorized.

3. Factorization and renormalization

In order to investigate the factorization, we define operators $O^{(1)}$ and $O^{(8)}$ as

$$\begin{aligned} O^{(1)} &= (\bar{c}_i \gamma_\mu P_- c_i) (\bar{s}_j \gamma_\mu P_- b_j), \\ O^{(8)} &= (\bar{c}_i [T^a]_{ij} \gamma_\mu P_- c_j) (\bar{s}_k [T^a]_{kl} \gamma_\mu P_- b_l). \end{aligned} \quad (3.1)$$

The operator with the color octet contraction $O^{(8)}$ includes the $SU(3)$ generators T^a .

Figure 2 illustrates the factorization of the $B \rightarrow K \ell^+ \ell^-$ four-point correlator. The contribution of $O^{(1)}$ is simply represented in the factorization approximation as,

$$\langle K J/\psi | O^{(1)} | B \rangle \simeq \langle K | \bar{s}_j \gamma_\mu P_- b_j | B \rangle \langle J/\psi | \bar{c} \gamma_\mu c | 0 \rangle. \quad (3.2)$$

Figure 3 is a typical example of the non-factorizable contribution. As we can see in the definition of $O^{(8)}$, the simple factorization is not allowed for this operator, because the factorized piece is color-octet, which vanishes when sandwiched by the physical states. Namely, factorization of the $O^{(8)}$ is represented as

$$\langle K J/\psi | O^{(8)} | B \rangle \simeq 0. \quad (3.3)$$

Since our lattice calculation is done in the O_1^c and O_2^c basis, we need to transform them to the $O^{(1)}$ and $O^{(8)}$ basis. The Firtz transformation $\bar{q}_1 \gamma_\mu P_- q_2 \bar{q}_3 \gamma_\mu P_- q_4 = \bar{q}_1 \gamma_\mu P_- q_4 \bar{q}_3 \gamma_\mu P_- q_2$ can be used to obtain the relation between O_1^c, O_2^c and $O^{(1)}, O^{(8)}$:

$$\begin{aligned} O_1^c &= O^{(1)}, \\ O_2^c &= \frac{1}{3} O^{(1)} + 2 O^{(8)}. \end{aligned} \quad (3.4)$$

β	a^{-1}	$L^3 \times T (\times L_s)$	meas.	am_{uds}	am_c	am_b
4.35	3.610(9)	$48^3 \times 96 (\times 8)$	400	0.025	0.27287	0.66619

Table 1: The parameters for our lattice calculation.

Here, we also consider the renormalization of the operator O_1^c and O_2^c . The renormalized operators $\langle O_1^c \rangle_R$ and $\langle O_2^c \rangle_R$ are written in terms of O_1^c and O_2^c with the renormalization constants Z_{11} and Z_{12} ,

$$\begin{aligned}\langle O_1^c \rangle_R &\equiv Z_{11} \langle O_1^c \rangle + Z_{12} \langle O_2^c \rangle \\ \langle O_2^c \rangle_R &\equiv Z_{12} \langle O_1^c \rangle + Z_{11} \langle O_2^c \rangle.\end{aligned}\tag{3.5}$$

The renormalization constant are determined through temporal moments of three-point correlators [9].

In order to test the factorization relation, we define the ratios R_1 and $R_{1/3}$ on the lattice of volume V ,

$$\begin{aligned}R_1 &\equiv \frac{V \langle K | J_V O_1^c | B \rangle_R}{\langle 0 | J_V J_\mu | 0 \rangle_R \langle K | \bar{s}_j \gamma_\mu P_- b_j | B \rangle_R}, \\ R_{1/3} &\equiv \frac{\langle K | J_V O_2^c | B \rangle_R}{\langle K | J_V O_1^c | B \rangle_R},\end{aligned}\tag{3.6}$$

which become 1 or $\frac{1}{3}$ when the factorization approximation is valid, respectively.

4. Preliminary results

Our lattice setup is summarized in Table 4. The lattice configurations are generated with $N_f = 2 + 1$ flavors of quarks, which are formulated by the Mobius domain-wall fermion [10]. The lattice spacing is $a^{-1} = 3.610(9)$ GeV, and each quark mass is set to $am_{uds} = 0.025$, $am_c = 0.27287$, and $am_b = 0.66619$. This choice yields the meson masses $m_\pi = 714(1)$ MeV and $am_B = 3.44(1)$ GeV. We insert two-different momenta $\mathbf{p}_{100} = (-\frac{2\pi}{L}, 0, 0)$ and $\mathbf{p}_{110} = (-\frac{2\pi}{L}, -\frac{2\pi}{L}, 0)$ for the final state of charmonium $c\bar{c}$. For the initial B meson state, we input a momentum $\mathbf{k} = (0, 0, 0)$. The momenta $\mathbf{p}_{100}$ and $\mathbf{p}_{110}$ are smaller than the physical one, but we focus on these two inputs as a first step. The energy spectrum with these input values are calculated as $E_K(\mathbf{p}_{100}) = 855(3)$ MeV, $E_K(\mathbf{p}_{110}) = 969(9)$ MeV, $E_{J/\psi}(\mathbf{p}_{100}) = 3.127(1)$ GeV, and $E_{J/\psi}(\mathbf{p}_{110}) = 3.158(1)$ GeV. As we discussed in the definition of the $B \rightarrow K$ decay amplitude, the spectrum satisfies the condition $m_B < E_{J/\psi} + E_K$. Namely, our setup does not suffer from artificial divergence, as we mentioned previously. The source operators are set at $t_K = 42$ for K meson source, $t_J = 27$ for electromagnetic coupling J_μ . The B-meson source is set at the $t = 0$. In this study, statistical uncertainty is estimated with 100 independent configurations with four different source points per configuration. Since the propagating mesons are heavy, the auto-correlation is not significant.

We use the renormalization constants determined through the moments of the corresponding three-point correlators [9]. They are $Z_{11} = 0.669(11)$ and $Z_{12} = 0.093(4)$.




Figure 4: The ratio $R_{1/3}$ are shown for each input momenta. The electromagnetic current is set at $t_I = 27$ as shown by the dashed line.




Figure 5: The ratio R_1 are shown for each input momenta. The electromagnetic current is set at $t_I = 27$ as shown by the dashed line.

Figure 4 shows the result for the ratio $R_{1/3}$, which should be equal to $1/3$ if the factorization is a good approximation. The result is almost consistent with $1/3$, and we do not see any significant violation of the approximation. On the other hand, the relation $R_1 \simeq 1$ is not satisfied, as shown in Figure 5. The size of the violation is as large as 30%. It suggests that the factorization approximation may underestimate the $\bar{c}c$ contribution to $B \rightarrow K\ell^+\ell^-$.

5. Discussions

A quantitative estimate of the $c\bar{c}$ contribution to $B \rightarrow K^{(*)}\ell^+\ell^-$ remains a notoriously difficult task, because of the non-perturbative dynamics of QCD. The first principle calculation of the lattice QCD can not be directly applied since there are many intermediate states that contribute to the real and imaginary parts of the amplitude. In this work, we simplify the problem by considering an unphysical setup with a smaller b quark mass, hoping that it captures the important part of the dynamics. We find a significant violation of the factorization ansatz, which may be used as inputs for phenomenological models to study more realistic situations. We also note that a large violation

of factorization was previously found for the $K \rightarrow \pi\pi$ amplitude [11], which suggests the need for fully non-perturbative calculation for similar processes.

Acknowledgements

The lattice QCD simulation has been performed on Blue Gene/Q supercomputer at the High Energy Accelerator Research Organization (KEK) under the Large Scale Simulation Program (Nos. 15/16-09, 16/17-14). Oakforest-PACS at JCAHPC under the support of the HPCI System Research Projects. K. N. is supported by the Grant-in-Aid for JSPS (Japan Society for the Promotion of Science) Research Fellow (No. 18J11457). This work is supported in part by the Grant-in-Aid of the Japanese Ministry of Education (No. 18H03710).

References

- [1] LHCb collaboration, R. Aaij et al., *Observation of a resonance in $B^+ \rightarrow K^+\mu^+\mu^-$ decays at low recoil*, *Phys. Rev. Lett.* **111** (2013) 112003 [1307.7595].
- [2] LHCb collaboration, R. Aaij et al., *Measurement of the phase difference between short- and long-distance amplitudes in the $B^+ \rightarrow K^+\mu^+\mu^-$ decay*, *Eur. Phys. J.* **C77** (2017) 161 [1612.06764].
- [3] M. Neubert and B. Stech, *Nonleptonic weak decays of B mesons*, *Adv. Ser. Direct. High Energy Phys.* **15** (1998) 294 [hep-ph/9705292].
- [4] M. Beneke, T. Feldmann and D. Seidel, *Systematic approach to exclusive $B \rightarrow V\ell^+\ell^-$, $V\gamma$ decays*, *Nucl. Phys.* **B612** (2001) 25 [hep-ph/0106067].
- [5] J. Lyon and R. Zwicky, *Resonances gone topsy turvy - the charm of QCD or new physics in $b \rightarrow s\ell^+\ell^-$?*, 1406.0566.
- [6] D. Du, A. X. El-Khadra, S. Gottlieb, A. S. Kronfeld, J. Laiho, E. Lunghi et al., *Phenomenology of semileptonic B-meson decays with form factors from lattice QCD*, *Phys. Rev.* **D93** (2016) 034005 [1510.02349].
- [7] RBC, UKQCD collaboration, N. H. Christ, X. Feng, A. Portelli and C. T. Sachrajda, *Prospects for a lattice computation of rare kaon decay amplitudes: $K \rightarrow \pi\ell^+\ell^-$ decays*, *Phys. Rev.* **D92** (2015) 094512 [1507.03094].
- [8] RBC, UKQCD collaboration, N. H. Christ, X. Feng, A. Portelli and C. T. Sachrajda, *Prospects for a lattice computation of rare kaon decay amplitudes II $K \rightarrow \pi\nu\bar{\nu}$ decays*, *Phys. Rev.* **D93** (2016) 114517 [1605.04442].
- [9] T. Ishikawa, K. Nakayama and S. Hashimoto, *Renormalization of bilinear and four-fermion operators through temporal moments*, *PoS LATTICE2019* **135** (2019).
- [10] R. C. Brower, H. Neff and K. Orginos, *The Möbius domain wall fermion algorithm*, *Comput. Phys. Commun.* **220** (2017) 1 [1206.5214].
- [11] RBC, UKQCD collaboration, P. A. Boyle et al., *Emerging understanding of the $\Delta I = 1/2$ Rule from Lattice QCD*, *Phys. Rev. Lett.* **110** (2013) 152001 [1212.1474].