# AGENDA-MANIPULATION IN RANKING\*

Gregorio Curello University of Bonn Ludvig Sinander University of Oxford

18 January 2022

#### Abstract

We study the susceptibility of committee governance (e.g. by boards of directors), modelled as the collective determination of a ranking of a set of alternatives, to manipulation of the order in which pairs of alternatives are voted on—agenda-manipulation. We exhibit an agenda strategy called *insertion sort* that allows a self-interested committee chair with no knowledge of how votes will be cast to do as well as if she had complete knowledge. Strategies with this 'regret-freeness' property are characterised by their *efficiency*, and by their avoidance of two intuitive errors. What distinguishes regret-free strategies from each other is how they prioritise among alternatives; insertion sort prioritises *lexicographically*.

# 1 Introduction

In many modern organisations, governance is entrusted not to a single individual, but to a committee: a group that reaches decisions by aggregating

<sup>&</sup>lt;sup>\*</sup>We are deeply grateful to Eddie Dekel for many detailed comments. We thank him, Péter Eső, Alessandro Pavan, John Quah and Bruno Strulovici for their guidance and support. We have profited from comments and suggestions provided by Georgy Egorov, Benny Moldovanu and Wojciech Olszewski (particularly fruitful), and by Nageeb Ali, Nemanja Antić, David Austen-Smith, Marco Battaglini, Steve Coate, Tommaso Denti, Francesc Dilmé, Piotr Dworczak, Jeff Ely, Matteo Escudé, Ben Golub, Sergiu Hart, Claudia Herresthal, Matt Jackson, Mathijs Janssen, Stephan Lauermann, Jiangtao Li, Shengwu Li, Elliot Lipnowski, Francisco Poggi, Marek Pycia, Sven Rady, Ariel Rubinstein, Eddie Schlee, Marciano Siniscalchi, Rani Spiegler, Francesco Squintani, Jean-Marc Tallon, Asher Wolinsky, Andrea Galeotti (as editor), four anonymous referees, and audiences at Bocconi, Bonn, Cornell, Hebrew/Tel Aviv, Hong Kong, Northwestern, Oxford, Paris, Penn State, Royal Holloway, UCL, Warwick and the 18<sup>th</sup> IO Theory Conference at UC Berkeley.

the individual votes of its members, usually via a majority or super-majority rule. Public firms and charities are run by boards of directors, for example.

To govern is to make decisions: to choose one alternative from among those that are available. These alternatives may be e.g. policies, candidates for a job, or investments, or they could be coarser things like *attributes* of policies, candidates or investments.

In practice, most governing committees meet too infrequently to be able to consider and solve each individual decision problem faced by their organisation. Lacking the ability or time to foresee which decision problems (sets of available alternatives) will arise at any given future date, the committee cannot specify what alternative ought to be chosen. Committees therefore tend to set only high-level priorities, while delegating the details of day-to-day decision-making to executive officers. Such committee governance may be captured in stylised fashion by thinking of the committee as determining a ranking of all the potentially-available alternatives, with executives instructed to choose the highest-ranked alternative from among those that turn out to be available.

As observed by Condorcet (1785), the majority will of a committee can easily fail to be transitive: there may be a majority for alternative  $\alpha$  over  $\beta$ , for  $\beta$  over  $\gamma$ , and for  $\gamma$  over  $\alpha$ . Whenever there are such (Condorcet) cycles, the order in which questions are considered by the committee will affect what decision is reached. The chair of the committee can thus influence collective decisions via her control of the agenda.

We model agenda-setting by the chair as her determining the order in which pairs of alternatives are voted on. We assume that the committee is sovereign, so that a majority vote for  $\alpha$  over  $\beta$  leads to  $\alpha$  being ranked above  $\beta$ . Suppose that  $\beta$  had already been ranked above  $\gamma$ . Since rankings are by their nature transitive, it is then necessary also that  $\alpha$  be ranked above  $\gamma$ . These are the rules: transitivity is imposed.

We assume that the chair is uncertain how voters will vote on any given pair of alternatives, so that the only way to find out whether  $\alpha$  would be at  $\beta$  or vice-versa is to offer a (binding) vote on this pair.

In this uncertain world, how much influence can the chair exert via her control of the agenda? And in doing so, what sorts of agenda-setting strategies should we expect to see the chair use?

Our first finding is that agenda-setting is surprisingly powerful. In particular, there exist *regret-free* agenda-setting strategies: no matter what the unknown voting behaviour of the committee, the ranking reached by such a strategy is not unambiguously worse (from the chair's perspective) than any other ranking that the chair could have reached had she perfectly known how voters would behave.

We prove this by exhibiting a regret-free strategy called *insertion sort*, which works as follows. Let's call the chair's  $k^{\text{th}}$ -favourite alternative simply 'k'. No votes involving k are offered until all of the alternatives that the chair considers strictly worse than k have been totally ranked. Once that has happened, insertion sort pits k against whichever of the worse alternatives is ranked highest; if k wins, then by transitivity, it is ranked above all of the worse alternatives. If not, then k is next pitted against the *second*-highestranked of the worse alternatives; if k wins, then it is ranked above all but the highest-ranked of the worse alternatives. If not, then k is pitted against the *third*-highest ranked of the worse alternatives; and so on.

Having established that regret-free agenda-setting is possible, we next ask what its qualitative properties are: that is, what do regret-free strategies have in common? We provide two characterisations of regret-free strategies. The first describes regret-free agenda-setting in terms of its outcomes: regret-free strategies are exactly those that (ex post) reach rankings that satisfy an *efficiency* property. The second is a qualitative description of behaviour: it identifies two intuitive types of *error*, and shows that avoiding these is both necessary and sufficient for regret-freeness.

Finally, we ask what it is that distinguishes one regret-free strategy from another. The answer is *prioritisation*: while one regret-free strategy focusses on getting a certain pair of alternatives ranked 'right' (i.e. the way around that the chair prefers), another strategy may prioritise another pair. We show that insertion sort is characterised by *lexicographic* prioritisation: it places the chair's favourite alternative as highly as possible, and subject to that, places her second-favourite alternative as well as possible, and so on.

#### 1.1 Related literature

We contribute to the agenda-manipulation literature initiated by Farquharson (1969) and Black (1958), which asks how a committee's choice can be influenced by varying the order in which binary questions are voted on (the *agenda*). Two classes of agenda are emphasised: the *amendment procedure* used in Anglo-Saxon and Scandinavian legislatures, and the *successive procedure* widely used in continental Europe. Under complete information, for both sincere and strategic voting, Miller (1977) and Banks (1985) characterise which alternatives an agenda-setter can induce a committee to choose using (i) amendment agendas, (ii) successive agendas, and (iii) arbitrary agendas.<sup>1,2</sup>
Extensions include super- and sub-majority voting rules (Barberà & Gerber, 2017) and random agendas (Roessler, Shelegia & Strulovici, 2018).

Specifically, this paper belongs to the literature on agenda-setting under incomplete information about voters' preferences. This literature long consisted of a single pioneering paper (Ordeshook & Palfrey, 1988), but has recently received interest from Kleiner and Moldovanu (2017), Gershkov, Moldovanu and Shi (2017, 2019, 2020) and Gershkov, Kleiner, Moldovanu and Shi (2020). We depart from these papers by considering a committee tasked not with choosing a single alternative, but rather with *ranking* all of the alternatives. We establish a link with the older literature by relating insertion sort to the amendment procedure (appendix G).

Also related is the social choice literature on ranking methods, meaning maps that assign to each (possibly cyclic) majority will a (transitive) ranking. 'Impossibility' results such as Arrow's (1950, 1951, 1963) theorem assert that certain normatively appealing properties are inconsistent. The literature beginning with Zermelo (1929), Wei (1952) and Kendall (1955) studies ranking methods with at least some attractive normative properties.<sup>3</sup> Our chair's problem can be formulated as a choice among ranking methods, but there is a feasibility constraint, and the objective reflects the chair's selfinterest. This suggests that solutions to our chair's problem will bear little relation to the normative-motivated ranking methods in the literature; we confirm this in supplemental appendix M.

Both the voting and social choice literatures presume the existence of Condorcet cycles. Cycles are certainly a priori plausible.<sup>4</sup> And they are empirically common, especially when there are few voters, as on a committee.<sup>5</sup>

Regret-based criteria for evaluating strategies appear in decision theory, including minimax regret (Savage, 1951) and 'regret theory' (Bell, 1982;

<sup>&</sup>lt;sup>1</sup>Part (i) under strategic voting due to Banks (1985); the rest are from Miller (1977). See Myerson (1991, §4.10) for a nice textbook treatment of (iii) under strategic voting.

 $<sup>^{2}</sup>$ Related work by Apesteguia, Ballester and Masatlioglu (2014) and Horan (2021) axiomatises the decision rules (which specify a choice for each choice set and preference profile) defined by various agendas under strategic voting.

<sup>&</sup>lt;sup>3</sup>For example, Copeland's (1951) method (Rubinstein, 1980), the Kemeny–Slater method (Kemeny, 1959; Slater, 1961; Young & Levenglick, 1978; Young, 1986, 1988) and the fairbets method (Daniels, 1969; Moon & Pullman, 1970; Slutzki & Volij, 2005). See Charon and Hudry (2010) and González-Díaz, Hendrick and Lohmann (2014) for an overview.

<sup>&</sup>lt;sup>4</sup>For example, the probability of a cycle with five (seven) voters and five (six) alternatives is 39% (61%). See Gehrlein (1989) for a table of such probabilities.

<sup>&</sup>lt;sup>5</sup>The authoritative survey of Gehrlein and Lepelley (2011) includes data on 127 elections, of which 32 had small electorates ( $\leq 25$  voters). A Condorcet cycle existed in 29% of all elections, and in 59% of small elections (see their Table 1.1).

Loomes & Sugden, 1982; Fishburn, 1983). The online learning literature (Gordon, 1999; Zinkevich, 2003) studies (asymptotic) regret-freeness.

# 1.2 Roadmap

We begin in §2 with two applications. We describe the environment and basic concepts in §3, and discuss interpretation and extensions in §4. We then (§5) establish that regret-free strategies exist by exhibiting one: insertion sort. In §6, we characterise regret-freeness in terms of efficiency (Theorem 2) and error-avoidance (Theorem 3), and show that both characterisations are tight (Propositions 1 and 2). In §7, we show that regret-free strategies differ in their prioritisation, and characterise insertion sort in terms of its lexicographic prioritisation (Theorem 4).

# 2 Two applications

Before describing the abstract model, we fix ideas with two applications.

**Hiring.** The alternatives  $\mathcal{X}$  are candidates for a job. Only some unknown subset of candidates would accept the job if offered it. The hiring committee decides the order in which offers should be made (a ranking): the job will be offered to the first candidate, then to the second if the first declined, and so on.

Relabelling, we may instead think of the alternatives as investment projects of unknown viability. A firm's board (or a lower-level committee) ranks the projects, whereupon a manager evaluates the first project (e.g. by commissioning market research) and implements it if viable, and otherwise evaluates the second project and implements that if viable, and so on.

Similarly, the committee could be a policy-making body, such as a ministerial cabinet or a parliamentary committee. Any given policy may turn out to be infeasible, for example due to a court ruling or political opposition. The committee therefore ranks the policies and tasks a bureaucrat with implementing the first policy if feasible, the second if not, and so on.

**Party lists.** A political party's leadership committee must draw up a party list, meaning a ranking of the party's parliamentary candidates  $\mathcal{X}$ . The K top-ranked candidates will earn parliamentary seats, where K is the (uncertain) number of seats won by the party in a subsequent election.

Electoral systems of this type, called *party-list* proportional representation, are used in most of the world's democracies. More precisely, we described the *closed-list* variant that gives voters no sway over party lists; this system is used in several dozen states.<sup>6</sup> Other countries allow the electorate to influence party lists.<sup>7</sup> In many of these, voters exert little influence on party lists in practice,<sup>8</sup> making the closed-list system a reasonable idealisation.

We may re-interpret this as a firm planning for downsizing. The firm will have to fire an uncertain number K of its workers  $\mathcal{X}$ . The board (or a lower-level committee) plans ahead by drawing up an order in which employees will (if necessary) be let go.

# **3** Environment

There is a finite set  $\mathcal{X}$  of alternatives. A ranking of the alternatives is a binary relation R, where x R y reads 'x is ranked above y according to R'. Formally, a ranking is a binary relation that is transitive, irreflexive (no alternative is ranked above itself) and total (each distinct pair is ranked).<sup>9</sup> To capture the notion of a 'possibly-incomplete ranking', we use the term *proto-ranking* for relations which are transitive and irreflexive, but not necessarily total.

There is a committee and its *chair*. The committee comprises an odd number I of voters,  $i \in \{1, \ldots, I\}$ . The committee meets to determine a ranking, and the chair sets the agenda.

# 3.1 Interaction

Initially, no alternatives are ranked. In each period, the chair offers a vote on an as-yet unranked pair of alternatives. The committee votes on this pair, and whichever alternative garners more votes wins. (The chair does not have a vote, though that can be allowed for: see §4.) The winning alternative is ranked above the losing one. In addition, transitivity is imposed: if alternative x is ranked above y and y above z, then x is considered ranked above z. The chair continues to offer votes until all alternatives are ranked.

More explicitly, the ranking decisions made by the end of a period t are captured by a proto-ranking  $R_t$ , where  $x \ R_t \ y$  if x has been ranked

 $<sup>^6{\</sup>rm For}$  example, Argentina, Germany, Japan, Russia, South Africa and Turkey.

 $<sup>^7\</sup>mathrm{E.g.}$  Brazil, Indonesia, Iraq, the Netherlands and Ukraine.

<sup>&</sup>lt;sup>8</sup>In many countries, such as Indonesia and the Netherlands, the party list is only altered if a candidate receives a large number of personal votes. Furthermore, voting for an individual candidate is typically optional, and many voters do not: in Sweden, only about a quarter do (Oscarsson, 2019). Of course there are exceptions, e.g. voting for individual candidates is mandatory and important in Finnish parliamentary elections.

<sup>&</sup>lt;sup>9</sup>Definitions of some standard order-theoretic terms are collected in appendix A.

above y, and  $x \not R_t y$  otherwise. In period t + 1, the chair offers a vote on an  $R_t$ -unranked pair x, y of distinct alternatives. If x is the winner, then the new proto-ranking  $R_{t+1}$  is the transitive closure of  $R_t \cup \{(x, y)\}$ .<sup>10</sup>

A history is a sequence of pairs offered for votes and a winner of each vote. A strategy of the chair specifies what pair to offer after each non-terminal history. We give formal versions of these definitions in appendix B.2.

#### 3.2 The majority will

The chair need not keep track of individual votes: all that matters for each pair of alternatives is which one wins. This essential information is captured by the binary relation W such that x W y iff a majority of voters vote for x over y when the pair x, y is offered. We call W the majority will.

We consider all possible majority wills W, meaning all total and asymmetric relations. Clearly only such relations need be contemplated (for each pair of alternatives, one must win and the other lose). Conversely, all such relations must be considered because each of them is the majority will of some committee, as we show in appendix B.1.

The outcome of a strategy under a majority will W is the ranking that results. If a history is visited by a strategy  $\sigma$  under some majority will, then we say that it belongs to the path of  $\sigma$ . We give formal definitions of outcomes and paths in appendix B.2.

#### 3.3 Reachable rankings

For a majority will W, we call a ranking W-reachable iff it is the outcome under W of some strategy of the chair. This captures ex-post feasibility: were the chair to have perfect knowledge of W, she could achieve all and only W-reachable rankings by offering some sequence of pairwise votes.

**Example 1.** There are three alternatives  $\mathcal{X} = \{\alpha, \beta, \gamma\}$ , and the majority will satisfies  $\alpha W \gamma W \beta W \alpha$ . Graphically:



The ranking  $\beta R \alpha R \gamma$  is W-reachable, achieved by offering  $\beta, \alpha$  and  $\alpha, \gamma$ . Similarly, the rankings  $\alpha R' \gamma R' \beta$  and  $\gamma R'' \beta R'' \alpha$  are W-reachable.

<sup>&</sup>lt;sup>10</sup>Equivalently,  $z R_{t+1} w$  iff either (1)  $z R_t w$ , or (2) both (a) z = x or  $z R_t x$  and (b) y = w or  $y R_t w$ . See appendix D.1 (Observation 3, p. 25) for a proof of equivalence.




Figure 1: 'More aligned with  $\succ$  than' for three alternatives  $\mathcal{X} = \{\alpha, \beta, \gamma\}$ , where  $\alpha \succ \beta \succ \gamma$ . In this ('Hasse') diagram, there is a directed path from one ranking to another iff the former is more aligned with  $\succ$ .

# 3.4 The chair's preferences

The chair has a strict preference over the alternatives  $\mathcal{X}$ , denoted  $\succ$ . We do not fully specify the chair's preferences over rankings. We assume only that she prefers a ranking over another whenever the former is *more aligned* with her own preference over alternatives.

**Definition 1.** For rankings  $\succ$ , R and R', we say that R is more aligned with  $\succ$  than R' iff for any pair  $x, y \in \mathcal{X}$  of alternatives with  $x \succ y$ , if x R' y then also x R y.

In words, whenever a pair  $x, y \in \mathcal{X}$  is ranked 'right' by R' (viz.  $x \succ y$  and x R' y), it is also ranked 'right' by R (i.e. x R y).

**Example 1** (continued). Let the chair's preference be  $\alpha \succ \beta \succ \gamma$ . Figure 1 depicts how all rankings are ordered by 'more aligned with  $\succ$  than'. Since R'' ranks no pairs 'right', every ranking is more aligned with  $\succ$  than R''. The rankings R and R' are incomparable to each other, because R ranks  $\beta, \gamma$  'right' (while R' does not), whereas R' ranks  $\alpha, \beta$  'right' (R doesn't).

#### 3.5 Regret-free strategies

Given a majority will W, we call a ranking W-unimprovable iff there is no other ranking that is both W-reachable and more aligned with  $\succ$ . Explicitly, R is W-unimprovable exactly if for any W-reachable ranking  $R' \neq R$ , some pair  $x, y \in \mathcal{X}$  is ranked 'right' by R and 'wrong' by R'.

**Example 1** (continued). The ranking R is W-unimprovable, since neither of the other two W-reachable rankings is more aligned with  $\succ (R'' \text{ is less} aligned, and <math>R'$  is incomparable). R' is also W-unimprovable; R'' is not.

The chair does not know the majority will. One would therefore expect her to face trade-offs: a strategy that does well against W may have a regrettable outcome under some W'. A regret-free strategy is one that has no such downside: its outcome under any majority will is unimprovable expost.

**Definition 2.** A strategy is regret-free iff for any majority will W, its outcome under W is W-unimprovable.

Regret-freeness is a highly demanding optimality property. Our first result (Theorem 1 in §5) will be that, surprisingly, regret-free strategies exist.

# 4 Discussion, interpretation, extensions

Various aspects of the model deserve comment.

## 4.1 'More aligned than'

'More aligned than' captures *unambiguous* superiority of one ranking over another with respect to the chair's preference  $\succ$ .<sup>11</sup> We illustrate by example:

**Hiring** (continued from §2). Recall that the top-ranked candidate in X will be hired, where  $X \subseteq \mathcal{X}$  is unknown. A ranking is more aligned with  $\succ$  exactly if it hires a weakly  $\succ$ -better candidate at every realisation of the uncertainty X, as we show in appendix B.4. Equivalently, a more aligned ranking is one that is preferred by every expected-utility preference consistent with  $\succ$ .

This application demonstrates two general points. First, more aligned rankings are unambiguously better for the chair, given her preference  $\succ$  over the alternatives. Secondly, these comparisons are the *only* unambiguous ones: any further comparisons would have to be based on extraneous cardinal information, rather than on  $\succ$  alone.

**Party lists** (continued from §2). Recall that the K highest-ranked candidates win parliamentary seats, where K is uncertain. If a ranking R is more aligned with  $\succ$  than R', then the  $k^{\text{th}} \succ$ -best candidate who wins a seat under R is weakly  $\succ$ -better than the  $k^{\text{th}} \succ$ -best under R'. The converse is true provided there is some uncertainty about which candidates can take up seats.<sup>12</sup>

<sup>&</sup>lt;sup>11</sup>It is an instance of *single-crossing dominance*, a general way of comparing rankings (or preferences); see Curello and Sinander (2020).

<sup>&</sup>lt;sup>12</sup>A party cannot alter its list after submitting it, but circumstances may render some of its candidates ineligible for parliamentary seats. For example, many countries disqualify candidates convicted of a serious crime, and all disqualify the deceased.

The 'more aligned than' criterion has its limitations. No extra credit is given for ranking *many* pairs of alternatives 'right', for example, nor for having pairs *near the top* ranked 'right'.

## 4.2 Unimprovability as optimality

Unimprovability is the strongest ex-post optimality concept available without further assumptions about the chair's preference over rankings.<sup>13</sup> It can therefore be thought of as optimality for a chair who is unable to make fine distinctions between rankings (which are complicated objects), or who is content to 'satisfice'.

Were we fully to specify the chair's preference over rankings, we could still break her problem under full information about W into two parts: first, reach the frontier (*W*-unimprovability), then choose among the frontier rankings.

**Hiring** (continued). A ranking R is W-unimprovable exactly if any W-reachable ranking  $R' \neq R$  hires a strictly  $\succ$ -worse candidate at some realisation  $X \subseteq \mathcal{X}$  of uncertainty.

# 4.3 The imposition of transitivity

To understand why we assume that transitivity is imposed automatically, observe that the purpose of the interaction is to turn the will of the majority, which may contain (Condorcet) cycles, into a (by definition transitive) ranking. Some pairs will thus necessarily be ranked by fiat. We require that transitivity be imposed immediately after each vote because this is necessary and sufficient for committee sovereignty, the requirement that x be ranked above y if x beat y in a vote.<sup>14</sup> Indeed, we contend that the protocol described in §3.1 is the only natural one, given that the interaction must end with a ranking: as shown in supplemental appendix H, any other protocol must violate either committee sovereignty or democratic legitimacy, the requirement that the chair offer enough votes to give the committee a fair say.

# 4.4 Fixed voting

By using the majority will, we implicitly assume fixed voting: x W y means that x garners a majority over y whenever x, y is offered, whatever the

<sup>&</sup>lt;sup>13</sup>It is a non-trivial optimality concept: we show in supplemental appendix J that not all W-reachable rankings are W-unimprovable exactly if W contains a (Condorcet) cycle, and that for a typical W, most W-reachable rankings are not W-unimprovable.

<sup>&</sup>lt;sup>14</sup>Sufficiency is obvious. For necessity, suppose the chair were allowed to offer x, z even though  $x R_t y R_t z$ ; then committee sovereignty is violated whenever z beats x.

strategy of the chair. This is reasonable if voters are non-strategic: if they are unsophisticated, say, or vote 'expressively' (to please their constituents, for example). Empirically, non-strategic voting appears to be the norm in many important institutions, such as the US Congress.<sup>15</sup>

If voters were strategic, then they could potentially benefit from voting in a non-fixed way, by tailoring their behaviour to (their conjecture about) the chair's strategy. There are limits to their gains from strategising, however: we show in supplemental appendix I that any insincerity by a voter risks producing a final ranking that is less aligned with her preference over the alternatives than the ranking that sincere voting would have delivered. By contrast, sincere voting carries no such risk: no strategising can ever improve on its outcome in the (strong) 'more aligned' sense.

# 4.5 Broader interpretations

The model admits interpretations in which W arises from something other than majority voting by a committee. It could be the (expressed) preference of a single individual, for example: in this case, cycles reflect inconsistencies in his judgment, and an advisor ('the chair') exploits these by asking him to make pairwise comparisons in a well-chosen order.

In practice, the chair may be more constrained in her choice of agendasetting strategy. In that case, our analysis provides an upper bound on how well she can do, and gives a qualitative indication of what incremental adjustments to agenda-setting might most benefit her.

Finally, there need not literally be a committee chair: more generally, our results speak to how institutional or procedural design can influence collective decisions, in the direction of a(ny) given priority  $\succ$ .

#### 4.6 Extensions

Varying the committee's rules merely alters W, and so our analysis continues to apply. For example, we can accommodate a super-majority voting rule, an even electorate  $I \in \mathbf{N}$ , and abstentions. To determine W, we need only specify which alternative wins in case of an indecisive vote, e.g. by assuming that there is a status quo ranking that prevails in such cases.

A more substantial variation is to permit the chair (sometimes) to decide how two alternatives are to be ranked following an indecisive vote on them.

 $<sup>^{15}</sup>$ See Ladha (1994), Poole and Rosenthal (1997) and Wilkerson (1999), as well as the survey by Groseclose and Milyo (2010).

This happens if the chair has a vote, for example. We extend all of our results to allow for this in supplemental appendix K.

Our definition of a strategy rules out conditioning on who voted how in the past, since a history records only which alternative in each pair wins (rather than the full vote tally).<sup>16</sup> This is merely to avoid uninteresting complications: we show in supplemental appendix L that our results hold for 'extended strategies' that can condition on past votes cast.

# 5 Do regret-free strategies exist?

In this section, we exhibit a regret-free strategy: insertion sort. We first ( $\S5.1$ ) introduce a notion of efficiency that implies regret-freeness, then (in  $\S5.2$ ) define insertion sort and show that it is efficient.

# 5.1 Efficiency

A ranking R is called W-efficient iff every pair on which the chair and committee agree is ranked accordingly: if  $x \succ y$  and x W y, then x R y.

**Example 1** (continued). Recall the details from pp. 7 and 8. Since  $\alpha \succ \beta \succ \gamma$ , *W*-efficiency requires precisely that  $\alpha$  be ranked above  $\gamma$ . There are three such rankings:  $\succ$  itself,  $\beta R \alpha R \gamma$  and  $\alpha R' \gamma R' \beta$ .

An efficient strategy is a strategy whose outcome under any majority will W is W-efficient. Efficiency matters because it is linked with regret-freeness:

**Lemma 1.** For any majority will W, every W-efficient ranking is W-unimprovable.

**Corollary 1.** Any efficient strategy is regret-free.

Proof of Lemma 1. Fix a majority will W, and let R be a W-efficient ranking. To establish that R is W-unimprovable, consider any W-reachable ranking  $R' \neq R$ ; we must show that R' is not more aligned with  $\succ$  than R.

Since  $R' \neq R$ , there are alternatives  $x, y \in \mathcal{X}$  such that x R' y and y R x. Enumerate all of the alternatives that R' ranks between x and y as

$$x = z_1 R' z_2 R' \cdots R' z_N = y.$$

Since R' is W-reachable, we must have  $z_m W z_{m+1}$  for each m < N. (This follows from Observation 1 in appendix B.3 (p. 22).) There has to be an n < N

 $<sup>^{16}\</sup>mathrm{Of}$  course, this is only a restriction if the chair can observe individual votes.

at which  $z_{n+1} R z_n$ , since otherwise we would have x R y by transitivity of R. And it must be that  $z_{n+1} \succ z_n$ , because otherwise the W-efficiency of R would require that  $z_n R z_{n+1}$ . Thus the pair  $z_n, z_{n+1}$  is ranked 'right' by R  $(z_{n+1} \succ z_n \text{ and } z_{n+1} R z_n)$  and 'wrong' by  $R' (z_n R' z_{n+1})$ , so that R' fails to be more aligned with  $\succ$  than R.

# 5.2 Insertion sort, a regret-free strategy

**Insertion sort.** Label the alternatives  $\mathcal{X} \equiv \{1, \ldots, n\}$  so that  $1 \succ \cdots \succ n$ . We proceed in rounds indexed by  $k \in \{n-1, \ldots, 2, 1\}$ , starting with k = n-1:

- By the  $k^{\text{th}}$  round, those alternatives which the chair considers strictly worse than alternative k (i.e. the alternatives  $\{k + 1, ..., n\}$ ) have already been totally ranked. Now 'insert' alternative k into that ranking:
- First, pit alternative k against the highest-ranked worse alternative. If k won, then  $\{k, \ldots, n\}$  are now totally ranked (with k on top).
- If k lost, pit k against the second-highest-ranked worse alternative. If k won, then  $\{k, \ldots, n\}$  are now totally ranked (with k second).
- If k lost again, pit k against the third-highest-ranked worse alternative.
- Now  $\{k, \ldots, n\}$  are totally ranked. Decrease k by 1 and repeat.<sup>17</sup>

Insertion sort is illustrated in Figure 2 for the case of three alternatives.

**Theorem 1.** Insertion sort is efficient, hence regret-free.

*Proof.* Fix a majority will W, and let R be the outcome of insertion sort under W; we must show that R is W-efficient. To that end, fix alternatives  $x, y \in \mathcal{X}$  with  $x \succ y$  and x W y; we shall prove that x R y.

Enumerate as  $z_1 \ R \ \cdots \ R \ z_K$  all of the alternatives that are strictly  $\succ$ -worse than x, and note that  $z_k = y$  for some  $k \leq K$ . By definition of insertion sort, x will be pitted against  $z_1, z_2, \ldots$  in turn until it wins a vote. If x loses against  $z_1, \ldots, z_{k-1}$ , then it is next pitted against  $z_k = y$ , and wins since  $x \ W \ y$  by hypothesis; thus  $x \ R \ y$ . If instead x wins against some  $z_\ell$  with  $\ell < k$ , then  $x \ R \ z_\ell \ R \ \cdots \ R \ z_k = y$ , so  $x \ R \ y$  by transitivity of R.

<sup>&</sup>lt;sup>17</sup>A detail: our definition of strategies requires them to specify behaviour after all histories, even those that cannot arise. (E.g. a strategy that first offers x, y must still specify behaviour after histories where only y, z are ranked.) Thus formally, insertion sort defines not a strategy, but rather an equivalence class of strategies having the same path.




Figure 2: Insertion sort with three alternatives  $\mathcal{X} = \{1, 2, 3\}$ , where the chair's preference is  $1 \succ 2 \succ 3$ . *R* denotes the (evolving) proto-ranking. The path taken depends on the majority will *W*.

Insertion sort is a standard sorting algorithm—see e.g. Knuth (1998, §5.2.1). In the sorting problem, there is a *transitive* and asymmetric relation W, and one seeks to reach a 'sorted' ranking R, meaning that x R y whenever x W y, by making (as few as possible) binary comparisons using W.

Insertion sort is closely related to the *amendment procedure*, a major focus of the agenda-setting literature; see appendix G.

# 6 What makes a strategy regret-free?

Insertion sort is regret-free, but it is not the only such strategy. Which of its characteristics are essential for regret-freeness, and which are extraneous?

We give two answers in this section. Our first characterisation, Theorem 2 (§6.1), is in terms of outcomes: the regret-free strategies are exactly the efficient ones, i.e. those reaching a W-efficient ranking under every majority will W. We show this characterisation to be tight (Proposition 1).

The second characterisation, Theorem 3 (§6.2), is in terms of behaviour: regret-freeness requires precisely that two intuitive errors be avoided. We show in addition that the advice offered by Theorem 3 can be operationalised myopically: avoiding errors ensures that a non-error pair will always be available to be offered next (Proposition 2).

## 6.1 The outcomes of regret-free strategies

Efficiency is not only sufficient for regret-freeness, but also necessary:

Theorem 2. A strategy is regret-free iff it is efficient.

The proof, in appendix D, establishes Theorem 2 jointly with Theorem 3 below (§6.2). The 'if' direction was already established in Corollary 1 (p. 12). The following gives a feel for the 'only if' direction:

**Example 2.** Consider alternatives  $\mathcal{X} = \{\alpha, \beta, \gamma, \delta\}$  with  $\alpha \succ \beta \succ \gamma \succ \delta$  and the following majority will W:<sup>18</sup>



There is exactly one *W*-reachable ranking that ranks  $\alpha$  above  $\beta$ , namely  $\alpha R \delta R \gamma R \beta$ .<sup>19</sup> Since no other *W*-reachable ranking ranks  $\alpha$  above  $\beta$ , *R* is *W*-unimprovable. But *R* fails to be *W*-efficient, for it ranks  $\delta$  above  $\beta$  despite the fact that  $\beta W \delta$ . (Thus the converse of Lemma 1 (p. 12) is false.)

Let  $\sigma$  be a strategy that (i) first offers  $\beta$ ,  $\gamma$ , (ii) then, in case  $\gamma$  won, offers  $\gamma$ ,  $\delta$ , and (iii) next, in case  $\gamma$  and then  $\delta$  won, offers  $\alpha$ ,  $\delta$ . This strategy has outcome R under W, so fails to be efficient. To see that it is not regret-free, consider a majority will W' that differs from W only in that  $\delta W' \alpha$ . The outcome of  $\sigma$  under W' is  $\delta R' \gamma R' \beta R' \alpha$ .<sup>20</sup> This ranking fails to be W'-unimprovable since  $\gamma R'' \beta R'' \delta R'' \alpha$  is W'-reachable and is more aligned with  $\succ$  by inspection. Thus  $\sigma$  fails to be regret-free.

The broader insight underlying the 'only if' part of Theorem 2 is that reaching a non-W-efficient outcome necessarily involves a sacrifice. (In the example,  $\sigma$  forgoes the opportunity to rank  $\beta$  above  $\delta$ .) For some majority wills, such as W, the sacrifice pays off. (It allows  $\alpha$  to be ranked above  $\beta$ , something that no W-reachable W-efficient ranking achieves.) But for other majority wills, such as W', no reward materialises, yielding a non-W'unimprovable outcome.

The characterisation in Theorem 2 is tight, in the following sense:

<sup>&</sup>lt;sup>18</sup>Symbolically,  $\alpha W \delta W \gamma W \beta W \alpha$ ,  $\gamma W \alpha$  and  $\beta W \delta$ .

<sup>&</sup>lt;sup>19</sup>This can be verified directly. Alternatively, since there is exactly one W-path from  $\alpha$  to  $\beta$  (namely  $(\alpha, \delta, \gamma, \beta)$ ), it follows by Observation 1 in appendix B.3 (p. 22).

<sup>&</sup>lt;sup>20</sup>First  $\delta R' \gamma R' \beta$  and  $\delta R' \alpha$  are determined. Then  $\alpha, \beta$  and  $\alpha, \gamma$  are offered (in some order that doesn't matter), and  $\alpha$  loses in both votes.

**Proposition 1.** For any majority will W and any W-reachable W-efficient ranking R, some regret-free strategy has outcome R under W.

Thus no statement sharper than Theorem 2 can be made about the outcomes under W of regret-free strategies: every W-reachable W-efficient ranking is admissible. We give the proof of Proposition 1 in appendix E. A non-trivial argument is required because a given W-reachable W-efficient ranking can be reached in many ways, not all of which form part of a regret-free strategy.<sup>21</sup>

# 6.2 The behaviour of regret-free strategies

To understand the behavioural content of regret-freeness, we begin with a simple intuition for why insertion sort is regret-free in the case of three alternatives (refer to Figure 2 on p. 14). Suppose first that the initial vote on 2, 3 went well (2 W 3). Offering 1, 2 next is a good idea because it affords an opportunity of a favourable imposition of transitivity: if the chair gets lucky (1 W 2), then she obtains 1 R 3 'for free' from 1 R 2 and 2 R 3. (Even though it may be that 3 W 1.) Offering 1, 3 would miss this opportunity.

Suppose instead that the initial vote on 2, 3 went badly (3 W 2). Offering 1, 2 next would risk an unfavourable imposition of transitivity: were the vote to go against her (2 W 1), then 3 would be ranked above 1 since 3 R 2 and 2 R 1. (Even though it may be that 1 W 3.)

To summarise, an intuition for why insertion sort is regret-free is that it never (1) misses an opportunity for a favourable imposition of transitivity nor (2) risks an unfavourable imposition of transitivity. Our second characterisation formalises this intuition.

**Definition 3.** Let R be a non-total proto-ranking, and let  $x \succ y$  be unranked alternatives. (I.e.  $x, y \in \mathcal{X}$  such that  $x \not R y \not R x$ .)

- (1) Offering x, y for a vote misses an opportunity (at R) iff there is an alternative  $z \in \mathcal{X}$  such that  $x \succ z \succ y$  and  $y \not R z \not R x$ .
- (2) Offering x, y for a vote takes a risk (at R) iff there is an alternative  $z \in \mathcal{X}$  such that either
  - (a)  $z \succ y$ , x R z and  $y \not R z$ , or

<sup>&</sup>lt;sup>21</sup>To see why, return to Example 1 (pp. 7 and 12). The W-efficient ranking  $\alpha R' \gamma R' \beta$  may be reached by offering  $\alpha, \gamma$  and then  $\gamma, \beta$ . But a strategy that does this cannot be regret-free, because it has outcome R' also under the majority will  $\alpha W' \beta W' \gamma W' \alpha$ , and R' is not a W'-unimprovable ranking (since the more aligned ranking  $\succ$  is W'-reachable).




Figure 3: Proof of Theorems 2 and 3.

(b)  $x \succ z$ , z R y and z R x.

The 'missed opportunity' in (1) is that x R y (the hoped-for outcome when offering x, y for a vote) could potentially have been obtained 'for free' by offering votes on x, z and z, y, via a 'favourable imposition of transitivity'. The 'risk' in (2)(a) is that if the vote on x, y were to go badly (so that y R x), then y R z would follow—an 'unfavourable imposition of transitivity'. Similarly for (2)(b).

We say that a strategy never misses an opportunity (never takes a risk) iff it does not miss an opportunity (take a risk) on the path.

**Theorem 3.** A strategy is regret-free iff it never misses an opportunity or takes a risk.

We give a joint proof of Theorems 2 and 3 in appendix D. The argument is illustrated in Figure 3.

The 'only if' part of Theorem 3 asserts that opportunity-missing and risk-taking are errors, in the sense that committing one of them will lead to a non-W-unimprovable ranking under some W. This is intuitive, but requires some work to show. To appreciate why, suppose that  $\sigma$  offers a pair  $x \succ y$ under W, and that doing so either misses an opportunity or takes a risk. It is then easy to find another majority will W' such that the outcome R of  $\sigma$  under W' ranks y above x, whereas some other W'-reachable ranking R'ranks x R' y. But that is not enough: if R' is to be more aligned with  $\succ$  than R, then every pair z, w ranked 'right' by  $R (z \succ w \text{ and } z R w)$  must also be ranked 'right' by R' (z R' w). We construct W' and R' with these properties in appendix D.2. The 'if' part of Theorem 3 asserts that these are the only errors. Thus to achieve regret-freeness, it suffices to avoid missing opportunities and taking risks, separately after each history. The proof in appendix D.1 begins with an arbitrary non-efficient strategy  $\sigma$ , meaning one whose outcome R under some majority will W ranks some pair  $x, y \in \mathcal{X}$  as y R x even though  $x \succ y$ and x W y. The pair x, y cannot have been voted on (else x would have prevailed), so must have been ranked by an imposition of transitivity. We show that avoiding the two errors suffices to preclude such 'unfavourable' impositions of transitivity; thus  $\sigma$  must have committed the one or the other.

Theorem 3 tells us that non-error pairs are attractive, but it does not promise that they exist. In particular, there could conceivably be histories along which the chair has committed no errors, but is now forced to do so because no unranked pairs remain that can be offered without missing an opportunity or taking a risk. The following rules out this scenario:

**Proposition 2.** After any history along which the chair has not missed an opportunity or taken a risk, if some pairs remain unranked, then there is a pair which can be offered without missing an opportunity or taking a risk.

The proof is in appendix E. Proposition 2 shows that the characterisation in Theorem 3 of regret-free behaviour is tight: for any W and sequence of pairs that is error-free under W, some regret-free strategy offers this sequence under W.<sup>22</sup>

Theorem 3, augmented by Proposition 2, allows us to give the following simple 'myopic' advice to the chair. After each history, inspect the current proto-ranking to identify an unranked pair of alternatives that would not miss an opportunity or take a risk; that such a pair exists is guaranteed by Proposition 2. Offer any such pair for a vote. By Theorem 3, the outcome will be W-unimprovable whatever the majority will W.

# 7 What distinguishes regret-free strategies?

In this section, we argue that regret-free strategies differ (only) in how they *prioritise* various pairs of alternatives. We show that insertion sort is characterised by a *lexicographic* prioritisation.

 $<sup>^{22}</sup>$ In particular, a strategy can be constructed which offers this sequence under W and commits no errors under other majority wills. Such a strategy is regret-free by Theorem 3.

# 7.1 Prioritisation

Given a majority will W, call a pair  $x \succ y$  an agreement pair iff x W y, and a disagreement pair otherwise. All regret-free strategies rank the agreement pairs 'right' (i.e. x R y) by Theorem 2. A disagreement pair is ranked 'wrong' (as y R x) if voted on, and ranked 'right' if instead ranked by transitivity, since regret-free strategies permit only favourable impositions of transitivity by Theorem 3.

What distinguishes regret-free strategies is thus precisely which disagreement pairs are voted on and which are held back. Held-back pairs have more opportunities to be ranked 'right' by transitivity.

**Example 1** (continued). Recall the details from pp. 7 and 8. The disagreement pairs are  $\alpha, \beta$  and  $\beta, \gamma$ . Insertion sort prioritises the former: its outcome is  $\alpha R' \gamma R' \beta$ . Other regret-free strategies prioritise  $\beta, \gamma$  instead, having outcome  $\beta R \alpha R \gamma$ .<sup>23</sup>

# 7.2 How does insertion sort prioritise?

To understand the priorities of insertion sort, label the alternatives  $\mathcal{X} \equiv \{1, \ldots, n\}$  so that  $1 \succ \cdots \succ n$ . Insertion sort leaves alternative 1 for last, by totally ranking the alternatives  $\{2, \ldots, n\}$  before offering any votes involving 1. This intuitively maximises opportunities for favourable impositions of transitivity involving 1, and so optimises the placement of this alternative.

Apart from alternative 1, alternative 2 is left for last by insertion sort, suggesting that this alternative is placed optimally, subject to optimal placement of alternative 1. This suggests that insertion sort prioritises lexicographically, by placing alternative 1 as well as possible, and subject to that placing alternative 2 as well as possible, and so on. We shall formalise this intuition.

A simple cardinal measure of how well a strategy  $\sigma$  places a given alternative  $x \in \mathcal{X}$  is how many  $\succ$ -worse alternatives y are ranked below x, i.e.

$$N_x^{\sigma}(W) \coloneqq |\{y \in \mathcal{X} : x \succ y \text{ and } x R^{\sigma}(W) y\}|_{\mathcal{X}}$$

where  $R^{\sigma}(W)$  denotes the outcome of  $\sigma$  under majority will W. We call a strategy  $\sigma$  better for x than another strategy  $\sigma'$  iff  $N_x^{\sigma}$  first-order stochastically dominates  $N_x^{\sigma'}$ , i.e.

$$|\{W: N_x^{\sigma}(W) \ge k\}| \ge |\{W: N_x^{\sigma'}(W) \ge k\}| \quad \text{for each } k \in \{1, \dots, n-1\}.$$

<sup>&</sup>lt;sup>23</sup>Some regret-free strategy has outcome R under W by Proposition 1 (p. 16). An example is 'reverse insertion sort': in its  $k^{\text{th}}$  round, the chair's top k-1 alternatives have already been ranked, and now her  $k^{\text{th}}$ -favourite is pitted in turn against the lowest-ranked of these  $\succ$ -better alternatives, against the second-lowest ranked, and so on until it loses a vote.

If a strategy  $\sigma$  belongs to a set  $\Sigma$  of strategies and is better for x than any other  $\sigma' \in \Sigma$ , then we say that  $\sigma$  is best for alternative x among  $\Sigma$ .

Call a strategy *lexicographic* iff among all strategies, it is best for the chair's favourite alternative; among such strategies, it is best for her second-favourite; among *such* strategies, it is best for her third-favourite; etc.

**Theorem 4.** A strategy is outcome-equivalent to insertion sort iff it is lexicographic.

('Outcome-equivalent' means having the same outcome under each majority will W.) The proof is in appendix F.

# Appendices

## A Standard definitions

This appendix collects the definitions of order-theoretic concepts used in this paper. Let  $\mathcal{A}$  be a non-empty set, and Q a binary relation on it. Recall that Q is formally a subset of  $\mathcal{A} \times \mathcal{A}$ , and that 'a Q b' is shorthand for ' $(a, b) \in Q$ '. For  $a, b \in \mathcal{A}$ , we write  $a \ Q \ b$  iff it is not the case that  $a \ Q \ b$ . For  $a, b \in \mathcal{A}$  such that  $a \ Q \ b$ , we denote by  $[a, b]_Q$  the Q-order interval

$$[a,b]_Q \coloneqq \{a,b\} \cup \{c \in \mathcal{A} : a \ Q \ c \ Q \ b\}.$$

Q is reflexive (irreflexive) iff  $a \ Q \ a$  ( $a \ Q \ a$ ) for every  $a \in A$ , total iff  $a \ Q \ b$  or  $b \ Q \ a$  for any distinct  $a, b \in A$ , complete iff it is reflexive and total, asymmetric iff  $a \ Q \ b$  implies  $b \ Q \ a$  for  $a, b \in A$ , and transitive iff  $a \ Q \ b \ Q \ c$  implies  $a \ Q \ c$  for  $a, b, c \in A$ .

The transitive closure of Q, denoted tr Q, is the smallest (in the sense of set inclusion) transitive relation that contains  $Q^{24}$ . The strict part of Qis the binary relation str Q such that  $a \operatorname{str} Q b$  iff a Q b and b Q a. For two binary relations Q and Q' on  $\mathcal{A}$ , Q' is an extension of Q iff both  $Q \subseteq Q'$  and str  $Q \subseteq \operatorname{str} Q'$ .

# B Additional material for §3

This appendix complements the exposition of the environment in §3. We show that all and only total and asymmetric relations can be majority wills (§B.1), formally define histories, strategies, outcomes and paths (§B.2), and provide characterisations of W-reachability (§B.3) and of 'more aligned with than' (§B.4).

<sup>&</sup>lt;sup>24</sup>Every relation possesses a transitive closure because the maximal relation  $\mathcal{A} \times \mathcal{A}$  is transitive and the intersection of transitive relations is transitive.

#### B.1 Which binary relations are majority wills? (§3.2)

In this appendix, we prove that all and only total and asymmetric relations are legitimate majority wills. A voting behaviour  $V_i$  specifies for each pair  $x, y \in \mathcal{X}$  whether voter *i* will vote for x ( $x V_i y$ ) or for y ( $y V_i x$ ). Formally:

**Definition 4.** A voting behaviour is a total and asymmetric relation. A voting profile is a collection  $(V_i)_{i=1}^{I}$  of voting behaviours.

The majority will of a voting profile  $(V_i)_{i=1}^I$  is the relation W such that  $x \ W \ y$  iff  $x \ V_i \ y$  for a majority of voters  $i \in \{1, \ldots, I\}$ . The following shows that all (and only) total and asymmetric relations are the majority will of some voting profile  $(V_i)_{i=1}^I$ , even if we insist that each voter's behaviour  $V_i$  be transitive (meaning that it can be rationalised as sincere):

**Fact 1.** For a binary relation W on  $\mathcal{X}$ , the following are equivalent:

- (1) W is total and asymmetric.
- (2) For every  $I \in \mathbf{N}$  odd, W is the majority will of some profile  $(V_i)_{i=1}^I$  of voting behaviours.
- (3) For some  $I \in \mathbf{N}$  odd, W is the majority will of some profile  $(V_i)_{i=1}^I$  of transitive voting behaviours.

Proof. It is immediate that (2) and (3) (separately) imply (1). To see that (1) implies (2), simply observe that a total and asymmetric relation W is the majority will of the voting profile  $(V_i)_{i=1}^I = (W)_{i=1}^I$  for any  $I \in \mathbf{N}$ . The fact that (1) implies (3) follows from McGarvey's (1953) theorem.

#### B.2 Formal definitions (§3.1–§3.2)

In this appendix, we formally define histories, strategies, outcomes and paths. A history is a sequence  $((x_t, y_t))_{t=1}^T$ , where  $\{x_t, y_t\}$  is the pair offered in period t, and  $x_t$  is the winner (i.e.  $x_t W y_t$ ):

**Definition 5.** There is exactly one history of length 0 (the 'empty history'). A history of length  $T \in \mathbf{N}$  is a sequence  $((x_t, y_t))_{t=1}^T$  of ordered pairs of alternatives satisfying  $x_t \neq y_t$  and  $x_t \not R_{t-1} y_t \not R_{t-1} x_t$  for each  $t \in \{1, \ldots, T\}$ , where  $R_0 \coloneqq \emptyset$  and  $R_t \coloneqq \operatorname{tr}(R_{t-1} \cup \{(x_t, y_t)\})$  for each  $t \in \{1, \ldots, T\}$ . (Here 'tr' denotes the transitive closure.) The history is terminal iff  $R_T$  is total.

Let  $\mathcal{H}_T$  be the set of all non-terminal histories of length  $T \geq 0$ , and write  $\mathcal{H} := \bigcup_{T=0}^{\infty} \mathcal{H}_T$  for all non-terminal histories.<sup>25</sup> A strategy offers, after each non-terminal history, a pair of alternatives that are unranked at that history:

**Definition 6.** A strategy of the chair is a map  $\sigma : \mathcal{H} \to 2^{\mathcal{X}}$  such that for each non-terminal history  $h \in \mathcal{H}$ , we have  $\sigma(h) = \{x, y\}$  for some alternatives  $x, y \in \mathcal{X}$  such that (h, (x, y)) and (h, (y, x)) are histories.

A strategy  $\sigma$  and a majority will W generate a terminal history  $((x_t, y_t))_{t=1}^T$ as follows: for each  $t \in \{1, \ldots, T\}$ ,  $(x_t, y_t)$  is given by

$$\{x_t, y_t\} \coloneqq \sigma \left( ((x_s, y_s))_{s=1}^{t-1} \right) \quad \text{and} \quad x_t \ W \ y_t$$

This history is associated with a sequence of proto-rankings  $(R_t)_{t=0}^T$ , as outlined in Definition 5 above.<sup>26</sup>

**Definition 7.** The *outcome* of a strategy  $\sigma$  under a majority will W is the ranking  $R_T$  associated with the terminal history they generate.

A strategy and a majority will also generate a sequence of non-terminal histories—namely, all truncations of their generated terminal history. If a non-terminal history is generated by  $\sigma$  and some majority will, we say that it belongs to the path of  $\sigma$ .

#### **B.3** A characterisation of *W*-reachability (§3.3)

This appendix contains a characterisation of reachability used in our proofs.

**Definition 8.** Let R be a ranking. Alternatives  $x, y \in \mathcal{X}$  with  $x \in R$  are *R*-adjacent iff there is no  $z \in \mathcal{X}$  such that x R z R y.

**Observation 1.** Let W be a majority will, and R a ranking. The following are equivalent:

(1) R is W-reachable.

(2) For any *R*-adjacent  $x, y \in \mathcal{X}$  with x R y, we have x W y.

Condition (2) admits a graph-theoretic interpretation. Think of W as a directed graph with vertices  $\mathcal{X}$  and a directed edge from x to y iff x W y (as in Example 1 on p. 7), and think of a ranking R as a sequence of alternatives: the highest-ranked, the second-highest-ranked, and so  $on.^{27}$  Condition (2) requires precisely that R be a directed path in W.

<sup>&</sup>lt;sup>25</sup>Histories can only be so long:  $\mathcal{H}_T$  is empty for (and only for)  $T \ge |\mathcal{X}| \times (|\mathcal{X}| - 1)/2$ .

<sup>&</sup>lt;sup>26</sup>Namely,  $R_0 = \emptyset$  and  $R_t = \operatorname{tr}(R_{t-1} \cup \{(x_t, y_t)\}) = \operatorname{tr}\left(\bigcup_{s=1}^t \{(x_s, y_s)\}\right)$  for  $t \ge 1$ . <sup>27</sup>Formally: identify R with the sequence  $(x_k)_{k=1}^{|\mathcal{X}|}$  such that  $x_1 R x_2 R \cdots R x_{|\mathcal{X}|}$ .

Proof. (2) implies (1): Let R satisfy (2). Then R is the outcome under W of any strategy that offers a vote on each R-adjacent pair of alternatives.

(1) implies (2): Let R be W-reachable, and let  $x, y \in \mathcal{X}$  be R-adjacent with x R y. By W-reachability, there is a strategy  $\sigma$  whose outcome under W is R. Along its induced history, it is determined that x R y. Since x, y are R-adjacent, this cannot be via an imposition of transitivity. So it must occur in a vote on  $\{x, y\}$ , in which x wins—thus x W y.

# B.4 A characterisation of 'more aligned with than' (§3.4)

In this appendix, we provide a characterisation of 'more aligned with than', and use it to prove the claims made in §3.4 (p. 9) about the applications.

**Observation 2.** For rankings  $\succ$ , R and R', the following are equivalent:<sup>28</sup>

- (1) R is more aligned with  $\succ$  than R'.
- (2) For every non-empty set  $X \subseteq \mathcal{X}$ , the *R*-highest alternative in *X* is identical to or  $\succ$ -better than the *R'*-highest alternative in *X*.

Proof. (1) implies (2): We prove the contra-positive. Suppose that R, R' do not satisfy (2), so that there is a non-empty  $X \subseteq \mathcal{X}$  whose R-highest element x is (strictly)  $\succ$ -worse than its R'-highest element x'. Then R' ranks x, x' 'right' ( $x' \succ x$  and x' R' x) whereas R ranks them 'wrong' (x R x'), so R is not more aligned with  $\succ$  than R'.

(2) implies (1): We prove the contra-positive. Suppose that R is not more aligned with  $\succ$  than R', so that there are alternatives  $x, x' \in \mathcal{X}$  with  $x' \succ x, x' R' x$  and x R x'. Then the R-highest alternative in  $X \coloneqq \{x, x'\}$  is (strictly)  $\succ$ -worse than the R'-highest.

**Hiring** (continued). We claimed that a more aligned ranking is precisely one that hires a weakly  $\succ$ -better candidate at every realisation of uncertainty. This follows immediately from Observation 2.

**Party lists** (continued). Enrich the model so that only a random subset  $X \subseteq \mathcal{X}$  of candidates is available.<sup>29</sup> We claim that R is more aligned with  $\succ$  than R' iff for every realisation (K, X) and every  $k \leq K$ , the  $k^{\text{th}} \succ$ -best candidate hired by R is weakly  $\succ$ -better than the  $k^{\text{th}} \succ$ -best hired by R'.

<sup>&</sup>lt;sup>28</sup>This is an instance of the Milgrom–Shannon (1994) comparative statics theorem: viewing  $(\mathcal{X}, \succ)$  as an ordered set of actions and R, R' as (strict) preferences, (1) says that R single-crossing dominates R', and (2) says that R chooses higher actions than R' does.

<sup>&</sup>lt;sup>29</sup>The grand set  $X = \mathcal{X}$  can occur with high probability, if desired.

To prove the 'only if' part, let R be more aligned with  $\succ$  than R', and fix an  $X \subseteq \mathcal{X}$  and a K. Assume without loss of generality that  $K \leq |X|$ . Label the candidates  $\{x_1, \ldots, x_K\}$  hired by R under X so that  $x_1 R \cdots R x_K$ , and similarly write  $x'_1 R' \cdots R' x'_K$  for those hired by R'. We must show that  $x_k \succeq x'_k$  for every  $k \in \{1, \ldots, K\}$ . We proceed by strong induction on k. The base case k = 1 is immediate from Observation 2.

For the induction step, suppose for some  $k \in \{2, \ldots, K\}$  that  $x_{\ell} \succeq x'_{\ell}$ for every  $\ell < k$ . Define  $Y \coloneqq X \setminus \{x_1, \ldots, x_{k-1}, x'_1, \ldots, x'_{k-1}\}$ . We have  $x_{\ell} \succeq x'_{\ell} \succ x'_k$  for every  $\ell < k$  by the induction hypothesis. It follows that  $x'_k \in Y$ , and hence that  $x'_k$  is the R'-highest alternative in Y. If  $x_k \in Y$ , then  $x_k$  is the R-highest alternative in Y, whence  $x_k \succeq x'_k$  by Observation 2. If instead  $x_k \notin Y$ , then  $x_k = x'_{\ell} \succ x'_k$  for some  $\ell < k$ .

For the 'if' part, we prove the contra-positive. Suppose that R is not more aligned with  $\succ$  than R'. Then by Observation 2, there is a subset  $X' \subseteq \mathcal{X}$ such that R hires a strictly  $\succ$ -worse candidate than R' at the realisation (K, X) = (1, X') of uncertainty.

# C An extension lemma

This appendix presents an extension lemma that is used in the proofs of Theorems 2 and 3 in appendix D. Recall from appendix A the definitions of 'extension' and of order intervals  $[x, y]_R$ .

**Extension lemma.** Let R be a proto-ranking, and let  $A \subseteq \mathcal{X}$  be such that  $[x, y]_R \subseteq A$  for all  $x, y \in A$  with x R y. Then the binary relation  $R \cup A^2$  admits a complete and transitive extension.

The extension lemma is used directly in the proof of Theorems 2 and 3 (in particular, in the proof of Lemma 4 in appendix D.2). To prove it, we use a theorem due to Suzumura (1976).

**Definition 9.** A binary relation Q on a set  $\mathcal{A}$  is Suzumura-consistent iff for  $a_1, \ldots, a_K \in \mathcal{A}, a_1 Q a_2 Q \cdots Q a_{K-1} Q a_K$  implies that either  $a_1 Q a_K$  or  $a_1 \mathcal{Q} a_K \mathcal{Q} a_1$ .

Suzumura's extension theorem. A binary relation admits a complete and transitive extension iff it is Suzumura-consistent.

See Bossert and Suzumura (2010, p. 45) for a proof.

Proof of the extension lemma. Let R and A satisfy the hypothesis; we seek a complete and transitive extension of the relation  $Q := R \cup A^2$ . By Suzumura's extension theorem, it suffices to show that for any finite sequence of alternatives  $(w_k)_{k=1}^K$  such that  $w_1 Q \cdots Q w_K$ , we have either  $w_1 Q w_K$  or  $w_1 Q w_K Q w_1$ . There are two cases.

Case 1:  $w_k R w_{k+1}$  for every k < K. Then  $w_1 R w_K$  since R is transitive (being a proto-ranking), so  $w_1 Q w_K$ .

Case 2:  $\{w_k, w_{k+1}\} \subseteq A$  for some k < K. Let k'(k'') be the smallest (largest) such k < K, so that  $w_1 R \cdots R w_{k'}$  if k' > 1 and  $w_{k''+1} R \cdots R w_K$  if k'' < K - 1. Assume toward a contradiction that  $w_K Q w_1$  and  $w_1 Q w_K$ . Since  $\{w_1, w_K\} \not\subseteq A$  (as otherwise  $w_1 Q w_K Q w_1$ ), it must be that  $w_K R w_1$ . It follows by transitivity of R that  $w_{k''+1} R w_K R w_1 R w_{k'}$ , and thus  $\{w_K, w_1\} \subseteq [w_{k''+1}, w_{k'}]_R$ . Note that  $[w_{k''+1}, w_{k'}]_R \subseteq A$  since  $w_{k''+1}, w_{k'} \in A$  by construction. Therefore  $\{w_K, w_1\} \subseteq A$ , which implies that  $w_K Q w_1 Q w_K$ —a contradiction.

The following corollary is used in supplemental appendix I.

**Extension corollary.** Let R be a proto-ranking, and let  $x, y \in \mathcal{X}$  be distinct alternatives such that  $x \not R y \not R x$ . Then there is a ranking  $R' \supseteq R$  such that x, y are R'-adjacent with x R' y.

Proof. Let R and  $x, y \in \mathcal{X}$  satisfy the hypothesis. Then  $R \cup \{x, y\}^2$  admits a complete and transitive extension Q by the extension lemma. Note that  $x \ Q \ y \ Q \ x$ . It follows that by appropriately breaking indifferences in Q, we may obtain a ranking  $R' \supseteq R \cup \{(x, y)\}$  such that x, y are R'-adjacent with  $x \ R' \ y$ .

# D Proof of Theorems 2 and 3 (\$6, pp. 15 and 17)

We prove Theorems 2 and 3 jointly, in the manner depicted in Figure 3 (p. 17). We already showed that efficiency implies regret-freeness (Corollary 1, p. 12). We shall establish the other two parts in §D.1 and §D.2.

#### D.1 Proof that error-avoiding strategies are efficient

The proof relies on two intermediate results, Lemma 2 and Corollary 2 below. We first require an abstract fact about the transitive closure operation:

**Observation 3.** Consider a proto-ranking R and unranked alternatives  $x, y \in \mathcal{X}$  (i.e.  $x \not R y \not R x$ ). Let R' be the transitive closure of  $R \cup \{(z, w)\}$ , and suppose that y R' x. Then (a) either y R z or y = z, and (b) either w R x or w = x.

Proof. Since R' is the transitive closure of  $R \cup \{(z, w)\}$  and y R' x, there must be a sequence  $(z_k)_{k=1}^K$  of alternatives with  $z_1 = y$ ,  $z_K = x$  and

$$(z_k, z_{k+1}) \in R \cup \{(z, w)\}$$
 for every  $k < K$ .

Since  $y \not R x$  and R is transitive, we must have  $(z_k, z_{k+1}) = (z, w)$  for some k < K. The result follows since R is transitive.

**Definition 10.** Let R be a proto-ranking. An ordered pair of alternatives  $(x, y) \in \mathcal{X}$  is a missed opportunity in R iff there is an alternative  $z \in \mathcal{X}$  such that  $x \succ z \succ y$ , y R x and y R z R x.

**Lemma 2.** Consider a proto-ranking R that contains no missed opportunities. Let  $x \succ y$  be alternatives with  $y \not R x$ . Suppose that offering  $\{z, w\}$  does not miss an opportunity or take a risk at R, and that doing this leads to a proto-ranking R' such that  $y \not R' x$ . Then  $\{z, w\} = \{x, y\}$ .

That is: if no errors are made, then a misfortune y R' x can occur only as the result of a vote on  $\{x, y\}$ , not via an imposition of transitivity. This captures the idea that avoiding errors is enough to preclude unfavourable impositions of transitivity.

*Proof.* Let R, x, y, z, w and R' satisfy the hypothesis of the lemma, and assume wlog that  $z \succ w$ . We must show that z = x and w = y.

# Claim. w R' z.

Proof of the claim. Suppose toward a contradiction that  $w \not R' z$ . We will show that R contains a missed opportunity.

Since R' is induced from R by offering  $\{z, w\}$ , and  $w \not R' z$ , it must be that R' is the transitive closure of  $R \cup \{(z, w)\}$ . Since  $y \not R x$  and y R' x, it follows by Observation 3 that (a) either y R z or y = z, and (b) either w R x or w = x. Now consider two cases.

Case 1:  $z \succ x$  or z = x. We will show that  $z \succ x \succ y$ ,  $y \not R z$ , and  $y \not R x \not R z$ , so that (z, y) is a missed opportunity in R. Both  $x \succ y$  and  $y \not R x$  hold by hypothesis. For  $x \not R z$ , suppose to the contrary that  $x \not R z$ ; then since  $w \ R x$  or w = x by property (b), we have  $w \ R z$  by transitivity of R, a contradiction.

To obtain  $y \ R \ z$ , observe that  $z \succ y$  since by hypothesis  $z \succ x$  or z = x, and we know that  $x \succ y$  and that  $\succ$  is transitive. Thus  $z \neq y$ , whence  $y \ R \ z$  follows by property (a). To see that  $z \succ x$ , simply note that z = x is impossible because  $y \ R \ z$  and  $y \ R \ x$ . Case 2:  $x \succ z$ . We will show that  $x \succ z \succ w$ ,  $w \mathrel{R} x$ , and  $w \mathrel{R} z \mathrel{R} x$ , so that (x, w) is a missed opportunity in R. Both  $x \succ z \succ w$  and  $w \mathrel{R} z$  hold by hypothesis. For  $z \mathrel{R} x$ , suppose to the contrary that  $z \mathrel{R} x$ ; then since  $y \mathrel{R} z$  or y = z by property (a), we have  $y \mathrel{R} x$  by transitivity of R, a contradiction.

To obtain  $w \ R \ x$ , observe that  $x \succ w$  since by hypothesis  $x \succ z$ , and we know that  $z \succ w$  and that  $\succ$  is transitive. Thus  $w \neq x$ , whence  $w \ R \ z$  follows by property (b).

In light of the claim, R' must be the transitive closure of  $R \cup \{(w, z)\}$ . Since  $y \not R x$  and y R' x, applying Observation 3 yields that (a) either y R w or y = w, and (b) either z R x or z = x.

We claim that

$$z \neq x \succ w \tag{1}$$

is impossible. Suppose toward a contradiction that (1) holds; we will show that offering  $\{z, w\}$  takes a risk at R, i.e. that  $x \succ w$ ,  $z \mathrel{R} x$  and  $w \mathrel{R} x$ . We have  $x \succ w$  by (1), and  $z \mathrel{R} x$  by (1) and property (a). To see that  $w \mathrel{R} x$ , suppose to the contrary that  $w \mathrel{R} x$ ; then since  $y \mathrel{R} w$  or y = w by property (a), it follows by transitivity of R that  $y \mathrel{R} x$ , a contradiction.

Now suppose toward a contradiction that  $\{z, w\} \neq \{x, y\}$ . We claim that

$$z \succ y \neq w. \tag{2}$$

If z = x, then  $z \succ y$  is immediate, and  $w \neq y$  follows since  $\{z, w\} \neq \{x, y\}$  by hypothesis. Suppose instead that  $z \neq x$ . Since (1) cannot hold, it must be that either  $w \succ x$  or w = x. Since  $x \succ y$ , it follows by transitivity of  $\succ$  that  $w \succ y$ , so that  $w \neq y$ . Furthermore, since  $z \succ w$ , transitivity of  $\succ$  yields  $z \succ y$ . So (2) holds.

It remains to derive a contradiction from  $\{z, w\} \neq \{x, y\}$ , using the fact that (2) must hold. We shall show that  $z \succ y$ ,  $y \mathrel{R} w$  and  $y \mathrel{R} z$ , so that offering  $\{z, w\}$  takes a risk at R. We obtain  $z \succ y$  from (2), and  $y \mathrel{R} w$  from (2) and property (a). And it must be that  $y \mathrel{R} z$  because  $y \mathrel{R} z$  together with property (b) and transitivity of R imply the contradiction  $y \mathrel{R} x$ .

**Corollary 2.** Suppose that R contains no missed opportunities, and that offering  $\{z, w\}$  (where  $z \not R w \not R z$ ) does not miss an opportunity or take a risk at R. Then the proto-ranking R' induced by offering  $\{z, w\}$  contains no missed opportunities.

*Proof.* Let R, z, w and R' be as in the hypothesis of the lemma, and suppose toward a contradiction that there is a missed opportunity (x, y) in R'.

We claim that  $y \not R x$  and y R' x, so that Lemma 2 is applicable. It must be that  $y \not R x$ , for otherwise (x, y) would be a missed opportunity in R. That y R' x is immediate from the fact that (x, y) is a missed opportunity in R'.

It follows by Lemma 2 that  $\{z, w\} = \{x, y\}$ . But since (x, y) is a missed opportunity in R', there is an alternative  $z' \in \mathcal{X}$  such that  $x \succ z' \succ y$  and  $y \mathbb{R}' z' \mathbb{R}' x$ , and thus  $y \mathbb{R} z' \mathbb{R} x$  since  $R \subseteq R'$ . Thus offering  $\{z, w\} = \{x, y\}$  misses an opportunity at R—a contradiction.

Armed with Lemma 2 and Corollary 2, we are ready to prove that error-avoiding strategies are efficient.

**Proposition.** A strategy that never misses an opportunity or takes a risk is efficient.

Proof. Take a strategy  $\sigma$  that is not efficient, and suppose that it never misses an opportunity or takes a risk; we shall derive a contradiction. Since  $\sigma$  is not efficient, there exists a majority will W such that the outcome R of  $\sigma$  under W fails to be W-efficient, which is to say that  $x \succ y$ , x W y and y R x for some alternatives  $x, y \in \mathcal{X}$ .

Write  $\emptyset = R_0 \subseteq R_1 \subseteq \cdots \subseteq R_{T'} = R$  for the sequence of proto-rankings associated with the terminal history generated by  $\sigma$  and W. Let  $T \leq T'$  be the first period in which x, y are ranked  $(y \not R_{T-1} x \not R_{T-1} y \text{ and } y R_T x)$ . Since x W y and  $y R_T x$ , it cannot be that  $\{x, y\}$  is voted on in period T.

Because  $R_0 = \emptyset$  contains no missed opportunities and  $\sigma$  never misses an opportunity or takes a risk, Corollary 2 promises that  $R_{T-1}$  contains no missed opportunities. Thus by Lemma 2,  $\sigma$  must offer  $\{x, y\}$  in period T—a contradiction.

#### D.2 Proof that regret-free strategies avoid errors

The proof relies on two lemmata. For the first, recall from appendix A the notation  $[x, y]_R$  for *R*-order intervals.

**Lemma 3.** Given a pair of alternatives  $x, y \in \mathcal{X}$ , let R' be a ranking such that x R' y, and let W be a majority will that agrees with R' on every pair  $\{z, w\} \nsubseteq [x, y]_{R'}$ . Then the outcome under W of any strategy agrees with R' on every pair  $\{z, w\} \nsubseteq [x, y]_{R'}$ .

*Proof.* Let x, y, R' and W satisfy the hypothesis, and let R be the outcome under W of some strategy of the chair.

**Claim.** Let  $\{z, w\} \not\subseteq [x, y]_{R'}$  be such that (a) z R' w and (b) x R' z or w R' y. Then z R w.

Proof of the claim. Assume that x R' z; we omit the similar argument for the case w R' y. Suppose toward a contradiction that w R z. Label the alternatives  $[w, z]_R \equiv \{x_1, \ldots, x_K\}$  so that

$$w = x_1 R \cdots R x_K = z.$$

Since R is W-reachable, we have  $x_1 W \cdots W x_K$  by Observation 1 (appendix B.3, p. 22).

It must be that x R' w, since otherwise z R' w and the transitivity of R' would produce the contradiction z R' x. We thus have x R' w and  $w \notin [x, y]_{R'}$ , which means that y R' w.

Suppose that  $y \ R' \ x_k$  for every k < K. Then  $\{x_k, x_{k+1}\} \notin [x, y]_{R'}$  for every k < K. Since R' agrees with W on pairs  $\{z', w'\} \notin [x, y]_{R'}$ , it follows that  $x_k \ R' \ x_{k+1}$  for every k < K, whence  $w \ R' \ z$  by transitivity of R'—a contradiction.

Suppose instead that  $x_{k'} R' y$  for some k' < K, and let k be the smallest such k'. Since  $y R' w = x_1$ , it must be that k > 1. By definition of k, we have  $x_k R' y R' x_{k-1}$ . On the one hand, the transitivity of R' demands that  $x_k R' x_{k-1}$ . On the other hand, since  $\{x_{k-1}, x_k\} \not\subseteq [x, y]_{R'}$  (because  $x_{k-1} \not R' y$  by definition of k) and  $x_{k-1} W x_k$ , we must have  $x_{k-1} R' x_k$ —a contradiction.

Now fix a pair  $\{z, w\} \nsubseteq [x, y]_{R'}$  such that  $z \ R' w$ ; we must show that  $z \ R w$ . If either  $x \ R' z$  or  $w \ R' y$ , then  $z \ R w$  follows from the claim.

Suppose instead that z R' x and y R' w. Observe that  $\{z, x\} \not\subseteq [x, y]_{R'}$ , (a) z R' x and (b) x R' y. We may therefore apply the claim to  $\{z, x\}$  to obtain z R x. Similarly applying the claim to  $\{x, w\}$  yields x R w, whence z R w follows by transitivity of R.

**Lemma 4.** Let R be a proto-ranking, and let  $x, y, z \in \mathcal{X}$  be such that

$$\{x, y, z\}^2 \cap R \subseteq \{(x, z)\}.$$

Then there exists a ranking  $R' \supseteq R \cup \{(x, z), (z, y)\}$  such that  $[x, y]_{R'} = [x, z]_R \cup \{y\}.$ 

To interpret the conclusion, observe that the properties of R' are equivalent to the following: (a) x R' z and  $[x, z]_{R'} = [x, z]_R$ , and (b) z R' y and

 $[z, y]_{R'} = \{z, y\}.^{30}$  In words, Lemma 4 runs as follows. Suppose that a protoranking R ranks x above z, and has nothing else to say about  $\{x, y, z\}.^{31}$ Call the (possibly empty) set of alternatives ranked below x and above z(i.e.  $[x, z]_R \setminus \{x, z\}$ ) the 'in-between set'. The lemma asserts that R may be extended to a ranking R' that (i) adds nothing to the in-between set ( $[x, z]_{R'} = [x, z]_R$ ) and that (ii) ranks y immediately below z (z R' y and  $[z, y]_{R'} = \{z, y\}$ ).

The proof of Lemma 4 relies on the extension lemma from appendix C.

Proof. Let a proto-ranking R and alternatives  $x, y, z \in \mathcal{X}$  satisfy the hypothesis. Define  $A := [x, z]_R \cup \{y\}$ .

**Claim.** For any  $x', y' \in A$  such that x' R y', we have  $[x', y']_R \subseteq A$ .

Proof of the claim. Fix alternatives  $x', y' \in A$  with x' R y'. By definition of A, it suffices to show that  $\{x', y'\} \not\ni y$ . So suppose toward a contradiction that x' = y; the other case is similar. We have  $y' \neq y$  since x' R y' and R is irreflexive (being a proto-ranking). Since  $y' \in A$ , it follows that  $y' \in [x, z]_R$ . By x' R y' and the transitivity of R, we obtain y = x' R z. But  $y \not R z$  by hypothesis—a contradiction.

By the claim, the extension lemma in appendix C (p. 24) is applicable, so there exists a complete and transitive extension Q of the binary relation  $R \cup A^2$ . Since z' Q w' Q z' for any  $z', w' \in A$ , we have in particular that w Q y Q w for any  $w \in [x, z]_R$ . We may therefore obtain the desired ranking R' by appropriately breaking indifferences in Q.

With Lemmata 3 and 4 in hand, we are ready to prove that regret-free strategies avoid errors.

**Proposition.** A regret-free strategy never misses an opportunity or takes a risk.

 $<sup>^{30}{\</sup>rm The}$  formal proof of equivalence is as follows. If a ranking R' has properties (a) and (b), then x~R'~z~R'~y by (a) and (b), and

$[x,y]_{R'} = [x,z]_{R'} \cup [z,y]_{R'}$	
$= [x,z]_{R'} \cup \{y\}$	since $[z, y]_{R'} = \{y\}$ by (b)
$= [x, z]_R \cup \{y\}$	since $[x, z]_{R'} = [x, z]_R$ by (a).

If a ranking R' has the properties in the lemma, then x R' z R' y, so

$$[x, z]_{R'} \cup [z, y]_{R'} = [x, y]_{R'} = [x, z]_R \cup \{y\}.$$

Hence  $[x, z]_{R'} = [x, z]_R$  (whence (a)) and  $[z, y]_{R'} = \{z, y\}$  (whence (b)). <sup>31</sup>For simplicity, neglect the case  $\{x, y, z\}^2 \cap R = \emptyset$ . Proof. We shall prove the contra-positive. To that end, fix a strategy  $\sigma$  and a majority will W such that  $\sigma$  misses an opportunity or takes a risk under W. We shall construct a majority will W' such that the outcome R of  $\sigma$  under W' fails to be W'-unimprovable. In particular, we shall find a W'-reachable ranking  $R' \neq R$  that is more aligned with  $\succ$  than R.

Let T be the first period in which  $\sigma$  either misses an opportunity or takes a risk under W. Write  $R_{T-1}$  for the associated end-of-period-(T-1) proto-ranking, and let  $\{x, y\}$  be the pair offered in period T.

We shall consider three cases, based on the behaviour of  $\sigma$  under W in period T. By hypothesis,  $\{x, y\}$  either misses an opportunity or takes a risk at  $R_{T-1}$ . If  $\{x, y\}$  misses an opportunity, there is an alternative  $z \in \mathcal{X}$  such that  $x \succ z \succ y$  and  $y \not R_{T-1} z \not R_{T-1} x$ . It cannot be that  $x R_{T-1} z R_{T-1} y$ , as this would imply that  $x R_{T-1} y$ , contradicting the fact that  $\{x, y\}$  is offered in period T. Thus one of the following must hold:

- (b)  $x R_{T-1} z \not\!\!R_{T-1} y$ , or

If instead  $\{x, y\}$  takes a risk, then there is a  $z \in \mathcal{X}$  such that either

- (d)  $z \succ y$ ,  $x R_{T-1} z$ , and  $y \not R_{T-1} z$ , or
- (e)  $x \succ z$ ,  $z R_{T-1} y$ , and  $z \mathbb{R}_{T-1} x$ .

This yields three cases, as follows. Case 1 is (a). Case 2 encompasses both (b) and (d) under the (slightly more general) hypothesis that 'there exists a  $z \in \mathcal{X}$  such that  $z \succ y$ ,  $z \not \mathbb{R}_{T-1} y \not \mathbb{R}_{T-1} z$ , and  $x \not \mathbb{R}_{T-1} z$ '. Finally, case 3 encompasses (c) and (e) under the hypothesis that 'there exists a  $z \in \mathcal{X}$ such that  $x \succ z$ ,  $z \not \mathbb{R}_{T-1} x \not \mathbb{R}_{T-1} z$  and  $z \not \mathbb{R}_{T-1} y$ . Since cases 2 and 3 are analogous, we omit the proof for the latter.

Case 1:  $\exists z \in \mathcal{X}$  such that  $x \succ z \succ y$  and  $\{x, z, y\}^2 \cap R_{T-1} = \emptyset$ . By Lemma 4, there is a ranking R' such that

$$R' \supseteq R_{T-1} \cup \{(x, z), (z, y)\}$$
 and  $[x, y]_{R'} = \{x, y, z\}.$ 

Define W' to equal R', except that y W' x. Clearly W' is a majority will (total and asymmetric), and R' is W'-reachable by Observation 1 (appendix B.3, p. 22) since x, y are not R'-adjacent. Denote by R the outcome of  $\sigma$  under W'. It remains to show that  $R \neq R'$ , and that R' is more aligned with  $\succ$  than R.

For the former, since  $x \ R' y$ , it suffices to show that  $y \ R x$ . To this end, observe that that  $R_{T-1} \subseteq W'$ . Thus the history of length T-1 generated by  $\sigma$  and W' is the same as that generated by  $\sigma$  and W, which means in particular that  $\{x, y\}$  is offered in period T. Since  $y \ W' x$ , it follows that  $y \ R x$ , as desired.

To show that R' is more aligned with  $\succ$  than R, observe that W' agrees with R' on every pair  $\{w, w'\} \not\subseteq \{x, y, z\} = [x, y]_{R'}$ . It follows by Lemma 3 that R and R' agree on every pair  $\{w, w'\} \not\subseteq \{x, y, z\}$ . Since  $x \succ z \succ y$  and x R' z R' y, it follows that R' is more aligned with  $\succ$  than R.

Case 2:  $\exists z \in \mathcal{X}$  such that  $z \succ y$ ,  $z \not R_{T-1} y \not R_{T-1} z$  and  $x R_{T-1} z$ . We shall begin with an auxiliary ranking R'', then use it to construct our majority will W' and ranking R'. By Lemma 4, there is a ranking

$$R'' \supseteq R_{T-1} \cup \{(x, z), (z, y)\}$$

such that

$$[x, y]_{R''} = [x, z]_{R_{T-1}} \cup \{y\}.$$
(3)

Define

$$X \coloneqq \{ w \in [x, z]_{R_{T-1}} \setminus \{x\} : w \succ y \},\$$

and let W' be such that

- (i) w W' y for every  $w \in X$ ,
- (ii) y W' w for every  $w \in [x, z]_{R_{T-1}} \setminus X$ , and
- (iii) W' agrees with R'' on every other pair.

Denote by R the outcome of  $\sigma$  under the majority will W'.

Observe that (i)  $y \not \mathbb{R}_{T-1} w$  for every  $w \in X$  (since otherwise  $y \not \mathbb{R}_{T-1} w \ \mathbb{R}_{T-1} z$ , contradicting the case-2 hypothesis), (ii)  $w \not \mathbb{R}_{T-1} y$  for every  $w \in [x, z]_{R_{T-1}} \setminus X$  (otherwise either  $x = w \not \mathbb{R}_{T-1} y$  or  $x \not \mathbb{R}_{T-1} w \not \mathbb{R}_{T-1} y$ , whence  $x \not \mathbb{R}_{T-1} y$ ), and (iii)  $\mathcal{R}_{T-1} \subseteq \mathcal{R}''$ . Thus  $\mathcal{R}_{T-1} \subseteq W'$  by definition of the latter. It follows that the history of length T-1 generated by  $\sigma$  and W' is the same as that generated by  $\sigma$  and W, which means in particular that  $\{x, y\}$  is offered in period T. Since y W' x, we thus have  $y \not \mathbb{R} x$ .

Since  $X \subseteq [x, y]_{R_{T-1}} \subseteq [x, y]_{R''}$  (by definition of X and (3)), W' agrees with R'' on every pair  $\{w, w'\} \not\subseteq [x, y]_{R''}$ . It follows by Lemma 3 that R agrees with R'' on every pair  $\{w, w'\} \not\subseteq [x, y]_{R''}$ . Therefore y, x are R-adjacent, so that

$$[y, z]_R = \{y\} \cup [x, z]_R = \{y\} \cup [x, z]_{R_{T-1}} = [x, y]_{R''}.$$
(4)

It follows that  $X \cup \{x\} \subseteq [y, z]_R$ .

Define  $X' \coloneqq X \cup \{x\}$ , and label its elements  $X' \equiv \{a_1, \ldots, a_K\}$  so that  $a_1 \ R \ \cdots \ R \ a_K$ . Similarly label  $[y, z]_R \setminus X' \equiv \{b_1, \ldots, b_L\}$  so that  $b_1 \ R \ \cdots \ R \ b_L$ .<sup>32</sup> Let R' be the ranking that

- (I) agrees with R on any pair  $\{w, w'\} \not\subseteq [y, z]_R$ , and
- (II) ranks the elements of  $[y, z]_R$  as  $a_1 R' \cdots R' a_K R' b_1 R' \cdots R' b_L$ .

We have now constructed a majority will W' and a ranking R'. Recall that R is the outcome of  $\sigma$  under W'. It remains to show that

(dist) R' is distinct from R,

(alig) R' is more aligned with  $\succ$  than R, and

(feas) R' is W'-reachable.

For (dist), observe that since  $x \in X'$  and  $y \in [y, z]_R \setminus X'$ , we have  $x = a_k R' b_\ell = y$  for some k and  $\ell$ .<sup>33</sup> Since y R x, it follows that  $R' \neq R$ .

For (alig), fix a pair  $w, w' \in \mathcal{X}$  with  $w \ R \ w'$  and  $w' \ R' \ w$ ; we must show that  $w' \succ w$ . By definition of R', it must be that  $w' \in X' = X \cup \{x\}$  and that  $w \in [y, z]_R \setminus X'$ . Thus  $w' \succ y$  (by  $x \succ y$  and the definition of X) and either y = w or  $y \succ w$ , whence  $w' \succ w$  by transitivity of  $\succ$ .

It remains to establish (feas). To this end (recalling Observation 1 in appendix B.3, p. 22), fix an R'-adjacent pair  $w, w' \in \mathcal{X}$  with w' R' w; we must show that w' W' w. There are two cases.

Sub-case (2)(a):  $\{w, w'\} \not\subseteq [y, z]_R$ . Then  $w' \ R \ w$  by part (I) of the definition of R'. Since R agrees with R'' on any pair  $\{z', z''\} \not\subseteq [y, z]_R$ , it follows that  $w' \ R'' \ w$ . It therefore suffices to show that W' agrees with R'' on  $\{w, w'\}$ . Recalling the definition (i)–(iii) of W',

- If  $\{w, w'\} \not\supseteq y$ , then W' agrees with R'' on  $\{w, w'\}$  by (iii).
- If  $w' = y \in [y, z]_R$ , then  $w \notin [y, z]_R = [x, z]_{R_{T-1}} \supseteq X$  by (4), so neither (i) nor (ii) applies to the pair  $\{w, y\} = \{w, w'\}$ . Thus by (iii), W' agrees with R'' on  $\{w, w'\}$ .
- If  $w = y \in [y, z]_R$ , then  $w' \notin [y, z]_R \supseteq [x, z]_{R_{T-1}} \supseteq X$  by (4), so neither (i) nor (ii) applies to the pair  $\{y, w'\} = \{w, w'\}$ . Thus by (iii), W' agrees with R'' on  $\{w, w'\}$ .

 $<sup>^{32}[</sup>y,z]_R \setminus X'$  is non-empty since y belongs to it.

<sup>&</sup>lt;sup>33</sup>In fact,  $k = \ell = 1$  since y is R-highest in  $[y, z]_R$  and (recall) y, x are R-adjacent.

Sub-case (2)(b):  $\{w, w'\} \subseteq [y, z]_R$ . Recall the definition (i)–(iii) of W'. Recall also part (II) of the definition of R'. Observe that  $y = b_1$  since y is R-highest in  $[y, z]_R$ , and  $y \notin X' = X \cup \{x\}$ . Furthermore,  $x = a_1$  since y, x are R-adjacent (recall (4)) and  $x \in X$ . Finally, remark that  $K \ge 2$  since  $z \in X' = X \cup \{x\}$  and  $z \neq x$ .

Suppose first that  $w' = y = b_1$ . Then since w', w are R'-adjacent with w' R' w, we have  $w = b_2 \notin X' \supseteq X$ . Thus part (i) does not apply to the pair  $\{w, y\} = \{w, w'\}$ . Thus part (ii) applies, yielding w' = y W' w.

Suppose instead that  $w = y = b_1$ . Then since w', w are R'-adjacent with w' R' w, we have  $w' = a_K \in X' = X \cup \{x\}$ . Since  $x = a_1$  and  $K \ge 2$ , we have  $w' \ne x$ . Thus  $w' \in X$ , so that (i) applies to the pair  $\{y, w'\} = \{w, w'\}$ , yielding w' W' y = w.

Finally, suppose that  $\{w, w'\} \not\supseteq y$ . Then W' and R'' agree on the pair  $\{w, w'\}$  by (iii), so it suffices to show that w' R'' w.

Since  $b_1 = y \notin \{w, w'\}$ , we have either  $\{w, w'\} \subseteq X'$  or  $\{w, w'\} \subseteq [y, z]_R \setminus X'$ . Thus R and R' agree on  $\{w, w'\}$  by part (I) of the definition of R', so that w' R w.

Now label  $[w', w]_R \equiv \{z_1, \ldots, z_J\}$  so that  $z_1 \ R \cdots R \ z_J$ . Since R is W'-reachable, we have  $z_1 \ W' \cdots W' \ z_J$  by Observation 1 (appendix B.3, p. 22). By the hypotheses  $w' \in [y, z]_R$  and  $w' \neq y$ , we must have  $y \ R \ w'$  and thus  $y \notin [w', w]_R$ . This together with the fact that  $z_j \ W' \ z_{j+1}$  for each j < J implies, via part (iii) of the definition of W', that  $z_j \ R'' \ z_{j+1}$  for each j < J. It follows by transitivity of R'' that  $w' = z_1 \ R'' \ z_J = w$ , as desired.

# E Proofs of Propositions 1 and 2 (§6, pp. 16 and 18)

In this appendix, we establish tightness for the characterisations of regretfreeness in Theorems 2 and 3. We begin in §E.1 with a lemma, then use it to deduce Proposition 2 (§E.2) and Proposition 1 (§E.3).

#### E.1 A lemma

**Definition 11.** Given a proto-ranking R and alternatives  $x \succ y$  and  $z \neq w$ , say that  $\{z, w\}$  makes  $\{x, y\}$  an error at R iff both  $x \not R y \not R x$  and  $z \not R w \not R z$ , and one of the following holds:

 $- x \succ z \succ y, y \not R z \not R x, \text{ and } w \in \{x, y\}.$  $- z \succ y, x R z \text{ and } w = y.$  $- x \succ z, z R y \text{ and } w = x.$ 

If  $\{z, w\}$  makes  $\{x, y\}$  an error, then offering  $\{x, y\}$  either misses an opportunity or takes a risk at R, and the chair 'should' offer  $\{z, w\}$  instead.<sup>34</sup>

Recall from appendix D.1 (p. 26) the definition of a missed opportunity.

**Lemma 5.** Let R be a proto-ranking, and let  $A \subseteq \mathcal{X}^2$  be a non-empty set of pairs of distinct alternatives. Suppose that for any pair  $\{x, y\} \in A$ , there is a pair  $\{z, w\} \in A$  that makes  $\{x, y\}$  an error at R. Then R contains a missed opportunity.

Proof. Let R and A satisfy the hypothesis. Then there is a pair  $\{z, w\} \in A$ and another pair  $\{z', w'\} \in A$  that makes  $\{z, w\}$  an error at R. Assume (wlog) that  $z \succ w$  and  $z' \succ w'$ . Since  $\{z, w\} \neq \{z', w'\}$ , we must have either  $z \neq z'$  or  $w \neq w'$ . Assume that  $z \neq z'$ ; the case  $w \neq w'$  is similar.

**First claim.** There exists a sequence  $(x_t)_{t=1}^T$  in  $\mathcal{X}$  with  $T \ge 2$  and  $x_1 \ne x_2$  such that for every  $t \le T$ , writing  $x_{T+1} \coloneqq x_1$ ,

- (i) if  $x_t \succ x_{t+1}$  then  $x_{t+1} \not \mathbb{R} x_t$ , and
- (ii) if  $x_{t+1} \succ x_t$  then  $x_t R x_{t+1}$ .

Proof of the first claim. Define  $\{x_1, y_1\} := \{z, w\}$  and  $\{x_2, y_2\} := \{z', w'\}$ . By the hypothesis of the lemma, there is a pair  $\{x_3, y_3\} \in A$  with (wlog)  $x_3 \succ y_3$  that makes  $\{x_2, y_2\}$  an error at R, a  $\{x_4, y_4\} \in A$  with  $x_4 \succ y_4$ that makes  $\{x_3, y_3\}$  an error at R, and so on. Since A is finite,  $\{x_1, y_1\}$ makes  $\{x_T, y_T\}$  an error for some  $T \in \mathbb{N}$ . We have  $T \ge 2$  and  $x_1 \ne x_2$  by construction, and (i)–(ii) must hold because  $\{x_{t+1}, y_{t+1}\}$  makes  $\{x_t, y_t\}$  an error at R.

Let  $(x_t)_{t=1}^T$  be a minimal sequence satisfying the conditions of the first claim (one with no strict subsequence that satisfies the conditions).

**Second claim.**  $x_t \neq x_s$  for all distinct  $t, s \in \{1, \ldots, T\}$ .

Proof of the second claim. Suppose toward a contradiction that  $x_t = x_{t+1}$ ; then the sequence  $x_1, \ldots, x_{t-1}, x_{t+1}, \ldots, x_T$  satisfies the conditions of the first claim, contradicting the minimality of  $(x_t)_{t=1}^T$ . Assume for the remainder that  $x_t \neq x_{t+1}$  for every  $t \in \{1, \ldots, T\}$ .

Suppose toward a contradiction that  $x_t = x_s$ , where t + 1 < s. Then the sequence  $x_{t+1}, \ldots, x_s$  satisfies the conditions of the first claim, which is absurd since  $(x_t)_{t=1}^T$  is minimal.

<sup>&</sup>lt;sup>34</sup>This is heuristic, as offering  $\{z, w\}$  might itself miss an opportunity or take a risk at R.

In light of the second claim, we may re-label the sequence  $(x_t)_{t=1}^T$  so that  $x_1 \succ x_t$  for every  $t \in \{2, \ldots, T\}$ . Let  $t' \leq T$  be the least  $t \geq 2$  such that  $x_T \succ \cdots \succ x_t$ . (So t' = T exactly if  $x_t \succ x_T$  for every t < T.) We shall show that  $\{x_1, x_{t'}\}$  is a missed opportunity in R; in particular, that  $t' \geq 3$  and

- (a)  $x_1 \succ x_{t'-1} \succ x_{t'}$ ,
- (b)  $x_{t'} R x_1$ , and
- (c)  $x_{t'} \not \mathbb{R} x_{t'-1} \not \mathbb{R} x_1$ .

For (b), if t' = T then  $x_{t'} = x_T R x_1$  by property (ii), and if not then  $x_{t'} R \cdots R x_T R x_1$  by property (ii), whence  $x_{t'} R x_1$  by transitivity of R. The second half  $x_{t'-1} \succ x_{t'}$  of (a) holds by definition of t'. The first half  $x_{t'} \not R x_{t'-1}$  of (c) then follows by property (i). Since  $x_{t'} R x_1$ , it follows that  $t' - 1 \neq 1$ , which is to say that  $t' \geq 3$ . The first half  $x_1 \succ x_{t'-1}$  of (a) then holds by construction. Finally, the second half  $x_{t'-1} \not R x_1$  of (c) must hold since otherwise the sequence  $(x_t)_{t=1}^{t'-1}$  would satisfy the conditions of the first claim, in contradiction with the minimality of  $(x_t)_{t=1}^T$ .

## E.2 Proof of Proposition 2 (p. 18)

At a history at which the chair has committed no errors, the proto-ranking clearly contains no missed opportunities. The following therefore implies Proposition 2.

**Proposition 2\*.** Let R be a non-total proto-ranking containing no missed opportunities. Then there exist distinct  $x, y \in \mathcal{X}$  such that  $x \not R y \not R x$  and offering a vote on  $\{x, y\}$  neither misses an opportunity nor takes a risk at R.

Proof. Let R be a non-total proto-ranking, and suppose that for any distinct  $x, y \in \mathcal{X}$  with  $x \not R y \not R x$ , offering a vote on  $\{x, y\}$  either misses an opportunity or takes a risk at R. We shall show that R contains a missed opportunity.

Let A be the set of all pairs  $\{x, y\} \subseteq \mathcal{X}$  with  $x \neq y$  and  $x \not R y \not R x$ . The set A is non-empty since R is not total. By hypothesis, for any  $\{x, y\} \in A$ , offering  $\{x, y\}$  either misses an opportunity or takes a risk at R, implying that some  $\{z, w\} \in A$  makes  $\{x, y\}$  an error at R. It follows by Lemma 5 (§E.1, p. 35) that R contains a missed opportunity.

#### E.3 Proof of Proposition 1 (p. 16)

**Lemma 6.** Fix a majority will W, let R be a W-efficient ranking, and let  $R' \subseteq R$  be a non-total proto-ranking containing no missed opportunities.

Then there exist distinct  $x, y \in \mathcal{X}$  such that  $x \not R' y \not R' x$ , W and R agree on  $\{x, y\}$ , and offering  $\{x, y\}$  does not miss an opportunity or take a risk at R'.

Proof of Proposition 1. Fix a majority will W and a W-efficient ranking R. By Proposition 2 (p. 18; already proved), it suffices to find a terminal history  $((x_t, y_t))_{t=1}^T$ , with associated proto-rankings  $(R_t)_{t=0}^T$ , <sup>35</sup> such that

- for every  $t \in \{1, \ldots, T\}$ ,  $x_t W y_t$  and  $x_t R y_t$ , and
- for every  $t \in \{2, \ldots, T\}$ , offering  $\{x_t, y_t\}$  does not miss an opportunity or take a risk at  $R_{t-1}$ .

Such a terminal history is obtained by repeatedly applying Lemma 6.

Proof of Lemma 6. We shall prove the contra-positive. Fix a majority will W and a W-efficient ranking R, and let  $R' \subseteq R$  be a non-total proto-ranking. Suppose that for any distinct  $x, y \in \mathcal{X}$  with  $x \not R' y \not R' x$  such that W and R agree on  $\{x, y\}$ , offering  $\{x, y\}$  misses an opportunity or takes a risk at R'. We will show that R' contains a missed opportunity.

Let A be the set of all pairs  $\{x, y\} \in \mathcal{X}^2$  such that  $x \succ y, x \not R' y \not R' x$ , and there is no  $z \in \mathcal{X}$  such that x R z R y. The set A is non-empty since it includes any R-adjacent pair  $\{x, y\}$  with  $x \not R' y \not R' x$ , and there must be such a pair since R' is non-total and  $R' \subseteq R$ . By Lemma 5 (§E.1, p. 35), it suffices to show that for any pair  $\{x, y\} \in A$ , there is a pair  $\{z, w\} \in A$  that makes  $\{x, y\}$  an error at R'.

So fix a pair  $\{x, y\} \in A$ . We claim that W and R must agree on  $\{x, y\}$ . If x, y are R-adjacent, then this holds by Observation 1 (appendix B.3, p. 22) since R is W-reachable. If x, y are not R-adjacent, then since no  $z \in \mathcal{X}$  satisfies x R z R y, it must be that y R x. Since  $x \succ y$  and R is W-efficient, it follows that y W x, so that W and R agree on  $\{x, y\}$ .

It follows from the (contra-positive) hypothesis that offering  $\{x, y\}$  either misses an opportunity or takes a risk at R'. Consider each in turn.

Case 1:  $\{x, y\}$  misses an opportunity. In this case there is a  $z \in \mathcal{X}$  satisfying  $x \succ z \succ y$  and  $y \not R' z \not R' x$ . Since  $\{x, y\} \in A$ , we must have either  $z \ R x$  or  $y \ R z$ . Assume that  $z \ R x$ ; the case  $y \ R z$  is analogous. Since  $R' \subseteq R$ , we have  $x \not R' z \not R x$ . Thus the pair  $\{x, z\}$  lives in A and makes  $\{x, y\}$  an error at R'.

Case 2:  $\{x, y\}$  takes a risk. Assume that there is a  $z \in \mathcal{X}$  such that  $z \succ y$ , x R' z and y R' z; the case in which  $x \succ z$ , z R' y and z R' x is similar. Then  $\{y, z\}$  makes  $\{x, y\}$  an error at R'. To see that  $\{y, z\}$  belongs to A, observe

<sup>&</sup>lt;sup>35</sup>Recall that  $R_0 = \emptyset$  and  $R_t = \operatorname{tr}\left(\bigcup_{s=1}^t \{(x_s, y_s)\}\right)$  for each  $t \in \{1, \ldots, T\}$ .

that (i) x R z since x R' z and  $R' \subseteq R$ , that (ii) y R z by (i) and  $\{x, y\} \in A$ , and finally that (iii) z R' y since otherwise x R' z and the transitivity of R' would imply the falsehood x R' y.

# F Proof of Theorem 4 (§7, p. 20)

In this appendix, we first prove Theorem 4 using two lemmata (§F.1), and then prove these lemmata (§F.2–§F.4). Throughout, we label the alternatives  $\mathcal{X} \equiv \{1, \ldots, n\}$  so that  $1 \succ \cdots \succ n$ .

## F.1 Proof using lemmata

**Definition 12.** Let  $\Sigma_0$  be the set of all strategies. For every integer  $k \in \{1, \ldots, n-2\}$ , let  $\Sigma_k$  be the set of all strategies  $\sigma$  with the following property: for any majority will W and alternative  $j \leq k$ , labelling the alternatives  $\{j+1,\ldots,n\} \equiv \{x_{j+1},\ldots,x_n\}$  as

$$x_{j+1} R^{\sigma}(W) \cdots R^{\sigma}(W) x_n,$$

the first vote involving j that  $\sigma$  offers under W is on  $\{j, x_{j+1}\}$ ; if j loses, then a second vote involving j is offered, namely on  $\{j, x_{j+2}\}$ ; if j loses again, then a third vote involving j is offered, namely on  $\{j, x_{j+3}\}$ ; and so on.

The definition of  $\Sigma_{n-2}$  describes a natural generalisation of insertion sort: for each alternative j, given how the  $\succ$ -worse alternatives  $\{x_{j+1}, \ldots, x_n\}$ are ultimately ranked, the same votes involving j are offered, in the same order, though not necessarily in adjacent periods.<sup>36</sup> Each  $\Sigma_k$  for k < n-2is defined by the same property restricted to those alternatives j that are  $\succ$ -better than or equal to k, so that  $\Sigma_0 \supseteq \Sigma_1 \supseteq \cdots \supseteq \Sigma_{n-2}$ .

**Lemma 7.** A strategy is outcome-equivalent to insertion sort iff it belongs to  $\Sigma_{n-2}$ .

**Lemma 8.** Given  $k \in \{1, \ldots, n-2\}$ , a strategy  $\sigma \in \Sigma_{k-1}$  is best for k among  $\Sigma_{k-1}$  iff it belongs to  $\Sigma_k$ .

Proof of Theorem 4. Let  $B_0 = \Sigma_0$  be the set of all strategies, and for each  $k \in \{1, \ldots, n-2\}$ , let  $B_k$  be the set of strategies in  $B_{k-1}$  that are best for k among  $B_{k-1}$ .<sup>37</sup> A lexicographic strategy is precisely one that lives in  $B_{n-2}$ .

<sup>&</sup>lt;sup>36</sup>Note a subtlety in the definition: although we label  $\{x_{j+1}, \ldots, x_n\}$  according to the outcome  $R^{\sigma}(W)$  of  $\sigma$  under W, a strategy in  $\Sigma_k$  need not (as insertion sort would) have totally ranked  $\{j + 1, \ldots, n\}$  before offering votes involving j.

<sup>&</sup>lt;sup>37</sup>Let  $B_k := \emptyset$  if  $B_{k-1}$  is empty. (It is in fact non-empty, but we haven't proved it yet.)

By Lemma 7, a strategy is outcome-equivalent to insertion sort iff it lives in  $\Sigma_{n-2}$ ; so what must be shown is that  $\Sigma_{n-2} = B_{n-2}$ . We shall prove the stronger claim that  $\Sigma_k = B_k$  for each  $k \in \{0, \ldots, n-2\}$  by (weak) induction on k. The base case k = 0 holds by definition of  $\Sigma_0$  and  $B_0$ . For the induction step, suppose that  $\Sigma_{k-1} = B_{k-1}$ ; then  $\Sigma_k = B_k$  by Lemma 8.

The remainder of this appendix is devoted to proving Lemmata 7 and 8. We begin in F.2 with two preliminary results, then prove Lemma 8 in F.3 and Lemma 7 in F.4.

#### F.2 Preliminary results

The following lemma is used in the proof of Lemma 7.

**Lemma 9.** Given a  $k \in \{1, ..., n-2\}$ , consider a strategy  $\sigma \in \Sigma_k$ , a majority will W, an alternative  $j \leq k$  and some  $m \in \{1, ..., n-j\}$ . Suppose that under W,  $\sigma$  offers at least m votes involving j, and that j loses the first m-1 of these. Let  $\ell$  be the  $m^{\text{th}}$  alternative pitted against j, and let R be the proto-ranking associated with the history after which the vote on  $\{j, \ell\}$ occurs. Then  $\ell R i$  for any  $i \neq \ell$  such that i > j and j did not lose against iprior to the vote on  $\{j, \ell\}$ .

Proof. Suppose toward a contradiction that  $\ell \not R$  *i* for some  $i \neq \ell$  with i > j such that *j* did not lose to *i* prior to the vote on  $\{j, \ell\}$ . Then there exists a ranking W' such that  $R \subseteq W'$  and  $i W' \ell$ . (Note that a ranking is precisely a transitive majority will.) Being a ranking, W' is the only W'-reachable ranking (by Observation 1 in appendix B.3, p. 22), so that  $R^{\sigma}(W') = W'$ , and in particular  $i R^{\sigma}(W') \ell$ .

Let T be the period in which  $\sigma$  offers  $\{j, \ell\}$  (i.e. the  $m^{\text{th}}$  vote involving j) under W. Since  $R \subseteq W'$ , the history of length T-1 generated by  $\sigma$  and W' is the same as that generated by  $\sigma$  and W. So in particular, under W',  $\sigma$  offers  $\{j, \ell\}$  in period T,  $\sigma$  does not offer  $\{j, i\}$  in an earlier period, and j does not win a vote in any earlier period. Since  $\sigma \in \Sigma_k$ , it follows that  $\ell R^{\sigma}(W') i$ , a contradiction.

The proof of Lemma 8 relies on the following.

**Lemma**  $\natural$ . Given  $k \in \{2, \ldots, n-2\}$ , if  $N_k^{\sigma}(W) > 0$  for some  $\sigma \in \Sigma_{k-1}$  and majority will W, then prior to winning its first vote, k is pitted only against  $\succ$ -worse alternatives.

Proof. We prove the contra-positive. Let  $k \in \{2, \ldots, n-2\}$ ,  $\sigma \in \Sigma_{k-1}$  and a majority will W be such that  $\sigma$  pits k against some j < k under W and k wins no vote before the one against j; we must show that  $N_k^{\sigma}(W) = 0$ . Let R be the proto-ranking associated with the history after which the vote on  $\{j, k\}$  occurs. It suffices to show that  $\ell R k$  for all  $\ell > k$ .

**Claim.** j wins no vote prior to the one against k.

Proof of the claim. Suppose toward a contradiction that the first alternative against which j wins is  $\ell \neq k$ . Then  $j \ R \ \ell$ . Since  $\sigma \in \Sigma_{k-1}$ , the vote on  $\{j, \ell\}$  is the  $m^{\text{th}}$  involving j, for some m < n - j. It follows by Lemma 9 (above) that  $\ell \ R \ k$ , which together with  $j \ R \ \ell$  and the transitivity of R yields  $j \ R \ k$ . On the other hand, since  $\{j, k\}$  is offered after a history with proto-ranking R, we must have  $j \ R \ k \ R \ j$ . Contradiction!

Fix an  $\ell > k$ ; we shall show that  $\ell \ R \ k$ . Since  $\sigma \in \Sigma_{k-1}$ , the vote on  $\{j, k\}$  is the  $m^{\text{th}}$  involving j, for some m < n - j. It must be that  $\sigma$  offers  $\{j, \ell\}$  prior to  $\{j, k\}$ , since otherwise Lemma 9 would yield  $k \ R \ \ell$ , contradicting the hypothesis that k wins no votes prior to the one against j. Since j wins no vote prior to the one against k and m < n - j, Lemma 9 yields  $\ell \ R \ k$ , as desired.

## F.3 Proof of Lemma 8 (§F.1, p. 38)

Let  $\mathcal{W}$  denote the set of all majority wills (total and asymmetric relations) on  $\mathcal{X}$ . We shall use the probabilistic notation  $\Pr(E|F) := |E \cap F|/|F|$  for the fraction of majority wills in  $F \subseteq \mathcal{W}$  that belong to  $E \cap F \subseteq F$ , and similarly  $\Pr(E) := \Pr(E|\mathcal{W})$ . This corresponds the thought experiment in which the majority will W is drawn uniformly at random from  $\mathcal{W}$ .

We must establish that membership of  $\Sigma_k$  is necessary and sufficient for being best for k among  $\Sigma_{k-1}$ . We prove sufficiency and necessity in turn, making use of Lemma  $\natural$  from the previous section.

Proof of sufficiency. Fix k, a strategy  $\sigma \in \Sigma_{k-1}$  and an  $m \in \{1, \ldots, n-k\}$ . We shall derive an upper bound for  $\Pr(N_k^{\sigma} \ge m)$ , then show that it is attained if  $\sigma \in \Sigma_k$ .

For each  $\ell \in \{1, \ldots, n-k\}$ , let  $F_{\ell} \subseteq \mathcal{W}$  be the set of majority wills under which  $\sigma$  offers at least  $\ell$  votes involving alternative k, with alternative k losing the first  $\ell - 1$  of these and winning the  $\ell^{\text{th}}$ .

**Claim.**  $\Pr(F_{\ell}) \leq 1/2^{\ell}$  for each  $\ell \in \{1, \ldots, n-k\}$ , with equality if  $\sigma \in \Sigma_k$ . Proof of the claim. A majority will W lies in  $F_{\ell}$  iff under  $\sigma$  and W,

- alternative k loses its first vote (probability 1/2),
- a second vote involving alternative k occurs (probability  $\leq 1$ , with equality if  $\sigma \in \Sigma_k$ ) and k loses (probability 1/2),
  - ...
- an  $(\ell 1)^{\text{th}}$  vote involving alternative k occurs (probability  $\leq 1$ , with equality if  $\sigma \in \Sigma_k$ ) and k again loses (probability 1/2), and
- an  $\ell^{\text{th}}$  vote involving alternative k occurs (probability  $\leq 1$ , with equality if  $\sigma \in \Sigma_k$ ) and k wins (probability 1/2).

Thus

$$\Pr(F_{\ell}) \leq \frac{1}{2} \times \left(1 \times \frac{1}{2}\right)^{\ell-1} = \frac{1}{2^{\ell}}, \quad \text{with equality if } \sigma \in \Sigma_k.$$

By Lemma  $\natural$  (§F.2, p. 39), if  $N_k^{\sigma}(W) > 0$ , then prior to winning its first vote, k was only pitted against  $\succ$ -worse alternatives. There are n-k of these:  $\{k+1,\ldots,n\}$ . Thus if  $N_k^{\sigma}(W) \ge m$  holds, then alternative k cannot have lost strictly more than n-m-k votes and must have won at least one—so in particular,  $W \in F_1 \cup \cdots \cup F_{n-k-m+1}$ . Thus

$$\Pr(N_1^{\sigma} \ge m) = \sum_{\ell=1}^{n-k-m+1} \Pr(F_\ell) \Pr(N_1^{\sigma} \ge m | F_\ell)$$
$$\le \sum_{\ell=1}^{n-k-m+1} \Pr(F_\ell) \qquad (\sharp)$$

$$\leq \sum_{\ell=1}^{n-k-m+1} \frac{1}{2^{\ell}},\tag{b}$$

where  $(\flat)$  holds by the claim.

Now suppose that  $\sigma \in \Sigma_k$ ; we shall show that  $(\sharp)$  and  $(\flat)$  hold with equality, so that  $\sigma$  attains the bound. For  $(\flat)$ , this follows from the claim. For  $(\sharp)$ , fix an  $\ell \in \{1, \ldots, n-k-m+1\}$  and a  $W \in F_\ell$ ; we must show that  $N_k^{\sigma}(W) \geq m$ . Label  $\{k+1, \ldots, n\} \equiv \{x_{k+1}, \ldots, x_n\}$  so that

$$x_{k+1} R^{\sigma}(W) \ldots R^{\sigma}(W) x_n$$

Since  $W \in F_{\ell}$ , we have by definition of  $\Sigma_k$  that  $k \ R^{\sigma}(W) \ x_{\ell+1}$ . Thus  $k \ R^{\sigma}(W) \ x_{\ell'}$  for each  $\ell' \in \{\ell+1,\ldots,n\}$ , so that  $N_k^{\sigma}(W) \ge n-\ell \ge m$ .

Proof of necessity. Take a strategy  $\sigma$  in  $\Sigma_{k-1} \setminus \Sigma_k$ . Since  $\sigma$  belongs to  $\Sigma_{k-1}$ , it satisfies the inequalities  $(\sharp)$  and  $(\flat)$  in the sufficiency argument. Suppose that one or the other holds strictly for some  $m \in \{1, \ldots, n-k\}$ , so that  $\sigma$  fails to attain the bound in the sufficiency proof. Since any  $\sigma' \in \Sigma_k$  attains the bound for every m by the (just-proved) sufficiency part, it follows that  $\sigma$  is not best for k among  $\Sigma_{k-1}$ . It therefore suffices to find an  $m \in \{1, \ldots, n-k\}$  such that either  $(\sharp)$  or  $(\flat)$  holds strictly.

Since  $\sigma \in \Sigma_{k-1} \setminus \Sigma_k$ , there is a majority will W such that, labelling the alternatives  $\{k+1,\ldots,n\} \equiv \{x_{k+1},\ldots,x_n\}$  so that

$$x_{k+1} R^{\sigma}(W) \ldots R^{\sigma}(W) x_n,$$

one of the following holds under W:

- (a)  $\sigma$  pits k against some  $j \neq x_{k+1}$  prior to pitting it against  $x_{k+1}$ .
- (b) For some  $\ell \in \{k+1, \ldots, n-k\}$ ,  $\sigma$  offers at least  $\ell$  votes involving k, the first  $\ell 1$  of which are against  $x_{k+1}, \ldots, x_{\ell}$  and are all lost by k, and the  $\ell^{\text{th}}$  of which is against some  $j \neq x_{\ell+1}$ .
- (c) For some  $\ell \in \{k+1, \ldots, n-k\}$ ,  $\sigma$  offers exactly  $\ell 1$  votes involving k, against  $x_{k+1}, \ldots, x_{\ell}$ , each of which is lost by k.

Case (a). We shall exhibit a majority will  $W' \in F_1$  such that  $N_k^{\sigma}(W') < n-k$ , so that  $(\sharp)$  holds strictly for m = n-k. This is trivial if  $N_k^{\sigma}(W) = 0$ (let W' := W), so suppose that  $N_k^{\sigma}(W) > 0$ . Let  $j \neq x_{k+1}$  be the first alternative pitted against k by  $\sigma$  under W, and let R be the proto-ranking associated with the history after which this occurs. Clearly  $j \not R k \not R x_{k+1}$ . Since  $\sigma \in \Sigma_{k-1}$  and  $N_k^{\sigma}(W) > 0$ , Lemma  $\natural$  (§F.2, p. 39) implies that j > k. So  $x_{k+1} R^{\sigma}(W) j$  (since  $j \in \{k+1,\ldots,n\} = \{x_{k+1},\ldots,n\}$  and  $j \neq x_{k+1}$ ), and thus  $j \not R x_{k+1}$ .

It follows that there is a ranking W' such that  $R \subseteq W'$  and  $x_{k+1} W'$  $k W' j.^{38}$  (Note that a ranking is precisely a transitive majority will.) Let Tbe the period in which  $\sigma$  offers  $\{j, k\}$  under W. Since  $R \subseteq W'$ , the history of length T-1 generated by  $\sigma$  and W' is the same as that generated by  $\sigma$ and W, and thus  $\{j, k\}$  is the first pair involving k that  $\sigma$  offers under W'. Thus  $W' \in F_1$  since k W' j. Being a ranking, W' is the only W'-reachable

<sup>&</sup>lt;sup>38</sup>To see why, observe that the transitive closure R' of  $R \cup \{(x_{k+1}, k)\}$  is a protoranking since  $k \not R x_{k+1}$ . Since  $j \neq x_{k+1}$  and  $j \not R x_{k+1}$ , we have  $j \not R' k$  by Observation 3 (appendix D.1, p. 25), and thus the transitive closure R'' of  $R' \cup \{(k, j)\}$  is also a protoranking. Let W' be any ranking that contains R''.

ranking (by Observation 1 in appendix B.3, p. 22), so  $R^{\sigma}(W') = W'$ . Thus  $N_k^{\sigma}(W') < n - k$  since  $x_{k+1} W' k$ .

Case (b). If j > k, then case-(a) argument yields a  $W' \in F_{\ell}$  such that  $N_k^{\sigma}(W') < n - k - \ell + 1$ , so that ( $\sharp$ ) holds strictly for  $m = n - k - \ell + 1$ . Suppose instead that j < k. Let W' be the majority will that agrees with W on every pair, except that k W' j. Clearly  $W' \in F_{\ell}$ . By (the contra-positive of) Lemma  $\sharp$ , we have  $N_k^{\sigma}(W') = 0$ . Thus  $\Pr(N_k^{\sigma} \ge 1|F_{\ell}) < 1$ , so that ( $\sharp$ ) holds strictly for m = 1.

Case (c). Let E be the set of majority wills under which  $\sigma$  offers at least  $\ell - 1$  votes involving k, the first  $\ell - 1$  of which k loses. Then  $\Pr(E) > 0$ , and the probability that  $\sigma$  offers at least  $\ell$  votes conditional on E is strictly less than 1. The argument for the claim in the sufficiency proof therefore yields  $\Pr(F_{\ell}) < 1/2^{\ell}$ , so that (b) holds strictly for e.g. m = 1.

## F.4 Proof of Lemma 7 (§F.1, p. 38)

We must show that membership of  $\Sigma_{n-2}$  is necessary and sufficient for outcome-equivalence to insertion sort. For the sufficiency part, we shall make use of Lemma 9 in §F.3.

Proof of sufficiency. Let  $\sigma \in \Sigma_{n-2}$ , fix a majority will W, and let R(R') be the outcome of  $\sigma$  (of insertion sort) under W. We will show for each  $k \in \{n-2,\ldots,1\}$  that R agrees with R' on  $\{k,\ldots,n\}$ , using induction on k. For the base case k = n-2, let  $\{j,\ell\}$  be the first pair offered by  $\sigma$ , where  $j < \ell$ ; it suffices to show that j = n - 1. Suppose to the contrary; then by Lemma 9 (§F.3, p. 39), prior to the first vote,  $\ell$  is already ranked above every i > j such that  $i \neq \ell$ , which is absurd. The induction step is immediate from the fact that  $\sigma \in \Sigma_{n-2}$ .

The proof of necessity refers to the argument for Lemma 8 in §F.3 above.

Proof of necessity. Let  $\sigma \notin \Sigma_{n-2}$ ; we must show that it is not outcomeequivalent to insertion sort. Note that there is a (unique)  $k \in \{1, \ldots, n-2\}$ such that  $\sigma \in \Sigma_{k-1} \setminus \Sigma_k$ . Recall the bound in the proof of the sufficiency part of Lemma 8 (§F.3 above). Since insertion sort belongs to  $\Sigma_k$ , it attains the bound for every  $m \in \{1, \ldots, n-k\}$  by the Lemma 8 sufficiency argument, and thus all of its outcome-equivalents do. By the Lemma 8 necessity argument,  $\sigma$  fails to attain the bound for some  $m \in \{1, \ldots, n-k\}$ , so is not among the outcome-equivalents of insertion sort.

## G The amendment procedure and insertion sort

In this appendix, we link insertion sort to the amendment procedure, which figures prominently in the agenda-setting literature (see §1.1 for references):

Amendment procedure. Label the alternatives  $\mathcal{X} \equiv \{1, \ldots, n\}$  so that  $1 \succ \cdots \succ n$ . First, pit n-1 against n. Then pit n-2 against the winner. Then pit n-3 against the previous round's winner. And so on. Call the winner of the final round the final winner.

The amendment procedure is designed to choose a single alternative. In particular, the final winner is ranked top, since any other alternative either lost a vote to it, or lost a vote to an alternative that lost to it, etc. The natural way to obtain a ranking is to apply amendment recursively:

**Recursive amendment.** Label the alternatives  $\mathcal{X} \equiv \{1, \ldots, n\}$  so that  $1 \succ \cdots \succ n$ . First, run the amendment procedure on  $\mathcal{X}$ , and write  $y_1$  for the final winner. Next, run the amendment procedure on  $\mathcal{X} \setminus \{y_1\}$ , writing  $y_2$  for the final winner.<sup>39</sup> Then run the amendment procedure on  $\mathcal{X} \setminus \{y_1, y_2\}$  to obtain a final winner  $y_3$ . And so on. The resulting ranking is  $y_1 R \cdots R y_{n-1} R y_n$ , where  $y_n$  denotes the unique element of  $\mathcal{X} \setminus \{y_1, \ldots, y_{n-1}\}$ .<sup>40</sup>

Recursive amendment is distinct from insertion sort.<sup>41</sup> Nonetheless:

**Proposition 3.** Recursive amendment is outcome-equivalent to insertion sort.

This follows from one of the lemmata in our proof of Theorem 4:

Proof. Recall from appendix F.1 (p. 38) the set  $\Sigma_{n-2}$  of strategies. Clearly recursive amendment belongs to  $\Sigma_{n-2}$ . Thus by Lemma 7 (p. 38), recursive amendment is outcome-equivalent to insertion sort.

In the sorting context, recursive amendment is called *selection sort*—see e.g. Knuth (1998, §5.2.3).

<sup>&</sup>lt;sup>39</sup>If the amendment procedure demands a vote on a pair x, y that is already ranked as (wlog) x R y, treat this as offering x, y and obtaining outcome x W y.

 $<sup>^{40}</sup>$ Like insertion sort, recursive amendment describes only a path, so formally defines an equivalence class of strategies; refer to footnote 17 on p. 13.

<sup>&</sup>lt;sup>41</sup>E.g. if  $\mathcal{X} = \{1, 2, 3, 4\}$  with  $1 \succ 2 \succ 3 \succ 4$ , 1 W 4 W 3 and 4 W 2 W 3, then insertion sort offers 3, 4, then 2, 4, then 2, 3, then 1, 4, whereas recursive amendment offers 3, 4, then 2, 4, then 2, 3. The two strategies *do* coincide when there are three alternatives.

# Supplemental appendices

# H A characterisation of our 'transitive' protocol

In this appendix, we show that among all possible rules of interaction between the chair and committee that lead to a ranking, the 'transitive' protocol studied in this paper (described in §3.1) is the only one (up to restriction) that denies the chair arbitrary power and that allows votes only on pairs. This protocol is thus the natural one, given the restriction to pairwise votes. Non-binary votes raise issues that are beyond the scope of this paper.<sup>42</sup>

A ballot is a set of two or more alternatives. An election is (B, V), where B is a ballot and V is a map  $\{1, \ldots, I\} \to B$  specifying what alternative each voter votes for. An electoral history is a finite sequence of elections with distinct ballots. For two (distinct) electoral histories h, h', we write  $h \sqsubseteq (\Box) h'$  iff h is a truncation of h'.

A protocol specifies for each (permitted) electoral history either (1) a set of ballots that the chair is permitted to offer or (2) a ranking. Formally:

## **Definition 13.** A protocol is $(\mathcal{H}, \rho)$ , where

- (1)  $\mathcal{H}$  is a non-empty set of electoral histories such that
  - if h' belongs to  $\mathcal{H}$ , then so does any  $h \sqsubseteq h'$ , and
  - if  $h = ((B_1, V_1), \dots, (B_t, V_t))$  belongs to  $\mathcal{H}$ , then so does  $h' = ((B_1, V_1), \dots, (B_t, V_t'))$  for any  $V_t' : \{1, \dots, n\} \to B_t$ .

Call  $h \in \mathcal{H}$  terminal (in  $\mathcal{H}$ ) iff there is no  $h' \supseteq h$  in  $\mathcal{H}$ .

(2)  $\rho$  is a map that assigns a ranking to each terminal  $h \in \mathcal{H}$ .

Call an electoral history binary iff each ballot has exactly two elements. A binary protocol  $(\mathcal{H}, \rho)$  is one whose  $\mathcal{H}$  consists of binary electoral histories. For any binary electoral history  $h = ((\{x_s, y_s\}, V_s))_{s=1}^t$ , where wlog  $|\{i : V_s(i) = x_s\}| > I/2$  for each  $s \in \{1, \ldots, t\}$ , let  $\mathbb{R}^h$  denote the transitive closure of  $\bigcup_{s=1}^t \{(x_s, y_s)\}$ .<sup>43</sup> The transitive protocol is the binary protocol that permits the chair to offer a ballot  $\{x, y\}$  after binary electoral history

<sup>&</sup>lt;sup>42</sup>Unlike in the binary case, there is no 'most natural' non-binary protocol. In particular, reasonable protocols can differ in what they deem the committee to have 'decided' in a vote on three or more alternatives in which none won an outright majority.

<sup>&</sup>lt;sup>43</sup>If h is the empty electoral history, then  $R^h = \emptyset$ .

h exactly if the pair x,y is unranked by  $\mathbb{R}^h,$  and assigns the ranking  $\mathbb{R}^h$  to each terminal  $h.^{44}$ 

To deny the chair arbitrary power, we focus on protocols that rank x above y whenever x won an outright majority and y was also on the ballot:

**Definition 14.** A protocol  $(\mathcal{H}, \rho)$  satisfies committee sovereignty iff for any terminal  $h = ((B_t, V_t))_{t=1}^T \in \mathcal{H}$  such that  $|\{i : V_t(i) = x\}| > I/2$  and  $y \in B_t \setminus \{x\}$  for some  $t \in \{1, \ldots, T\}$ , we have  $x \ \rho(h) \ y$ .

For binary protocols, committee sovereignty is equivalent to imposing transitivity after every vote:

**Observation 4.** A binary protocol  $(\mathcal{H}, \rho)$  satisfies committee sovereignty iff  $\rho(h) \supseteq R^h$  for every terminal  $h \in \mathcal{H}$ .

That is, any pair linked by a chain of majorities  $(x \ R^h \ y)$  must be ranked accordingly  $(x \ \rho(h) \ y)$ , and so cannot be offered for a vote.<sup>45</sup>

Proof. Let  $(\mathcal{H}, \rho)$  be binary and satisfy committee sovereignty, and take a terminal  $h = ((\{x_t, y_t\}, V_t))_{t=1}^T \in \mathcal{H}$ , where wlog  $x_t \ R^h \ y_t$  for each  $t \in \{1, \ldots, T\}$ . Then  $\rho(h) \supseteq \bigcup_{t=1}^T \{(x_t, y_t)\}$  by committee sovereignty, whence  $\rho(h) \supseteq R^h$  because  $\rho(h)$  is transitive and  $R^h$  is by definition the smallest transitive relation containing  $\bigcup_{t=1}^T \{(x_t, y_t)\}$ .

For the converse, let  $(\mathcal{H}, \rho)$  be binary with  $\rho(h) \supseteq \mathbb{R}^h$  for every terminal  $h \in \mathcal{H}$ . Take any terminal  $h = ((\{x_t, y_t\}, V_t))_{t=1}^T \in \mathcal{H}$  and suppose that  $|\{i : V_t(i) = x_t\}| > I/2$ ; we must show that  $x_t \ \rho(h) \ y_t$ . Since  $x_t \ \mathbb{R}^h \ y_t$ , this follows immediately from  $\rho(h) \supseteq \mathbb{R}^h$ .

More is needed to deny the chair excessive power: she must also be required to offer enough ballots to give the committee a fair say. To formalise this, write  $x S^h y$  for an electoral history  $h = ((B_t, V_t))_{t=1}^T$  iff

$$x, y \in B_t$$
 and  $|\{i : V_t(i) = x\}| \ge |\{i : V_t(i) = y\}|$ 

for some  $t \in \{1, \ldots, T\}$ , and say that *h* gives the committee *a* say on *x*, *y* iff  $\{z_1, z_L\} = \{x, y\}$  for some sequence  $z_1 S^h z_2 S^h \cdots S^h z_L$  of alternatives.

<sup>44</sup>Explicitly it is  $(\mathcal{H}^*, \rho^*)$ , where  $\mathcal{H}^*$  consists of all binary electoral histories h' such that

$$h \sqsubset ((\{x_1, y_1\}, V_1), \dots, (\{x_t, y_t\}, V_t)) \sqsubseteq h' \text{ implies } x_t \not \mathbb{R}^h y_t \not \mathbb{R}^h x_t,$$

(so that  $h \in \mathcal{H}$  is terminal iff  $\mathbb{R}^h$  is a ranking,) and  $\rho^{\star}(h) := \mathbb{R}^h$  for each terminal  $h \in \mathcal{H}^{\star}$ . <sup>45</sup>Formally: if  $x \mathbb{R}^h y$ , then no terminal  $h' \supseteq h$  can feature the ballot  $\{x, y\}$  (except

in h). For otherwise there would be a terminal  $h' \supseteq h'$  can back the ballot [x,y] (except in h). For otherwise there would be a terminal h' in which y beats x in a vote, so that  $x R^{h'} y R^{h'} x$ , which is impossible since  $\rho(h') \supseteq R^{h'}$  and  $\rho(h')$  is a ranking.

**Definition 15.** A protocol  $(\mathcal{H}, \rho)$  satisfies democratic legitimacy iff every terminal  $h \in \mathcal{H}$  gives the committee a say on each pair of alternatives.

Write  $\tau(\mathcal{H})$  for the terminal elements of  $\mathcal{H}$ . A protocol  $(\mathcal{H}, \rho)$  is a restriction of  $(\mathcal{H}', \rho')$  iff  $\tau(\mathcal{H}) \subseteq \tau(\mathcal{H}')$  and  $\rho = \rho'|_{\tau(\mathcal{H})}$ .<sup>46</sup> To wit, anything the chair can do under  $(\mathcal{H}, \rho)$ , she can also do under  $(\mathcal{H}', \rho')$ .

**Proposition 4.** A protocol is binary and satisfies committee sovereignty and democratic legitimacy iff it is a restriction of the transitive protocol.

Thus any binary protocol that does not give the chair arbitrary power must be the transitive protocol, possibly with limitations on what unranked pairs the chair may offer at some histories. Neglecting such limitations as ad hoc, we arrive at the transitive protocol.

Proof. Any restriction of the transitive protocol  $(\mathcal{H}^*, \rho^*)$  satisfies the three properties since  $(\mathcal{H}^*, \rho^*)$  does and the properties are preserved under restriction. For the converse, let  $(\mathcal{H}, \rho)$  satisfy the three properties; we must show that  $\tau(\mathcal{H}) \subseteq \tau(\mathcal{H}^*)$  and  $\rho = \rho^*|_{\tau(\mathcal{H})}$ .

To establish  $\tau(\mathcal{H}) \subseteq \tau(\mathcal{H}^*)$ , we show separately that  $\mathcal{H} \subseteq \mathcal{H}^*$  and that any  $h \in \tau(\mathcal{H}) \subseteq \mathcal{H}^*$  is terminal in  $\mathcal{H}^*$ . For the former, fix a pair of electoral histories  $h \sqsubset h' = ((\{x_s, y_s\}, V_s))_{s=1}^t \in \mathcal{H}$ . We must show that the pair  $x_t, y_t$ is unranked by  $\mathbb{R}^h$ , so suppose toward a contradiction that  $x_t \mathbb{R}^h y_t$ . Then we must have  $x_t \rho(h'') y_t$  for any terminal  $h'' \in \mathcal{H}$  such that  $h'' \supseteq h$  since  $\rho(h'') \supseteq \mathbb{R}^{h''} \supseteq \mathbb{R}^h$  by Observation 4. In particular, this must hold for any terminal  $h'' \in \mathcal{H}$  with first t-1 elections  $(\{x_1, y_1\}, V_1), \ldots, (\{x_{t-1}, y_{t-1}\}, V_{t-1})$  and  $t^{\text{th}}$ election  $(\{x_t, y_t\}, V_t')$ , where  $V_t'$  satisfies  $|\{i : V_t'(i) = y_t\}| > I/2$ . But for such an h'', committee sovereignty of  $(\mathcal{H}, \rho)$  clearly demands that  $y_t \rho(h'') x_t$ —a contradiction.

For the latter, let  $h \in \tau(\mathcal{H}) \subseteq \mathcal{H}^*$ ; we must show that h is terminal in  $\mathcal{H}^*$ , meaning precisely that  $\mathbb{R}^h$  is total. Since h is binary, a pair x, y is ranked by  $\mathbb{R}^h$  iff h gives the committee a say on x, y. And h gives the committee a say on every pair since  $(\mathcal{H}, \rho)$  satisfies democratic legitimacy.

To show that  $\rho = \rho^*|_{\tau(\mathcal{H})}$ , fix an  $h \in \tau(\mathcal{H})$ . Then  $\rho(h) \supseteq R^h = \rho^*(h)$  by Observation 4 and the definition of  $\rho^*$ , and the containment must be an equality since  $\rho(h)$  and  $\rho^*(h)$  are both rankings.

### I The limits of strategic voting

In this appendix, we show that sincere voting is the unique regret-free strategy of each voter. In fact, we show something stronger: deviating from sincere

 $<sup>^{46}\</sup>tau(\mathcal{H}) \subseteq \tau(\mathcal{H}')$  holds exactly if  $\mathcal{H} \subseteq \mathcal{H}'$  and any  $h \in \tau(\mathcal{H})$  is terminal in  $\mathcal{H}'$ .

voting results in a no better (a worse) outcome against any (some) strategies of the chair and the other voters, in the 'more aligned' sense.

Let each voter  $i \in \{1, \ldots, I\}$  have a strict preference  $\succ_i$  over the alternatives  $\mathcal{X}$ . A strategy  $\sigma_i$  of a voter specifies, after each history and for every offered pair x, y, whether x or y should be voted for. (Since a history records only which pairs were offered and which alternative won in each pair, not who voted how, this definition of a strategy rules out complex path-dependence. We shall relax that stricture below.)

A voter's sincere strategy is the one that always instructs her to vote for whichever alternative she likes better. For a strategy  $\sigma$  of the chair and strategies  $\sigma_1, \ldots, \sigma_I$  of the voters, write  $R(\sigma, \sigma_1, \ldots, \sigma_I)$  for the outcome (the ranking that results).

**Definition 16.** Let  $\sigma_i, \sigma'_i$  be strategies of voter *i*, and  $\sigma, \sigma_{-i}$  strategies of the chair and the other voters.  $\sigma'_i$  is obviously better than  $\sigma_i$  against  $\sigma, \sigma_{-i}$  iff  $R(\sigma, \sigma'_i, \sigma_{-i})$  is distinct from, and more aligned with  $\succ_i$  than,  $R(\sigma, \sigma_i, \sigma_{-i})$ .

When one strategy is obviously better than another, it yields a better outcome no matter what voter *i*'s exact preference over rankings, given only the weak assumption that voter *i* weakly prefers rankings more aligned with her preference  $\succ_i$  over alternatives. By contrast, comparing strategies that are not related by 'obviously better than' involves trade-offs.

**Definition 17.** A strategy  $\sigma_i$  of a voter is *dominant* iff for any alternative strategy  $\sigma'_i$ ,

- (∄) there exist no strategies  $\sigma, \sigma_{-i}$  of the chair and other voters against which  $\sigma'_i$  is obviously better than  $\sigma_i$ , and
- ( $\exists$ ) there exist strategies  $\sigma, \sigma_{-i}$  of the chair and other voters against which  $\sigma_i$  is obviously better than  $\sigma'_i$ .

Dominance is strong. (Albeit not as strong as conventional dominance, since 'obviously better' is only a partial ordering.) Observe that there can be at most one dominant strategy. In fact, there is exactly one:

**Proposition 5.** For each voter, the sincere strategy is (uniquely) dominant.

Proposition 5 remains true, with the same proof, if the definition of dominance is strengthened to allow the alternative strategy  $\sigma'_i$  to be an 'extended strategy' that can condition on who voted how in previous periods.

*Proof.* Fix a voter i, and let  $\sigma_i^*$  be her sincere strategy. We must establish properties ( $\nexists$ ) and ( $\exists$ ) in the definition of dominance. For the latter, we shall make use of the extension corollary in appendix C (p. 25).

Property  $(\nexists)$ : Fix strategies  $\sigma, \sigma_{-i}$  of the chair and other voters and a non-sincere strategy  $\sigma'_i$  of voter i, and suppose that  $R' \coloneqq R(\sigma, \sigma'_i, \sigma_{-i})$  is distinct from  $R \coloneqq R(\sigma, \sigma^*_i, \sigma_{-i})$ ; we must show that R' is not more aligned with  $\succ_i$  than R. Let T be the first period in which the proto-rankings  $R_T$ and  $R'_T$  differ, and let  $\{x, y\}$  be the pair voted on in this period, where (wlog)  $x \ R_T \ y$  and  $y \ R'_T \ x$ . The two strategy profiles generate the same length-(T-1) history h (by definition of T), and thus the same period-Tvotes  $\sigma_j(h)$  by the other voters  $j \neq i$ . So voter i is pivotal after history h, and since  $\sigma^*_i$  is sincere it must be that  $x \succ_i y$ . Thus  $R' \supseteq R'_T$  is not more aligned with  $\succ_i$  than  $R \supseteq R_T$ .

Property ( $\exists$ ): Take any non-sincere strategy  $\sigma'_i$ . Choose strategies  $\sigma', \sigma'_{-i}$  such that  $\sigma'_i$  votes non-sincerely along the terminal history induced by the strategy profile  $(\sigma', \sigma'_i, \sigma'_{-i})$ , and let T be the first period in which this occurs. Then the proto-ranking in period T-1 is the same under the strategy profiles  $(\sigma', \sigma'_i, \sigma'_{-i})$  and  $(\sigma', \sigma^*_i, \sigma'_{-i})$ ; call it  $R_{T-1}$ . Write  $\{x, y\}$  for the pair of alternatives that are voted on in period T, where (wlog)  $x \succ_i y$ .

By the extension corollary in appendix C (p. 25), there exists a ranking  $R \supseteq R_{T-1}$  with  $x \ R \ y$  and  $x, y \ R$ -adjacent. Let R' be exactly R, except with the positions of x and y reversed. Clearly R is more aligned with  $\succ_i$  than R', and the two are distinct.

It thus suffices to find strategies  $\sigma$  and  $\sigma_{-i}$  such that  $R(\sigma, \sigma_i^*, \sigma_{-i}) = R$ and  $R(\sigma, \sigma_i', \sigma_{-i}) = R'$ . For the chair, let  $\sigma \coloneqq \sigma'$ . As for  $\sigma_{-i}$ , let half of the other voters  $j \in I \setminus \{i\}$  vote according R (i.e. vote for x over y iff x R y), and the rest vote according to R'.

## J How many *W*-reachable rankings are *W*-unimprovable?

This appendix contains two results. In §J.1, we show that for a given majority will W, every W-reachable ranking is W-unimprovable iff W is transitive. In §J.2, we show that on average across majority wills W, only a small fraction of W-reachable rankings are W-unimprovable if there are many alternatives.

#### J.1 When is agenda-setting valuable?

Given the majority will W, the value of agenda-setting lies in being able to reach a W-unimprovable ranking rather than some (necessarily W-reachable) ranking that is not W-unimprovable. This motivates the following definition: **Definition 18.** Given her preference  $\succ$  (a ranking), the chair benefits from agenda-setting under a majority will W iff there exists a W-reachable ranking that is not W-unimprovable.

**Proposition 6.** For a majority will W, the following are equivalent:

- (1) W is not a ranking (i.e. is not transitive).
- (2) For some  $\succ$ , the chair benefits from agenda-setting under W.
- (3) For every  $\succ$ , the chair benefits from agenda-setting under W.

In words, agenda-setting is valuable precisely because it allows the chair to exploit Condorcet cycles: the chair benefits whenever there is a cycle in W, and otherwise does not benefit.

Proof. (3) immediately implies (2). To see that (2) implies (1), consider the contra-positive: if W is a ranking, then it is clearly the only W-reachable ranking, so the chair does not benefit from agenda-setting for any  $\succ$ .

To prove that (1) implies (3), fix any ranking  $\succ$  and any majority will W that is not a ranking; it suffices to exhibit distinct W-reachable rankings R and R' such that R is more aligned with  $\succ$  than R'. To that end let R be a W-efficient ranking (these are easily seen to exist). Similarly let R' be a W-anti-efficient ranking, i.e. one such that  $x \prec y$  and x W y implies x R y.

To show that R is more aligned with  $\succ$  than R', take  $x, y \in \mathcal{X}$  with  $x \succ y$ . If  $x \ W \ y$ , then  $x \ R \ y$  since R is W-efficient. If instead  $y \ W \ x$ , then  $y \ R' \ x$  since R' is W-anti-efficient. Thus  $x \ R' \ y$  implies  $x \ R \ y$ .

It remains only to show that R and R' are distinct. Since W is not a ranking, there must be  $x, y, z \in \mathcal{X}$  such that  $x \ W \ y \ W \ z \ W \ x$ . Suppose wlog that  $x \succ z$ . There are three cases. If  $x \succ y \succ z$ , then  $x \ R \ y \ R \ z$  and  $z \ R' \ x \ R' \ y$ , which are distinct. If  $y \succ x \succ z$ , then  $y \ R \ z \ R' \ x \ R' \ y$ , which are distinct. If  $x \succ z \succ y$ , then  $x \ R \ y \ R \ z \ R' \ x$ , which are distinct. If  $x \succ z \succ y$ , then  $x \ R \ y \ R \ z \ R' \ x$ , which are distinct.

#### J.2 Most *W*-reachable rankings are not *W*-unimprovable

The following asserts that when there are enough alternatives, only a small fraction of a typical W's W-reachable rankings are W-unimprovable.

**Proposition 7.** For each  $n \in \mathbf{N}$ , let  $\mathbb{R}^n$   $(W^n)$  denote a uniform random draw from the set of all rankings (majority wills) on  $\mathcal{X}_n := \{1, \ldots, n\}$ , with  $\mathbb{R}^n$  and  $W^n$  independent. Then

 $\Pr(\mathbb{R}^n \text{ is } W^n \text{-unimprovable} | \mathbb{R}^n \text{ is } W^n \text{-reachable}) \to 0 \text{ as } n \to \infty.$ 

Proof. Fix any  $n \ge 5$ , and define  $K_n := \lfloor (n-1)/4 \rfloor$ . Further fix a ranking R and a majority will W on  $\mathcal{X}_n$ , and label the alternatives  $\mathcal{X}_n = \{x_1, \ldots, x_n\}$  so that  $x_1 R \ldots R x_n$ . Given  $k \in \{1, \ldots, K_n\}$ , say that R admits a local W-improvement at  $(x_{4k-2}, x_{4k-1}, x_{4k})$  iff both

 $-x_{4k-3} W x_{4k} W x_{4k-2}$  and  $x_{4k-1} W x_{4k+1}$ , and

$$-x_{4k} \succ x_{4k-1}$$
 and  $x_{4k} \succ x_{4k-2}$ 

If R admits a local W-improvement at  $(x_{4k-2}, x_{4k-1}, x_{4k})$ , then it fails to be W-unimprovable since the ranking

$$x_1 R' \cdots R' x_{4k-3} R' x_{4k} R' x_{4k-2} R' x_{4k-1} R' x_{4k+1} R' \cdots R' x_n$$

is then W-reachable (by Observation 1 in appendix B.3 (p. 22)) and more aligned with  $\succ$ .

For each  $n \ge 5$ , let  $(X_k^n)_{k=1}^n$  be random variables such that

$$\{X_1^n, \dots, X_n^n\} = \mathcal{X}_n$$
 and  $X_1^n R^n \dots R^n X_n^n$  a.s.

The events  $X_{4k}^n \succ X_{4k-1}^n$  and  $X_{4k}^n \succ X_{4k-2}^n$  are independent across  $k \in \{1, \ldots, K_n\}$  and each have probability 1/4. It follows by Observation 1 that conditional on  $\mathbb{R}^n$  being  $W^n$ -reachable, the events

 $R^{n}$  admits a local  $W^{n}$ -improvement at  $(X_{4k-2}^{n}, X_{4k-1}^{n}, X_{4k}^{n})$ 

are independent across  $k \in \{1, \ldots, K_n\}$  and have probability  $(1/2)^5$ . Thus

$$\Pr(\mathbb{R}^n \text{ is } W^n \text{-unimprovable} | \mathbb{R}^n \text{ is } W^n \text{-reachable}) \le \left(1 - (1/2)^5\right)^{K_n},$$

which vanishes as  $n \to \infty$  since  $K_n = \lfloor (n-1)/4 \rfloor$  diverges.

### K Indecisive votes

In this appendix, we allow the vote on a pair of alternatives to be indecisive, in which case the chair may choose how they are ranked. (This occurs e.g. when the chair is a voting member of the committee.) To that end, we re-interpret x W y to mean that the chair is permitted to rank x above y, and allow for the possibility that both x W y and y W x. A vote on  $\{x, y\}$ with x W y is indecisive if also y W x, and decisive otherwise.

The 'majority will' W must still be total and irreflexive, but not necessarily asymmetric. By appeal to an argument similar to that for Fact 1

(appendix B.1, p. 21), any total and irreflexive relation W should be considered.

A history still records what pairs were offered and how each pair was ranked, and a *strategy* now specifies not only what pair to offer after each history, but also how to rank them if the vote is indecisive. Note that a history does not record whether a vote was decisive or not, and thus that we rule out strategies that condition on this information. We show in supplemental appendix L how this restriction may be dropped.

Regret-free and efficient strategies are defined as before, with 'for any majority will W' replaced by 'for any total and irreflexive W'. By Lemma 1 (p. 12), efficiency still implies regret-freeness.

When the chair offers  $\{x, y\}$  with  $x \succ y$  and the vote is indecisive, we say that she ranks in her interest iff she ranks x above y, and against her interest otherwise. Augment the definition of insertion sort in §5 so that the chair ranks in her interest whenever a vote is indecisive. Theorem 1 (§5, p. 13) remains true, with the same proof: insertion sort is efficient, and thus regret-free.

The characterisations of regret-free strategies (Theorems 2 and 3 in §6, pp. 15 and 17) extend as follows:

**Theorem (2+3)'.** For a strategy  $\sigma$ , the following are equivalent:

- (a)  $\sigma$  is regret-free.
- (b)  $\sigma$  is efficient.
- (c)  $\sigma$  never misses an opportunity, takes a risk, or ranks against the chair's interest.

*Proof.* We establish the implications depicted in Figure 3 (p. 17). The proof that (c) implies (b) given in appendix D.1 applies essentially unchanged. That (b) implies (a) follows from Lemma 1 (p. 12).

To show that (a) implies (c), observe that the proof in appendix D.2 establishes that regret-free strategies never miss an opportunity or take a risk. It therefore remains only to show that a regret-free strategy must not rank against the chair's interest.

We prove the contra-positive. Let  $\sigma$  be a strategy that ranks against the chair's interest under some majority will W; we shall find a majority will W' such that the outcome R of  $\sigma$  under W' fails to be W'-unimprovable. In particular, we shall exhibit a W'-reachable ranking  $R' \neq R$  that is more aligned with  $\succ$  than R.

Let T be the first period in which  $\sigma$  ranks against the chair's interest under W. Write  $R_{T-1}$  for the associated end-of-period-(T-1) proto-ranking, and let  $\{x, y\}$  be the pair offered in period T. By hypothesis, x W y W x, and the chair chooses to rank y above x.

By the extension corollary in appendix C (p. 25), there exists a ranking  $R' \supseteq R_{T-1} \cup \{(x, y)\}$  such that x, y are R'-adjacent. Define a majority will W' by  $W' := R' \cup \{(y, x)\}$ , and denote by R the outcome of  $\sigma$  under W'. Clearly R' is W'-reachable. It remains to show that  $R \neq R'$  and that R' is more aligned with  $\succ$  than R.

For the former, since  $x \ R' \ y$ , it suffices to show that  $y \ R \ x$ . To this end, observe that that  $R_{T-1} \subseteq R' \subseteq W'$ . Thus the history of length T-1generated by  $\sigma$  and W' is the same as that generated by  $\sigma$  and W, which means in particular that  $\{x, y\}$  is offered in period T, and that y is ranked above x if the vote is indecisive. Under W', the vote is indeed indecisive  $(x \ W' \ y \ W' \ x)$ , and thus  $y \ R \ x$  as desired.

To show that R' is more aligned with  $\succ$  than R, observe that W' agrees with R' on every pair  $\{z, w\} \not\subseteq \{x, y\} = [x, y]_{R'}$ . It follows by Lemma 3 in appendix D.2 (p. 28) that R and R' agree on every pair  $\{z, w\} \neq \{x, y\}$ . Since  $x \succ y$  and x R' y, it follows that R' is more aligned with  $\succ$  than R.

All of the remaining results also extend: the characterisations of regretfreeness are tight (Propositions 1 and 2 in §6, pp. 16 and 18), the outcomeequivalents of insertion sort are (include) the lexicographic (amendment) strategies (Theorem 4 and Proposition 3 in §7, pp. 20 and 44), and sincere voting is dominant (Proposition 5 in supplemental appendix I).

#### L Strategies with extended history-dependence

By definition, a strategy does not condition on who voted how in the past. To relax this restriction, let an *extended history* be a sequence of pairs offered and votes cast by each member on each pair, and let an *extended strategy* assign to each extended history an unranked pair of alternatives.

Recall from appendix B.1 (p. 21) the definition of voting profiles. The outcome of an extended strategy  $\sigma$  under a voting profile  $(V_i)_{i=1}^I$  is the final ranking that results. A strategy is regret-free (efficient) iff its outcome under every voting profile  $(V_i)_{i=1}^I$  is W-unimprovable (W-efficient), where W denotes the majority will of  $(V_i)_{i=1}^I$ .

Insertion sort is clearly an extended strategy, so is efficient by Theorem 1 (§5, p. 13). Our characterisation of regret-freeness (Theorems 2 and 3 in §6, pp. 15 and 17) remains valid:

**Theorem (2+3)**". For an extended strategy  $\sigma$ , the following are equivalent:

- (a)  $\sigma$  is regret-free.
- (b)  $\sigma$  is efficient.
- (c)  $\sigma$  never misses an opportunity or takes a risk.

*Proof.* We prove the implications depicted in Figure 3 (p. 17). That (c) implies (b) follows from the argument in appendix D.1, which applies unchanged to extended strategies. That (b) implies (a) follows from Lemma 1 (p. 12).

To show that (a) implies (c), we prove the contra-positive by augmenting the argument in appendix D.2. Take an extended strategy  $\sigma$  that misses an opportunity or takes a risk under some voting profile  $(V_i)_{i=1}^I$ , and let t be the first period in which this occurs. Let W be the majority will of  $(V_i)_{i=1}^I$ . Construct an alternative majority will W' exactly as in the proof in appendix D.2. Construct in addition a voting profile  $(V'_i)_{i=1}^I$  whose majority will is W', and such that the extended history up to time t under  $\sigma$  and  $(V'_i)_{i=1}^I$  is the same as under  $\sigma$  and  $(V_i)_{i=1}^I$ . The argument in appendix D.2 ensures that the outcome of  $\sigma$  under  $(V'_i)_{i=1}^I$  fails to be W'-unimprovable. Thus  $\sigma$  fails to be regret-free.

#### M Relation to ranking methods

In this appendix, we investigate the link with the social choice literature mentioned in §1.1 (p. 3). We recast the chair's problem as a choice among ranking methods, characterise the constraint set of this problem, and compare its solutions to ranking methods in the literature.

A ranking method is a map that assigns to each majority will a ranking. Each strategy  $\sigma$  induces a ranking method, namely the one whose value at a majority will W is the outcome of  $\sigma$  under W. Call a ranking method  $\rho$  feasible iff it is induced by some strategy, and regret-free iff  $\rho(W)$  is W-unimprovable for every W. Clearly the chair's problem in §3 can be re-formulated as a choice between ranking methods, where the constraint set consists of the feasible ranking methods and the objective is to choose a regret-free one.

For a majority will W and rankings R, R', say that R is more aligned with W than R' iff for any pair  $x, y \in \mathcal{X}$  of alternatives with x W y, if x R' ythen also x R y. This is exactly the definition in the text (§3.4, p. 8), except that we allow W to be any majority will (not necessarily a ranking). **Definition 19.** A ranking method  $\rho$  is *faithful* iff for every majority will W, no ranking  $R \neq \rho(W)$  is more aligned with W than  $\rho(W)$ .

Faithfulness clearly admits a normative interpretation. It is a natural strengthening of Condorcet consistency, the requirement that  $\rho(W)$  rank x highest if x W y for every alternative  $y \neq x$ . The following shows that it also has a positive interpretation:

**Observation 5.** A ranking method  $\rho$  is faithful iff  $\rho(W)$  is W-reachable for every majority will W.

Proof. Fix a ranking method  $\rho$  and a majority will W, and write  $R \coloneqq \rho(W)$ . If R is W-reachable, then any  $R' \neq R$  fails to be more aligned with W since it must rank some R-adjacent pair  $x \ R \ y$  as  $y \ R' \ x$ , where  $x \ W \ y$  by Observation 1 (appendix B.3, p. 22). If R is not W-reachable, then by Observation 1 there is an R-adjacent pair  $x \ R \ y$  such that  $y \ W \ x$ , so the ranking  $R' \neq R$  that agrees with R on every pair but x, y is more aligned with W.

By Observation 5, any feasible ranking method must be faithful. The converse does not hold, because feasibility also imposes restrictions across majority wills. To describe these constraints, we introduce a second property:

**Definition 20.** A ranking method  $\rho$  is consistent iff whenever  $\rho(W) \neq \rho(W')$  for two majority wills W and W', there are alternatives  $x, y \in \mathcal{X}$  such that x W y W' x and

 $x \ \rho(W'') \ y$  iff  $x \ W'' \ y$  for every majority will  $W'' \supseteq W \cap W'$ .

This property is mathematically natural, but we do not think that it has any normative appeal. Instead, it captures constraints that the rules of the interaction impose on the chair:

**Proposition 8.** A ranking method is feasible iff it is faithful and consistent.

Call a ranking method  $\rho$  efficient iff  $\rho(W)$  is W-efficient for every majority will W. (W-efficiency is defined in §5, p. 12.) By Theorem 2 (p. 15), a feasible ranking method is regret-free iff it is efficient. Thus:

**Corollary 3.** A ranking method is feasible and regret-free iff it is faithful, consistent and efficient.

While faithfulness has a normative interpretation, consistency and efficiency are 'positive' in nature: the former is a constraint imposed by the rules of the game, while the latter is defined in terms of the chair's self-interested preference  $\succ$ . This makes feasible and regret-free ranking methods quite different from those studied in the literature, which are characterised by purely normative axioms (e.g. Rubinstein (1980) for the Copeland method). Indeed, standard ranking methods such as those of Copeland and Kemeny–Slater are neither consistent nor efficient, though the latter is faithful.

Proof of Proposition 8. For necessity, let  $\rho$  be feasible. Then  $\rho$  is faithful by Observation 5. To show that it is consistent, let  $\sigma$  be a strategy inducing  $\rho$ , and fix majority wills W and W' such that  $\rho(W) \neq \rho(W')$ . Let t be the first period in which the history generated by  $\sigma$  and W differs from that generated by  $\sigma$  and W', and let  $\{x, y\}$  be the pair offered in this period. Then W and W' disagree on  $\{x, y\}$ . Furthermore, the pair  $\{x, y\}$  is clearly offered in period t of the history generated by  $\sigma$  and any  $W'' \supseteq W \cap W'$ , so that  $x \rho(W'') y$  iff x W'' y.

For sufficiency, let  $\rho$  be faithful and consistent; we shall construct a strategy that induces  $\rho$ . For each history h, let  $W_h$  and  $W'_h$  be majority wills such that

- (a) if  $h = ((x_t, y_t))_{t=1}^T$ , then  $x_t W_h y_t$  and  $x_t W'_h y_t$  for each  $t \in \{1, \dots, T\}$ , and
- (b)  $W_h$  and  $W'_h$  disagree on any pair that is not voted on in h.

Since  $\rho$  is faithful,  $\rho(W_h)$  is  $W_h$ -reachable and  $\rho(W'_h)$  is  $W'_h$ -reachable by Observation 5. Thus by Observation 1 (appendix B.3, p. 22), we have  $\rho(W_h) = \rho(W'_h)$  iff h is terminal. Since  $\rho$  is consistent, we may for each non-terminal history h choose a pair  $\sigma(h) \coloneqq \{x, y\} \subseteq \mathcal{X}$  that satisfies

- (c)  $x W_h y W'_h x$  and
- (d) x W'' y iff  $x \rho(W'') y$  for any majority will  $W'' \supseteq W_h \cap W'_h$ .

**Claim.** Let  $h = ((x_t, y_t))_{t=1}^T \neq \emptyset$  be a history such that  $\{x_1, y_1\} = \sigma(\emptyset)$  and

$$\{x_t, y_t\} = \sigma\Big(((x_s, y_s))_{s=1}^{t-1}\Big)$$
 for each  $t \in \{2, \dots, T\}$ .

Then (i) for any majority will W'' with  $x_t W'' y_t$  for each  $t \in \{1, \ldots, T\}$ , we have  $x_t \rho(W'') y_t$  for each  $t \in \{1, \ldots, T\}$ , and (ii) the pair  $\sigma(h)$  is unranked by the transitive closure of  $\bigcup_{t=1}^T \{(x_t, y_t)\}$ .

Proof of the claim. For the first part, fix a  $t \in \{1, \ldots, T\}$  and a majority will W'' such that  $x_s W'' y_s$  for each  $s \in \{1, \ldots, T\}$ . Define  $h' \coloneqq ((x_s, y_s))_{s=1}^{t-1}$  (meaning  $h' = \emptyset$  if t = 1), noting that  $\sigma(h') = \{x_t, y_t\}$ . We have  $W'' \supseteq W_{h'} \cap W'_{h'}$  since  $W_{h'}$  and  $W'_{h'}$  satisfy (b), whence  $x_t \rho(W'') y_t$  by (d).

For the second part, we have  $x_t \ \rho(W_h) \ y_t$  and  $x_t \ \rho(W'_h) \ y_t$  for every  $t \in \{1, \ldots, T\}$  by (a) and the first part of the claim, implying that  $\rho(W_h)$  and  $\rho(W'_h)$  (being transitive) agree on every pair ranked by the transitive closure of  $\bigcup_{t=1}^T \{(x_t, y_t)\}$ . Since  $\rho(W_h)$  and  $\rho(W'_h)$  disagree on the pair  $\sigma(h)$  by (c) and (d), it follows that  $\sigma(h)$  is unranked by the transitive closure.  $\Box$ 

By the second part of the claim,  $\sigma$  is a well-defined strategy.<sup>47</sup> To show that it induces  $\rho$ , fix a majority will W, and let  $h = ((x_t, y_t))_{t=1}^T$  be the terminal history generated by  $\sigma$  and W; we must demonstrate that  $\rho(W)$  is the transitive closure of  $\bigcup_{t=1}^T \{(x_t, y_t)\}$ . Since both are rankings, it suffices to show that  $x_t \rho(W) y_t$  for every  $t \in \{1, \ldots, T\}$ . And this follows from the claim since  $x_t W y_t$  for every  $t \in \{1, \ldots, T\}$ .

# References

- Apesteguia, J., Ballester, M. A., & Masatlioglu, Y. (2014). A foundation for strategic agenda voting. Games and Economic Behavior, 87, 91–99. https://doi.org/10.1016/j.geb.2014.05.006
- Arrow, K. J. (1950). A difficulty in the concept of social welfare. Journal of Political Economy, 58(4), 328–346. https://doi.org/10.1086/256963
- Arrow, K. J. (1951). Social choice and individual values. Wiley.
- Arrow, K. J. (1963). Social choice and individual values (2nd). Yale University Press.
- Banks, J. S. (1985). Sophisticated voting outcomes and agenda control. Social Choice and Welfare, 1(4), 295–306. https://doi.org/10.1007/ BF00649265
- Barberà, S., & Gerber, A. (2017). Sequential voting and agenda manipulation. *Theoretical Economics*, 12(1), 211–247. https://doi.org/10.3982/ TE2118
- Bell, D. E. (1982). Regret in decision making under uncertainty. Operations Research, 30(5), 961–981. https://doi.org/10.1287/opre.30.5.961
- Black, D. (1958). The theory of committees and elections. Cambridge University Press.

<sup>&</sup>lt;sup>47</sup>We actually defined  $\sigma$  only on the path. Off the path, any behaviour will do.

- Bossert, W., & Suzumura, K. (2010). Consistency, choice, and rationality. Harvard University Press.
- Charon, I., & Hudry, O. (2010). An updated survey on the linear ordering problem for weighted or unweighted tournaments. Annals of Operations Research, 175(1), 107–158. https://doi.org/10.1007/s10479-009-0648-7
- The marquis de Condorcet. (1785). Essai sur l'application de l'analyse à la probabilité des décisions rendus à la pluralité des voix. L'imprimerie royale.
- Copeland, A. (1951). A reasonable social welfare function [notes from University of Michigan seminar on applications of mathematics to the social sciences].
- Curello, G., & Sinander, L. (2020). The preference lattice [working paper, 9 Jul 2020].
- Daniels, H. E. (1969). Round-robin tournament scores. *Biometrika*, 56(2), 295–299. https://doi.org/10.2307/2334422
- Farquharson, R. (1969). Theory of voting. Yale University Press.
- Fishburn, P. C. (1983). Transitive measurable utility. Journal of Economic Theory, 31(2), 293–317. https://doi.org/10.1016/0022-0531(83)90079-0
- Gehrlein, W. V. (1989). The probability of intransitivity of pairwise comparisons in individual preference. Mathematical Social Sciences, 17(1), 67–75. https://doi.org/10.1016/0165-4896(89)90017-6
- Gehrlein, W. V., & Lepelley, D. (2011). Voting paradoxes and group coherence: The Condorcet efficiency of voting rules. Springer.
- Gershkov, A., Kleiner, A., Moldovanu, B., & Shi, X. (2020). The art of compromising: Voting with interdependent values and the flag of the Weimar Republic [working paper, 4 Nov 2020].
- Gershkov, A., Moldovanu, B., & Shi, X. (2017). Optimal voting rules. Review of Economic Studies, 84(2), 688–717. https://doi.org/10.1093/restud/ rdw044
- Gershkov, A., Moldovanu, B., & Shi, X. (2019). Voting on multiple issues: What to put on the ballot? *Theoretical Economics*, 14(2), 555–596. https://doi.org/10.3982/TE3193
- Gershkov, A., Moldovanu, B., & Shi, X. (2020). Monotonic norms and orthogonal issues in multidimensional voting. *Journal of Economic Theory*, 189. https://doi.org/10.1016/j.jet.2020.105103
- González-Díaz, J., Hendrick, R., & Lohmann, E. (2014). Paired comparisons analysis: An axiomatic approach to ranking methods. Social Choice

and Welfare, 42(1), 139–169. https://doi.org/10.1007/s00355-013-0726-2

- Gordon, G. J. (1999). Regret bounds for prediction problems. Proceedings of the twelfth annual Conference on Computational Learning Theory, 29–40. https://doi.org/10.1145/307400.307410
- Groseclose, T., & Milyo, J. (2010). Sincere versus sophisticated voting in Congress: Theory and evidence. *Journal of Politics*, 72(1), 60–73. https://doi.org/10.1017/s0022381609990478
- Horan, S. (2021). Agendas in legislative decision-making. Theoretical Economics, 16(1), 235–274. https://doi.org/10.3982/TE2618
- Kemeny, J. G. (1959). Mathematics without numbers. *Dædalus*, 88(4), 577–591.
- Kendall, M. G. (1955). Further contributions to the theory of paired comparisons. Biometrics, 11(1), 43–62. https://doi.org/10.2307/3001479
- Kleiner, A., & Moldovanu, B. (2017). Content-based agendas and qualified majorities in sequential voting. American Economic Review, 107(6), 1477–1506. https://doi.org/10.1257/aer.20160277
- Knuth, D. E. (1998). The art of computer programming, volume 3: Sorting and searching (2nd). Addison–Wesley.
- Ladha, K. K. (1994). Coalitions in congressional voting. Public Choice, 78(1), 43–63. https://doi.org/10.1007/BF01053365
- Loomes, G., & Sugden, R. (1982). Regret theory: An alternative theory of rational choice under uncertainty. The Economic Journal, 92(368), 805–824. https://doi.org/10.2307/2232669
- McGarvey, D. C. (1953). A theorem on the construction of voting paradoxes. Econometrica, 21(4), 608–610. https://doi.org/10.2307/1907926
- Milgrom, P., & Shannon, C. (1994). Monotone comparative statics. *Econo*metrica, 62(1), 157–180. https://doi.org/10.2307/2951479
- Miller, N. R. (1977). Graph-theoretical approaches to the theory of voting. American Journal of Political Science, 21(4), 769–803. https://doi. org/10.2307/2110736
- Moon, J. W., & Pullman, N. J. (1970). On generalized tournament matrices. SIAM Review, 12(3), 384–399. https://doi.org/10.1137/1012081
- Myerson, R. B. (1991). Game theory. Harvard University Press.
- Ordeshook, P. C., & Palfrey, T. R. (1988). Agendas, strategic voting, and signaling with incomplete information. American Journal of Political Science, 32(2), 441–466. https://doi.org/10.2307/2111131
- Oscarsson, H. (2019). Rekordlåg personröstning. In U. Andersson, B. Rönnerstrand, P. Öhberg, & A. Bergström (Eds.), Storm och stiltje (pp. 107– 116). SOM-institutet.

- Poole, K. T., & Rosenthal, H. (1997). Congress: A political-economic history of roll call voting. Oxford University Press.
- Roessler, C., Shelegia, S., & Strulovici, B. (2018). Collective commitment. Journal of Political Economy, 126(1), 347–380. https://doi.org/10. 1086/694294
- Rubinstein, A. (1980). Ranking the participants in a tournament. SIAM Journal on Applied Mathematics, 38(1), 108–111. https://doi.org/10. 1137/0138009
- Savage, L. J. (1951). The theory of statistical decision. Journal of the American Statistical Association, 46(253), 55–67. https://doi.org/10.2307/ 2280094
- Slater, P. (1961). Inconsistencies in a schedule of paired comparisons. Biometrika, 48(3–4), 303–312. https://doi.org/10.1093/biomet/48.3-4.303
- Slutzki, G., & Volij, O. (2005). Ranking participants in generalized tournaments. International Journal of Game Theory, 33(2), 255–270. https://doi.org/10.1007/s00182-005-0197-5
- Suzumura, K. (1976). Remarks on the theory of collective choice. *Economica*, 43(172), 381–390. https://doi.org/10.2307/2553273
- Wei, T.-H. (1952). Algebraic foundations of ranking theory (doctoral thesis). University of Cambridge.
- Wilkerson, J. D. (1999). "Killer" amendments in Congress. American Political Science Review, 93(3), 535–552. https://doi.org/10.2307/2585573
- Young, H. P. (1986). Optimal ranking and choice from pairwise comparisons. In B. Grofman & G. Owen (Eds.), *Information pooling and group decision making* (pp. 113–122). JAI Press.
- Young, H. P. (1988). Condorcet's theory of voting. American Political Science Review, 82(4), 1231–1244. https://doi.org/10.2307/1961757
- Young, H. P., & Levenglick, A. (1978). A consistent extension of Condorcet's election principle. SIAM Journal on Applied Mathematics, 35(2), 285–300. https://doi.org/10.1137/0135023
- Zermelo, E. (1929). Die Berechnung der Turnier-Ergebnisse als ein Maximumproblem der Wahrscheinlichkeitsrechnung. Mathematische Zeitschrift, 29(1), 436–460. https://doi.org/10.1007/BF01180541
- Zinkevich, M. (2003). Online convex programming and generalized infinitesimal gradient ascent. Proceedings of the twentieth International Conference on Machine Learning, 928–935.