

LIMIT THEOREMS OF STOCHASTIC DIFFERENTIAL EQUATIONS WITH JUMPS*

HUIJIE QIAO

*School of Mathematics, Southeast University
Nanjing, Jiangsu 211189, China*

*Department of Mathematics, University of Illinois at Urbana-Champaign
Urbana, IL 61801, USA
hjgiaogean@seu.edu.cn*

ABSTRACT. In this paper, we study the convergence for solutions to a sequence of (possibly degenerate) stochastic differential equations with jumps, when the coefficients converge in some appropriate sense. Our main tools are the superposition principles. We also give sufficient conditions for the existence and uniqueness of weak solutions to the corresponding non-local Fokker-Planck equations.

1. INTRODUCTION

Stochastic differential equations (SDEs in short) with jumps appear naturally in various applied fields. They have attracted the attention of many people. For example, in [4], Jacod systematically discussed the martingale problems associated with SDEs with jumps. In [5], Jacod and Shiryaev studied limit theorems of SDEs driven by càdlàg processes under Lipschitz conditions. Qiao and Zhang [10] proved that, under non-Lipschitz conditions, for almost all sample point ω , the solutions to a certain SDE with jumps form a homeomorphism flow. Recently, Fournier and Xu [2] established the equivalence between SDEs with jumps and the corresponding non-local Fokker-Planck equations under only linear growth conditions.

In this paper, we study the convergence for solutions to a sequence of (possibly degenerate) stochastic differential equations with jumps, when the coefficients converge in some appropriate sense. More precisely, we fix a $T > 0$ and the following sequence of SDEs with jumps:

$$dX_t^n = b^n(t, X_t^n)dt + \sigma^n(t, X_t^n)dB_t + \gamma^n \int_{\mathbb{U}} g(t, X_{t-}^n, u)N(dt, du), \quad t \in [0, T], \quad (1)$$

where (B_t) is an m -dimensional Brownian motion and $N(dt, du)$ is a Poisson random measure with the intensity $dt\nu(du)$. Here ν is a finite measure defined on $(\mathbb{U}, \mathcal{U})$, where $(\mathbb{U}, \mathcal{U})$ is any measurable space. The coefficients $b^n : [0, T] \times \mathbb{R}^d \mapsto \mathbb{R}^d$, $\sigma^n : [0, T] \times \mathbb{R}^d \mapsto \mathbb{R}^{d \times m}$ are Borel measurable functions, $\{\gamma^n\}$ is a sequence of real numbers and $g : [0, T] \times \mathbb{R}^d \times \mathbb{U} \mapsto \mathbb{R}^d$ is Borel measurable. Under some pretty weak conditions, we show that, when $b^n \rightarrow b$, $\sigma^n \rightarrow \sigma$, $\gamma^n \rightarrow \gamma$ in some sense as $n \rightarrow \infty$, where $b : [0, T] \times \mathbb{R}^d \mapsto \mathbb{R}^d$,

AMS Subject Classification(2010): 60H10, 60J75.

Keywords: Weak solutions, martingale solutions, Fokker-Planck equations, superposition principles.

*This work was partly supported by NSF of China (No. 11001051, 11371352) and China Scholarship Council under Grant No. 201906095034.

$\sigma : [0, T] \times \mathbb{R}^d \mapsto \mathbb{R}^{d \times m}$ are Borel measurable and γ is a real number, the martingale solutions of Eq.(1) converge to that of the following equation

$$dX_t = b(t, X_t)dt + \sigma(t, X_t)dB_t + \gamma \int_{\mathbb{U}} g(t, X_{t-}, u)N(dt, du), \quad t \in [0, T]. \quad (2)$$

This kind of convergence results are very useful in approximation theory and statistics ([5]). We also give some sufficient conditions for the existence and uniqueness of weak solutions to the corresponding non-local Fokker-Planck equations (FPEs in short).

We will prove our convergence theorems using the superposition principle. And we will prove the superposition principle for the following SDE:

$$dX_t = b(t, X_t)dt + \sigma(t, X_t)dB_t + \int_{\mathbb{U}} f(t, X_{t-}, u)N(dt, du), \quad t \in [0, T], \quad (3)$$

where $f : [0, T] \times \mathbb{R}^d \times \mathbb{U} \mapsto \mathbb{R}^d$ is Borel measurable.

The paper is arranged as follows. In the next section, we introduce some concepts, such as weak solutions and martingale solutions of SDEs with jumps and weak solutions of FPEs, and their relationship. We study limits of SDEs with jumps in Section 3. In Section 4, the existence and uniqueness of weak solutions to the corresponding non-local FPEs are investigated. Finally, we prove Remark 3.3 in the appendix.

The following convention will be used throughout the paper: C with or without indices will denote different positive constants whose values may change from one place to another.

2. PRELIMINARY

2.1. Notation. In this subsection, we introduce some notation used in the sequel.

We use $|\cdot|$ and $\|\cdot\|$ for the norms of vectors and matrices, respectively. We use $\langle \cdot, \cdot \rangle$ to denote the scalar product in \mathbb{R}^d . Let A^* denote the transpose of the matrix A .

$C^2(\mathbb{R}^d)$ stands for the space of continuous functions on \mathbb{R}^d which have continuous partial derivatives of order up to 2, and $C_b^2(\mathbb{R}^d)$ stands for the subspace of $C^2(\mathbb{R}^d)$, consisting functions whose derivatives up to order 2 are bounded. $C_c^2(\mathbb{R}^d)$ is the collection of all functions in $C^2(\mathbb{R}^d)$ with compact support and $C_c^\infty(\mathbb{R}^n)$ denotes the collection of all real-valued C^∞ functions of compact support.

Let $\mathcal{P}(\mathbb{R}^d)$ be the space of all probability measures on $\mathcal{B}(\mathbb{R}^d)$, equipped with the topology of weak convergence. Let $\mathcal{P}_1(\mathbb{R}^d)$ be the collection of all the probability measures μ on $\mathcal{B}(\mathbb{R}^d)$ satisfying

$$\mu(|\cdot|) := \int_{\mathbb{R}^d} |x| \mu(dx) < \infty.$$

Let $L^\infty([0, T], \mathcal{P}_1(\mathbb{R}^d))$ be the collection of all measurable families $(\mu_t)_{t \in [0, T]}$ of probability measures on $\mathcal{B}(\mathbb{R}^d)$ satisfying $\sup_{t \in [0, T]} \mu_t(|\cdot|) < \infty$.

2.2. Weak solutions and martingale solutions for SDEs with jumps. In this subsection, we introduce the concepts of weak solutions and martingale solutions for SDEs with jumps, and prove their equivalence.

First of all, we recall the definition of weak solutions to Eq.(3). [6] is a good reference for the definition below, and it does not deal with SDEs with jumps.

Definition 2.1. (*Weak solutions*) By a weak solution to Eq.(3), we mean a quadruple $\{(\Omega, \mathcal{F}, \mathbb{P}; (\mathcal{F}_t)_{t \in [0, T]}), (B, N, X)\}$, where $(\Omega, \mathcal{F}, \mathbb{P}; (\mathcal{F}_t)_{t \in [0, T]})$ is a complete filtered probability space, (B_t) is an (\mathcal{F}_t) -adapted Brownian motion, $N(dt, du)$ is an (\mathcal{F}_t) -adapted Poisson random measure, independent of B_t , with the intensity $dt\nu(du)$, and (X_t) is an (\mathcal{F}_t) -adapted process such that for all $t \in [0, T]$,

$$\mathbb{P} \left(\int_0^t \left(|b(s, X_s)| + \|\sigma\sigma^*(s, X_s)\| + \int_{\mathbb{U}} |f(s, X_{s-}, u)| \nu(du) \right) ds < \infty \right) = 1,$$

and

$$X_t = X_0 + \int_0^t b(s, X_s) ds + \int_0^t \sigma(s, X_s) dB_s + \int_0^t \int_{\mathbb{U}} f(s, X_{s-}, u) N(ds, du), \text{ a.s. } \mathbb{P}.$$

If any two weak solutions to Eq.(3) with the same initial distribution have the same law, then we say uniqueness in law holds for Eq.(3).

Let $D_T := D([0, T], \mathbb{R}^d)$ be the collection of càdlàg functions from $[0, T]$ to \mathbb{R}^d . The generic element in D_T is denoted by w . We equip D_T with the Skorokhod topology and then D_T is a Polish space. For any $t \in [0, T]$, set

$$e_t : D_T \rightarrow \mathbb{R}^d, \quad e_t(w) = w_t, \quad w \in D_T.$$

Let $\mathcal{B}_t := \sigma\{w_s : s \in [0, t]\}$, $\bar{\mathcal{B}}_t := \cap_{s>t} \mathcal{B}_s$, and $\mathcal{B} := \mathcal{B}_T$. For $\phi \in C_b^2(\mathbb{R}^d)$, set

$$\begin{aligned} (\mathcal{A}_t \phi)(x) &:= b_i(t, x) \partial_i \phi(x) + a_{ij}(t, x) \partial_{ij} \phi(x), \\ (\mathcal{B}_t \phi)(x) &:= \int_{\mathbb{U}} \left[\phi(x + f(t, x, u)) - \phi(x) \right] \nu(du), \end{aligned}$$

where $a(t, x) = \frac{1}{2} \sigma \sigma^*(t, x)$. In the following, we define martingale solutions of Eq.(3). (c.f. [6, 12])

Definition 2.2. (*Martingale solutions*) For $\mu_0 \in \mathcal{P}(\mathbb{R}^d)$. A probability measure \mathbb{P} on (D_T, \mathcal{B}) is called a martingale solution of Eq.(3) with the initial law μ_0 at time 0, if

- (i) $\mathbb{P} \circ e_0^{-1} = \mu_0$,
- (ii) For any $\phi \in C_c^2(\mathbb{R}^d)$,

$$\mathcal{M}_t^\phi := \phi(w_t) - \phi(w_0) - \int_0^t (\mathcal{A}_s \phi + \mathcal{B}_s \phi)(w_s) ds \quad (4)$$

is a $\bar{\mathcal{B}}_t$ -adapted martingale under the probability measure \mathbb{P} . The uniqueness of the martingale solutions to Eq.(3) means that, if $\mathbb{P}, \tilde{\mathbb{P}}$ are two martingale solutions to Eq.(3) with $\mathbb{P} \circ e_0^{-1} = \tilde{\mathbb{P}} \circ e_0^{-1}$, then $\mathbb{P} \circ e_t^{-1} = \tilde{\mathbb{P}} \circ e_t^{-1}$ for any $t \in [0, T]$.

Next, we assume:

(**H** _{b, σ}) There is a constant C_1 such that for all $(t, x) \in [0, T] \times \mathbb{R}^d$,

$$|b(t, x)| + \|\sigma(t, x)\| \leq C_1(1 + |x|).$$

(**H** _{f}) There is a constant C_2 such that for all $(t, x) \in [0, T] \times \mathbb{R}^d$,

$$\int_{\mathbb{U}} |f(t, x, u)|^2 \nu(du) \leq C_2(1 + |x|)^2.$$

By the Hölder inequality, it is easy to see that

(\mathbf{H}'_f)

$$\int_{\mathbb{U}} |f(t, x, u)| \nu(du) \leq C(1 + |x|).$$

The relationship between martingale solutions and weak solutions is as follows.

Proposition 2.3. *Assume that $(\mathbf{H}_{b,\sigma})$ and (\mathbf{H}_f) hold, and that $\mu_0 \in \mathcal{P}(\mathbb{R}^d)$.*

(i) *The existence of a weak solution $(X_t)_{t \in [0, T]}$ to Eq.(3) with $\mathcal{L}_{X_0} = \mu_0$ is equivalent to the existence of a martingale solution \mathbb{P} to Eq.(3) with the initial law μ_0 . Moreover, $\mathcal{L}_{X_t} = \mathbb{P} \circ e_t^{-1}$ for any $t \in [0, T]$.*

(ii) *The uniqueness in law of the weak solution $(X_t)_{t \in [0, T]}$ to Eq.(3) with $\mathcal{L}_{X_0} = \mu_0$ is equivalent to the uniqueness of the martingale solutions \mathbb{P} to Eq.(3) with the initial law μ_0 .*

Proof. We only prove (i). Assume that $\{(\hat{\Omega}, \hat{\mathcal{F}}, \hat{\mathbb{P}}; (\hat{\mathcal{F}}_t)_{t \in [0, T]}), (\hat{B}, \hat{N}, \hat{X})\}$ is a weak solution of Eq.(3) with $\mathcal{L}_{\hat{X}_0} = \mu_0$. It follows from the Itô formula, that for any $\phi \in C_c^2(\mathbb{R}^d)$,

$$\begin{aligned} & \phi(\hat{X}_t) - \phi(\hat{X}_0) - \int_0^t (\mathcal{A}_s \phi + \mathcal{B}_s \phi)(\hat{X}_s) ds \\ &= \int_0^t \partial_i \phi(\hat{X}_s) \sigma_{ij}(s, \hat{X}_s) d\hat{B}_s + \int_0^t \int_{\mathbb{U}} (\phi(\hat{X}_s + f(s, \hat{X}_s, u)) - \phi(\hat{X}_s)) \tilde{N}(dsdu), \end{aligned}$$

where $\tilde{N}(dsdu) := \hat{N}(dsdu) - \nu(du)ds$ is the compensated martingale measure of $\hat{N}(dsdu)$. Note that

$$\int_0^T |\partial_i \phi(\hat{X}_s) \sigma_{ij}(s, \hat{X}_s)|^2 ds \leq \int_0^T C I_{|\hat{X}_s| \leq M} (1 + |\hat{X}_s|)^2 ds \leq CT(1 + M)^2$$

and

$$\begin{aligned} & \int_0^T \int_{\mathbb{U}} |\phi(\hat{X}_s + f(s, \hat{X}_s, u)) - \phi(\hat{X}_s)|^2 \nu(du) ds \\ & \leq \|\phi\|_{C_c^2(\mathbb{R}^d)}^2 \int_0^T \int_{\mathbb{U}} I_{|\hat{X}_s| \leq M} |f(s, \hat{X}_s, u)|^2 \nu(du) ds \\ & \quad + \int_0^T \int_{\mathbb{U}} I_{|\hat{X}_s| > M} |\phi(\hat{X}_s + f(s, \hat{X}_s, u))|^2 \nu(du) ds \\ & \leq \|\phi\|_{C_c^2(\mathbb{R}^d)}^2 \int_0^T I_{|\hat{X}_s| \leq M} (1 + |\hat{X}_s|)^2 ds \\ & \quad + \|\phi\|_{C_c^2(\mathbb{R}^d)}^2 \nu(\mathbb{U}) T \\ & \leq \|\phi\|_{C_c^2(\mathbb{R}^d)}^2 T(1 + M)^2 + \|\phi\|_{C_c^2(\mathbb{R}^d)}^2 \nu(\mathbb{U}) T, \end{aligned}$$

where $M > 0$ is a number such that $\text{supp}(\phi) \subset B_M := \{y \in \mathbb{R}^d; |y| \leq M\}$. Thus,

$$\int_0^t \partial_i \phi(\hat{X}_s) \sigma_{ij}(s, \hat{X}_s) d\hat{B}_s + \int_0^t \int_{\mathbb{U}} (\phi(\hat{X}_s + f(s, \hat{X}_s, u)) - \phi(\hat{X}_s)) \tilde{N}(dsdu)$$

is an $(\hat{\mathcal{F}}_t)_{t \in [0, T]}$ -adapted martingale and then

$$\phi(\hat{X}_t) - \phi(\hat{X}_0) - \int_0^t (\mathcal{A}_s \phi + \mathcal{B}_s \phi)(\hat{X}_s) ds$$

is an $(\hat{\mathcal{F}}_t)_{t \in [0, T]}$ -adapted martingale. Set $\mathbb{P} := \hat{\mathbb{P}} \circ \hat{X}^{-1}$, the argument above shows that \mathbb{P} is a martingale solution of Eq.(3).

Conversely, assume that \mathbb{P} is a martingale solution of Eq.(3). For any $n \geq 1$, we take $\phi_n \in C_c^2(\mathbb{R}^d)$ so that $\phi_n(x) = x^j$, j -th component of x , for all $|x| \leq n$, and define $\tau_n := \inf\{t \geq 0, |w_t| \geq n\}$. It follows from (4) that

$$\begin{aligned} \mathcal{M}_t^{\phi_n} &= \phi_n(w_{t \wedge \tau_n}) - \phi_n(w_0) - \int_0^{t \wedge \tau_n} (\mathcal{A}_s \phi_n + \mathcal{B}_s \phi_n)(w_s) ds \\ &= w_{t \wedge \tau_n}^j - w_0^j - \int_0^{t \wedge \tau_n} \left(b_j(s, w_s) + \int_{\mathbb{R}^d} y_j \nu_{s, w_s}(dy) \right) ds \end{aligned}$$

is a martingale under the probability measure \mathbb{P} , where $\nu_{s, w_s}(dy) := \nu(df^{-1}(s, w_s, \cdot)(y))$. Thus $w_{t \wedge \tau_n}$ is a semimartingale and w_t has local characteristics $(\tilde{b}, \tilde{a}, \tilde{\nu}_{\cdot, w_{\cdot}})$ (c.f. [4, 5]). By the definition of the characteristics, we have for any $\xi \in \mathbb{R}^d$,

$$e^{i\langle \xi, w_t \rangle} - e^{i\langle \xi, w_0 \rangle} - \int_0^t e^{i\langle \xi, w_s \rangle} \left\{ i\tilde{b}_j \xi_j - \tilde{a}_{ij} \xi_i \xi_j + \int_{\mathbb{R}^d} (e^{i\langle \xi, y \rangle} - 1 - i\langle \xi, y \rangle) \tilde{\nu}_{s, w_s}(dy) \right\} ds \quad (5)$$

is a local martingale. On the other side, by applying $\phi(x) = e^{i\langle \xi, x \rangle}$ to (4), we have that

$$\begin{aligned} e^{i\langle \xi, w_t \rangle} - e^{i\langle \xi, w_0 \rangle} - \int_0^t e^{i\langle \xi, w_s \rangle} \left\{ i b_j \xi_j + i \int_{\mathbb{R}^d} y_j \xi_j \nu_{s, w_s}(dy) - a_{lj} \xi_l \xi_j \right. \\ \left. + \int_{\mathbb{R}^d} (e^{i\langle \xi, y \rangle} - 1 - i\langle \xi, y \rangle) \nu_{s, w_s}(dy) \right\} ds \end{aligned} \quad (6)$$

is a local martingale. Comparing (5) with (6), one gets that $\tilde{b}(s, w_s) = b(s, w_s) + \int_{\mathbb{R}^d} y \nu_{s, w_s}(dy)$, $\tilde{a}(s, w_s) = a(s, w_s)$ and $\tilde{\nu}_{s, w_s} = \nu_{s, w_s}$. Put

$$\kappa_t := w_t - w_{t-}, \quad N_\kappa((0, t], A) := \sum_{0 < s \leq t} I_A(\kappa_s), \quad A \in \mathcal{B}(\mathbb{R}^d \setminus \{0\}),$$

and then $\tilde{N}_\kappa((0, t], A) := N_\kappa((0, t], A) - \int_0^t \nu_{s, w_s}(A) ds$ is the compensated martingale measure of N_κ . Define

$$M_t := w_t - w_0 - \int_0^t \left(b(s, w_s) + \int_{\mathbb{R}^d} y \nu_{s, w_s}(dy) \right) ds - \int_0^t \int_{\mathbb{R}^d} y \tilde{N}_\kappa(ds dy),$$

and then by the definition of the characteristics, it holds that M_t is a continuous local martingale with $\langle M_l, M_j \rangle_t = \int_0^t 2a_{lj}(s, w_s) ds$.

Next, we take another filtered probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}}; (\tilde{\mathcal{F}}_t)_{t \in [0, T]})$ and a d -dimensional Brownian motion \tilde{B} on it. Define

$$(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}}; (\tilde{\mathcal{F}}_t)_{t \in [0, T]}) := (\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}}; (\tilde{\mathcal{F}}_t)_{t \in [0, T]}) \times (D_T, \mathcal{B}, \mathbb{P}; (\tilde{B}_t)_{t \in [0, T]}),$$

and

$$\pi : \tilde{\Omega} \mapsto D_T, \quad \pi(\tilde{\omega}) = w, \quad \tilde{\omega} = (\tilde{\omega}, w) \in \tilde{\Omega}.$$

So, on $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}}; (\tilde{\mathcal{F}}_t)_{t \in [0, T]})$, $\tilde{M}_t(\tilde{\omega}) = M_t(w)$ is still a local martingale and $\tilde{B}_t(\tilde{\omega}) = \tilde{B}_t(\tilde{\omega})$ is still a Brownian motion. By the similar method to that in [6, 4.6 Proposition, Page 315], we can construct a d -dimensional Brownian motion B on $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}}; (\tilde{\mathcal{F}}_t)_{t \in [0, T]})$ so that

$$\tilde{M}_t = \int_0^t \sigma(s, \pi(\tilde{\omega})_s) dB_s,$$

i.e.

$$\pi(\check{\omega})_t = \pi(\check{\omega})_0 + \int_0^t b(s, \pi(\check{\omega})_s) ds + \int_0^t \sigma(s, \pi(\check{\omega})_s) dB_s + \int_0^t \int_{\mathbb{U}} f(s, \pi(\check{\omega})_s, u) N(dsdu),$$

where $N(dsdu)$ is a Poisson random measure on $(\check{\Omega}, \check{\mathcal{F}}, \check{\mathbb{P}}; (\check{\mathcal{F}}_t)_{t \in [0, T]})$ with the intensity measure $\nu(du)ds$. Therefore, $\{(\check{\Omega}, \check{\mathcal{F}}, \check{\mathbb{P}}; (\check{\mathcal{F}}_t)_{t \in [0, T]}), (B, N, \pi)\}$ is a weak solution of Eq.(3). Thus, the proof is complete. \square

2.3. Weak solutions of Fokker-Planck equations. In this subsection, we introduce weak solutions of the FPEs and prove a property about them.

Consider the FPE associated with Eq.(3):

$$\partial_t \mu_t = (\mathcal{A}_t + \mathcal{B}_t)^* \mu_t, \quad (7)$$

where $(\mathcal{A}_t + \mathcal{B}_t)^*$ is the adjoint operator of $\mathcal{A}_t + \mathcal{B}_t$, and $(\mu_t)_{t \in [0, T]}$ is a family of probability measures on \mathbb{R}^d . Weak solutions of Eq.(7) are defined as follows.

Definition 2.4. *A measurable family $(\mu_t)_{t \in [0, T]}$ of probability measures is called a weak solution of the non-local FPE (7) if for any $R > 0$ and $t \in [0, T]$,*

$$\int_0^t \int_{\mathbb{R}^d} I_{B_R}(x) \left(|b(s, x)| + \|a(s, x)\| + \int_{\mathbb{U}} |f(s, x, u)| \nu(du) \right) \mu_s(dx) ds < \infty, \quad (8)$$

and for all $\phi \in C_c^2(\mathbb{R}^d)$ and $t \in [0, T]$,

$$\mu_t(\phi) = \mu_0(\phi) + \int_0^t \mu_s(\mathcal{A}_s \phi + \mathcal{B}_s \phi) ds, \quad (9)$$

where $\mu_t(\phi) := \int_{\mathbb{R}^d} \phi(x) \mu_t(dx)$. The uniqueness of the weak solutions to Eq.(7) means that, if $(\mu_t)_{t \in [0, T]}$ and $(\tilde{\mu}_t)_{t \in [0, T]}$ are two weak solutions to Eq.(7) with $\mu_0 = \tilde{\mu}_0$, then $\mu_t = \tilde{\mu}_t$ for any $t \in [0, T]$.

We claim that the definition above makes sense. That is, under (8), it holds that $\int_0^t |\mu_s(\mathcal{A}_s \phi + \mathcal{B}_s \phi)| ds < \infty$ for all $\phi \in C_c^2(\mathbb{R}^d)$. Indeed, for any $\phi \in C_c^2(\mathbb{R}^d)$, we assume that the support of ϕ is in some ball B_R . Then

$$\begin{aligned} \int_0^t |\mu_s(\mathcal{A}_s \phi + \mathcal{B}_s \phi)| ds &\leq \int_0^t \int_{\mathbb{R}^d} \left[|b_i(s, x) \partial_i \phi(x)| + \frac{1}{2} |(\sigma \sigma^*)_{ij}(s, x) \partial_{ij} \phi(x)| \right. \\ &\quad \left. + \int_{\mathbb{U}} |\phi(x + f(s, x, u)) - \phi(x)| \nu(du) \right] \mu_s(dx) ds \\ &\leq \|\phi\|_{C_c^2(\mathbb{R}^d)} \int_0^t \int_{\mathbb{R}^d} I_{B_R}(x) \left(|b(s, x)| + \|a(s, x)\| \right. \\ &\quad \left. + \int_{\mathbb{U}} |f(s, x, u)| \nu(du) \right) \mu_s(dx) ds \\ &\quad + \|\phi\|_{C_c^2(\mathbb{R}^d)} \int_0^t \int_{\mathbb{R}^d} I_{B_R^c}(x) \nu(\mathbb{U}) \mu_s(dx) ds \\ &< \infty. \end{aligned}$$

If a weak solution $(\mu_t)_{t \in [0, T]}$ of the non-local FPE (7) is absolutely continuous with respect to the Lebesgue measure, then there exists a non-negative measurable function

v with $\int_{\mathbb{R}^d} v(t, x) dx = 1$ such that $\mu_t(dx) = v(t, x) dx$. Thus, v satisfies the following equation in the distributional sense

$$\partial_t v = -\partial_i (b_i v) + \partial_{ij} (a_{ij} v) + \mathcal{B}_t^* v. \quad (10)$$

Set

$$\mathcal{L}_+ := \left\{ v \geq 0 : \int_{\mathbb{R}^d} v(t, x) dx = 1, \text{ and } \sup_{t \in [0, T]} \int_{\mathbb{R}^d} |x| v(t, x) dx < \infty \right\}.$$

If there exists a $v \in \mathcal{L}_+$ satisfying Eq.(10) in the distributional sense, we say Eq.(10) has a weak solution in \mathcal{L}_+ .

Proposition 2.5. *Assume that $(\mathbf{H}_{b, \sigma})$ and (\mathbf{H}'_f) hold, and that $\mu_0(dx) = v_0(x) dx \in \mathcal{P}_1(\mathbb{R}^d)$. If there exists a non-negative measurable function v with $\int_{\mathbb{R}^d} v(t, x) dx = 1$ and $v(0, x) = v_0(x)$ such that v is a weak solution of the non-local FPE (10), then $v \in \mathcal{L}_+$.*

Proof. By the assumption and the definition of \mathcal{L}_+ , we only need to prove

$$\sup_{t \in [0, T]} \int_{\mathbb{R}^d} |x| v(t, x) dx < \infty.$$

By Definition 2.4, it holds that for all $\phi \in C_c^2(\mathbb{R}^d)$ and $t \in [0, T]$

$$\begin{aligned} \int_{\mathbb{R}^d} v(t, x) \phi(x) dx &= \int_{\mathbb{R}^d} v_0(x) \phi(x) dx + \int_0^t \int_{\mathbb{R}^d} (\mathcal{A}_s \phi(x) + \mathcal{B}_s \phi(x)) v(s, x) dx ds \\ &= \int_{\mathbb{R}^d} v_0(x) \phi(x) dx + \int_0^t \int_{\mathbb{R}^d} \left[b_i(s, x) \partial_i \phi(x) + a_{ij}(s, x) \partial_{ij} \phi(x) \right. \\ &\quad \left. + \int_{\mathbb{U}} [\phi(x + f(s, x, u)) - \phi(x)] \nu(du) \right] v(s, x) dx ds. \end{aligned} \quad (11)$$

Take

$$\phi(x) = \lambda_n(x) - 2n, \quad \lambda_n(x) := n \lambda \left(\frac{\varrho(x)}{n} \right), \quad n \in \mathbb{N},$$

where $\varrho(x) = (1 + |x|^2)^{1/2}$ and $\lambda : \mathbb{R}_+ \mapsto \mathbb{R}_+$ is a twice continuously differentiable, increasing concave function with

$$\lambda(r) = \begin{cases} r, & 0 \leq r \leq 1, \\ 2, & r \geq 2. \end{cases}$$

It is easy to see that $\phi \in C_c^2(\mathbb{R}^d)$. Inserting $\phi(x)$ in (11), one can obtain that

$$\begin{aligned} \int_{\mathbb{R}^d} v(t, x) \lambda_n(x) dx &= \int_{\mathbb{R}^d} v_0(x) \lambda_n(x) dx + \int_0^t \int_{\mathbb{R}^d} \left[b_i(s, x) \partial_i \lambda_n(x) + a_{ij}(s, x) \partial_{ij} \lambda_n(x) \right. \\ &\quad \left. + \int_{\mathbb{U}} [\lambda_n(x + f(s, x, u)) - \lambda_n(x)] \nu(du) \right] v(s, x) dx ds \\ &\leq \int_{\mathbb{R}^d} v_0(x) \lambda_n(x) dx + \int_0^t \int_{\mathbb{R}^d} \left[|b_i(s, x)| |\partial_i \lambda_n(x)| + |a_{ij}(s, x)| |\partial_{ij} \lambda_n(x)| \right. \\ &\quad \left. + \int_{\mathbb{U}} |\lambda_n(x + f(s, x, u)) - \lambda_n(x)| \nu(du) \right] v(s, x) dx ds \\ &\leq C \int_{\mathbb{R}^d} v_0(x) (1 + |x|) dx + \int_0^t \int_{\mathbb{R}^d} \left[C \varrho(x) |D \lambda_n(x)| + C \varrho(x)^2 \|D^2 \lambda_n(x)\| \right] v(s, x) dx ds \end{aligned}$$

$$+ \int_{\mathbb{U}} |\lambda_n(x + f(s, x, u)) - \lambda_n(x)| \nu(du) \Big] v(s, x) dx ds, \quad (12)$$

where $\lambda_n(x) \leq C(1 + |x|)$ is used in the last inequality and $D\lambda_n(x), D^2\lambda_n(x)$ are denoted as the gradient and Hessian matrix of $\lambda_n(x)$, respectively.

Next, by some elementary computations, it holds that

$$|D\lambda_n(x)| \leq C, \quad |D^2\lambda_n(x)| \leq C \frac{\lambda_n(x)}{\varrho(x)}, \quad \|D^2\lambda_n(x)\| \leq C \frac{\lambda_n(x)}{\varrho(x)^2}. \quad (13)$$

We now consider $\int_{\mathbb{U}} |\lambda_n(x + f(s, x, u)) - \lambda_n(x)| \nu(du)$. When $\varrho(x) > 2n$, $\lambda_n(x) = 2n$, $\lambda_n(x + f(s, x, u)) \leq 2n$, and

$$\int_{\mathbb{U}} |\lambda_n(x + f(s, x, u)) - \lambda_n(x)| \nu(du) \leq 4n\nu(\mathbb{U}) = 2\lambda_n(x)\nu(\mathbb{U}). \quad (14)$$

When $\varrho(x) \leq 2n$, $n\frac{\varrho(x)}{n} \leq \lambda_n(x)$, $\frac{\lambda_n(x)}{\varrho(x)} \geq 1$ and

$$\int_{\mathbb{U}} |\lambda_n(x + f(s, x, u)) - \lambda_n(x)| \nu(du) \leq C \int_{\mathbb{U}} |f(s, x, u)| \nu(du) \leq C\varrho(x) \leq C\lambda_n(x). \quad (15)$$

So, (14) and (15) yield that

$$\int_{\mathbb{U}} |\lambda_n(x + f(s, x, u)) - \lambda_n(x)| \nu(du) \leq C\lambda_n(x). \quad (16)$$

Finally, combining (13) and (16) with (12), we have that

$$\int_{\mathbb{R}^d} v(t, x) \lambda_n(x) dx \leq C \int_{\mathbb{R}^d} v_0(x) (1 + |x|) dx + C \int_0^t \int_{\mathbb{R}^d} \lambda_n(x) v(s, x) dx ds.$$

Thus, the Gronwall inequality implies that

$$\int_{\mathbb{R}^d} v(t, x) \lambda_n(x) dx \leq C,$$

where C is independent of n, t . Note that $\lim_{n \rightarrow \infty} \lambda_n(x) = \varrho(x)$. Therefore, by the Fatou lemma, we have

$$\sup_{t \in [0, T]} \int_{\mathbb{R}^d} v(t, x) |x| dx \leq \sup_{t \in [0, T]} \int_{\mathbb{R}^d} v(t, x) \varrho(x) dx \leq C.$$

The proof is complete. \square

2.4. The superposition principle for SDEs with jumps and non-local FPEs. In this subsection, we state two superposition principles for SDEs with jumps and non-local FPEs.

It is well known that under suitable assumptions, Eq.(3) has a weak solution $\{(\hat{\Omega}, \hat{\mathcal{F}}, \hat{\mathbb{P}}; (\hat{\mathcal{F}}_t)_{t \in [0, T]}, (\hat{B}, \hat{N}, \hat{X})\}$. By the Itô formula, it holds that $\hat{\mathbb{P}} \circ \hat{X}_t^{-1} =: \mathcal{L}_{\hat{X}_t}$ is a weak solution of Eq.(7). The natural question is whether the converse result is right. The answer is affirmative. The following proposition describes in detail the relationship between weak solutions to Eq.(3) and weak solutions to Eq.(7).

Proposition 2.6. *Suppose that $(\mathbf{H}_{b,\sigma})$ and (\mathbf{H}'_f) hold, and that $\mu_0 \in \mathcal{P}_1(\mathbb{R}^d)$.*

(i) The existence of a weak solution $(X_t)_{t \in [0, T]}$ to Eq.(3) with $\mathcal{L}_{X_0} = \mu_0$ is equivalent to the existence of a weak solution $(\mu_t)_{t \in [0, T]}$ to Eq.(7) starting from μ_0 . Moreover, $\mathcal{L}_{X_t} = \mu_t$ for any $t \in [0, T]$.

(ii) The uniqueness in law of the weak solutions $(X_t)_{t \in [0, T]}$ to Eq.(3) with $\mathcal{L}_{X_0} = \mu_0$ is equivalent to the uniqueness of the weak solutions $(\mu_t)_{t \in [0, T]}$ to Eq.(7) starting from μ_0 .

Proof. Thanks to [2, Corollary 1.5], we only show that if the weak solution $(\mu_t)_{t \in [0, T]}$ to Eq.(7) starting from μ_0 is unique, the weak solution $(X_t)_{t \in [0, T]}$ to Eq.(3) with $\mathcal{L}_{X_0} = \mu_0$ is unique in law. Assume that $\{(\Omega, \mathcal{F}, \mathbb{P}; (\mathcal{F}_t)_{t \in [0, T]}), (B, N, X)\}$ and $\{(\hat{\Omega}, \hat{\mathcal{F}}, \hat{\mathbb{P}}; (\hat{\mathcal{F}}_t)_{t \in [0, T]}), (\hat{B}, \hat{N}, \hat{X})\}$ are weak solutions of Eq.(3) with $\mathbb{P} \circ X_0^{-1} = \hat{\mathbb{P}} \circ \hat{X}_0^{-1} = \mu_0$. By (i), we know that $\mathbb{P} \circ X_t^{-1}$ and $\hat{\mathbb{P}} \circ \hat{X}_t^{-1}$ are weak solutions to Eq.(7) starting from μ_0 . Since the weak solutions to Eq.(7) starting from μ_0 are unique, $\mathbb{P} \circ X_t^{-1} = \hat{\mathbb{P}} \circ \hat{X}_t^{-1}$ for $t \in [0, T]$. That is, the weak solution $(X_t)_{t \in [0, T]}$ to Eq.(3) with $\mathcal{L}_{X_0} = \mu_0$ is unique in law. \square

Combining Proposition 2.6 with Proposition 2.3, we have the following result.

Theorem 2.7. *Suppose that $(\mathbf{H}_{b,\sigma})$ and (\mathbf{H}_f) hold, and that $\mu_0 \in \mathcal{P}_1(\mathbb{R}^d)$.*

(i) The existence of a martingale solution \mathbb{P} to Eq.(3) with the initial law μ_0 is equivalent to the existence of a weak solution $(\mu_t)_{t \in [0, T]}$ to Eq.(7) starting from μ_0 . Moreover, $\mathbb{P} \circ e_t^{-1} = \mu_t$ for any $t \in [0, T]$.

(ii) The uniqueness of the martingale solutions \mathbb{P} to Eq.(3) with the initial law μ_0 is equivalent to the uniqueness of the weak solutions $(\mu_t)_{t \in [0, T]}$ to Eq.(7) starting from μ_0 .

Remark 2.8. *Proposition 2.6 and Theorem 2.7 are usually called as superposition principles.*

3. THE LIMITS OF SDES WITH JUMPS

In this section, we take $f(t, x, u) = \gamma g(t, x, u)$ for $\gamma \in \mathbb{R}$. And then Eq.(3) becomes

$$dX_t = b(t, X_t)dt + \sigma(t, X_t)dB_t + \gamma \int_{\mathbb{U}} g(t, X_t, u)N(dt, du), \quad t \in [0, T]. \quad (17)$$

We consider the following sequence of SDEs with jumps: for any $n \in \mathbb{N}$,

$$dX_t^n = b^n(t, X_t^n)dt + \sigma^n(t, X_t^n)dB_t + \gamma^n \int_{\mathbb{U}} g(t, X_t, u)N(dt, du), \quad t \in [0, T], \quad (18)$$

where $b^n : [0, T] \times \mathbb{R}^d \mapsto \mathbb{R}^d$, $\sigma^n : [0, T] \times \mathbb{R}^d \mapsto \mathbb{R}^{d \times m}$ are Borel measurable functions and $\{\gamma^n\}$ is a real sequence. We study the relationship between martingale solutions of Eq.(17) and that of Eq.(18) when $b^n \rightarrow b, a^n \rightarrow a, \gamma^n \rightarrow \gamma$ in some sense, where $a^n := \frac{1}{2}\sigma^n \sigma^{n*}$.

The main result in the section is the following theorem.

Theorem 3.1. *Suppose that b^n, b, σ^n, σ satisfy $(\mathbf{H}_{b,\sigma})$ uniformly, $\{\gamma^n\}$ is uniformly bounded, g satisfies (\mathbf{H}_f) , and that Eq.(10) has a unique weak solution in \mathcal{L}_+ . Let $\mu_0(dx) = v_0(x)dx \in \mathcal{P}_1(\mathbb{R}^d)$, and \mathbb{P}^n, \mathbb{P} be the martingale solutions of Eq.(18) and Eq.(17) with the initial law μ_0 , respectively. Assume that*

(i) $b^n \rightarrow b, a^n \rightarrow a$ in $L^1_{loc}([0, T] \times \mathbb{R}^d)$, $\gamma^n \rightarrow \gamma$ as $n \rightarrow \infty$;

(ii) $\mathbb{P}^n \circ e_t^{-1}$ is absolutely continuous with respect to the Lebesgue measure on \mathbb{R}^d and $v^n(t, x)$ denotes the density, i.e., $v^n(t, x) := \frac{(\mathbb{P}^n \circ e_t^{-1})(dx)}{dx}$ for any $t \in [0, T]$, and

$$\sup_{x \in \mathbb{R}^d} |v^n(t, x)| \leq C,$$

where the constant $C > 0$ only depends on T .

Then $\mathbb{P}^n \rightarrow \mathbb{P}$ in $\mathcal{P}(D_T)$.

Proof. Step 1. We prove that $\{\mathbb{P}^n\}_{n \in \mathbb{N}}$ is tight in $\mathcal{P}(D_T)$.

By Theorem 4.5 in [5, Page 356], it is sufficient to check that

$$(iii) \lim_{K \rightarrow \infty} \sup_n \mathbb{P}^n \left(\sup_{t \in [0, T]} |w_t| > K \right) = 0,$$

(iv) For any stopping time τ , it holds that

$$\lim_{\theta \rightarrow 0} \sup_n \sup_{0 \leq \tau < \tau + \theta \leq T} \mathbb{P}^n (|w_{\tau + \theta} - w_\tau| \geq N) = 0, \quad \forall N > 0.$$

Since for any $n \in \mathbb{N}$, \mathbb{P}^n is a martingale solution of Eq.(18) with the initial law μ_0 , Proposition 2.3 yields that there exists a weak solution $\{(\hat{\Omega}^n, \hat{\mathcal{F}}^n, \hat{\mathbb{P}}^n; (\hat{\mathcal{F}}_t^n)_{t \in [0, T]}), (\hat{B}^n, \hat{N}^n, \hat{X}^n)\}$ of Eq.(18) with $\mathcal{L}_{\hat{X}_t^n} = \mathbb{P}^n \circ e_t^{-1}$ and $\mathcal{L}_{\hat{X}_0^n} = \mu_0$. So, by Definition 2.1, it holds that for any $t \in [0, T]$,

$$\hat{X}_t^n = \hat{X}_0^n + \int_0^t b^n(s, \hat{X}_s^n) ds + \int_0^t \sigma^n(s, \hat{X}_s^n) d\hat{B}_s^n + \int_0^t \int_{\mathbb{U}} \gamma^n g(s, \hat{X}_s^n, u) \hat{N}^n(ds, du). \quad (19)$$

And then the BDG inequality furthermore gives

$$\begin{aligned} \mathbb{E}^{\hat{\mathbb{P}}^n} \left(\sup_{s \in [0, t]} |\hat{X}_s^n| \right) &\leq \mathbb{E}^{\hat{\mathbb{P}}^n} |\hat{X}_0^n| + \mathbb{E}^{\hat{\mathbb{P}}^n} \left(\int_0^t |b^n(r, \hat{X}_r^n)| dr \right) + \mathbb{E}^{\hat{\mathbb{P}}^n} \left(\sup_{s \in [0, t]} \left| \int_0^s \sigma^n(r, \hat{X}_r^n) d\hat{B}_r^n \right| \right) \\ &\quad + \mathbb{E}^{\hat{\mathbb{P}}^n} \left(\int_0^t \int_{\mathbb{U}} |\gamma^n| |g(s, \hat{X}_s^n, u)| \hat{N}^n(dr, du) \right) \\ &\leq \mu_0(|\cdot|) + \mathbb{E}^{\hat{\mathbb{P}}^n} \left(\int_0^t |b^n(r, \hat{X}_r^n)| dr \right) + C \mathbb{E}^{\hat{\mathbb{P}}^n} \left(\int_0^t \|\sigma^n(r, \hat{X}_r^n)\|^2 dr \right)^{1/2} \\ &\quad + \mathbb{E}^{\hat{\mathbb{P}}^n} \left(\int_0^t \int_{\mathbb{U}} |\gamma^n| |g(s, \hat{X}_s^n, u)| \nu(du) dr \right) \\ &\leq \mu_0(|\cdot|) + \mathbb{E}^{\hat{\mathbb{P}}^n} \left(\int_0^t C(1 + |\hat{X}_r^n|) dr \right) + C \mathbb{E}^{\hat{\mathbb{P}}^n} \left(\int_0^t C(1 + |\hat{X}_r^n|)^2 dr \right)^{1/2}, \end{aligned}$$

and

$$\mathbb{E}^{\hat{\mathbb{P}}^n} \left(\sup_{s \in [0, t]} (1 + |\hat{X}_s^n|) \right) \leq \mu_0(1 + |\cdot|) + C(t + t^{1/2}) \mathbb{E}^{\hat{\mathbb{P}}^n} \left(\sup_{s \in [0, t]} (1 + |\hat{X}_s^n|) \right).$$

By taking t_0 with $C(t_0 + t_0^{1/2}) < 1/2$, we obtain that

$$\mathbb{E}^{\hat{\mathbb{P}}^n} \left(\sup_{s \in [0, t_0]} |\hat{X}_s^n| \right) \leq 2\mu_0(|\cdot|) + 1.$$

On $[t_0, 2t_0], [2t_0, 3t_0], \dots, [\lfloor \frac{T}{t_0} \rfloor t_0, T]$, by the same way to the above we deduce and conclude that

$$\mathbb{E}^{\hat{\mathbb{P}}^n} \left(\sup_{t \in [0, T]} |\hat{X}_t^n| \right) \leq 2^{\lfloor \frac{T}{t_0} \rfloor + 1} \mu_0(|\cdot|) + 2^{\lfloor \frac{T}{t_0} \rfloor + 1} - 1, \quad (20)$$

where $\lfloor \frac{T}{t_0} \rfloor$ stands for the largest integer no more than $\frac{T}{t_0}$. Thus, it follows from (20) that $\{\mathbb{P}^n\}_{n \in \mathbb{N}}$ satisfies (iii).

Next, we still consider Eq.(19). For any stopping time τ and $\theta > 0$ with $0 \leq \tau < \tau + \theta \leq T$, it holds that

$$\hat{X}_{\tau+\theta}^n - \hat{X}_\tau^n = \int_\tau^{\tau+\theta} b^n(s, \hat{X}_s^n) ds + \int_\tau^{\tau+\theta} \sigma^n(s, \hat{X}_s^n) d\hat{B}_s^n + \int_\tau^{\tau+\theta} \int_{\mathbb{U}} \gamma^n g(s, \hat{X}_s^n, u) \hat{N}^n(ds, du).$$

By the similar deduction to the above, we have that

$$\begin{aligned} \mathbb{E}^{\hat{\mathbb{P}}^n} |\hat{X}_{\tau+\theta}^n - \hat{X}_\tau^n| &\leq \mathbb{E}^{\hat{\mathbb{P}}^n} \int_\tau^{\tau+\theta} C(1 + |\hat{X}_s^n|) ds + C \mathbb{E}^{\hat{\mathbb{P}}^n} \left(\int_\tau^{\tau+\theta} C(1 + |\hat{X}_s^n|)^2 ds \right)^{1/2} \\ &\leq C \mathbb{E}^{\hat{\mathbb{P}}^n} \left(\sup_{s \in [0, T]} (1 + |\hat{X}_s^n|) \right) (\theta + \theta^{1/2}) \\ &\leq C \left(2^{\lfloor \frac{T}{t_0} \rfloor + 1} \mu_0(|\cdot|) + 2^{\lfloor \frac{T}{t_0} \rfloor + 1} \right) (\theta + \theta^{1/2}), \end{aligned}$$

where the last inequality is based on (20). Thus, by some elementary computations, it holds that $\{\mathbb{P}^n\}_{n \in \mathbb{N}}$ satisfies (iv). And then, $\{\mathbb{P}^n\}_{n \in \mathbb{N}}$ is relatively weakly compact. That is, there exists a weak convergence subsequence still denoted as $\{\mathbb{P}^n\}_{n \in \mathbb{N}}$.

Step 2. We show that the limit point of $\{\mathbb{P}^n\}_{n \in \mathbb{N}}$ is \mathbb{P} .

Assume that the limit point of $\{\mathbb{P}^n\}_{n \in \mathbb{N}}$ is $\bar{\mathbb{P}}$. Since \mathbb{P} is a martingale solution of Eq.(17) with the initial law μ_0 , and Eq.(10) has a unique weak solution, by Theorem 2.7 we only need to prove that $\bar{\mathbb{P}}$ is a martingale solution of Eq.(17) with the initial law μ_0 . That is, it is sufficient to check that for $0 \leq s < t \leq T$ and a bounded continuous $\bar{\mathcal{B}}_s$ -measurable functional $\chi_s : D_T \mapsto \mathbb{R}$,

$$\int_{D_T} \left[\phi(w_t) - \phi(w_s) - \int_s^t (\mathcal{A}_r \phi + \mathcal{B}_r \phi)(w_r) dr \right] \chi_s(w) \bar{\mathbb{P}}(dw) = 0, \quad \forall \phi \in C_c^2(\mathbb{R}^d). \quad (21)$$

Next, note that $\mathbb{P}^n \circ e_t^{-1} \rightarrow \bar{\mathbb{P}} \circ e_t^{-1}$ in $\mathcal{P}(\mathbb{R}^d)$ and $\mathbb{P}^n \circ e_0^{-1} = \mu_0 = \bar{\mathbb{P}} \circ e_0^{-1}$. Thus, by (ii), there exists a $\bar{v}(t, x) \geq 0$ with $\int_{\mathbb{R}^d} \bar{v}(t, x) dx = 1$ such that $\bar{\mathbb{P}} \circ e_t^{-1}(dx) = \bar{v}(t, x) dx$ and $v^n(t, \cdot) \rightarrow \bar{v}(t, \cdot)$ in $w^* - L^\infty(\mathbb{R}^d)$, where $w^* - L^\infty(\mathbb{R}^d)$ is the dual space of $C_c(\mathbb{R}^d)$, and $\bar{v}(0, x) = v_0(x)$. Besides, (20) admits us to obtain that

$$\sup_n \sup_{t \in [0, T]} \int_{\mathbb{R}^d} |x| v^n(t, x) dx \leq C. \quad (22)$$

By suitable approximation, it holds that

$$\sup_{t \in [0, T]} \int_{\mathbb{R}^d} |x| \bar{v}(t, x) dx < \infty.$$

In the following, set $\nu_{t,x}(dz) := \nu(dg^{-1}(t, x, \cdot)(z))$, and then

$$\mathcal{B}_t \phi(x) = \int_{\mathbb{R}^d} [\phi(x + \gamma z) - \phi(x)] \nu_{t,x}(dz).$$

Thus, based on Lemma 4.1 in [2] and Lemma 3.2 below, we know that for any $\varepsilon > 0$ and the coefficients b, a , there exist $\tilde{b} : [0, T] \times \mathbb{R}^d \mapsto \mathbb{R}^d$, $\tilde{a} : [0, T] \times \mathbb{R}^d \mapsto \mathbb{S}_+(\mathbb{R}^d)$, where $\mathbb{S}_+(\mathbb{R}^d)$ is the set of nonnegative definite symmetric $d \times d$ real matrices, and a family of measures $\tilde{\nu}_\cdot$ such that

(v) \tilde{b}, \tilde{a} are continuous and compactly supported;

(vi) for any $\phi \in C_c^2(\mathbb{R}^d)$, $(t, x) \mapsto \tilde{\mathcal{B}}_t \phi(x)$ is continuous, where $\tilde{\mathcal{B}}_t \phi(x) := \int_{\mathbb{R}^d} [\phi(x + \gamma z) - \phi(x)] \tilde{\nu}_{t,x}(dz)$, and $\sup_{t \in [0, T], x \in \mathbb{R}^d} |\tilde{\mathcal{B}}_t \phi(x)| < \infty$;

(vii)

$$\int_0^T \int_{\mathbb{R}^d} \left(|b(t, x) - \tilde{b}(t, x)| + \frac{\|a(t, x) - \tilde{a}(t, x)\|}{1 + |x|} + |\mathcal{B}_t \phi(x) - \tilde{\mathcal{B}}_t \phi(x)| \right) \bar{v}(t, x) dx dt < \varepsilon.$$

And then the operators with respect to $\tilde{b}, \tilde{a}, \tilde{\nu}_\cdot$ are denoted as $\tilde{\mathcal{A}}_t + \tilde{\mathcal{B}}_t$.

Now, we treat (21). Inserting $\tilde{\mathcal{A}}_t + \tilde{\mathcal{B}}_t$, one can estimate (21) to get

$$\begin{aligned} & \left| \int_{D_T} \left[\phi(w_t) - \phi(w_s) - \int_s^t (\mathcal{A}_r \phi + \mathcal{B}_r \phi)(w_r) dr \right] \chi_s(w) \bar{\mathbb{P}}(dw) \right| \\ & \leq \left| \int_{D_T} \left[\phi(w_t) - \phi(w_s) - \int_s^t (\tilde{\mathcal{A}}_r \phi + \tilde{\mathcal{B}}_r \phi)(w_r) dr \right] \chi_s(w) \bar{\mathbb{P}}(dw) \right| \\ & \quad + \left| \int_{D_T} \left[\int_s^t \left((\tilde{\mathcal{A}}_r \phi + \tilde{\mathcal{B}}_r \phi)(w_r) - (\mathcal{A}_r \phi + \mathcal{B}_r \phi)(w_r) \right) dr \right] \chi_s(w) \bar{\mathbb{P}}(dw) \right| \\ & =: I_1 + I_2. \end{aligned} \tag{23}$$

To deal with I_1 , we recall that \mathbb{P}^n is a martingale solution of Eq.(18) with the initial law μ_0 , which means that

$$\int_{D_T} \left[\phi(w_t) - \phi(w_s) - \int_s^t (\mathcal{A}_r^n \phi + \mathcal{B}_r^n \phi)(w_r) dr \right] \chi_s(w) \mathbb{P}^n(dw) = 0,$$

where $\mathcal{A}_r^n + \mathcal{B}_r^n$ stands for the generator of Eq.(18). So,

$$\begin{aligned} & \int_{D_T} \left[\phi(w_t) - \phi(w_s) - \int_s^t (\tilde{\mathcal{A}}_r \phi + \tilde{\mathcal{B}}_r \phi)(w_r) dr \right] \chi_s(w) \mathbb{P}^n(dw) \\ & = \int_{D_T} \left[\int_s^t \left((\mathcal{A}_r^n \phi + \mathcal{B}_r^n \phi)(w_r) - (\tilde{\mathcal{A}}_r \phi + \tilde{\mathcal{B}}_r \phi)(w_r) \right) dr \right] \chi_s(w) \mathbb{P}^n(dw), \end{aligned}$$

and furthermore

$$\begin{aligned} & \left| \int_{D_T} \left[\phi(w_t) - \phi(w_s) - \int_s^t (\tilde{\mathcal{A}}_r \phi + \tilde{\mathcal{B}}_r \phi)(w_r) dr \right] \chi_s(w) \mathbb{P}^n(dw) \right| \\ & = \left| \int_{D_T} \left[\int_s^t \left((\mathcal{A}_r^n \phi + \mathcal{B}_r^n \phi)(w_r) - (\tilde{\mathcal{A}}_r \phi + \tilde{\mathcal{B}}_r \phi)(w_r) \right) dr \right] \chi_s(w) \mathbb{P}^n(dw) \right| \\ & \leq C \int_{D_T} \int_s^t \left| (\mathcal{A}_r^n \phi + \mathcal{B}_r^n \phi)(w_r) - (\tilde{\mathcal{A}}_r \phi + \tilde{\mathcal{B}}_r \phi)(w_r) \right| dr \mathbb{P}^n(dw) \\ & \leq C \int_s^t \int_{\mathbb{R}^d} \left[|(b_i^n(r, x) - \tilde{b}_i(r, x)) \partial_i \phi(x)| + |(a_{ij}^n(r, x) - \tilde{a}_{ij}(r, x)) \partial_{ij} \phi(x)| \right] \end{aligned}$$

$$\begin{aligned}
& + |\mathcal{B}_r^n \phi(x) - \mathcal{B}_r \phi(x)| + |\mathcal{B}_r \phi(x) - \tilde{\mathcal{B}}_r \phi(x)| \Big] v^n(r, x) dx dr \\
\leq & C \int_s^t \int_{\mathbb{R}^d} \left[|(b_i^n(r, x) - \tilde{b}_i(r, x)) \partial_i \phi(x)| + |(a_{ij}^n(r, x) - \tilde{a}_{ij}(r, x)) \partial_{ij} \phi(x)| \right. \\
& \left. + \int_{\mathbb{R}^d} |\phi(x + \gamma^n z) - \phi(x + \gamma z)| \nu_{r,x}(dz) + |\mathcal{B}_r \phi(x) - \tilde{\mathcal{B}}_r \phi(x)| \right] v^n(r, x) dx dr \\
\leq & C \int_s^t \int_{\mathbb{R}^d} \left[|(b_i^n(r, x) - \tilde{b}_i(r, x)) \partial_i \phi(x)| + |(a_{ij}^n(r, x) - \tilde{a}_{ij}(r, x)) \partial_{ij} \phi(x)| \right. \\
& \left. + \|\phi\|_{C_c^2(\mathbb{R}^d)} \int_{\mathbb{R}^d} I_{|x+\gamma z| \leq M} |\gamma^n - \gamma| |z| \nu_{r,x}(dz) + \int_{\mathbb{R}^d} I_{|x+\gamma z| > M} |\phi(x + \gamma^n z)| \nu_{r,x}(dz) \right. \\
& \left. + |\mathcal{B}_r \phi(x) - \tilde{\mathcal{B}}_r \phi(x)| \right] v^n(r, x) dx dr, \tag{24}
\end{aligned}$$

where the fact $\text{supp}(\phi) \subset B_M$ for $M > 0$ is applied in the last inequality. As $n \rightarrow \infty$, based on (i) (v) (vi) and $v^n(r, \cdot) \rightarrow \bar{v}(r, \cdot)$ in $w^* - L^\infty(\mathbb{R}^d)$, (24) yields that

$$\begin{aligned}
I_1 & \leq C \int_s^t \int_{\mathbb{R}^d} \left[|(b_i(r, x) - \tilde{b}_i(r, x)) \partial_i \phi(x)| + |(a_{ij}(r, x) - \tilde{a}_{ij}(r, x)) \partial_{ij} \phi(x)| \right. \\
& \quad \left. + |\mathcal{B}_r \phi(x) - \tilde{\mathcal{B}}_r \phi(x)| \right] \bar{v}(r, x) dx dr \\
& \leq C \int_s^t \int_{\mathbb{R}^d} \left[|b(r, x) - \tilde{b}(r, x)| + \|a(r, x) - \tilde{a}(r, x)\| I_{|x| \leq M} \right. \\
& \quad \left. + |\mathcal{B}_r \phi(x) - \tilde{\mathcal{B}}_r \phi(x)| \right] \bar{v}(r, x) dx dr \\
& \leq C \int_s^t \int_{\mathbb{R}^d} \left[|b(r, x) - \tilde{b}(r, x)| + \frac{\|a(r, x) - \tilde{a}(r, x)\|}{1 + |x|} \right. \\
& \quad \left. + |\mathcal{B}_r \phi(x) - \tilde{\mathcal{B}}_r \phi(x)| \right] \bar{v}(r, x) dx dr \\
& < C\varepsilon, \tag{25}
\end{aligned}$$

where we use $I_{|x| \leq M} \leq (1 + M) \frac{1}{1 + |x|}$ and (vii) in the third and fourth inequality, respectively.

In the following, we treat I_2 . By the similar deduction to that in (25), one can obtain that

$$\begin{aligned}
I_2 & \leq C \int_{D_T} \int_s^t \left| (\tilde{\mathcal{A}}_r \phi + \tilde{\mathcal{B}}_r \phi)(w_r) - (\mathcal{A}_r \phi + \mathcal{B}_r \phi)(w_r) \right| dr \bar{\mathbb{P}}(dw) \\
& \leq C \int_s^t \int_{\mathbb{R}^d} \left[|(b_i(r, x) - \tilde{b}_i(r, x)) \partial_i \phi(x)| + |(a_{ij}(r, x) - \tilde{a}_{ij}(r, x)) \partial_{ij} \phi(x)| \right. \\
& \quad \left. + |\mathcal{B}_r \phi(x) - \tilde{\mathcal{B}}_r \phi(x)| \right] \bar{v}(r, x) dx dr \\
& < C\varepsilon. \tag{26}
\end{aligned}$$

Combining (25) (26) with (23), we get that

$$\left| \int_{D_T} \left[\phi(w_t) - \phi(w_s) - \int_s^t (\mathcal{A}_r \phi + \mathcal{B}_r \phi)(w_r) dr \right] \chi_s(w) \bar{\mathbb{P}}(dw) \right| < C\varepsilon.$$

Letting $\varepsilon \rightarrow 0$, we finally have (21). The proof is complete. \square

Lemma 3.2. *For any $\varepsilon > 0$, there is a family of measures $\tilde{\nu}_\cdot$ such that for any $\phi \in C_c^2(\mathbb{R}^d)$,*

$$(i) \quad (t, x) \mapsto \tilde{\mathcal{B}}_t \phi(x) \text{ is continuous and } \sup_{t \in [0, T], x \in \mathbb{R}^d} |\tilde{\mathcal{B}}_t \phi(x)| < \infty;$$

(ii)

$$\int_0^T \int_{\mathbb{R}^d} |\mathcal{B}_t \phi(x) - \tilde{\mathcal{B}}_t \phi(x)| \bar{v}(t, x) dx dt < \varepsilon.$$

Proof. The method comes from [11, Lemma 3.8]. By [4, Lemma 14.50, Page 469], it holds that there is a measurable function

$$h_{t,x}(\theta) : [0, T] \times \mathbb{R}^d \times [0, \infty) \mapsto \mathbb{R}^d \cup \{\infty\},$$

such that for $t \in [0, T]$ and $x \in \mathbb{R}^d$

$$\nu_{t,x}(A) = \int_0^\infty I_A(h_{t,x}(\theta)) d\theta, \quad \forall A \in \mathcal{B}(\mathbb{R}^d).$$

And then for any $\phi \in C_c^2(\mathbb{R}^d)$,

$$\mathcal{B}_t \phi(x) = \int_{\mathbb{R}^d} [\phi(x + \gamma z) - \phi(x)] \nu_{t,x}(dz) = \int_0^\infty [\phi(x + \gamma h_{t,x}(\theta)) - \phi(x)] d\theta.$$

Next, by the theory of functional analysis, we know that there exists a sequence of measurable functions $\{h_{t,x}^n(\theta), n \in \mathbb{N}\}$ such that for any $\theta \geq 0$ and $n \in \mathbb{N}$, $(t, x) \mapsto h_{t,x}^n(\theta)$ is continuous with compact support, $|h_{t,x}^n(\theta)| \leq |h_{t,x}(\theta)|$ and

$$\lim_{n \rightarrow \infty} \int_0^T \int_{\mathbb{R}^d} \int_0^\infty (|h_{t,x}^n(\theta) - h_{t,x}(\theta)|^2 \wedge 1) \bar{v}(t, x) d\theta dx dt = 0.$$

Thus, for any $\varepsilon > 0$, there exists a $N \in \mathbb{N}$ such that

$$\int_0^T \int_{\mathbb{R}^d} \int_0^\infty (|h_{t,x}^N(\theta) - h_{t,x}(\theta)|^2 \wedge 1) \bar{v}(t, x) d\theta dx dt < \varepsilon. \quad (27)$$

Now, for $t \in [0, T]$ and $x \in \mathbb{R}^d$, set

$$\tilde{\nu}_{t,x}(A) := \int_0^\infty I_A(h_{t,x}^N(\theta)) d\theta, \quad \forall A \in \mathcal{B}(\mathbb{R}^d),$$

and then $(t, x) \mapsto \tilde{\nu}_{t,x}(A)$ is continuous. So, it holds that for any $\phi \in C_c^2(\mathbb{R}^d)$,

$$(t, x) \mapsto \tilde{\mathcal{B}}_t \phi(x) = \int_{\mathbb{R}^d} [\phi(x + \gamma z) - \phi(x)] \tilde{\nu}_{t,x}(dz)$$

is continuous. Besides, note that

$$|\tilde{\mathcal{B}}_t \phi(x)| \leq \int_0^\infty |\phi(x + \gamma h_{t,x}^N(\theta)) - \phi(x)| d\theta \leq C \int_0^\infty (|\gamma h_{t,x}^N(\theta)| \wedge 1) d\theta.$$

Since $h_{t,x}^N(\theta)$ has a compact support in (t, x) , we have that

$$\sup_{t \in [0, T], x \in \mathbb{R}^d} |\tilde{\mathcal{B}}_t \phi(x)| < \infty.$$

Thus, (i) is proved.

For (ii), we compute that

$$\begin{aligned} |\mathcal{B}_t \phi(x) - \tilde{\mathcal{B}}_t \phi(x)| &\leq \int_0^\infty |\phi(x + \gamma h_{t,x}(\theta)) - \phi(x + \gamma h_{t,x}^N(\theta))| d\theta \\ &\leq C \int_0^\infty \left(I_{B_l}(h_{t,x}(\theta)) I_{B_{R+|\gamma|l}}(x) + I_{B_{l \vee \frac{|x|-R}{|\gamma|}}}^c(h_{t,x}(\theta)) \right) \\ &\quad \times (|h_{t,x}^N(\theta) - h_{t,x}(\theta)| \wedge 1) d\theta \\ &\leq C \left(\int_0^\infty \left(I_{B_l}(h_{t,x}(\theta)) I_{B_{R+|\gamma|l}}(x) + I_{B_{l \vee \frac{|x|-R}{|\gamma|}}}^c(h_{t,x}(\theta)) \right) d\theta \right)^{1/2} \\ &\quad \times \left(\int_0^\infty (|h_{t,x}^N(\theta) - h_{t,x}(\theta)|^2 \wedge 1) d\theta \right)^{1/2} \\ &\leq C \left(I_{B_{R+|\gamma|l}}(x) \nu_{t,x}(B_l) + \nu_{t,x}(B_{l \vee \frac{|x|-R}{|\gamma|}}^c) \right)^{1/2} \\ &\quad \times \left(\int_0^\infty (|h_{t,x}^N(\theta) - h_{t,x}(\theta)|^2 \wedge 1) d\theta \right)^{1/2} \\ &\leq C \left(\int_0^\infty (|h_{t,x}^N(\theta) - h_{t,x}(\theta)|^2 \wedge 1) d\theta \right)^{1/2}, \end{aligned}$$

where $l > 0$ is a constant, and $\text{supp}(\phi) \subset B_R$ and $\nu_{t,x}(\mathbb{R}^d) < \infty$ are used in the second and fifth inequality, respectively. Therefore, the Hölder inequality admits us to obtain that

$$\begin{aligned} &\int_0^T \int_{\mathbb{R}^d} |\mathcal{B}_t \phi(x) - \tilde{\mathcal{B}}_t \phi(x)| \bar{v}(t, x) dx dt \\ &\leq C \int_0^T \int_{\mathbb{R}^d} \left(\int_0^\infty (|h_{t,x}^N(\theta) - h_{t,x}(\theta)|^2 \wedge 1) d\theta \right)^{1/2} \bar{v}(t, x) dx dt \\ &\leq CT^{1/2} \left(\int_0^T \int_{\mathbb{R}^d} \int_0^\infty (|h_{t,x}^N(\theta) - h_{t,x}(\theta)|^2 \wedge 1) d\theta \bar{v}(t, x) dx dt \right)^{1/2} \\ &\leq CT^{1/2} \varepsilon^{1/2}, \end{aligned}$$

where (27) is used in the last inequality. The proof is complete. \square

Remark 3.3. Assume that $b : [0, T] \times \mathbb{R}^d \mapsto \mathbb{R}^d$, $\sigma : [0, T] \times \mathbb{R}^d \mapsto \mathbb{S}_+(\mathbb{R}^d)$ are continuous and satisfy $(\mathbf{H}_{b,\sigma})$, $\gamma \in \mathbb{R}$, g satisfies (\mathbf{H}_f) , and that Eq.(10) has a unique weak solution in \mathcal{L}_+ , and \mathbb{P} is a martingale solution of Eq.(17) with the initial law $\mu_0 = v_0(x) dx \in \mathcal{P}_1(\mathbb{R}^d)$. Set

$$b_i^n(t, x) := \int_{\mathbb{R}^d} \varphi_n(x - y) b_i(t, y) dy,$$

$$\begin{aligned}
a_{ij}^n(t, x) &:= \int_{\mathbb{R}^d} \varphi_n(x-y) a_{ij}(t, y) dy, \quad i, j = 1, 2, \dots, d, \\
\sigma^n(t, x) &:= \sqrt{2a^n(t, x)}, \\
\gamma^n &:= \frac{n}{n+1} \gamma,
\end{aligned}$$

where $\varphi \in C_c^\infty(\mathbb{R}^d)$ is a nonnegative mollifier with the support in B_1 and $\int_{\mathbb{R}^d} \varphi(x) dx = 1$, $\varphi_n(x) = n^d \varphi(nx)$. And then we also assume that \mathbb{P}^n is a martingale solution of the corresponding Eq.(18) with the initial law μ_0 , $\mathbb{P}^n \circ e_t^{-1}$ is absolutely continuous with respect to the Lebesgue measure on \mathbb{R}^d and $v^n(t, x)$ denotes the density, i.e., $v^n(t, x) := \frac{(\mathbb{P}^n \circ e_t^{-1})(dx)}{dx}$ for any $t \in [0, T]$, and

$$\sup_{x \in \mathbb{R}^d} |v^n(t, x)| \leq C,$$

where the constant $C > 0$ only depends on T . So, $b^n, \sigma^n, \gamma^n, a^n, \mathbb{P}^n$ satisfy the conditions of Theorem 3.1. Thus, it holds that $\mathbb{P}^n \rightarrow \mathbb{P}$ in $\mathcal{P}(D_T)$.

For the readers' convenience, we put the verification of b^n, σ^n, a^n satisfying the conditions of Theorem 3.1 in the appendix.

Remark 3.4. If $\gamma^n = \gamma = 0$, b^n, b, σ^n, σ are uniformly bounded, Theorem 3.1 reduces to [1, Theorem 3.7]. Therefore, Theorem 3.1 is more general.

In the following, we give some special cases. If we remove the jump term from Eq.(17), i.e.

$$dX_t = b(t, X_t)dt + \sigma(t, X_t)dB_t, \quad t \in [0, T]. \quad (28)$$

The following corollary describes the relationship between martingale solutions of Eq.(18) and that of Eq.(28).

Corollary 3.5. Suppose that $b^n, b : [0, T] \times \mathbb{R}^d \mapsto \mathbb{R}^d$, $\sigma^n, \sigma : [0, T] \mapsto \mathbb{R}^{d \times m}$, $\{\gamma^n\}$ are uniformly bounded, g satisfies (\mathbf{H}_f) , and that $b \in L^1([0, T], BV_{loc}(\mathbb{R}^d, \mathbb{R}^d))$, $\partial_i b_i \in L^1_{loc}([0, T] \times \mathbb{R}^d)$, $(\partial_i b_i)^- \in L^1([0, T], L^\infty(\mathbb{R}^d))$. Let $\mu_0(dx) = v_0(x)dx \in \mathcal{P}_1(\mathbb{R}^d)$ with $\|v_0\|_\infty < \infty$, and \mathbb{P}^n, \mathbb{P} be martingale solutions of Eq.(18) and Eq.(28) with the initial law μ_0 , respectively. Assume that

- (i) $b^n \rightarrow b$ in $L^1_{loc}([0, T] \times \mathbb{R}^d)$, $a^n \rightarrow a$ in $L^1_{loc}([0, T])$, $\gamma^n \rightarrow 0$ as $n \rightarrow \infty$;
- (ii) $\mathbb{P}^n \circ e_t^{-1}$ is absolutely continuous with respect to the Lebesgue measure on \mathbb{R}^d and $v^n(t, x)$ denotes the density, i.e., $v^n(t, x) := \frac{(\mathbb{P}^n \circ e_t^{-1})(dx)}{dx}$ for any $t \in [0, T]$, and

$$\sup_{x \in \mathbb{R}^d} |v^n(t, x)| \leq C,$$

where the constant $C > 0$ only depends on T .

Then $\mathbb{P}^n \rightarrow \mathbb{P}$ in $\mathcal{P}(D_T)$.

Proof. By [1, Theorem 4.12] and Proposition 2.5, it holds that Eq.(10) with $\mathcal{B}_t = 0$ has a unique weak solution in

$$\tilde{\mathcal{L}}_+ := \left\{ v \geq 0 : \int_{\mathbb{R}^d} v(t, x) dx = 1, \sup_{t \in [0, T]} \|v(t, \cdot)\|_\infty < \infty, \text{ and } \sup_{t \in [0, T]} \int_{\mathbb{R}^d} |x| v(t, x) dx < \infty \right\}.$$

Note that $\tilde{\mathcal{L}}_+ \subset \mathcal{L}_+$. Thus, the remaining proof is similar to that of Theorem 3.1. \square

Remark 3.6. *This corollary means that SDEs with jumps can converge to SDEs without jumps in some sense.*

If we remove the continuous diffusion term from Eq.(17) and take $\mathbb{U} \in \mathcal{B}(\mathbb{R}^d)$, $f(t, x, u) = \gamma u$, i.e.

$$dX_t = b(t, X_t)dt + \gamma \int_{\mathbb{U}} uN(dt, du), \quad t \in [0, T]. \quad (29)$$

The following corollary characterizes the relationship between martingale solutions of (18) and ones of (29).

Corollary 3.7. *Suppose that $b^n, b : [0, T] \times \mathbb{R}^d \mapsto \mathbb{R}^d$, $\sigma^n : [0, T] \times \mathbb{R}^d \mapsto \mathbb{R}^{d \times m}$, $\{\gamma^n\}$ are uniformly bounded and for some $q > 1$, $|\nabla b| \in L^\infty([0, T], L^q_{loc}(\mathbb{R}^d, \mathbb{R}^d))$, $(\partial_i b_i)^- \in L^\infty([0, T] \times \mathbb{R}^d)$, and for any $p \geq 1$,*

$$\int_{\mathbb{U}} |u|^2 (1 + |u|)^p \nu(du) < \infty.$$

Let $\mu_0(dx) = v_0(x)dx \in \mathcal{P}_1(\mathbb{R}^d)$ with

$$\int_{\mathbb{R}^d} v(0, x)^r (1 + |x|^2)^{(r-1)d} dx < \infty, \quad r > \frac{q}{q-1} = q^*,$$

and \mathbb{P}^n, \mathbb{P} be martingale solutions of Eq.(18) and Eq.(29) with the initial law μ_0 , respectively. Assume that

(i) $b^n \rightarrow b, a^n \rightarrow 0$ in $L^1_{loc}([0, T] \times \mathbb{R}^d)$, $\gamma^n \rightarrow \gamma$ as $n \rightarrow \infty$;

(ii) $\mathbb{P}^n \circ e_t^{-1}$ is absolutely continuous with respect to the Lebesgue measure on \mathbb{R}^d and $v^n(t, x)$ denotes the density, i.e., $v^n(t, x) := \frac{(\mathbb{P}^n \circ e_t^{-1})(dx)}{dx}$ for any $t \in [0, T]$, and

$$\sup_{x \in \mathbb{R}^d} |v^n(t, x)| \leq C,$$

where the constant $C > 0$ only depends on T .

Then $\mathbb{P}^n \rightarrow \mathbb{P}$ in $\mathcal{P}(D_T)$.

Proof. By [13, Theorem 5.7] and Proposition 2.5, we know that Eq.(10) with $a = 0$ has a unique weak solution in

$$\begin{aligned} \hat{\mathcal{L}}_+ := \left\{ v \geq 0 : \int_{\mathbb{R}^d} v(t, x) dx = 1, \sup_{t \in [0, T]} \int_{\mathbb{R}^d} v(t, x)^{q^*} (1 + |x|^2)^{(q^*-1)d} dx < \infty, \right. \\ \left. \text{and } \sup_{t \in [0, T]} \int_{\mathbb{R}^d} |x| v(t, x) dx < \infty \right\}. \end{aligned}$$

Note that $\hat{\mathcal{L}}_+ \subset \mathcal{L}_+$. Therefore, the remaining proof continues as that of Theorem 3.1. \square

Remark 3.8. *This corollary means that SDEs with jumps can converge to SDEs with pure jumps in some sense.*

If Eq.(17) has no continuous diffusion term and no jump term, i.e.

$$dX_t = b(t, X_t)dt, \quad t \in [0, T], \quad (30)$$

Corollary 3.5 yields the following corollary on martingale solutions of Eq.(18) and Eq.(30).

Corollary 3.9. *Suppose that $b^n, b : [0, T] \times \mathbb{R}^d \mapsto \mathbb{R}^d$, $\sigma^n : [0, T] \mapsto \mathbb{R}^{d \times m}$, $\{\gamma^n\}$ are uniformly bounded, g satisfies (\mathbf{H}_f) , and that $b \in L^1([0, T], BV_{loc}(\mathbb{R}^d, \mathbb{R}^d))$, $\partial_i b_i \in L^1_{loc}([0, T] \times \mathbb{R}^d)$, $(\partial_i b_i)^- \in L^1([0, T], L^\infty(\mathbb{R}^d))$. Let $\mu_0(dx) = v_0(x)dx \in \mathcal{P}_1(\mathbb{R}^d)$ with $\|v_0\|_\infty < \infty$, and \mathbb{P}^n, \mathbb{P} be martingale solutions of Eq.(18) and Eq.(30) with the initial law μ_0 , respectively.*

Assume that

- (i) $b^n \rightarrow b, a^n \rightarrow 0$ in $L^1_{loc}([0, T] \times \mathbb{R}^d)$, $\gamma^n \rightarrow 0$ as $n \rightarrow \infty$;
- (ii) $\mathbb{P}^n \circ e_t^{-1}$ is absolutely continuous with respect to the Lebesgue measure on \mathbb{R}^d and $v^n(t, x)$ denotes the density, i.e., $v^n(t, x) := \frac{(\mathbb{P}^n \circ e_t^{-1})(dx)}{dx}$ for any $t \in [0, T]$, and

$$\sup_{x \in \mathbb{R}^d} |v^n(t, x)| \leq C,$$

where the constant $C > 0$ only depends on T .

Then $\mathbb{P}^n \rightarrow \mathbb{P}$ in $\mathcal{P}(D_T)$.

Remark 3.10. *This corollary means that SDEs with jumps can converge to ordinary differential equations in some sense.*

Remark 3.11. *If $\gamma^n = 0$ and $b^n = b, \sigma^n = \frac{1}{n}I_d$, the above corollary is just right [1, Corollary 3.9].*

4. WEAK SOLUTIONS OF FPEs IN \mathcal{L}_+

In this section, we take $\mathbb{U} \in \mathcal{B}(\mathbb{R}^d)$ and $f(t, x, u) = \gamma u$ and assume that for any $p \geq 1$

$$\int_{\mathbb{U}} |u|^2 (1 + |u|)^p \nu(du) < \infty.$$

That is, Eq.(10) becomes

$$\partial_t v = -\partial_i (b_i v) + \partial_{ij} (a_{ij} v) + \int_{\mathbb{U}} [v(\cdot - \gamma u) - v(\cdot)] \nu(du). \quad (31)$$

We give some sufficient conditions for Eq.(31) to have a unique weak solution in \mathcal{L}_+ .

The main result in the section is the following theorem.

Theorem 4.1. *Suppose that $b : [0, T] \times \mathbb{R}^d \mapsto \mathbb{R}^d$, $\sigma : [0, T] \times \mathbb{R}^d \mapsto \mathbb{S}_+(\mathbb{R}^d)$ satisfy $(\mathbf{H}_{b, \sigma})$ and that for some $q > 1$, $|\nabla b| \in L^1([0, T], L^q_{loc}(\mathbb{R}^d, \mathbb{R}^d))$, $(\partial_i b_i)^- \in L^1([0, T], L^\infty(\mathbb{R}^d))$, $\|\nabla \sigma\|^2 \in L^1([0, T], L^\infty(\mathbb{R}^d))$. Then for $\mu_0(dx) = v_0(x)dx \in \mathcal{P}_1(\mathbb{R}^d)$ with*

$$\int_{\mathbb{R}^d} v_0(x)^r (1 + |x|^2)^{(r-1)d} dx < \infty$$

for some $r > 1$, Eq.(31) has a unique weak solution in \mathcal{L}_+ .

Proof. First of all, we take a complete filtered probability space $(\check{\Omega}, \check{\mathcal{F}}, \check{\mathbb{P}}; (\check{\mathcal{F}}_t)_{t \in [0, T]})$, an $(\check{\mathcal{F}}_t)$ -adapted Brownian motion \check{B}_t and an $(\check{\mathcal{F}}_t)$ -adapted Poisson random measure $\check{N}(dt, du)$ independent of \check{B}_t with the intensity measure $dt\nu(du)$. We consider the following equation:

$$\begin{aligned} \check{X}_t(x) &= x + \int_0^t \left(b(s, \check{X}_s(x)) - \int_{\mathbb{U}} \gamma u \nu(du) \right) ds + \int_0^t \sigma(s, \check{X}_s(x)) d\check{B}_s \\ &\quad + \int_0^t \int_{\mathbb{U}} \gamma u \check{N}(dsdu), \quad x \in \mathbb{R}^d, \quad t \in [0, T], \end{aligned} \quad (32)$$

where $\tilde{N}(dsdu) := \tilde{N}(dsdu) - \nu(du)ds$ is the compensated martingale measure of $\tilde{N}(dsdu)$. By [13, Theorem 4.2], one can obtain that there exists a unique weak solution $\tilde{X}_t(x)$ of Eq.(32) satisfying

$$\sup_{t \in [0, T]} \tilde{\mathbb{E}} \left(\int_{\mathbb{R}^d} |\psi(\tilde{X}_t(x))|^{r^*} \rho(dx) \right) \leq C \|\psi\|_{L_{\rho}^{r^*}}^{r^*}, \quad \forall \psi \in C_c^\infty(\mathbb{R}^d), \quad (33)$$

where C is independent of ψ , $\tilde{\mathbb{E}}$ denotes the expectation under the probability measure $\tilde{\mathbb{P}}$ and

$$\frac{1}{r} + \frac{1}{r^*} = 1, \quad \rho(dx) = \frac{1}{(1 + |x|^2)^d} dx, \quad \|\psi\|_{L_{\rho}^{r^*}}^{r^*} := \int_{\mathbb{R}^d} |\psi(x)|^{r^*} \rho(dx).$$

Next, we choose a $\tilde{\mathcal{F}}_0$ -measurable d -dimensional random vector \tilde{X}_0 such that $\tilde{\mathbb{P}} \circ \tilde{X}_0^{-1} = \mu_0$. Thus, $\bar{X}_t := \tilde{X}_t(\tilde{X}_0)$ uniquely solves the following equation

$$\begin{aligned} \bar{X}_t &= \tilde{X}_0 + \int_0^t \left(b(s, \bar{X}_s) - \int_{\mathbb{U}} \gamma u \nu(du) \right) ds + \int_0^t \sigma(s, \bar{X}_s) d\tilde{B}_s \\ &\quad + \int_0^t \int_{\mathbb{U}} \gamma u \tilde{N}(dsdu). \end{aligned}$$

Moreover, it follows from the Hölder inequality and the Jensen inequality that for any $\psi \in C_c^\infty(\mathbb{R}^d)$,

$$\begin{aligned} \left| \int_{\mathbb{R}^d} \psi(x) \mathcal{L}_{\bar{X}_t}(dx) \right| &= |\tilde{\mathbb{E}} \psi(\bar{X}_t)| = |\tilde{\mathbb{E}} [\tilde{\mathbb{E}}[\psi(\tilde{X}_t(x)) | x = \tilde{X}_0]]| = \left| \int_{\mathbb{R}^d} \tilde{\mathbb{E}}[\psi(\tilde{X}_t(x))] v_0(x) dx \right| \\ &\leq \left(\int_{\mathbb{R}^d} |\tilde{\mathbb{E}}[\psi(\tilde{X}_t(x))]|^{r^*} \rho(dx) \right)^{1/r^*} \left(\int_{\mathbb{R}^d} v_0(x)^r (1 + |x|^2)^{rd} \rho(dx) \right)^{1/r} \\ &\leq \left(\tilde{\mathbb{E}} \int_{\mathbb{R}^d} |\psi(\tilde{X}_t(x))|^{r^*} \rho(dx) \right)^{1/r^*} \left(\int_{\mathbb{R}^d} v_0(x)^r (1 + |x|^2)^{(r-1)d} dx \right)^{1/r} \\ &\leq C \|\psi\|_{L_{\rho}^{r^*}}, \end{aligned}$$

where the last inequality is based on (33). By the theory of functional analysis, we know that there exists a $v(t, \cdot)(1 + |\cdot|^2)^d \in L_{\rho}^r$ such that

$$\int_{\mathbb{R}^d} \psi(x) v(t, x) (1 + |x|^2)^d \rho(dx) = \int_{\mathbb{R}^d} \psi(x) v(t, x) dx = \int_{\mathbb{R}^d} \psi(x) \mathcal{L}_{\bar{X}_t}(dx).$$

So, [6, 4.25 Problem, Page 325] gives $\mathcal{L}_{\bar{X}_t}(dx) = v(t, x) dx$ for any $t \in [0, T]$. By Proposition 2.6, it holds that v uniquely solves the FPE (31) in the distribution sense. Finally, using $\mu_0(dx) = v_0(x) dx \in \mathcal{P}_1(\mathbb{R}^d)$ and Proposition 2.5, we conclude that $v \in \mathcal{L}_+$. The proof is complete. \square

Remark 4.2. *These conditions in Theorem 4.1 are not optimal, because under them a pathwise unique strong solution of Eq.(32) exists ([13, Theorem 4.2]). Thus, if one tries to weaken them to assure existence and uniqueness of weak solutions to the FPE (31), some techniques from the theory of partial differential equations are needed ([1]). Here, we do not pursue this.*

5. APPENDIX

In this section, we show that $b^n, \sigma^n, a^n, \mathbb{P}^n$ satisfy the conditions in Theorem 3.1.

Proof of Remark 3.3.

(i) b^n, σ^n satisfy $(\mathbf{H}_{b,\sigma})$.

First of all, we show that b^n satisfies $(\mathbf{H}_{b,\sigma})$. For any $t \in [0, T]$, $x \in \mathbb{R}^d$,

$$\begin{aligned} |b^n(t, x)| &\leq \int_{\mathbb{R}^d} \varphi_n(x-y) |b(t, y)| dy \leq C \int_{\mathbb{R}^d} \varphi_n(x-y) (1+|y|) dy \\ &\leq C \int_{\mathbb{R}^d} \varphi_n(x-y) (1+|x-y|+|x|) dy \\ &\leq C(1+|x|) + C \int_{\mathbb{R}^d} \varphi_n(x-y) |x-y| dy. \end{aligned}$$

Note that

$$\int_{\mathbb{R}^d} \varphi_n(x-y) |x-y| dy = \int_{\mathbb{R}^d} n^d \varphi(n(x-y)) |x-y| dy = \int_{\mathbb{R}^d} \varphi(v) \frac{|v|}{n} dv \leq 1.$$

Thus, all the above computation yields that

$$|b^n(t, x)| \leq C(1+|x|),$$

where C is independent of n .

For σ^n , we only need to verify that $\|a^n(t, x)\| \leq C(1+|x|)^2$. Note that $\|a(t, y)\| \leq C(1+|y|)^2$ and $\text{supp}(\varphi) = B_1$. The remaining proof is similar to that of b^n .

(ii) $b^n \rightarrow b, a^n \rightarrow a$ in $L^1_{loc}([0, T] \times \mathbb{R}^d)$.

For any $0 \leq s < t \leq T$ and B_R ,

$$\begin{aligned} &\int_s^t \int_{B_R} (|b^n(r, x) - b(r, x)| + \|a^n(r, x) - a(r, x)\|) dx dr \\ &\leq \int_s^t \int_{B_R} \int_{\mathbb{R}^d} \varphi_n(x-y) (|b(r, y) - b(r, x)| + \|a(r, y) - a(r, x)\|) dy dx dr \\ &= \int_s^t \int_{B_R} \int_{\mathbb{R}^d} \varphi(v) \left(|b(r, x - \frac{v}{n}) - b(r, x)| + \|a(r, x - \frac{v}{n}) - a(r, x)\| \right) dv dx dr. \end{aligned}$$

Thus, continuity of b, σ in the spatial variable and the dominated convergence theorem allow us to obtain that the above limit is zero as $n \rightarrow \infty$.

Acknowledgements:

The author is very grateful to Professor Xicheng Zhang and Renming Song for valuable discussions.

REFERENCES

- [1] A. Figalli: Existence and uniqueness of martingale solutions for SDEs with rough or degenerate coefficients. *J. Funct. Anal.*, 254 (2008)109153.
- [2] N. Fournier and L. Xu: On the equivalence between some jumping SDEs with rough coefficients and some non-local PDEs. *Ann. Inst. Henri Poincaré Probab. Stat.*, 55 (2019)1163-1178.
- [3] N. Ikeda and S. Watanabe: *Stochastic Differential Equations and Diffusion Processes*, 2nd ed., North-Holland/Kodansha, Amsterdam/Tokyo, 1989.
- [4] J. Jacod: *Calcul stochastique et problèmes de martingales*. Lecture Notes in Mathematics 714. Springer, Berlin, 1979.

- [5] J. Jacod and A.N. Shiryaev: *Limit Theorems for Stochastic Processes*. Springer-Verlag, Berlin, 1987.
- [6] I. Karatzas and S. E. Shreve: *Brownian motion and stochastic calculus*. 2nd ed., Graduate Texts in Mathematics, vol. 113, Springer-Verlag, New York, 2005: 284-295.
- [7] H. Qiao: Exponential ergodicity for SDEs with jumps and non-Lipschitz coefficients, *Journal of Theoretical Probability*, 27(2014)137-152.
- [8] H. Qiao: Euler-Maruyama approximation for SDEs with jumps and non-Lipschitz coefficients, *Osaka Journal of Mathematics*, 51(2014)47-66.
- [9] H. Qiao and J. Duan: Stationary measure for stochastic differential equations with jumps, *Stochastics*, 88(2016)864-883.
- [10] H. Qiao and X. Zhang: Homeomorphism flows for non-Lipschitz stochastic differential equations with jumps, *Stochastic Processes and their Applications*, 118(2008)2254-2268.
- [11] M. Röckner, L. Xie and X. Zhang: Superposition principle for non-local Fokker-Planck operators. *arXiv*: 1910.11022v1.
- [12] D.W. Stroock, S.R.S. Varadhan: *Multidimensional Diffusion Processes*, Grundlehren Math. Wiss., vol. 233, Springer-Verlag, Berlin, 1979.
- [13] X. Zhang: Degenerate irregular SDEs with jumps and application to integro-differential equations of Fokker-Planck type. *Electron. J. Probab.*, 18(2013)1-25.