

# THE CONE THEOREM AND THE VANISHING OF CHOW COHOMOLOGY

DAN EDIDIN AND RYAN RICHEY

ABSTRACT. We show that a cone theorem for  $\mathbb{A}^1$ -homotopy invariant contravariant functors implies the vanishing of the positive degree part of the operational Chow cohomology rings of a large class of affine varieties. We also discuss how this vanishing relates to a number of questions about representing Chow cohomology classes of GIT quotients in terms of equivariant cycles.

## 1. INTRODUCTION

In [FM, Ful], Fulton and MacPherson define for any scheme  $X$  a graded cohomology ring  $A_{\text{op}}^*(X)$  which equals the classical intersection ring when  $X$  is non-singular. An element of  $A_{\text{op}}^*(X)$  is a collection of operations on Chow groups of  $X$ -schemes compatible with basic operations in intersection theory. The product structure is given by composition. By construction there is a pullback of operational rings  $A_{\text{op}}^*(Y) \rightarrow A_{\text{op}}^*(X)$  for any morphism  $X \rightarrow Y$ .

For an arbitrary singular scheme elements of  $A_{\text{op}}^*(X)$  do not have natural interpretations in terms of algebraic cycles, although Kimura [Kim] showed how  $A_{\text{op}}^*(X)$  can be related to the intersection ring of a resolution of singularities of  $X$ . Despite the formal structure, there is a class of singular varieties where the operational Chow groups are readily computable. Specifically, if  $X$  is a complete linear variety then Totaro [Tot] proved that the pairing

$$A_{\text{op}}^k(X) \times A_k(X) \rightarrow A_0(X) = \mathbb{Z}, (c, \alpha) \mapsto c \cap \alpha$$

is perfect so  $A_{\text{op}}^k(X) = \text{Hom}(A_k(X), \mathbb{Z})$ , where  $A_k(X)$  denotes the classical Chow group of  $k$ -dimensional cycles. When  $X$  is a complete toric variety Fulton and Sturmfels [FS] gave an explicit description of the operational product in terms of Minkowski weights. This was generalized by Payne [Pay] who showed that the  $T$ -equivariant operational Chow ring of a toric variety  $X$  can be identified with the ring of integral piecewise polynomial functions on the fan of  $X$ .

The Cox construction expresses any toric variety as a good quotient  $(\mathbb{A}^n \setminus B)/G$  where  $G$  is a diagonalizable group. In [ES2] the first author and Satriano showed that if  $X = Z/G$  is the good quotient of a smooth variety by a linearly reductive group then the rational operational Chow ring  $A_{\text{op}}^*(X)_{\mathbb{Q}}$  naturally embeds in the equivariant Chow ring  $A_G^*(Z)_{\mathbb{Q}}$ . Moreover, the image of an element in  $A_{\text{op}}^k(X)$  is represented by a class of the form  $\sum c_i [Z_i]$  where the  $Z_i \subset Z$  are codimension- $k$   $G$ -invariant subvarieties of  $Z$  which are saturated with respect to the quotient map  $\pi: Z \rightarrow X$ .

By the étale slice theorem [Lun] the local model for the good quotient of a smooth variety at a closed orbit  $Gx \subset Z$  is the quotient  $V \rightarrow V/G_x$  where  $V = T_{x,Z}/T_{x,Gx}$  is the normal

---

2010 *Mathematics Subject Classification.* 14C15, 14F43, 14M25.

The first author was supported by Simons Collaboration Grant 315460.

space to the orbit  $Gx$  at  $x$ , and  $G_x$  is the stabilizer of  $x$ . Therefore, a natural problem is to compute the operational Chow rings of good quotients  $V/G$  where  $V$  is a representation of a linearly reductive group  $G$ . Examples of such quotients are affine toric varieties associated to maximal dimensional strongly convex polyhedral cones. In [Ric, ER1] it is shown that  $A^*(X) = \mathbb{Z}$  for any affine toric variety  $X$ , and likewise that  $\text{op } K^0(X) = \mathbb{Z}$  where  $\text{op } K^0$  is the operational  $K$ -theory defined by Anderson and Payne [AP].

The purpose of this paper is to show that stronger results hold. The contravariant functors  $A_{\text{op}}^*$  and  $\text{op } K^0$  are both  $\mathbb{A}^1$ -homotopy invariants. The cone theorem (Theorem 3.1) implies the vanishing of such functors on a large class of naturally occurring varieties including affine toric varieties and quotients of representations of reductive groups.

Since quotients of the form  $V/G$  are also the local models for good moduli spaces of Artin stacks [AHR] we conclude with a discussion of how the vanishing of  $A_{\text{op}}^*(V/G)$  relates to questions about the image of  $A_{\text{op}}^*(X)$  in  $A^*\mathcal{X}$  when  $X$  is the good moduli space of a smooth algebraic stack  $\mathcal{X}$ .

**Dedication.** It is a pleasure to dedicate this work to William Fulton on the occasion of his 80th birthday.

## 2. HOMOTOPY INVARIANT FUNCTORS

Fix a ground field  $k$  and let  $Sch/k$  denote the category of  $k$ -schemes of finite type.

**Definition 2.1.** A homotopy invariant functor is a contravariant functor  $H: Sch/k \rightarrow Ab$  such that the pullback  $H(X) \xrightarrow{\pi^*} H(X \times \mathbb{A}^1)$  is an isomorphism for all  $X$  in  $Sch/k$ . Likewise if  $G$  is an algebraic group then a  $G$ -homotopy invariant functor is a contravariant functor  $H^G: Sch^G/k \rightarrow Ab$  such that for any  $G$ -scheme  $X$  the pullback  $H^G(X) \xrightarrow{\pi^*} H^G(X \times \mathbb{A}^1)$  is an isomorphism, where the action of  $G$  on  $\mathbb{A}^1$  is trivial.

In this paper we focus on several homotopy invariant functors – the operational Chow cohomology ring defined in [FM] and [Ful, Chapter 17] as well as its equivariant counterpart defined in [EG] and the (equivariant) operational  $K$ -theory defined by Anderson and Payne in [AP].

**2.1. Chow cohomology.** Let  $X$  be a scheme. Following [Ful], let  $A_k(X)$  denote the group of dimension  $k$  cycle classes modulo rational equivalence, and if  $X$  is equidimensional of dimension  $n$  we let  $A^k(X)$  denote the group of  $(n - k)$ -dimensional cycle classes modulo rational equivalence. If  $X$  is smooth and equidimensional, then the intersection product on  $A_k(X)$  as constructed in [Ful, Chapter 6.1] makes  $A^*(X)$  into a commutative, graded ring.

For general schemes [Ful, Chapter 17] defines a graded operational Chow cohomology ring  $A_{\text{op}}^*(X) := \bigoplus_{k \geq 0} A_{\text{op}}^k(X)$ : an element  $c \in A_{\text{op}}^k(X)$  is a collection of homomorphisms of groups:

$$c_g^{(k)} : A_p X' \rightarrow A_{p-k} X'$$

for every morphism  $g : X' \rightarrow X$  which are compatible with respect to proper pushforward, and pullbacks along flat morphisms and regular embeddings (see [Ful, Definitions 17.1 and 17.3]). The product is given by composition and turns  $A_{\text{op}}^*(X)$  into a graded ring called the **Chow cohomology ring** of  $X$ . Moreover, if  $X$  has a resolution of singularities (e.g. if the characteristic of the ground field is zero or if  $X$  is a toric variety) then  $A_{\text{op}}^*(X)$  is known to be commutative. If  $X$  is smooth, then [Ful, Corollary 17.4] proves that the Poincaré duality

map  $A_{\text{op}}^k(X) \rightarrow A^k(X) = A_{n-k}(X)$  is an isomorphism of rings where the intersection product agrees with the product given by composition.

**Remark 2.1.** In [Ful, Chapter 17] the Chow cohomology ring is also denoted  $A^*(X)$  without the inclusion of the subscript 'op'.

**Proposition 2.2.** Chow cohomology is an  $\mathbb{A}^1$ -homotopy invariant functor.

*Proof.* Consider the pullback  $\pi^*: A_{\text{op}}^*(X) \rightarrow A_{\text{op}}^*(X \times \mathbb{A}^1)$  where  $\pi: X \times \mathbb{A}^1 \rightarrow X$  is the projection. First note that injectivity of  $\pi^*$  is a formal consequence of the functoriality; the composition  $X \xrightarrow{\iota} X \times \mathbb{A}^1 \xrightarrow{\pi} X$  is the identity, where  $\iota(x) = (x, 0)$ .

Suppose that  $c \in A_{\text{op}}^k(X \times \mathbb{A}^1)$ . We wish to show that  $c = \pi^*d$  for some  $d \in A_{\text{op}}^k(X)$ . Given a morphism  $Y \xrightarrow{f} X$  let  $g = f \times \text{id}: Y \times \mathbb{A}^1 \rightarrow X \times \mathbb{A}^1$ . Since the flat pullback  $\pi^*: A_*(Y) \rightarrow A_*(Y \times \mathbb{A}^1)$  is an isomorphism and this isomorphism is compatible with other operations on Chow homology we can define a class  $d \in A_{\text{op}}^*(X)$  such that  $\pi^*d = c$  by the formula  $d_f(\alpha) = (\pi^*)^{-1}c_g(\pi^*\alpha)$ .  $\square$

2.1.1. *Equivariant Chow cohomology.* An equivariant version of operational Chow cohomology was defined in [EG]. An element  $c \in A_{\text{op},G}^k(X)$  is a collection operations on equivariant Chow groups  $c_f: A_*^G(X') \rightarrow A_{*-k}^G(X')$  for every equivariant morphism  $X' \xrightarrow{f} X$  compatible with equivariant proper pushforward and equivariant flat maps and equivariant regular embeddings maps. [EG, Corollary 2] states that if  $X$  admits a resolution of singularities then  $A_{\text{op},G}^k(X)$  can be identified with the operational Chow group  $A_{\text{op}}^k(X_G)$  where  $X_G$  is an algebraic space of the form  $X \times^G U$ . Here  $U$  is an open set in a representation  $V$  of  $G$  on which  $G$  acts freely and  $\text{codim}(V \setminus U) > k$ . It follows from this identification that the equivariant Chow cohomology groups  $A_{\text{op},G}^k(X)$  enjoy all of the formal properties of ordinary operational Chow cohomology. In particular, the functor  $A_{\text{op},G}^*$  is a homotopy invariant functor on the category of schemes or algebraic spaces with a  $G$ -action.

2.2. **Operational  $K$ -theory.** Following [AP], if  $X$  is a scheme we denote by  $K_0(X)$  the Grothendieck group of coherent sheaves, and  $K^0(X)$  the Grothendieck group of perfect complexes. If  $X$  has an ample family of line bundles, then  $K^0(X)$  is the same as the naive Grothendieck group of vector bundles.

For any scheme  $X$ , Anderson and Payne define the **operational  $K$ -theory**  $\text{op} K^0(X)$  of  $X$  as follows. An element  $c \in \text{op} K^0(X)$  is a collection of operators  $c_f: K_0(X') \rightarrow K_0(X')$  indexed by morphisms  $X' \xrightarrow{f} X$  compatible with proper pushforward, flat pullback and pullback along regular embeddings.

For any scheme  $X$ , there is a canonical map  $\text{op} K^0(X) \rightarrow K_0(X)$  given by  $c \mapsto c_{\text{id}_X}(\mathcal{O}_X)$ . If  $X$  is smooth, then [AP, Corollary 4.5] states that this map is an isomorphism.

**Theorem 2.3.** [AP, Theorem 1.1]  $\text{op} K^0$  is a homotopy invariant functor.

2.2.1. *Equivariant operational  $K$ -theory.* Anderson and Payne also define the equivariant operational  $K$ -theory ring as the ring of operations on the equivariant Grothendieck group of coherent sheaves,  $K_0^G(X)$ . Since the equivariant Grothendieck group is an  $\mathbb{A}^1$ -homotopy invariant the proof of [AP, Theorem 1.1] goes through and we conclude that  $\text{op} K_G^0$  is a homotopy invariant functor. If  $G = T$  is a torus and  $X$  is smooth, then Anderson and Payne also prove that  $\text{op} K_T^0(X)$  can be identified with  $K_0^T(X)$ .

## 3. THE CONE THEOREM

Fix a base scheme  $X$  of finite type defined over a field  $k$ .

**Definition 3.1.** An  $X$ -cone is a scheme of the form  $C = \text{Spec}_X S$  where  $S = \bigoplus_{n=0}^{\infty} S_n$  is a finitely generated graded  $\mathcal{O}_X$ -algebra such that  $S_0 = \mathcal{O}_X$ . (Note that we do not require that  $S$  be locally generated in degree one.)

More generally, if  $G$  is an algebraic group and  $X$  is a  $G$ -scheme then we say that  $C = \text{Spec } S$  is a  $G$ -cone if the  $S_n$  are sheaves of  $G$ - $\mathcal{O}_X$  modules and multiplication of local sections is  $G$ -equivariant.

The inclusion  $S_0 \rightarrow S$  defines a projection  $\rho: C \rightarrow X$  and the identification of  $S_0 = S/S^+$  defines an inclusion  $\iota: X \rightarrow C$ . Clearly,  $\rho \circ \iota = \text{id}_X$ .

The key property of homotopy invariant functors is the following cone theorem.

**Theorem 3.1.** [Wei, cf. Exercise IV.11.5] The pullbacks  $\rho^*$  and  $\iota^*$  are inverses. In particular  $H(X) = H(C)$ . Likewise if  $H^G$  is a  $G$ -homotopy invariant functor and  $C = \text{Spec } S$  is a  $G$ -cone then  $H^G(X) = H^G(Y)$ .

*Proof.* We give the proof in the non-equivariant case as the proof in the equivariant case is identical.

Since  $\rho \circ \iota = \text{id}_X$ , we know that  $\iota^* \circ \rho^*: X \rightarrow X$  is the identity. In particular,  $\rho^*$  is injective. Thus it suffices to prove that  $\rho^* \circ \iota^*: C \rightarrow C$  is an isomorphism.

Since  $C$  is a cone over  $X = \text{Spec } S_0$ , the map of graded rings  $S \rightarrow S[t]$ , sending  $S_n$  to  $t^n S_n$ , defines an  $\mathbb{A}^1$  action  $\sigma: C \times \mathbb{A}^1 \rightarrow C$  with fixed scheme  $X = \text{Spec } S_0$ .

Let  $s_t: C \rightarrow C$  be the map  $x \mapsto tx$ . For  $t \neq 0$ ,  $s_t$  is an isomorphism with inverse  $s_{t^{-1}}$  and  $s_0: C \rightarrow C$  is the composition  $\iota \circ \rho$ . The map  $s_t$  is itself a composition

$$C \xrightarrow{i_t} C \times \mathbb{A}^1 \xrightarrow{\sigma} C$$

where  $i_t: X \hookrightarrow X \times \mathbb{A}^1$  is the inclusion  $x \mapsto (x, t)$ .

Now if  $\pi: X \times \mathbb{A}^1 \rightarrow X$  is the projection, then for any  $t$ ,  $\pi \circ i_t = \text{id}$ . Since  $\pi^*$  is assumed to be an isomorphism,  $i_t^*$  must also be an isomorphism for any  $t$ . Since, for  $t \neq 0$  the composite  $s_t = \sigma \circ i_t$  is an isomorphism we see that  $\sigma^*$  must also be an isomorphism. Hence,  $s_0^* = (\sigma \circ i_0)^*$  is an isomorphism. But  $s_0 = (\iota \circ \rho)$ , so  $(\iota \circ \rho)^*$  is an isomorphism as claimed.  $\square$

**Example 3.2.** Let  $X \subset Y$  be a closed subscheme and let  $C_X Y$  be the normal cone of  $X$  in  $Y$ . Theorem 3.1 implies that the pullback  $A_{\text{op}}^*(X) \rightarrow A_{\text{op}}^*(C_X Y)$  is an isomorphism. In particular if  $X$  is smooth then  $A_{\text{op}}^*(C_X Y)$  is identified with the Chow ring of  $X$ . If the closed embedding  $X \hookrightarrow Y$  is a regular embedding (for example if  $X$  and  $Y$  are both smooth) then  $C_X Y$  is the normal bundle to  $X$  in  $Y$  and this identification follows from the usual homotopy invariance of operational Chow rings.

**3.1. Cone theorem for bivariant groups.** The operational Chow and  $K$ -theory rings defined by Fulton–MacPherson and Anderson–Payne are part of a more general construction associated to the covariant (for proper morphisms) functors  $K_0$  and  $A_*$ . Given a morphism of schemes  $Y \rightarrow X$  the bivariant Chow group  $A_{\text{op}}^k(Y \rightarrow X)$  is the graded abelian group consisting of a collection of operators  $c_f: A_*(Y') \rightarrow A_{*-k}^*(X')$  for each morphism  $X' \xrightarrow{f} X$  compatible with proper pushforward and flat pullbacks and pullback and pullback along regular embeddings, where  $Y' = Y \times_X X'$ . The group  $\text{op } K^0(Y \rightarrow X)$  is defined analogously

but the compatibility is with proper pushforward, flat pullback and pullback along regular embeddings.

The groups  $A_{\text{op}}^*(Y \rightarrow X)$  and  $\text{op } K^0(Y \rightarrow X)$  are contravariant functors on the category whose objects are morphisms of schemes  $Y \rightarrow X$  and whose morphisms are cartesian dia-

$$\begin{array}{ccc} Y' & \rightarrow & X' \\ \text{grams } \downarrow & & \downarrow \\ Y & \rightarrow & X \end{array}$$

$$Y \times \mathbb{A}^1 \rightarrow X \times \mathbb{A}^1$$

It is easy to show that pullback along the diagram

$$\begin{array}{ccc} & \downarrow & \downarrow \\ & Y & \rightarrow X \end{array}$$

induces isomorphisms  $A_{\text{op}}^*(Y \rightarrow X) \rightarrow A_{\text{op}}^*(Y \times \mathbb{A}^1 \rightarrow X \times \mathbb{A}^1)$ ,  $\text{op } K^0(Y \rightarrow X) \rightarrow \text{op } K^0(Y \times \mathbb{A}^1 \rightarrow X \times \mathbb{A}^1)$ . As a corollary we obtain a cone isomorphism theorem for these bivariant groups.

**Corollary 3.3.** Given a morphism,  $Y \rightarrow X$  and cone  $C \rightarrow X$  let  $C_Y \rightarrow Y$  be the cone obtained by base change. Then the pullbacks  $A_{\text{op}}^*(Y \rightarrow X) \rightarrow A_{\text{op}}^*(C_Y \rightarrow C)$  and  $\text{op } K^0(Y \rightarrow X) \rightarrow \text{op } K^0(C_Y \rightarrow C)$  are isomorphisms.

#### 4. AFFINE TORIC VARIETIES

**Theorem 4.1.** If  $X = X(\sigma)$  is an affine toric variety defined by a strongly convex rational cone  $\sigma$  in a lattice  $N$ , then for any homotopy invariant functor  $H$  on  $\text{Sch}/k$ ,  $H(X) = H(T_0)$  where  $T_0$  is an algebraic torus.

*Proof.* Since  $X(\sigma)$  is an affine toric variety the proof of [CLS, Proposition 3.3.9] shows that we can decompose  $X = X(\bar{\sigma}) \times T_0$  where  $T_0$  is a torus and  $\bar{\sigma}$  is a full dimensional cone. Since  $\bar{\sigma}$  is full-dimensional the semi-group  $S_{\bar{\sigma}}$  is generated in positive degree, so  $R = k[X(\bar{\sigma})]$  is a positively graded ring with  $R_0 = k$ .

Hence  $S = k[X(\sigma)] = k[T_0] \otimes_k R$  is a positively graded ring with  $S_0 = k[T_0]$ . Hence by the cone theorem,  $H(X) = H(T_0)$   $\square$

**Corollary 4.2.** If  $X$  is an affine toric variety then  $A^0(X) = \mathbb{Z}$  and  $A_{\text{op}}^k(X) = 0$  for  $k > 0$ . Likewise,  $\text{op } K^0(X) = \mathbb{Z}$ .

*Proof.* By the theorem we know that  $A_{\text{op}}^*(X) = A_{\text{op}}^*(T_0)$  and  $\text{op } K^0(X) = \text{op } K^0(T_0)$ . Since a torus is an open subset of  $\mathbb{A}^n$ ,  $A_{\text{op}}^k(T_0) = A^k(T_0) = 0$  if  $k > 0$  and  $A^0(X) = \mathbb{Z}$  for any  $X$ . Likewise,  $\text{op } K^0(X) = \text{op } K^0(T_0) = K_0(T_0) = \mathbb{Z}$ .  $\square$

**Example 4.3.** When  $X$  is a complete toric variety then we know that  $A_{\text{op}}^k(X) = \text{Hom}(A_k(X), \mathbb{Z})$ . However, for non-complete toric variety this result fails. For example if  $X = \mathbb{A}^1$  then  $A_0(X) = 0$  but  $A^0(X) = \mathbb{Z}$ . Another example is to let  $\sigma$  denote the cone generated by  $\{(1, 0, 1), (0, -1, 1), (-1, 0, 1), (0, 1, 1)\}$  in  $\mathbb{R}^3$ . One can compute that  $A_2(X(\sigma)) = \mathbb{Z}/2 \oplus \mathbb{Z}$ . Thus,  $\text{Hom}(A_2(X(\sigma)), \mathbb{Z}) = \mathbb{Z}$  but by Corollary 4.2,  $A_{\text{op}}^2(X(\sigma)) = 0$ .

**Example 4.4.** In the equivariant case we have an analogous result for the  $T$ -equivariant operational Chow ring and  $K$ -theory. To simplify the notation we assume that the cone  $\sigma$  is full dimensional.

**Corollary 4.5.** If  $X$  is an affine toric variety associated to a full dimensional cone  $\sigma$  then  $A_{\text{op},T}^*(X) = \text{Sym}(X(T))$  and  $\text{op}K_T^0(X) = R(T)$ . Here  $\text{Sym}(X(T))$  is the polynomial algebra generated by the character group of  $T$  and  $R(T)$  is the representation ring of  $T$ .

*Proof.* In this case  $S_0 = \text{Spec} k$ , so  $A_{\text{op},T}^*(X) = A_T^*(\text{pt}) = \text{Sym}(X(T))$  by [EG, Section 3.2] and  $\text{op}K_T^0(X) = K_T^0(\text{pt}) = R(T)$ .  $\square$

**4.1. An alternative proof of the vanishing of Chow cohomology and operational  $K$ -theory on affine toric varieties.** In [Ric, ER1] a more involved proof that  $A_{\text{op}}^*(X) = \text{op}K^0(X) = \mathbb{Z}$  when  $X$  is an affine toric variety is given. The proof of both of these statements rests on the fact that both  $A_{\text{op}}^*(X)$  and  $\text{op}K^0(X)$  satisfy the following descent property for proper surjective morphisms.

If  $X' \rightarrow X$  is a proper surjective morphism and if  $H$  denotes either functor  $A_{\text{op}}^*$  or  $\text{op}K^0$  then the sequence

$$0 \rightarrow H(X) \otimes \mathbb{Q} \rightarrow H(X') \otimes \mathbb{Q} \xrightarrow{p_1^* - p_2^*} H(X' \times_X X') \otimes \mathbb{Q}$$

is exact where  $p_1, p_2$  are the two projections  $X' \times_X X' \rightarrow X'$ .

This descent property does not hold for arbitrary homotopy invariant functors – for example it need not hold for the functor  $\text{op}K_G^0$  when  $G$  is not a torus. However when the descent property holds for a functor  $H$ , it can be used as a tool to calculate  $H$  on singular schemes [Kim, AP].

## 5. OPERATIONAL CHOW RINGS OF GOOD MODULI SPACES

The goal of this section is to explain how the cone theorem for homotopy invariant functors can shed light on questions about the structure of the operational Chow ring for quotients of smooth varieties and, more generally, good moduli spaces of smooth Artin stacks.

**5.1. Strong cycles on good moduli spaces of Artin stacks.** Let  $G$  be a linearly algebraic group acting on a scheme  $X$ . We say that a scheme  $Y$  equipped with a  $G$ -invariant morphism  $p: X \rightarrow Y$  is a *good quotient* if  $p$  is affine and  $(p_*\mathcal{O}_X)^G = \mathcal{O}_Y$ . The basic example is the quotient  $X^{ss} \rightarrow X^{ss}/G$  where  $X^{ss}$  is the set of semi-stable points (with respect to a choice of linearization) for the action of a linearly reductive group on a projective variety  $X$ . This definition was extended to Artin stacks by Alper.

**Definition 5.1** ([Alp, Definition 4.1]). Let  $\mathcal{X}$  be an Artin stack and let  $X$  be an algebraic space. We say that  $X$  is a *good moduli space of  $\mathcal{X}$*  if there is a morphism  $\pi: \mathcal{X} \rightarrow X$  such that

- (1)  $\pi$  is *cohomologically affine* meaning that the pushforward functor  $\pi_*$  on the category of quasi-coherent  $\mathcal{O}_{\mathcal{X}}$ -modules is exact.
- (2)  $\pi$  is *Stein* meaning that the natural map  $\mathcal{O}_X \rightarrow \pi_*\mathcal{O}_{\mathcal{X}}$  is an isomorphism.

**Remark 5.1.** If  $\mathcal{X} = [Z/G]$  where  $G$  is a linearly reductive algebraic group then the statement that  $X$  is a good moduli space for  $\mathcal{X}$  is equivalent to the statement that  $X$  is the good quotient of  $Z$  by  $G$ .

**Definition 5.2** ([ER2]). Let  $\mathcal{X}$  be an Artin stack with good moduli space  $X$  and let  $\pi: \mathcal{X} \rightarrow X$  be the good moduli space morphism. We say that a closed point  $x$  of  $\mathcal{X}$  is *stable* if  $\pi^{-1}(\pi(x)) = x$  under the induced map of topological spaces  $|\mathcal{X}| \rightarrow |X|$ . A closed point  $x$  of  $\mathcal{X}$  is *properly stable* if it is stable and the stabilizer of  $x$  is finite.

We say  $\mathcal{X}$  is stable (resp. properly stable) if there is a good moduli space  $\pi: \mathcal{X} \rightarrow X$  and the set of stable (resp. properly stable) points is non-empty. Likewise we say that  $\pi$  is a stable (resp. properly stable) good moduli space morphism.

**Remark 5.2.** Again this definition is modeled on GIT. If  $G$  is a linearly reductive group and  $X^{ss}$  is the set of semistable points for a linearization of the action of  $G$  on a projective variety  $X$  then a (properly) stable point of  $[X^{ss}/G]$  corresponds to a (properly) stable orbit in the sense of GIT. The stack  $[X^{ss}/G]$  is stable if and only if  $X^s \neq \emptyset$ . Likewise  $[X^{ss}/G]$  is properly stable if and only if  $X^{ps} \neq \emptyset$ . As is the case for GIT quotients, the set of stable (resp. properly stable points) is open [ER2].

**Definition 5.3.** [ES1, ES2] Let  $\mathcal{X}$  be an irreducible Artin stack with stable good moduli space  $\pi: \mathcal{X} \rightarrow X$ . A closed integral substack  $\mathcal{Z} \subseteq \mathcal{X}$  is *strong* if  $\text{codim}_{\mathcal{X}} \mathcal{Z} = \text{codim}_X \pi(\mathcal{Z})$  and  $\mathcal{Z}$  is saturated with respect to  $\pi$ , i.e.  $\pi^{-1}(\pi(\mathcal{Z})) = \mathcal{Z}$  as stacks. We say  $\mathcal{Z}$  is *topologically strong* if  $\text{codim}_{\mathcal{X}} \mathcal{Z} = \text{codim}_X \pi(\mathcal{Z})$  and  $\pi^{-1}(\pi(\mathcal{Z}))_{red} = \mathcal{Z}$ .

Let  $A_{\text{tst}}^*(\mathcal{X}/X)$  be the subgroup of  $A^*(\mathcal{X})$  generated by topologically strong cycles and  $A_{\text{st}}^*(\mathcal{X}/X)$  be the subgroup generated by by strong cycles. (Here the Chow group  $A^*(\mathcal{X})$  is the Chow group defined by Kresch [Kre]. When  $\mathcal{X} = [Z/G]$  it can be identified with the equivariant Chow group  $A_G^*(Z)$  of [EG].)

The main result of [ES2] is the following theorem which says that if  $X$  is smooth then any operational class  $c \in A^k(X)$  can be represented by a codimension  $k$ -cycle on the stack  $\mathcal{X}$  which is saturated with respect to the good moduli space morphism  $\mathcal{X} \rightarrow X$ .

**Theorem 5.3.** [ES2, Theorem 1.1] Let  $\mathcal{X}$  be a properly stable smooth Artin stack with good moduli space  $\mathcal{X} \xrightarrow{\pi} X$ . Then there is a pullback  $\pi: A_{\text{op}}^*(X)_{\mathbb{Q}} \rightarrow A^*(\mathcal{X})_{\mathbb{Q}}$  which is injective and factors through the subgroup  $A_{\text{tst}}^*(\mathcal{X})_{\mathbb{Q}}$ .

[ES2, Example 3.24] shows that not every topologically strong cycle is in the image of  $A_{\text{op}}^*(X)$ . However, [ES2, Theorem 1.7c] states that any strong lci cycle on  $\mathcal{X}$  is in the image of  $A_{\text{op}}^*(X)_{\mathbb{Q}}$ . (A cycle  $\sum_i a_i [\mathcal{Z}_i]$  is lci if the  $\mathcal{Z}_i$  are closed substacks of  $\mathcal{X}$  such that the inclusion  $\mathcal{Z}_i \hookrightarrow \mathcal{X}$  is an lci morphism.) This leads to a number of successively weaker questions about the operational Chow rings of good moduli spaces of smooth Artin stacks. They can be viewed as analogues for quotients of smooth varieties by reductive groups of Conjectures 2 and 3 of [MV].

**Question 5.4.** Is the image of  $A_{\text{op}}^*(X)_{\mathbb{Q}}$  contained in the subgroup of  $A^*(\mathcal{X})_{\mathbb{Q}}$  generated by strong lci cycles?

**Question 5.5.** Is the image of  $A_{\text{op}}^*(X)_{\mathbb{Q}}$  equal to the subring of  $A^*(\mathcal{X})_{\mathbb{Q}}$  generated by strong lci cycles?

**Question 5.6.** Is  $A_{\text{op}}^*(X)_{\mathbb{Q}}$  generated by Chern classes of perfect complexes on  $X$ ?

**Remark 5.7.** Note that Question 5.6 is an analogue of the question raised by Anderson and Payne about the surjectivity of the map  $K^0(X) \rightarrow \text{op } K^0(X)$  where  $K^0(X)$  is the Grothendieck group of perfect complexes. Anderson and Payne prove that for 3-dimensional complete toric varieties this map is in fact surjective. By comparison in [ES2] the authors prove that  $A^*(X)_{\mathbb{Q}} = A_{\text{st}}^*(\mathcal{X}/X)_{\mathbb{Q}}$  and in the case of 3-dimensional toric varieties [Ric] shows that  $A^*(X)_{\mathbb{Q}}$  is generated by strong lci cycles.

Given the relation between the operational Chow ring and strong cycles leads to the following additional question.

**Question 5.8.** Is  $A_{\text{st}}^*(\mathcal{X}/X)$  (resp.  $A_{\text{Lst}}^*(\mathcal{X}/X)$ ) a subring of  $A^*(\mathcal{X})$ ; i.e., is the product of strong (resp. topologically strong) cycles strong?

**5.2. The cone theorem and local models for good moduli.** The étale slice theorem of Alper, Hall and Rydh [AHR] states if  $\mathcal{X}$  is a smooth stack then at a closed point  $x$  of  $\mathcal{X}$  the good moduli morphism  $\mathcal{X} \rightarrow X$  is étale locally isomorphic to the quotient  $[V/G_x] \rightarrow \text{Spec } k[V]^{G_x}$  where  $G_x$  is the inertia group of  $x$  in  $\mathcal{X}$  and  $V$  is a representation of  $G_x$ . Thus we may view stacks of the form  $[V/G]$  with their good moduli spaces  $\text{Spec } k[V]^G$  as the “affine models” of smooth stacks with good moduli spaces.

The cone theorem has the following corollary.

**Corollary 5.9.** Let  $V$  be a representation of a reductive group  $G$  and let  $X = \text{Spec } k[V]^G$  be the quotient. Then for any homotopy invariant functor  $H(X) = H(\text{Spec } k)$ . In particular,  $A_{\text{op}}^*(X) = \mathbb{Z}$  and  $\text{op } K^0(X) = \mathbb{Z}$ .

*Proof.* Since  $G$  acts linearly on  $V$  the action of  $G$  on  $k[V]$  preserves the natural grading. Hence the invariant ring  $k[V]^G$  is also graded. Thus by Theorem 3.1  $H(X) = H(\text{Spec } k)$  since  $k$  is the 0-th graded piece of  $k[V]^G$ .  $\square$

Combining Corollary 5.9 with [ES2, Theorem 1.7c] yields the following result.

**Corollary 5.10.** If  $\mathcal{Z} \subset \mathcal{X} = [V/G]$  is a proper closed substack which is strongly regularly embedded then  $[\mathcal{Z}]$  is torsion in  $A^*(\mathcal{X}) = A_G^*(\text{pt})$ . In particular if  $A_G^*(\text{pt})$  is torsion free (for example if  $G$  is torus or  $\text{GL}_n$ ) then  $[\mathcal{Z}] = 0$ .

Note that any integral strong divisor  $\mathcal{D}$  on  $\mathcal{X} = [V/G]$  is necessarily defined by a single  $G$ -invariant equation. In this case  $\mathcal{O}(\mathcal{D})$  is an equivariantly trivial line bundle so  $[\mathcal{D}] = c_1(\mathcal{O}(\mathcal{D})) = 0$  in the equivariant Chow ring  $A_G^*(V)$ . More generally, if an integral substack  $\mathcal{Z} \subset [V/G]$  is a complete intersection of strong divisors, then its class in  $A_G^*(V)$  is also 0. In particular it shows that in the “local case” the image of  $A_{\text{op}}^*(X)$  is contained in the subgroup generated by strong global complete intersections.

This leads to the following question.

**Question 5.11.** Are there examples of strong integral substacks  $\mathcal{Z} \subset [V/G]$  such that  $[\mathcal{Z}] \neq 0$  in  $A^*([V/G])_{\mathbb{Q}}$ ?

**5.3. Acknowledgments.** The authors are grateful to Sam Payne and Angelo Vistoli for helpful comments.

## REFERENCES

- [Alp] Jarod Alper, *Good moduli spaces for Artin stacks*, Ann. Inst. Fourier (Grenoble) **63** (2013), no. 6, 2349–2402.
- [AHR] Jarod Alper, Jack Hall, and David Rydh, *A Luna étale slice theorem for algebraic stacks*, arXiv:1504.06467.
- [AP] Dave Anderson and Sam Payne, *Operational K-theory*, Doc. Math. **20** (2015), 357–399.
- [CLS] David A. Cox, John B. Little, and Henry K. Schenck, *Toric varieties*, Graduate Studies in Mathematics, vol. 124, American Mathematical Society, Providence, RI, 2011.
- [EG] Dan Edidin and William Graham, *Equivariant intersection theory*, Invent. Math. **131** (1998), no. 3, 595–634.



- [ER1] Dan Edidin and Ryan Richey, *The Chow cohomology of affine toric varieties*, arXiv:1904.08476.
- [ER2] Dan Edidin and David Rydh, *Canonical reduction of stabilizers for Artin stacks with good moduli spaces*, arXiv:1710.03220.
- [ES1] Dan Edidin and Matthew Satriano, *Strong cycles and intersection products on good moduli spaces, K-Theory—Proceedings of the International Colloquium, Mumbai, 2016*, Hindustan Book Agency, New Delhi, 2018, pp. 223–238.
- [ES2] ———, *Towards an intersection Chow cohomology on GIT quotients*, arXiv:1707.05890, to appear in *Transform. Groups*.
- [Ful] William Fulton, *Intersection theory*, Springer-Verlag, Berlin, 1984.
- [FM] William Fulton and Robert MacPherson, *Categorical framework for the study of singular spaces*, *Mem. Amer. Math. Soc.* **31** (1981), no. 243, vi+165.
- [FS] William Fulton and Bernd Sturmfels, *Intersection theory on toric varieties*, *Topology* **36** (1997), no. 2, 335–353.
- [Kim] Shun-ichi Kimura, *Fractional intersection and bivariant theory*, *Comm. Algebra* **20** (1992), no. 1, 285–302.
- [Kre] Andrew Kresch, *Cycle groups for Artin stacks*, *Invent. Math.* **138** (1999), no. 3, 495–536.
- [Lun] Domingo Luna, *Slices étales*, *Sur les groupes algébriques*, *Soc. Math. France, Paris, 1973*, pp. 81–105. *Bull. Soc. Math. France, Paris, Mémoire* 33.
- [MV] Marco Maggesi and Gabriele Vezzosi, *Some elementary remarks on lci algebraic cycles*, arXiv:1612.04570.
- [Pay] Sam Payne, *Equivariant Chow cohomology of toric varieties*, *Math. Res. Lett.* **13** (2006), no. 1, 29–41.
- [Ric] Ryan Richey, Ph.D. thesis, University of Missouri, May 2019.
- [Tot] Burt Totaro, *Chow groups, Chow cohomology, and linear varieties*, *Forum Math. Sigma* **2** (2014), e17, 25pages.
- [Wei] Charles A. Weibel, *The K-book*, *Graduate Studies in Mathematics*, vol. 145, American Mathematical Society, Providence, RI, 2013, An introduction to algebraic K-theory.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA MO 65211

*E-mail address:* edidind@missouri.edu

*E-mail address:* rrtf3@mail.missouri.edu