Mass prediction for the last discovered member of the axial-vector nonet with quantum numbers $\mathbf{J}^{PC} = 1^{+-*}$

M.V. Chizhov¹ and M.N. Naydenov¹

¹Department of Atomic Physics, Faculty of Physics, Sofia University, 1164 Sofia, Bulgaria

In this paper we have shown that the novel mass relation among the vector states, $\phi(1020)$ and $\phi(1680)$, with quantum numbers $J^{PC} = 1^{--}$ and axial-vector strangeonium state $h_1(s\bar{s})$ with quantum numbers 1^{+-} is also valid for nonzero current quark mass. This relation predicts the mass of recently discovered $h_1(s\bar{s})$ state by the Collaboration BESIII within experimental accuracy.

PACS numbers: 11.30.Qc, 12.39.-x, 12.40.Yx

Keywords: Nambu-Jona-Lasinio model, mass relation

^{*}Corresponding author: mih@phys.uni-sofia.bg

I. INTRODUCTION

It is known that there exist the two lowest-lying axial-vector nonets of meson states with quantum numbers for their neutral components $J^{PC} = 1^{++}$ and 1^{+-} . The first nonet, "A", includes isotriplet of a_1 mesons, two isodoublets of K_{1A} strange mesons and two isosinglet states f_1 . The second nonet, "B", consists of isotriplet of b_1 mesons¹, two isodoublets of K_{1B} strange mesons and two isosinglet states h_1 .

The two isosinglets of spin-1 states have nearly pure $u\bar{u} + d\bar{d}$ and $s\bar{s}$ structures. There is still discussion about identification of the two f_1 states among three observed resonances $f_1(1285)$, $f_1(1420)$ and $f_1(1510)$. While identification of the last from two $h_1(1170)$ and $h_1(1415)$ states has been established in last two years. The Collaboration BESIII discovered axial-vector strangeonium state $h_1(s\bar{s})$ [1, 2] and confirmed its previous observations by two collaborations LASS [3] and Crystal Barrel [4].

There is interesting situation with the measured and the predicted mass of the axialvector strangeonium state with quantum numbers 1^{+-} . For the first time this state has been observed by the Collaboration LASS in 1988 with mass $m_{h_1(s\bar{s})}^{\text{LASS}} = (1380\pm20)$ MeV. Therefore, this state got the name $h_1(1380)$. The latest measurements by the Collaboration BESIII $m_{h_1(s\bar{s})}^{\text{BESIII}} = (1423.2\pm2.1\pm7.3)$ MeV and PDG average $m_{h_1(s\bar{s})}^{\text{PDG}} = (1416\pm8)$ MeV [5] required changing the name of this resonance to $h_1(1415)$ in the last year. However, there was no accurate prediction of the mass of this state among cited theoretical papers in the first version of BESIII publication [1]. The precise prediction of $h_1(s\bar{s})$ mass $m_{h_1(s\bar{s})}^{\text{theor}} = (1415\pm13)$ MeV has been given in 2003 in the paper [6].

The detail consideration of this prediction and new investigation will be presented below.

II. THE MODEL

Explanation of the spontaneous chiral symmetry breaking [7] and the introduction of quarks [8, 9] give us a principal possibility to describe the whole variety of the light hadron states, in particular the quark-antiquark meson states. The most theoretically and experimentally well studied is the nonet of pseudoscalar mesons, which arise as pseudo-Goldstone

¹ The b_1 mesons consist of the light u and d quarks. Its name denotes the type of the nonet and has no relation to b quark.

bosons in result of spontaneous and explicit chiral symmetry breaking. The properties of pseudoscalar nonet can be accurately investigated using Nambu – Jona-Lasinio (NJL) model [10] or chiral perturbation theory [11]. Meanwhile, theoretical and experimental situation with identification and explaning properties of members of scalar nonet as quarkantiquark states is unsatisfactory.

In this paper we consider a model for spin-1 nonets in the framework of paper [6]. It is well known, that besides the two considered lowest-lying axial-vector nonets, a nonet of vector mesons exists. The latest consists of isotriplet ρ mesons, two isodoublets K^* mesons and two isosinglets ω and ϕ with quantum numbers 1⁻⁻, which have nearly pure $u\bar{u} + d\bar{d}$ and $s\bar{s}$ structures, correspondingly. So, there is an obvious *asymmetry* between the numbers of axial-vector and vector nonets.

In paper [6] in the framework of extended U(1) massless quark NJL model a new approach to restoring the symmetry has been suggested. In this way new mass relations among spin-1 mesons from different nonets have been derived, which are confirmed experimentally. The basic idea was in the consideration of all possible Lorentz invariant *local* Yukawa interactions between quark currents: $\overline{\psi}\psi$, $\overline{\psi}\gamma^5\psi$, $\overline{\psi}\gamma_\mu\psi$, $\overline{\psi}\gamma_\mu\gamma^5\psi$, $\overline{\psi}\sigma_{\mu\nu}\psi$ and the corresponding meson fields S, P, V_{μ} , A_{μ} , $T_{[\mu\nu]}^2$ with quantum numbers 0^{++} , 0^{-+} , 1^{--} , 1^{++} , $(1^{--}$, $1^{+-})$. The quark current $\overline{\psi}\sigma_{\mu\nu}\psi$ and the corresponding second-rank antisymmetric tensor field $T_{[\mu\nu]}$ possess two types of different quantum numbers: 1^{--} and 1^{+-} . On the mass-shell $T_{[\mu\nu]}$ can be decomposed into vector, R_{μ} , and axial-vector, B_{μ} , fields

$$T_{[\mu\nu]} = (\hat{\partial}_{\mu}R_{\nu} - \hat{\partial}_{\nu}R_{\mu}) - \frac{1}{2}\epsilon_{\mu\nu\alpha\beta}(\hat{\partial}^{\alpha}B^{\beta} - \hat{\partial}^{\beta}B^{\alpha})$$
(1)

with corresponding quantum numbers 1⁻⁻ and 1⁺⁻. Here we introduce the definition $\hat{\partial}_{\mu} = \partial_{\mu}/\sqrt{-\partial^2}$. Vice versa the R_{μ} and B_{μ} fields can be expressed through $T_{[\mu\nu]}$:

$$R_{\mu} = \hat{\partial}^{\nu} T_{[\mu\nu]}, \qquad B_{\mu} = \frac{1}{2} \epsilon_{\mu\nu\alpha\beta} \hat{\partial}^{\nu} T^{[\alpha\beta]}, \qquad (2)$$

which due to antisymmetry of $T_{\mu\nu}$ obey the obvious identities

$$\partial^{\mu}R_{\mu} \equiv 0, \qquad \partial^{\mu}B_{\mu} \equiv 0. \tag{3}$$

Therefore, the existing $h_1(s\bar{s})$ meson with quantum numbers 1^{+-} from axial-vector "B" nonet can be described by axial-vector field B_{μ} and requires the inclusion in the model the

² The notation $T_{[\mu\nu]} = \frac{1}{2}(T_{\mu\nu} - T_{\nu\mu})$ means antisymmetrization on Lorentz indices.

quark current $\overline{\psi}\sigma_{\mu\nu}\psi$ and the corresponding second-rank antisymmetric tensor field $T_{[\mu\nu]}$. Since the two vector fields V_{μ} and R_{μ} have the same quantum numbers 1⁻⁻ they could mix, which leads to the two physical states, ϕ and ϕ' for U(1) $s\bar{s}$ quark structure. So, extending this suggestion to U(3) model, it can be proposed that the lowest-lying vector nonet of physical states with quantum numbers 1⁻⁻ is produced from a mixing of vector and tensor nonets, similar to the two $K_1(1270)$ and $K_1(1400)$ physical states, which are superpositions of K_{1A} and K_{1B} states from the corresponding axial-vector nonets.

The new relation between the masses of $\phi(1020)$, $\phi' = \phi(1680)$ and $h_1(s\bar{s})$ physical states [6]

$$2m_{\phi}^2 - m_{\phi}m_{\phi'} + 2m_{\phi'}^2 = 3m_{h_1(s\bar{s})}^2,\tag{4}$$

which predicts the mass of undiscovered then $h_1(s\bar{s})$ state, has been obtained in approximation of zero current quark mass. However, it can be rather applied to $\omega(782)$, $\omega' = \omega(1650)$ and $h_1(1170)$ physical states, which consist of the light u and d quarks. Therefore, to confirm the relation (4), here we investigate the case of nonzero current quark mass m_0 .


In order to obtain relation (4) we consider only interactions of vector, V_{μ} , tensor, $T_{[\mu\nu]}$, and scalar, S, fields with the field of strange quark, ψ . Linearized NJL Lagrangian with auxiliary (without kinetic terms) boson fields, has the form

$$\mathcal{L}_{0} = \overline{\psi}(\not{q} - m_{0})\psi + g_{S}\overline{\psi}\psi S - \frac{\mu_{S}^{2}}{2}S^{2} + g_{V}\overline{\psi}\gamma^{\mu}\psi V_{\mu} + \frac{\mu_{V}^{2}}{2}V_{\mu}^{2} + \frac{g_{T}}{2}\overline{\psi}\sigma^{\mu\nu}\psi T_{[\mu\nu]} + \frac{\mu_{T}^{2}}{4}\left(4\,\hat{\partial}^{\mu}T_{[\lambda\mu]}\hat{\partial}_{\nu}T^{[\lambda\nu]} - T_{[\mu\nu]}T^{[\mu\nu]}\right), \qquad (5)$$

where μ_S^2 , μ_V^2 and μ_T^2 are bare mass terms³. Using relations (1,2,3) the last line in eq. (5) can be rewritten as

$$g_T \,\overline{\psi} \sigma^{\mu\nu} \psi \,\hat{\partial}_\mu R_\nu + i g_T \,\overline{\psi} \sigma^{\mu\nu} \gamma^5 \psi \,\hat{\partial}_\mu B_\nu + \frac{\mu_T^2}{2} \left(R_\mu^2 + B_\mu^2 \right). \tag{6}$$

The corresponding Feynman diagrams are presented in Fig. 1.



 $^{^{3}}$ The form of the mass term for the antisymmetric tensor field is suggested in [12].

Figure 1: Feynman diagrams for Yukawa interactions. Here $\hat{q}_{\mu} = q_{\mu}/\sqrt{q^2}$, where q_{μ} is incoming four-momentum of R_{μ} and B_{μ} states.

III. QUANTUM CORRECTIONS AND SYMMETRY BREAKING

Kinetic terms for the meson fields can be obtained through radiative quantum corrections. Let's calculate one-loop self-energy contributions for all the boson fields. For example, the contribution from the self-energy diagram of the scalar field (Fig. 2a) has the form




Figure 2: Quantum corrections to the self-energy part (a) and the self-interactions (b,c) of the scalar fields.

$$\Pi^{SS}(q) = ig_S^2 N_C \int \frac{\mathrm{d}^4 p}{(2\pi)^4} \mathrm{Tr} \left[(\not p - m_0)^{-1} (\not p - \not q - m_0)^{-1} \right]$$

= $4ig_S^2 N_C \int \frac{\mathrm{d}^4 p}{(2\pi)^4} \frac{1}{p^2 - m_0^2} - 2ig_S^2 N_C \int \frac{\mathrm{d}^4 p}{(2\pi)^4} \frac{q^2 - 4m_0^2}{(p^2 - m_0^2)^2} + \text{finite terms}$
= $4g_S^2 N_C I_2 - 8g_S^2 N_C m_0^2 I_0 + 2g_S^2 N_C I_0 q^2 + \text{finite terms},$ (7)

where

$$I_2 \equiv i \int \frac{\mathrm{d}^4 p}{(2\pi)^4} \frac{1}{p^2 - m_0^2} = \int \frac{\mathrm{d}^4 p_E}{(2\pi)^4} \frac{1}{p_E^2 + m_0^2} > 0 \tag{8}$$

is the quadratically divergent integral and

$$I_0 \equiv -i \int \frac{\mathrm{d}^4 p}{(2\pi)^4} \frac{1}{\left(p^2 - m_0^2\right)^2} = \int \frac{\mathrm{d}^4 p_E}{(2\pi)^4} \frac{1}{\left(p_E^2 + m_0^2\right)^2} > 0 \tag{9}$$

is the logarithmically divergent integral.

The first two terms in the last line of eq. (7) give contribution into the mass term of the scalar field $m_S^2 = \mu_S^2 + 4g_S^2 N_C (2m_0^2 I_0 - I_2)$. The third term indicates the appearance of the kinetic term. To normalize the scalar field wave function correctly, we must require fulfillment of the condition

$$2g_S^2 N_C I_0 = 1. (10)$$

Applying the similar procedure to all boson fields introduced in (5) and (6)

$$\Pi_{\mu\nu}^{VV} = \frac{4}{3} g_V^2 N_C I_0 (q_\mu q_\nu - q^2 g_{\mu\nu}) - 2g_V^2 N_C (m_0^2 I_0 + I_2) g_{\mu\nu} + \text{finite terms}, \qquad (11)$$

$$\Pi_{\mu\nu}^{VR} = 4g_V g_T N_C I_0 \frac{m_0}{\sqrt{q^2}} (q_\mu q_\nu - q^2 g_{\mu\nu}) + \text{finite terms},$$
(12)

$$\Pi_{\mu\nu}^{RR} = \frac{2}{3} g_T^2 N_C I_0 (q_\mu q_\nu - q^2 g_{\mu\nu}) + 4 g_T^2 N_C I_0 \frac{m_0^2}{q^2} (q_\mu q_\nu - q^2 g_{\mu\nu}) + \text{finite terms}, \quad (13)$$

$$\Pi^{BB}_{\mu\nu} = \frac{2}{3} g_T^2 N_C I_0 (q_\mu q_\nu - q^2 g_{\mu\nu}) - 4 g_T^2 N_C I_0 \frac{m_0^2}{q^2} (q_\mu q_\nu - q^2 g_{\mu\nu}) + \text{finite terms}, \quad (14)$$

the following useful relations among various Yukawa coupling constants from normalization of kinetic terms are obtained:

$$3g_S^2 = 2g_V^2 = g_T^2 = \frac{3}{2N_C I_0}.$$
(15)

In other words, due to the dynamic origin of the kinetic terms, all interactions in the model are described by only one coupling constant, for example, $g = g_S$.

The self-interactions in Fig. 2(b,c) lead to the following effective potential for the scalar field

$$V_{\rm eff} = \frac{m_S^2}{2}S^2 - 2gm_0S^3 + \frac{g^2}{2}S^4.$$
 (16)

The extremum condition

$$\left. \frac{\mathrm{d}V_{\mathrm{eff}}}{\mathrm{d}S} \right|_{S=\langle S \rangle} = m_S^2 \langle S \rangle - 6gm_0 \langle S \rangle^2 + 2g^2 \langle S \rangle^3 = 0 \tag{17}$$

at negative m_S^2 leads always to the absolute minimum of the effective potential with nontrivial solution $g\langle S \rangle = (3m_0 - \sqrt{9m_0^2 - 2m_S^2})/2 < 0$. It corresponds to the spontaneous symmetry breaking, which provides the strange quark with positive constituent mass

$$m_s = m_0 - g\langle S \rangle > 0. \tag{18}$$

It also leads to additional contributions to mass terms and mixing between vector bosons (Fig. 3).

There are no additional contribution to the mass of the vector field V_{μ} , $m_V^2 = \mu_V^2 - 2g_V^2 N_C (m_0^2 I_0 + I_2)$, while its mixing (12) with the vector field R_{μ} gets such contribution after symmetry breaking

$$\Delta \Pi^{VR}_{\mu\nu} = -4g_V g_T N_C I_0 \frac{g\langle S \rangle}{\sqrt{q^2}} (q_\mu q_\nu - q^2 g_{\mu\nu}) + \text{finite terms}$$
(19)




Figure 3: Quantum contributions to the mass terms and mixing between vector bosons after spontaneous symmetry breaking. Here the double lines correspond simultaneously to any of the R_{μ} or B_{μ} bosons.

in such a way that it depends only on physical constituent mass of the strange quark:

$$\Pi_{\mu\nu}^{VR} + \Delta \Pi_{\mu\nu}^{VR} = \sqrt{\frac{18m_s^2}{q^2}} \left(q_\mu q_\nu - q^2 g_{\mu\nu} \right) + \text{finite terms.}$$
(20)

In its turn R_{μ} and B_{μ} fields get additional contributions

$$\Delta \Pi^{RR}_{\mu\nu} = 4g_T^2 N_C I_0 \frac{g^2 \langle S \rangle^2 - 2m_0 g \langle S \rangle}{q^2} (q_\mu q_\nu - q^2 g_{\mu\nu}) + \text{finite terms}, \qquad (21)$$

$$\Delta \Pi^{BB}_{\mu\nu} = -4g_T^2 N_C I_0 \frac{g^2 \langle S \rangle^2 - 2m_0 g \langle S \rangle}{q^2} (q_\mu q_\nu - q^2 g_{\mu\nu}) + \text{finite terms}, \qquad (22)$$

to their mass terms $m_R^2 = \mu_T^2 - 4g_T^2 N_C I_0(m_0^2 + g^2 \langle S \rangle^2 - 2m_0 g \langle S \rangle) = \mu_T^2 - 6m_s^2$ and $m_B^2 = \mu_T^2 + 4g_T^2 N_C I_0(m_0^2 + g^2 \langle S \rangle^2 - 2m_0 g \langle S \rangle) = \mu_T^2 + 6m_s^2$, which also depend on the physical constituent mass of strange quark.

As a result the effective Lagrangian for the spin-1 bosons has the form

$$\mathcal{L}_{\text{eff}} = -\frac{1}{2} \left(V_{\mu} \ R_{\mu} \right) \begin{pmatrix} q^2 - m_V^2 & \sqrt{18m_s^2 q^2} \\ \sqrt{18m_s^2 q^2} \ q^2 - m_B^2 + 12m_s^2 \end{pmatrix} \begin{pmatrix} V^{\mu} \\ R^{\mu} \end{pmatrix} - \frac{1}{2} B_{\mu} \left(q^2 - m_B^2 \right) B^{\mu}.$$
(23)

In paper [6] it has been shown that the masses of spin-1 isovector states ρ , $\rho' = \rho(1450)$, b_1 and spin-1 light isosinglets ω , ω' , $h_1(1170)$ correspond to the hypothesis of the maximal mixing between V_{μ} and R_{μ} states. It leads to the additional constraint

$$m_V^2 = m_B^2 - 12m_s^2. (24)$$

Zeros of determinant of the matrix between $(V_{\mu} R_{\mu})^T$ doublets correspond to the masses for the physical ϕ and ϕ' bosons, while $h_1(s\bar{s})$ meson has m_B mass. Using Vieta's formulas

$$m_{\phi}^2 + m_{\phi'}^2 = 2(m_V^2 + 9m_s^2), \qquad m_{\phi}m_{\phi'} = m_V^2$$
 (25)

for quadratic equation $(q^2)^2 - 2(m_V^2 + 9m_s^2)q^2 + m_V^4 = 0$ and relation (24) we have reproduced the mass formula (4) without the assumption of initial massless quarks.

IV. CONCLUSION

In this paper we have shown that novel mass relation (4) for U(1) case of the strange quark with nonzero current mass, m_0 , has the same form as for initially massless quark. Therefore, the prediction of the mass of axial-vector strangeonium state $h_1(s\bar{s})$ with quantum numbers 1^{+-} in [6] is valid. This is also confirmed experimentally by the Collaboration BESIII [2].

- [1] M. Ablikim et al. (BESIII Collaboration), arXiv:1804.05536v1 [hep-ex]
- [2] M. Ablikim et al. (BESIII Collaboration), Phys. Rev. D98 072005 (2018).
- [3] D. Aston *et al.* (LASS Collaboration), Phys. Lett. B201 573 (1988).
- [4] A. Abele et al. (Crystal Barrel Collaboration), Phys. Lett. B415 280 (1997).
- [5] M. Tanabashi et al. (Particle Data Group), Phys. Rev. D98 030001 (2018) and 2019 update.
- [6] M. Chizhov, JETP Lett. 80 73 (2004), hep-ph/0307100.
- [7] Y. Nambu, "A 'Superconductor' Model of Elementary Particles and its Consequencies", Talk given at a conference at Purdue (1960), reprinted in "Broken Symmetries, Selected Papers by Y. Nambu", eds. T. Eguchi and K. Nishijima, World Scientific (1995).
- [8] M. Gell-Mann, Phys. Lett. 8 214 (1964).
- [9] G. Zweig, preprint CERN-TH-401, January 17 (1964).
- [10] Nambu Y., Jona-Lasinio G., Phys. Rev., 1961, **122**, 345; *ibid.* **124**, 246.
- [11] G. Ecker, Prog. Part. Nucl. Phys. **35** 1 (1995).
- [12] M.V. Chizhov, Mod. Phys. Lett. A 8 2753 (1993).