

Non-associative central extensions of null-filiform associative algebras*

Ivan Kaygorodov^{a, b}, Samuel A. Lopes^c, and Pilar Páez-Guillán^d

^a*CMCC, Universidade Federal do ABC, Santo André, Brazil.*

^b*Siberian Federal University, Krasnoyarsk, Russia.*

^c*CMUP, Faculdade de Ciências da Universidade do Porto, Rua do Campo Alegre 687, 4169-007 Porto, Portugal.*

^d*IMAT, Universidade de Santiago de Compostela, Santiago de Compostela, Spain.
Email addresses: kaygorodov.ivan@gmail.com, slopes@fc.up.pt, pilar.paez@usc.es*

Abstract

We give the algebraic classification of alternative, left alternative, Jordan, bicommutative, left commutative, assosymmetric, Novikov and left symmetric central extensions of null-filiform associative algebras.

MSC Numbers (2010): 16D70, 16S70.

Keywords: central extension, algebraic classification, nilpotent algebras, null-filiform algebra.

Introduction

The subject of algebra extensions has long been a focal point of interest in mathematics and physics. In short, an algebra A is an extension of another algebra B by K if there exists a short exact sequence $0 \rightarrow K \rightarrow A \rightarrow B \rightarrow 0$. The easiest example is the direct sum $B \oplus K$, along with the inclusion and projection maps. Imposing new conditions on the extensions, we find important special types, such as the split extensions, the HNN-extensions, which have been used to prove numerous theorems of embeddability in different varieties of non-associative algebras [19, 20, 22], and many others. In this paper, we will be concerned with central extensions, i.e. extensions in which the annihilator of A contains K . Several important algebras can be constructed as central extensions; for example, the Virasoro algebra is the universal central extension of the Witt algebra, and the Heisenberg algebra is a central extension of a commutative Lie algebra.

The algebraic study of central extensions of different varieties of non-associative algebras plays an important role in the classification problem in such varieties (for example, see [23]).

*The first part of this work is supported by the Russian Science Foundation under grant 18-71-10007. The second part of this work was supported by MTM 2016-79661-P; FPU scholarship (Spain); and CMUP (UID/MAT/00144/2019), which is funded by FCT (Portugal) with national (MCTES) and European structural funds through the programs FEDER, under the partnership agreement PT2020.

Since Skjelbred and Sund in 1978 [21] devised a method for classifying nilpotent Lie algebras, making crucial use of central extensions, it has been adapted to many other varieties of algebras, including associative [8], Malcev [14, 15], Jordan [12, 13], Novikov [16], anticommutative [4], binary Lie [1] or bicommutative [18], among others (see, [5, 6, 7, 10, 11]).

The study of central extensions of null-filiform algebras was initiated in [2], where all Leibniz central extensions of null-filiform Leibniz algebras were described. The associative central extensions of the associative null-filiform algebra of dimension n , which we will denote by μ_0^n , were studied in [17] within the framework of associative algebras, and it was proven that the only associative central extension of μ_0^n is μ_0^{n+1} . However, the null-filiform algebras can be considered as elements of more general varieties of algebras, such as alternative, left alternative, Jordan, bicommutative, left commutative, assosymmetric, Novikov or left symmetric, among others (note that the right alternative, right commutative and right symmetric cases are analogous to their respective left counterparts). In particular, it was proven in [3, 18] that, up to isomorphism, the null-filiform algebra μ_0^3 admits the trivial extension μ_0^4 and also another nontrivial bicommutative extension. Then, it is reasonable to wonder whether there will be nontrivial extensions in the aforementioned varieties of algebras.

Our main result is the classification of the isomorphism classes of central extensions of the associative null-filiform algebra over several varieties of non-associative algebras, as the ones mentioned above. This is in part summarized in Table 1, at the end of Section 5.

The structure of the paper is as follows. Section 1 is devoted to formalizing the method of Skjelbred and Sund over any variety of non-associative algebras defined by a set of polynomial identities. Section 2 presents a quick review of null-filiform algebras. Respectively, Sections 3, 4 and 5 deal with left alternative and alternative, Jordan, and left commutative and bicommutative central extensions. Finally, Section 6 deals with the assosymmetric, Novikov and left symmetric cases, which happen to come out as a trivial corollary of the left commutative and bicommutative cases.

Notation and underlying assumptions Throughout the paper, \mathbb{F} will denote a field. Unless otherwise specified, this field will be arbitrary and all vector spaces, tensor products, (multi)linear maps and automorphism groups will be taken over \mathbb{F} . Given a nonnegative integer n , $[n]$ will denote the set $\{i \in \mathbb{Z} \mid 1 \leq i \leq n\}$. Also, we fix the following notation:

- AS : variety of associative algebras;
- AL : variety of left alternative algebras;
- J : variety of Jordan algebras;
- LC : variety of left commutative algebras;
- RC : variety of right commutative algebras;
- BC : variety of bicommutative algebras.

1 Non-associative central extensions

A variety M of non-associative algebras over \mathbb{F} is defined by a set of identities $\{E_i\}_{i \in I}$ of the form

$$E_i: \sum_{j=1}^{t_i} p_{i,j}(z_1, \dots, z_\ell) = 0,$$

where $Z = \{z_1, \dots, z_\ell\}$ is a finite alphabet, $p_{i,j} = p_{i,j}(z_1, \dots, z_\ell)$ is a non-associative word in Z of length $n_{i,j} \geq 2$, with a coefficient of either 1 or -1 and $Z_{i,j} \subseteq Z$ is the set of letters which occur in $p_{i,j}$.

As $n_{i,j} \geq 2$, each $p_{i,j}$ can be expressed uniquely as a product

$$p_{i,j} = p_{i,j}^1 p_{i,j}^2,$$

where $Z_{i,j} = Z_{i,j}^1 \cup Z_{i,j}^2$ and each $p_{i,j}^k$, with $k \in \{1, 2\}$, either has length one or is another concatenation of products. Recursively, we will obtain $n_{i,j} - 1$ factors $p_{i,j}^\alpha$ with $|Z_{i,j}^\alpha| = 1$ and α of the form $\alpha = \alpha_1, \alpha_2, \alpha_3 \dots, \alpha_\ell$, with $\alpha_k \in \{1, 2\}$ for all k .

Let A be an algebra in the variety M , and let V be a vector space over \mathbb{F} . Following the Skjelbred-Sund method, we introduce the *cocycles* of A with respect to V as the bilinear maps $\theta: A \times A \rightarrow V$ satisfying the set of identities $\{\tilde{E}_i\}_{i \in I}$, where

$$\tilde{E}_i: \sum_{j=1}^{t_i} \theta(p_{i,j}^1, p_{i,j}^2) = 0.$$

These elements form a vector space over \mathbb{F} , which we will denote by $Z_M^2(A, V)$.

We also define the *coboundaries* of A with respect to V as follows. Let f be a linear map from A to V , and set $\delta f: A \times A \rightarrow V$ with $(\delta f)(x, y) = f(xy)$. It is clear from

$$\sum_{j=1}^{t_i} \delta f(p_{i,j}^1, p_{i,j}^2) = \sum_{j=1}^{t_i} f(p_{i,j}) = f(0) = 0,$$

for all $i \in I$, that $\delta f \in Z_M^2(A, V)$. We define $B^2(A, V) = \{\delta f : f \in \text{Hom}(A, V)\}$; it is a linear subspace of $Z_M^2(A, V)$. The quotient space $Z_M^2(A, V)/B^2(A, V)$ is called the *second cohomology space* and is denoted by $H_M^2(A, V)$.

Consider the group $\text{Aut}(A)$ of automorphisms of the algebra A . If θ is a cocycle and $\phi \in \text{Aut}(A)$, we define $\phi \cdot \theta: A \times A \rightarrow V$ by $\phi \cdot \theta(x, y) = \theta(\phi(x), \phi(y))$. The automorphism ϕ preserves the product, so $\phi(p_{i,j}(x_1, \dots, x_\ell)) = p_{i,j}(\phi(x_1), \dots, \phi(x_\ell))$; from this, we have that $\phi \cdot \theta \in Z_M^2(A, V)$. This induces an action of $\text{Aut}(A)$ on $Z_M^2(A, V)$. Moreover, $\theta = \delta f \iff \phi \cdot \theta = \delta(f \circ \phi)$, so the action is inherited by the quotient $H_M^2(A, V)$. Although this is a right action, we will write it on the left to follow the usual convention. The orbit of an element $\theta \in Z_M^2(A, V)$ will be denoted by $\text{Orb}(\theta)$, and the orbit of $[\theta] \in H_M^2(A, V)$, by $\text{Orb}([\theta])$.

For every bilinear map $\theta: A \times A \rightarrow V$, we can define the algebra $A_\theta = A \oplus V$ with the product $[x + v, y + w]_\theta = xy + \theta(x, y)$.

Lemma 1.1. *The algebra A_θ belongs to the variety M if and only if $\theta \in Z_M^2(A, V)$.*

Proof. By definition, A_θ belongs to M if it satisfies the identities $\{E_i\}_{i \in I}$, i.e., if it satisfies

$$\sum_{j=1}^{t_i} p_{i,j}(x_1 + v_1, \dots, x_\ell + v_\ell) = 0,$$

for all $i \in I$ and all $x_k \in A$, $v_k \in V$, with $k \in [\ell]$. For the sake of brevity, we will write

$$\sum_{j=1}^{t_i} p_{i,j}(X + V) = 0;$$

also, when we make reference to these identities in the algebra A , we will write

$$\sum_{j=1}^{t_i} p_{i,j}(X) = 0.$$

It holds that

$$p_{i,j}(X + V) = [p_{i,j}^1(X + V), p_{i,j}^2(X + V)]_\theta.$$

An easy induction in the cardinal $|Z_{i,j}|$ shows that

$$p_{i,j}(X + V) = p_{i,j}(X) + \theta(p_{i,j}^1(X), p_{i,j}^2(X)).$$

As A belongs to the variety M , it follows that A_θ satisfies the identities $\{E_i\}_{i \in I}$ if and only if θ satisfies $\{\tilde{E}_i\}_{i \in I}$. \square

In all that follows, we will consider A_θ just for $\theta \in Z_M^2(A, V)$. Then, it is easy to check that the algebra A_θ of the variety M is a central extension of A by V . We define the dimension of the extension as the dimension of V .

The particular identities of the variety M are not involved any more in the development of the method. Thus, we will just give a brief overview of the main definitions and results, and refer the reader to other texts in which this method is carefully detailed, such as [12, 14].

One fundamental definition is the *annihilator* of θ , $\text{Ann}(\theta) = \{x \in A \mid \theta(x, A) + \theta(A, x) = 0\}$. Recalling that the annihilator of an algebra A is $\text{Ann}(A) = \{x \in A \mid xA + Ax = 0\}$, it is clear that $\text{Ann}(A_\theta) = (\text{Ann}(A) \cap \text{Ann}(\theta)) \oplus V$.

Other important notions are the following. Let A be an algebra and I be a nonzero subspace of $\text{Ann}(A)$. If $A = A_0 \oplus I$ for some ideal A_0 , then I is called an *annihilator component* of A . The central extensions of A with no annihilator component are called *non-split central extensions*.

We show now that any algebra A of the variety M with nonzero annihilator is isomorphic to a central extension of some other suitable algebra A' in M . The demonstration can be found in [14, Lemma 5].

Lemma 1.2. *Let A be an n -dimensional algebra in the variety M such that $\dim(\text{Ann}(A)) = s \neq 0$. Then there exist, up to isomorphism, a unique $(n - s)$ -dimensional algebra A' in M and a cocycle $\theta \in Z_M^2(A, V)$ for some vector space V of dimension s , such that $\text{Ann}(A) \cap \text{Ann}(\theta) = 0$, $A \cong A'_\theta$ and $A/\text{Ann}(A) \cong A'$.*

Then, in order to decide when two algebras with nonzero annihilator of the variety M are isomorphic, it suffices to find criteria in terms of the cocycles.

Let us fix a basis $\{e_1, \dots, e_s\}$ of V . For every algebra A and every cocycle $\theta \in Z_M^2(A, V)$, there exist s unique cocycles $\theta_i \in Z_M^2(A, \mathbb{F})$, $i \in [s]$, such that $\theta(x, y) = \sum_{i=1}^s \theta_i(x, y)e_i$, for all $x, y \in A$. Straightforward facts about these cocycles are that $\theta \in B^2(A, V)$ if and only if $\theta_i \in B^2(A, \mathbb{F})$ for $i \in [s]$, and that $\text{Ann}(\theta) = \text{Ann}(\theta_1) \cap \dots \cap \text{Ann}(\theta_s)$. Also, it is not difficult to prove (see [14, Lemma 13]) that if $\text{Ann}(A) \cap \text{Ann}(\theta) = 0$, then A_θ has an annihilator component if and only if $[\theta_1], \dots, [\theta_s]$ are linearly dependent in $H_M^2(A, \mathbb{F})$. In other words, if $\text{Ann}(A) \cap \text{Ann}(\theta) = 0$, then A_θ is a non-split central extension of A if and only if $\langle [\theta_1], \dots, [\theta_s] \rangle \in G_s(H_M^2(A, \mathbb{F}))$, where the Grassmanian $G_s(H_M^2(A, \mathbb{F}))$ is the set of s -dimensional linear subspaces of $H_M^2(A, \mathbb{F})$. Finally (see [14, Lemma 15]), if $\langle [\theta_1], \dots, [\theta_s] \rangle =$

$\langle [\vartheta_1], \dots, [\vartheta_s] \rangle \in G_s(\mathbf{H}_M^2(\mathbf{A}, \mathbb{F}))$, we have that $\bigcap_{i=1}^s \text{Ann}(\theta_i) \cap \text{Ann}(\mathbf{A}) = \bigcap_{i=1}^s \text{Ann}(\vartheta_i) \cap \text{Ann}(\mathbf{A})$, and therefore the set

$$T_s(\mathbf{A}) = \{W = \langle [\theta_1], \dots, [\theta_s] \rangle \in G_s(\mathbf{H}_M^2(\mathbf{A}, \mathbb{F})) \mid \bigcap_{i=1}^s \text{Ann}(\theta_i) \cap \text{Ann}(\mathbf{A}) = 0\}$$

is well-defined. Then, the set of all non-split central extensions of \mathbf{A} with s -dimensional annihilator, by some s -dimensional vector space \mathbf{V} , can be written as

$$E(\mathbf{A}, \mathbf{V}) = \{A_\theta \mid \theta(x, y) = \sum_{i=1}^s \theta_i(x, y)e_i \text{ and } \langle [\theta_1], \dots, [\theta_s] \rangle \in T_s(\mathbf{A})\}.$$

Now, if ϕ is an automorphism of \mathbf{A} and $W = \langle [\theta_1], \dots, [\theta_s] \rangle \in G_s(\mathbf{H}_M^2(\mathbf{A}, \mathbb{F}))$ it holds that $\phi W = \langle [\phi\theta_1], \dots, [\phi\theta_s] \rangle$ also has dimension s . This induces an action of $\text{Aut } \mathbf{A}$ on $G_s(\mathbf{H}_M^2(\mathbf{A}, \mathbb{F}))$; the orbit of W will be denoted by $\text{Orb}(W)$. The set $T_s(\mathbf{A})$ is stable under this action ([14, Lemma 16]).

Having established these results, we can determine whether two s -dimensional non-split central extensions A_θ and A_ϑ in the variety \mathbf{M} are isomorphic or not. For the proof, see [14, Lemma 17].

Lemma 1.3. *Let $A_\theta, A_\vartheta \in E(\mathbf{A}, \mathbf{V})$. Suppose that $\theta(x, y) = \sum_{i=1}^s \theta_i(x, y)e_i$ and $\vartheta(x, y) = \sum_{i=1}^s \vartheta_i(x, y)e_i$. Then the algebras A_θ and A_ϑ of the variety \mathbf{M} are isomorphic if and only if*

$$\text{Orb}(\langle [\theta_1], \dots, [\theta_s] \rangle) = \text{Orb}(\langle [\vartheta_1], \dots, [\vartheta_s] \rangle).$$

Therefore, given an algebra \mathbf{A} of dimension $n - s$ in the variety \mathbf{M} , we can construct all non-split central extensions A_θ of \mathbf{A} according to the following procedure:

1. Determine $\mathbf{H}_M^2(\mathbf{A}, \mathbb{F})$, $\text{Ann}(\mathbf{A})$ and $\text{Aut } \mathbf{A}$.
2. Determine the set of $(\text{Aut } \mathbf{A})$ -orbits on $T_s(\mathbf{A})$.
3. For each orbit, construct the algebra of the variety \mathbf{M} associated with a representative of it.

Then, thanks to this procedure and to Lemma 1.2, we can classify all the nilpotent algebras of dimension n of a variety \mathbf{M} , provided that the nilpotent algebras of dimension $n - 1$ are known.

Let \mathbf{A} be an algebra and fix a basis $\{e_1, e_2, \dots, e_n\}$ of \mathbf{A} . We define the bilinear form $\Delta_{i,j}: \mathbf{A} \times \mathbf{A} \rightarrow \mathbb{F}$ by $\Delta_{i,j}(e_l, e_m) = \delta_{i,l}\delta_{j,m}$. Then the set $\{\Delta_{i,j} \mid i, j \in [n]\}$ is a basis for the linear space of bilinear forms on \mathbf{A} ; in particular, every $\theta \in Z^2(\mathbf{A}, \mathbf{V})$ can be uniquely written as $\theta = \sum_{i,j=1}^n c_{i,j}\Delta_{i,j}$, with $c_{i,j} \in \mathbb{F}$.

2 Null-filiform associative algebras

For an algebra A of an arbitrary variety M , we consider the series

$$A^1 = A, \quad A^{i+1} = \sum_{k=1}^i A^k A^{i+1-k}, \quad i \geq 1.$$

An n -dimensional algebra A is called *null-filiform* if $\dim A^i = (n+1) - i$, $i \in [n+1]$. All null-filiform associative algebras were described in [9].

Theorem 2.1. *An arbitrary n -dimensional null-filiform associative algebra is isomorphic to the algebra:*

$$\mu_0^n : \quad e_i e_j = e_{i+j}, \quad i, j \in [n],$$

where $\{e_1, e_2, \dots, e_n\}$ is a basis of the algebra μ_0^n and we define $e_m = 0$ for all $m > n$.

Using the procedure explained in Section 1, we can easily find all associative central extensions of μ_0^n . Let $\nabla_j = \sum_{k=1}^j \Delta_{k, j+1-k}$, for $j \in [n]$. We need the following result from [17].

Proposition 2.2. *Let μ_0^n be the null-filiform associative algebra of dimension n . Then*

$$\begin{aligned} Z_{AS}^2(\mu_0^n, \mathbb{F}) &= \langle \nabla_j \mid j \in [n] \rangle, & B^2(\mu_0^n, \mathbb{F}) &= \langle \nabla_j \mid j \in [n-1] \rangle, \\ H_{AS}^2(\mu_0^n, \mathbb{F}) &= Z_A^2(\mu_0^n, \mathbb{F}) / B^2(\mu_0^n, \mathbb{F}) = \langle [\nabla_n] \rangle. \end{aligned}$$

Remark 2.3. The above result appears in [17] for $\mathbb{F} = \mathbb{C}$, the field of complex numbers. Nevertheless, it is immediate to check, e.g. using the methods appearing shortly in this paper, that it does hold over an arbitrary field.

As $\dim H_{AS}^2(\mu_0^n, \mathbb{F}) = 1$, the next result follows easily.

Theorem 2.4. *Every non-split 1-dimensional associative central extension of μ_0^n is isomorphic to μ_0^{n+1} .*

In the following sections we will study non-associative central extensions of μ_0^n . It is easy to see that μ_0^{n+1} is an associative central extension of μ_0^n . All extensions of this type will be called trivial. The basis $\{e_1, \dots, e_n\}$ of μ_0^n will always be assumed to satisfy the relations from Theorem 2.1.

3 Alternative and left alternative central extensions

Throughout this section, we assume that the characteristic of the field \mathbb{F} satisfies $\text{char}(\mathbb{F}) \neq 2$. Recall that an algebra A is said to be left alternative (respectively, right alternative) if it satisfies the identity

$$x(xy) = (xx)y \quad (\text{respectively, } (xy)y = x(yy)),$$

for all $x, y \in A$. Also, if A is both left alternative and right alternative, it is called alternative.

Let us consider μ_0^n as a left alternative algebra. Note also that the linearization of the left alternative identity for μ_0^n leads to

$$e_i(e_j e_k) + e_j(e_i e_k) = (e_i e_j) e_k + (e_j e_i) e_k,$$

for $i, j, k \in [n]$. Then, its space of cocycles is formed by all the bilinear maps $\theta: \mu_0^n \times \mu_0^n \rightarrow \mathbb{F}$ satisfying $\theta(e_i, e_j e_k) + \theta(e_j, e_i e_k) = \theta(e_i e_j, e_k) + \theta(e_j e_i, e_k)$. This can be expressed as

$$\theta(e_i, e_{j+k}) + \theta(e_j, e_{i+k}) = 2\theta(e_{i+j}, e_k), \quad (3.1)$$

for $i, j, k \in [n]$, and considering that $e_m = 0$ for $m > n$.

Theorem 3.2. *Assume that $\text{char}(\mathbb{F}) \neq 2$. Then all left alternative and all alternative central extensions of μ_0^n are trivial.*

Proof. The first step is to compute $Z_{LA}^2(\mu_0^n, \mathbb{F})$. Let $\theta = \sum_{i,j} c_{i,j} \Delta_{i,j}$ be an arbitrary cocycle of μ_0^n considered as a left alternative algebra. The identity (3.1) leads to

$$c_{i,j+k} + c_{j,i+k} = 2c_{i+j,k}$$

for $i, j, k \in [n]$, with the assumption that $c_{i,j} = 0$ if $i > n$ or $j > n$. Given integers m, s such that $m, s - m \in [n]$ and $m \geq 2$, and taking $i = m - 1$, $j = 1$ and $k = s - m$ in the above equation, we get

$$2c_{m,s-m} = c_{m-1,s-(m-1)} + c_{1,s-1}. \quad (3.3)$$

Claim: For all $i, j \in [n]$, $c_{i,j} = c_{1,i+j-1}$. In particular, $c_{i,j} = 0$ if $i + j \geq n + 2$.

The claim will follow by induction on i , the case $i = 1$ being trivial. So assume that $i \geq 2$ and that the claim holds for $i - 1$. Taking $m = i$ and $s = i + j$ in (3.3), we get

$$2c_{i,j} = c_{i-1,j+1} + c_{1,i+j-1}.$$

If $j = n$, then $j + 1, i + j - 1 \geq n + 1$ so $c_{i-1,j+1} = 0 = c_{1,i+j-1}$ and it follows from the identity above that $c_{i,j} = 0$, as $\text{char}(\mathbb{F}) \neq 2$. In particular, $c_{i,j} = c_{1,i+j-1}$ holds. Otherwise, if $j < n$, then we can use the induction hypothesis to obtain $2c_{i,j} = c_{i-1,j+1} + c_{1,i+j-1} = 2c_{1,i+j-1}$, whence the claim.

So, it is clear that $Z_{LA}^2(\mu_0^n, \mathbb{F})$ is in the linear space spanned by $\{\nabla_j\}_{j=1}^n$; it is also immediate to see that every element from $Z_{AS}^2(\mu_0^n, \mathbb{F})$ is a left alternative cocycle. Thus, $H_{LA}^2(\mu_0^n, \mathbb{F}) = H_{AS}^2(\mu_0^n, \mathbb{F})$, and every left alternative central extension is trivial. Since any alternative extension is left alternative, it follows that every alternative central extension is trivial as well. □

4 Jordan central extensions

Throughout this section, we assume that the characteristic of the field \mathbb{F} satisfies $\text{char}(\mathbb{F}) \neq 2, 3$. Recall that a commutative algebra A is said to be Jordan if it satisfies

$$x^2(yx) = (x^2y)x,$$

for all $x, y \in A$. It is immediate to check that the commutative algebra μ_0^n satisfies the previous identity and therefore is a Jordan algebra.

The space of Jordan cocycles of μ_0^n is formed by all the bilinear maps $\theta: \mu_0^n \times \mu_0^n \rightarrow \mathbb{F}$ satisfying

$$\theta(x, y) = \theta(y, x), \quad (4.1)$$

$$\theta(x^2, yx) = \theta(x^2y, x). \quad (4.2)$$

Equivalently, since $\text{char}(\mathbb{F}) \neq 2, 3$, we have

$$\theta(e_i, e_j) = \theta(e_j, e_i),$$

$$\theta(e_{i+\ell}, e_{j+k}) + \theta(e_{j+\ell}, e_{i+k}) + \theta(e_{k+\ell}, e_{i+j}) = \theta(e_i, e_{j+k+\ell}) + \theta(e_j, e_{i+k+\ell}) + \theta(e_k, e_{i+j+\ell}),$$

for $i, j, k, \ell \in [n]$. In particular, taking $i = j$, we see that every cocycle must verify

$$2\theta(e_{i+\ell}, e_{i+k}) + \theta(e_{\ell+k}, e_{2i}) = 2\theta(e_i, e_{i+k+\ell}) + \theta(e_k, e_{2i+\ell}). \quad (4.3)$$

Theorem 4.4. *All Jordan central extensions of μ_0^n are trivial.*

Proof. We will prove that $\{\nabla_j\}_{j=1}^n$ is a basis for $Z_J^2(\mu_0^n, \mathbb{F})$.

Let $\theta = \sum_{i,j} c_{i,j} \Delta_{i,j}$ be an arbitrary cocycle in $Z_J^2(\mu_0^n, \mathbb{F})$. Note that $c_{i,j} = c_{j,i}$, for all $i, j \in [n]$. By (4.3), we also have

$$c_{k,2a+b} = 2c_{a+b,a+k} + c_{b+k,2a} - 2c_{a,a+b+k}, \quad (4.5)$$

for all $a, b, k \in [n]$, with the assumption that $c_{i,j} = 0$ in case $i > n$ or $j > n$.

Claim: For all $i, j \in [n]$, $c_{i,j} = c_{1,i+j-1}$. In particular, $c_{i,j} = 0$ if $i + j \geq n + 2$.

The proof is by induction on i . If either $i = 1$ or $j = 1$, the claim is trivial, by commutativity. So we can assume that $i, j \geq 2$. If $i = 2$, then

$$c_{2,j} = \theta(e_1^2, e_{j-1}e_1) = \theta(e_{j+1}, e_1) = \theta(e_1, e_{j+1}) = c_{1,j+1}.$$

Let us assume thus that $i \geq 3$ and $j \geq 2$. There are unique integers a, b with $b \in \{1, 2\}$ such that $i = 2a + b$. Notice that $1 \leq a < a + b < i \leq n$. Then, using (4.5) with $k = j$, we get

$$c_{i,j} = c_{2a+b,j} = 2c_{a+b,a+j} + c_{b+j,2a} - 2c_{a,a+b+j}.$$

There are two cases to consider.

Case1: $a + b + j \leq n$. In this case, we can use the inductive hypothesis and the commutativity to get

$$\begin{aligned} 2c_{a+b,a+j} + c_{b+j,2a} - 2c_{a,a+b+j} &= 2c_{1,2a+b+j-1} + c_{1,2a+b+j-1} - 2c_{1,2a+b+j-1} \\ &= c_{1,2a+b+j-1} = c_{1,i+j-1}. \end{aligned}$$

Case2: $a + b + j \geq n + 1$. Then $c_{a,a+b+j} = 0$ and we have $c_{i,j} = 2c_{a+b,a+j} + c_{2a,b+j}$. Either $a + j \geq n + 1$ and hence $c_{a+b,a+j} = 0$, or $a + j \leq n$ and the inductive hypothesis says that $c_{a+b,a+j} = c_{1,2a+b+j-1} = 0$, because $2a + b + j - 1 \geq a + b + j \geq n + 1$. In any case, $c_{a+b,a+j} = 0$. Similarly, $c_{2a,b+j} = 0$ and we conclude that $c_{i,j} = 0 = c_{1,i+j-1}$, because $i + j - 1 \geq a + b + j \geq n + 1$.

The claim is thus established. It remains to observe that the $\{\nabla_j\}_{j=1}^n$ are indeed Jordan cocycles. In [17], it is proved that $\nabla_j \in Z_{\text{AS}}^2(\mu_0^n, \mathbb{F})$, i.e. they verify $\nabla_j(xy, z) = \nabla_j(x, yz)$. Clearly, also $\nabla_j(x, y) = \nabla_j(y, x)$. These conditions are indeed stricter than conditions (4.1) and (4.2), so it is clear that $\nabla_j \in Z_J^2(\mu_0^n, \mathbb{F})$. This implies that $H_J^2(\mu_0^n, \mathbb{F}) = H_{\text{AS}}^2(\mu_0^n, \mathbb{F})$ and, according to [17], the only Jordan central extension of μ_0^n is μ_0^{n+1} . \square

5 Left commutative and bicommutative central extensions

Recall that an algebra A is said to be left (respectively, right) commutative if it satisfies

$$x(yz) = y(xz) \quad (\text{respectively, } (xy)z = (xz)y),$$

for all $x, y, z \in A$. Equivalently, A is left (respectively, right) commutative if and only if the left (respectively, right) multiplication operators commute. In case A is both left and right commutative, we say that A is bicommutative.

The main results in this section will require the field \mathbb{F} to be algebraically closed and of sufficiently large characteristic (which for simplicity we will assume to be 0, when necessary), while others hold for arbitrary fields. Unless otherwise is stated, it should be assumed that the field \mathbb{F} is arbitrary.

Proposition 5.1. *Let $n \geq 2$ and recall the bilinear forms $\nabla_j = \sum_{k=1}^j \Delta_{k,j+1-k}$, defined for $j \in [n]$. Then the following hold:*

- (a) $\dim Z_{\text{LC}}^2(\mu_0^n, \mathbb{F}) = 2n - 1$ and $\{\Delta_{i,1} \mid 2 \leq i \leq n\} \cup \{\nabla_j \mid j \in [n]\}$ is a basis of $Z_{\text{LC}}^2(\mu_0^n, \mathbb{F})$.
- (b) $\dim H_{\text{LC}}^2(\mu_0^n, \mathbb{F}) = n$ and the classes $\{[\Delta_{i,1}] \mid 2 \leq i \leq n\} \cup \{[\nabla_n]\}$ form a basis of $H_{\text{LC}}^2(\mu_0^n, \mathbb{F})$.
- (c) In the bicommutative case we have that $\dim H_{\text{BC}}^2(\mu_0^n, \mathbb{F}) = 2$, with basis $\{[\Delta_{2,1}], [\nabla_n]\}$.

Proof. The space $Z_{\text{LC}}^2(\mu_0^n, \mathbb{F})$ consists of the bilinear forms $\theta = \sum_{i,j=1}^n c_{i,j} \Delta_{i,j}$, with $c_{i,j} \in \mathbb{F}$, satisfying $\theta(e_i, e_j e_k) = \theta(e_j, e_i e_k)$, for all $i, j, k \in [n]$.

Claim: If $j \geq 2$, then $c_{i,j} = c_{1,i+j-1}$. In particular, if $i + j \geq n + 2$ then $c_{i,j} = 0$.

For $j \geq 2$ we have

$$c_{i,j} = \theta(e_i, e_j) = \theta(e_i, e_1 e_{j-1}) = \theta(e_1, e_i e_{j-1}) = c_{1,i+j-1}. \quad (5.2)$$

Thus, if $i + j \geq n + 2$, then necessarily $j \geq 2$ and (5.2) gives $c_{i,j} = 0$. This establishes the claim.

Hence, $c_{i,2} = c_{i-1,3} = \cdots = c_{2,i} = c_{1,i+1}$ for all $i \in [n-1]$ and we can write

$$\theta = \sum_{i=1}^n c_{i,1} \Delta_{i,1} + \sum_{j=2}^n c_{1,j} (\nabla_j - \Delta_{j,1}) = \sum_{i=2}^n (c_{i,1} - c_{1,i}) \Delta_{i,1} + \sum_{j=1}^n c_{1,j} \nabla_j, \quad (5.3)$$

which shows that $Z_{\text{LC}}^2(\mu_0^n, \mathbb{F}) \subseteq \langle \{\Delta_{i,1} \mid 2 \leq i \leq n\} \cup \{\nabla_j \mid j \in [n]\} \rangle$.

It remains to prove the reverse inclusion, i.e. that the $\Delta_{i,1}$ and the ∇_j are left commutative cocycles. Let $k, \ell, m \in [n]$. Then

$$\Delta_{i,1}(e_k, e_\ell e_m) = 0 = \Delta_{i,1}(e_\ell, e_k e_m),$$

so indeed $\Delta_{i,1} \in Z_{\text{LC}}^2(\mu_0^n, \mathbb{F})$. As $\langle \nabla_j \mid j \in [n-1] \rangle = \mathbf{B}^2(\mu_0^n, \mathbb{F}) \subseteq Z_{\text{LC}}^2(\mu_0^n, \mathbb{F})$, it remains to show that ∇_n is a cocycle. This follows immediately from the fact that

$$\nabla_n(e_k, e_\ell e_m) = \begin{cases} 1 & \text{if } k + \ell + m = n + 1; \\ 0 & \text{otherwise.} \end{cases}$$

This concludes the proof of (a). Part (b) is clear by $\mathbf{B}^2(\mu_0^n, \mathbb{F}) = \langle \nabla_j \mid j \in [n-1] \rangle$.

We precede the proof of (c) with some general considerations. Let A be an algebra and denote by $\nu : A \times A \rightarrow A$ its multiplication map. The opposite algebra A^{op} is the algebra with the same underlying vector space and with multiplication $\nu \circ \tau$, where $\tau : A \times A \rightarrow A \times A$, $(a, b) \mapsto (b, a)$ is the flip. Clearly, A is right commutative if and only if A^{op} is left commutative and $\theta \in \mathbf{Z}_{\text{RC}}^2(A, \mathbb{F})$ if and only if $\theta \circ \tau \in \mathbf{Z}_{\text{LC}}^2(A^{\text{op}}, \mathbb{F})$. Moreover, $\mathbf{Z}_{\text{BC}}^2(A, \mathbb{F}) = \mathbf{Z}_{\text{LC}}^2(A, \mathbb{F}) \cap \mathbf{Z}_{\text{RC}}^2(A, \mathbb{F})$.

We return now to the case of the algebra μ_0^n . This algebra is commutative, so it coincides with its opposite algebra. Let $\theta \in \mathbf{Z}_{\text{BC}}^2(\mu_0^n, \mathbb{F})$. Then $\theta \in \mathbf{Z}_{\text{LC}}^2(\mu_0^n, \mathbb{F})$ and by (a) we can write θ as in (5.3). What is more, for such a θ we have $\theta \in \mathbf{Z}_{\text{BC}}^2(\mu_0^n, \mathbb{F})$ if and only if $\theta \circ \tau \in \mathbf{Z}_{\text{LC}}^2(\mu_0^n, \mathbb{F})$. Seeing that $\Delta_{i,j} \circ \tau = \Delta_{j,i}$, we obtain $\nabla_j \circ \tau = \nabla_j$, for all $j \in [n]$.

Thence, in view of the results in (a), and the observation that $\Delta_{1,2} = \nabla_2 - \Delta_{2,1}$, we have that $\theta \circ \tau \in \mathbf{Z}_{\text{LC}}^2(\mu_0^n, \mathbb{F})$ if and only if $c_{i,1} = c_{1,i}$ for all $3 \leq i \leq n$. This proves that $\{\nabla_j \mid j \in [n]\} \cup \{\Delta_{2,1}\}$ is a basis of $\mathbf{Z}_{\text{BC}}^2(\mu_0^n, \mathbb{F})$. Finally, given our description of $\mathbf{B}^2(\mu_0^n, \mathbb{F})$, we immediately obtain the basis $\{[\Delta_{2,1}], [\nabla_n]\}$ of $\mathbf{H}_{\text{BC}}^2(\mu_0^n, \mathbb{F})$. \square

5.1 The automorphism group of μ_0^n

We denote by $\text{Aut}(\mu_0^n)$ the automorphism group of μ_0^n . Let $\phi \in \text{Aut}(\mu_0^n)$. Then we identify ϕ with its matrix $(\phi_{i,j})_{i,j \in [n]}$ relative to the basis $\{e_1, \dots, e_n\}$. It is easy to see that the automorphisms of μ_0^n are precisely those linear endomorphisms $\phi = (\phi_{i,j})_{i,j \in [n]}$ with $\phi_{1,1} \neq 0$, $\phi_{2,1}, \dots, \phi_{n,1} \in \mathbb{F}$ arbitrary and $\phi_{i,j}$ with $2 \leq j \leq n$ determined by $\phi(e_j) = \phi(e_1)^j$. In other words,

$$\phi_{i,j} = \sum_{k_1 + \dots + k_j = i} \phi_{k_1,1} \cdots \phi_{k_j,1},$$

for all $i, j \in [n]$.

It follows that $\phi_{i,i} = \phi_{1,1}^i$ and $\phi_{i,j} = 0$ if $j > i$. For $j < i$ we also have

$$\phi_{i,j} = j\phi_{1,1}^{j-1}\phi_{i-j+1,1} + p_{i,j}(\phi_{1,1}, \phi_{2,1}, \dots, \phi_{i-j,1}), \quad (5.4)$$

for some polynomial $p_{i,j}$ with coefficients in \mathbb{F} which depends only on i and j .

Write $\phi = \sum_{i,j=1}^n \phi_{i,j} E_{i,j}$, where we think of $E_{i,j}$ both as the matrix unit with all entries equal to 0 except for the entry (i, j) which equals 1, and also as the linear endomorphism of μ_0^n defined by $E_{i,j}(e_k) = \delta_{j,k} e_i$. Then,

$$\phi \cdot \Delta_{s,t}(e_k, e_\ell) = \Delta_{s,t}(\phi(e_k), \phi(e_\ell)) = \sum_{k \leq i} \sum_{\ell \leq j} \phi_{i,k} \phi_{j,\ell} \Delta_{s,t}(e_i, e_j) = \sum_{k \leq s, \ell \leq t} \phi_{s,k} \phi_{t,\ell}.$$

Thence, the formula for the action is given as

$$\phi \cdot \Delta_{s,t} = \sum_{k \leq s, \ell \leq t} \phi_{s,k} \phi_{t,\ell} \Delta_{k,\ell}. \quad (5.5)$$

5.2 The orbit decomposition of $\mathbf{H}_{\text{LC}}^2(\mu_0^n, \mathbb{F})$

To state the main result of this subsection we need an additional definition. For $0 \leq i \leq n$, let $R(i, n)$ be the multiplicative subgroup of \mathbb{F}^* consisting of all $(i+1)$ -st powers of all $(n+1)$ -st roots of unity in \mathbb{F}^* . In case \mathbb{F} is algebraically closed of characteristic 0, there exists some

primitive $(n+1)$ -st root of unity ζ and $R(i, n) = \langle \zeta^{i+1} \rangle$ is the cyclic group generated by ζ^{i+1} . Denote the quotient group by

$$\mathbb{F}_{(i,n)} = \mathbb{F}^*/R(i, n), \quad (5.6)$$

and for $\mu \in \mathbb{F}^*$, let $\bar{\mu} = \mu R(i, n)$ be the corresponding coset.

Theorem 5.7. *Assume that \mathbb{F} is algebraically closed of characteristic 0 and let $n \geq 2$. The following elements of $Z_{\text{LC}}^2(\mu_0^n, \mathbb{F})$ give a complete list of distinct representatives of the orbits of the automorphism group $\text{Aut}(\mu_0^n)$ on $H_{\text{LC}}^2(\mu_0^n, \mathbb{F})$:*

- (a) 0;
- (b) $\{\Delta_{i,1} \mid 2 \leq i \leq n\}$;
- (c) $\{\nabla_n + \mu\Delta_{n,1} \mid \mu \in \mathbb{F}\}$;
- (d) $\{\nabla_n + \bar{\mu}\Delta_{i,1} \mid 2 \leq i \leq n-1, \bar{\mu} \in \mathbb{F}_{(i,n)}\}$.

From this result and Proposition 5.1, we immediately deduce the orbit space decomposition in the bicommutative case.

Corollary 5.8. *Assume that \mathbb{F} is algebraically closed of characteristic 0 and let $n \geq 2$. The following elements of $Z_{\text{BC}}^2(\mu_0^n, \mathbb{F})$ give a complete list of distinct representatives of the orbits of the automorphism group $\text{Aut}(\mu_0^n)$ on $H_{\text{BC}}^2(\mu_0^n, \mathbb{F})$:*

(Case $n = 2$)

- (a) 0;
- (b) $\Delta_{2,1}$;
- (c) $\{\nabla_2 + \mu\Delta_{2,1} \mid \mu \in \mathbb{F}\}$.

(Case $n > 2$)

- (a) 0;
- (b) $\Delta_{2,1}$;
- (c) ∇_n ;
- (d) $\{\nabla_n + \bar{\mu}\Delta_{2,1} \mid \bar{\mu} \in \mathbb{F}_{(2,n)}\}$.

We devote the remainder of this section to the proof of Theorem 5.7.

Lemma 5.9. *Assume that \mathbb{F} is algebraically closed of characteristic 0. Fix $i \geq 2$. Given scalars $a_{1,1}, \dots, a_{i-1,1} \in \mathbb{F}$ with $a_{1,1} \neq 0$, we define, for $2 \leq \ell, m \leq i$:*

$$a_{\ell,m} = \sum_{j_1 + \dots + j_m = \ell} a_{j_1,1} \cdots a_{j_m,1}. \quad (5.10)$$

Note that $a_{\ell,m}$ depends only on $a_{1,1}, \dots, a_{\ell-m+1,1}$. Then, for any $0 \leq k \leq i-1$, the map

$$\begin{aligned} \rho_k : \quad \mathbb{F}^* \times \mathbb{F}^k &\longrightarrow \mathbb{F}^* \times \mathbb{F}^k \\ (a_{1,1}, \dots, a_{k+1,1}) &\mapsto (a_{i,i}a_{1,1}, a_{i,i-1}a_{1,1}, \dots, a_{i,i-k}a_{1,1}) \end{aligned}$$

is onto and $(i+1)$ -to-1, i.e. $|\rho_k^{-1}(\lambda_1, \dots, \lambda_{k+1})| = i+1$, for all $(\lambda_1, \dots, \lambda_{k+1}) \in \mathbb{F}^* \times \mathbb{F}^k$.

Proof. The proof is by induction on $0 \leq k \leq i - 1$. First, notice that (5.10) implies that $a_{\ell,\ell} = a_{1,1}^\ell$. If $k = 0$, then $\rho_0(a_{1,1}) = a_{1,1}^{i+1}$ and the relation $|\rho_0^{-1}(\lambda)| = i + 1$ for any $\lambda \in \mathbb{F}^*$ follows from both our assumptions on \mathbb{F} .

Now assume that $k \geq 1$ and take $(\lambda_1, \dots, \lambda_{k+1}) \in \mathbb{F}^* \times \mathbb{F}^k$. By the induction hypothesis, $|\rho_{k-1}^{-1}(\lambda_1, \dots, \lambda_k)| = i + 1$, so choose $(a_{1,1}, \dots, a_{k,1}) \in \mathbb{F}^* \times \mathbb{F}^{k-1}$ such that $\rho_{k-1}(a_{1,1}, \dots, a_{k,1}) = (\lambda_1, \dots, \lambda_k)$. As in (5.4), we have $a_{i,i-k}a_{1,1} = (i-k)a_{1,1}^{i-k}a_{k+1,1} + a_{1,1}p_{i,i-k}(a_{1,1}, \dots, a_{k,1})$, with $p_{i,i-k}$ a polynomial in $a_{1,1}, \dots, a_{k,1}$. Thus, we can set

$$a_{k+1,1} = \frac{\lambda_{k+1} - a_{1,1}p_{i,i-k}(a_{1,1}, \dots, a_{k,1})}{(i-k)a_{1,1}^{i-k}}, \quad (5.11)$$

and, by construction, $\rho_k(a_{1,1}, \dots, a_{k+1,1}) = (\rho_{k-1}(a_{1,1}, \dots, a_{k,1}), a_{i,i-k}a_{1,1}) = (\lambda_1, \dots, \lambda_{k+1})$.

The above shows that the $i + 1$ distinct solutions of the problem for $k - 1$ give rise to $i + 1$ distinct solutions of the problem for k . Conversely, each solution $(a_{1,1}, \dots, a_{k+1,1})$ of the latter determines a solution $(a_{1,1}, \dots, a_{k,1})$ of the former and, by (5.11), $a_{k+1,1}$ is completely determined by $a_{1,1}, \dots, a_{k,1}$, so there are exactly $i + 1$ solutions of the problem for k . By induction, the proof is complete. \square

Now we can start computing orbits of $\text{Aut}(\mu_0^n)$ on $Z_{\text{LC}}^2(\mu_0^n, \mathbb{F})$.

Proposition 5.12. *Assume that \mathbb{F} is algebraically closed of characteristic 0. For $i \in [n]$ we have*

$$\text{Orb}(\Delta_{i,1}) = \{\lambda_1\Delta_{i,1} + \lambda_2\Delta_{i-1,1} + \dots + \lambda_i\Delta_{1,1} \mid \lambda_1, \dots, \lambda_i \in \mathbb{F}, \lambda_1 \neq 0\}.$$

Proof. Let $\phi = (\phi_{i,j}) \in \text{Aut}(\mu_0^n)$. Then, by (5.5), $\phi \cdot \Delta_{i,1} = \phi_{1,1}^{i+1}\Delta_{i,1} + \sum_{k=1}^{i-1} \phi_{i,k}\phi_{1,1}\Delta_{k,1}$ and as $\phi_{1,1} \neq 0$, the direct inclusion in the statement is proved.

Conversely, given $(\lambda_1, \dots, \lambda_i) \in \mathbb{F}^* \times \mathbb{F}^{i-1}$, Lemma 5.9 gives $(a_{1,1}, \dots, a_{i,1}) \in \mathbb{F}^* \times \mathbb{F}^{i-1}$ such that $\lambda_{i-k+1} = a_{i,k}a_{1,1}$, for all $k \in [i]$. Since $a_{1,1} \neq 0$, there exists $\phi \in \text{Aut}(\mu_0^n)$ such that $\phi_{k,1} = a_{k,1}$, for all $k \in [i]$. For any such $\phi \in \text{Aut}(\mu_0^n)$ and $j \in [i]$, we have

$$a_{i,j} = \sum_{k_1 + \dots + k_j = i} a_{k_1,1} \cdots a_{k_j,1} = \sum_{k_1 + \dots + k_j = i} \phi_{k_1,1} \cdots \phi_{k_j,1} = \phi_{i,j},$$

so $\phi \cdot \Delta_{i,1} = \sum_{k=1}^i \phi_{i,k}\phi_{1,1}\Delta_{k,1} = \sum_{k=1}^i a_{i,k}a_{1,1}\Delta_{k,1} = \sum_{k=1}^i \lambda_{i-k+1}\Delta_{k,1}$. This proves the reverse inclusion. \square

Recall that the ∇_j , for $j \in [n - 1]$, form a basis of $B^2(\mu_0^n, \mathbb{F})$.

Lemma 5.13. *Let $\phi = (\phi_{i,j}) \in \text{Aut}(\mu_0^n)$. In $H_{\text{LC}}^2(\mu_0^n, \mathbb{F})$ we have $\phi \cdot [\nabla_n] = \phi_{1,1}^{n+1}[\nabla_n]$.*

Proof. If $i + j > n + 1$, then $\Delta_{i,j}$ does not occur in $\phi \cdot \nabla_n$. Otherwise, for $i + j \leq n + 1$, the coefficient of $\Delta_{i,j}$ in the expression for $\phi \cdot \nabla_n$ is

$$\alpha_{i,j} = \sum_{k=i}^{n+1-j} \phi_{k,i}\phi_{n+1-k,j},$$

so that $\phi \cdot \nabla_n = \sum_{i+j \leq n+1} \alpha_{i,j}\Delta_{i,j}$.

Since $\text{Aut}(\mu_0^n)$ acts on $Z_{\text{LC}}^2(\mu_0^n, \mathbb{F})$ and on $B^2(\mu_0^n, \mathbb{F})$, we must have $\phi \cdot \nabla_n \in Z_{\text{LC}}^2(\mu_0^n, \mathbb{F}) \setminus B^2(\mu_0^n, \mathbb{F})$. Thus, by the proof of Proposition 5.1, we have $\alpha_{1,k} = \alpha_{2,k-1} = \dots = \alpha_{k-1,2}$ for

every $2 \leq k \leq n$. Moreover, by reparameterizing the summation index, we obtain $\alpha_{i,j} = \sum_{\ell=j}^{n+1-i} \phi_{n+1-\ell,i} \phi_{\ell,j} = \alpha_{j,i}$; in particular, $\alpha_{k,1} = \alpha_{1,k}$. It follows that

$$\phi \cdot \nabla_n = \sum_{k=1}^n \alpha_{1,k} \nabla_k.$$

Now, as $[\nabla_k] = 0$, for all $k \in [n-1]$, we get $\phi \cdot [\nabla_n] = \alpha_{1,n} [\nabla_n] = \phi_{1,1}^{n+1} [\nabla_n]$. \square

Proposition 5.14. *Assume that \mathbb{F} is algebraically closed of characteristic 0. Let $\mu \in \mathbb{F}^*$. We have*

$$\text{Orb}([\nabla_n + \mu \Delta_{n,1}]) = \left\{ \lambda_1 [\nabla_n] + \mu \lambda_1 [\Delta_{n,1}] + \sum_{k=2}^{n-1} \lambda_{n+1-k} [\Delta_{k,1}] \mid \lambda_1, \dots, \lambda_{n-1} \in \mathbb{F}, \lambda_1 \neq 0 \right\}.$$

Proof. Let $\phi = (\phi_{i,j}) \in \text{Aut}(\mu_0^n)$. Then, using Lemma 5.13 and recalling that $[\Delta_{1,1}] = [\nabla_1] = 0$,

$$\phi \cdot [\nabla_n + \mu \Delta_{n,1}] = \phi_{1,1}^{n+1} [\nabla_n] + \mu \phi_{n,n} \phi_{1,1} [\Delta_{n,1}] + \mu \sum_{k=2}^{n-1} \phi_{n,k} \phi_{1,1} [\Delta_{k,1}]. \quad (5.15)$$

Since $\phi_{n,n} = \phi_{1,1}^n$ and $\phi_{1,1} \neq 0$, the direct inclusion in the statement follows.

Conversely, let $\lambda_1, \dots, \lambda_{n-1} \in \mathbb{F}$, with $\lambda_1 \neq 0$. Consider the map ρ_{n-1} from Lemma 5.9. By that result, for any $\mu \in \mathbb{F}^*$, there are $(a_{1,1}, \dots, a_{n,1}) \in \mathbb{F}^* \times \mathbb{F}^{n-1}$ such that $\rho_{n-1}(a_{1,1}, \dots, a_{n,1}) = \left(\lambda_1, \frac{\lambda_2}{\mu}, \dots, \frac{\lambda_{n-1}}{\mu}, 0 \right)$. Let $\phi \in \text{Aut}(\mu_0^n)$ be determined by $\phi_{k,1} = a_{k,1}$, for all $k \in [n]$. Then $\phi_{i,j} = a_{i,j}$ for all $i, j \in [n]$. It follows that $\phi_{1,1}^{n+1} = \phi_{n,n} \phi_{1,1} = \lambda_1$ and similarly $\mu \phi_{n,k} \phi_{1,1} = \lambda_{n+1-k}$ for all $2 \leq k \leq n-1$. Thence, by (5.15),

$$\phi \cdot [\nabla_n + \mu \Delta_{n,1}] = \lambda_1 [\nabla_n] + \mu \lambda_1 [\Delta_{n,1}] + \sum_{k=2}^{n-1} \lambda_{n+1-k} [\Delta_{k,1}],$$

proving the reverse inclusion. \square

Proposition 5.16. *Assume that \mathbb{F} is algebraically closed of characteristic 0. For $2 \leq i \leq n-1$ and $\mu \in \mathbb{F}^*$ we have*

$$\text{Orb}([\nabla_n + \mu \Delta_{i,1}]) = \left\{ \lambda^{n+1} [\nabla_n] + \lambda^{i+1} \mu [\Delta_{i,1}] + \sum_{j=2}^{i-1} \lambda_{i+1-j} [\Delta_{j,1}] \mid \lambda_2, \dots, \lambda_{i-1} \in \mathbb{F}, \lambda \in \mathbb{F}^* \right\}.$$

Proof. Let $\phi = (\phi_{i,j}) \in \text{Aut}(\mu_0^n)$. Then,

$$\phi \cdot [\nabla_n + \mu \Delta_{i,1}] = \phi_{1,1}^{n+1} [\nabla_n] + \sum_{k=2}^i \phi_{i,k} \phi_{1,1} \mu [\Delta_{k,1}].$$

So, as before, to prove the result it suffices to assume that $i \geq 3$ and to show that given $\phi_{1,1}, \mu \in \mathbb{F}^*$, the elements $\phi_{i,k} \phi_{1,1} \mu$, with $2 \leq k \leq i-1$, can take on arbitrary values in \mathbb{F} , for appropriate choices of $\phi_{2,1}, \dots, \phi_{i-1,1} \in \mathbb{F}$.

Considering ρ_{i-2} , we know that for any $\lambda_2, \dots, \lambda_{i-1} \in \mathbb{F}$ there are $i + 1$ solutions to the equation

$$\rho_{i-2}(a_{1,1}, \dots, a_{i-1,1}) = \left(\phi_{1,1}^{i+1}, \frac{\lambda_2}{\mu}, \dots, \frac{\lambda_{i-1}}{\mu} \right).$$

As $a_{1,1}^{i+1} = \phi_{1,1}^{i+1}$, exactly one of the above solutions satisfies $a_{1,1} = \phi_{1,1}$ and the remainder of the proof goes as before. \square

The final ingredient in the proof of Theorem 5.7 explains the relevance of the factor group $\mathbb{F}_{(i,n)}$, defined in (5.6).

Lemma 5.17. *Fix $2 \leq i \leq n - 1$. For $\mu, \mu' \in \mathbb{F}^*$ we have*

$$\text{Orb}([\nabla_n + \mu\Delta_{i,1}]) = \text{Orb}([\nabla_n + \mu'\Delta_{i,1}]) \iff \bar{\mu} = \overline{\mu'},$$

where $\bar{\mu}, \overline{\mu'} \in \mathbb{F}_{(i,n)}$.

Remark 5.18. In view of this result, it makes sense to write $\text{Orb}([\nabla_n + \bar{\mu}\Delta_{i,1}])$, for $\mu \in \mathbb{F}^*$, and we can parameterize the orbits of the form above by the elements of $\mathbb{F}_{(i,n)}$.

Proof. For the direct implication, suppose that $[\nabla_n + \mu'\Delta_{i,1}] \in \text{Orb}([\nabla_n + \mu\Delta_{i,1}])$. Then there is some $\lambda \in \mathbb{F}^*$ such that $\lambda^{n+1} = 1$ and $\mu' = \lambda^{i+1}\mu$, so $\mu'/\mu \in R(i, n)$.

Conversely, if $\mu' = \lambda^{i+1}\mu$ for some $\lambda \in \mathbb{F}^*$ with $\lambda^{n+1} = 1$, then

$$[\nabla_n + \mu'\Delta_{i,1}] = \lambda^{n+1}[\nabla_n] + \lambda^{i+1}\mu[\Delta_{i,1}] \in \text{Orb}([\nabla_n + \mu\Delta_{i,1}]),$$

which shows that the respective orbits coincide. \square

We are now ready to conclude our main result of this section.

Proof of Theorem 5.7. Let $0 \neq \theta = \lambda_1[\nabla_n] + \sum_{j=2}^n \lambda_j[\Delta_{j,1}] \in \mathbf{H}_{\text{LC}}^2(\mu_0^n, \mathbb{F})$. Then:

- If $\lambda_1 = 0$, then $\theta \in \text{Orb}([\Delta_{i,1}])$, where $i = \max\{j \mid \lambda_j \neq 0\}$. (See Proposition 5.12.)
- If $\lambda_1 \neq 0$ and $\lambda_n \neq 0$, then $\theta \in \text{Orb}([\nabla_n + \mu\Delta_{n,1}])$, where $\mu = \lambda_n/\lambda_1 \neq 0$. (See Proposition 5.14.)
- If $\lambda_1 \neq 0$ and $\lambda_j = 0$ for all $2 \leq j \leq n$, then $\theta \in \text{Orb}([\nabla_n])$. (See Lemma 5.13.)
- If $\lambda_1 \neq 0$, $\lambda_n = 0$ and $\lambda_j \neq 0$ for some $2 \leq j \leq n - 1$, then $\theta \in \text{Orb}([\nabla_n + \bar{\mu}\Delta_{i,1}])$, where $i = \max\{2 \leq j \leq n - 1 \mid \lambda_j \neq 0\}$ and $\mu \in \mathbb{F}^*$ is defined as follows: choose some $\lambda \in \mathbb{F}^*$ such that $\lambda^{n+1} = \lambda_1$; then take $\mu = \lambda_i/\lambda^{i+1}$. (See Proposition 5.16.) By Lemma 5.17, $\bar{\mu}$ is uniquely determined by θ .

The fact that distinct elements given in the statement of the Theorem yield disjoint orbits follows easily from the description of the orbits in Propositions 5.12, 5.14, 5.16 and Lemmas 5.13, 5.17. \square

5.3 The orbit decomposition of $T_1(\mu_0^n)$ in the left commutative and bicommutative varieties

Observe that $\text{Ann}(\mu_0^n) = \mathbb{F}e_n$, so in order to get a non-split central extension of μ_0^n with 1-dimensional annihilator associated with a cocycle θ , we must have $e_n \notin \text{Ann}(\theta)$. This excludes the orbit representatives 0 and $\Delta_{i,1}$, for $i < n$. Notice also that for nonzero cocycles θ and θ' , the 1-dimensional spaces $\langle[\theta]\rangle$ and $\langle[\theta']\rangle$ are in the same $\text{Aut}(\mu_0^n)$ -orbit in the corresponding Grassmannian if and only if there is $\lambda \in \mathbb{F}^*$ such that $\lambda[\theta']$ is in the orbit of $[\theta]$. In particular, if the $\text{Aut}(\mu_0^n)$ -orbit of $[\theta]$ is closed under the multiplicative action of \mathbb{F}^* , then $\langle[\theta]\rangle$ and $\langle[\theta']\rangle$ are in the same orbit if and only if $[\theta]$ and $[\theta']$ are in the same orbit.

In our context, all orbits are closed under multiplication by nonzero scalars, except for the orbits of the form $\text{Orb}([\nabla_n + \bar{\mu}\Delta_{i,1}])$, with $2 \leq i \leq n-1$ and $\bar{\mu} \in \mathbb{F}_{(i,n)}$. Given $\epsilon \in \mathbb{F}^*$, Proposition 5.16 shows that $\epsilon^{n+1}[\nabla_n + \bar{\mu}\Delta_{i,1}] = [\nabla_n + \overline{\epsilon^{n-i}\bar{\mu}}\Delta_{i,1}]$. Since \mathbb{F} is algebraically closed, for a fixed $i \leq n-1$, ϵ^{n-i} can take on any nonzero value as $\epsilon \in \mathbb{F}^*$ varies. Hence, $[\nabla_n + \bar{\mu}\Delta_{i,1}]$ and $[\nabla_n + \bar{\mu}'\Delta_{j,1}]$ define 1-dimensional spaces in the same $\text{Aut}(\mu_0^n)$ -orbit of the Grassmannian if and only if $i = j$.

Combining all the results from Section 5, we obtain our main result below.

Theorem 5.19. *Assume that \mathbb{F} is algebraically closed of characteristic 0 and let $n \geq 2$. The following elements of $\mathbf{Z}_{\text{LC}}^2(\mu_0^n, \mathbb{F})$ parameterize the distinct orbits of $\text{Aut}(\mu_0^n)$ on the subspace $T_1(\mu_0^n)$ of the Grassmannian $G_1(\mathbf{H}_{\text{LC}}^2(\mu_0^n, \mathbb{F}))$, i.e. they parameterize the distinct isomorphism classes of non-split left commutative central extensions of the n -dimensional null-filiform algebra μ_0^n with 1-dimensional annihilator:*

- (a) $\Delta_{n,1}$;
- (b) $\{\nabla_n + \mu\Delta_{n,1} \mid \mu \in \mathbb{F}\}$ ($\mu = 0$ gives the trivial extension μ_0^{n+1});
- (c) $\{\nabla_n + \Delta_{i,1} \mid 2 \leq i \leq n-1\}$.

In the bicommutative case, the corresponding representatives are:

(Case $n = 2$)

- (a) $\Delta_{2,1}$;
- (b) $\{\nabla_2 + \mu\Delta_{2,1} \mid \mu \in \mathbb{F}\}$
($\mu = 0$ gives the trivial extension μ_0^3).

(Case $n > 2$)

- (a) ∇_n (trivial extension μ_0^{n+1});
- (b) $\nabla_n + \Delta_{2,1}$.

Remark 5.20. Note that Theorem 5.19 and the previous discussion only make reference to non-split central extensions with 1-dimensional annihilator. However, it is also possible to construct non-split left commutative central extensions of μ_0^n with 2-dimensional annihilator, whose isomorphism classes are parameterized by the cocycles $\Delta_{k,1}$ for $2 \leq k \leq n-1$. In the bicommutative case for $n > 2$, also $\Delta_{2,1}$ is a representative of an isomorphism class of non-split central extensions with 2-dimensional annihilator.

The explicit description of the multiplication in the central extensions referenced in Theorems 5.7 and 5.19, in Corollary 5.8 and in Remark 5.20 can be found in the following table. Only the nonzero products of the basis elements $\{e_1, \dots, e_{n+1}\}$ are displayed.

cocycle	multiplication, $i, j \in [n]$	
$\Delta_{n,1}$	$e_i e_j = e_{i+j}$ $e_n e_1 = e_{n+1}$	if $i + j \leq n$
$\Delta_{k,1}$ ($2 \leq k \leq n-1$)	$e_i e_j = e_{i+j}$ $e_k e_1 = e_{k+1} + e_{n+1}$	if $i + j \leq n$ and $(i, j) \neq (k, 1)$
$\nabla_n + \Delta_{k,1}$ ($2 \leq k \leq n-1$)	$e_i e_j = e_{i+j}$ $e_k e_1 = e_{k+1} + e_{n+1}$	if $i + j \leq n+1$ and $(i, j) \neq (k, 1)$
$\nabla_n + \mu \Delta_{n,1}$ ($\mu \in \mathbb{F}$)	$e_i e_j = e_{i+j}$ $e_n e_1 = (1 + \mu)e_{n+1}$	if $i + j \leq n+1$ and $i \neq n$ ($\mu = 0$ gives the trivial extension)

Table 1: Isomorphism classes of 1-dimensional non-split left commutative and bicommutative central extensions of the n -dimensional null-filiform algebra μ_0^n .

6 Assosymmetric, Novikov, and left symmetric central extensions

6.1 Assosymmetric central extensions

Recall that an algebra A is said to be assosymmetric if it satisfies the identities

$$(xy)z - x(yz) = (yx)z - y(xz) = (xz)y - x(zy),$$

for all $x, y, z \in A$. Now, to find the assosymmetric central extensions of μ_0^n , we need cocycles $\theta: \mu_0^n \times \mu_0^n \rightarrow \mathbb{F}$ satisfying

$$\theta(e_i e_j, e_k) - \theta(e_i, e_j e_k) = \theta(e_j e_i, e_k) - \theta(e_j, e_i e_k) = \theta(e_i e_k, e_j) - \theta(e_i, e_k e_j),$$

for all $i, j, k \in [n]$. Note that for the algebra μ_0^n , these two equalities reduce to

$$\theta(e_i, e_{j+k}) = \theta(e_j, e_{i+k}) \quad \text{and} \quad \theta(e_{i+j}, e_k) = \theta(e_{i+k}, e_j),$$

for all $i, j, k \in [n]$, with the usual convention that $e_m = 0$ if $m > n$. This means that every assosymmetric cocycle of μ_0^n is in $Z_{\text{BC}}^2(\mu_0^n, \mathbb{F})$. On the other hand, it is easy to see that every element from $Z_{\text{BC}}^2(\mu_0^n, \mathbb{F})$ is an assosymmetric cocycle.

6.2 Novikov central extensions

Recall that an algebra A is said to be Novikov if it satisfies the identities

$$(xy)z = (xz)y \quad \text{and} \quad (xy)z - x(yz) = (yx)z - y(xz),$$

for all $x, y, z \in A$. Now, to find all the Novikov central extensions of μ_0^n , we require cocycles $\theta: \mu_0^n \times \mu_0^n \rightarrow \mathbb{F}$ satisfying

$$\theta(e_i e_j, e_k) = \theta(e_i e_k, e_j) \quad \text{and} \quad \theta(e_i e_j, e_k) - \theta(e_i, e_j e_k) = \theta(e_j e_i, e_k) - \theta(e_j, e_i e_k),$$

for all $i, j, k \in [n]$. For the algebra μ_0^n , these identities reduce to

$$\theta(e_{i+j}, e_k) = \theta(e_{i+k}, e_j) \quad \text{and} \quad \theta(e_i, e_{j+k}) = \theta(e_j, e_{i+k}),$$

with $e_m = 0$ if $m > n$. As these are exactly the identities for the bicommutative cocycles of $Z_{\text{BC}}^2(\mu_0^n, \mathbb{F})$, this case also reduces to the bicommutative case.

6.3 Left symmetric central extensions

Recall that an algebra A is said to be left symmetric if it satisfies the identity

$$(xy)z - x(yz) = (yx)z - y(xz),$$

for all $x, y, z \in A$. To find all the left symmetric central extensions of μ_0^n , we need cocycles $\theta: \mu_0^n \times \mu_0^n \rightarrow \mathbb{F}$ satisfying

$$\theta(e_i e_j, e_k) - \theta(e_i, e_j e_k) = \theta(e_j e_i, e_k) - \theta(e_j, e_i e_k),$$

for all $i, j, k \in [n]$. Note that for the algebra μ_0^n , we obtain just the relation

$$\theta(e_i, e_{j+k}) = \theta(e_j, e_{i+k}), \quad \text{for all } i, j, k \in [n],$$

where $e_m = 0$ if $m > n$. Thus, θ is the same as a left commutative cocycle from $Z_{\text{LC}}^2(\mu_0^n, \mathbb{F})$, and it follows that this case reduces to the left commutative case.

References

- [1] Abdelwahab H., Calderón A.J., Kaygorodov I., The algebraic and geometric classification of nilpotent binary Lie algebras, *International Journal of Algebra and Computation*, 29 (2019), 6, 1113–1129.
- [2] Adashev J., Camacho L., Omirov B., Central extensions of null-filiform and naturally graded filiform non-Lie Leibniz algebras, *Journal of Algebra*, 479 (2017), 461–486.
- [3] Burde D., Dekimpe K., Deschamps S., LR-algebras. New developments in Lie theory and geometry, 125–140, *Contemp. Math.*, 491 (2009), Amer. Math. Soc., Providence, RI, 2009.
- [4] Calderón Martín A., Fernández Ouaridi A., Kaygorodov I., The classification of n -dimensional anticommutative algebras with $(n - 3)$ -dimensional annihilator, *Communications in Algebra*, 47 (2019), 1, 173–181.
- [5] Cical S., De Graaf W., Schneider C., Six-dimensional nilpotent Lie algebras, *Linear Algebra and its Applications*, 436 (2012), 1, 163–189.
- [6] Darijani I., Usefi H., The classification of 5-dimensional p -nilpotent restricted Lie algebras over perfect fields, I, *Journal of Algebra*, 464 (2016), 97–140.
- [7] De Graaf W., Classification of 6-dimensional nilpotent Lie algebras over fields of characteristic not 2, *Journal of Algebra*, 309 (2007), 2, 640–653.

- [8] De Graaf W., Classification of nilpotent associative algebras of small dimension, *International Journal of Algebra and Computation*, 28 (2018), 1, 133–161.
- [9] Dekimpe K., Ongenaes V., Filiform left-symmetric algebras, *Geometriae Dedicata*, 74 (1999), no. 2, 165–199.
- [10] Fernández Ouaridi A., Kaygorodov I., Khrypchenko M., Volkov Yu., Degenerations of nilpotent algebras, arXiv:1905.05361
- [11] Gorshkov I., Kaygorodov I., Khrypchenko M., The algebraic classification of Tortkara algebras, arXiv:1904.00845.
- [12] Hegazi A., Abdelwahab H., Classification of five-dimensional nilpotent Jordan algebras, *Linear Algebra and its Applications*, 494 (2016), 165–218.
- [13] Hegazi A., Abdelwahab H., The classification of n -dimensional non-associative Jordan algebras with $(n - 3)$ -dimensional annihilator, *Communications in Algebra*, 46 (2018), 2, 629–643.
- [14] Hegazi A., Abdelwahab H., Calderón Martín A., The classification of n -dimensional non-Lie Malcev algebras with $(n - 4)$ -dimensional annihilator, *Linear Algebra and its Applications*, 505 (2016), 32–56.
- [15] Hegazi A., Abdelwahab H., Calderón Martín A., Classification of nilpotent Malcev algebras of small dimensions over arbitrary fields of characteristic not 2, *Algebras and Representation Theory*, 21 (2018), 1, 19–45.
- [16] Karimjanov I., Kaygorodov I., Khudoyberdiyev K., The algebraic and geometric classification of nilpotent Novikov algebras, *Journal of Geometry and Physics*, 143 (2019), 11–21.
- [17] Karimjanov I., Kaygorodov I., Ladra M., Central extensions of filiform associative algebras, *Linear and Multilinear Algebra*, (2019), DOI: 10.1080/03081087.2019.1620674
- [18] Kaygorodov I., Páez-Guillán P., Voronin V., The algebraic and geometric classification of nilpotent bicommutative algebras, *Algebras and Representation Theory*, 2020, DOI: 10.1007/s10468-019-09944-x
- [19] Ladra M., Shahryari M., Zargheh C., HNN-extensions of Leibniz algebras, *Journal of Algebra*, 532 (2019), 183–200.
- [20] Lichtman A. I., Shirvani M., HNN-extensions of Lie algebras, *Proceedings of the American Mathematical Society*, 125 (1997), 12, 3501–3508.
- [21] Skjelbred T., Sund T., Sur la classification des algèbres de Lie nilpotentes, *Comptes Rendus Mathématique Académie des Sciences, Paris. Ser. A-B*, 286 (1978), 5, A241–A242.
- [22] Wasserman A., A derivation HNN construction for Lie algebras, *Israel Journal of Mathematics*, 106 (1998), 76–92.
- [23] Zusmanovich P., Central extensions of current algebras, *Transactions of the American Mathematical Society*, 334 (1992), 1, 143–152.