

High probability decoupling via approximate unitary designs and efficient relative thermalisation

Aditya Nema and Pranab Sen

Abstract

We prove a new concentration result for non-catalytic decoupling by showing that, for suitably large t , applying a unitary chosen uniformly at random from an approximate t -design on a quantum system followed by a fixed quantum operation almost decouples, with high probability, the given system from another reference system to which it may initially have been correlated. Earlier works either did not obtain high decoupling probability, or used provably inefficient unitaries, or required catalytic entanglement for decoupling. In contrast, our approximate unitary designs always guarantee decoupling with exponentially high probability and, under certain conditions, lead to computationally efficient unitaries. As a result we conclude that, under suitable conditions, efficiently implementable approximate unitary designs achieve relative thermalisation in quantum thermodynamics with exponentially high probability. We also show the scrambling property of black hole, when the black hole evolution is according to pseudorandom approximate unitary t -design, as opposed to the Haar random evolution considered earlier by Hayden-Preskill.

I. INTRODUCTION

A peculiar characteristic of quantum information theory is that many information transmission protocols, be it compression of quantum messages or sending quantum information through unassisted quantum channels, can be constructed by first removing correlations of a particular system from some other system around it. This behooves us to prove general theorems that take a bipartite quantum state shared between a system A (e.g. the “particular system” above) and a reference R (e.g. the “some other system” above), apply a local operation on A , and then, if suitable conditions are met, prove that the resulting state is close to a product state between the output system B and the untouched reference R . This process of removing quantum correlations, i.e. obtaining a state close to a product state, is referred to as decoupling. Decoupling theorems play a vital role in proving achievability bounds for several quantum information theory protocols as well as thermalization results in quantum thermodynamics. In particular, the so-called Fully Quantum Slepian Wolf (FQSW) protocol [1], which has been hailed as the mother protocol of quantum information theory, is constructed via a decoupling argument. In the FQSW problem, the system A is thought of as a bipartite system $A = A_1 \otimes A_2$ and the fixed superoperator is nothing but tracing out A_2 . The FQSW protocol is used as a building block for many other important protocols in quantum information theory in the asymptotic iid setting e.g. noisy teleportation, noisy super

A. Nema is currently with the Institute of Quantum Information, RWTH Aachen University, Germany. The first draft of this work was submitted while pursuing PhD at School of Technology and Computer Science, Tata Institute of Fundamental Research, Mumbai and first two revisions were submitted during postdoc at the Department of Mathematical Informatics, Nagoya University, Japan. e-mail: aditya.nema.30@gmail.com.

P. Sen is currently with Centre for Quantum Technologies, National University of Singapore and School of Technology and was at Computer Science, Tata Institute of Fundamental Research, Mumbai India during earlier submissions. e-mail: pranab.sen.73@gmail.com.

Manuscript received June 13, 2021.

dense coding, distributed compression, entanglement unassisted and assisted quantum channel coding, one way entanglement distillation, reverse Shannon theorem etc. Asymptotic iid setting means that the given messages / channels are of the tensor power form $(\cdot)^{\otimes n}$ for large n . However the basic decoupling and FQSW results are actually one-shot results where the given message / channel is to be used only once. The one shot FQSW result can be extended to obtain a one-shot relative thermalization result in quantum thermodynamics [2], where a system $\Omega \leq S \otimes E$, with S being the subsystem of physical interest and E being the so-called ‘environment’ or ‘bath’ subsystem, initially starts out in a correlated state together with a reference system R but very soon evolves into something close to a so-called ‘relative thermal state’ on S tensored with the reduced state on the reference R .

In this paper, we build on the following important decoupling theorem proved by Dupuis in his doctoral thesis [3].

Fact 1. Consider a quantum state ρ^{AR} shared between a system A and a reference R . Let $\mathcal{T}^{A \rightarrow B}$ be a completely positive trace preserving superoperator (aka CPTP map aka quantum operation) with input system A and output system B . Let U be a unitary on the system A . Define the function

$$f(U) := \|(\mathcal{T}^{A \rightarrow B} \otimes \mathbb{I}^R)((U^A \otimes I^R)\rho^{AR}(U^{A\dagger} \otimes I^R)) - \omega^B \otimes \rho^R\|_1,$$

where \mathbb{I}^R is the identity superoperator on R and I^R is the identity operator on R . Let A' be a new system having the same dimension as A . Define the Choi-Jamiołkowski state $\omega^{A'B} := (\mathcal{T}^{A \rightarrow B} \otimes \mathbb{I}^{A'})(|\Phi\rangle\langle\Phi|^{AA'})$, of $\mathcal{T}^{A \rightarrow B}$ where $|\Phi\rangle^{AA'} := |A|^{-1/2} \sum_a |a\rangle^A \otimes |a\rangle^{A'}$ is the standard normalized EPR state on system AA' . Then, $\mathbb{E}_{U \sim \text{Haar}}[f(U)] \leq 2^{-\frac{1}{2}H_2(A|R)_\rho - \frac{1}{2}H_2(A'|B)_\omega}$, where the expectation is taken over the Haar measure on unitary operators on A . The quantity $H_2(\cdot|\cdot)$ is the conditional Rényi 2-entropy defined in Definition 1 below. We remark that $H_2(A|R)_\rho = -2\log\|\tilde{\rho}^{AR}\|_2$ and $H_2(A'|B)_\omega = -2\log\|\tilde{\omega}^{A'B}\|_2$, where $\tilde{\rho}^{AR}$ and $\tilde{\omega}^{A'B}$ are certain positive semidefinite matrices defined in Definition 1.

Informally speaking, the above theorem states that if some entropic conditions are met then, in expectation, the state σ^{BR} obtained by first applying a Haar random unitary U^A on the initial state ρ^{AR} followed by a CPTP map $\mathcal{T}^{A \rightarrow B}$ is close to the decoupled state $\omega^B \otimes \rho^R$. Here, $\omega^B = \mathcal{T}^{A \rightarrow B}(\frac{I^A}{|A|})$ is the state obtained by applying \mathcal{T} to the completely mixed state on A . In fact, $\omega^{A'B}$ defined in Fact 1 above is nothing but the Choi-Jamiołkowski state corresponding to the CPTP map $\mathcal{T}^{A \rightarrow B}$. Intuitively, a Haar random unitary U^A ‘randomizes’ or ‘scrambles’ the state on A to give the completely mixed state which is then sent to ω^B by $\mathcal{T}^{A \rightarrow B}$. So it is reasonable to believe that the local state on B should be ω^B . Notice that the local state on R after applying U^A and $\mathcal{T}^{A \rightarrow B}$ is always ρ^R . The punch of the decoupling theorem is that the global state is close to the desired tensor product state.

The distance of the actual global state from the desired tensor product state is upper bounded by two quantities. The first quantity $H_2(A|R)_\rho$ is usually negative, which signifies that A and R are entangled in the initial state ρ . To decouple A from R we start by applying a Haar random unitary U to the system A . A single unitary cannot decouple A from R , and that is why the decoupling theorem above also has the CPTP map \mathcal{T} . Now in an intuitive sense, the EPR state $\Phi^{AA'}$ is the ‘most entangled state’. So if a Haar random unitary U on the system A of $\Phi^{AA'}$ followed by the CPTP map \mathcal{T} can decouple the output system B from R , then it must be able to decouple B from R when the input is any entangled state ρ^{AR} , provided that the ‘amount of entanglement’ $H_2(A|R)_\rho$ between A and R in ρ is less than the ‘amount of entanglement’ $H_2(A'|B)_\omega$ between A' and B in ω . This explains the expression $H_2(A'|B)_\omega + H_2(A|R)_\rho$ in the above upper bound. To counteract a negative $H_2(A|R)_\rho$, the quantity $H_2(A'|B)_\omega$ had better be positive which signifies that A' is mostly decoupled from B in the state ω .

Dupuis showed in his doctoral thesis how the decoupling theorem above can be used to recover in a unified fashion several previously known results as well as obtain some totally new results in quantum

information theory. Szehr et al. [4] extended the decoupling theorem by showing that the expectation can be taken over approximate unitary 2-designs (defined formally in Definition 11 below) instead of over Haar random unitaries. The advantage of unitary 2-designs is that efficient constructions for them exist unlike the case with Haar random unitaries. Szehr et al. also upper bounded the expected trace distance in terms of smooth entropic quantities which have better mathematical properties compared to the non-smooth ones. In particular, in the asymptotic iid limit, the smooth entropic quantities are suitably bounded by n times the corresponding Shannon entropies which is not the case with the non-smooth quantities. Their result (adapted to our notations) is stated below.

Fact 2. *Under the setting of Fact 1 above,*

$$\mathbb{E}_{U^A \sim \text{Haar}}[f(U)] \leq 2^{-\frac{1}{2}H_2^\epsilon(A|R)_\rho - \frac{1}{2}H_2^\epsilon(A'|B)_\omega} + 12\epsilon,$$

where the expectation is taken over the Haar measure on unitary operators on A . The same result holds if the expectation is taken over the uniform choice of a unitary from an exact 2-design. If the expectation is taken over the uniform choice of a unitary from a δ -approximate 2-design, the upper bound gets multiplied by a multiplicative factor dependent on the dimension of A and δ . The smooth conditional Rényi 2-entropy terms appearing in the bound are defined in [3].

In a different vein Anshu and Jain [5] showed, extending earlier work by Ambainis and Smith [6], that it is possible to add a small ancilla C in tensor product with A , apply an efficient unitary to $A \otimes C$ and then trace out C so that A is now decoupled from R even before applying the CPTP map \mathcal{T} . The difference between Ambainis and Smith's or Anshu and Jain's works on one hand, and Dupuis', Szehr et al.'s or our works on the other hand is that we want a single unitary on the system A to achieve decoupling and not the average of a number of unitaries on A or, more generally, a unitary on a larger system $A \otimes C$. A single unitary cannot decouple A from R . That is why the decoupling theorem above also has the CPTP map \mathcal{T} . The single-unitary-followed-by-CPTP-map form of the decoupling theorem is required for quantum Shannon theory applications where there is no entanglement assistance e.g. sending quantum information over an unassisted quantum channel.

After obtaining the decoupling result in expectation above, it is natural to ask whether such a theorem also holds with high probability over the choice of the random unitary U^A . Dupuis [3] answered this question in the affirmative for the Haar measure. That result, adapted to our notation, is as follows:

Fact 3. *Under the setting of Fact 1 above, we have*

$$\mathbb{P}_{U^A \sim \text{Haar}}[f(U) > 2^{-\frac{1}{2}H_2^\epsilon(A|R)_\rho - \frac{1}{2}H_2^\epsilon(A'|B)_\omega} + 16\epsilon + \delta] \leq 2 \exp\left(-\frac{|A|\delta^2}{2^{H_{\min}^\epsilon(A)_\rho + 4}}\right),$$

where the smooth min-entropy $H_{\min}^\epsilon(\cdot)$ is defined in Definition 2 below, and the probability is taken over the Haar measure on \mathbb{U}^A .

The concentration of measure result for the decoupling theorem above immediately implies an exponential concentration result for the FQSW problem, which further implies that relative thermalization occurs for a system in contact with a heat bath for all but an exponentially small fraction of unitary evolutions of the system as long as the system is assumed to evolve according to a Haar random unitary. However this is not a very satisfactory explanation from a physical and computational point of view as Haar random unitaries are provably impossible to implement by quantum circuits with size polylogarithmic in the dimension of the system. Also, Haar random unitaries on a system A require $\Omega(|A|^2 \log |A|)$ number of random bits for a precise description. This leads us to wonder if relative thermalization can be achieved with high probability by simpler unitary evolutions of the system A . Nakata et al. [7] gave an

affirmative answer by showing that decoupling can be achieved by choosing products of random unitaries diagonal in the Pauli X and Z bases, but even they are not efficiently implementable and in addition, require $\Omega(|A| \log |A|)$ random bits for a precise description. Moreover the fraction of such unitaries which achieve decoupling is not strongly concentrated near one.

A. Our results

In this paper we prove for the first time that, for suitable values of t , *approximate unitary t -designs achieve decoupling with probability exponentially close to one*. An exact t -design of $n \times n$ unitaries can be described using $O(t \log n)$ random bits [8] as opposed to $\Omega(n^2 \log n)$ random bits required to describe a Haar random unitary to reasonable precision. Thus for many applications our result implies a substantial saving in the number of random bits compared to Dupuis' result. However, the concentration guaranteed by our result is less than that guaranteed by Dupuis even though it is exponential. Our concentration bound for decoupling via unitary designs is expressed in terms of smooth entropic quantities. An informal version is stated below:

Result 1. *Consider a quantum state ρ^{AR} shared between a system A and a reference R . Let $\mathcal{T}^{A \rightarrow B}$ be a completely positive trace preserving superoperator with input system A and output system B . Let U be a unitary on the system A . Define the function*

$$f(U) := \|(\mathcal{T}^{A \rightarrow B} \otimes \mathbb{I}^R)((U^A \otimes I^R)\rho^{AR}(U^{A\dagger} \otimes I^R)) - \omega^B \otimes \rho^R\|_1,$$

where \mathbb{I}^R is the identity superoperator on R and I^R is the identity operator on R . Let A' be a new system having the same dimension as A . Define the Choi-Jamiołkowski state $\omega^{A'B} := (\mathcal{T}^{A \rightarrow B} \otimes \mathbb{I}^{A'}) (|\Phi\rangle\langle\Phi|^{AA'})$, of $\mathcal{T}^{A \rightarrow B}$ where $|\Phi\rangle^{AA'} := |A|^{-1/2} \sum_a |a\rangle^A \otimes |a\rangle^{A'}$ is the standard EPR state on system AA' . Let $0 < \epsilon, \delta < 1/3$. Let $\kappa > 0$. Then,

$$\mathbb{P}_{U \sim \text{design}}[f(U) > 2^{-\frac{1}{2}H_2^\epsilon(A|R)_\rho - \frac{1}{2}\mathbb{H}_2^{\epsilon,\delta}(A'|B)_\omega + 1} + 14\sqrt{\epsilon} + 2\kappa] \leq 7 \cdot 2^{-a\kappa^2}. \quad (1)$$

where the matrix U^A is chosen uniformly at random from an approximate t -design of unitaries, $a := |A| \cdot 2^{-(1+\delta)\mathbb{H}_{\max}^\epsilon(B)_\omega + H_2^\epsilon(A|R)_\rho - 9}$, $t := 8a\kappa^2$. The smooth entropies $\mathbb{H}_{\max}^\epsilon(\cdot)$, $\mathbb{H}_2^{\epsilon,\delta}(\cdot)$, $H_2^\epsilon(\cdot)$ are defined in Definitions 3, 4, 1 respectively below. Since the result holds for all $\kappa > 0$, which is the parameter for deviation from the expectation value of $f(U)$, one can replace κ in terms of the parameter t of our approximate unitary t -design by $\sqrt{\frac{t}{8a}}$. This can be done once the analysis is done for a particular value of $\kappa > 0$. Further, we chose parameter $\beta := \sqrt{\frac{1}{a}}$. We thus get probability of the decoupling Equation 1 exponential in ' t ', as:

$$\mathbb{P}_{U \sim \text{design}}[f(U) > 2^{-\frac{1}{2}H_2^\epsilon(A|R)_\rho - \frac{1}{2}\mathbb{H}_2^{\epsilon,\delta}(A'|B)_\omega + 1} + 14\sqrt{\epsilon} + \sqrt{t/2}\beta] \leq 7 \cdot 2^{-\frac{t}{8}}. \quad (2)$$

The three smooth one-shot entropic terms used in Result 1 approach the standard Shannon entropic terms in the asymptotic iid limit.

Remark 1. *Note that Equations 1 and 2 are exactly the same. Moreover, Equation 1 represents a measure concentration result or the so-called tail probability (commonly used for large deviation analysis) for the decoupling function f . Equation 2 provides a more intuitive relation of the decoupling phenomenon with approximate unitary t -designs. This way of writing also indicates that we have exponentially high concentration in the decoupling theorem, exponential in t .*

We can thus infer the following corollary of our main result in the asymptotic iid setting. We state an informal version of the corollary below.

Result 2. Consider the setting of Result 1 above. Let n be a large enough positive integer. Consider the n -fold tensor powers $\omega^{(A')^n B^n} := (\omega^{A'B})^{\otimes n}$, $\rho^{A^n R^n} := (\rho^{AR})^{\otimes n}$. Let $\epsilon' := 8(n + |A||B|)^{|A||B|}\epsilon^{1/4}$. Let $\kappa > 0$. Then,

$$\mathbb{P}_U \left[f(U) > 2^{-\frac{n}{2}(H(A|R)_\rho - \delta(3H(AR)_\rho + 7H(R)_\rho)) - \frac{n}{2}(H(A'|B)_\omega - \delta(3H(A'B)_\omega + 7H(B)_\omega))} + 28(\epsilon')^{1/4} + 2\kappa \right] \leq 7 \cdot 2^{-a\kappa^2},$$

which can also be expressed as (following the explanation from Remark 1):

$$\mathbb{P}_U \left[f(U) > 2^{-\frac{n}{2}(H(A|R)_\rho - \delta(3H(AR)_\rho + 7H(R)_\rho)) - \frac{n}{2}(H(A'|B)_\omega - \delta(3H(A'B)_\omega + 7H(B)_\omega))} + 28(\epsilon')^{1/4} + \sqrt{t/2}\beta \right] \leq 7 \cdot 2^{-\frac{t}{8}},$$

where the unitary U^{A^n} is chosen uniformly at random from an approximate t -design,

$$\begin{aligned} a &:= |A|^n \cdot 2^{n(H(A|R)_\rho - \delta(3H(AR)_\rho + 7H(R)_\rho)) - nH(B)_\omega(1+7\delta) - 9}, \\ t &:= |A|^n \kappa^2 2^{n(H(A|R)_\rho + 32\sqrt{\epsilon'}) + \log(\epsilon')^{-1} - nH(B)_\omega(1-5\delta) - 6}, \\ \beta &:= \sqrt{\frac{1}{a}}. \end{aligned}$$

The proof of our main result and the analysis of its iid limit requires us to define two novel one-shot entropic quantities that we call *smooth modified conditional Rényi 2-entropy* $\mathbb{H}_2^{\epsilon, \delta}(\cdot|\cdot)$ and *smooth modified max-entropy* $\mathbb{H}_{\max}^\epsilon(\cdot)$. Their definitions and techniques used in our proofs should be of independent interest.

Our concentration result for decoupling immediately implies that approximate unitary $|A_1|$ -designs decouple a quantum system in the Fully Quantum Slepian Wolf (FQSW) theorem with probability $1 - \exp(-\Theta(|A_1|))$, where the system A is expressed as a tensor product $A_1 \otimes A_2$ and the superoperator simply traces out A_2 .

Result 3 (Partial trace (or FQSW) concentration under design). Consider the setting of Result 1. Consider the FQSW decoupling function

$$f(U) = f_{\text{FQSW}}(U^{A_1 A_2}) := \|\text{Tr}_{A_2}[(U^{A_1 A_2} \otimes I^R) \circ \rho^{A_1 A_2 R}] - \pi^{A_1} \otimes \rho^R\|_1.$$

Let $H_2^\epsilon(A|R) = -\log \|(\tilde{\rho}')^{AR}\|_2^2$. Suppose we are promised that $\|(\tilde{\rho}')^R\|_2^2 < 0.9|A_1||A_2|\|(\tilde{\rho}')^{AR}\|_2^2$, $|A_1| \geq 2$, $|A_2| > |A_1|$ and $|A_2|2^{H_2^\epsilon(A_1 A_2|R)_\rho - 8} - 4 > 2\log|A_1| + 3\log|A_2|$. Let $\kappa > 0$. The following concentration inequality holds:

$$\mathbb{P}_{U \sim \text{design}}[f(U) > \sqrt{\frac{|A_1|}{|A_2|}} \cdot 2^{-\frac{1}{2}H_2^\epsilon(A|R)_\rho + 1} + 14\sqrt{\epsilon} + 2\kappa] \leq 7 \cdot 2^{-a\kappa^2}, \quad (3)$$

where the unitary U^A is chosen uniformly at random from an approximate t -design, $a := |A_2|2^{H_2^\epsilon(A|R)_\rho - 9}$ and $t := 8a\kappa^2$. Moreover, if $|A_1| \leq \text{polylog}(|A_2|)$ and $\kappa = \sqrt{\frac{|A_1|}{|A_2|}} \cdot 2^{-\frac{1}{2}H_2^\epsilon(A|R)_\rho + O(1)}$ which further

implies that $t = O(|A_1|)$, then efficient constructions for such approximate t -designs exist and the concentration Equation 3 can also be expressed simply as:

$$\mathbb{P}_{U \sim \text{design}}[f(U) > \sqrt{\frac{|A_1|}{|A_2|}} \cdot 2^{-\frac{1}{2}H_2^\epsilon(A|R)_\rho+1} + 14\sqrt{\epsilon} + \beta \sqrt{|A_1|/2}] \leq 7 \cdot 2^{-\frac{|A_1|}{2}}, \quad (4)$$

where $\beta := \sqrt{1/a}$.

The statement just above Result 3 can be obtained by setting $\kappa = \sqrt{\frac{|A_1|}{|A_2|}} \cdot 2^{-\frac{1}{2}H_2^\epsilon(A|R)_\rho+O(1)}$. This immediately leads to a saving in the number of random bits to $O(|A_1| \log(|A_1||A_2|))$ for approximate t -design from $\Omega(|A_1|^2|A_2|^2 \log(|A_1||A_2|))$ required by Haar random unitaries. If $|A_1| = \text{polylog}|A_2|$, then efficient algorithms exist for implementing approximate unitary $|A_1|$ -designs [9], [10]. Thus, for small values of $|A_1|$ our result shows that FQSW decoupling can indeed be achieved by efficiently implementable unitaries with probability exponentially close to one. This result can be extended to show that *for small systems S , relative thermalisation can be achieved by efficiently implementable unitaries with probability exponentially close to one for a wide range of parameters, the first result of this kind.*

We remark that the task of replacing Haar random unitary operator via randomly chosen unitary operator from an approximate unitary design is fairly non-trivial in the case of the decoupling theorem that we consider in this work. This is because of the following reasons:

- 1) The function $f(U)$ in Result 1 above is not a polynomial in the entries of the unitary operator U . So, even though Dupuis proved a concentration result for $f(U)$ under Haar measure in Fact 3, a similar statement for U chosen uniformly from a unitary design is not straightforward. Hence we upper bound $f(U)$ by a function $g^2(U)$, defined in Section I-B below using Fact 12, which is a polynomial in entries of U . The methodology of replacing Haar measure with unitary design can be applied to $g^2(U)$. However we first have to prove a concentration result for $g(U)$ under the Haar measure, which calls for the evaluation of a ‘good’ Lipschitz constant of $g(U)$, which is another challenging task that we carry out here. We then have to prove a concentration result for $g^2(U)$ from the concentration result for $g(U)$, which is also a non-trivial task.
To have an analogous ‘tight’ Lipschitz constant in asymptotic iid limit so that we can recover Dupuis result, we need to define new smooth max entropy! Known definition of smoothed max entropy does not suffice for this purpose due to weighting operators in Cauchy-Schwarz.
- 2) In order to define $g(U)$ appropriately, we have to perturb the CPTP map $\mathcal{T}^{A \rightarrow B}$ to a CP map $\hat{\mathcal{T}}^{A \rightarrow B}$ (see Equation 8 for precise definition) in the diamond norm in order to obtain tail bounds involving smooth conditional entropies. This ensures that, for any input state ρ^{AR} , the operator $(\mathcal{T}^{A \rightarrow B} \otimes \mathbb{I}^R)(\rho^{AR})$ is close to the operator $(\hat{\mathcal{T}}^{A \rightarrow B} \otimes \mathbb{I}^R)(\rho^{AR})$ in the trace distance. The smooth conditional entropies defined in earlier works like [3] and [4] do not quite suffice for this purpose; they only manage to show that the positive semidefinite matrices obtained by applying CP maps $\mathcal{T}^{A \rightarrow B}$, $\hat{\mathcal{T}}^{A \rightarrow B}$ to a certain ‘averaged state’ are close. Additionally, in order to obtain a good Lipschitz constant for $g(U)$, we have to cleverly design the weighting operator arising from the weighted Cauchy-Schwarz inequality required to upper bound Schatten 1-norm of an operator with its Schatten 2-norm. We also want the smooth one-shot entropic quantities to approach their natural Shannon entropic analogues in the asymptotic iid regime. It is challenging to meet all three requirements simultaneously, and for this we need to define a *novel one-shot smooth conditional modified Rényi 2-entropy*.

Addressing the above two issues forms the new technical advancement towards the decoupling literature.

B. Proof technique

We now give a high level description of the proof of our main result. For a unitary U on the system A , we define the value taken by the decoupling function at U as follows:

$$f(U) := \|(\mathcal{T}^{A \rightarrow B} \otimes \mathbb{I}^R)((U^A \otimes I^R)\rho^{AR}(U^{A\dagger} \otimes I^R)) - \omega^B \otimes \rho^R\|_1.$$

We wish to prove a tail bound for $f(U)$ where U is chosen uniformly from a unitary design. For this, it is easier to first prove a tail bound for a related function $g(U)$:

$$g(U) := \|((\tilde{\mathcal{T}}')^{A \rightarrow B} \otimes \mathbb{I}^R)((U^A \otimes I^R)(\tilde{\rho}')^{AR}(U^{A\dagger} \otimes I^R)) - (\tilde{\omega}')^B \otimes (\tilde{\rho}')^R\|_2,$$

where $(\tilde{\mathcal{T}}')^{A \rightarrow B}$, $(\tilde{\rho}')^{AR}$, $(\tilde{\omega}')^{A'B}$ will be defined formally, later in Section III. For the discussion in this section, consider $\tilde{\mathcal{T}}'$ to be a perturbed version of \mathcal{T} within a distance of $O(\sqrt{\epsilon})$, $(\tilde{\rho}')^{AR} = 2^{-\frac{1}{2}\mathbb{H}_2^{(\epsilon, \delta)}(A|R)_\rho}$ and $(\tilde{\omega}')^{A'B} = 2^{-\frac{1}{2}\mathbb{H}_2^\epsilon(A|B)_\omega}$. We will have, for all probability distributions on U^A ,

$$\begin{aligned} \mathbb{P}_{U^A}[f(U) > 2^{-\frac{1}{2}H_2^\epsilon(A|R)_\rho - \frac{1}{2}\mathbb{H}_2^{\epsilon, \delta}(A'|B)_\omega + 1} + 14\sqrt{\epsilon} + 2\theta] \\ \leq \mathbb{P}_{U^A}[g(U) > \|(\tilde{\rho}')^{AR}\|_2 \cdot \|(\tilde{\omega}')^{A'B}\|_2 + \theta]. \end{aligned}$$

We then bound $\mathbb{P}_{U^A}[g(U) > \|(\tilde{\rho}')^{AR}\|_2 \cdot \|(\tilde{\omega}')^{A'B}\|_2 + \theta]$ where U^A is chosen according to the Haar measure. For this we need to upper bound the Lipschitz constant of $g(U)$, which we do in Lemma 2. Then Levy's lemma (Fact 20) gives an exponential concentration result for $g(U)$ under the Haar measure. Using techniques from [11], [12], we obtain upper bounds on the centralised moments of $(g(U))^2$ under the Haar measure. Observe now that $(g(U))^2$ is a balanced degree two polynomial (for the precise meaning see Definition 10) in the matrix entries of U . We then use Low's [11] derandomisation technique in order to obtain an exponential concentration result for $(g(U))^2$ when the unitary U^A is chosen uniformly from t -designs with the value of t stated above. This then leads to a similar exponential concentration result for $f(U)$ when U^A is chosen uniformly from a t -design, completing the proof of Theorem 1.¹

C. Organisation of the paper

Section II describes some notations, definitions and basic facts required for the paper. Section III proves the main result on one-shot decoupling with exponentially high concentration using unitary t -designs. The bounds obtained are described using smooth versions of variants of one-shot Rényi 2-entropies and max entropies. Section IV considers the main decoupling result in the iid limit and obtains bounds in terms of the more familiar Shannon entropic quantities. Section V shows how to apply the main result in order to obtain an exponential concentration for FQSW theorem for unitary designs. Section VI discuss the implications of FQSW concentration to relative thermalisation in quantum thermodynamics and to the Hayden-Preskill model for the black hole information paradox. Section VII concludes the paper and discusses directions for further research.

II. PRELIMINARIES

A. Notation

All vector spaces considered in the paper are finite dimensional inner product spaces, aka finite dimensional Hilbert spaces, over the complex field. We use $|V|$ to denote the dimension of a Hilbert space V . Letters $c_1, c_2, c'_1, c'_2, \dots$ denote positive universal constants. Logarithms are all taken in base two. We tacitly assume that the ceiling is taken of any formula that provides dimension or value of t

¹A part of this work was presented in the online workshop Beyond IID in Information Theory-8.

in unitary t -design. The symbols \mathbb{E} , \mathbb{P} denote expectation and probability respectively. The abbreviation "iid" is used to mean identically and independently distributed, which just means taking the tensor power of the identical copies of the underlying state. The notation "：“ is used to denote the definitions of the underlying mathematical quantities.

The notation $\mathcal{L}(A_1, A_2)$ denotes the Hilbert space of all linear operators from Hilbert space A_1 to Hilbert space A_2 with the inner product being the Hilbert-Schmidt inner product $\langle M, N \rangle := \text{Tr}[M^\dagger N]$. For the special case when $A_1 = A_2$ we use the phrase operator on A_1 and the symbol $\mathcal{L}(A_1)$. Further, when $A_1 = A_2 = \mathbb{C}^m$, \mathcal{M}_m denotes vector space of all $m \times m$ matrices. The symbol I^A denotes the identity operator on vector space A . The matrix π^A denotes the so-called completely mixed state on system A , i.e., $\pi^A := \frac{I^A}{|A|}$. We use the notation $U \circ A$ as a short hand to denote the conjugation of the operator U on the operator A , that is, $U \circ A := UAU^\dagger$.

The symbol ρ usually denotes a quantum state aka density matrix which is nothing but a Hermitian positive semidefinite matrix with unit trace, and $\mathcal{D}(\mathbb{C}^d)$ denotes the set of all $d \times d$ density matrices. The symbol $\text{Pos}(\mathbb{C}^d)$ denotes the set of all $d \times d$ positive semidefinite matrices, and the symbol $\mathbb{U}(d)$ denotes the set of all $d \times d$ unitary matrices with complex entries. For a positive semidefinite matrix σ , we use σ^{-1} to denote the operator which is the orthogonal direct sum of the inverse of σ on its support and the zero operator on the orthogonal complement of the support. This definition of σ^{-1} is also known as the *Moore-Penrose pseudoinverse*. The symbol $|v\rangle$ denotes a vector v of unit ℓ_2 -norm, and $\langle v|$ denotes the corresponding linear functional. A rank one density matrix is called a pure quantum state. Often, in what is a loose notation, a pure quantum state $|v\rangle\langle v|$ is denoted by just the vector $|v\rangle$ or, if we want to emphasise the density matrix formalism, by the notation v . For two Hermitian matrices A, B of the same dimension, we use $A \geq B$ as a shorthand for the statement that $A - B$ is positive semidefinite.

We use the notation $H(\cdot)$ to denote the usual Shannon or von Neumann entropy of the underlying state and the notation $\mathbb{H}(\cdot)$ to denote the modified or the new defined entropic quantities in this work.

Let $M \in \mathcal{L}(A)$. The symbol $\text{Tr } M$ denotes the trace of operator M . Trace is a linear map from $\mathcal{L}(A)$ to \mathbb{C} . Let A, B be two vector spaces. The partial trace $\text{Tr}_B[\cdot]$ obtained by tracing out B is defined to be the unique linear map from $\mathcal{L}(A \otimes B)$ to $\mathcal{L}(A)$ satisfying $\text{Tr}_B[M \otimes N] = (\text{Tr } N)M$ for all operators $M \in \mathcal{L}(A)$, $N \in \mathcal{L}(B)$.

A linear map $\mathcal{T} : \mathcal{M}_m \rightarrow \mathcal{M}_d$ is called a superoperator. A superoperator \mathcal{T} is said to be *positive* if it maps positive semidefinite matrices to positive semidefinite matrices, and *completely positive* if $\mathcal{T} \otimes \mathbb{I}$ is a positive superoperator for all identity superoperators \mathbb{I} . A superoperator \mathcal{T} is said to be *trace preserving* if $\text{Tr}[\mathcal{T}(M)] = \text{Tr } M$ for all $M \in \mathcal{M}_m$. Completely positive and trace preserving (abbreviated as CPTP) superoperators are called *quantum operations* or *quantum channels*. In this paper we only consider completely positive and trace non-increasing superoperators. Note that both trace and partial trace defined in the previous paragraph are quantum channels.

The adjoint of a superoperator is defined with respect to the Hilbert-Schmidt inner product on matrices. In other words, if $\mathcal{T} : \mathcal{M}_m \rightarrow \mathcal{M}_d$ is a superoperator, then its adjoint $\mathcal{T}^\dagger : \mathcal{M}_d \rightarrow \mathcal{M}_m$ is a superoperator uniquely defined by the property that $\langle \mathcal{T}^\dagger(A), B \rangle = \langle A, \mathcal{T}(B) \rangle$ for all $A \in \mathcal{M}_d$, $B \in \mathcal{M}_m$.

We will be using the Stinespring representation of a superoperator, which we state as the following fact:

Fact 4. Any superoperator $\mathcal{T}^{A \rightarrow B}$ can be represented as:

$$\mathcal{T}^{A \rightarrow B}(M^A) = \text{Tr}_Z \{ V_{\mathcal{T}}^{AC \rightarrow BZ} (M^A \otimes (|0\rangle\langle 0|)^C) (W_{\mathcal{T}}^{AC \rightarrow BZ})^\dagger \}$$

where $V_{\mathcal{T}}$, $W_{\mathcal{T}}$ are operators that map vectors from $A \otimes C$ to vectors in $B \otimes Z$. Systems C and Z are considered as the input and output ancillary systems respectively, such that $|A||C| = |B||Z|$. Without

loss of generality, $|C| \leq |B|$ and $|Z| \leq |A|$. Furthermore, in the following special cases $V_{\mathcal{T}}$, $W_{\mathcal{T}}$ have additional properties.

- 1) \mathcal{T} is completely positive if and only if $V_{\mathcal{T}} = W_{\mathcal{T}}$.
- 2) \mathcal{T} is trace preserving if and only if $V_{\mathcal{T}}^{-1} = W_{\mathcal{T}}^{\dagger}$. Thus, \mathcal{T} is completely positive and trace preserving if and only if $V_{\mathcal{T}} = W_{\mathcal{T}}$ and are unitary operators.
- 3) \mathcal{T} is completely positive and trace non-decreasing if and only if $V_{\mathcal{T}} = W_{\mathcal{T}}$ and $\|V_{\mathcal{T}}\|_{\infty} \leq 1$.

For $p \geq 1$, Schatten p -norm for any operator $M \in \mathcal{L}(A_1, A_2)$ is defined as $\|M\|_p \triangleq [\text{Tr}((M^{\dagger}M)^{p/2})]^{1/p}$. In other words, $\|M\|_p$ is nothing but the ℓ_p -norm of the tuple of singular values of M . The Schatten ∞ -norm is defined by taking the limit $p \rightarrow \infty$, and turns out to be the largest singular value of M . The Schatten 2-norm, aka the Hilbert Schmidt norm, is nothing but the ℓ_2 -norm of the tuple obtained by stretching out the entries of the matrix into a vector. The Schatten ∞ -norm is nothing but the operator norm $\|M\|_{\infty} = \max_{\|v\|_2=1} \|Mv\|_2$. The Schatten 1-norm is also known as the trace norm. We have the norm properties $|\text{Tr } M| \leq \|M\|_1$, $\|M\|_1 \leq \sqrt{\text{Tr } I} \|M\|_2$, $\|M\|_p \leq (\text{Tr } I)^{1/p} \|M\|_{\infty}$, $\|M \otimes N\|_p = \|M\|_p \|N\|_p$, $\|M\|_p \leq \|M\|_q$ if $p \geq q$ and $\|MN\|_p \leq \min\{\|M\|_p \|N\|_{\infty}, \|M\|_{\infty} \|N\|_p\}$.

The distance between two CP maps $\mathcal{T}_1^{A \rightarrow B}$ and $\mathcal{T}_2^{A \rightarrow B}$ can be measured in terms of the *diamond norm* [13] defined as follows:

$$\|\mathcal{T}_1 - \mathcal{T}_2\|_{\diamond} := \max_{\rho^{AA'}} \|(\mathcal{T}_1 \otimes \mathbb{I}^{A'}) (\rho^{AA'}) - (\mathcal{T}_2 \otimes \mathbb{I}^{A'}) (\rho^{AA'})\|_1,$$

where A' is a new Hilbert space of the same dimension as A and the maximisation is over all quantum states $\rho^{AA'}$.

B. Matrix manipulation

Fact 5. For Hilbert spaces \mathcal{H}_X , \mathcal{H}_Y suppose that vectors $|\psi\rangle, |\phi\rangle \in \mathcal{H}_X \otimes \mathcal{H}_Y$ satisfy $\text{Tr}_Y(|\psi\rangle\langle\psi|) = \text{Tr}_Y(|\phi\rangle\langle\phi|)$. Then there exists a unitary operator U on \mathcal{H}_Y such that $|\psi\rangle = (I^X \otimes U^Y)|\phi\rangle$.

Fix an orthonormal basis $\{|a\rangle^A\}_a$ of A and $\{|z\rangle^Z\}_z$ of Z . Consider the tensor basis $\{|a\rangle^A \otimes |z\rangle^Z\}_{a,z}$ of the Hilbert space $A \otimes Z$. The isometric linear map $\text{vec}^{A,Z} : \mathcal{L}(Z, A) \rightarrow A \otimes Z$ is defined as the unique linear map satisfying $\text{vec}^{A,Z}(|a\rangle^A \langle z|^Z) := |a\rangle^A \otimes |z\rangle^Z$ [14]. The inverse linear map is denoted by $(\text{vec}^{A,Z})^{-1}$. It is also an isometry. We will be using the following property of the vec^{-1} map which we state as a fact here. A simpler version of this fact was used in [15].

Fact 6. For any two vectors $|x\rangle^{AZ}$, $|y\rangle^{AZ}$ on a bipartite Hilbert space $A \otimes Z$,

$$(\text{Tr}_Z(|x\rangle\langle y|^{AZ}))^{A \times A} = (\text{vec}^{-1}(|x\rangle))^{A \times Z} \left[(\text{vec}^{-1}(|y\rangle))^{A \times Z} \right]^{\dagger}$$

where $\text{vec}^{-1} : A \otimes Z \rightarrow A \times Z := \mathcal{L}(Z, A)$.

Proof. Fix orthonormal bases $\{|a\rangle^A\}_a$, $\{|z\rangle^Z\}_z$ for A , Z . We can write

$$|x\rangle^{AZ} = \sum_{az} x_{az} |a\rangle^A |z\rangle^Z, \quad |y\rangle^{AZ} = \sum_{az} y_{az} |a\rangle^A |z\rangle^Z.$$

This gives

$$\begin{aligned} \text{vec}^{-1}(|x\rangle) &= \sum_{az} x_{az} |a\rangle^A \langle z|^Z, \quad \text{vec}^{-1}(|y\rangle) = \sum_{az} y_{az} |a\rangle^A \langle z|^Z, \\ \Rightarrow (\text{vec}^{-1}(|x\rangle)) (\text{vec}^{-1}(|y\rangle))^{\dagger} &= \sum_{aa'} \sum_z x_{az} y_{a'z}^* |a\rangle^A \langle a'|^A. \end{aligned}$$

On the other hand

$$\text{Tr}_Z(|x\rangle\langle y|^{AZ}) = \text{Tr}_Z\left(\sum_{aa'}\sum_{zz'}x_{az}y_{a'z'}^*|a\rangle\langle a'|^A\otimes|z\rangle\langle z'|^Z\right) = \sum_{aa'}x_{az}y_{a'z}^*|a\rangle\langle a'|^A.$$

This completes the proof. \square

We now state the so called polar decomposition of any linear operator.

Fact 7. Any operator M can be expressed as $M = VQ$, known as the left polar decomposition, where V is a unitary matrix and Q is a positive semidefinite matrix. Also, M can be expressed as $M = PU$, where P is a positive semidefinite matrix and U is a unitary matrix. This is known as the right polar decomposition.

Next, we state four useful facts from Dupuis' thesis [3].

Fact 8 ([3, Lemma I.1]). Let ρ, ρ' and σ be positive semidefinite operators on \mathcal{H} such that $\text{Tr}[\rho'] \leq \text{Tr}[\sigma]$ and $\rho' \geq \rho$. Then, $\|\rho' - \sigma\|_1 \leq 2\|\rho - \sigma\|_1$.

Fact 9 ([3, Lemma I.2]). Let ρ^{AB} be a positive semidefinite operator, and let $0 \leq P^B \leq I^B$. Then, $\text{Tr}_B[(P^B \otimes I^A)\rho^{AB}(P^B \otimes I^A)] \leq \rho^A$.

Proof. We give a more direct and elementary proof of this fact than what was given in [3]. The proof is a simple application of the definition of the partial trace and the fact that $P^B \leq I^B \Rightarrow (P^B)^2 \leq P^B$. By spectral theorem for positive semidefinite matrices, we express P^B in its eigenbasis as $P^B = \sum_{i=1}^{|B|} p_i |b_i\rangle\langle b_i|^B$. Since $P^B \leq I^B$, therefore $p_i \leq 1, \forall i$. Now we express ρ^{AB} in block diagonal form with $\{|b_j\rangle\}_{j=1}^{|B|}$ as the orthonormal basis for \mathcal{H}_B :

$$\rho^{AB} = \sum_{j,j'=1}^{|B|} A_{j,j'}^A \otimes |b_j\rangle\langle b_{j'}|^B \Rightarrow \rho^A = \sum_j |A_{j,j}^A|.$$

The block matrices $A_{j,j}$ are positive semidefinite. Now evaluating $\text{Tr}_B[P^B \cdot \rho^{AB}]$:

$$\text{Tr}_B[P^B \cdot \rho^{AB}] = \sum_{j,j',k,l=1}^{|B|} p_k p_l A_{j,j'}^A \langle b_k | b_j \rangle \langle b_{j'} | b_l \rangle \langle b_l | b_k \rangle = \sum_{j=1}^{|B|} p_j^2 A_{j,j}^A \stackrel{a}{\leq} \sum_{j=1}^{|B|} A_{j,j}^A = \rho^A,$$

where (a) holds since $p_j \leq 1$ and $A_{j,j}$ are positive semidefinite matrices for all j . This completes the proof. \square

Fact 10 ([3, Lemma I.3]). Let $|\psi\rangle^{AB} \in A \otimes B$, $\rho^A \in \text{Pos}(A)$ such that $\rho^A \leq \psi^A$. Then, there exists an operator P^B on B such that $0^B \leq P^B \leq I^B$ and $\text{Tr}_B[(P^B \otimes I^A)|\psi\rangle^{AB}\langle\psi|(P^B \otimes I^A)] = \rho^A$.

Proof. We give a simpler and more direct proof of this fact than given in [3]. Since $\psi^A \geq \rho^A$, there exists a positive semidefinite matrix σ^A such that $\psi^A = \rho^A + \sigma^A$. Let the vector $|\rho\rangle^{AB}$ be a purification of ρ^A and the vector $|\sigma\rangle^{AB}$ be a purification of σ^A . The squares of the lengths of the purifying vectors are equal to the traces of the respective matrices. Now let $Q = \mathbb{C}^2$ be the system representing a qubit. We define the unit length pure state $|\theta\rangle^{ABQ}$ as:

$$|\theta\rangle^{ABQ} \triangleq |\rho\rangle^{AB} \otimes |0\rangle^Q + |\sigma\rangle^{AB} \otimes |1\rangle^Q$$

It follows that $|\theta\rangle^{ABQ}$ is a purification of the state ψ^A and so is the state $|\psi\rangle^{AB} \otimes |0\rangle^Q$. Thus by Fact 5 there exists a unitary matrix U^{BQ} on the composite system BQ satisfying:

$$|\theta\rangle^{ABQ} = (I^A \otimes U^{BQ})(|\psi\rangle^{AB} |0\rangle^Q)$$

Now we define a POVM measurement that first appends the ancilla Q initialized to state $|0\rangle^Q$ to the state $|\psi\rangle^{AB}$, followed by applying the unitary $I^A \otimes U^{BQ}$ on the state $|\psi\rangle^{AB}|0\rangle^Q$ and finally measuring the ancilla system Q of the resultant state in computational basis $\{|0\rangle\langle 0|^Q, |1\rangle\langle 1|^Q\}$. The measurement succeeds if we get the outcome 0 in the ancilla register. Formally, the outcome ‘success’ is described by an operator $M^B := (I^B \otimes \langle 0|^Q)U^{BQ}(I^B \otimes |0\rangle^Q)$. Clearly, $\|M^B\|_\infty \leq 1$. We thus have:

$$\begin{aligned} & \text{Tr}_B[(I^A \otimes M^B)|\psi\rangle\langle\psi|^{AB}(I^A \otimes M^{\dagger B})] \\ &= \text{Tr}_B[(I^{AB} \otimes \langle 0|^Q)(I^A \otimes U^{BQ})(|\psi\rangle\langle\psi|^{AB}|0\rangle^Q)(\langle\psi|^{AB}\langle 0|^Q)(I^A \otimes U^{\dagger BQ})(I^{AB} \otimes |0\rangle^Q)] \\ &= \text{Tr}_B[(I^{AB} \otimes \langle 0|^Q)|\theta\rangle\langle\theta|^{AB}(I^{AB} \otimes |0\rangle^Q)] = \text{Tr}_B[\rho^{AB}\langle\theta|] = \rho^A. \end{aligned}$$

Now, to come up with P^B as mentioned in the statement of the fact we express $M^B = U_M^B P^B$, using the polar decomposition from Fact 7, with $P_B \geq 0$. Since $\|M\|_\infty \leq 1$, therefore $P^B \leq I^B$. Thus we get,

$$\begin{aligned} \rho^A &= \text{Tr}_B[(I^A \otimes M^B)|\psi\rangle\langle\psi|^{AB}(I^A \otimes M^{\dagger B})] \\ &= \text{Tr}_B[(I^A \otimes U_M^B)(I^A \otimes P^B)|\psi\rangle\langle\psi|^{AB}(I^A \otimes P^B)(I^A \otimes U_M^{\dagger B})] \\ &\stackrel{a}{=} \sum_i (I^A \otimes \langle i|^B)(I^A \otimes U_M^B)(I^A \otimes P^B)|\psi\rangle\langle\psi|^{AB}(I^A \otimes P^B)(I^A \otimes U_M^{\dagger B})(I^A \otimes |i\rangle^B) \\ &= \sum_i (I^A \otimes (U_M^{\dagger}|i\rangle)^{B\dagger})(I^A \otimes P^B)|\psi\rangle\langle\psi|^{AB}(I^A \otimes P^B)(I^A \otimes (U_M^{\dagger}|i\rangle)^B) \\ &\stackrel{b}{=} \sum_i (I^A \otimes \langle u_i|^B)(I^A \otimes P^B)|\psi\rangle\langle\psi|^{AB}(I^A \otimes P^B)(I^A \otimes |u_i\rangle^B) \\ &= \text{Tr}_B[(I^A \otimes P^B)|\psi\rangle\langle\psi|^{AB}(I^A \otimes P^B)] \end{aligned}$$

where (a) follows by the basis dependent definition of Tr_B by fixing $\{|i\rangle\}_{i=1}^{|B|}$ as an orthonormal basis for system B ; (b) holds since U_M^B is a unitary matrix that maps orthonormal basis $\{|i\rangle\}_{i \in B}$ to orthonormal basis $\{|u_i\rangle\}_{i \in B}$. We thus have a $0 \leq P^B \leq I^B$ satisfying the fact. This completes the proof. \square

Fact 11 ([3, Lemma 3.5]). *Let $\rho^{AB} \in \text{Pos}(A \otimes B)$ and let $\rho^B := \text{Tr}_A \rho^{AB}$. Then,*

$$|A|^{-1} \leq \frac{\|\rho^{AB}\|_2^2}{\|\rho^B\|_2^2} \leq |A|.$$

In order to upper bound Schatten 1-norm of an operator, sometimes it is more convenient to upper bound Schatten 2-norm of a slightly modified operator. The following fact, which is nothing but an application of the Cauchy-Schwarz inequality, allows us to do so.

Fact 12. *Let $M \in \mathcal{L}(\mathcal{H})$ and $\sigma \in \mathcal{D}(\mathcal{H})$, $\sigma > 0$. Then $\|M\|_1 \leq \|\sigma^{-1/4} M \sigma^{-1/4}\|_2$.*

We will also need Winter’s gentle measurement lemma [16].

Fact 13. *Let P be a positive operator such that $P \leq I$. For any density matrix ρ , satisfying $\text{Tr}[P\rho P] \geq 1 - \epsilon$, it holds that $\|\rho - P\rho P\|_1 \leq 2\sqrt{\epsilon}$.*

We now state an important geometric fact about how a pair of subspaces of a Hilbert space interact. This fact, first discovered by Jordan a hundred and fifty years ago but which has since been independently rediscovered many times, defines *canonical angles* between a pair of subspaces. These angles are sometimes called as *chordal angles*.

Fact 14. [17, Fact 6] *Let A, B be subspaces of a Hilbert space \mathcal{H} . Then there is a decomposition of \mathcal{H} as an orthogonal direct sum of the following types of subspaces:*

- 1) *One dimensional spaces orthogonal to both A and B ;*
- 2) *One dimensional spaces contained in both A and B ;*

- 3) One dimensional spaces contained in A and orthogonal to B ;
- 4) One dimensional spaces contained in B and orthogonal to A ;
- 5) Two dimensional spaces intersecting A, B each in one dimensional spaces.

Moreover, the one dimensional spaces in (2) and (3) above together with the one dimensional intersections of the spaces in (5) with A form an orthonormal basis of A . A similar statement holds for B .

We end this section by stating two properties of the so-called *swap trick* that will be useful later on.

Fact 15 ([3, Lemma 3.3]). For two operators $M^A, N^A \in \mathcal{L}(A)$, we have $\text{Tr}[(MN)^A] = \text{Tr}[(M^{A_1} \otimes N^{A_2})F^{A_1 A_2}]$, where A_1, A_2 are two Hilbert spaces of the same dimension as A and $F^{A_1 A_2}$ swaps the tensor multiplicand systems A_1 and A_2 .

Fact 16. For an operator $M^{AR} \in \mathcal{L}(A \otimes R)$, we have

$$\left\| \text{Tr}_{R_1 R_2} [(I^{A_1 A_2} \otimes F^{R_1 R_2})(M^{A_1 R_1} \otimes (M^\dagger)^{A_2 R_2})] \right\|_1 \leq |A| \left\| M^{AR} \right\|_2^2.$$

Proof. Fix an orthonormal basis $\{|r\rangle\}_{r \in R}$ for the system R . Let $M^{AR} = \sum_{rr'} M_{rr'}^A \otimes |r\rangle\langle r'|^R$, where $M_{rr'}^A$ is an operator in A for every r, r' . Then,

$$\begin{aligned} & \text{Tr}_{R_1 R_2} [(I^{A_1 A_2} \otimes F^{R_1 R_2})(M^{A_1 R_1} \otimes (M^\dagger)^{A_2 R_2})] \\ &= \sum_{rr' r'' r'''} \left(M_{rr'}^{A_1} \otimes (M_{r'' r'''}^\dagger)^{A_2} \right) \text{Tr}_{R_1 R_2} [F^{R_1 R_2} (|r\rangle\langle r'|^{R_1} \otimes |r''\rangle\langle r'''|^{R_2})] \\ &= \sum_{rr' r'' r'''} \left(M_{rr'}^{A_1} \otimes (M_{r'' r'''}^\dagger)^{A_2} \right) \text{Tr}_{R_1 R_2} [F^{R_1 R_2} |r\rangle^{R_1} |r''\rangle^{R_2} \langle r'|^{R_1} \langle r'''|^{R_2}] \\ &= \sum_{rr' r'' r'''} \left(M_{rr'}^{A_1} \otimes (M_{r'' r'''}^\dagger)^{A_2} \right) \text{Tr}_{R_1 R_2} [|r''\rangle^{R_1} |r\rangle^{R_2} \langle r'|^{R_1} \langle r'''|^{R_2}] = \sum_{rr'} M_{rr'}^{A_1} \otimes (M_{rr'}^\dagger)^{A_2}. \end{aligned}$$

So,

$$\begin{aligned} \left\| \sum_{rr'} M_{rr'}^{A_1} \otimes (M_{rr'}^\dagger)^{A_2} \right\|_1 &\leq \sum_{rr'} \left\| M_{rr'}^{A_1} \otimes (M_{rr'}^\dagger)^{A_2} \right\|_1 = \sum_{rr'} \left\| M_{rr'}^A \right\|_1^2 \\ &\leq \sum_{rr'} |A| \left\| M_{rr'}^A \right\|_2^2 \\ &= |A| \left\| M^{AR} \right\|_2^2. \end{aligned}$$

This completes the proof. \square

C. Entropic quantities

The Shannon entropy of a random variable X with probability distribution $(p_x)_x$ is defined as $H(X)_p := -\sum_x p_x \log p_x$. For a quantum system B in a state ω^B , the Shannon entropy is defined analogously as $H(B)_\omega := -\text{Tr}[\omega \log \omega]$. For a bipartite quantum system AB in a state ω^{AB} , the conditional Shannon entropy is defined as $H(A|B)_\omega := H(AB)_\omega - H(B)_\omega$.

We recall the definition of the smooth conditional Rényi 2-entropy from [3].

Definition 1. Let $0 \leq \epsilon < 1$. The ϵ -smooth conditional Rényi 2-entropy for a bipartite positive semidefinite operator ρ^{AR} on systems A and R is defined as:

$$H_2^\epsilon(A|R)_\rho := -2 \log \min_{\substack{\sigma^{AR} \in \text{Pos}(AR): \\ \|\rho^{AR} - \sigma^{AR}\|_1 \leq \epsilon \\ \omega^R \in \mathcal{D}(R): \omega^R > 0^R}} \{ \|(\omega^R \otimes \mathbb{I}^A)^{-1/4} \sigma^{AR} (\omega^R \otimes \mathbb{I}^A)^{-1/4}\|_2 \}.$$

When $\epsilon = 0$, we simply refer to the above quantity as conditional Rényi 2-entropy and denote it by $H_2(A|R)_\rho$ and define $\tilde{\rho}^{AR} := (\omega^R \otimes \mathbb{I}^A)^{-1/4} \rho^{AR} (\omega^R \otimes \mathbb{I}^A)^{-1/4}$.

We also recall the definition of ϵ -smooth max-entropy defined in [18].

$$H_{\max}^\epsilon(B)_\rho := 2 \log \min_{\substack{\sigma^B \in \text{Pos}(B): \\ \|\rho^B - \sigma^B\|_1 \leq \epsilon}} \{ \text{Tr} \sqrt{\sigma} \}$$

Next, we recall the definition of ϵ -smooth min-entropy defined in [3].

Definition 2. Let $0 \leq \epsilon < 1$. The ϵ -smooth min-entropy of ρ^B is defined as:

$$H_{\min}^\epsilon(B)_\rho := -\log \min_{\substack{\sigma^B \in \text{Pos}(B): \\ \|\sigma^B - \rho^B\|_1 \leq \epsilon}} \|\sigma^B\|_\infty.$$

We now define a new quantity that we call the *smooth modified max-entropy*.

Definition 3. The ϵ -smooth modified max-entropy of system B under a quantum state ω^B is defined as:

$$\mathbb{H}_{\max}^\epsilon(B)_\omega := \log \|(\omega_\epsilon'')^B\|_\infty^{-1},$$

where $(\omega_\epsilon'')^B$ is the positive semidefinite matrix obtained by zeroing out those smallest eigenvalues of ω^B that sum to less than or equal to ϵ . The ϵ -smooth modified max-entropy of a probability distribution can be defined similarly.

It is easy to see that $H_{\max}^\epsilon(B)_\omega \leq \mathbb{H}_{\max}^\epsilon(B)_\omega \leq \log(|B|/\epsilon)$ for any state ω^B .

A slightly finer linear algebraic analysis would give us that $\mathbb{H}_{\max}^\epsilon(B)_\omega \leq \log \left(\frac{|B| - \text{supp}((\omega_\epsilon'')^B) + 1}{\epsilon} \right)$. One of the applications of this smoothed modified max-entropy is found in a recent work on one-shot purity distillation in [19].

We next define a novel entropic quantity called *smooth modified conditional Rényi 2-entropy*.

Definition 4. Let $0 \leq \epsilon, \delta < 1$. The (ϵ, δ) -smooth modified conditional Rényi 2-entropy for a bipartite positive semidefinite operator ω^{AB} on systems A and B is defined as:

$$\begin{aligned} \mathbb{H}_{2,\delta}^{\epsilon,\delta}(A|B)_\omega &:= -2 \log \min_{\substack{\eta^{AB}: 0^{AB} \leq \eta^{AB} \leq \omega^{AB}, \|\omega^{AB} - \eta^{AB}\|_1 \leq \epsilon \\ \forall |v\rangle \in \text{supp}(\eta^{AB}): \|(I^A \otimes \Pi_{\text{supp}(\omega_{\epsilon,\delta}''')^B})|v\rangle\|_2^2 \geq 1 - \epsilon}} \| (I^A \otimes (\omega_{\epsilon,\delta}''')^B)^{-1/4} \eta^{AB} (I^A \otimes (\omega_{\epsilon,\delta}''')^B)^{-1/4} \|_2, \end{aligned}$$

where $(\omega_{\epsilon,\delta}''')^B$ is the positive semidefinite operator obtained by zeroing out those eigenvalues of ω^B that are smaller than $2^{-(1+\delta)\mathbb{H}_{\max}^\epsilon(B)_\omega}$.

Observe that $(\omega_{\epsilon,\delta}''')^B \geq (\omega_\epsilon'')^B$, where $(\omega_\epsilon'')^B$ is defined in Definition 3 above.

We remark that the above modified definitions of the smoothed max and the conditional 2-entropy is inspired (not directly inferred though) from the so-called information spectrum methods used in the

analysis of asymptotic independent but not identical classical and quantum information theory. We refer the reader to [20]–[22] for a detailed exposition.

It is easy to see, via Fact 13, that for any state ω^{AB} , $H_2^{4\sqrt{\epsilon}}(A|B)_\omega \geq \mathbb{H}_2^{\epsilon,\delta}(A|B)_\omega$ for any $\delta > 0$. Also the following fact holds.

Fact 17. *Let $\epsilon > 0$. Then, $H_2^\epsilon(A|B)_\omega \leq H(A|B)_\omega + 8\epsilon \log |A| + 2 + 2 \log \epsilon^{-1}$.*

Proof. The proof follows by combining Equation 8 of [23] with Theorem 7, Lemma 2 and Equation 33 of [18] and then applying the Alicki-Fannes inequality [24]. \square

D. Types and typicality

The smooth entropic quantities defined in the previous section are suitably bounded by the standard Shannon entropic quantities in the iid limit, as will be shown in Section IV. In order to lay the groundwork for the proofs in Section IV, we recall the definitions of types, typical sequences and subspaces.

Definition 5. *Let X be a finite set. Fix a probability distribution p on X . Let n be a positive integer. Let X^n denote the random variable corresponding to n independent copies of X . The notation x^n shall represent a sequence of length n over the alphabet X . Let $N(a|x^n)$ denote the number of occurrences of the symbol $a \in X$ in the sequence x^n . The multiset $\{N(a|x^n)\}_{a \in X}$ is called the type of x^n . The set of all possible types is nothing but the set of all possible $|X|$ -tuples of non-negative integers summing up to n .*

Definition 6. *Let $0 < \delta < 1$. The set of strongly δ -typical types of length n over the alphabet X pertaining to the distribution p is defined as [25] $\left\{ (m_a)_{a \in X} : \forall a \in X, m_a \in np(a)(1 \pm \delta) \right\}$. A sequence x^n is said to be strongly δ -typical if its type is strongly δ -typical. The set of strongly δ -typical sequences is denoted by $T_{p,\delta}^{X^n}$.*

Let p^n denote the n -fold tensor power of probability distribution p . The strongly typical sequences satisfy the following property which is called as Asymptotic Equipartition Property (AEP) in classical Shannon theory.

Fact 18 ([17]). *The number of types is $\binom{n+|X|-1}{|X|-1}$. The set of all possible sequences X^n is partitioned into a disjoint union, over all possible types, of sequences having a given type. Let $0 < \epsilon, \delta < 1/2$. Define $p_{\min} := 2^{-\mathbb{H}_{\max}^{\epsilon/2}(X)_p}$. Let $n \geq 4p_{\min}^{-1} \delta^{-2} \log(|X|/\epsilon)$. Then,*

$$\begin{aligned} \sum_{x^n \in T_{p,\delta}^{X^n}} p^n(x^n) &\geq 1 - \epsilon, \\ \forall x^n \in T_{p,\delta}^{X^n} : 2^{-nH(X)(1+\delta)} &\leq p^n(x^n) \leq 2^{-nH(X)(1-\delta)}, \\ 2^{nH(X)(1-\delta)}(1 - \epsilon) &\leq |T_{p,\delta}^{X^n}| \leq 2^{nH(X)(1+\delta)}. \end{aligned}$$

In the quantum setting, we extend the notion of types and typical sequences with respect to a particular distribution to the notion of type subspaces and typical subspaces with respect to the n -fold tensor product of a quantum state.

Definition 7. *Let ρ be a density matrix over a Hilbert space B . Consider a canonical eigenbasis $\mathcal{B} = \{|\chi_1\rangle, \dots, |\chi_{|B|}\rangle\}$ of ρ . Consider the diagonalisation $\rho = \sum_{\chi \in \mathcal{B}} q(\chi) |\chi\rangle\langle\chi|$, where the set of eigenvalues $\{q(\chi)\}_\chi$ can be treated as a probability distribution over \mathcal{B} . Given a type $(m(\chi))_{\chi \in \mathcal{B}}$, which is nothing*

but a $|B|$ -tuple of non-negative integers summing to n , we define the corresponding type subspace to be the span of all n -fold tensor products of vectors from \mathcal{B} having the given type.

Definition 8. Let $0 < \epsilon, \delta < 1/2$. The strongly δ -typical subspace of $B^{\otimes n}$ corresponding to the n -fold tensor power operator $\rho^{\otimes n}$, $T_{\rho, \delta}^{B^n}$, is defined as the orthogonal direct sum of all type subspaces with strongly δ -typical types with respect to the probability distribution q on \mathcal{B} .

Let $\Pi_{\rho, \delta}^{B^n}$ denote the orthogonal projection onto $T_{\rho, \delta}^{B^n}$. The typical projector satisfies the following so called quantum AEP analogous to that of Fact 18:

Fact 19 ([17]). The number of types is $\binom{n+|B|-1}{|B|-1}$. The Hilbert space $B^{\otimes n}$ can be decomposed into an orthogonal direct sum, over all possible types, of type subspaces. Let $0 < \epsilon, \delta < 1/2$. Let ρ be a quantum state. Define $q_{\min} \cong q_{\min}(\rho) := 2^{-\mathbb{H}_{\max}^{\epsilon/2}(B)_\rho}$. Suppose that $n \geq 4q_{\min}^{-1}\delta^{-2}\log(|B|/\epsilon)$. Then,

$$\begin{aligned} \text{Tr}[\rho^{\otimes n} \Pi_{\rho, \delta}^{B^n}] &\geq 1 - \epsilon, \\ 2^{-nH(X)(1+\delta)} \Pi_{\rho, \delta}^{B^n} &\leq \Pi_{\rho, \delta}^{B^n} \rho^{\otimes n} = \rho^{\otimes n} \Pi_{\rho, \delta}^{B^n} = \Pi_{\rho, \delta}^{B^n} \rho^{\otimes n} \Pi_{\rho, \delta}^{B^n} \leq 2^{-nH(X)(1-\delta)} \Pi_{\rho, \delta}^{B^n}, \\ 2^{nH(X)(1-\delta)}(1 - \epsilon) &\leq \text{Tr} \Pi_{\rho, \delta}^{B^n} \leq 2^{nH(X)(1+\delta)}. \end{aligned}$$

E. Concentration of measure

We state the main tool for concentration of measure of Lipschitz functions defined on the sphere or on the unitary group in high dimensions.

Definition 9. A complex valued function f defined on a subset $S \subseteq \mathbb{C}^n$ is said to be L -Lipschitz, or with Lipschitz constant L , if $\forall x, y \in S$ it satisfies the following inequality:

$$|f(x) - f(y)| \leq L\|x - y\|_2. \quad (5)$$

Remark 2. In order to find Lipschitz constant of a complex valued Lipschitz function f with the domain as the set of unitary operators, we find an upper bound on the ratio $\frac{|f(U) - f(V)|}{\|U - V\|_2}$. We recall that the norm $\|U - V\|_2$ is the Schatten-2 norm of the operator $U - V$ defined as $\sqrt{\text{Tr}[(U - V)^\dagger(U - V)]}$.

Fact 20. (Levy's Lemma [26], [27]) Let f be an L -Lipschitz function on $\mathbb{U}(n)$ where the metric on $\mathbb{U}(n)$ is induced by the embedding of $\mathbb{U}(n)$ into \mathbb{C}^{n^2} . In other words, the metric on $\mathbb{U}(n)$ is taken to be the Schatten 2-norm. Consider the Haar probability measure on $\mathbb{U}(n)$. Let the mean of f be μ . Then:

$$\mathbb{P}_{U \sim \text{Haar}}(|f(U) - \mu| \geq \lambda) \leq 2 \exp\left(-\frac{n\lambda^2}{4L^2}\right).$$

Remark 3. Lévy's lemma is widely used tool in high dimensional probability and asymptotic geometric functional analysis. While the original form can be found in [26], several simplifications have been developed in the literature and one such classic reference is [28]. Here we use a much simplified form derived for sub-Gaussian distribution described in [29, Theorem 5.1.4].

The following fact can be used to compute upper bounds on the centralised moments of Lipschitz functions. The proof follows Bellare and Rompel's seminal work on concentration for sums of t -wise independent random variables [30, Lemma A.1] and its quantum adaptation by Low [11, Lemma 3.3]. However, inspired by a technique from [12], we extend the earlier results in an important and essential way by computing upper bounds on the centralised moments of squares of Lipschitz functions also, which will be required in Section III-B.

Fact 21. Let X be a non-negative random variable. Suppose there is a number μ satisfying, for any $\kappa > 0$, the tail bound $\mathbb{P}[|X - \mu| > \kappa] \leq C \exp(-a\kappa^2)$ for some positive constants C, a . Let m be a positive integer. Then

$$\mathbb{E}[(X - \mu)^{2m}] \leq C \left(\frac{m}{a}\right)^m, \quad \mathbb{E}[(X^2 - \mu^2)^{2m}] \leq \begin{cases} 3C \left(\frac{9m\mu^2}{a}\right)^m & 1 \leq m \leq \frac{9}{64}a\mu^2 \\ 3C \left(\frac{64m^2}{a^2}\right)^m & \text{otherwise} \end{cases}.$$

Proof. Let Ω with a probability measure $d\omega$ be the sample space serving as the domain of the measurable function X . Then,

$$\begin{aligned} \mathbb{E}[(X - \mu)^{2m}] &= \int_{\Omega} (X(\omega) - \mu)^{2m} d\omega = \int_{\Omega} \int_{0 \leq x \leq (X(\omega) - \mu)^{2m}} dx d\omega = \int_0^{\infty} \int_{(X(\omega) - \mu)^{2m} \geq x} d\omega dx \\ &= \int_0^{\infty} \mathbb{P}[(X - \mu)^{2m} \geq x] dx = \int_0^{\infty} \mathbb{P}[|X - \mu| \geq x^{\frac{1}{2m}}] dx \leq C \int_0^{\infty} \exp(-ax^{1/m}) dx \\ &= Cma^{-m} \int_0^{\infty} e^{-y} y^{m-1} dy = Ca^{-m} \Gamma(m+1) \leq C \left(\frac{m}{a}\right)^m. \end{aligned}$$

Let $A := \{\omega : 0 \leq X(\omega) < 2\mu\}$ and \bar{A} denote its complement in Ω . We thus have

$$\mathbb{P}(\bar{A}) \leq C \exp(-a\mu^2) \quad (6)$$

and,

$$\mathbb{E}[(X^2 - \mu^2)^{2m}] = \int_{\Omega} (X(\omega)^2 - \mu^2)^{2m} d\omega = \int_A (X(\omega)^2 - \mu^2)^{2m} d\omega + \int_{\bar{A}} (X(\omega)^2 - \mu^2)^{2m} d\omega,$$

We now separately upper bound the two integrals on the right hand side above. For the first integral,

$$\begin{aligned} \int_A (X(\omega)^2 - \mu^2)^{2m} d\omega &= \int_A (X(\omega) - \mu)^{2m} (X(\omega) + \mu)^{2m} d\omega \leq (3\mu)^{2m} \int_{\Omega} (X(\omega) - \mu)^{2m} d\omega \\ &\leq C(3\mu)^{2m} \left(\frac{m}{a}\right)^m = C \left(\frac{9m\mu^2}{a}\right)^m. \end{aligned}$$

Now if $m > \frac{9}{64}a\mu^2$, the last term above can be further upper bounded by $C \left(\frac{64m^2}{a^2}\right)^m$.

For the second integral,

$$\begin{aligned} \int_{\bar{A}} (X(\omega)^2 - \mu^2)^{2m} d\omega &= 2m \int_{\bar{A}} \int_{0 \leq x \leq (X(\omega)^2 - \mu^2)} x^{2m-1} dx d\omega \\ &= 2m \int_0^{3\mu^2} \int_{\bar{A}} x^{2m-1} d\omega dx + 2m \int_{3\mu^2}^{\infty} \int_{X(\omega)^2 \geq \mu^2 + x} x^{2m-1} d\omega dx \\ &= (3\mu^2)^{2m} \mathbb{P}(\bar{A}) + 2m \int_{3\mu^2}^{\infty} \int_{X(\omega)^2 \geq \mu^2 + x} x^{2m-1} d\omega dx \\ &\stackrel{a}{\leq} C(3\mu^2)^{2m} \exp(-a\mu^2) + 2m \int_{3\mu^2}^{\infty} \int_{X(\omega) \geq \mu + (\sqrt{x}/2)} x^{2m-1} d\omega dx \end{aligned}$$

$$\begin{aligned}
&\stackrel{b}{\leq} C \left(\frac{9m\mu^2}{a} \right)^m + 2m \int_{3\mu^2}^{\infty} \mathbb{P}[X - \mu \geq \sqrt{x}/2] x^{2m-1} dx \\
&\leq C \left(\frac{9m\mu^2}{a} \right)^m + 2Cm \int_{3\mu^2}^{\infty} \exp(-ax/4) x^{2m-1} dx \\
&\leq C \left(\frac{9m\mu^2}{a} \right)^m + 2Cm \left(\frac{4}{a} \right)^{2m} \int_0^{\infty} e^{-y} y^{2m-1} dy \\
&= C \left(\frac{9m\mu^2}{a} \right)^m + C \left(\frac{4}{a} \right)^{2m} \Gamma(2m+1) \\
&\leq C \left(\frac{9m\mu^2}{a} \right)^m + C \left(\frac{8m}{a} \right)^{2m}.
\end{aligned}$$

where (a) follows from Equation 6 and the observation that for any $x \geq 3\mu^2$, $\sqrt{\mu^2 + x} > \mu + \sqrt{x}/2$; and (b) follows from the observation that

$$\exp(-a\mu^2) \leq \left(\frac{m}{a\mu^2} \right)^m \quad (7)$$

, which can be seen from the following cases:

Case 1: For all $m \geq a\mu^2$, it holds that $\left(\frac{m}{a\mu^2} \right)^m \geq 1$ and $\exp(-a\mu^2) \leq 1$. Thus Equation 7 holds trivially.

Case 2: For $m < a\mu^2$, choose a $\delta < 1$ and let $m = \delta a\mu^2$. Then $\left(\frac{m}{a\mu^2} \right)^m = \exp(-\delta \ln(\frac{1}{\delta}) a\mu^2) > \exp(-a\mu^2)$ holds since $\delta < 1$. As δ was chosen arbitrarily, Equation 7 holds.

Thus, Equation 7 implies:

$$C(3\mu^2)^{2m} \exp(-a\mu^2) \leq C \left(\frac{9m\mu^2}{a} \right)^m$$

and (b) holds.

Now if $m \leq \frac{9}{64}a\mu^2$, the last term above can be further upper bounded by $C \left(\frac{9m\mu^2}{a} \right)^m$. Thus $\mathbb{E}[(X^2 - \mu^2)^{2m}] \leq 3C \left(\frac{9m\mu^2}{a} \right)^m$ if $m \leq \frac{9}{64}a\mu^2$, and $\mathbb{E}[(X^2 - \mu^2)^{2m}] \leq 3C \left(\frac{64m^2}{a^2} \right)^m$ if $m > \frac{9}{64}a\mu^2$. This completes the proof of the fact. \square

F. Unitary t -designs

Definition 10. A monomial in elements of a matrix U is of degree (r, s) if it contains r conjugated elements and s unconjugated elements of U . We call it balanced if $r = s$ and will simply say a balanced monomial has degree t if it is of degree (t, t) . A polynomial is of degree t if it is a sum of balanced monomials of degree at most t .

Definition 11. A probability distribution ν on a finite set of $d \times d$ unitary matrices is said to be an ϵ -approximate unitary t -design if for all balanced monomials M of degree at most t , the following holds [11]:

$$\left| \mathbb{E}_{\nu} [M(U)] - \mathbb{E}_{\text{Haar}} [M(U)] \right| \leq \frac{\epsilon}{d^t}$$

If $\epsilon = 0$, we say that ν is an exact unitary t -design, or just unitary t -design.

For technical ease, we use quantum tensor product expanders (qTPEs) in place of unitary designs in our actual proofs. The formal definition of a qTPE follows.

Definition 12. A quantum t -tensor product expander (t -qTPE) in \mathcal{H} , $|\mathcal{H}| = d$, of degree s can be defined as a quantum operation $\mathcal{G} : L(\mathcal{H}^{\otimes t}) \rightarrow L(\mathcal{H}^{\otimes t})$ that can be expressed as $\mathcal{G}(M) = \frac{1}{s} \sum_{i=1}^s (U_i)^{\otimes t} M (U_i^{-1})^{\otimes t}$, for any matrix $M \in L(\mathcal{H}^{\otimes t})$, where $\{U_i\}_{i=1}^s$ are $d \times d$ unitary matrices. The qTPE is said to have second singular value λ if $\|\mathcal{G} - \mathcal{I}\|_\infty \leq \lambda$, where \mathcal{I} is the ‘ideal’ quantum operation defined by its action on a matrix M by $\mathcal{I}(M) := \int_{U \in \mathbb{U}(D)} U^{\otimes t} M (U^\dagger)^{\otimes t} d\text{Haar}(U)$. In other words, if $M \in L(\mathcal{H}^{\otimes t})$, then $\|\mathcal{G}(M) - \mathcal{I}(M)\|_2 \leq \lambda \|M\|_2$. We use the notation (d, s, λ, t) -qTPE to denote such a quantum tensor product expander.

A (d, s, λ, t) -qTPE can be sequentially iterated $O(\frac{t \log d + \log \epsilon^{-1}}{\log \lambda - 1})$ times to obtain an ϵ -approximate unitary t -design [11, Lemma 2.7]. For $t = \text{polylog}(d)$, efficient construction of t -qTPEs are known [9], [10].

III. PROOF OF THE MAIN THEOREM

We now state our main theorem in full detail.

Theorem 1. Consider a quantum state ρ^{AR} shared between a system A and a reference R . Let $\mathcal{T}^{A \rightarrow B}$ be a completely positive trace preserving superoperator with input system A and output system B . Let U be a unitary on the system A . Define the function

$$f(U) := \left\| [\mathcal{T}^{A \rightarrow B} \otimes \mathbb{I}^R] ((U^A \otimes I^R) \rho^{AR} (U^{A\dagger} \otimes I^R)) - \omega^B \otimes \rho^R \right\|_1,$$

where \mathbb{I}^R is the identity superoperator on R and I^R is the identity operator on R . Let A' be a new system having the same dimension as A . Define the Choi-Jamiołkowski state $\omega^{A'B} := (\mathcal{T}^{A \rightarrow B} \otimes \mathbb{I}^{A'}) (|\Phi\rangle\langle\Phi|^{AA'})$, of $\mathcal{T}^{A \rightarrow B}$ where $|\Phi\rangle^{AA'} := |A|^{-1/2} \sum_a |a\rangle^A \otimes |a\rangle^{A'}$ is the standard EPR state on system AA' . Let $0 < \epsilon, \delta < 1/3$. Let $\kappa > 0$. Then,

$$\mathbb{P}_{U^A \sim \text{TPE}} \left[f(U) > 2^{-\frac{1}{2} H_2^\epsilon(A|R)_\rho - \frac{1}{2} \mathbb{H}_2^{\epsilon, \delta}(A'|B)_\omega + 1} + 14\sqrt{\epsilon} + 2\kappa \right] \leq 7 \cdot 2^{-a\kappa^2}.$$

where U^A is a unitary matrix chosen uniformly at random from a $(|A|, s, \lambda, t)$ -quantum tensor product expander, $a := |A| \cdot 2^{-(1+\delta)\mathbb{H}_{\max}^\epsilon(B)_\omega + H_2^\epsilon(A|R)_\rho - 9}$, $t := 8a\kappa^2$, $\lambda := (|A|^{-8}|B|^{-6} \cdot \mu^2)^t$. As mentioned in Remark 1, by choosing parameters $\alpha := 2^{-\frac{1}{2} H_2^\epsilon(A|R)_\rho - \frac{1}{2} \mathbb{H}_2^{\epsilon, \delta}(A'|B)_\omega + 1}$ and $\beta := \sqrt{\frac{1}{a}}$ the above concentration inequality can also be expressed as:

$$\mathbb{P}_{U^A \sim \text{design}} [f(U) > \alpha + 14\sqrt{\epsilon} + \sqrt{t/2}\beta] \leq 7 \cdot 2^{-\frac{t}{8}}.$$

The quantity μ is defined as $\mu := \mathbb{E}_{\text{Haar}}[g(U)]$ for a related function $g(U)$ defined in Equation 11 below. We require that $\mu < 1$.

Remark 4. Such qTPEs can be obtained by sequentially iterating $O(t(\log |A| + \log |B| + \log \mu^{-1}))$ times an $(|A|, s, O(1), t)$ -qTPE, which is polynomial in t , $\log \mu^{-1}$ and the number of input and output qubits of the CPTP map \mathcal{T} [9], [10].

The proof is broken into three subsections. In the first subsection, we show that, for any probability distribution on U^A , instead of proving a tail bound for the given random variable $f(U)$, it suffices to prove a tail bound for a related random variable $g(U)$, where $f(U)$, $g(U)$ were informally defined in Section I-B. In the second subsection, we first obtain an upper bound on the Lipschitz constant of $g(U)$

which by Levy's lemma (Fact 20) leads to a tail bound for $g(U)$ where U is chosen from the Haar measure. We then obtain upper bounds on the centralised moments of $(g(U))^2$ under the Haar measure. Now $(g(U))^2$ is a balanced degree two polynomial in the matrix entries of U . In the final subsection, we apply Low's method to finally obtain a tail bound for $(g(U))^2$ for a uniformly random U chosen from a unitary design. This finishes the proof of Theorem 1.

A. From $f(U)$ to $g(U)$

We summarize various operators superoperators and their Choi-Jamiołkowski states that we will be working with, in order to establish the proof of Theorem 1, respectively in the following Table I and II

TABLE I
VARIOUS SUPEROPERATORS USED IN GOING FROM $f(U)$ TO $g(U)$.

$\mathcal{T}^{A \rightarrow B} :$	unperturbed decoupling CPTP superoperator with Stinespring unitary: $V_{\mathcal{T}}^{AC \rightarrow BZ}$
$\hat{\mathcal{T}}^{A \rightarrow B}(\cdot) :=$	$\text{Tr}_Z [(P^Z \otimes I^B) V_{\mathcal{T}} \circ (\cdot)]$: CP Trace non-increasing map; P comes from Lemma 10
$(\mathcal{T}')^{A \rightarrow B}(\cdot) :=$	$\Pi_{\omega_{\epsilon, \delta}^{B, \prime \prime \prime}} \circ \hat{\mathcal{T}}$: CP Trace non-increasing map

TABLE II
CHOI-JAMIOŁKOWSKI STATES OF SUPEROPERATORS IN TABLE I.

$\omega^{A'B} :=$	$(\mathcal{T} \otimes \mathcal{I}^{A'}) (\Phi^{AA'})$; $\text{Tr}(\omega^{A'B}) = 1$; Desired smoothing: $(\omega_{\epsilon, \delta}^{B, \prime \prime \prime})^B :=$ positive semidefinite operator obtained by zeroing out those eigenvalues of ω^B that are smaller than $2^{-(1+\delta)\mathbb{H}_{\max}^f(B)_\omega}$
$\eta^{A'B} :=$	$(\hat{\mathcal{T}} \otimes \mathcal{I}^{A'}) (\Phi^{AA'})$; $\text{Tr}(\eta) \leq 1$
$(\omega')^{A'B} :=$	$(\mathcal{T} \otimes \mathcal{I}^{A'}) (\Phi^{AA'})$; $= \Pi_{\omega_{\epsilon, \delta}^{B, \prime \prime \prime}} \circ \eta^{A'B}$ and $\text{Tr}[\omega'] \leq 1$

Recall that for a unitary U on the system A , we define the value taken by the decoupling function at U as follows:

$$f(U) := \left\| [\mathcal{T}^{A \rightarrow B} \otimes \mathbb{I}^R] ((U \otimes I^R) \rho^{AR} (U^\dagger \otimes I^R)) - \omega^B \otimes \rho^R \right\|_1.$$

Let $\eta^{AB} \leq \omega^{AB}$ be the positive semidefinite operator achieving the optimum in Definition 4. Let $V_{\mathcal{T}}^{AC \rightarrow BZ}$ be a unitary Stinespring dilation of the CPTP map $\mathcal{T}^{A \rightarrow B}$ provided by Fact 4. Thus

$$(\mathcal{T}^{A \rightarrow B} \otimes \mathbb{I}^{A'}) (M^{AA'}) = \text{Tr}_Z \left[(V_{\mathcal{T}}^{AC \rightarrow BZ} \otimes I^{A'}) (M^{AA'} \otimes |0\rangle\langle 0|^C) (V_{\mathcal{T}}^{AC \rightarrow BZ} \otimes I^{A'})^\dagger \right]$$

for any $M^{AA'} \in \mathcal{L}(A \otimes A')$, where A' is a new Hilbert space of the same dimension as A . Recall that $\omega^{A'B} := (\mathcal{T}^{A \rightarrow B} \otimes \mathbb{I}^{A'}) (\Phi^{AA'})$, where $\Phi^{AA'}$ is the standard EPR pure state on AA' . By Fact 10, there exists a POVM element on Z , $0^Z \leq P^Z \leq I^Z$, such that

$$(\hat{\mathcal{T}}^{A \rightarrow B} \otimes \mathbb{I}^{A'}) (\Phi^{AA'}) \tag{8}$$

$$\begin{aligned}
&:= \text{Tr}_Z \left[(P^Z \otimes I^{BA'}) (V_{\mathcal{T}}^{AC \rightarrow BZ} \otimes I^{A'}) (\Phi^{AA'} \otimes |0\rangle\langle 0|^C) (V_{\mathcal{T}}^{AC \rightarrow BZ} \otimes I^{A'})^\dagger (P^Z \otimes I^{BA'}) \right] \\
&= \eta^{BA'}.
\end{aligned}$$

The superoperator $\hat{\mathcal{T}}^{A \rightarrow B}$ is completely positive and trace non-increasing. Define the function

$$\hat{f}(U) := \left\| \left[\hat{\mathcal{T}}^{A \rightarrow B} \otimes \mathbb{I}^R \right] ((U^A \otimes I^R) \rho^{AR} (U^{A\dagger} \otimes I^R)) - \omega^B \otimes \rho^R \right\|_1.$$

By Fact 9,

$$(\hat{\mathcal{T}}^{A \rightarrow B} \otimes \mathbb{I}^R)((U \otimes I^R) \rho^{AR} (U^\dagger \otimes I^R)) \leq (\mathcal{T}^{A \rightarrow B} \otimes \mathbb{I}^R)((U \otimes I^R) \rho^{AR} (U^\dagger \otimes I^R)). \quad (9)$$

Hence by Fact 8, $f(U) \leq 2\hat{f}(U)$.

Let $\Pi_{\omega_{\epsilon, \delta}'''}^B$ be the projector onto the support of $(\omega_{\epsilon, \delta}''')^B$, where we recall from Definition 4 that $(\omega_{\epsilon, \delta}''')^B$ is the positive semidefinite operator obtained by zeroing out those eigenvalues of ω^B that are smaller than $2^{-(1+\delta)\mathbb{H}_{\max}^\epsilon(B)_\omega}$. Define the completely positive trace non-increasing superoperator $(\mathcal{T}')^{A \rightarrow B} := \Pi_{\omega_{\epsilon, \delta}'''}^B \circ \hat{\mathcal{T}}^{A \rightarrow B}$. From Definition 4 and Fact 13, we can conclude

$$\begin{aligned}
&\left\| \left[\hat{\mathcal{T}}^{A \rightarrow B} \otimes \mathbb{I}^R \right] ((U \otimes I^R) \rho^{AR} (U^\dagger \otimes I^R)) - \left[(\mathcal{T}')^{A \rightarrow B} \otimes \mathbb{I}^R \right] ((U \otimes I^R) \rho^{AR} (U^\dagger \otimes I^R)) \right\|_1 \\
&\leq 2\sqrt{\epsilon}
\end{aligned}$$

in fact, we can conclude that $\|\hat{\mathcal{T}}^{A \rightarrow B} - (\mathcal{T}')^{A \rightarrow B}\|_\diamond \leq 2\sqrt{\epsilon}$. Define the function

$$f'(U) := \left\| \left[(\mathcal{T}')^{A \rightarrow B} \otimes \mathbb{I}^R \right] ((U \otimes I^R) \rho^{AR} (U^\dagger \otimes I^R)) - \omega^B \otimes \rho^R \right\|_1.$$

By triangle inequality, $|f'(U) - \hat{f}(U)| \leq 2\sqrt{\epsilon}$ which further implies that $f(U) \leq 2f'(U) + 4\sqrt{\epsilon}$.

Define the states $(\rho')^{AR}$, ξ^R to be the ones achieving the optimum in Definition 1 of $H_2^\epsilon(A|R)_\rho$ i.e.

$$\|(I^A \otimes \xi^R)^{-1/4} (\rho')^{AR} (I^A \otimes \xi^R)^{-1/4}\|_2 = 2^{-\frac{1}{2}H_2^\epsilon(A|R)_\rho}. \quad (10)$$

Define the function

$$f''(U) := \left\| \left[(\mathcal{T}')^{A \rightarrow B} \otimes \mathbb{I}^R \right] ((U^A \otimes I^R) (\rho')^{AR} (U^{A\dagger} \otimes I^R)) - \omega^B \otimes (\rho')^R \right\|_1.$$

By triangle inequality, $|f''(U) - f'(U)| \leq 2\epsilon$ which implies that

$$f(U) \leq 2f''(U) + 8\sqrt{\epsilon}.$$

Now define the Hermitian positive semidefinite matrix $(\omega')^{A'B} := \left[(\mathcal{T}')^{A \rightarrow B} \otimes \mathbb{I}^{A'} \right] (\Phi^{AA'})$. Observe that $(\omega')^{A'B} = \Pi_{\omega_{\epsilon, \delta}'''}^B \circ \eta^{A'B}$. From Definition 4 and Fact 13, we have $\|(\omega')^{A'B} - \eta^{A'B}\|_1 \leq 2\sqrt{\epsilon}$ which further implies that $\|(\omega')^{A'B} - \omega^{A'B}\|_1 \leq 3\sqrt{\epsilon}$. Define the function

$$f'''(U) := \left\| \left[(\mathcal{T}')^{A \rightarrow B} \otimes \mathbb{I}^R \right] ((U \otimes I^R) (\rho')^{AR} (U^\dagger \otimes I^R)) - (\omega')^B \otimes (\rho')^R \right\|_1.$$

Again by triangle inequality, $|f'''(U) - f''(U)| \leq 3\sqrt{\epsilon}$ which implies that

$$f(U) \leq 2f'''(U) + 14\sqrt{\epsilon}.$$

Observe now that the range space of $(\mathcal{T}')^{A \rightarrow B}$ is contained in the support of $(\omega_{\epsilon, \delta}''')^B$. By Fact 12, we can upper bound $f'''(U)$ by the function $g(U)$ defined by

$$g(U) := \left\| \left[(\tilde{\mathcal{T}}')^{A \rightarrow B} \otimes \mathbb{I}^R \right] ((U^A \otimes I^R) (\tilde{\rho}')^{AR} (U^{A\dagger} \otimes I^R)) - (\tilde{\omega}')^B \otimes (\tilde{\rho}')^R \right\|_2, \quad (11)$$

where $(\tilde{\mathcal{T}}')^{A \rightarrow B} := ((\omega'''_{\epsilon, \delta})^{-1/4})^B \circ (\mathcal{T}')^{A \rightarrow B}$,

$$(\tilde{\rho}')^{AR} := (I^A \otimes \xi^R)^{-1/4} (\rho')^{AR} (I^A \otimes \xi^R)^{-1/4}, \quad (12)$$

and

$$(\tilde{\omega}')^{A'B} := ((\tilde{\mathcal{T}}')^{A \rightarrow B} \otimes \mathbb{I}^{A'}) (\Phi^{AA'}) = (I^{A'} \otimes (\omega'''_{\epsilon, \delta})^B)^{-1/4} (\omega')^{A'B} (I^{A'} \otimes (\omega'''_{\epsilon, \delta})^B)^{-1/4}. \quad (13)$$

Thus, $f(U) \leq 2g(U) + 14\sqrt{\epsilon}$. Recall from Definitions 1 and 4 respectively, that

$$2^{-\frac{1}{2}H_2^\epsilon(A|R)_\rho} = \|(\tilde{\rho}')^{AR}\|_2, \quad \text{and} \quad (14)$$

$$2^{-\frac{1}{2}\mathbb{H}_2^{\epsilon, \delta}(A'|B)_\rho} = \|(\tilde{\omega}')^{A'B}\|_2. \quad (15)$$

We have thus shown the following lemma.

Lemma 1. *Let $\mu, \kappa > 0$. For all probability distributions on U^A ,*

$$\mathbb{P}_{U^A}[f(U) > 2\mu + 14\sqrt{\epsilon} + 2\kappa] \leq \mathbb{P}_{U^A}[g(U) > \mu + \kappa].$$

In particular this holds for $\mu = 2^{-\frac{1}{2}H_2^\epsilon(A|R)_\rho - \frac{1}{2}\mathbb{H}_2^{\epsilon, \delta}(A'|B)_\omega} = \|(\tilde{\rho}')^{AR}\|_2 \cdot \|(\tilde{\omega}')^{A'B}\|_2$.

B. Bounding centralised moments of $(g(U))^2$ under Haar measure

We now upper bound the tail of $g(U)$ when U^A is chosen from the Haar measure. For this we need to upper bound the Lipschitz constant of $g(U)$ as follows.

Lemma 2. *The Lipschitz constant of $g(U)$, that is for unitary operators U, V , the ratio $\frac{|g(U) - g(V)|}{\|U - V\|_2}$ is less than $2^{\frac{1+\delta}{2}\mathbb{H}_{\max}^\epsilon(B)_\omega - \frac{1}{2}H_2^\epsilon(A|R)_\rho + 1}$.*

Proof. From Equations 12, 10 and Definition 1 recall that $\|(\tilde{\rho}')^{AR}\|_2 = 2^{-1/2H_2^\epsilon(A|R)}$. Write $(\tilde{\rho}')^{AR}$ in any canonical tensor basis for $A \otimes R$:

$$\begin{aligned} (\tilde{\rho}')^{AR} &= \sum_{ij} \sum_{kl} \tilde{\rho}'_{ijkl} |i\rangle\langle j|^A \otimes |k\rangle\langle l|^R = \sum_{kl} (\tilde{M}'_{kl})^A \otimes |k\rangle\langle l|^R \\ &= \sum_{kl} \sum_x s_x^{kl} |a_x^{kl}\rangle\langle b_x^{kl}|^A \otimes |k\rangle\langle l|^R, \end{aligned}$$

where $\tilde{M}'_{kl} := \sum_{ij} \tilde{\rho}'_{ijkl} |i\rangle\langle j|^R$, and $\tilde{M}'_{kl} = \sum_x s_x^{kl} |a_x^{kl}\rangle\langle b_x^{kl}|^A$ is the singular value decomposition of \tilde{M}'_{kl} .

Let $W_{\tilde{\mathcal{T}}'}^{AC \rightarrow BZ}$ be a Stinespring dilation of the completely positive trace non-increasing map $\tilde{\mathcal{T}}'^{A \rightarrow B}$ provided by Fact 4.

Thus $(\tilde{\mathcal{T}}')^{A \rightarrow B}(M^A) = \text{Tr}_Z[W_{\tilde{\mathcal{T}}'} \circ (M^A \otimes 0^C)]$, where $0^C := |0\rangle\langle 0|^C$. Note that

$$W_{\tilde{\mathcal{T}}'}^{AC \rightarrow BZ} = ((\omega'''_{\epsilon, \delta})^B \otimes I^Z)^{-1/4} (\Pi_{\omega'''_{\epsilon, \delta}}^B \otimes I^Z) (I^B \otimes P^Z) V_{\tilde{\mathcal{T}}'}^{AC \rightarrow BZ}, \quad (16)$$

where $V_{\tilde{\mathcal{T}}'}^{AC \rightarrow BZ}$ is the unitary Stinespring dilation of $\tilde{\mathcal{T}}'^{A \rightarrow B}$ provided by Fact 4 and P^Z , $\Pi_{\omega'''_{\epsilon, \delta}}^B$ and $(\omega'''_{\epsilon, \delta})^B$ are defined in Section III-A. We have

$$\|W_{\tilde{\mathcal{T}}'}^{AC \rightarrow BZ}\|_\infty \leq \|((\omega'''_{\epsilon, \delta})^B \otimes I^Z)^{-1/4}\|_\infty = \|((\omega'''_{\epsilon, \delta})^B)^{-1}\|_\infty^{1/4} \leq 2^{\frac{(1+\delta)}{4}\mathbb{H}_{\max}^\epsilon(B)_\omega}, \quad (17)$$

where the last inequality follows from the definition of $(\omega'''_{\epsilon, \delta})^B$ given in Definition 4. Moreover,

$$\|W_{\tilde{\mathcal{T}}'}^{AC \rightarrow BZ}\|_2 \leq \sqrt{|B||Z|} \|W_{\tilde{\mathcal{T}}'}^{AC \rightarrow BZ}\|_\infty \leq \sqrt{|B||A|} \cdot 2^{\frac{(1+\delta)}{4}\mathbb{H}_{\max}^\epsilon(B)_\omega}, \quad (18)$$

where we used that $|Z| \leq |A|$ guaranteed by Fact 4.

Let U^A, V^A be two unitaries on A . Then,

$$|g(U) - g(V)| \quad (19)$$

$$\begin{aligned} &\leq \left\| \left[(\tilde{T}')^{A \rightarrow B} \otimes \mathbb{I}^R \right] ((U^A \otimes I^R) \circ (\tilde{\rho}')^{AR}) - \left[(\tilde{T}')^{A \rightarrow B} \otimes \mathbb{I}^R \right] ((V^A \otimes I^R) \circ (\tilde{\rho}')^{AR}) \right\|_2 \\ &= \left\| \left[(\tilde{T}')^{A \rightarrow B} \otimes \mathbb{I}^R \right] ((U^A \otimes I^R) \circ (\tilde{\rho}')^{AR} - (V^A \otimes I^R) \circ (\tilde{\rho}')^{AR}) \right\|_2 \\ &\leq \left\| \left[(\tilde{T}')^{A \rightarrow B} \otimes \mathbb{I}^R \right] ((U^A \otimes I^R)(\tilde{\rho}')^{AR}(U^A \otimes I^R)^\dagger - (U^A \otimes I^R)(\tilde{\rho}')^{AR}(V^A \otimes I^R)^\dagger) \right\|_2 \\ &\quad + \left\| \left[(\tilde{T}')^{A \rightarrow B} \otimes \mathbb{I}^R \right] ((U^A \otimes I^R)(\tilde{\rho}')^{AR}(V^A \otimes I^R)^\dagger - (V^A \otimes I^R)(\tilde{\rho}')^{AR}(V^A \otimes I^R)^\dagger) \right\|_2. \end{aligned} \quad (20)$$

We now upper bound

$$\begin{aligned} &\left\| \left[(\tilde{T}')^{A \rightarrow B} \otimes \mathbb{I}^R \right] ((U^A \otimes I^R)(\tilde{\rho}')^{AR}(U^A \otimes I^R)^\dagger - (U^A \otimes I^R)(\tilde{\rho}')^{AR}(V^A \otimes I^R)^\dagger) \right\|_2 \\ &= \left\| \sum_{kl} \left(\sum_x s_x^{kl} (\tilde{T}')^{A \rightarrow B} ((U|a_x^{kl}\rangle^A \langle b_x^{kl}|U^\dagger) - (U|a_x^{kl}\rangle^A \langle b_x^{kl}|V^\dagger)) \right) \otimes |k\rangle^R \langle l| \right\|_2 \\ &= \sqrt{\sum_{kl} \left\| \sum_x s_x^{kl} (\tilde{T}')^{A \rightarrow B} ((U|a_x^{kl}\rangle^A \langle b_x^{kl}|U^\dagger) - (U|a_x^{kl}\rangle^A \langle b_x^{kl}|V^\dagger)) \right\|_2^2}. \end{aligned}$$

Fix k, l . For ease of notation drop the superscript kl below. We now upper bound

$$\begin{aligned} &\left\| \sum_x s_x (\tilde{T}')^{A \rightarrow B} ((U|a_x\rangle^A \langle b_x|U^\dagger) - (U|a_x\rangle^A \langle b_x|V^\dagger)) \right\|_2 \\ &= \left\| \sum_x s_x \left(\text{Tr}_Z [W_{\tilde{T}'} \circ ((U|a_x\rangle^A \langle b_x|U^\dagger) \otimes 0^C)] - \text{Tr}_Z [W_{\tilde{T}'} \circ ((U|a_x\rangle^A \langle b_x|V^\dagger) \otimes 0^C)] \right) \right\|_2 \\ &\stackrel{a}{=} \left\| \sum_x s_x \left(P_{x,U}^{B \times Z} (Q_{x,U}^{B \times Z})^\dagger - P_{x,U}^{B \times Z} (Q_{x,V}^{B \times Z})^\dagger \right) \right\|_2 \\ &\stackrel{b}{=} \left\| P_U^{BQ \times ZA} (Q_U^{BQ \times ZA})^\dagger - P_U^{BQ \times ZA} (Q_V^{BQ \times ZA})^\dagger \right\|_2 \\ &\leq \left\| P_U^{BQ \times ZA} \right\|_2 \left\| Q_U^{BQ \times ZA} - Q_V^{BQ \times ZA} \right\|_2 \\ &\stackrel{c}{\leq} 2^{\frac{1+\delta}{2} \mathbb{H}_{\max}^\epsilon(B)_\omega} \|\tilde{M}'_{kl}\|_2 \|U - V\|_2, \end{aligned}$$

where

(a)

$$\begin{aligned} P_{x,U}^{B \times Z} &:= (\text{vec}^{B,Z})^{-1} (W_{\tilde{T}'}^{AC \rightarrow BZ} (U^A \otimes I^C) (|a_x\rangle^A \otimes |0\rangle^C)), \\ Q_{x,U}^{B \times Z} &:= (\text{vec}^{B,Z})^{-1} (W_{\tilde{T}'}^{AC \rightarrow BZ} (U^A \otimes I^C) (|b_x\rangle^A \otimes |0\rangle^C)), \end{aligned}$$

and $Q_{x,V}^{B \times Z}$ is defined similarly. The above operators map system Z to system B or are $B \times Z$ matrices for fixed bases of B and Z . The equality holds due to Fact 6.

(b) Let Q be a single qubit register and x range over the computational basis of A .

$$P_U^{BQ \times ZA} := \sum_x s_x (P_{x,U}^{B \times Z} \otimes |0\rangle^Q \langle x|^A), \quad Q_U^{BQ \times ZA} := \sum_x (Q_{x,U}^{B \times Z} \otimes |0\rangle^Q \langle x|^A),$$

and $Q_V^{BQ \times ZA}$ is defined similarly. The equality follows by inspection.

(c) We have

$$\begin{aligned} \|P_{U,kl}^{BQ \times ZA}\|_2^2 &= \sum_x (s_x^{kl})^2 \|P_{x,U,kl}^{B \times Z}\|_2^2 = \sum_x (s_x^{kl})^2 \|W_{\tilde{T}'}^{AC \rightarrow BZ}(U^A \otimes I^C)(|a_x^{kl}\rangle^A \otimes |0\rangle^C)\|_2^2 \\ &\leq \|W_{\tilde{T}'}^{AC \rightarrow BZ}\|_\infty^2 \sum_x (s_x^{kl})^2 \leq 2^{\frac{1+\delta}{2} \mathbb{H}_{\max}^\epsilon(B)\omega} \|\tilde{M}'_{kl}\|_2^2, \end{aligned}$$

where the last inequality follows from Equation 17. Again using Equation 17 in the second and third inequality below, we get

$$\begin{aligned} &\|Q_U^{BQ \times ZA} - Q_V^{BQ \times ZA}\|_2^2 \\ &= \sum_x \|Q_{x,U}^{B \times Z} - Q_{x,V}^{B \times Z}\|_2^2 \\ &= \sum_x \|W_{\tilde{T}'}^{AC \rightarrow BZ}(U^A \otimes I^C)(|b_x\rangle^A \otimes |0\rangle^C) - W_{\tilde{T}'}^{AC \rightarrow BZ}(V^A \otimes I^C)(|b_x\rangle^A \otimes |0\rangle^C)\|_2^2 \\ &= \sum_x \|W_{\tilde{T}'}^{AC \rightarrow BZ}((U^A \otimes I^C) - (V^A \otimes I^C))(|b_x\rangle^A \otimes |0\rangle^C)\|_2^2 \\ &\leq \|W_{\tilde{T}'}^{AC \rightarrow BZ}\|_\infty^2 \sum_x \|((U^A \otimes I^C) - (V^A \otimes I^C))(|b_x\rangle^A \otimes |0\rangle^C)\|_2^2 \\ &\leq \|((\omega_{\epsilon,\delta}''')^B)^{-1}\|_\infty^{1/2} \sum_x \|((U^A \otimes I^C) - (V^A \otimes I^C))(|b_x\rangle^A \otimes |0\rangle^C)\|_2^2 \\ &= \|((\omega_{\epsilon,\delta}''')^B)^{-1}\|_\infty^{1/2} \sum_x \|(U - V)^A |b_x\rangle\|_2^2 \\ &= \|((\omega_{\epsilon,\delta}''')^B)^{-1}\|_\infty^{1/2} \sum_x \langle b_x | (U - V)^\dagger (U - V) | b_x \rangle \\ &= \|((\omega_{\epsilon,\delta}''')^B)^{-1}\|_\infty^{1/2} \text{Tr}[(U - V)^\dagger (U - V)] \leq 2^{\frac{1+\delta}{2} \mathbb{H}_{\max}^\epsilon(B)\omega} \|U - V\|_2^2. \\ &\Rightarrow \|P_U^{BQ \times ZA}\|_2 \|Q_U^{BQ \times ZA} - Q_V^{BQ \times ZA}\|_2 \leq 2^{\frac{1+\delta}{2} \mathbb{H}_{\max}^\epsilon(B)\omega} \|\tilde{M}'_{kl}\|_2 \|U - V\|_2, \text{ thereby proving the} \\ &\text{desired inequality.} \end{aligned}$$

This implies that

$$\begin{aligned} &\left\| \left[(\tilde{T}')^{A \rightarrow B} \otimes \mathbb{I}^R \right] ((U \otimes I)(\tilde{\rho}')^{AR}(U \otimes I)^\dagger - (U \otimes I)(\tilde{\rho}')^{AR}(V \otimes I)^\dagger) \right\|_2 \\ &\leq \sqrt{\sum_{kl} 2^{(1+\delta) \mathbb{H}_{\max}^\epsilon(B)\omega} \|\tilde{M}'_{kl}\|_2^2 \|U - V\|_2^2} \\ &= 2^{\frac{1+\delta}{2} \mathbb{H}_{\max}^\epsilon(B)\omega} \|U - V\|_2 \sqrt{\sum_{kl} \|\tilde{M}'_{kl}\|_2^2} = 2^{\frac{1+\delta}{2} \mathbb{H}_{\max}^\epsilon(B)\omega} \|U - V\|_2 \|(\tilde{\rho}')^{AR}\|_2. \end{aligned}$$

Hence $|g(U) - g(V)| \leq 2 \cdot 2^{\frac{1+\delta}{2} \mathbb{H}_{\max}^\epsilon(B)\omega} \|(\tilde{\rho}')^{AR}\|_2 \|U - V\|_2$. This completes the proof of the lemma. \square

Note 1. The proof technique to find a tighter Lipschitz constant of the function g (as opposed to the naive one) entails the evaluation of the Schatten-2 norm of the partial isometry $W_{\tilde{T}'}^{AC \rightarrow BZ}$. This is imperative in finding the closeness between the higher order moments of the function $g(U)^2$ under Haar measure and quantum tensor product expander in Lemma 6. Hence we re-write Equation 18 below:

$$\|W_{\tilde{T}'}^{AC \rightarrow BZ}\|_2 \leq \sqrt{|B||Z|} \|W_{\tilde{T}'}^{AC \rightarrow BZ}\|_\infty \leq \sqrt{|B||A|} \cdot 2^{\frac{(1+\delta)}{4} \mathbb{H}_{\max}^\epsilon(B)_\omega}.$$

Lemma 3. For any unitary $U \in \mathbb{U}(A)$, $g(U) \leq (2|A|)^{1/2} \cdot 2^{\frac{1+\delta}{2} \mathbb{H}_{\max}^\epsilon(B)_\omega - \frac{1}{2} H_2^\epsilon(A|R)_\rho}$.

Proof. Define the Hermitian matrix $\gamma^{AR} := (\tilde{\rho}')^{AR} - \pi^A \otimes (\tilde{\rho}')^R$. We have

$$\begin{aligned} \|\gamma^{AR}\|_2^2 &= \|(\tilde{\rho}')^{AR}\|_2^2 + \|\pi^A \otimes (\tilde{\rho}')^R\|_2^2 - \langle (\tilde{\rho}')^{AR}, (\pi^A \otimes (\tilde{\rho}')^R) \rangle - \langle (\pi^A \otimes (\tilde{\rho}')^R), (\tilde{\rho}')^{AR} \rangle \\ &\leq \|(\tilde{\rho}')^{AR}\|_2^2 + \|\pi^A \otimes (\tilde{\rho}')^R\|_2^2 \leq 2\|(\tilde{\rho}')^{AR}\|_2^2, \end{aligned} \quad (21)$$

where we used the fact that $(\tilde{\rho}')^{AR}$, $(\pi^A \otimes (\tilde{\rho}')^R)$ are positive semidefinite matrices in the first inequality and Fact 11 in the second inequality.

Observe that $g(U) = \left\| [(\tilde{T}')^{A \rightarrow B} \otimes \mathbb{I}^R] ((U^A \otimes I^R) \circ \gamma^{AR}) \right\|_2$. Arguing similarly as in the proof of Lemma 2 (proceeding along the similar lines as in Equations 19, 20 without the second term inside the norm and similarly leading to the inequalities (a), (b) and (c) without involving their respective second terms which are subtracted), we can conclude that

$$g(U) \leq 2^{\frac{1+\delta}{2} \mathbb{H}_{\max}^\epsilon(B)_\omega} \|\gamma^{AR}\|_2 \cdot \|U\|_2 \leq (2|A|)^{1/2} \cdot 2^{\frac{1+\delta}{2} \mathbb{H}_{\max}^\epsilon(B)_\omega - \frac{1}{2} H_2^\epsilon(A|R)_\rho},$$

where we have used $\|U\|_2 = |A|^{1/2}$ and Equation 15 in the last inequality. This completes the proof. \square

We now apply Levy's Lemma (Fact 20) to obtain an exponential upper bound on the deviation of $g(U)$ about its expectation μ when U is chosen from the Haar measure.

Proposition 1. Define $\mu := \mathbb{E}_{U \sim \text{Haar}}[g(U)]$. Define $a := |A| 2^{H_2^\epsilon(A|R)_\rho - (1+\delta) \mathbb{H}_{\max}^\epsilon(B)_\omega - 4}$. Let $\kappa > 0$. Then

$$\mathbb{P}_{U \sim \text{Haar}}[|g(U) - \mu| > \kappa] \leq 2 \exp(-a\kappa^2). \quad (22)$$

Additionally,

$$\mu^2 \leq \mathbb{E}_{U \sim \text{Haar}}[(g(U))^2] = \alpha \|(\tilde{\rho}')^R\|_2^2 + \beta \|(\tilde{\rho}')^{AR}\|_2^2 - \|(\tilde{\omega}')^B\|_2^2 \|(\tilde{\rho}')^R\|_2^2 \quad (23)$$

$$< \|(\tilde{\omega}')^{A'B}\|_2^2 \cdot \|(\tilde{\rho}')^{AR}\|_2^2 = 2^{-\mathbb{H}_2^{\epsilon, \delta}(A'|B)_\omega - H_2^\epsilon(A|R)_\rho}, \quad (24)$$

where $(\tilde{\omega}')^{A'B}$, $(\tilde{\rho}')^{AR}$ are defined in Equations 13, 12 respectively,

$$\delta_1 := \|(\tilde{\omega}')^B\|_2^2 \frac{|A|^2 - |A|\eta}{|A|^2 - 1}, \quad \delta_2 := \|(\tilde{\omega}')^{A'B}\|_2^2 \frac{|A|^2 - |A|\eta^{-1}}{|A|^2 - 1}, \quad \eta := \frac{\|(\tilde{\omega}')^{A'B}\|_2^2}{\|(\tilde{\omega}')^B\|_2^2}.$$

Moreover if

$$a \cdot \mathbb{E}_{U \sim \text{Haar}}[(g(U))^2] + \log \mathbb{E}_{U \sim \text{Haar}}[(g(U))^2] > \log |A| + \mathbb{H}_{\max}^\epsilon(B)_\omega - H_2^\epsilon(A|R)_\rho, \text{ then}$$

$$\mathbb{E}_{U \sim \text{Haar}}[(g(U))^2] \leq 8\mu^2. \quad (25)$$

Proof. Equations (3.31, 3.32, 3.33, 3.37 and 3.43) in [3] and Equations (3.44, 3.45 and 3.46) used in the proof of Fact 1 (in [3]) implies Equation 23 and Equation 24 respectively. Fact 20 applied to the function $g(U)$ with upper bound on the Lipschitz constant given by Lemma 2 gives Equation 22. Further,

$$\mathbb{E}_{U \sim \text{Haar}}[(g(U))^2]$$

$$\begin{aligned}
&\stackrel{a}{=} \mathbb{E}_{U \sim \text{Haar}} \left[(g(U))^2 \mathbf{1} \left\{ g(U) \leq \left(\mu + 2^{-1} \left\{ \mathbb{E}_{U \sim \text{Haar}} [(g(U))^2] \right\}^{1/2} \right) \right\} \right] \\
&\quad + \mathbb{E}_{U \sim \text{Haar}} \left[(g(U))^2 \mathbf{1} \left\{ g(U) \leq \left(\mu + 2^{-1} \left\{ \mathbb{E}_{U \sim \text{Haar}} [(g(U))^2] \right\}^{1/2} \right) \right\} \right] \\
&\stackrel{b}{\leq} \left(\mu + 2^{-1} \left\{ \mathbb{E}_{U \sim \text{Haar}} [(g(U))^2] \right\}^{1/2} \right)^2 \\
&\quad + \left[(2|A|)^{1/2} \cdot 2^{\frac{1+\delta}{2} \mathbb{H}_{\max}^{\epsilon}(B)_{\omega} - \frac{1}{2} H_2^{\epsilon}(A|R)_{\rho}} \right]^2 \cdot \mathbb{P}_{U \sim \text{Haar}} \left[g(U) - \mu > 2^{-1} \left\{ \mathbb{E}_{U \sim \text{Haar}} [(g(U))^2] \right\}^{1/2} \right] \\
&\leq 2\mu^2 + 2^{-1} \mathbb{E}_{U \sim \text{Haar}} [(g(U))^2] \\
&\quad + \left[(2|A|)^{1/2} \cdot 2^{\frac{1+\delta}{2} \mathbb{H}_{\max}^{\epsilon}(B)_{\omega} - \frac{1}{2} H_2^{\epsilon}(A|R)_{\rho}} \right]^2 \cdot 2 \exp(-2^{-2} a \cdot \mathbb{E}_{U \sim \text{Haar}} [(g(U))^2]) \\
&\leq 2\mu^2 + \frac{3}{4} \mathbb{E}_{U \sim \text{Haar}} [(g(U))^2].
\end{aligned}$$

where in (a) the notation $\mathbf{1} \left\{ g(U) \leq \left(\mu + 2^{-1} \left[\mathbb{E}_{U \sim \text{Haar}} [(g(U))^2] \right]^{1/2} \right) \right\}$ denotes the indicator function of the event in its argument, taking a value one if the event occurs and zero otherwise; (b) holds because the first summand uses the fact $\mathbb{E}(\mathbf{1}(\text{event})) = \mathbb{P}(\text{event}) \leq 1$ and we use Lemma 3 to upper bound the second summand.

This implies that $\mathbb{E}_{U \sim \text{Haar}} [(g(U))^2] \leq 8\mu^2$, completing the proof of Equation 25 and hence of the proposition. \square

We now evaluate the higher order moments of the functions $g(U) - \mu$ and $(g(U))^2 - \mu^2$.

Lemma 4. Define $\mu := \mathbb{E}_{U \sim \text{Haar}} [g(U)]$. Let m be a positive integer. Then the $(2m)$ -th moments of the functions $g(U) - \mu$ and $(g(U))^2 - \mu^2$ are upper bounded by

$$\mathbb{E}_{U \sim \text{Haar}} (|g(U) - \mu|^{2m}) \leq 2(m 2^{(1+\delta) \mathbb{H}_{\max}^{\epsilon}(B)_{\omega} - H_2^{\epsilon}(A|R)_{\rho} + 4} |A|^{-1})^m,$$

$$\begin{aligned}
&\mathbb{E}_{U \sim \text{Haar}} (|(g(U))^2 - \mu^2|^{2m}) \\
&\leq \begin{cases} 6(9m\mu^2 2^{(1+\delta) \mathbb{H}_{\max}^{\epsilon}(B)_{\omega} - H_2^{\epsilon}(A|R)_{\rho} + 4} |A|^{-1})^m & ; m \leq \frac{9}{64} |A| \mu^2 \cdot 2^{-(1+\delta) \mathbb{H}_{\max}^{\epsilon}(B)_{\omega} + H_2^{\epsilon}(A|R)_{\rho} - 4} \\ 6(64m^2 2^{2(1+\delta) \mathbb{H}_{\max}^{\epsilon}(B)_{\omega} - 2H_2^{\epsilon}(A|R)_{\rho} + 8} |A|^{-2})^m & ; \text{otherwise} \end{cases}.
\end{aligned}$$

Proof. Applying Fact 21 to the non-negative random variable $|g(U) - \mu|$ with concentration given by Proposition 1 gives the bounds of the lemma. \square

Note 2. It might seem apparent at this point that one can directly apply [11, Lemma 3.4] to obtain the expectation of the function $[g^2(U) - \mu^2]^{2m}$ and then apply Markov's inequality to conclude the result. However, there is a technical subtlety here. Low's Lemma 3.4 in [11] has an additive parameter ' α ', which is the sum of the coefficients in the polynomial expansion of $g^2(U)$. Since the function g depends the perturbed superoperator \mathcal{T}' (and not \mathcal{T}), finding the value of α is more complicated than our subsequent analysis (especially Lemma 6). Furthermore, applying Low's technique at this point with crude approximation (still not so easy) will give a much larger value of t than what we get by appealing to the definition of quantum tensor product expanders. It might not be feasible to conclude that under any assumptions computationally efficient designs can lead to decoupling. In contrast, our analysis results in computationally efficient decoupling under the setting Theorem 2, for $|A_1| = \text{poly}(\log |A_2|)$ and $\kappa = O(\mu)$. Hence our subsequent method that leads to Lemma 6 and 7 are needed to obtain a handle

on the parameters λ , m of the quantum tensor product expanders. This provides a way to analyze how large a value of the unitary design parameter t is required to obtain an exponentially decaying tail for any deviation κ .

C. Concentration of $(g(U))^2$ under t -design

In this section we finally obtain an exponential concentration for $(g(U))^2$ when U is chosen uniformly at random from a unitary t -design for suitable t . We first prove the following lemma.

Lemma 5. *Let $\mathcal{T}^{A \rightarrow B}$ be a completely positive superoperator with Stinespring dilation $W_{\mathcal{T}}^{AC \rightarrow BZ}$, where $|A||C| = |B||Z|$, the input ancillary system is C and the output ancillary system is D . Let $F^{A_1 A_2}$ and $F^{B_1 B_2}$ be the appropriate swap operators. Then*

$$\|((\mathcal{T}^\dagger)^{B_1 \rightarrow A_1} \otimes (\mathcal{T}^\dagger)^{B_2 \rightarrow A_2})(F^{B_1 B_2})\|_2 = \|(\mathcal{T}^{A_1 \rightarrow B_1} \otimes \mathcal{T}^{A_2 \rightarrow B_2})(F^{A_1 A_2})\|_2 \leq \|W_{\mathcal{T}}\|_2^4.$$

Proof. By Stinespring representation of \mathcal{T} as given in Fact 4, $\mathcal{T}^{A \rightarrow B}(M^A) = \text{Tr}_Z [W_{\mathcal{T}}(M^A \otimes |0\rangle\langle 0|^C)W_{\mathcal{T}}]$ for any $M^A \in \mathcal{L}(A)$. Expressing the swap operator $F^{A_1 A_2}$ in computational basis, we have

$$F^{A_1 A_2} = \sum_{aa'} \{|a'\rangle\langle a|\} \{|a\rangle\langle a'|\}^{A_1 A_2} = \sum_{aa'} |a'\rangle\langle a|^{A_1} \otimes |a\rangle\langle a'|^{A_2}.$$

Note that the swap operator is Hermitian. Further, observe that

$$\begin{aligned} & \|[(\mathcal{T}^\dagger)^{B_1 \rightarrow A_1} \otimes (\mathcal{T}^\dagger)^{B_2 \rightarrow A_2}](F^{B_1 B_2})\|_2^2 \\ & \stackrel{a}{=} \text{Tr} \left[\{((\mathcal{T}^\dagger)^{B_1 \rightarrow A_1} \otimes (\mathcal{T}^\dagger)^{B_2 \rightarrow A_2})(F^{B_1 B_2})\}^2 \right] \\ & \stackrel{b}{=} \text{Tr} \left[\{((\mathcal{T}^\dagger \otimes \mathcal{T}^\dagger)F^{B_1 B_2}) \otimes ((\mathcal{T}^\dagger \otimes \mathcal{T}^\dagger)F^{B'_1 B'_2})\} F^{(A_1 A_2)(A'_1 A'_2)} \right] \\ & \stackrel{c}{=} \text{Tr} \left[(F^{B_1 B_2} \otimes F^{B'_1 B'_2}) \left\{ [(\mathcal{T}^{A_1 \rightarrow B_1} \otimes \mathcal{T}^{A_2 \rightarrow B_2}) \otimes (\mathcal{T}^{A'_1 \rightarrow B'_1} \otimes \mathcal{T}^{A'_2 \rightarrow B'_2})] F^{(A_1 A_2)(A'_1 A'_2)} \right\} \right] \\ & \stackrel{d}{=} \text{Tr} \left[F^{(B_1 B'_1)(B_2 B'_2)} \left\{ [(\mathcal{T}^{A_1 \rightarrow B_1} \otimes \mathcal{T}^{A'_1 \rightarrow B'_1})] \otimes [\mathcal{T}^{A_2 \rightarrow B_2} \otimes \mathcal{T}^{A'_2 \rightarrow B'_2}](F^{A_1 A'_1} \otimes F^{A_2 A'_2}) \right\} \right] \\ & = \text{Tr} \left[F^{(B_1 B'_1)(B_2 B'_2)} \left\{ [\mathcal{T}^{A_1 \rightarrow B_1} \otimes \mathcal{T}^{A'_1 \rightarrow B'_1}](F^{A_1 A'_1}) \otimes [\mathcal{T}^{A_2 \rightarrow B_2} \otimes \mathcal{T}^{A'_2 \rightarrow B'_2}](F^{A_2 A'_2}) \right\} \right] \\ & \stackrel{e}{=} \text{Tr} \left[\left\{ [\mathcal{T}^{A_1 \rightarrow B_1} \otimes \mathcal{T}^{A'_1 \rightarrow B'_1}](F^{A_1 A'_1}) \right\}^2 \right] \\ & = \|[\mathcal{T}^{A_1 \rightarrow B_1} \otimes \mathcal{T}^{A_2 \rightarrow B_2}](F^{A_1 A_2})\|_2^2, \end{aligned}$$

where in

- (a) we use the fact that \mathcal{T}^\dagger is completely positive as \mathcal{T} is completely positive and the fact that the swap operator is Hermitian, implying that $((\mathcal{T}^\dagger)^{B_1 \rightarrow A_1} \otimes (\mathcal{T}^\dagger)^{B_2 \rightarrow A_2})(F^{B_1 B_2})$ is Hermitian,
- (b) we take A'_1, A'_2 to be two new systems of the same dimension as A , $F^{(A_1 A_2)(A'_1 A'_2)}$ as the operator swapping $(A_1 A_2)$ with $(A'_1 A'_2)$ and Fact 15,
- (c) we use the definition of the adjoint of a superoperator under the Hilbert-Schmidt inner product,
- (d) we use a property of the swap operator,
- (e) we use Fact 15.

This proves the first of the equalities asserted above.

Finally,

$$\|[\mathcal{T}^{A_1 \rightarrow B_1} \otimes \mathcal{T}^{A_2 \rightarrow B_2}](F^{A_1 A_2})\|_2$$

$$\begin{aligned}
&= \left\| \sum_{aa'} \left\{ \left(\text{Tr}_Z \left[W_{\mathcal{T}}(|a'\rangle\langle a|^A \otimes |0\rangle\langle 0|^C) W_{\mathcal{T}}^\dagger \right] \right)^{B_1} \otimes \left(\text{Tr}_Z \left[W_{\mathcal{T}}(|a\rangle\langle a'|^{A_2} \otimes |0\rangle\langle 0|^C) W_{\mathcal{T}}^\dagger \right] \right)^{B_2} \right\} \right\|_2 \\
&= \left\| \sum_{aa'} \left\{ \left(\text{Tr}_Z \left[W_{\mathcal{T}}(|a'\rangle\langle 0|)(\langle a|\langle 0|)^{A_1 C} W_{\mathcal{T}}^\dagger \right] \right)^{B_1} \otimes \left(\text{Tr}_Z \left[W_{\mathcal{T}}(|a\rangle\langle 0|)(\langle a'|\langle 0|)^{A_2 C} W_{\mathcal{T}}^\dagger \right] \right)^{B_2} \right\} \right\|_2 \\
&\stackrel{a}{=} \left\| \sum_{aa'} (P_{a'} P_a^\dagger)^{B_1} \otimes (P_a P_{a'}^\dagger)^{B_2} \right\|_2 \leq \sum_{aa'} \left\| (P_{a'} P_a^\dagger)^{B_1} \otimes (P_a P_{a'}^\dagger)^{B_2} \right\|_2 \\
&= \sum_{aa'} \left\| (P_{a'} P_a^\dagger)^{B_1} \right\|_2 \cdot \left\| (P_a P_{a'}^\dagger)^{B_2} \right\|_2 \leq \sum_{aa'} \|P_{a'}^{B \times Z}\|_2^2 \cdot \|P_a^{B \times Z}\|_2^2 = \left(\sum_a \|P_a^{B \times Z}\|_2^2 \right)^2 \\
&= \left(\sum_a \left\| (W_{\mathcal{T}}(|a\rangle^A \otimes |0\rangle^C))^{BZ} \right\|_2^2 \right)^2 \leq \left(\sum_{ac} (\langle a|^A \langle c|^C) W_{\mathcal{T}}^\dagger W_{\mathcal{T}} (|a\rangle^A |c\rangle^C) \right)^2 \\
&= \left(\text{Tr} [W_{\mathcal{T}}^\dagger W_{\mathcal{T}}] \right)^2 = \|W_{\mathcal{T}}\|_2^4,
\end{aligned}$$

where in (a) we define $P_a^{B \times Z} := (\text{vec}^{B, Z})^{-1}((W_{\mathcal{T}}(|a\rangle^A |0\rangle^C))^{BZ})$ and use Fact 6. This completes the proof of the lemma. \square

Note that $(g(U))^2$ is a balanced degree two polynomial in the matrix entries of U . We now find out how close the moments of $(g(U))^2$ under Haar measure are to their counterparts under t -design.

We do this by using the following Proposition 2.

Proposition 2. *The $(2i)^{\text{th}}$ power of the function $g(U)$ which is equivalent to the simplification of the balanced degree two polynomial $[g(U)^2]^i$ is given by:*

$$\begin{aligned}
&(g(U)^2)^i \\
&= \text{Tr} \left[\{(\gamma')^{A_1 A_2}\}^{\otimes i} \cdot \left\{ (U^\dagger)^{A_1} \otimes (U^\dagger)^{A_2} \right\}^{\otimes i} \circ \left[((\tilde{\mathcal{T}}')^\dagger)^{B_1 \rightarrow A_1} \otimes ((\tilde{\mathcal{T}}')^\dagger)^{B_2 \rightarrow A_2} \right] (F^{B_1 B_2}) \right\}^{\otimes i} \Big],
\end{aligned}$$

where we define the Hermitian matrices $\gamma^{AR} := (\tilde{\rho}')^{AR} - \pi^A \otimes (\tilde{\rho}')^R$ and $(\gamma')^{A_1 A_2} := \text{Tr}_{R_1 R_2} \{ (I^{A_1 A_2} \otimes F^{R_1 R_2}) (\gamma^{A_1 R_1} \otimes \gamma^{A_2 R_2}) \}$.

Proof. Let A_1, A_2 denote two Hilbert spaces of the same dimension as A ; similarly for B_1, B_2 and R_1, R_2 . Further, $A_1(j), 1 \leq j \leq i$ denote Hilbert spaces of the same dimension as A_1 , likewise for $A_2(j), B_1(j), B_2(j), R_1(j)$ and $R_2(j)$. Observe that

$$\begin{aligned}
&(g(U)^2)^i \\
&= \left(\left\| [(\tilde{\mathcal{T}}')^{A \rightarrow B} \otimes \mathbb{I}^R] ((U^A \otimes I^R) \circ (\tilde{\rho}')^{AR}) - (\tilde{\omega}'^B) \otimes (\tilde{\rho}')^R \right\|_2^2 \right)^i \\
&= \left(\text{Tr} \left[\left\{ [(\tilde{\mathcal{T}}')^{A \rightarrow B} \otimes \mathbb{I}^R] ((U^A \otimes I^R) \circ \gamma^{AR}) \right\}^2 \right] \right)^i \\
&= \left(\text{Tr} \left[\left\{ [(\tilde{\mathcal{T}}')^{A_1 \rightarrow B_1} \otimes \mathbb{I}^{R_1}] ((U^{A_1} \otimes I^{R_1}) \circ \gamma^{A_1 R_1}) \right\} \right. \right. \\
&\quad \left. \left. \otimes \left\{ [(\tilde{\mathcal{T}}')^{A_2 \rightarrow B_2} \otimes \mathbb{I}^{R_2}] ((U^{A_2} \otimes I^{R_2}) \circ \gamma^{A_2 R_2}) \right\} (F^{B_1 B_2} \otimes F^{R_1 R_2}) \right] \right)^i
\end{aligned}$$

$$\begin{aligned}
&= \left(\text{Tr} \left[(I^{A_1 A_2} \otimes F^{R_1 R_2}) (\gamma^{A_1 R_1} \otimes \gamma^{A_2 R_2}) \right. \right. \\
&\quad \left. \left. \left\{ \left((U^\dagger)^{A_1} \otimes (U^\dagger)^{A_2} \right) \circ \left([((\tilde{T}')^\dagger)^{B_1 \rightarrow A_1} \otimes ((\tilde{T}')^\dagger)^{B_2 \rightarrow A_2}] (F^{B_1 B_2}) \right) \right\} \otimes I^{R_1 R_2} \right] \right)^i \\
&= \text{Tr} \left[\bigotimes_{j=1}^i \left((I^{A_1(j) A_2(j)} \otimes F^{R_1(j) R_2(j)}) (\gamma^{A_1(j) R_1(j)} \otimes \gamma^{A_2(j) R_2(j)}) \right. \right. \\
&\quad \left. \left. \left\{ \left((U^\dagger)^{A_1(j)} \otimes (U^\dagger)^{A_2(j)} \right) \circ \left([((\tilde{T}')^\dagger)^{B_1(j) \rightarrow A_1(j)} \otimes ((\tilde{T}')^\dagger)^{B_2(j) \rightarrow A_2(j)}] (F^{B_1(j) B_2(j)}) \right) \right\} \right. \right. \\
&\quad \left. \left. \otimes I^{R_1(j) R_2(j)} \right\} \right) \Big] \\
&= \text{Tr} \left[\left\{ \bigotimes_{j=1}^i \left((I^{A_1(j) A_2(j)} \otimes F^{R_1(j) R_2(j)}) (\gamma^{A_1(j) R_1(j)} \otimes \gamma^{A_2(j) R_2(j)}) \right) \right\} \right. \\
&\quad \left. \left\{ \bigotimes_{j=1}^i \left(\left((U^\dagger)^{A_1(j)} \otimes (U^\dagger)^{A_2(j)} \right) \circ \left([((\tilde{T}')^\dagger)^{B_1(j) \rightarrow A_1(j)} \otimes ((\tilde{T}')^\dagger)^{B_2(j) \rightarrow A_2(j)}] (F^{B_1(j) B_2(j)}) \right) \right) \right. \right. \\
&\quad \left. \left. \otimes \bigotimes_{j=1}^i I^{R_1(j) R_2(j)} \right\} \right] \\
&= \text{Tr} \left[\text{Tr}_{\bigotimes_{j=1}^i (R_1(j) R_2(j))} \left\{ \bigotimes_{j=1}^i \left((I^{A_1(j) A_2(j)} \otimes F^{R_1(j) R_2(j)}) (\gamma^{A_1(j) R_1(j)} \otimes \gamma^{A_2(j) R_2(j)}) \right) \right\} \right. \\
&\quad \left. \bigotimes_{j=1}^i \left\{ \left((U^\dagger)^{A_1(j)} \otimes (U^\dagger)^{A_2(j)} \right) \circ \left([((\tilde{T}')^\dagger)^{B_1(j) \rightarrow A_1(j)} \otimes ((\tilde{T}')^\dagger)^{B_2(j) \rightarrow A_2(j)}] (F^{B_1(j) B_2(j)}) \right) \right\} \right] \\
&= \text{Tr} \left[\left\{ \text{Tr}_{R_1 R_2} \left((I^{A_1 A_2} \otimes F^{R_1 R_2}) (\gamma^{A_1 R_1} \otimes \gamma^{A_2 R_2}) \right) \right\}^{\otimes i} \right. \\
&\quad \left. \left\{ \left((U^\dagger)^{A_1} \otimes (U^\dagger)^{A_2} \right) \circ \left([((\tilde{T}')^\dagger)^{B_1 \rightarrow A_1} \otimes ((\tilde{T}')^\dagger)^{B_2 \rightarrow A_2}] (F^{B_1 B_2}) \right) \right\}^{\otimes i} \right] \\
&= \text{Tr} \left[\left\{ (\gamma')^{A_1 A_2} \right\}^{\otimes i} \right. \\
&\quad \left. \left\{ (U^\dagger)^{A_1} \otimes (U^\dagger)^{A_2} \right\}^{\otimes i} \circ \left[[((\tilde{T}')^\dagger)^{B_1 \rightarrow A_1} \otimes ((\tilde{T}')^\dagger)^{B_2 \rightarrow A_2}] (F^{B_1 B_2}) \right] \right\}^{\otimes i} \Big]
\end{aligned}$$

This completes the proof of the proposition. \square

Lemma 6. Let i be a positive integer. Consider a $(|A|, s, \lambda, 4i)$ -qTPE for some positive integer s and $\lambda \geq 0$. Then,

$$\left| \mathbb{E}_{U \sim \text{TPE}} [(g(U))^{2i}] - \mathbb{E}_{U \sim \text{Haar}} [(g(U))^{2i}] \right| \leq (2|A|^3|B|^2)^i \cdot \lambda \cdot 2^{i(1+\delta)\mathbb{H}_{\max}^c(B)_\omega - iH_2^c(A|R)_\rho}.$$

Proof. We evaluate $[g(U)]^{2i}$ using Proposition 2 as:

$$(g(U)^2)^i$$

$$= \text{Tr} \left[\{(\gamma')^{A_1 A_2}\}^{\otimes i} \cdot \left\{ (U^\dagger)^{A_1} \otimes (U^\dagger)^{A_2} \right\}^{\otimes i} \circ \left[((\tilde{\mathcal{T}}')^\dagger)^{B_1 \rightarrow A_1} \otimes ((\tilde{\mathcal{T}}')^\dagger)^{B_2 \rightarrow A_2} \right] (F^{B_1 B_2}) \right\}^{\otimes i} \right].$$

Hence by taking expectation with respect to Haar and TPE and using the linearity of expectation we get,

$$\begin{aligned} & \left| \mathbb{E}_{U \sim \text{TPE}} [(g(U))^{2i}] - \mathbb{E}_{U \sim \text{Haar}} [(g(U))^{2i}] \right| \\ &= \left| \mathbb{E}_{U \sim \text{TPE}} \left[\text{Tr} \left[\{(\gamma')^{A_1 A_2}\}^{\otimes i} \cdot \left\{ (U^\dagger)^{A_1} \otimes (U^\dagger)^{A_2} \right\}^{\otimes i} \circ \left\{ \left[((\tilde{\mathcal{T}}')^\dagger)^{B_1 \rightarrow A_1} \otimes ((\tilde{\mathcal{T}}')^\dagger)^{B_2 \rightarrow A_2} \right] (F^{B_1 B_2}) \right\}^{\otimes i} \right] \right] \right. \right. \\ & \quad \left. \left. - \mathbb{E}_{U \sim \text{Haar}} \left[\text{Tr} \left[\{(\gamma')^{A_1 A_2}\}^{\otimes i} \cdot \left\{ (U^\dagger)^{A_1} \otimes (U^\dagger)^{A_2} \right\}^{\otimes i} \circ \left\{ \left[((\tilde{\mathcal{T}}')^\dagger)^{B_1 \rightarrow A_1} \otimes ((\tilde{\mathcal{T}}')^\dagger)^{B_2 \rightarrow A_2} \right] (F^{B_1 B_2}) \right\}^{\otimes i} \right] \right] \right] \right| \\ &= \left| \text{Tr} \left[\{(\gamma')^{A_1 A_2}\}^{\otimes i} \cdot \left\{ \mathbb{E}_{U \sim \text{TPE}} \left[\left\{ (U^\dagger)^{A_1} \otimes (U^\dagger)^{A_2} \right\}^{\otimes i} \circ \left(\left[((\tilde{\mathcal{T}}')^\dagger)^{B_1 \rightarrow A_1} \otimes ((\tilde{\mathcal{T}}')^\dagger)^{B_2 \rightarrow A_2} \right] (F^{B_1 B_2}) \right)^{\otimes i} \right] \right. \right. \right. \right. \\ & \quad \left. \left. \left. - \mathbb{E}_{U \sim \text{Haar}} \left[\left\{ (U^\dagger)^{A_1} \otimes (U^\dagger)^{A_2} \right\}^{\otimes i} \circ \left(\left[((\tilde{\mathcal{T}}')^\dagger)^{B_1 \rightarrow A_1} \otimes ((\tilde{\mathcal{T}}')^\dagger)^{B_2 \rightarrow A_2} \right] (F^{B_1 B_2}) \right)^{\otimes i} \right] \right] \right\} \right] \right| \\ &\leq \left\| ((\gamma')^{A_1 A_2})^{\otimes i} \right\|_1 \cdot \left\| \mathbb{E}_{U \sim \text{TPE}} \left[\left\{ (U^\dagger)^{A_1} \otimes (U^\dagger)^{A_2} \right\}^{\otimes i} \circ \left\{ \left[((\tilde{\mathcal{T}}')^\dagger)^{B_1 \rightarrow A_1} \otimes ((\tilde{\mathcal{T}}')^\dagger)^{B_2 \rightarrow A_2} \right] (F^{B_1 B_2}) \right\}^{\otimes i} \right] \right. \right. \\ & \quad \left. \left. - \mathbb{E}_{U \sim \text{Haar}} \left[\left\{ (U^\dagger)^{A_1} \otimes (U^\dagger)^{A_2} \right\}^{\otimes i} \circ \left\{ \left[((\tilde{\mathcal{T}}')^\dagger)^{B_1 \rightarrow A_1} \otimes ((\tilde{\mathcal{T}}')^\dagger)^{B_2 \rightarrow A_2} \right] (F^{B_1 B_2}) \right\}^{\otimes i} \right] \right] \right\|_\infty \\ &\leq \left\| (\gamma')^{A_1 A_2} \right\|_1^i \cdot \left\| \mathbb{E}_{U \sim \text{TPE}} \left[\left\{ (U^\dagger)^{A_1} \otimes (U^\dagger)^{A_2} \right\}^{\otimes i} \circ \left\{ \left[((\tilde{\mathcal{T}}')^\dagger)^{B_1 \rightarrow A_1} \otimes ((\tilde{\mathcal{T}}')^\dagger)^{B_2 \rightarrow A_2} \right] (F^{B_1 B_2}) \right\}^{\otimes i} \right] \right. \right. \\ & \quad \left. \left. - \mathbb{E}_{U \sim \text{Haar}} \left[\left\{ (U^\dagger)^{A_1} \otimes (U^\dagger)^{A_2} \right\}^{\otimes i} \circ \left\{ \left[((\tilde{\mathcal{T}}')^\dagger)^{B_1 \rightarrow A_1} \otimes ((\tilde{\mathcal{T}}')^\dagger)^{B_2 \rightarrow A_2} \right] (F^{B_1 B_2}) \right\}^{\otimes i} \right] \right] \right\|_2 \\ &\stackrel{a}{\leq} (|A| \|\gamma^{AR}\|_2^2)^i \cdot \lambda \cdot \left\| \left\{ \left[((\tilde{\mathcal{T}}')^\dagger)^{B_1 \rightarrow A_1} \otimes ((\tilde{\mathcal{T}}')^\dagger)^{B_2 \rightarrow A_2} \right] (F^{B_1 B_2}) \right\}^{\otimes i} \right\|_2 \\ &\stackrel{b}{\leq} (2|A| \|(\tilde{\rho}')^{AR}\|_2^2)^i \cdot \lambda \cdot \left\| \left[((\tilde{\mathcal{T}}')^\dagger)^{B_1 \rightarrow A_1} \otimes ((\tilde{\mathcal{T}}')^\dagger)^{B_2 \rightarrow A_2} \right] (F^{B_1 B_2}) \right\|_2^i \\ &\stackrel{c}{\leq} (2|A| \|(\tilde{\rho}')^{AR}\|_2^2)^i \cdot \lambda \cdot \|W_{\tilde{\mathcal{T}}'}\|_2^{4i} \stackrel{d}{\leq} (2|A|)^i 2^{-iH_2^\epsilon(A|R)_\rho} \cdot \lambda \cdot 2^{i(1+\delta)\mathbb{H}_{\max}^\epsilon(B)_\omega} (|A|^2 |B|^2)^i \\ &= (2|A|^3 |B|^2)^i \cdot \lambda \cdot 2^{i(1+\delta)\mathbb{H}_{\max}^\epsilon(B)_\omega - iH_2^\epsilon(A|R)_\rho}, \end{aligned}$$

where

- (a) follows from Fact 16 and Definition 12,
- (b) follows from Equation 21.

(c) follows from Lemma 5,

(d) follows from Note 1 and Equations 18 and 12.

This completes the proof of the lemma. \square

Now we upper bound the centralised $(2m)$ -th moment of $(g(U))^2$ under the approximate unitary design.

Lemma 7. *Let m be a positive integer. Suppose $\mu := \mathbb{E}_{U \sim \text{Haar}}[g(U)]$ and $H_2^\epsilon(A|R)_\rho < 0$. Consider a $(|A|, s, \lambda, 4m)$ -qTPE for some positive integer s and*

$$\lambda^{1/m} \leq \begin{cases} 2^4 m \cdot |A|^{-7} |B|^{-4} \mu^2 \cdot 2^{-(1+\delta)\mathbb{H}_{\max}^\epsilon(B)_\omega + H_2^\epsilon(A|R)_\rho}, & m < \frac{9}{64} |A| \mu^2 \cdot 2^{-(1+\delta)\mathbb{H}_{\max}^\epsilon(B)_\omega + H_2^\epsilon(A|R)_\rho - 4} \\ 2^{10} m^2 \cdot |A|^{-8} |B|^{-4} & \text{otherwise} \end{cases}.$$

Then,

$$\begin{aligned} & \mathbb{E}_{U \sim \text{TPE}} [((g(U))^2 - \mu^2)^{2m}] \\ & \leq \begin{cases} 7(9m\mu^2 2^{(1+\delta)\mathbb{H}_{\max}^\epsilon(B)_\omega - H_2^\epsilon(A|R)_\rho + 4} |A|^{-1})^m & m < \frac{9}{64} |A| \mu^2 \cdot 2^{-(1+\delta)\mathbb{H}_{\max}^\epsilon(B)_\omega + H_2^\epsilon(A|R)_\rho - 4} \\ 7(64m^2 2^{2(1+\delta)\mathbb{H}_{\max}^\epsilon(B)_\omega - 2H_2^\epsilon(A|R)_\rho + 8} |A|^{-2})^m & \text{otherwise} \end{cases}. \end{aligned}$$

Proof. From Lemma 6, we get

$$\begin{aligned} & \left| \mathbb{E}_{U \sim \text{TPE}} [((g(U))^2 - \mu^2)^{2m}] - \mathbb{E}_{U \sim \text{Haar}} [((g(U))^2 - \mu^2)^{2m}] \right| \\ & = \left| \sum_{i=0}^{2m} \binom{2m}{i} (-\mu^2)^{2m-i} \left(\mathbb{E}_{U \sim \text{TPE}} [(g(U))^{2i}] - \mathbb{E}_{U \sim \text{Haar}} [(g(U))^{2i}] \right) \right| \\ & \leq \sum_{i=0}^{2m} \binom{2m}{i} (\mu^2)^{2m-i} \left| \mathbb{E}_{U \sim \text{TPE}} [(g(U))^{2i}] - \mathbb{E}_{U \sim \text{Haar}} [(g(U))^{2i}] \right| \\ & \leq \lambda \sum_{i=0}^{2m} \binom{2m}{i} (\mu^2)^{2m-i} (2|A|^3 |B|^2)^i \cdot 2^{i(1+\delta)\mathbb{H}_{\max}^\epsilon(B)_\omega - iH_2^\epsilon(A|R)_\rho} \\ & = \lambda (\mu^2 + (2|A|^3 |B|^2) \cdot 2^{(1+\delta)\mathbb{H}_{\max}^\epsilon(B)_\omega - H_2^\epsilon(A|R)_\rho})^{2m} \\ & \stackrel{a}{\leq} \lambda ((3|A|^3 |B|^2) \cdot 2^{(1+\delta)\mathbb{H}_{\max}^\epsilon(B)_\omega - H_2^\epsilon(A|R)_\rho})^{2m}. \end{aligned}$$

where (a) follows from the observation that $\mu^2 \leq \mathbb{E}_U[g(U)^2]$ (from Jensen's inequality for the function x^2) and using the upper bound on $g(U)$ from Lemma 3 and the fact that $|A|, |B| \geq 1$.

Using Lemma 4 and the above inequality, we get

$$\begin{aligned} & \mathbb{E}_{U \sim \text{TPE}} [((g(U))^2 - \mu^2)^{2m}] \\ & \leq \lambda \left((3|A|^3 |B|^2)^2 \cdot 2^{2(1+\delta)\mathbb{H}_{\max}^\epsilon(B)_\omega - 2H_2^\epsilon(A|R)_\rho} \right)^m \\ & \quad + \begin{cases} 6(9m\mu^2 2^{(1+\delta)\mathbb{H}_{\max}^\epsilon(B)_\omega - H_2^\epsilon(A|R)_\rho + 4} |A|^{-1})^m & ; m < \frac{9}{64} |A| \mu^2 \cdot 2^{-(1+\delta)\mathbb{H}_{\max}^\epsilon(B)_\omega + H_2^\epsilon(A|R)_\rho - 4} \\ 6(64m^2 2^{2(1+\delta)\mathbb{H}_{\max}^\epsilon(B)_\omega - 2H_2^\epsilon(A|R)_\rho + 8} |A|^{-2})^m & ; \text{otherwise} \end{cases} \\ & \leq \begin{cases} 7(9m\mu^2 2^{(1+\delta)\mathbb{H}_{\max}^\epsilon(B)_\omega - H_2^\epsilon(A|R)_\rho + 4} |A|^{-1})^m & ; m < \frac{9}{64} |A| \mu^2 \cdot 2^{-(1+\delta)\mathbb{H}_{\max}^\epsilon(B)_\omega + H_2^\epsilon(A|R)_\rho - 4} \\ 7(64m^2 2^{2(1+\delta)\mathbb{H}_{\max}^\epsilon(B)_\omega - 2H_2^\epsilon(A|R)_\rho + 8} |A|^{-2})^m & ; \text{otherwise} \end{cases}. \end{aligned}$$

This completes the proof of the lemma. \square

Proof of Theorem 1. Define the positive integer $m := \lfloor |A| \kappa^2 2^{-(1+\delta) \mathbb{H}_{\max}^\epsilon(B)_\omega + H_2^\epsilon(A|R)_\rho - 8} \rfloor$. Using Lemma 7, we get

$$\begin{aligned}
\mathbb{P}_{U \sim \text{TPE}}[g(U) - \mu > \kappa] &\leq \mathbb{P}_{U \sim \text{TPE}}[(g(U))^2 > (\mu + \kappa)^2] \\
&\leq \begin{cases} \mathbb{P}_{U \sim \text{TPE}}[(g(U))^2 - \mu^2 > 2\mu\kappa] & ; \kappa \leq \mu \\ \mathbb{P}_{U \sim \text{TPE}}[(g(U))^2 - \mu^2 > \kappa^2] & ; \text{otherwise} \end{cases} \\
&\leq \begin{cases} \mathbb{P}_{U \sim \text{TPE}}[((g(U))^2 - \mu^2)^{2m} > (2\mu\kappa)^{2m}] & ; \kappa \leq \mu \\ \mathbb{P}_{U \sim \text{TPE}}[((g(U))^2 - \mu^2)^{2m} > \kappa^{4m}] & ; \text{otherwise} \end{cases} \\
&\leq \begin{cases} \frac{\mathbb{E}_{U \sim \text{TPE}}[(g(U))^2 - \mu^2]^{2m}}{(2\mu\kappa)^{2m}} & ; \kappa \leq \mu \\ \frac{\mathbb{E}_{U \sim \text{TPE}}[(g(U))^2 - \mu^2]^{2m}}{\kappa^{4m}} & ; \text{otherwise} \end{cases} \\
&\leq \begin{cases} \frac{7(9m\mu^2 2^{(1+\delta) \mathbb{H}_{\max}^\epsilon(B)_\omega - H_2^\epsilon(A|R)_\rho + 4} |A|^{-1})^m}{(2\mu\kappa)^{2m}} & ; \kappa \leq \mu, m < \frac{9}{64} |A| \mu^2 \cdot 2^{-(1+\delta) \mathbb{H}_{\max}^\epsilon(B)_\omega + H_2^\epsilon(A|R)_\rho - 4} \\ \frac{7(8m 2^{(1+\delta) \mathbb{H}_{\max}^\epsilon(B)_\omega - H_2^\epsilon(A|R)_\rho + 4} |A|^{-1})^{2m}}{\kappa^{4m}} & ; \text{otherwise} \end{cases} \\
&\leq \begin{cases} 7(3m\kappa^{-2} 2^{(1+\delta) \mathbb{H}_{\max}^\epsilon(B)_\omega - H_2^\epsilon(A|R)_\rho + 4} |A|^{-1})^m & ; \kappa \leq \mu \\ 7(8m\kappa^{-2} 2^{(1+\delta) \mathbb{H}_{\max}^\epsilon(B)_\omega - H_2^\epsilon(A|R)_\rho + 4} |A|^{-1})^{2m} & ; \text{otherwise} \end{cases} \\
&\leq 7 \cdot 2^{-a\kappa^2},
\end{aligned}$$

where $a := |A| \cdot 2^{-(1+\delta) \mathbb{H}_{\max}^\epsilon(B)_\omega + H_2^\epsilon(A|R)_\rho - 9}$.

Since $0 < \delta < 1$, $\mathbb{H}_{\max}^\epsilon(B)_\omega \leq \log |B|$ and $H_2^\epsilon(A|R)_\rho \geq -\log |A|$, therefore $(m \cdot |A|^{-8} |B|^{-6} \cdot \mu^2)^m \leq (2^4 \cdot m \cdot |A|^{-7} |B|^{-4} \mu^2 \cdot 2^{-(1+\delta) \mathbb{H}_{\max}^\epsilon(B)_\omega + H_2^\epsilon(A|R)_\rho - 4})^m$

And thus, the above probability analysis requires us to use a $(|A|, s, \lambda', 4m)$ -qTPE with

$$\lambda' \leq (m \cdot |A|^{-8} |B|^{-6} \cdot \mu^2)^m,$$

if $\kappa < \mu$ and

$$\lambda' \leq (2^{10} \cdot m^2 \cdot |A|^{-8} |B|^{-4})^m \quad (26)$$

otherwise. Define $t := |A| \kappa^2 2^{H_2^\epsilon(A|R)_\rho - \mathbb{H}_{\max}^\epsilon(B)_\omega - 6}$. Then $t \geq 4m$ and a $(|A|, s, \lambda, t)$ -qTPE suffices for the derandomization where $\lambda := (|A|^{-8} |B|^{-6} \cdot \mu^2)^t$.

Note that this value of λ is smaller than that obtained in Equation 26 and hence a unitary chosen at random from this qTPE shall also achieve exponential concentration. For a qTPE to achieve this smaller λ requires slightly more number of iterations of a qTPE with constant singular value gap, as compared to a qTPE with λ' in Equation 26. Combined with Lemma 1, we finally get

$$\mathbb{P}_{U \sim \text{TPE}} \left[f(U) > 2^{-\frac{1}{2} H_2^\epsilon(A|R)_\rho - \frac{1}{2} \mathbb{H}_{\max}^\epsilon(A'|B)_\omega + 1} + 14\sqrt{\epsilon} + 2\kappa \right] \leq 7 \cdot 2^{-a\kappa^2}.$$

Combining Proposition 1 with the above analysis finishes the proof of Theorem 1. \square

IV. THE ASYMPTOTIC IID CASE

In this section we first show that the smooth one-shot entropies defined in Section II-C approach their natural Shannon entropic analogues in the asymptotic iid limit. We then use those results to prove the asymptotic iid version of the main theorem. The asymptotic iid version will be stated in terms of Shannon entropies.

A. Asymptotic smoothing of $\mathbb{H}_{\max}^\epsilon$ and $\mathbb{H}_2^{\epsilon,\delta}$

In this section, we use the properties of typical sequences and subspaces to find an upper bound on $\mathbb{H}_{\max}^\epsilon$ and a lower bound on $\mathbb{H}_2^{\epsilon,\delta}$ in the asymptotic limit of many iid copies of the underlying quantum states. The bounds obtained will be the Shannon entropic quantities that one would expect. We first prove a few essential lemmas.

Lemma 8. Suppose we have a density matrix ω on the system AB . Let $|w_j\rangle^{AB}$, $j \in [|A||B|]$ be the eigenvectors of ω^{AB} with eigenvalues q_j . For $j \in [|A||B|]$, define $\theta_j^B := \text{Tr}_A[|w_j\rangle^{AB}\langle w_j|]$. Let $p_j := \{p_j(i)\}_{i \in [|B|]}$, $j \in [|A||B|]$ be the probability distribution on $|B|$ obtained by measuring θ_j in the eigenbasis of ω^B . Let $0 < \epsilon, \delta < 1/3$. Define $q_{\min} := 2^{-\mathbb{H}_{\max}^{\epsilon/2}(AB)\omega}$, $p_{\min} := \min_{j \in [|A||B|]} 2^{-\mathbb{H}_{\max}^{\epsilon/2}(|B|)p_j}$. Let $n \geq 2^5 q_{\min}^{-1} p_{\min}^{-1} \delta^{-2} \log(|A||B|/\epsilon)$. Consider the n -fold tensor power $\omega^{A^n B^n} := (\omega^{AB})^{\otimes n}$. Let τ be a strongly δ -typical type of an eigenvector sequence of $\omega^{A^n B^n}$. Let $(|v_1\rangle \otimes \dots \otimes |v_n\rangle)^{A^n B^n}$ be an eigenvector sequence of type τ . Let $\sigma^{B^n} := \Pi_{\omega, 3\delta}^{B^n} \omega^{B^n} \Pi_{\omega, 3\delta}^{B^n}$. Then,

$$\text{Tr}[(I^{A^n} \otimes \Pi_{\omega, 3\delta}^{B^n})(|v_1\rangle \dots |v_n\rangle \langle v_1| \dots \langle v_n|)^{A^n B^n}] \geq 1 - \epsilon.$$

Proof. Since τ is a strongly δ -typical type, the number of occurrences n_j of each $|w_j\rangle^{AB}$ in the sequence $|v_1\rangle^{AB}, \dots, |v_n\rangle^{AB}$ is $nq_j(1 \pm \delta)$. After a suitable rearranging, we can write

$$(|v_1\rangle \otimes \dots \otimes |v_n\rangle)^{A^n B^n} = (|w_1\rangle^{\otimes n_1} \otimes \dots \otimes |w_{|A||B|}\rangle^{\otimes n_{|A||B|}})^{A^n B^n}.$$

To prove the lemma, it suffices to show that

$$\text{Tr}[\Pi_{\omega, 3\delta}^{B^n}((\theta_1^B)^{\otimes n_1} \otimes \dots \otimes (\theta_{|A||B|}^B)^{\otimes n_{|A||B|}})] \geq 1 - \epsilon. \quad (27)$$

Let $|x_1\rangle^B, \dots, |x_{|B|}\rangle^B$ be the eigenbasis of ω^B with eigenvalues $r_1, \dots, r_{|B|}$. Observe that we have the operator equality $\sum_{j=1}^{|A||B|} q_j \theta_j^B = \omega^B$. Now consider the matrices θ_j^B in the basis $|x_1\rangle^B, \dots, |x_{|B|}\rangle^B$. Thus for any $i \in [|B|]$, $\sum_{j=1}^{|A||B|} q_j p_j(i) = r_i$.

Let $\Pi_j^{B^{n_j}}$ be the projector onto the eigenvectors of $(\omega^B)^{\otimes n_j}$ that are strongly δ -typical according to p_j (and not with respect to the eigenvalues $\{r_i\}_{i \in [|B|]}$ of ω^B). Note that this is not the same as projecting onto the eigenvectors of $(\theta_j^B)^{\otimes n_j}$ and also $(\theta_j^B)^{\otimes n_j}$ does not commute with $\omega_B^{\otimes n_j}$.

Since $\{p_j(i)\}_{i \in [|B|]}$ is the probability distribution on B obtained by measuring θ_j^B in the eigenbasis of ω^B for all $j \in [|A||B|]$ and $\Pi_j^{B^{n_j}}$ is the typical projector with respect to the probability distribution $\{p_j\}$ (not on the eigenbasis of $\omega^{B^{n_j}}$), therefore by Fact 19 we have, $\text{Tr}[\Pi_j^{B^{n_j}}(\theta_j^B)^{\otimes n_j}] \geq 1 - \frac{\epsilon}{|A||B|}$. Thus,

$$\begin{aligned} \text{Tr}[(\Pi_1^{B^{n_1}} \otimes \dots \otimes \Pi_{|A||B|}^{B^{n_{|A||B|}}})(\theta_1^B)^{\otimes n_1} \otimes \dots \otimes (\theta_{|A||B|}^B)^{\otimes n_{|A||B|}}] &\geq \prod_{j=1}^{|A||B|} \left(1 - \frac{\epsilon}{|A||B|}\right) \\ &\geq 1 - \epsilon. \end{aligned} \quad (28)$$

Fix an eigenvector $(|v_1\rangle \otimes \dots \otimes |v_n\rangle)^{A^n B^n}$ in the support of $\Pi_1^{B^{n_1}} \otimes \dots \otimes \Pi_{|A||B|}^{B^{n_{|A||B|}}}$; the eigenvector can be viewed as a sequence of length n_1 (typical with respect to probability p_1), \dots , $n_{|A||B|}$ (typical with respect to probability distribution $p_{|A||B|}$) such that $\sum_{j=1}^{|A||B|} n_j = n$. Then the number of occurrences of $|x_i\rangle$ in the sequence is

$$\sum_{j=1}^{|A||B|} nq_j(1 \pm \delta)p_j(i)(1 \pm \delta) = r_i(1 \pm 3\delta).$$

This shows that the eigenvector is strongly (3δ) -typical for the state ω^{B^n} . Also, since eigenvectors corresponding to distinct eigenvalues are orthogonal, therefore $\bigotimes_{j=1}^{|A||B|} \Pi_j^{B^{n_j}}$ commutes with $\Pi_{\omega, 3\delta}^{B^n}$. In other words,

$$\Pi_1^{B^{n_1}} \otimes \cdots \otimes \Pi_{|A||B|}^{B^{n_{|A||B|}}} \leq \Pi_{\omega, 3\delta}^{B^n} \quad (29)$$

Combining Equation 28 with Equation 29 and then Equation 27 completes the proof of the lemma. \square

Note: The non trivial take away message of the above lemma is that a typical vector of ω^{AB} still retains the typicality property when projected onto the marginal ω^B (in the sense that any typical eigenvector of $(\omega^{AB})^{\otimes n}$ has a large projection onto the subspace spanned by $I^{A^n} \otimes \Pi_{\omega, 3\delta}^{B^n}$; even though the eigenvectors of ω^{AB} and ω^B are not related), but the probability distribution is not the one obtained by eigenvalues of ω^B . Although the above statement might look believable in analogy with the classical typical sequences, it is still non-trivial to prove as is suggested by the proof above. The proof centrally uses the property of strongly typical types.

Lemma 9. *Consider the setting of Lemma 8. Let $V_\tau \leq A^n B^n$ denote the type subspace corresponding to type τ . Then there is a subspace $\hat{V}_\tau \leq V_\tau$, $|\hat{V}_\tau| \geq (1 - \sqrt{\epsilon})|V_\tau|$ such that for every vector $|v\rangle \in \hat{V}_\tau$, $\|(I^{A^n} \otimes \Pi_\sigma^{B^n})|v\rangle\|_2^2 \geq 1 - \sqrt{\epsilon}$.*

Proof. We invoke Fact 14 with $A := V_\tau$ and $B := I^{A^n} \otimes \Pi_\sigma^{B^n}$ in order to prove this lemma. Take the basis for A provided by Fact 14. Call it $\{|a\rangle\}_a$. Observe that the vectors $\Pi_B|a\rangle$ are pairwise orthogonal (some of them may be the zero vector). From Lemma 8, we know that $\text{Tr}[\Pi_B \frac{\Pi_A}{|A|}] \geq 1 - \epsilon$. By Markov's inequality, there is a subset S of the basis vectors of A , $|S| \geq (1 - \sqrt{\epsilon})|A|$ such that for all $a \in S$, $\|\Pi_B|a\rangle\|_2^2 = \text{Tr}[\Pi_B|a\rangle\langle a|] \geq 1 - \sqrt{\epsilon}$. Define the subspace $\hat{A} := \text{span}_{a \in S}|a\rangle$. From the above observation, for any vector $|v\rangle \in \hat{A}$, $\|\Pi_B|v\rangle\|_2^2 \geq 1 - \sqrt{\epsilon}$. This subspace \hat{A} serves as the subspace \hat{V}_τ required by the lemma. \square

Lemma 10. *Let $0 < \epsilon < 1$. Let $|v_1\rangle, \dots, |v_t\rangle$ be orthonormal vectors lying in a Hilbert space \mathcal{H} . Suppose there is a subspace $B \leq \mathcal{H}$ with the property that $\|\Pi_B|v_i\rangle\|_2^2 \geq 1 - \epsilon$ for all $i \in [t]$. Let $|v\rangle$ be a unit vector lying in the span of the vectors $|v_i\rangle$. Then, $\|\Pi_B|v\rangle\|_2^2 \geq 1 - 8t\sqrt{\epsilon}$.*

Proof. Let $|v\rangle = \sum_{i=1}^t \alpha_i |v_i\rangle$ where $\sum_{i=1}^t |\alpha_i|^2 = 1$. Define the column t -tuple $\alpha := (\alpha_1, \dots, \alpha_t)^T$, and the $t \times t$ -matrix M with $M_{ij} := \langle v_i | \Pi_B | v_j \rangle$. Note that M is Hermitian. Then,

$$\|\Pi_B|v\rangle\|_2^2 = \sum_{i,j=1}^t \alpha_i^* \alpha_j \langle v_i | \Pi_B | v_j \rangle = \alpha^\dagger M \alpha.$$

We have $M_{ii} \geq 1 - \epsilon$. For $i \neq j$, we use triangle inequality and Fact 13 to obtain

$$\|\Pi_B|v_i\rangle - \Pi_B|v_j\rangle\|_2 \geq \| |v_i\rangle - |v_j\rangle \|_2 - \|\Pi_B|v_i\rangle - |v_i\rangle\|_2 - \|\Pi_B|v_j\rangle - |v_j\rangle\|_2 \geq \sqrt{2} - 4\sqrt{\epsilon},$$

which implies that $2 - 8\sqrt{2\epsilon} \leq \|\Pi_B|v_i\rangle - \Pi_B|v_j\rangle\|_2^2 \leq 2 - 2\Re(M_{ij})$, which further implies that $\Re(M_{ij}) \leq 4\sqrt{2\epsilon}$. Arguing similarly with $\|\Pi_B|v_i\rangle - \sqrt{-1} \cdot \Pi_B|v_j\rangle\|_2$, we conclude that $\Im(M_{ij}) \leq 4\sqrt{2\epsilon}$. Thus, $|M_{ij}| \leq 8\sqrt{\epsilon}$. By Gershgorin's theorem, the smallest eigenvalue of M is larger than $1 - \epsilon - 8(t-1)\sqrt{\epsilon} \geq 1 - 8t\sqrt{\epsilon}$. So $\|\Pi_B|v\rangle\|_2^2 = \alpha^\dagger M \alpha \geq 1 - 8t\sqrt{\epsilon}$, completing the proof of the lemma. \square

Proposition 3. *Suppose we have a density matrix ω on the system B . Let $0 < \epsilon, \delta < 1/3$ and define $q_{\min} := 2^{-\mathbb{H}_{\max}^{\epsilon/2}(B)_\omega}$. Let $n \geq 4q_{\min}^{-1} \delta^{-2} \log(|B|/\epsilon)$. Consider the n -fold tensor power $\omega^{B^n} := (\omega^B)^{\otimes n}$. Then, $n(1 - \delta)H(B)_\omega \leq \mathbb{H}_{\max}^\epsilon(B^n)_\omega \leq n(1 + \delta)H(B)_\omega$.*

Proof. Consider the eigenvalues of ω^{B^n} that are not strongly δ -typical; call them atypical. By Fact 19, the atypical eigenvalues sum to less than or equal to ϵ and the smallest typical eigenvalue is at least $2^{-n(1+\delta)H(B)_\omega}$. Hence the eigenvalues less than $2^{-n(1+\delta)H(B)_\omega}$ add up to less than or equal to ϵ . Moreover, the eigenvalues less than $2^{-n(1-\delta)H(B)_\omega}$ add up to more than $1 - \epsilon > \epsilon$. This completes the proof the proposition. \square

Proposition 4. Suppose we have a density matrix ω on the system AB . Let $|w_j\rangle^{AB}$, $j \in [|A||B|]$ be the eigenvectors of ω^{AB} with eigenvalues q_j . For $j \in [|A||B|]$, define $\theta_j^B := \text{Tr}_A[|w_j\rangle^{AB}\langle w_j|]$. Let $p_j := \{p_j(i)\}_{i \in [|B|]}$, $j \in [|A||B|]$ be the probability distribution on $[|B|]$ obtained by measuring θ_j in the eigenbasis of ω^B . Let $0 < \epsilon, \delta < 1/3$. Define $q_{\min} := 2^{-\mathbb{H}_{\max}^{\epsilon/2}(AB)_\omega}$, $p_{\min} := \min_{j \in [|A||B|]} 2^{-\mathbb{H}_{\max}^{\epsilon/2}(|B|)_{p_j}}$. Let $n := 2^5 q_{\min}^{-1} p_{\min}^{-1} \delta^{-2} \log(|A||B|/\epsilon)$. Consider the n -fold tensor power $\omega^{A^n B^n} := (\omega^{AB})^{\otimes n}$. Let $\epsilon' := 8(n + |A||B|)^{|A||B|} \epsilon^{1/4}$. Then,

$$\begin{aligned} nH(A|B)_\omega + 32n\sqrt{\epsilon'} \log |A| + \log(\epsilon')^{-1} &\geq H_2^{4\sqrt{\epsilon'}}(A^n|B^n)_\omega \geq (H_2^{\epsilon', 5\delta}(A^n|B^n)_\omega \\ &\geq nH(A|B)_\omega - n\delta(3H(AB)_\omega + 7H(B)_\omega). \end{aligned}$$

Proof. For a type τ of $\omega^{A^n B^n}$, define $p_\tau := \text{Tr}[\Pi_{V_\tau} \omega^{A^n B^n}]$. By Fact 19,

$$\omega^{A^n B^n} = \bigoplus_{\tau} \Pi_{V_\tau} \omega^{A^n B^n} \Pi_{V_\tau} = \bigoplus_{\tau} p_\tau \frac{\Pi_{V_\tau}}{|V_\tau|},$$

where the direct sum is over all types τ . Now define $\eta^{A^n B^n} := \bigoplus_{\tau: \text{typical}} p_\tau \frac{\Pi_{V_\tau}}{|V_\tau|}$, where the sum is only over strongly δ -typical types τ . By Fact 19 and Lemma 9,

$$\eta^{A^n B^n} \leq \omega^{A^n B^n}, \quad \|\eta^{A^n B^n} - \omega^{A^n B^n}\|_1 \leq 2\sqrt{\epsilon}.$$

Let $\sigma^{B^n} := \Pi_{\omega, 3\delta}^{B^n} \omega^{B^n} \Pi_{\omega, 3\delta}^{B^n}$. By Fact 19, we have

$$\sigma^{B^n} \leq \omega^{B^n}, \quad \|\omega^{B^n} - \sigma^{B^n}\|_1 \leq \epsilon, \quad \|(\sigma^{B^n})^{-1}\|_\infty \leq 2^{nH(B)_\omega(1+3\delta)}.$$

From Lemma 9, we already know that for any strongly δ -typical type τ , for any vector $|w_\tau\rangle \in \hat{V}_\tau$, $\|(I^{A^n} \otimes \Pi_{\omega, 3\delta}^{B^n})|w_\tau\rangle\|_2^2 \geq 1 - \sqrt{\epsilon}$. We now have to show a similar result for an arbitrary linear combination of vectors $|w_\tau\rangle$ over all strongly δ -typical types τ . For this we invoke Lemma 10 and Fact 19. We thus conclude that for any vector $|v\rangle \in \text{supp}(\eta^{A^n B^n})$,

$$\|(I^{A^n} \otimes \Pi_{\omega, 3\delta}^{B^n})|v\rangle\|_2 \geq 1 - 8 \binom{n + |A||B| - 1}{|A||B| - 1} \epsilon^{1/4} \geq 1 - \epsilon'.$$

By Fact 19, the smallest non-zero eigenvalue of $(\omega_{\epsilon, \delta}''')^{B^n}$ (which is defined by setting $\epsilon = 0$ in $\omega_{\epsilon, \delta}'''$ from Definition 4) is smaller than $2^{-nH(B)_\omega(1-\delta)}$. Again invoking Fact 19, we conclude that $\text{supp}((\omega_{\epsilon, 5\delta}''')^{B^n}) \geq \text{supp}(\sigma^{B^n})$. Thus, for any vector $|v\rangle \in \text{supp}(\eta^{A^n B^n})$, $\|(I^{A^n} \otimes \Pi_{\omega_{\epsilon, 5\delta}'''}^{B^n})|v\rangle\|_2 \geq 1 - \epsilon'$. Moreover, by Proposition 3

$$\begin{aligned} \log\|(\omega_{\epsilon, 5\delta}''')^{B^n}\|_\infty^{-1} &= (1 + 5\delta) \log\|(\omega_{\epsilon}''')^{B^n}\|_\infty^{-1} = (1 + 5\delta) \mathbb{H}_{\max}^\epsilon(B^n)_\omega \\ &\leq n(1 + 7\delta)H(B)_\omega. \end{aligned}$$

Again using Fact 19, we get

$$\mathbb{H}_2^{\epsilon', 5\delta}(A^n|B^n)_\omega$$

$$\begin{aligned}
&\geq -2 \log \left\| \left(I^{A^n} \otimes (\omega'''_{\epsilon, 5\delta})^{B^n} \right)^{-1/4} \eta^{A^n B^n} \left(I^{A^n} \otimes (\omega'''_{\epsilon, 5\delta})^{B^n} \right)^{-1/4} \right\|_2 \\
&\geq -2 \log \left(\left\| \left(I^{A^n} \otimes (\omega'''_{\epsilon, 5\delta})^{B^n} \right)^{-1/4} \right\|_\infty^2 \cdot \left\| \eta^{A^n B^n} \right\|_2 \right) \\
&= -2 \log \left(\left\| (\omega'''_{\epsilon, 5\delta})^{B^n} \right\|_\infty^{-1/2} \cdot \left(\sum_{\tau: \text{typical}} \frac{p_\tau^2 |\hat{V}_\tau|}{|V_\tau|^2} \right)^{1/2} \right) \\
&\geq -\log \left\| (\omega'''_{\epsilon, 5\delta})^{B^n} \right\|_\infty^{-1} - 2 \log \left(\sum_{\tau: \text{typical}} \frac{p_\tau^2}{|V_\tau|} \right)^{1/2} \\
&= -\log \left\| (\omega'''_{\epsilon, 5\delta})^{B^n} \right\|_\infty^{-1} - 2 \log \left\| \Pi_{\omega, \delta}^{A^n B^n} \omega^{A^n B^n} \Pi_{\omega, \delta}^{A^n B^n} \right\|_2 \\
&\geq -nH(B)_\omega (1 + 7\delta) - 2 \log(2^{-nH(AB)_\omega (1-\delta)} \cdot \left\| \Pi_{\omega, \delta}^{A^n B^n} \right\|_2) \\
&= -nH(B)_\omega (1 + 7\delta) - 2 \log(2^{-nH(AB)_\omega (1-\delta)} \cdot (\text{Tr } \Pi_{\omega, \delta}^{A^n B^n})^{1/2}) \\
&\geq -nH(B)_\omega (1 + 7\delta) - 2 \log 2^{-\frac{n}{2} H(AB)_\omega (1-3\delta)} \\
&\geq nH(A|B)_\omega - n\delta(3H(AB)_\omega + 7H(B)).
\end{aligned}$$

Combining with Fact 17 completes the proof of the proposition. \square

Remark 5. The importance of using smooth one shot entropic quantities has been remarked in many earlier works. In particular, it is only the smooth quantities that approach their natural Shannon entropic counterparts in asymptotic iid limit, not their unsmoothed version. $H_2^\epsilon(\cdot|\cdot)$ is also shown to be bounded by its natural Shannon counterpart in the asymptotic iid limit. However we would like to emphasise here the importance of the additional smoothing parameter δ , which is crucial in this work.

We note that the smooth decoupling theorem was first proved by Szehr et al. [4] in terms of the conditional H_{\min}^ϵ quantity. Actually their proof works without any change for the conditional H_2^ϵ quantity also, which is a slightly stronger statement. However an important step in their proof was to replace the original superoperator in the statement of the decoupling theorem by another superoperator whose Choi matrix is a smoothed version of the Choi matrix of the original superoperator. They showed that on average over the Haar random unitary U applied to the input state, the new superoperator gives an output state close to the output state of the original superoperator. However this does not mean that for every unitary U , the output of the new superoperator is close to the output of the original superoperator. However, in order to prove a concentration result for decoupling, we need to bound a certain Lipschitz constant. By its very definition the Lipschitz constant should work for all unitaries not just for average Haar random ones. This stringent requirement leads us to the second smoothing parameter δ . Existence of this δ implies that for every unitary U , the output state of the new superoperator is δ -close in trace distance to the output state of the original superoperator. The parameter δ is the one shot analogue of the (non-trivial) observation that there is a large subspace of the jointly typical subspace of $(\rho^{AB})^{\otimes n}$ such that every vector in this subspace has projection length of at most $1 - \delta$ on the tensor product of the marginal typical subspaces on $(\rho^A)^{\otimes n}$ and $(\rho^B)^{\otimes n}$.

Remark 6. Consider fixed systems A, B and a fixed state ω^{AB} . For a fixed δ , divide the smooth modified conditional Rényi 2-entropy, the smooth conditional Rényi 2-entropy and the smooth modified max-entropy by n and let $\epsilon \rightarrow 0$. This implies that $n \rightarrow \infty$ and $\epsilon' \rightarrow 0$. Finally, let $\delta \rightarrow 0$. This shows that in the asymptotic iid limit, the smooth conditional Rényi 2-entropy divided by n is lower bounded by the smooth modified conditional Rényi 2-entropy divided by n which is further lower bounded by the conditional

Shannon entropy, and the smooth modified max-entropy divided by n is upper bounded by the Shannon entropy.

B. Proof of the iid extension of Theorem 1

In this section we take our main one-shot concentration result and apply it in the asymptotic iid setting. That is, we take the n -fold tensor product copy of the channel \mathcal{T} and the state ρ^{AR} , apply Theorem 1 to it, and obtain bounds in terms of the standard Shannon entropies.

Corollary 1. *Consider the setting of Theorem 1 above. Consider the density matrix $\omega^{A'B}$. Let $|w_j\rangle^{A'B}$, $j \in [|A||B|]$ be the eigenvectors of $\omega^{A'B}$ with eigenvalues q_j . For $j \in [|A||B|]$, define $\theta_j^B := \text{Tr}_{A'}[|w_j\rangle\langle w_j|^{AB}]$. Let p_j , $j \in [|A||B|]$ be the probability distribution on $[|B|]$ obtained by measuring θ_j in the eigenbasis of ω^B . Define $q_{\min} := 2^{-\mathbb{H}_{\max}^{\epsilon/2}(A'B)_\omega}$, $p_{\min} := \min_{j \in [|A||B|]} 2^{-\mathbb{H}_{\max}^{\epsilon/2}(|B|)_{p_j}}$, $n := 2^5 q_{\min}^{-1} p_{\min}^{-1} \delta^{-2} \log \frac{|A||B|}{\epsilon}$. Consider the n -fold tensor powers $\omega^{(A')^n B^n} := (\omega^{A'B})^{\otimes n}$, $\rho^{A^n R^n} := (\rho^{AR})^{\otimes n}$. Let $\epsilon' := 8(n + |A||B|)^{|A||B|} \epsilon^{1/4}$. Let $\kappa > 0$. Then,*

$$\mathbb{P}_U \left[f(U) > 2^{-\frac{n}{2}(H(A|R)_\rho - \delta(3H(AR)_\rho + 7H(R)_\rho)) - \frac{n}{2}(H(A'|B)_\omega - \delta(3H(A'B)_\omega + 7H(B)_\omega))} + 28(\epsilon')^{1/4} + 2\kappa \right] \leq 7 \cdot 2^{-a\kappa^2},$$

which can also be expressed as:

$$\mathbb{P}_U \left[f(U) > 2^{-\frac{n}{2}(H(A|R)_\rho - \delta(3H(AR)_\rho + 7H(R)_\rho)) - \frac{n}{2}(H(A'|B)_\omega - \delta(3H(A'B)_\omega + 7H(B)_\omega))} + 28(\epsilon')^{1/4} + \sqrt{t/2} \beta \right] \leq 7 \cdot 2^{-a\kappa^2},$$

where the unitary U^{A^n} is chosen uniformly at random from a $(|A|^n, s, \lambda, t)$ -qTPE, and the parameters a, t, α, β are defined as

$$\begin{aligned} a &:= |A|^n \cdot 2^{n(H(A|R)_\rho - \delta(3H(AR)_\rho + 7H(R)_\rho)) - nH(B)_\omega(1+7\delta) - 9}, \\ t &:= |A|^n \kappa^2 2^{n(H(A|R)_\rho + 32\sqrt{\epsilon'}) + \log(\epsilon')^{-1} - nH(B)_\omega(1-5\delta) - 6}, \\ \beta &:= \sqrt{1/a}, \end{aligned}$$

and μ, λ are defined in Theorem 1 above.

Proof. The proof follows by a direct application of Theorem 1 and Propositions 3 and 4. We just need to keep in mind that the ‘weighting state’ in the definition of $\mathbb{H}_2^{\epsilon', 5\delta}((A')^n | B^n)_{\omega \otimes n}$ is $(\omega_{\epsilon, 5\delta}'')^{B^n}$ as in the proof of Proposition 4, and the function $g(U)$ is defined with respect to the perturbed Choi-Jamiołkowski state $\eta^{(A')^n B^n}$ defined in Proposition 4 and a perturbed input state $(\rho')^{A^n R^n}$ which is $4\sqrt{\epsilon'}$ -close to the original input state $\rho^{A^n R^n}$. We get

$$\begin{aligned} \mu &\leq 2^{-\frac{1}{2}\mathbb{H}_2^{\epsilon', 5\delta}(A'^n | B^n)_{\omega \otimes n} - \frac{1}{2}H_2^{4\sqrt{\epsilon'}}(A^n | R^n)_{\rho \otimes n}} \\ &\leq 2^{-\frac{n}{2}(H(A|R)_\rho - \delta(3H(AR)_\rho + 7H(R)_\rho)) - \frac{n}{2}(H(A'|B)_\omega - \delta(3H(A'B)_\omega + 7H(B)_\omega))}, \\ a &= |A|^n 2^{H_2^{4\sqrt{\epsilon'}}(A^n | R^n)_{\rho \otimes n} - (1+\delta)\mathbb{H}_{\max}^\epsilon(B^n)_{\omega \otimes n} - 9} \end{aligned}$$

$$\begin{aligned}
&\geq |A|^n 2^{n(H(A|R)_\rho - \delta(3H(AR)_\rho + 7H(R)_\rho)) - nH(B)_\omega(1+7\delta) - 9}, \\
t &= |A|^n \kappa^2 2^{H_2^{\sqrt{\epsilon'}}(A^n|R^n)_{\rho^{\otimes n}} - \mathbb{H}_{\max}^\epsilon(B^n)_{\omega^{\otimes n}} - 6} \\
&\leq |A|^n \kappa^2 2^{n(H(A|R)_\rho + 32\sqrt{\epsilon'}) + \log(\epsilon')^{-1} - nH(B)_\omega(1-5\delta) - 6}.
\end{aligned}$$

Substituting the above expressions in Theorem 1 proves the desired corollary. \square

V. DECOUPLING UNDER PARTIAL TRACE (ALSO REFERRED TO AS THE FULLY QUANTUM SLEPIAN WOLF (FQSW) [1])

In this section, we prove the concentration result for the Fully Quantum Slepian Wolf (FQSW) problem with respect to unitary designs.

Theorem 2. *Consider the setting of Theorem 1. Consider the FQSW decoupling function*

$$f(U) = f_{\text{FQSW}}(U^{A_1 A_2}) := \|\text{Tr}_{A_2}[(U^{A_1 A_2} \otimes I^R) \circ \rho^{A_1 A_2 R}] - \pi^{A_1} \otimes \rho^R\|_1.$$

Suppose we are promised that

$$\begin{aligned}
&\|(\tilde{\rho}')^R\|_2^2 < 0.9|A_1||A_2| \|(\tilde{\rho}')^{AR}\|_2^2, |A_1| \geq 2, |A_2| > |A_1|, \\
&|A_2| 2^{H_2^\epsilon(A_1 A_2|R)_\rho - 8} - 4 > -H_2^\epsilon(A_1 A_2|R) + \log|A_1| + 2\log|A_2|.
\end{aligned}$$

In other words $\rho^{A_1 A_2 R}$ is not too close to a tensor product state on $A_1 A_2 \otimes R$ or $\rho^{A_1 A_2}$ is not too close to the completely mixed state $\pi^{A_1 A_2}$. The following concentration inequality holds:

$$\mathbb{P}_{U \sim \text{design}} \left[f(U) > \sqrt{\frac{|A_1|}{|A_2|}} \cdot 2^{-\frac{1}{2}H_2^\epsilon(A|R)_\rho + 1} + 14\sqrt{\epsilon} + 2\kappa \right] \leq 7 \cdot 2^{-a\kappa^2},$$

where the unitary U^A is chosen uniformly at random from an $(|A_1||A_2|, s, \lambda, t)$ -qTPE $a := |A_2| 2^{H_2^\epsilon(A|R)_\rho - 9}$ and $t := 8a\kappa^2$. The quantity λ , defined in Theorem 1 satisfies the inequality

$$\left(0.008|A_2|^{-9}|A_1|^{-13} 2^{-H_2^\epsilon(A_1 A_2|R)_\rho} \right)^t < \lambda < \left(|A_2|^{-9}|A_1|^{-13} 2^{-H_2^\epsilon(A_1 A_2|R)_\rho} \right)^t.$$

for values of $\kappa = O(\mu)$, $t = O(|A_1|)$. Above concentration inequality can also be expressed as:

$$\mathbb{P}_{U \sim \text{design}} [f(U) > \sqrt{\frac{|A_1|}{|A_2|}} \cdot 2^{-\frac{1}{2}H_2^\epsilon(A|R)_\rho + 1} + 14\sqrt{\epsilon} + \beta \sqrt{|A_1|/2}] \leq 7 \cdot 2^{-\frac{|A_1|}{2}}, \quad (30)$$

where $\beta := \sqrt{1/a}$. Thus, if $|A_1| \leq \text{polylog}(|A_2|)$ and $\kappa = O(\mu)$ then efficient constructions for such qTPEs exist.

Proof. In order to obtain the desired concentration result, we apply Theorem 1 with the following parameters:

- The input system A to the superoperator is $A := A_1 \otimes A_2$. Output system $B := A_1$ and superoperator $\mathcal{T}^{A \rightarrow A_1} := \text{Tr}_{A_2}$;
- The state $\omega^{A' A_1} = (\mathcal{T}^{A \rightarrow A_1} \otimes \mathbb{I}^{A'}) (\Phi^{A A'}) = \Phi^{A_1 A'_1} \otimes \pi^{A'_2}$;
- Take $\delta = 0$. We get $(\omega'''_{\epsilon,0})^{A_1} = \pi^{A_1}$. Hence $\mathbb{H}_{\max}^\epsilon(A_1)_\omega = \log|A_1|$ as the reduced state $\omega^{A_1} = \pi^{A_1}$;
- The matrix $(\tilde{\omega}')^{A' A_1} := (I^{A'_1} \otimes I^{A'_2} \otimes (\omega'''_{\epsilon,0})^{A_1})^{-1/4} \circ (\Phi^{A_1 A'_1} \otimes \pi^{A'_2}) = |A_1|^{1/2} (\Phi^{A_1 A'_1} \otimes \pi^{A'_2})$. Note that $(H_2^{\epsilon,0}(A'|A_1)_\omega = -2\log\|(\tilde{\omega}')^{A' A_1}\|_2$. We have,

$$\|(\tilde{\omega}')^{A' A_1}\|_2 = \||A_1|^{1/2} \Phi^{A_1 A'_1} \otimes \pi^{A'_2}\|_2 = \sqrt{\frac{|A_1|}{|A_2|}},$$

$$\begin{aligned}
\|(\tilde{\omega}')^{A_1}\|_2 &= \| |A_1|^{1/2} \Phi^{A_1} \|_2 = 1, \\
\eta &= \frac{\|(\tilde{\omega}')^{A'_1 A_1}\|_2^2}{\|(\tilde{\omega}')^{A_1}\|_2^2} = \frac{|A_1|}{|A_2|}, \\
\delta_1 &= \|(\tilde{\omega}')^{A_1}\|_2^2 \frac{|A|^2 - |A|\eta}{|A|^2 - 1} = \frac{|A_1|^2 |A_2|^2 - |A_1|^2}{|A_1|^2 |A_2|^2 - 1}, \\
\delta_2 &= \|(\tilde{\omega}')^{A'_1 A_1}\|_2^2 \frac{|A|^2 - |A|\eta^{-1}}{|A|^2 - 1} = \frac{|A_1|}{|A_2|} \cdot \frac{|A_1|^2 |A_2|^2 - |A_2|^2}{|A_1|^2 |A_2|^2 - 1};
\end{aligned}$$

- The function $g(U) = |A_1|^{1/2} \|(\text{Tr}_{A_2} \otimes \mathbb{I}^R)((U^A \otimes I^R) \circ (\tilde{\rho}')^{AR}) - \pi^{A_1} \otimes (\tilde{\rho}')^R\|_2$. Then,

$$\begin{aligned}
\mathbb{E}_{U \sim \text{Haar}}[(g(U))^2] &= \delta_1 \|(\tilde{\rho}')^R\|_2^2 + \delta_2 \|(\tilde{\rho}')^{AR}\|_2^2 - \|(\tilde{\omega}')^{A_1}\|_2^2 \|(\tilde{\rho}')^R\|_2^2 \\
&= -\frac{|A_1|^2 - 1}{|A_1|^2 |A_2|^2 - 1} \|(\tilde{\rho}')^R\|_2^2 + \frac{|A_1|}{|A_2|} \cdot \frac{|A_1|^2 |A_2|^2 - |A_2|^2}{|A_1|^2 |A_2|^2 - 1} \|(\tilde{\rho}')^{AR}\|_2^2 \\
&\leq \frac{|A_1|}{|A_2|} \cdot \|(\tilde{\rho}')^{AR}\|_2^2, \\
\mathbb{E}_{U \sim \text{Haar}}[(g(U))^2] &= -\frac{|A_1|^2 |A_2|^2 - |A_2|^2}{(|A_1|^2 |A_2|^2 - 1)|A_2|^2} \|(\tilde{\rho}')^R\|_2^2 + \frac{|A_1|}{|A_2|} \cdot \frac{|A_1|^2 |A_2|^2 - |A_2|^2}{|A_1|^2 |A_2|^2 - 1} \|(\tilde{\rho}')^{AR}\|_2^2 \\
&\geq \frac{0.1|A_1|}{|A_2|} \cdot \frac{|A_1|^2 |A_2|^2 - |A_2|^2}{|A_1|^2 |A_2|^2 - 1} \|(\tilde{\rho}')^{AR}\|_2^2 \\
&\geq \frac{0.1|A_1|}{|A_2|} \left(1 - \frac{|A_2|^2}{|A_1|^2 |A_2|^2}\right) \|(\tilde{\rho}')^{AR}\|_2^2 \\
&\geq \frac{0.07|A_1|}{|A_2|} \cdot \|(\tilde{\rho}')^{AR}\|_2^2;
\end{aligned}$$

- The tail probability exponent a becomes $a = |A||A_1|^{-1} 2^{H_2^\epsilon(A|R)_\rho - 9} = |A_2| 2^{H_2^\epsilon(A|R)_\rho - 9}$.
- Define $\mu := \mathbb{E}_{U \sim \text{Haar}}[g(U)]$. Since $a \cdot \mathbb{E}_{U \sim \text{Haar}}[(g(U))^2] + \log \mathbb{E}_{U \sim \text{Haar}}[(g(U))^2] > \log |A_1| + \log |A_2| + \mathbb{H}_{\max}^\epsilon(A_1)_\omega - H_2^\epsilon(A_1 A_2 | R)_\rho$, we get $\mu^2 \leq \mathbb{E}[(g(U))^2] \leq 8\mu^2$;
- The qTPE parameter t becomes $t = |A|\kappa^2 \cdot 2^{H_2^\epsilon(A|R)_\rho - 6} \cdot |A_1|^{-1} = |A_2|\kappa^2 \cdot 2^{H_2^\epsilon(A|R)_\rho - 6}$. For values of $\kappa = O(\mu)$, $t = O(|A_1|)$;
- The qTPE parameter λ becomes $\lambda = (|A|^{-8} |A_1|^{-6} \mu^2)^t$, which satisfies

$$(0.008|A_2|^{-9} |A_1|^{-13} 2^{-H_2^\epsilon(A_1 A_2 | R)_\rho})^t < \lambda < (|A_2|^{-9} |A_1|^{-13} 2^{-H_2^\epsilon(A_1 A_2 | R)_\rho})^t.$$

Sequentially iterating $O(t(\log |A_1| + \log |A_2|))$ times an $(|A_1||A_2|, s, O(1), t)$ -qTPE gives us the desired $(|A_1||A_2|, s, \lambda, t)$ -qTPE for derandomisation. Observe that if $t \leq \text{polylog}(|A_2|)$, then efficient constructions for such qTPEs exist [9], [10].

Now substituting these parameters in Theorem 1 we get

$$\mathbb{P}_{U^{A_1 A_2} \sim \text{TPE}} \left[f(U) > 2\sqrt{\frac{|A_1|}{|A_2|}} 2^{-\frac{1}{2} H_2^\epsilon(A|R)_\rho} + 14\sqrt{\epsilon} + 2\kappa \right] \leq 7 \cdot 2^{-a(\min\{\mu^2, \kappa^2\})},$$

for U chosen uniformly at random from a $(|A_1||A_2|, s, \lambda, t)$ -qTPE. This completes the proof. \square

In the following section we give two applications of a simple form of our general decoupling, via Fully Quantum Slepian Wolf Theorem 2. These applications are widely studied in theoretical physics and also gives an intuitive understanding of our theorem and a simple calculation of the various entropic quantities that govern the concentration of the decoupling theorem. These further aids the understanding that the

nature is not as random as Haar measure rather performs efficient computation, at times the parameters oblivious to us makes it more random.

VI. APPLICATIONS OF MEASURE CONCENTRATION OF THE DECOUPLING THEOREM WITH UNITARY DESIGNS VIA FQSW

A. Relative Thermalization

An immediate application of measure concentration of FQSW is in the area of quantum thermodynamics, in describing a process called relative thermalization [2]. One of the most fundamental questions in quantum thermodynamics is how a small system starting out in a particular quantum state spontaneously thermalizes when brought in contact with a much larger environment e.g. a bath. More precisely when brought in contact with a bath, the small system decouples from any other system, which we may call as the reference system, it may be initially entangled with. The formal definition of relative thermalization is as follows:

Definition 13. Let system S , environment E and reference R be quantum systems and $\Omega \subseteq S \otimes E$ be a subspace corresponding to a physical constraint such as total energy. The global system is in a state $\rho^{\Omega R}$, supported in the Hilbert space $\Omega \otimes R$. The time evolution is described by a unitary on $S \otimes E$. The state after time evolution is denoted by $\sigma^{\Omega R}$. The system S is said to be κ -thermalized relative to R in state $\sigma^{\Omega R}$ if:

$$\|\sigma^{SR} - \omega^S \otimes \sigma^R\|_1 \leq \kappa$$

where $\sigma^{SR} := \text{Tr}_E[\sigma^{\Omega R}]$ and $\omega^S \triangleq \text{Tr}_E[\frac{I_\Omega}{|\Omega|}]$ is the so called local microcanonical state.

Thus, relative thermalization requires that, after the environment E is traced out, the system S should be close to the state ω^S and should not have strong correlations with the reference R . If the time-evolution of $S \otimes E$ is modelled by a Haar random unitary on Ω , then Fact 1 guarantees that relative thermalization occurs in expectation over the Haar measure. Furthermore, Fact 3 says that $1 - \exp\left(-\frac{|\Omega|\kappa^2}{2^{-H_{\min}^\epsilon(\Omega)_\rho+6}}\right)$ fraction of Haar random unitaries achieve relative thermalization, if $2^{-\frac{1}{2}H_2^\epsilon(\Omega|R)_\rho - \frac{1}{2}H_2^\epsilon(\Omega'|S)_\omega} + 16\epsilon \leq \frac{\kappa}{2}$.

Since Haar random unitaries are computationally inefficient, it is natural to wonder whether nature truly evolves via Haar random unitary. Hence, the work of Nakata et al. [7] investigates what happens if the evolution of system plus environment is modelled by a unitary chosen from an efficiently implementable approximate unitary 2-design. Their unitary acts on the subspace Ω only. They show that relative thermalization indeed takes place for the same parameter regime as Haar random unitaries, but for a much smaller fraction $1 - \exp\left(-\frac{\kappa^4}{|\Omega|^{32-4H_{\min}^\epsilon(\Omega)_\rho+22}}\right)$ of design unitaries. It is reasonable to expect that $H_{\min}^\epsilon(\Omega)_\rho \leq \log |\Omega|/2$, i.e. ρ^Ω is not highly mixed over Ω . In this case the fraction of unitaries achieving relative thermalization is only guaranteed to be at least $1 - \exp\left(-\frac{\kappa^4}{2^{22}|\Omega|}\right)$, which is almost zero for large $|\Omega|$.

We now analyze relative thermalization using the lens of unitary designs. We follow the proof technique of Theorem 2.

- The superoperator is Tr_E with input system Ω and output system S ;
- The state $\omega^{\Omega'S} = (\text{Tr}_E \otimes \mathbb{I}^{\Omega'}) (\Phi^{\Omega\Omega'})$ and $(\tilde{\rho}')^{\Omega R} := (\xi^R)^{-1/4} \circ \rho^{\Omega R}$;
- Take $\epsilon = \frac{\kappa^2}{60}$ and $\delta = 0$. Let $2^{-\frac{1}{2}H_2^\epsilon(\Omega|R)_\rho - \frac{1}{2}H_2^{\epsilon,0}(\Omega'|S)_\omega} \leq \frac{\kappa}{4}$, $\mathbb{H}_2^{\epsilon,0}(\Omega'|S)_\omega \leq \log |\Omega| - \log |S|$, $|S| > 2$, $\mathbb{H}_{\max}^\epsilon(S)_\omega = O(\log(|S|))$, and $\|(\tilde{\rho}')^R\|_2^2 < 0.9|\Omega|\|(\tilde{\rho}')^{\Omega R}\|_2^2$;
- The matrix $(\tilde{\omega}')^{\Omega'S} := (I^{\Omega'} \otimes (\omega_{\epsilon,\delta}''')^S)^{-1/4} \circ \omega^{\Omega'S}$. Note that $(H_2')^{\epsilon,0}(\Omega'|S)_\omega = -2 \log \|(\tilde{\omega}')^{\Omega'S}\|_2$, $H_2^\epsilon(\Omega|R)_\rho = -2 \log \|(\tilde{\rho}')^{\Omega R}\|_2$. We have:

$$1 \geq \|(\tilde{\omega}')^S\|_2 \geq \sqrt{1-\epsilon},$$

$$\begin{aligned}\eta &= \frac{\|(\tilde{\omega}')^{\Omega'S}\|_2^2}{\|(\tilde{\omega}')^S\|_2^2}, \quad |\Omega|^{-1} \leq 2^{-\mathbb{H}_2^{\epsilon,0}(\Omega'|S)_\omega} \leq \eta \leq (1-\epsilon)^{-1} 2^{-\mathbb{H}_2^{\epsilon,0}(\Omega'|S)_\omega} \leq 1, \\ \delta_1 &= \|(\tilde{\omega}')^S\|_2^2 \frac{|\Omega|^2 - |\Omega|\eta}{|\Omega|^2 - 1}, \quad \|(\tilde{\omega}')^S\|_2^2(1 - |\Omega|^{-1}) \leq \alpha \leq \|(\tilde{\omega}')^S\|_2^2, \\ \delta_2 &= \|(\tilde{\omega}')^{\Omega'S}\|_2^2 \frac{|\Omega|^2 - |\Omega|\eta^{-1}}{|\Omega|^2 - 1}, \quad \|(\tilde{\omega}')^{\Omega'S}\|_2^2(1 - |S|^{-1}) \leq \beta \leq \|(\tilde{\omega}')^{\Omega'S}\|_2^2;\end{aligned}$$

- The function

$$g(U) := \left\| (I^R \otimes (\omega'''_{\epsilon,\delta})^S)^{-1/4} \circ ((\text{Tr}_E \otimes \mathbb{I}^R)((U^\Omega \otimes I^R) \circ (\tilde{\rho}')^{\Omega R}) - \omega^S \otimes (\tilde{\rho}')^R) \right\|_2. \text{ Then,}$$

$$\begin{aligned}\mathbb{E}_{U \sim \text{Haar}}[(g(U))^2] &= \delta_1 \|(\tilde{\rho}')^R\|_2^2 + \delta_2 \|(\tilde{\rho}')^{\Omega R}\|_2^2 - \|(\tilde{\omega}')^S\|_2^2 \|(\tilde{\rho}')^R\|_2^2 \\ &\leq \|(\tilde{\omega}')^{\Omega'S}\|_2^2 \|(\tilde{\rho}')^{\Omega R}\|_2^2, \\ \mathbb{E}_{U \sim \text{Haar}}[(g(U))^2] &\geq \|(\tilde{\omega}')^{\Omega'S}\|_2^2 \|(\tilde{\rho}')^{\Omega R}\|_2^2 (1 - |S|^{-1}) - \|(\tilde{\omega}')^S\|_2^2 \|(\tilde{\rho}')^R\|_2^2 |\Omega|^{-1} \\ &\geq 0.05 \|(\tilde{\omega}')^{\Omega'S}\|_2^2 \|(\tilde{\rho}')^{\Omega R}\|_2^2\end{aligned}$$

- The tail probability exponent a becomes $a \geq |\Omega||S|^{-1} 2^{H_2^\epsilon(\Omega|R)_\rho - 9}$;
- Define $\mu := \mathbb{E}_{\text{Haar}}[g(U)]$. We ensure that $a \cdot \mathbb{E}_{U \sim \text{Haar}}[(g(U))^2] + \log \mathbb{E}_{U \sim \text{Haar}}[(g(U))^2] > \log |\Omega| + \mathbb{H}_{\max}^\epsilon(S)_\omega - H_2^\epsilon(\Omega|R)_\rho$, so that we get $\mu^2 \leq \mathbb{E}[(g(U))^2] \leq 8\mu^2$;
- The qTPE parameter t becomes $t = 8a\kappa^2$. If $\kappa^2 = \text{poly}(|S|)|\Omega|^{-1} \cdot 2^{-H_2^\epsilon(\Omega|R)_\rho}$ (which is the case when κ is close to μ in FQSW), then $t \leq \text{poly}(|S|)$;
- The qTPE parameter λ becomes $\lambda = (|\Omega|^{-8}|S|^{-6}\mu^2)^t$. Sequentially iterating $O(t \log |\Omega|)$ times an $(|\Omega|, s, O(1), t)$ -qTPE gives us the desired $(|\Omega|, s, \lambda, t)$ -qTPE for derandomisation. Observe that if $t \leq \text{polylog}(|\Omega|)$, then efficient constructions for such qTPEs exist [9], [10].

We have thus proved the following theorem for relative thermalization.

Theorem 3. Consider the setting of Theorem 1 and Definition 13. Suppose we are promised that

$$\begin{aligned}2^{-\frac{1}{2}H_2^\epsilon(\Omega|R)_\rho - \frac{1}{2}\mathbb{H}_2^{\epsilon,0}(\Omega'|S)_\omega} &\leq \frac{\kappa}{4}, \mathbb{H}_2^{\epsilon,0}(\Omega'|S)_\omega \leq \log |\Omega| - \log |S|, |S| > 2, \mathbb{H}_{\max}^\epsilon(S)_\omega = O(\log(|S|)), \\ \|(\tilde{\rho}')^R\|_2^2 &< 0.9|\Omega| \|(\tilde{\rho}')^{\Omega R}\|_2^2, 2^{-14}|\Omega||S|^{-1} \cdot 2^{H_2^\epsilon(\Omega|R)_\rho} - \mathbb{H}_2^{\epsilon,0}(\Omega'|S)_\omega > 2 \log |\Omega|.\end{aligned}$$

Then S is κ -thermalized relative to R in state $\sigma^{\Omega R}$ for an $1 - 5 \cdot 2^{-a\kappa^2}$ fraction of unitaries U on Ω where U is chosen uniformly at random from a $(|\Omega|, s, \lambda, t)$ -qTPE $a := |\Omega||S|^{-1} \cdot 2^{H_2^\epsilon(\Omega|R)_\rho - 9}$, $t := 8a\kappa^2$ and $\lambda := (|\Omega|^{-8}|S|^{-6}\mu^2)^t$. For $\kappa^2 = \text{poly}(|S|)|\Omega|^{-1} \cdot 2^{-H_2^\epsilon(\Omega|R)_\rho}$, $t \leq \text{poly}(|S|)$. For such κ , which includes the case where $\kappa = O(\mu)$, if $|S| \leq \text{polylog}(|\Omega|)$, then efficient constructions for such qTPEs exist.

Theorem 3 achieves better performance than the result of Dupuis (Fact 3) and the result of Nakata et al. [7] in the following sense:

- 1) In our result, the system plus environment evolves according to a unitary chosen uniformly at random from an approximate unitary t -design for moderate values of t . Our unitary acts on the subspace Ω only. For a wide range of parameters, our unitaries require $O(|S| \log |\Omega|)$ random bits for a precise description which is less than the $\Omega(|\Omega|^2 \log |\Omega|)$ random bits required by the Haar random unitaries of Dupuis, as well as less than the $\Omega(|\Omega| \log |\Omega|)$ random bits required by the approach of Nakata et al. Moreover, the random unitaries used by Dupuis and by Nakata et al. are not efficiently implementable, whereas our unitaries are efficiently implementable when $|S| \leq \text{polylog}(|\Omega|)$.
- 2) Our Theorem 3 shows that relative thermalisation still takes place for the fraction $1 - 5 \cdot 2^{-a\kappa^2}$ of unitaries, where $a = |\Omega||S|^{-1} \cdot 2^{H_2^\epsilon(\Omega|R)_\rho - 9}$. Note that $H_2^\epsilon(\Omega|R)_\rho \geq -\log |\Omega|$ for any state $\rho^{\Omega R}$.

The equality is achieved when $\rho^{\Omega R}$ is maximally entangled on Ω . Under the reasonable assumption that $H_2^\epsilon(\Omega|R)_\rho \geq -0.5 \log |\Omega|$, i.e., $\rho^{\Omega R}$ is not highly entangled on Ω , the fraction of unitaries that achieve relative thermalisation is at least $1 - 5 \cdot 2^{-2^{-9} \cdot |\Omega|^{1/2} |S|^{-1} \kappa^2}$. As Ω is generally of a much larger dimension than the system S , it is reasonable to expect that $|S| < |\Omega|^{1/4}$. In that case, the fraction of unitaries that achieve relative thermalisation in our result is guaranteed to be at least $1 - 5 \cdot 2^{-2^{-9} \cdot |\Omega|^{1/4} \kappa^2}$ which is nearly one for large $|\Omega|$. We also assume that $H_{\min}^\epsilon(\Omega) \leq \log |\Omega|/2$, i.e. the state ρ^Ω is not highly mixed on Ω . For this range of parameters, our decoupling result is much better than that of Nakata et al. which can only guarantee that $1 - \exp\left(-\frac{\kappa^4}{2^{22}|\Omega|}\right) \approx 0$ fraction of unitaries achieve relative thermalisation. However, our result is worse than that of Dupuis which guarantees that $1 - \exp\left(-\frac{|\Omega|^{3/2} \kappa^2}{2^6}\right)$ fraction of Haar random unitaries achieve relative thermalisation. Note that the unitaries of Nakata et al. and Dupuis used for obtaining concentration of measure for decoupling, and consequently for relative thermalisation, are not efficient to implement. Only the unitaries used to obtain decoupling in expectation by Nakata et al. can be implemented efficiently.

We summarise our relative thermalization result in Table III for a clear comparison with the known results of Dupuis and Nakata et al.

	Dupuis [3]	Nakata et al. [7]	Theorem 3
Fraction	$1 - \exp\left(- \Omega ^{3/2} \kappa^2\right)$	$1 - \exp\left(- \Omega ^{-1} \kappa^2\right)$	$1 - \exp\left(- \Omega ^{1/2} \kappa^2\right)$
Randomness	$ \Omega ^2 \log \Omega $	$ \Omega \log \Omega $	$\text{poly}(S) \log \Omega $
Random unitary	Haar	X and Z -diagonal	Approx. $\text{poly}(S)$ -design
Efficiency	Always inefficient	Always inefficient	Efficient for $ S = \text{polylog}(\Omega)$

TABLE III

ACHIEVING κ -RELATIVE THERMALIZATION, $\kappa^2 = \text{poly}(|S|)|\Omega|^{-1} \cdot 2^{-H_2^\epsilon(A|R)_\rho}$, $H_{\min}^\epsilon(\Omega)_\rho = \log |\Omega|/2$, $H_2^\epsilon(\Omega|R)_\rho \geq -0.5 \log |\Omega|$, CONSTANT FACTORS IGNORED.

Remark 7. Nakata et al.'s work and other works on relative thermalisation shows that for some well known Hamiltonians in physics relative thermalisation eventually occurs. Concentration bounds, even if given, could not answer our everyday observation in nature that thermalisation occurs with overwhelmingly high probability. Prior to this work, only Haar random unitaries were known to achieve thermalisation with overwhelmingly high probability. Our work is the first one to show that relative thermalisation can indeed take place with overwhelmingly high probability using unitaries that are simpler than Haar random in a precise sense. Moreover, for a particular range of parameters, relative thermalisation can even be achieved by computationally efficient unitaries. Physically this work lays out a clear direction of further research, that simple unitaries made up of a few basic building blocks can achieve relative thermalisation with overwhelmingly high probability.

B. Application to the black hole information paradox model by Hayden-Preskill [31]

A brief description of Hayden-Preskill toy model for the black hole information paradox [31]: Hayden-Preskill in their seminal paper [31] described the evolution of black hole as a unitary operator. Further they assumed that an old black hole which has emitted half of its information via Hawking's radiation can be assumed to be in a maximally entangled state with its environment. Alice throws a quantum register of $\log |M|$ qubits in a maximally entangled state with its reference N , in such an old black hole. Subsequently,

after the black hole evolves unitarily and emits Hawking radiation, a receiver say Bob, can intercept the radiation and can decode Alice's information by waiting for no longer than $O(\log |M|)$ qubits of radiation. Hayden-Preskill applied the decoupling theorem, by assuming black hole evolution as a Haar random unitary, to conclude that for a 'typical' black hole Alice's information can be decoded by Bob provided Bob has access to the initial black hole's environment, and emitted Hawking radiation. This is because if a black hole's internal state, after interaction with Alice's register and evolution, is decoupled with Alice's purifying environment, then Uhlmann's theorem gives a decoder that maps black hole's initial and received radiations to purifying register for Alice's reference. This information retrieval property, similar to channel coding problem was first observed by Hayden-Preskill with the above mentioned assumptions in [31]. This phenomenon was termed as information mirror or scrambling.

Now, if we assume that black holes must scramble efficiently, is this information retrieval property still generic to a typical but efficiently evolving black hole? That is, even assuming efficiently implementable dynamics, is the fraction of black holes that do not behave like "information mirrors" still exponentially small? We provide an affirmative answer to the above question.

Hayden-Preskill [31] toy model setup:

- Alice's initial state: A register of $\log |M|$ qubits. It is assumed to be in a maximally entangled state with a purifying reference (N), denoted by: $|\Phi\rangle^{MN} := \frac{1}{|M|} \sum_{i=1}^{|M|} |i\rangle^M |i\rangle^N$.
- Black hole is denoted by system B_1 (before interacting with Alice's register). It is assumed to be maximally entangled with an environment E (initial radiations), denoted by: $|\Phi\rangle^{B_1 E} := \frac{1}{|B_1|} \sum_{j=1}^{|B_1|} |j\rangle^{B_1} |j\rangle^E$.
- Evolution of black hole is modelled by a unitary operator $U^{MB_1 \rightarrow RB_2}$. Thus $|M| \times |B_1| = |R| \times |B_2|$.
- Emitted radiations from black hole after evolution are denoted by quantum system R .
- Final state of black hole is denoted by quantum system B_2 . B_2 is inaccessible to Bob.
- The decoupling superoperator $\mathcal{T}^{RB_2 E \rightarrow B_2} := \text{Tr}_{RE}$, as the systems E and N are in product state from the beginning.
- The input state to the "decoupling theorem" $\Phi^{MN} \otimes \Phi^{B_1 E}$.
- The decoupling step works by analyzing the distance of

$$\begin{aligned} & [\mathcal{T}^{REB_2 \rightarrow B_2} \otimes \mathbb{I}^N] \{ (U^{MB_1 \rightarrow RB_2} \otimes I^{NE}) \circ (\Phi^{MN} \otimes \Phi^{B_1 E}) \} \\ &= [\text{Tr}_{RE} \otimes \mathbb{I}^N] \{ (U^{MB_1 \rightarrow RB_2} \otimes I^{NE}) \cdot (\Phi^{MN} \otimes \Phi^{B_1 E}) \} \\ &= [\text{Tr}_R \otimes \mathbb{I}^N] \{ (U^{MB_1 \rightarrow RB_2} \otimes I^N) \circ (\Phi^{MN} \otimes \Phi^{B_1 E}) \} \end{aligned}$$

with its expected value equal to $\pi^{B_2} \otimes \pi^N$. This shows that the 'average' state on the joint system N, E, R is a pure state and hence there exists a decoding isometry $V_{\mathcal{D}}^{RE \rightarrow M}$ can map this pure state to Φ^{MN} , thereby giving Bob the access to Alice's initial quantum information.

For the rest of this section we work with the following decoupling function:

$$f(U) := \left\| [\text{Tr}_R \otimes \mathbb{I}^N] \{ (U^{MB_1 \rightarrow RB_2} \otimes I^N) \cdot (\Phi^{MN} \otimes \pi^{B_1}) \} - \pi^{B_2} \otimes \pi^N \right\|_1 \quad (31)$$

We follow the proof technique of Theorem 2.

- The input states in the notation of the Theorem 2 are defined as $\rho^{MB_1 N} := \Phi^{MN} \otimes \pi^N$ and $(\tilde{\rho}')^{MB_1 N} := (\xi^N)^{-1/4} \circ \rho^{MB_1 N}$. Note that $|M| = |N|$.
The choice of the weighting matrix $\xi^N = \pi^N$ gives $H_2^\epsilon(A|R)_\rho = -\log \|(\tilde{\rho}')^2\|_2^2 = \log \frac{|B_1|}{|M|}$.
- The state $\omega^{R' B_2 B_2} := [\text{Tr}_R \otimes \mathbb{I}^{R' B_2}] (\Phi^{RR'} \otimes \Phi^{B_2 B_2}) = \pi^R \otimes \Phi^{B_2 B_2}$,
- Take $\delta = 0$. We get $(\omega_{\epsilon,0}'')^{B_2} = \pi^{B_2}$. Hence $\mathbb{H}_{\max}^\epsilon(B_2)_\omega = \log |B_2|$ as the reduced state $\omega^{B_2} = \pi^{B_2}$.

- The matrix $(\tilde{\omega}')^{R'B_2'B_2} := (I^{B_2'} \otimes I^{R'} \otimes (\omega_{\epsilon,0}''')^{B_2})^{-1/4} \circ (\Phi^{B_2B_2'} \otimes \pi^{R'}) = |B_2|^{1/2}(\Phi^{B_2B_2'} \otimes \pi^{R'})$. Note that $\mathbb{H}_2^{\epsilon,0}(R'B_2'|B_2)_\omega = -2 \log \|(\tilde{\omega}')^{R'B_2'B_2}\|_2$. We have,

$$\begin{aligned} \|(\tilde{\omega}')^{R'B_2'B_2}\|_2 &= \| |B_2|^{1/2} \Phi^{B_2B_2'} \otimes \pi^{R'} \|_2 = \sqrt{\frac{|B_2|}{|R|}}, \\ \|(\tilde{\omega}')^{B_2}\|_2 &= \| |B_2|^{1/2} \Phi^{B_2} \|_2 = 1, \\ \eta &= \frac{\|(\tilde{\omega}')^{R'B_2'B_2}\|_2^2}{\|(\tilde{\omega}')^{B_2}\|_2^2} = \frac{|B_2|}{|R|}, \\ \delta_1 &= \|(\tilde{\omega}')^{B_2}\|_2^2 \frac{|R|^2|B_2|^2 - |R||B_2|\eta}{|B_2|^2|R|^2 - 1} = \frac{|B_2|^2|R|^2 - |B_2|^2}{|B_2|^2|R|^2 - 1}, \\ \delta_2 &= \|(\tilde{\omega}')^{B_2'R'B_2}\|_2^2 \frac{|B_2|^2|R|^2 - |B_2||R|\eta^{-1}}{|B_2|^2|R|^2 - 1} = \frac{|B_2|}{|R|} \cdot \frac{|B_2|^2|R|^2 - |R|^2}{|B_2|^2|R|^2 - 1}; \end{aligned}$$

- The function $g(U) = |B_2|^{1/2} \|\text{Tr}_R \otimes \mathbb{I}^N\|((U^{MB_1 \rightarrow RB_2} \otimes I^N) \circ (\tilde{\rho}')^{MB_1N}) - \pi^{B_2} \otimes (\tilde{\rho}')^N\|_2$. Then,

$$\begin{aligned} \mathbb{E}_{U \sim \text{Haar}}[(g(U))^2] &= \delta_1 \|(\tilde{\rho}')^N\|_2^2 + \delta_2 \|(\tilde{\rho}')^{MB_1N}\|_2^2 - \|(\tilde{\omega}')^{B_2}\|_2^2 \|(\tilde{\rho}')^N\|_2^2 \\ &= -\frac{|B_2|^2 - 1}{|B_2|^2|R|^2 - 1} \|(\tilde{\rho}')^N\|_2^2 + \frac{|B_2|}{|R|} \cdot \frac{|B_2|^2|R|^2 - |R|^2}{|B_2|^2|R|^2 - 1} \|(\tilde{\rho}')^{MB_1N}\|_2^2 \\ &\leq \frac{|B_2|}{|R|} \cdot \|(\tilde{\rho}')^{MB_1N}\|_2^2 = \frac{|M|^2}{|R|^2}, \\ \mathbb{E}_{U \sim \text{Haar}}[(g(U))^2] &= -\frac{|B_2|^2|R|^2 - |R|^2}{(|B_2|^2|R|^2 - 1)|R|^2} \|(\tilde{\rho}')^N\|_2^2 + \frac{|B_2|}{|R|} \cdot \frac{|B_2|^2|R|^2 - |R|^2}{|B_2|^2|R|^2 - 1} \|(\tilde{\rho}')^{MB_1N}\|_2^2 \\ &\geq \frac{0.1|B_2|}{|R|} \cdot \frac{|B_2|^2|R|^2 - |R|^2}{|B_2|^2|R|^2 - 1} \|(\tilde{\rho}')^{MB_1N}\|_2^2 \\ &\geq \frac{0.1|B_2|}{|R|} \left(1 - \frac{|R|^2}{|B_2|^2|R|^2}\right) \|(\tilde{\rho}')^{MB_1N}\|_2^2 \\ &\geq \frac{0.07|B_2|}{|R|} \cdot \|(\tilde{\rho}')^{MB_1N}\|_2^2 = 0.07 \frac{|M|^2}{|R|^2}; \end{aligned}$$

- The tail probability exponent a becomes $a = |RB_2||B_2|^{-1} 2^{H_2^\epsilon(MB_1|N)_\rho - 9} = 2^{-9} \frac{|R||B_1|}{|M|}$.
- Define $\mu := \mathbb{E}_{\text{Haar}}[g(U)]$. By a direct calculation we have $a \cdot \mathbb{E}_{U \sim \text{Haar}}[(g(U))^2] + \log \mathbb{E}_{U \sim \text{Haar}}[(g(U))^2] > \log |B_2| + \log |R| + \mathbb{H}_{\max}^\epsilon(B_2)_\omega - H_2^\epsilon(MB_1|N)_\rho$, and thus we get $\mu^2 \leq \mathbb{E}[(g(U))^2] \leq 8\mu^2$;
- The qTPE parameter t becomes $t = |M||B_1|\kappa^2 \cdot 2^{H_2^\epsilon(MB_1|N)_\rho - 6} \cdot |B_2|^{-1}$. For values of $\kappa = O(\mu)$, $t = O(|B_2|)$;
- The qTPE parameter λ becomes $\lambda = (|MB_1|^{-8}|B_2|^{-6}\mu^2)^t$, which satisfies

$$(0.008|R|^{-9}|B_2|^{-13}2^{-H_2^\epsilon(MB_1|N)_\rho})^t < \lambda < (|R|^{-9}|B_2|^{-13}2^{-H_2^\epsilon(MB_1|N)_\rho})^t.$$

Sequentially iterating $O(t(\log |M| + \log |B_1|))$ times an $(|M||B_1|, s, O(1), t)$ -qTPE gives us the desired $(|M||B_1|, s, \lambda, t)$ -qTPE for efficient evolution. Observe that if $t \leq \text{polylog}(|R|)$, then efficient constructions for such qTPEs exist [9], [10].

We summarize our conclusion as follows:

Theorem 4 (FQSW concentration and black hole as mirror). *If an old black hole, that has already radiated half of its matter, evolves according to an approximate unitary $O(|B_2|)$ -design then it satisfies the following concentration inequality:*

$$\mathbb{P}_{U \sim \text{design}} \left[f(U) > \frac{|M|}{|R|} + 14\sqrt{\epsilon} + 2\kappa \right] \leq 7 \cdot 2^{-a\kappa^2},$$

where the unitary $U_{BH}^{MB_1 \rightarrow RB_2}$ is chosen uniformly at random from an approximate t -design, $a := 2^{-9} \frac{|R||B_1|}{|M|}$ and $t := 8a\kappa^2$. The above concentration inequality can also be expressed directly in terms of the parameter t of the evolution of the black hole according to a unitary chosen uniformly at random from an approximate unitary t -design as:

$$\mathbb{P}_{U \sim \text{design}} \left[f(U) > \frac{|M|}{|R|} + 14\sqrt{\epsilon} + \sqrt{t/2} \frac{|M|}{|R||B_1|} \right] \leq 7 \cdot 2^{-\frac{t}{8}},$$

This theorem in turn implies that an overwhelming fraction of such black holes act as an information mirror. Further, one has to wait no longer than the size of Alice's register in order to decode Alice's information scrambled by the such a black hole. Such a unitary can be described by using $O(|B_2| \log(|M||B_1|))$ random bits as opposed to a Haar random black hole that needs $O(|M|^2 |B_1|^2 \log(|M||B_1|))$ for a reasonable approximation.

Moreover, if $|B_2| \leq O(\text{polylog}(|R|))$ and $\kappa = O\left(\frac{|M|}{|R|}\right)$ which further implies that $t = O(|B_2|) = O(\text{poly log } |R|)$, then a typical black hole not only acts as an information mirror but can also be described in a computationally efficient manner (or simulated efficiently on a quantum computer).

VII. CONCLUSION

In this work we obtain a novel concentration result for one-shot non-catalytic decoupling via approximate unitary t -designs for moderate values of t . Our bounds are stated in terms of one-shot smooth variants of Rényi 2-entropies and max-entropies. We then consider the asymptotic iid limit of our concentration result and show that the bounds reduce to the standard Shannon entropies. Finally, we apply our concentration result to a special case when the superoperator \mathcal{T} is just the partial trace. This case is also referred to as the Fully Quantum Slepian Wolf theorem in the literature, for its application to lossless quantum source compression demonstrated in [1]. This leads to a new result on relative thermalisation of quantum systems. In particular for systems that are much smaller than their ambient spaces, we show that for a wide range of parameters relative thermalisation can be achieved with probability exponentially close to one using efficiently implementable unitaries. This is the first result of this kind.

We also apply our FQSW result to the Hayden-Preskill toy model for the black hole information paradox. We show that the information mirror property observed by Hayden and Preskill for a Haar random unitary black hole continues to hold with probability exponentially close to one even when the black hole evolution is restricted to be a uniformly random unitary picked from a small discrete set of 'simple' i.e. computationally efficient unitaries.

For larger systems, it is unknown whether suitable efficient approximate t -designs exist. Hence the question of whether relative thermalisation can be achieved by efficiently implementable unitaries with exponentially high probability in the general case still remains open.

Several applications of the original decoupling theorem in expectation are known in the literature. Our result can be applied to many of them obtaining, for the first time, corresponding concentration results via approximate unitary t -designs. Whether these concentration results have any operational significance is a topic left for future research.

ACKNOWLEDGEMENT

AN would like to thank Prof. Francesco Buscemi for pointing out Hayden-Preskill toy model for black hole information paradox as a potential application to our main theorem. We acknowledge support of the Department of Atomic Energy, Government of India, under project no. 12-R&D-TFR-5.01-0500, for carrying out this research work. AN acknowledges support from the European Research Council (ERC Grant Agreement No. 948139) and also from MEXT Quantum Leap Flagship Program (MEXT QLEAP) Grant No. JPMXS0120319794.

REFERENCES

- [1] A. Abeyesinghe, I. Devetak, P. Hayden, and A. Winter, “The mother of all protocols: Restructuring quantum information’s family tree,” *Proceedings of the Royal Society of London Series A*, vol. 465, pp. 2537–2563, 2009. 1, 37, 44
- [2] L. del Rio, A. Hutter, R. Renner, and S. Wehner, “Relative thermalization,” *Physical Review E*, vol. 94, pp. 022 104:1–022 104:15, 2016. 2, 39
- [3] F. Dupuis, “The decoupling approach to quantum information theory,” Ph.D. dissertation, Université de Montréal, Canada, 2010. 2, 3, 6, 10, 11, 12, 13, 24, 41
- [4] O. Szechr, F. Dupuis, M. Tomamichel, and R. Renner, “Decoupling with unitary approximate two-designs,” *New Journal of Physics*, vol. 15, pp. 053 022:1–053 022:20, 2013. 3, 6, 35
- [5] A. Anshu and R. Jain, “Efficient methods for one-shot quantum communication,” 2018, available at arXiv:1809.07056. 3
- [6] A. Ambainis and A. Smith, “Small pseudo-random families of matrices: Derandomizing approximate quantum encryption,” in *Proceedings of the 8th International Workshop on Randomization and Computation (RANDOM)*, 2004, pp. 249–260, lecture Notes in Computer Science (LNCS), vol. 3122, Springer-Verlag. 3
- [7] Y. Nakata, C. Hirche, C. Morgan, and A. Winter, “Decoupling with random diagonal unitaries,” *Quantum*, vol. 1, pp. 18–34, 2017. 3, 39, 40, 41
- [8] G. Kuperberg, “Numerical cubature from Archimedes’ Hat-Box theorem,” *SIAM Journal on Numerical Analysis*, vol. 44, no. 3, pp. 908–935, 2006. 4
- [9] F. Brandao, A. Harrow, and M. Horodecki, “Local random quantum circuits are approximate polynomial-designs,” *Communications in Mathematical Physics*, vol. 346, pp. 397–434, 2016. 6, 18, 38, 40, 43
- [10] P. Sen, “Efficient quantum tensor product expanders and unitary t -designs via the zigzag product,” 2018, arXiv:1808.10521. 6, 18, 38, 40, 43
- [11] R. Low, “Large deviation bounds for k -designs,” *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, vol. 465, no. 2111, pp. 3289–3308, 2009. 7, 15, 17, 18, 25
- [12] P. Sen, “A quantum Johnson-Lindenstrauss lemma via unitary t -designs,” 2018, available at arXiv:1807.08779. 7, 15
- [13] A. Kitaev and J. Watrous, “Parallelization, amplification, and exponential time simulation of quantum interactive proof systems,” in *32nd Annual ACM Symposium on Theory of Computing (STOC)*, 2000, pp. 608–617. 9
- [14] J. Watrous, “Notes on super-operator norms induced by Schatten norms,” 2004, available at arXiv:quant-ph/0411077. 9
- [15] G. Aubrun, S. Szarek, and E. Werner, “Nonadditivity of Rényi entropy and Dvoretzky’s theorem,” *Journal of Mathematical Physics*, vol. 51, pp. 022 102:1–022 102:7, 2010. 9
- [16] A. Winter, “Coding theorem and strong converse for quantum channels,” *IEEE Transactions on Information Theory*, vol. 45, no. 7, pp. 2481–2485, 1999. 11
- [17] P. Sen, “Achieving the Han-Kobayashi inner bound for the quantum interference channel,” in *IEEE International Symposium on Information Theory (ISIT)*, 2012, pp. 736–740, full version at arXiv:1109.0802. 11, 14, 15
- [18] M. Tomamichel, R. Colbeck, and R. Renner, “A fully quantum asymptotic equipartition property,” *IEEE Transactions on Information Theory*, vol. 55, no. 12, pp. 5840–5847, 2009. 13, 14
- [19] S. Chakraborty, A. Nema, and F. Buscemi, “One-shot purity distillation with local noisy operations and one-way classical communication,” 2022. [Online]. Available: <https://arxiv.org/abs/2208.05628> 13
- [20] M. Hayashi and H. Nagaoka, “General formulas for capacity of classical-quantum channels,” *IEEE Transactions on Information Theory*, vol. 49, no. 7, pp. 1753–1768, 2003. 14
- [21] T. S. Han, *Information-Spectrum Methods in Information Theory*. Springer Publishing Company, Incorporated, 2014. 14
- [22] V. Y. F. Tan and A. Roth, 2014. 14
- [23] M. Tomamichel, M. Berta, and M. Hayashi, “Relating different quantum generalizations of the conditional Rényi entropy,” *Journal of Mathematical Physics*, vol. 55, pp. 082 206:1–082 206:10, 2014. 14
- [24] R. Alicki and M. Fannes, “Continuity of quantum mutual information,” 2003, available at arXiv:quant-ph/0312081. 14
- [25] El Gamal, A. and Kim, Y., *Network Information Theory*. Cambridge University Press, 2012. 14
- [26] P. Lévy and F. Pellegrino, *Problèmes concrets d’analyse fonctionnelle*, ser. Collection de monographies sur la théorie des fonctions. Gauthier-Villars, 1951. [Online]. Available: <https://books.google.co.jp/books?id=g0cPMQAACAAJ> 15
- [27] Anderson, G., Guionnet, A., and Zeitouni, O., *An introduction to random matrices*. Cambridge University Press, 2009. 15
- [28] M. Ledoux and M. Talagrand, *Probability in Banach Spaces: Isoperimetry and Processes*, ser. A Series of Modern Surveys in Mathematics Series. Springer, 1991. [Online]. Available: <https://books.google.co.jp/books?id=cyKYDfvxRjsC> 15

- [29] R. Vershynin, *High-Dimensional Probability: An Introduction with Applications in Data Science*, ser. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 2018. [Online]. Available: <https://books.google.co.jp/books?id=J-VjswEACAAJ> 15
- [30] M. Bellare and J. Rompel, “Randomness-efficient oblivious sampling,” in *35th Annual Symposium on Foundations of Computer Science (FOCS)*, 1994, pp. 276–287. 15
- [31] P. Hayden and J. Preskill, “Black holes as mirrors: quantum information in random subsystems,” *Journal of High Energy Physics*, vol. 2007, no. 09, p. 120–120, Sep 2007. [Online]. Available: <http://dx.doi.org/10.1088/1126-6708/2007/09/120> 41, 42