

The group of rational points on the Holm curve is torsion-free

Fredrick M. Nelson

February 7, 2021

Contents

1	Introduction	1
2	Definitions	2
3	Isomorphism	2
4	Three Lemmas	3
5	Division Polynomials	4
6	Proofs	5

Abstract

Using the division polynomials for elliptic curves in Weierstrass form, it is shown that the group of rational points on the curve $H : k(y^3 - y) = l(x^3 - x)$ is torsion-free.

1 Introduction

The equation of the Holm curve is $H : k(y^3 - y) = l(x^3 - x)$ where k and l are distinct, positive, relatively prime, square-free integers. Holm [1] used H to show how to construct rational right triangles whose areas have ratio k/l .

A positive integer n is a congruent number if there exists a rational right triangle with area n . Rajan and Ramarason [2] used H to show that there are infinitely many congruent numbers with ratio k/l .

It will be shown that the group of rational points on H has no points of finite order.

2 Definitions

\mathbb{Z} is the set of integers and \mathbb{Q} is the set of rational numbers.

If p is prime and $x \in \mathbb{Q} - \{0\}$, then $v_p(x)$ is the index of p in the prime factorization of x .

$$H(\mathbb{Q}) = \{(x, y) \in \mathbb{Q}^2 : k(y^3 - y) = l(x^3 - x)\} \text{ and}$$

$$E(\mathbb{Q}) = \{(x, y) \in \mathbb{Q}^2 : y^2 = x^3 - 3k^2l^2x + k^2l^2(k^2 + l^2)\} \cup \{\infty\}.$$

If $P = (a, b) \in E(\mathbb{Q})$, then $x(P) = a$ and $y(P) = b$.

3 Isomorphism

By Mordell's theorem, $H(\mathbb{Q})$ and $E(\mathbb{Q})$ are finitely generated abelian groups. Choose $(0, 0)$ as the identity element for $H(\mathbb{Q})$ and the point at infinity ∞ as the identity element for $E(\mathbb{Q})$.

Then the map $\gamma : H(\mathbb{Q}) \rightarrow E(\mathbb{Q})$ given by

$$\gamma(x, y) = \left(\frac{kl(kx - ly)}{lx - ky}, \frac{kl(k^2 - l^2)}{lx - ky} \right) \quad (1)$$

is an isomorphism. The inverse map is given by

$$\gamma^{-1}(x, y) = \left(\frac{k(x - l^2)}{y}, \frac{l(x - k^2)}{y} \right). \quad (2)$$

4 Three Lemmas

$E(\mathbb{Q})$ is in Weierstrass form so the points of finite order of $E(\mathbb{Q})$ are integral by the Nagell-Lutz Theorem. Since γ is an isomorphism, we need only show that each non-identity element of $E(\mathbb{Q})$ has a multiple which is not integral. For this, it suffices to prove the following three lemmas.

Lemma 1:

If (x, y) is an integral point in $E(\mathbb{Q})$ and 2 divides k or l , then

$$v_2(x(2(x, y))) = \begin{cases} 0 & \text{if } v_2(x) \geq 2, \\ 2 & \text{if } v_2(x) = 1, \\ -2 & \text{if } v_2(x) = 0. \end{cases}$$

Hence, $8(x, y)$ is not an integral point of $E(\mathbb{Q})$.

Lemma 2:

If (x, y) is an integral point in $E(\mathbb{Q})$ and p is an odd prime dividing k or l , then

$$v_p(x(3(x, y))) \leq \begin{cases} -2 & \text{if } p = 3, \\ 0 & \text{if otherwise.} \end{cases}$$

Hence, $3(x, y)$ is not an integral point of $E(\mathbb{Q})$.

Lemma 3: If (x, y) is an integral point in $E(\mathbb{Q})$ and $p \geq 5$ is a prime dividing k or l , then $v_p(x(3p(x, y))) \leq -2$.

Hence, $3p(x, y)$ is not an integral point of $E(\mathbb{Q})$.

5 Division Polynomials

Let $a, b \in \mathbb{Z}$ where $4a^3 + 27b^2 \neq 0$. Then $y^2 = x^3 + ax + b$ is an elliptic curve in Weierstrass form.

The division polynomials ψ_n are defined as follows:

$$\psi_0 = 0,$$

$$\psi_1 = 1,$$

$$\psi_2 = 2y,$$

$$\psi_3 = 3x^4 + 6ax^2 + 12bx - a^2,$$

$$\psi_4 = y(4x^6 + 20ax^4 + 80bx^3 - 20a^2x^2 - 16abx - 4a^3 - 32b^2),$$

$$\psi_{2n+1} = \psi_{n+2}\psi_n^3 - \psi_{n-1}\psi_{n+1}^3 \text{ for } n \geq 2, \text{ and}$$

$$\psi_{2n} = \frac{\psi_n(\psi_{n+2}\psi_{n-1}^2 - \psi_{n-2}\psi_{n+1}^2)}{2y} \text{ for } n \geq 3.$$

Auxiliary functions ϕ_n and ω_n are defined from ψ_n by

$$\phi_n = x\psi_n^2 - \psi_{n-1}\psi_{n+1} \text{ for } n \geq 1 \text{ and}$$

$$\omega_n = \frac{\psi_{n-1}^2\psi_{n+2} - \psi_{n-2}\psi_{n+1}^2}{4y} \text{ for } n \geq 2.$$

From these definitions it follows that $n(x, y) = \left(\frac{\phi_n}{\psi_n^2}, \frac{\omega_n}{\psi_n^3}\right)$ if $n \geq 2$ and (x, y) is a point on the curve $y^2 = x^3 + ax + b$.

The following proposition follows from the properties of division polynomials described in [3].

Proposition

$$(i) \ 2(x, y) = \left(\frac{x^4 - 2ax^2 - 8bx + a^2}{4y^2}, \frac{x^6 + 5ax^4 + 20bx^3 - 5a^2x^2 - 4abx - a^3 - 8b^2}{8y^3}\right).$$

$$(ii) \ x(3(x, y)) = \frac{x^9 - 12ax^7 - 96bx^6 + 30a^2x^5 - 24abx^4 + 12(3a^3 + 4b^2)x^3 + 48a^2bx^2 + 3a(3a^3 + 32b^2)x + 8b(a^3 + 8b^2)}{(3x^4 + 6ax^2 + 12bx - a^2)^2}.$$

(iii) The degree and leading coefficient of ψ_n^2 are $n^2 - 1$ and n^2 , respectively.

- (iv) The degree and leading coefficient of ϕ_n are n^2 and 1, respectively.
- (v) The coefficient of x^{n^2-2} in ψ_n^2 and the coefficient of x^{n^2-1} in ϕ_n are both 0.
- (vi) If a positive integer d divides both a and b , then d divides both $\psi_n^2 - n^2x^{n^2-1}$ and $\phi_n - x^{n^2}$.
- (vii) Let d, n, w, z be integers with $d \geq 1, n \geq 2$ and $w, z \neq 0$. Let $x = \frac{w}{z}$. Then $z^{n^2-3}(\psi_n^2 - n^2x^{n^2-1})$ and $z^{n^2-2}(\phi_n - x^{n^2})$ are integers. Furthermore, if d divides both a and b then d divides both $z^{n^2-3}(\psi_n^2 - n^2x^{n^2-1})$ and $z^{n^2-2}(\phi_n - x^{n^2})$.

6 Proofs

For the remainder of the paper, let $a = -3k^2l^2$ and $b = k^2l^2(k^2 + l^2)$.

Then $E(\mathbb{Q}) = \{(x, y) \in \mathbb{Q}^2 : y^2 = x^3 + ax + b\} \cup \{\infty\}$.

Note that k^2l^2 is a positive integer that divides both a and b .

Also, if $(x, y) \in E(\mathbb{Q}) - \{\infty\}$, then both x and y are nonzero. For if $x = 0$, then $y^2 = k^2l^2(k^2 + l^2)$, $k^2 + l^2$ is a square, and one of k and l has a square factor, a contradiction. Hence, $x \neq 0$. If $y = 0$, then $\exists (u, v) \in E(\mathbb{Q})$ such that $\left(\frac{kl(ku-lv)}{lu-kv}, \frac{kl(k^2-l^2)}{lu-kv}\right) = \gamma(u, v) = (x, 0)$, $\frac{kl(k^2-l^2)}{lu-kv} = 0$, $kl(k^2 - l^2) = 0$, $k^2 - l^2 = 0$, and $k = l$, a contradiction. Hence, $y \neq 0$.

Proof of Lemma 1

We have $k, l \in \mathbb{Z}$, $k, l > 0$, $k \neq l$, $\gcd(k, l) = 1$, and k and l are square-free. We may assume without loss of generality that $2|k$. Then $v_2(k) = 1$ and $v_2(l) = 0$. Since $v_2(k) = 1$, $\exists j \in \mathbb{Z} - \{0\}$ such that $k = 2j$ and $v_2(j) = 0$. Then $y^2 = x^3 - 3k^2l^2x + k^2l^2(k^2 + l^2) = x^3 - 3(2j)^2l^2x + (2j)^2l^2((2j)^2 + l^2) = x^3 - 12j^2l^2x + 4j^2l^2(4j^2 + l^2)$ and $x(2(x, y)) = \frac{x^4 + 6k^2l^2x^2 - 8k^2l^2(k^2 + l^2)x + 9k^4l^4}{4y^2} = \frac{x^4 + 6(2j)^2l^2x^2 - 8(2j)^2l^2((2j)^2 + l^2)x + 9(2j)^4l^4}{4(x^3 - 12j^2l^2x + 4j^2l^2(4j^2 + l^2))} = \frac{x^4 + 24j^2l^2x^2 - 32j^2l^2(4j^2 + l^2)x + 144j^4l^4}{4(x^3 - 12j^2l^2x + 4j^2l^2(4j^2 + l^2))}$.

Case 1: $v_2(x) \geq 2$. Then $(\exists n \geq 2)(\exists u \in \mathbb{Z} - \{0\})$ such that $x = 2^n u$ and

$$v_2(u) = 0. \text{ Then } x(2(x, y)) = \frac{(2^n u)^4 + 24j^2 l^2 (2^n u)^2 - 32j^2 l^2 (4j^2 + l^2)(2^n u) + 144j^4 l^4}{4((2^n u)^3 - 12j^2 l^2 (2^n u) + 4j^2 l^2 (4j^2 + l^2))} =$$

$$\frac{2^{n+1}(2^{3n-5} u^4 + 2^{n-2} 3j^2 l^2 u^2 - j^2 l^2 (4j^2 + l^2)u) + 9j^4 l^4}{2^n(2^{2n-2} u^3 - 3j^2 l^2 u) + j^2 l^2 (4j^2 + l^2)},$$

the numerator and denominator of $x(2(x, y))$ are both odd, and $v_2(x(2(x, y))) = 0$.

Case 2: $v_2(x) = 1$. Then $\exists u \in \mathbb{Z} - \{0\}$ such that $x = 2u$ and $v_2(u) = 0$.

Since $v_2(u) = v_2(j) = v_2(l) = 0$, $\exists f, g, h \in \mathbb{Z}$ such that $u = 2f + 1$, $j = 2g + 1$, and $l = 2h + 1$.

$$\text{Then } x(2(x, y)) = \frac{(2u)^4 + 24j^2 l^2 (2u)^2 - 32j^2 l^2 (4j^2 + l^2)(2u) + 144j^4 l^4}{4((2u)^3 - 12j^2 l^2 (2u) + 4j^2 l^2 (4j^2 + l^2))}$$

$$= \frac{(2(2f+1))^4 + 24(2g+1)^2(2h+1)^2(2(2f+1))^2 - 32(2g+1)^2(2h+1)^2(4(2g+1)^2 + (2h+1)^2)(2(2f+1)) + 144(2g+1)^4(2h+1)^4}{4((2(2f+1))^3 - 12(2g+1)^2(2h+1)^2(2(2f+1)) + 4(2g+1)^2(2h+1)^2(4(2g+1)^2 + (2h+1)^2))}.$$

This expression for $x(2(x, y))$ simplifies to a fraction whose numerator is 4 times an odd number and whose denominator is odd. Hence, $v_2(x(2(x, y))) = v_2(4) = 2$.

Case 3: $v_2(x) = 0$.

$$\text{Then } \exists u \in \mathbb{Z} \text{ such that } x = 2u+1 \text{ and } x(2(x, y)) = \frac{x^4 + 24j^2 l^2 x^2 - 32j^2 l^2 (4j^2 + l^2)x + 144j^4 l^4}{4(x^3 - 12j^2 l^2 x + 4j^2 l^2 (4j^2 + l^2))}$$

$$= \frac{(2u+1)^4 + 24j^2 l^2 (2u+1)^2 - 32j^2 l^2 (4j^2 + l^2)(2u+1) + 144j^4 l^4}{4((2u+1)^3 - 12j^2 l^2 (2u+1) + 4j^2 l^2 (4j^2 + l^2))}$$

$$= \frac{8(u+3j^2 l^2 - 4j^2 l^4 - 16j^4 l^2 + 18j^4 l^4 + 3u^2 + 4u^3 + 2u^4 + 12j^2 l^2 u^2 + 12j^2 l^2 u - 8j^2 l^4 u - 32j^4 l^2 u) + 1}{4(2(3u - 6j^2 l^2 + 2j^2 l^4 + 8j^4 l^2 + 6u^2 + 4u^3 - 12j^2 l^2 u) + 1)},$$

the numerator of $x(2(x, y))$ is odd, the denominator of $x(2(x, y))$ is 4 times an odd number, and $v_2(x(2(x, y))) = -2$.

If $v_2(x) = 0$, then $v_2(x(2_E(x, y))) = -2$ and $2_E(x, y)$ is not an integral point.

If $v_2(x) = 1$, then $v_2(x(2_E(x, y))) = 2$, $v_2(x(4(x, y))) = 0$, $v_2(x(8(x, y))) = -2$, and $8(x, y)$ is not an integral point.

If $v_2(x) \geq 2$, then $v_2(x(2_E(x, y))) = 0$, $v_2(x(4(x, y))) = -2$, and $4(x, y)$ is not an integral point.

In all three cases, $8(x, y)$ is not an integral point.

Proof of Lemma 2

We have $k, l \in \mathbb{Z}$, $k, l > 0$, $k \neq l$, $\gcd(k, l) = 1$, and k and l are square-free.

We may assume without loss of generality that $p|k$. Then $v_p(k) = 1$ and $v_p(l) = 0$. Since $v_p(k) = 1$, $\exists j \in \mathbb{Z} - \{0\}$ such that $k = pj$, $v_p(j) = v_p(l) = 0$. Then $a = -3k^2l^2 = -3p^2j^2l^2$ and $b = k^2l^2(k^2 + l^2) = p^2j^2l^2(p^2j^2 + l^2)$. Also, there exist unique $r, u \in \mathbb{Z}$ such that $x = pu + r$ and $r \in \{0, 1, \dots, p-1\}$. Substituting these values for a, b , and x into the formula for $x(3(x, y))$ given by part (ii) of the Proposition, we obtain $x(3(x, y)) = \frac{r\alpha + p^6(64j^6l^{12} + p\beta)}{9r\gamma + 9p^6\delta^2} = \frac{p\epsilon + r^9}{9r\gamma + 9p^6\delta^2}$, where $\alpha, \beta, \gamma, \delta$, and ϵ are integers. If $r = 0$, then $x(3(x, y)) = \frac{p^6(64j^6l^{12} + p\beta)}{9p^6\delta^2} = \frac{p\beta + 64j^6l^{12}}{9\delta^2}$. If $r \neq 0$, then $x(3(x, y)) = \frac{p\epsilon + r^9}{9r\gamma + 9p^6\delta^2}$. In both cases, p does not divide the numerator of $x(3(x, y))$ and $v_p(x(3(x, y))) \leq \begin{cases} -2 & \text{if } p = 3, \\ 0 & \text{if otherwise.} \end{cases}$

Proof of Lemma 3

We have $k, l \in \mathbb{Z}$, $k, l > 0$, $k \neq l$, $\gcd(k, l) = 1$, and k and l are square-free. We may assume without loss of generality that $p|k$. Then $v_p(k) = 1$ and $v_p(l) = 0$. By Lemma 2, $v_p(x(3(x, y))) \leq 0$. Hence, $\exists w, z \in \mathbb{Z} - \{0\}$ such that $x(3(x, y)) = \frac{w}{z}$ and $v_p(w) = 0$. Since k^2l^2 divides both a and b , part (vii) of the Proposition implies that $z^{p^2-3}(\psi_p^2 - p^2(\frac{w}{z})^{p^2-1})$ and $z^{p^2-2}(\phi_p - (\frac{w}{z})^{p^2})$ are integers, each of which is divisible by k^2l^2 . Hence, $x(3p(x, y)) = x(p(3(x, y))) = x(p(x(3(x, y)), y(3(x, y)))) = x(p(\frac{w}{z}, y(3(x, y)))) = \frac{\phi_p}{\psi_p^2} = \frac{z^{p^2-2}\phi_p}{z^{p^2-2}\psi_p^2} = \frac{z^{p^2-2}\phi_p}{z^{2p^2-3}\psi_p^2} = \frac{z^{p^2-2}(\phi_p - (\frac{w}{z})^{p^2} + \frac{wp^2}{z^{p^2}})}{z^{2p^2-3}(\psi_p^2 - p^2(\frac{w}{z})^{p^2-1} + \frac{p^2wp^2-1}{z^{p^2-1}})} = \frac{z^{p^2-2}(\phi_p - (\frac{w}{z})^{p^2}) + z^{p^2-2}(\frac{wp^2}{z^{p^2}})}{z(z^{p^2-3}(\psi_p^2 - p^2(\frac{w}{z})^{p^2-1}) + z^{p^2-3}(\frac{p^2wp^2-1}{z^{p^2-1}}))} = \frac{k^2l^2\sigma + \frac{wp^2}{z^2}}{z(k^2l^2\tau + \frac{p^2wp^2-1}{z^2})}$, where $\sigma, \tau \in \mathbb{Z}$. Since $v_p(k) = 1$, $\exists j \in \mathbb{Z}$ with $j \geq 1$ such that $k = pj$ and $v_p(j) = 0$. Then $x(3p(x, y)) = \frac{(pj)^2l^2\sigma + \frac{wp^2}{z^2}}{z((pj)^2l^2\tau + \frac{p^2wp^2-1}{z^2})} = \frac{p^2j^2l^2\sigma z^2 + wp^2}{p^2z(j^2l^2\tau z^2 + wp^2-1)}$ and $v_p(x(3p(x, y))) \leq -2$.

References

- [1] A. Holm, Some points in Diophantine analysis, *Proc. Edinburgh Math. Soc.* 22 (1903), 40-48.
- [2] A. Rajan and F. Ramaroson, Ratios of congruent numbers, *Acta Arithmetica* 128.2 (2007), 101-106.
- [3] J. McKee, Computing Division Polynomials, *Mathematics of Computation* 63.208 (October 1994), 767-771.